Unravelling processes between phenotypic plasticity and population

dynamics in migratory birds

Jin Liu[1, 2], Weipan Lei[1], Xunqiang Mo[3], Chris J. Hassell[4], Zhengwang Zhang[1*], Tim

Coulson[2]

1. Key Laboratory for Biodiversity Science and Ecological Engineering, College of

Life Sciences, Beijing Normal University, Beijing, 100875, China.

2. Department of Zoology, University of Oxford, Oxford, OX1 3SZ, UK

3. School of Geographic and Environmental Sciences, Tianjin Normal University,

Tianjin, 300387, China

4. Global Flyway Network, PO Box 3089, Broome, WA 6725, Australia

*Corresponding author: zzw@bnu.edu.cn

Abstract

1. Populations can rapidly respond to environmental change via adaptive phenotypic plasticity, which can also modify interactions between individuals and their environment, affecting population dynamics. Bird migration is a highly plastic resource-tracking strategy in seasonal environments. However, the link between the population dynamics of migratory birds and migration strategy plasticity is not well understood.

2. The quality of staging habitats affects individuals' migration timing and energy budgets in the course of migration, and can consequently affect individuals' breeding and overwintering performance, and impact population dynamics. Given staging habitats being lost in many parts of the world, our goal is to investigate responses of individual migration strategies and population dynamics in the face of loss of staging habitat, and to identify the key processes connecting them.

3. We started by constructed and analysed a general full-annual-cycle individual-based model with a stylized migratory population to generate hypotheses on how changes in the size of staging habitat might drive changes in individual stopover duration and population dynamics, and to identify the key processes connecting them. Next, through the interrogation of census data, we tested these hypotheses by analysing population trends and stopover duration of migratory waterbirds experiencing loss of staging habitat.

4. We found empirical support for our modelling-identified hypotheses: the loss of staging habitat generates plasticity in migration strategies, with individuals

remaining on the staging habitat for longer to obtain food due to a reduction in per

capita food availability. The subsequent increasing population density on the

staging habitat has knock on effects on population dynamics in the breeding and

overwintering stage.

5. Our results demonstrate how environmental change that impacts one energetically

costly life history stage in migratory birds can have population dynamics impacts

across the entire annual cycle via phenotypic plasticity.

Key words: annual cycle, carrying capacity, density-dependent, loss of staging

habitat, individual-based model, migratory birds, population dynamics, stopover

duration

Introduction

Populations can rapidly respond to environmental change via adaptive phenotypic plasticity, and this allows them to cope with profound environmental impacts (Pigliucci 2001, Piersma and Drent 2003, Coulson et al. 2017). Plasticity modifies interactions between individuals and their environment, ultimately affecting population dynamics (Miner et al. 2005). Migration can be an adaptive plastic strategy in seasonal environments (Lack 1968, Newton 2010), that allows individuals to increase reproductive output by avoiding unsuitable ecological conditions (Hedenström 2008, Winkler et al. 2014). Plasticity of migration strategies enables migratory species to respond to environmental changes in multiple ways, such as by altering migratory routes (Sutherland and Crockford 1993, Dolman and Sutherland 1995), timing of migration (Gienapp et al. 2007, Balbontín et al. 2009), and through diet (Parrish 2000). However, the link between the population dynamics of migratory species and migration strategy plasticity is not well understood.

Bird migration is a resource-tracking strategy that aims to optimize a bird's energy budget in the face of fluctuating resources in seasonal environments, and in the face of strong competition (Cox 1968, Alerstam et al. 2003, Somveille et al. 2018, Winger et al. 2019). Migration is energetically costly, so birds build up fat reserves. However, carrying a large energy reserve increases flight costs and can also attract predators (Alerstam and Lindström 1990). One strategy to minimise such costs is to stop over several times during the journeys between breeding and wintering sites to refuel (Piersma 1987). For individuals to remain in favourable environments across

their migration route, they must carefully manage the timing of departure and arrival

(Alerstam and Lindström 1990, Alerstam et al. 2003, Winkler et al. 2014). In general,

individuals that arrive at breeding grounds earlier have higher reproductive success

than those that arrive later (Marra et al. 1998, Norris et al. 2004), and selection

favours individuals that minimize the time spent travelling during the northward

migration (Lindstrom and Alerstam 1992). Migratory birds usually spend much longer

accumulating energy reserves in staging areas than in flying (Hedenström and

Alerstam 1997). Therefore, the total time spent on migration is consequently strongly

influenced by the quality of, and an individual's behaviour at, staging areas

(Hedenström and Alerstam 1997, Erni et al. 2002).

For migratory species, all stages of the annual cycle are closely linked at both the

individual- and population-level, through carry-over and density-dependent effects

(Newton 2010, Harrison et al. 2011). The individual state in one stage can influence

individual performance in subsequent stages, and the change in population size in one

stage can influence per capita rates and consequently regulate population size in later

stages (Marra et al. 1998, Studds and Marra 2005, Ryan Norris and Marra 2007,

Ratikainen et al. 2008). The strategy an individual follows while at the staging area

can consequently affect breeding and overwintering performance, and impact

population dynamics.

Staging habitat for migratory waterbirds in the Yellow Sea is being lost in

significant quantities, primarily due to land reclamation for infrastructure

development and aquaculture (Yang et al. 2011, Bi et al. 2012, Murray et al. 2014,

Wang et al. 2014). It has been suggested as the main contribution to population

declines of migratory birds along the East Asian Australasian Flyway (EAAF)

(Amano et al. 2012, Ma et al. 2014, Piersma et al. 2016, Studds et al. 2017),

presumably because staging habitat in this system is the stage of the annual cycle

where density-dependence is strongest (Sutherland 1996b). Habitat loss in staging

habitats can reduce food resources, decrease foraging and fat-accumulation rates of

migrants (Baker et al. 2004, Morrision 2006, Verkuil et al. 2012), increase

competition and interference in the population, and can have significant consequences

for population regulation (Sutherland 1996b, Newton 2010). However, the way in

which individuals respond to such changes, as well as the processes and mechanisms

that cause population declines, are yet to be generally established. Migratory

waterbirds along the EAAF provides a unique system to explore how individuals

respond to changes in one life history stage, and how this response would influence

their populations.

In this study, we use theoretical modelling to generate hypotheses that we next

test with empirical data to examine the effects of habitat loss within staging areas on

individual migration strategies and population dynamics, and the process that connect

them. First, we conducted an individual-based modelling (IBM) exercise, building a

stylized full-annual-cycle model in which individuals follow the same migrating rules,

to investigate how variation in the size of staging habitat drives changes on individual

stopover duration and energy reserves, population density, and affects life-history

process including reproduction and survival at different life cycle stages. We used this

model to make predictions about changes in strategy might influence population

dynamics in the staging, breeding, and overwintering grounds. We did this by

examined where in the life cycle density-dependence operated most strongly. Next,

using 13 years of census data, we examined whether observed empirical trends were

consistent with hypotheses we generated from our individual-based model. Our

empirical analyses were consistent with the theoretical hypotheses our model

suggested: we found that the loss of staging habitat generates plasticity in migration

strategy, with individuals remaining in the staging habitat for longer to obtain food

due to a reduction in per capita food availability. The subsequent increasing

population density in the staging habitat has knock on effects on breeding and

overwintering stage that impact the population dynamics, via modified survival and

reproduction rates. We conclude that environmental change effects on one life history

stage in migratory birds can consequently have population dynamics impacts across

the entire annual cycle via phenotypic plasticity.

Materials and Methods

The individual-based model (IBM)

The IBM we constructed is a stylized model. The basic assumptions of our model are:

1) the stages of the annual cycle in our model includes northward migration, staging in

the course of northward migration, breeding, southward migration, and overwintering;

2) all individuals followed the same set of rules in our models; 3) individuals who meet

the condition for reproduction produce once each year; 4) males are not limiting and

can be ignored such that we construct a female-only model.

Model description

Our IBMs includes three types of habitats in the model landscape, which are

wintering habitat, breeding habitat and staging habitat. The staging habitat split into

two habitat types – S1 and S2 area. In order to simulate the process of habitat loss in

the staging habitat, the size of S1 area remained constant across all simulations, while

the habitat size of S2 area was adjusted (Fig.S1a-c). The rest of the grid cells in the

landscape are non-habitat, where individuals pass by during migration but do not stop.

Each grid cell within those three types of habitats contained renewing food resources,

food resources renewed each time step at the habitat-specified food recovery rate after

consumption. Each time step in this model represented one day such that one year was

comprised of 365 steps.

We only considered the female component of the population and characterised

each individual by identity, age, reproduction status, and energy reserves. The

behaviours of each individual in the model included: fly, move, search for food, eat,

orient, mature, reproduce and die. Fly was the movement across habitats during the

course of migration, the speed of "fly" was three grid cells per time step. Move was

the movement within each habitats during the course of overwintering, staging and

breeding, the speed of "move" was one grid cell per time step. Birds lost energy

reserves through either type of movement in our model. Birds searched for food by

following a random movement rule, towards the grid cell with the highest food value,

ate and increased their energy reserves when food was available, but there was no

randomness of food acquisition, as long as the food resources was available, the

behaviour of "eat" happened and energy was stored. When the energy reserve of an

individual reached the energy threshold for departure, or the time reached for the

latest possible departure arrived, the individual first oriented, adjusting its facing to

the central grid cell of the next destination habitat in the next time step, then flied

towards it. Individuals whose energy reserves reached the threshold for reproduction

got matured and reproduced once each year. The order of "mature" and "fly to the

breeding habitat" is different in the models with different breeding strategies (see

below). Reproduction also consumed individuals' energy reserves. Newborns were set

as an initial value of age, reproduction status, and energy reserves. Individuals aged

15, or with zero energy reserves, died and were removed from the population. Details

of events and decisions of the models are provided in Appendix S2 (supporting

information).

The energy reserves of each individual was assumed to be dependent on their

initial energy, energy gained, and energy expended. The expected energy gained from

food relied on both population density and food density (Goss-Custard et al. 2002),.

The stopover duration of a bird in the staging habitat was related to the energy

requirement for migration, and the rate of energy acquisition (Hedenström and

Alerstam 1997). In our model, the conditions for leaving the staging habitat were

either the energy reserves exceeded the energy threshold or time passed the time

threshold. Parameter values are provided in Table S1 and Table S2 in Appendix1,

details of individual energy reserves and stopover duration are provided in Appendix

S2 (supporting information).

Model implementation

Our main goal is to examine the effects of habitat change in the staging habitat

on individual stopover duration in the course of northward migration and population

dynamics across the entire life cycle, and to identify the processes that connect them.

To examine the role of the staging in the course of northward migration across the

whole life cycle to the hypothesized processes, we test impacts of reducing carrying

capacity at other life cycle stages (breeding stage and wintering stage). In addition,

since individuals with different breeding strategies (capital breeding and income

breeding) has different energy budgets along the life cycle, we also tested effects of

breeding strategies on model outputs.

Five types of models were constructed in our study, with models differing in the

size of the S2 area, the size of the breeding habitat and the size of the wintering

habitat, the food recovery rate in each of these three habitats, and the breeding

strategies individuals followed (Fig S1, Table S1 in Appendix S1). More specifically:

- Model 1 is the null model, with the same size and the food recovery rate

at all three habitats, and a "capital breeding" strategy for reproduction.

- Model 2 examined the effects of the size of the staging habitat on

individual strategies and population dynamics, with eight scenarios where

the size of S2 area was varied were run, with the lowest food recovery

rate at the staging habitat and a "capital breeding" strategy for

reproduction.

•   Model 3 examined effects of the lowest carrying capacity at the breeding

habitat to individual strategies and population dynamics, with the lowest

size and food recovery rate at the breeding habitat and a "capital

breeding" strategy for reproduction.

•   Model 4 examined effects of the lowest carrying capacity at the wintering

habitat to individual strategies and population dynamics, with the lowest

size and food recovery rate at the wintering habitat and a "capital

breeding" strategy for reproduction.

•   Model 5 examine the effects of breeding strategies to individual stopover

strategies and population dynamics, with the same settings of habitat size

and food recovery rate with model 2, but with an "income breeding"

strategy for reproduction.

The average daily population density and the total number of individuals,

individual stopover duration, individual energy reserves and energy accumulation rate

during stopover, were recorded at equilibrium, and were examined by comparing the

mean value of simulation results. The model was run 10 times for each scenario, and

for 30 years in each simulation, which was sufficient to converge to stationary

dynamics. Results were obtained from year five to year 30 of the simulation (Fig S3,

Appendix S1).

The average daily population density was the average of daily bird number in the

habitat during the period of the current life cycle stage each year, it was recorded at

the S1 area. The total number of individuals was recorded for the population and three

types of habitats each year respectively. Individuals were recorded in age classes and

reproduction status (juveniles, breeding adults, nonbreeding adult).

The per capita reproduction rate for adults, the reproduction rate among breeding

adults and the survival rate were all calculated. The per capita reproduction rate was

the ratio of the number of juveniles and the number of all adults in the breeding stage;

the reproduction rate among breeding adults was the ratio of the number of juveniles

and the number of breeding adults; survival rate was split into two periods, one is the

survival rate of the northward migration (the ratio between the number of adults at the

breeding habitat and the number of individuals on the first day of each year), one is

the survival rate of the southward migration (the ratio between the number of

individuals at the wintering habitat and the number of individuals at the breeding

habitat).

Mean individual stopover duration in the S1 area was recorded each year.

Individual departure energy reserves was recorded for three types of habitats

respectively, as the energy reserves at the last day before the individual left the habitat

range. Individual energy reserves during the stopover period was recorded at each

time step when the individual stayed in the staging habitat. The energy accumulation

rate during stopover was calculated by dividing energy gained from the staging

habitat by stopover duration.

Sensitivity analysis

To examine the impacts of parameter values on the outputs of model 2 and to test

the robustness of the model results were to variation in parameter values, we

conducted a local sensitivity analysis (methods and results see Appendix S3 in the

supporting information).

Empirical data

Our empirical study was conducted in the wetlands in the north of Bohai Bay,

between 38°36'-39°13'N and 117°11'-118.22'E, located in the north-west of the

Yellow Sea. Census data of migratory waterbirds were collected at boreal spring

between 2004 and 2018 (details of study area and data collection are provided in

Appendix S5, supporting information).

Statistical analyses for bird number trends

The abundance of all waterbirds, the abundance of the most common species,

and waterbird species richness were analyzed in this study. Survey sites that were

surveyed on less than 30% of survey dates were excluded from the analysis.

To estimate total waterbird abundance, we summed the counts of all species i

(including unidentified species) observed from all of the survey sites k in each day j,

denoting it as N_a. To identify the most common species, we used two methods: the

"Frequency Based Method" and the "Distribution Based Method" to select 25

common species (Appendix S5). The number of these 25 species observed on each

day j and survey site k was denoted as N_b. To estimate waterbird richness, we

calculated the number of species observed each day j each survey site k, excluding

unidentified species, denoting it as M.

We calculated survey effort as the number of observers each day at each site as

$E_{j,k}$. We transformed date to Julian date t, and calculated t^2 as visual examination of

the data revealed a quadratic relationship. We treated 'year' (y) as a continuous

variable in our models, and we treated 'survey site' (k) as a categorical variable.

Waterbird abundance and richness (including N_a, N_b and M) were used as

response variables, $E_{j,k}$, k, t, t^2, i and y were explanatory variables. The distribution of

N_a, N_b and M was well described as an over-dispersed Poisson (variance greater than

mean), so we fitted Generalized Linear model (GLM) with a "quasi-Poisson" error

structure in program R. The regression equations were of the form:

$N_a = \exp(\alpha_1 + \beta_{11}(y) + \beta_{12}(E_{j,k}) + \beta_{13}(y \times E_{j,k}) + \beta_{14}(k) + \beta_{15}(y \times k) + \beta_{16}(t) + \beta_{17}(t^2) +$

$\beta_{18}(y \times t) + \beta_{19}(y \times t^2) + \varepsilon_1)$ (1)

$N_b = \exp(\alpha_2 + \beta_{21}(y) + \beta_{22}(E_{j,k}) + \beta_{23}(y \times E_{j,k}) + \beta_{24}(k) + \beta_{25}(y \times k) + \beta_{26}(t) + \beta_{27}(t^2) +$

$\beta_{28}(y \times t) + \beta_{29}(y \times t^2) + \beta_{210}(i) + \beta_{211}(y \times i) + \varepsilon_2)$ (2)

$M = \exp(\alpha_3 + \beta_{31}(y) + \beta_{32}(E_{j,k}) + \beta_{33}(y \times E_{j,k}) + \beta_{34}(k) + \beta_{35}(y \times k) + \beta_{36}(t) + \beta_{37}(t^2) +$

$\varepsilon_3)$ (3)

Statistical analyses for stopover duration trends

We estimated the stopover duration for each common species by using our

census data. Since the quadratic relationship between Julian date and bird abundance

revealed from our census data (Fig. S4 a), we first estimated normal distributions of

bird abundance within the period of stopover for each species each year (Fig. S4b). To

do this, we extracted the mean date and variance in date for each species each year

from census data, and scaled the curves by the bird number from the census data.

Then we estimated the date by which each quantile of the distribution of bird

abundance is reached, date at 2.5% quantile and 97.5% quantile was the date of

arrival at staging habitat and the date of departure from staging habitat for each

species each year within 95% confidential interval respectively (Fig. S4c). We

calculated the stopover duration based on the arrival date and departure date for each

species each year, denoting it as $T_{i,y}$ (details are provided in S5.4 Appendix S5).

The distribution of $T_{i,y}$ was normal distribution, so we fitted Linear model in

program R to test the stopover duration trends across years for each common species,

the regression equation was of the form:

$$T_{i,y} = \alpha_4 + \beta_{41}(y) + \beta_{42}(i) + \beta_{43}(y \times i) + \varepsilon_4 \quad (4)$$

The correlation between stopover duration and bird abundance

We fitted a regression between bird abundance and stopover duration to test the

correlation between them. We calculated the mean abundance during the stopover

period for each species i each year y, denoting it as N_c. N_c was used as the response

variable, y, i, $T_{i,y}$ were explanatory variables. The distribution of N_c was an over-

dispersed Poisson, so we fitted GLM with a "quasi-Poisson" error structure in

program R. The regression equations were of the form:

$N_c = \exp(\alpha_5 + \beta_{51}(y) + \beta_{52}(i) + \beta_{53}(y \times i) + \beta_{54}(T_{i,y}) + \beta_{55}(y \times T_{i,y}) +$

$\beta_{56}(i \times T_{i,y}) + \varepsilon_5) \quad (5)$

We used the ANOVA command in R to assess the significance of each variable,

used adjusted-R^2 to assess the goodness of fit of the linear model, and used 1-

(Residual Deviance/Null Deviance) to assess the goodness of fit of the GLMs.

Results

IBM: population dynamics and individual stopover duration at the staging

habitat

As the size of S2 area decreased through eight scenarios in the model 2

simulations, there was a decrease in the total number of individuals, and an increase in

average daily population density at the S1 area (Fig. 1a). The curve of daily

population density at the S1 area became steeper in the early and late phases of the

stopover period, and higher and wider when the population reached peak numbers

(Fig 1b), showing that the population took less time to reach the peak number and

remained at the peak number for a longer period of time. Individual's annual stopover

duration at the S1 area increased (Fig 1c), suggesting individuals stayed longer in the

S1 area as the size of S2 decreased.

Figure 1. Population dynamics and stopover duration as the size of the staging habitat

decreased across the eight scenarios in model 2 simulations. a) The relationship

between average daily population density at the S1 area and the total number of

individuals; b) The curves of population density at the S1 area during the period of

stopover; c) Individual annual stopover duration in relation to the size of staging

habitat.

IBM: Individual energy reserves

As the size of S2 area decreased in model 2 simulations, the individual energy

accumulation rate when birds remained in the staging habitat decreased (Fig 2a), the

distribution shifted towards lower energy reserves with fewer individuals having

reached the energy threshold for departure from the staging habitat (Fig 2b). The

energy reserves when individuals left the staging habitat was decreased, however, the

departure energy reserves from the breeding habitat and wintering habitat was

increased, both in adults and juveniles (Fig 2c).

Figure 2. Individual energy reserves across the annual cycle in model 2 simulations.

a) The energy accumulating rate during the period of stopover when individuals at the

staging habitat. b) The distribution of individual departure energy reserves from the

staging habitat. c) Individual departure energy reserves at each annual cycle stage.

IBM: Population structure and demography

As the size of S2 area decreased in model 2 simulations, the proportion of

breeding adults and juveniles in the population decreased significantly, while the

proportion of nonbreeding adults increased in each life cycle stage (Fig 3a). The per

capita reproduction rate decreased because of an increased proportion of nonbreeding

adults, but the reproduction rate among breeding adults increased. The survival rate in

both the northward migration and southward migration increased (Fig 3b).

Figure 3. Population structure and demography as the size of the staging habitat

decreased across the eight scenarios in model 2 simulations. a) The proportion of

individuals in different age groups and maturity status in three stages of the annual

cycle. b) per capita rates of reproduction for all adults, reproduction for breeding

adults, survival during southward migration and survival during northward migration.

IBM: the comparison between different model settings

By comparing the results from models (model 2, 3, 4) with the lowest carrying

capacity at different life cycle stages and the results from the null model (model 1),

we found only when the lowest carrying capacity was at the staging habitat (model 2),

population density at the S1 area increased, individuals remained longer at the staging

habitat, with lower energy reserves when leaving the staging habitat, leading to fewer

breeding adults and juveniles in the population and lower per capita reproduction rate.

Individuals do breed had better performance comparing to other models, with higher

energy reserves when leaving the breeding and wintering habitat, and higher survival

rate during migration. However, when the lowest carrying capacity was at other life

cycle stages (model 3 - breeding stage or model 4 - wintering stage), the processes

was completely different. By comparing the results from models with different

breeding strategy settings (model 2 – capital breeding, model 5 – income breeding),

we found different breeding strategies did not influence the processes linking

migration strategy and population dynamics across the life cycle (Fig 4, details are

provided in Appendix S4).

Figure 4. Comparison of models outputs. The name of results are in the form of "the

type of results_habitat type (or the whole population)_age classes". Six types of

results are listed, "density" is the average daily population density, "duration" is the

individual stopover duration, "energy" is the energy reserves when leaving the habitat,

"individual" is the number of individuals, "proportion*100" is the one hundred fold

proportion of individuals of each age class in the population, "rate*100" is the one

hundred fold vital rates. Four types of habitat are listed, including the S1 area, the

staging habitat, the breeding habitat, and the wintering habitat. Age classes includes

juveniles and adults, adults were split into two types: breeding adults and non-

breeding adults. The vertical black dotted line represents the position of the zero

value. The distance from the point to the zero value line represents the difference

between the results of the model (model 2-5) and the results of the null model (model

1).

Empirical data: overall abundance

Analysis of the abundance of all recorded species revealed a significant increase

(Fig.5a) in numbers of waterbirds at a rate of 61.85% per year ($F_{1, 809} = 13.60$, $p <$

0.001). Even though effort significantly affected abundance estimates ($F_{1, 808} = 11.04$,

$p < 0.001$) with more birds counted as effort increased; there was no interaction

between year and effort ($F_{1, 787} = 0.094$, $p = 0.76$). There was a positive relationship

between waterbird abundance and Julian date ($F_{1, 789} = 5.98$, $p = 0.015$). This slowed,

and became negative with time, as the quadric term of date ($F_{1, 788} = 10.34$, $p =$

0.0014) indicated a parabola of waterbird abundance within the migration season.

There was also a significant interaction between year and the quadric term of date

($F_{1,768} = 17.96$, $p < 0.001$), showing that the parabola shape within the migration

season changed across years. The average waterbird abundance ($F_{18, 790} = 30.81$, $p <$

0.001), and the annual trend ($F_{18, 769} = 3.59$, $p < 0.001$) differed between survey sites

(Fig.S8a). The goodness of fit of the GLM was 0.511.

Figure 5. Trends of waterbirds abundance, richness and stopover duration from the

census data. a) Overall waterbirds abundance as a function of year; b) the abundance

of the most common species as a function of year; c) waterbirds richness as a function

of year; d) stopover duration of common species as a function of year; e) The

correlation between stopover duration and the abundance in common species

Empirical data: the abundance of the most common species

In the analysis of the most common species, the abundance of common species

increased at a rate of 35.13% per year ($F_{1, 18418} = 6.92$, p = 0.009) (Fig.5b), after

correcting for effects from observers effort ($F_{1, 18417} = 47.63$, p < 0.001) and the

interaction between year and effort ($F_{1,18372}=6.94$, p=0.008), survey sites ($F_{18,18399} =$

95.28, p< 0.001) and the interaction between year and survey sites ($F_{18,18354} = 5.96$, p<

0.001) (Fig. S8b), species ($F_{24,18373} = 80.14$, p < 0.001) and the interaction between

year and species ($F_{24,18328} = 6.96$, p < 0.001). There was a quadratic association

between Julian date and the number of birds (linear term, $F_{1, 18398} = 25.77$, p < 0.001;

quadratic term $F_{1, 18397} = 85.23$, p < 0.001), but the interaction between year and Julian

date was not significant (linear term, $F_{1, 18352} = 0.52$, p = 0.470; quadratic term $F_{1, 18353}$

= 2.57, p = 0.109). The goodness of fit of the GLM was 0.443.

Empirical data: waterbird richness

Waterbird richness significantly increased with time ($F_{1, 805} = 28.20$, p < 0.001)

having corrected for effort (Fig.5c), with a rate of 7.93% per year. Effort was not

statistically significant on richness ($F_{1, 804} = 0.26$, p = 0.61), although a weak

interaction between year and effort was observed ($F_{1, 783} = 4.69$, p = 0.031). The

relationship between richness and Julian date within a year was positive, with an

initial rate of increase of 3.76% per day ($F_{1, 785} = 34.50$, p < 0.001). This rate slowed

as Julian date increased, as the relationship between richness and the quadric term of

Julian date was negative ($F_{1, 784} = 63.55$, p < 0.001), indicating an arched parabola in

each migration season. The waterbird richness ($F_{18, 786} = 29.16$, p < 0.001) and yearly

trend ($F_{18, 765} = 4.69$, p < 0.001) differed among survey sites (Fig S8c). The goodness

of fit was 0.487.

Empirical data: stopover duration

Analysis of the stopover duration of common species revealed a significant increasing

trend at the rate of 1.39 days per year ($F_{1,245} = 14.58$, p < 0.001) (Fig.5d). The

stopover duration ($F_{24,221} = 7.20$, $p < 0.001$) and its temporal trends ($F_{24,197} = 1.90$, $p =$

0.009) were differed among species (Fig S8d). The adjust R-square of the linear

model was 0.428.

In the analysis of the correlation between bird abundance and the stopover duration,

abundance was positively related with stopover duration ($F_{1,220} = 18.36$, $p < 0.001$)

(Fig.5e), and also positively related with year ($F_{1,245} = 56.41$, $p<0.001$). The

abundance ($F_{24,221} = 35.93$, $p < 0.001$) was differed across species. The relation

between year and abundance ($F_{24,196} = 3.28$, $p < 0.001$), and the relation between

stopover duration and abundance ($F_{24, 172} = 2.24$, $p = 0.002$) was differed across

species. The interaction between year and stopover duration was not significant ($F_{1,171}$

$= 0.70$, $p = 0.404$). The goodness of fit of the GLM was 0.857.

Discussion

By building a full-annual-cycle IBM of a stylized migratory population, we

identify the critical role of stopover stage of northward migration in the influence of

migration strategies and population dynamics of migratory birds across the whole

annual cycle, and also identify the key processes linking individual migration strategy

and population dynamics (Fig 6). Our empirical data provides evidence to support the

mechanisms we showed from our theoretical model. Specifically, our results are

consistent with the loss of staging habitat generating plasticity in migration strategy

via increased intraspecific competition during migration stopovers. As the size of the

staging area is reduced, individuals need to remain in the staging area for longer to

obtain sufficient food to continue on their way due to an increase in the density of

competitors. Our model shows that the consequence of this is individuals depart later,

and often in poor condition, and fewer individuals make it to the breeding area.

However, those that do make it fare well. The dynamics at the staging area can

consequently have knock on effects on populations in the overwintering and breeding

areas that impact the population dynamics across the annual cycle, by altering the

component of the life history where population dynamics are regulated.

Figure 6. The processes linking individual migration strategy and population

dynamics in our study. a) The changes in carrying capacity and population density as

a function of the size of the staging habitat. S1 and S2 represents the two areas of the

staging habitat, B represents the breeding habitat, and W represents the wintering

habitat. The circle sizes represent the carrying capacity of each habitat, with the gray

rectangles representing maximum population size across the annual cycle. In (a1) the

size of the wintering and breeding habitats determine carrying capacity, while in (a2)

the size of the staging habitat determines it. The line charts shows how population

density in the staging areas changes during the period of migration stopover for

scenarios (a1) and (a2) respectively. b) The feedback process between individual

stopover duration and population dynamics.

Habitat loss in the staging habitat reduces the carrying capacity of the flyway,

leading to population declines along it. As a consequence, the part of the annual cycle

that determines the carrying capacity has been switched from the breeding habitat and

wintering habitat to the staging habitat, as the spatial extent of staging habitat

decreases. The staging habitat in the course of northward migration becomes the stage

where the strongest density-dependence operates, and the population becomes

regulated by the carrying capacity of staging habitat (Fig 6a). In contrast, competition

on the breeding or wintering habitat is reduced, with processes operating in these

areas no longer playing a major role in regulating the population dynamics. The

contributions of different life history stages to the population dynamics can

consequently vary with the spatial extent of the staging area. Such changes have the

potential to alter selection on traits associated with competition for resources, and the

entire life history.

Both in our simulations and empirical analysis, population density changes

spatio-temporally following habitat loss in the staging habitat, with a positive

correlation between bird numbers and stopover duration (Fig. 1b, Fig. S9 in Appendix

S1). This contrasts the "buffer effect" hypothesis, which proposes that population

density changes only in space (Brown 1969, Sutherland 1996a, Gill et al. 2001). Our

results show that in addition to birds becoming more concentrated in the remaining

area, a decrease in the extent of the staging area can also result in an increase in time

spent there, leading to intensified competition during staging period. Therefore,

during the time-limited northward migration, the reduced staging area not only leads

to a higher population density in the staging habitat as individual birds stay longer, but

also that the high population density is maintained for a longer period of time during

this part of the life cycle (Fig.6a). And this can alter the strength of population

regulation in either the wintering or breeding areas (Fig. 6b). How do we reach this

conclusions?

In our models, individuals adjust their migration strategy to prolong the stopover

duration when food becomes scarce. As a consequence, individuals often arrive at the

breeding ground late and with low energy reserves, or they fail to reach the breeding

grounds. This results in fewer adults arriving in the breeding ground, fewer adults

reproducing, and a decline in total reproductive output. However, those individuals

that do arrive experience less competition, and have higher departure energy reserves

at the breeding habitat, breeding adults have higher per capita reproductive rate. All

individuals have higher survival rate during southward migration and northward

migration. Previous studies have reported evidence that support parts of the processes

we describe here including: longer stopover duration is related to habitat loss, scarce

food or high density of competitors (Moore and Yong 1991, Kelly et al. 2002, Conklin

et al. 2021), decreased refuelling rate causes poor departure body mass in the staging

area (Baker et al. 2004), limited refuelling time reducing survival rates in the

northward migration (Rakhimberdiev et al. 2018), and reductions in total population-

level reproductive output (Newton 2006, Desprez et al. 2018).

All of this is due to increased competition in the staging habitat, which reduces

per capita food availability, and generates behavioural plasticity in migratory strategy.

As individual behaviour changes, so to do the population dynamics. The altered

population dynamics, in turn, affect individual behaviour (Miner et al. 2005). This

process continues until an equilibrium is reached, when the behaviour settles down to

equilibrium, as do the population dynamics (Fig. 6b). The connection between

individual phenotypic plasticity and population dynamics is the result of feedback

process across the annual cycle that can result in eco-evolutionary feedbacks are

argued by Coulson (2020).

If the part of the annual cycle that determines carrying capacity is not switched to

staging habitat, patterns and processes of individual strategies and population

dynamics can be different (Fig. 4). The staging habitat was no longer the stage

regulating population dynamics, the population density at the staging habitat no

longer affected the trade-off between time and energy budgets in the course of

stopover period, then influencing the life history process. The strongest density-

dependent effects at the breeding habitat or the wintering habitat can cause a decrease

in reproduction rate or survival rate at either stage, leading to a lower population

density at the staging habitat, and a shortened stopover duration. Evidence has been

reported by (Holmes et al. 1996, Rockwell et al. 2017) who observed reduced survival

rates when the wintering habitat is limiting, and (Marra and Holmes 2001,

Rodenhouse et al. 2003, Tomotani et al. 2018) who reported poor physical condition

of juveniles and decreased reproduction when the breeding habitat is limiting. In

addition, even though different breeding strategies may influence individuals' energy

budgets across the life cycle stages, our simulation results from model 2 and model 5

showed similar patterns and processes, indicating both capital breeders and income

breeders would increase their stopover duration when facing habitat loss at staging

habitats. Therefore, whether the changes in the extent of the staging habitat alters the

part of the life cycle where carrying capacity is lowest will determine the direction of

change in life-history processes. Our proposed process can only occur when the

staging habitat become the stage that determine the carrying capacity of the whole

annual cycle.

Our empirical and theoretical results align well, and are consistent with, patterns

reported in the existing literature on the EAAF migration flyway, which are

decreasing trends of total population size along the flyway, increasing trends of

population density (Wilson et al. 2011, Yang et al. 2011, Clemens et al. 2016, Piersma

et al. 2016, Studds et al. 2017) and increasing stopover duration at the remaining

staging area(Conklin et al. 2021). Our study reveals the underlying mechanisms

behind a seemingly positive phenomena observed in migratory birds, while also

highlighting the crucial role of one life history stage to population dynamics across

the whole annual cycle.

We call for further investigation on the connection between individual timing,

energy reserves and demography across the whole annual cycle. Migratory birds need

to be studied in depth at their breeding, wintering and staging areas to allow us to

fully understand their dynamics. Focusing on a single area in detail, with limited data

elsewhere, makes dynamical inference challenging, and may also lead researchers to

reach erroneous conclusions. In addition, migratory species play a key role for a

number of other ecological processes by transporting energy and nutrients between

spatially distinct areas, or via their impacts on the ecological networks they form with

other species in geographically separated areas (Bauer and Hoye 2014), understanding

their dynamics is also important to further extend our knowledge on their roles in

ecosystem stability. Although logistically difficult, and costly, our work suggests that

studies of migratory species across the entire annual cycle are necessary if we are

going to understand the dynamics of species that exhibit one of the most remarkable

behaviours in the animal kingdom.

576
577

Acknowledgements

We thank Hongyan Yang, Pinjia Que, Yajing Chang, Bingrun Zhu, Jingsheng Ma, Jianmin Wang, Paul Holt, Adrian Boyle and the Global Flyway Network team, and many volunteers for help with the fieldwork. This work was supported by the grants from the Natural Science Foundation of China (31830089 and 31572288 to ZZ, 31801985 to WL), the Paulson Institute grant and the SEE Foundation grant to ZZ, the China Scholarship Council grant to JL, the Fundamental Research Funds for the Central Universities to WL. CJH has been funded by GFN with grants secured by Theunis Piersma via BirdLife Netherlands (2007-2012), WWF Netherlands (2010-2014, 2016, 2018), Spinoza Premium of Netherlands Organisation Prize for Scientific Research to Theunis Piersma (2014-2017) and MAVA (foundation pour la nature) in 2018.

Conflict of interest

None of the authors have a conflict of interest.

Authors' contributions

JL, TC and ZZ conceived the ideas and designed methodology. JL, WL, XM, and CJH collected the data. JL built the model, analysed the data and led the writing of the manuscript with significant contributions from TC. All authors contributed critically to the drafts and gave final approval for publication.

Data availability statement

Code and data will be archived if the paper is accepted for publication.

Reference

Alerstam, T., A. Hedenström, and S. Åkesson. 2003. Long-distance migration: evolution and determinants. Oikos **103**:247-260.

Alerstam, T., and Å. Lindström. 1990. Optimal bird migration: the relative importance of time, energy, and safety. Pages 331-351 Bird migration. Springer.

Amano, T., T. Székely, K. Koyama, H. Amano, and W. J. Sutherland. 2012. A framework for monitoring the status of populations: An example from wader populations in the East Asian–Australasian flyway. Biological conservation **145**:278-295.

Baker, A. J., P. M. González, T. Piersma, L. J. Niles, I. de Lima Serrano do Nascimento, P. W. Atkinson, N. A. Clark, C. D. T. Minton, M. K. Peck, and G. Aarts. 2004. Rapid population decline in red knots: fitness consequences of decreased refuelling rates and late arrival in Delaware Bay. Proceedings of the Royal Society of London. Series B: Biological Sciences **271**:875-882.

Balbontín, J., A. P. Møller, I. G. Hermosell, A. Marzal, M. Reviriego, and F. De Lope. 2009. Individual responses in spring arrival date to ecological conditions during winter and migration in a migratory bird. Journal of Animal Ecology **78**:981-989.

Bauer, S., and B. J. Hoye. 2014. Migratory animals couple biodiversity and ecosystem functioning worldwide. Science **344**:1242552.

Bi, X., F. Liu, and X. Pan. 2012. Coastal projects in China: from reclamation to restoration. Environmental Science & Technology **46**:4691-4692.

Brown, J. L. 1969. The buffer effect and productivity in tit populations. The American

Naturalist **103**:347-354.

Clemens, R., D. I. Rogers, B. D. Hansen, K. Gosbell, C. D. Minton, P. Straw, M. Bamford, E. J. Woehler, D. A. Milton, M. A. Weston, B. Venables, D. Weller, C. Hassell, B. Rutherford, K. Onton, A. Herrod, C. E. Studds, C.-Y. Choi, K. L. Dhanjal-Adams, N. J. Murray, G. A. Skileter, and R. A. Fuller. 2016. Continental-scale decreases in shorebird populations in Australia. Emu-Austral Ornithology **116**:119-135.

Conklin, J. R., S. Lisovski, and P. F. J. N. C. Battley. 2021. Advancement in long-distance bird migration through individual plasticity in departure. Nature Communications **12**:1-9.

Coulson, T. 2020. Environmental perturbations and transitions between ecological and evolutionary equilibria: An eco-evolutionary feedback framework. bioRxiv **ver.4 peer-reviewed and recommended by PCI Ecology**:509067.

Coulson, T., B. E. Kendall, J. Barthold, F. Plard, S. Schindler, A. Ozgul, and J.-M. Gaillard. 2017. Modeling adaptive and nonadaptive responses of populations to environmental change. The American Naturalist **190**:313-336.

Cox, G. W. 1968. The role of competition in the evolution of migration. Evolution:180-192.

Desprez, M., S. Jenouvrier, C. Barbraud, K. Delord, and H. Weimerskirch. 2018. Linking oceanographic conditions, migratory schedules and foraging behaviour during the non-breeding season to reproductive performance in a long-lived seabird. Functional ecology **32**:2040-2053.

Dolman, P. M., and W. J. Sutherland. 1995. The response of bird populations to habitat

loss. Ibis **137**:S38-S46.

Erni, B., F. Liechti, and B. Bruderer. 2002. Stopover strategies in passerine bird

migration: a simulation study. Journal of theoretical biology **219**:479-493.

Gienapp, P., R. Leimu, and J. Merilä. 2007. Responses to climate change in avian

migration time—microevolution versus phenotypic plasticity. Climate Research

35:25-35.

Gill, J. A., K. Norris, P. M. Potts, T. G. Gunnarsson, P. W. Atkinson, and W. J.

Sutherland. 2001. The buffer effect and large-scale population regulation in

migratory birds. Nature **412**:436-438.

Goss-Custard, J. D., R. A. Stillman, A. D. West, R. W. G. Caldow, and S. McGrorty.

2002. Carrying capacity in overwintering migratory birds. Biological

conservation **105**:27-41.

Harrison, X. A., J. D. Blount, R. Inger, D. R. Norris, and S. Bearhop. 2011. Carry-over

effects as drivers of fitness differences in animals. Journal of Animal Ecology

80:4-18.

Hedenström, A. 2008. Adaptations to migration in birds: behavioural strategies,

morphology and scaling effects. Philosophical Transactions of the Royal

Society B: Biological Sciences **363**:287-299.

Hedenström, A., and T. Alerstam. 1997. Optimum fuel loads in migratory birds:

distinguishing between time and energy minimization. Journal of theoretical

biology **189**:227-234.

Holmes, R. T., P. P. Marra, and T. W. Sherry. 1996. Habitat-specific demography of

breeding black-throated blue warblers (Dendroica caerulescens): implications

for population dynamics. Journal of Animal Ecology:183-195.

Kelly, J. F., L. S. DeLay, and D. M. Finch. 2002. Density-dependent mass gain by

Wilson's Warblers during stopover. The Auk **119**:210-213.

Lack, D. 1968. Bird migration and natural selection. Oikos:1-9.

Lindstrom, A., and T. Alerstam. 1992. Optimal fat loads in migrating birds: a test of the

time-minimization hypothesis. The American Naturalist **140**:477-491.

Ma, Z., D. S. Melville, J. Liu, Y. Chen, H. Yang, W. Ren, Z. Zhang, T. Piersma, and B.

Li. 2014. Rethinking China's new great wall. Science **346**:912-914.

Marra, P. P., K. A. Hobson, and R. T. Holmes. 1998. Linking winter and summer events

in a migratory bird by using stable-carbon isotopes. Science **282**:1884-1886.

Marra, P. P., and R. T. Holmes. 2001. Consequences of dominance-mediated habitat

segregation in American Redstarts during the nonbreeding season. The Auk

118:92-104.

Miner, B. G., S. E. Sultan, S. G. Morgan, D. K. Padilla, and R. A. Relyea. 2005.

Ecological consequences of phenotypic plasticity. Trends in Ecology and

Evolution **20**:685-692.

Moore, F. R., and W. Yong. 1991. Evidence of food-based competition among passerine

migrants during stopover. Behavioral Ecology and Sociobiology **28**:85-90.

Morrision, R. I. G. 2006. Body transformations, condition, and survival in Red Knots

Calidris canutus travelling to breed at Alert, Ellesmere Island, Canada. Ardea

94:607-618.

Murray, N. J., R. S. Clemens, S. R. Phinn, H. Possingham, and R. A. Fuller. 2014. Tracking the rapid loss of tidal wetlands in the Yellow Sea. Frontiers in Ecology and the Environment **12**:267-272.

Newton, I. 2006. Can conditions experienced during migration limit the population levels of birds? Journal of Ornithology **147**:146-166.

Newton, I. 2010. The migration ecology of birds. Elsevier, UK.

Norris, D. R., P. P. Marra, T. K. Kyser, T. W. Sherry, and L. M. Ratcliffe. 2004. Tropical winter habitat limits reproductive success on the temperate breeding grounds in a migratory bird. Proceedings of the Royal Society of London. Series B: Biological Sciences **271**:59-64.

Parrish, J. D. 2000. Behavioral, energetic, and conservation implications of foraging plasticity during migration. Studies in Avian Biology **20**:53-70.

Piersma, T. 1987. Hop, skip, or jump? Constraints on migration of arctic waders by feeding, fattening, and flight speed. Nederlandse Ornithologische Unie (NOU).

Piersma, T., and J. Drent. 2003. Phenotypic flexibility and the evolution of organismal design. Trends in Ecology & Evolution **18**:228-233.

Piersma, T., T. Lok, Y. Chen, C. J. Hassell, H. Y. Yang, A. Boyle, M. Slaymaker, Y. C. Chan, D. S. Melville, and Z. W. Zhang. 2016. Simultaneous declines in summer survival of three shorebird species signals a flyway at risk. Journal of Applied Ecology **53**:479-490.

Pigliucci, M. 2001. Phenotypic plasticity: beyond nature and nurture. JHU Press.

Rakhimberdiev, E., S. Duijns, J. Karagicheva, C. J. Camphuysen, A. Dekinga, R.

Dekker, A. Gavrilov, J. Ten Horn, J. Jukema, A. Saveliev, M. Soloviev, T. L.

Tibbits, J. A. van Gils, and T. Piersma. 2018. Fuelling conditions at staging sites

can mitigate Arctic warming effects in a migratory bird. Nature

Communications **9**:1-10.

Ratikainen, I. I., J. A. Gill, T. G. Gunnarsson, W. J. Sutherland, and H. Kokko. 2008.

When density dependence is not instantaneous: theoretical developments and

management implications. Ecology Letters **11**:184-198.

Rockwell, S. M., J. M. Wunderle, T. S. Sillett, C. I. Bocetti, D. N. Ewert, D. Currie, J.

D. White, and P. P. Marra. 2017. Seasonal survival estimation for a long-

distance migratory bird and the influence of winter precipitation. Oecologia

183:715-726.

Rodenhouse, N. L., T. Scott Sillett, P. J. Doran, and R. T. Holmes. 2003. Multiple

density–dependence mechanisms regulate a migratory bird population during

the breeding season. Proceedings of the Royal Society of London. Series B:

Biological Sciences **270**:2105-2110.

Ryan Norris, D., and P. P. Marra. 2007. Seasonal interactions, habitat quality, and

population dynamics in migratory birds. The Condor **109**:535-547.

Somveille, M., A. S. Rodrigues, and A. Manica. 2018. Energy efficiency drives the

global seasonal distribution of birds. Nature ecology & evolution **2**:962-969.

Studds, C. E., B. E. Kendall, N. J. Murray, H. B. Wilson, D. I. Rogers, R. S. Clemens,

K. Gosbell, C. J. Hassell, R. Jessop, D. S. Melville, D. A. Milton, C. D. T.

Minton, H. P. Possingham, A. C. Riegen, P. Straw, E. J. Woehler, and R. A. Fuller. 2017. Rapid population decline in migratory shorebirds relying on Yellow Sea tidal mudflats as stopover sites. Nature Communications **8**:14895.

Studds, C. E., and P. P. Marra. 2005. Nonbreeding habitat occupancy and population processes: an upgrade experiment with a migratory bird. Ecology **86**:2380-2385.

Sutherland, W. J. 1996a. From individual behaviour to population ecology. Oxford University Press on Demand.

Sutherland, W. J. 1996b. Predicting the consequences of habitat loss for migratory populations. Proceedings of the Royal Society of London. Series B: Biological Sciences **263**:1325-1327.

Sutherland, W. J., and N. J. Crockford. 1993. Factors affecting the feeding distribution of red-breasted geese Branta ruficollis wintering in Romania. Biological conservation **63**:61-65.

Tomotani, B. M., H. van der Jeugd, P. Gienapp, I. de la Hera, J. Pilzecker, C. Teichmann, and M. E. Visser. 2018. Climate change leads to differential shifts in the timing of annual cycle stages in a migratory bird. Global change biology **24**:823-835.

Verkuil, Y. I., N. Karlionova, E. N. Rakhimberdiev, J. Jukema, J. J. Wijmenga, J. C. Hooijmeijer, P. Pinchuk, E. Wymenga, A. J. Baker, and T. Piersma. 2012. Losing a staging area: Eastward redistribution of Afro-Eurasian ruffs is associated with deteriorating fuelling conditions along the western flyway. Biological conservation **149**:51-59.

Wang, W., H. Liu, Y. Li, and J. Su. 2014. Development and management of land

reclamation in China. Ocean & Coastal Management **102**:415-425.

Wilson, H. B., B. E. Kendall, R. A. Fuller, D. A. Milton, and H. P. Possingham. 2011.

Analyzing variability and the rate of decline of migratory shorebirds in Moreton

Bay, Australia. Conservation Biology **25**:758-766.

Winger, B. M., G. G. Auteri, T. M. Pegan, and B. C. Weeks. 2019. A long winter for the

Red Queen: rethinking the evolution of seasonal migration. Biological Reviews

94:737-752.

Winkler, D. W., C. Jørgensen, C. Both, A. I. Houston, J. M. McNamara, D. J. Levey, J.

Partecke, A. Fudickar, A. Kacelnik, and D. Roshier. 2014. Cues, strategies, and

outcomes: how migrating vertebrates track environmental change. Movement

Ecology **2**:1-15.

Yang, H.-Y., B. Chen, M. Barter, T. Piersma, C.-F. Zhou, F.-S. Li, and Z.-W. Zhang.

2011. Impacts of tidal land reclamation in Bohai Bay, China: ongoing losses of

critical Yellow Sea waterbird staging and wintering sites. Bird Conservation

International **21**:241-259.

770