The hippocampus shows an own-age bias during unfamiliar face viewing

Joshua D. Koen[1]*, Nedra Hauck[2], Michael D. Rugg[2,3]

[1]Department of Psychology, University of Notre Dame

[2]Center for Vital Longevity and School of Behavioral and Brain Sciences, University of Texas at
Dallas

[3]School of Psychology, University of East Anglia, UK

*Correspondence: jkoen@nd.edu

Acknowledgements: This work was supported by an award to M.D.R. from the National Institute
on Aging [grant number AG039103], and by awards to J.D.K. from the National Institute on Aging
[grant number AG049583] and the Aging Mind Foundation.

Data Availability Statement: The behavioral data are available on the Open Science Framework
(https://osf.io/bpk2d/?view_only=1f70deef4ca64fb38bdc30d9602d8189). Un-threshold statistical
maps from the voxel-wise fMRI contrasts are available on NeuroVault
(https://neurovault.org/collections/PVADJEBH/).

Conflict of Interest Statement: The authors declare no conflicts of interest.

Word Count: 3,495

Number of Tables: 2

Number of Figures: 3

Abstract

The present study investigated the neural correlates of the own-age bias for face recognition in a
repetition suppression paradigm. Healthy young and older adults viewed upright and inverted
unfamiliar faces. Some of the upright faces were repeated following one of two delays (lag 0 or lag
11). Repetition suppression effects were observed in bilateral fusiform cortex. However, there were
no significant effects indicating an own-age bias in repetition suppression. The absence of these
effects is arguably inconsistent with perceptual expertise accounts of own-age biases in face
processing. By contrast, the right anterior hippocampus showed an own-age bias (greater activity for
own- than other-age faces) when viewing an unfamiliar face for the first time. Given the importance
of the hippocampus for episodic memory encoding, we conjecture that the increased hippocampal
activity for own-age relative to other-age faces reflects differential engagement of neural processes
supporting the episodic encoding of faces and might provide insight into the neural underpinnings
of own-age biases in face recognition memory.

Introduction

The ways in which we evaluate, attend to, and remember human faces are susceptible to own-age biases (e.g., Bartlett and Fulton, 1991; Ebner, 2008; Wiese et al., 2008; Rhodes and Anastasi, 2012). Notably, both young and older adults show better recognition memory for own-age faces (Rhodes and Anastasi, 2012). These findings have motivated research that has demonstrated own-age biases in neural correlates of face processing (e.g., Ebner et al., 2011a, 2011b, 2013; Neumann et al., 2015; Wiese et al., 2008; Wolff et al., 2012; Ziaei et al., 2019; for review, see Wiese et al., 2013). Identifying the neuroanatomical correlates of the own-age bias will play an important role in understanding the processes contributing to this phenomenon. There is a growing literature describing different brain regions associated with own-age biases (Ebner et al., 2013, 2011a; Wright et al., 2008; Ziaei et al., 2019; see also Golarai et al., 2017). These studies have reported increased BOLD signal for own-age versus other-age faces in the amygdala, medial prefrontal cortex, orbitofrontal cortex, and insula across a variety of conditions. Although the above studies did not report own-age effects in the fusiform gyrus, a canonical face-processing region, Golarai et al. (2017) recently reported such effects in the fusiform gyrus in both children (7-10 years) and young adults (18-40 years).

The present study had two goals. First, building on prior research, we investigated if own-age biases are present in repetition suppression effects elicited by unfamiliar faces. Repeating a face elicits a 'repetition suppression' effect whereby the BOLD response is reduced for repeated compared to first presentation faces (see Henson, 2016; Henson and Rugg, 2003). Repetition suppression in regions such as the occipital and fusiform face areas are proposed to reflect modulation of processes contributing to the identification of individual faces (Goh et al., 2010; Hermann et al., 2017). This proposal aligns with perceptual expertise theories of the own-age bias arguing that own-age faces are processed more efficiently and are better individuated than other-age faces due to more extensive experience with own-age peers. Second, we aimed to conceptually replicate prior studies examining own-age biases when viewing unfamiliar faces for the first time in the absence of task demands (see Wright et al., 2008).

Materials and Methods

Ethics Statement

This study was approved by Institutional Review Board of the University of Texas at Dallas and University of Texas Southwestern Medical Center. All participants provided written informed consent prior to participation.

Participants

A sample of 24 young and 26 older participants contributed to the analyses reported here. Participants were recruited from the University of Texas at Dallas and the greater Dallas metropolitan area and were financially compensation for their time ($30/hour). The sample sizes were determined by the requirements of a primary experiment in which the aim was to obtain usable data from 24 young and 24 older adults for a memory encoding task (Koen et al., 2019). These were the largest samples that could be accommodated given the resources available for the project. The higher number of participants contributing to the analyses reported here results from the fact some participants were excluded from the primary experiment but retained in the present study.

All participants were right-handed, reported having normal or corrected-to-normal vision
and had no contraindications to MRI scanning. Exclusion criteria included a history of
cardiovascular disease (other than treated hypertension), diabetes, psychiatric disorder, illness, or
trauma affecting the central nervous system, substance abuse, and self-reported current or recent use
of psychotropic medication or sleeping aids. All participants were considered cognitively normal as
determined by performance on a test battery (Table 1; for details of the battery see Koen et al.,
2019). All participants scored 27 or more on the Mini-Mental State Examination (MMSE; Folstein et
al., 1975) and no more than 1.5 standard deviations below age-normalized scores on any one
memory measure or on two (or more) non-memory measures. Data from an additional young adult
male and one older adult male were excluded due to excessive in-scanner motion (> 8 mm
maximum frame-wise displacement) during the task.

Materials and Procedure

The critical stimuli comprised 208 faces from the CAL/PAL database (Ebner, 2008; Minear
and Park, 2004). All face stimuli were 640x480 pixels with a grey background. Half of the face
stimuli depicted younger adults (age range: 18-34 years) and half depicted older adults (age range: 60-
91 years). The 104 stimuli of each face age group were randomly assigned to the lag 0 repeat (24
faces), lag 11 repeat (24 faces), control (32 faces) and inverted (24 faces) conditions. The stimuli
were further split into two lists, with half of the stimuli from each of the face age by trial type
conditions. There were an equal number of male and females faces in each condition. An additional
8 images were used in a practice task completed outside of the scanner. Stimuli were presented to
participants via a mirror mounted to the head coil. Cogent software
(www.vislab.ucl.ac.uk/cogent_2000.php) as implemented in Matlab 2011b (www.mathworks.com)
was used for stimulus control and response logging.

The experimental procedure is depicted in Figure 1. Participants were shown a series of faces
(1 sec duration followed by a 1.25 sec white fixation cross) and were instructed to press a key with
their right index finger whenever an inverted face was presented. Speed was emphasized on the
button press to inverted faces. There were 48 null trials dispersed throughout the block to jitter the
stimuli, with no more than one null trial occurring consecutively. Responses were recorded until the
beginning of the next trial.

Each session of the task comprised 48 null trials and 152 face trials (80 first presentation
trials, 24 lag 0 repeats; 24 lag 11 repeats, 24 inverted faces; equally split across faces depicting young
and older adults). Inverted faces were presented once only, and no more than two inverted trials
occurred in succession. Faces assigned to the repeat condition were presented twice with the
repetition occurring either on the immediately succeeding trial (lag 0) or after 11 intervening trials
(lag 11). Faces in the control condition were presented on only one occasion. Control faces and the
first presentations of repeated faces were collapsed into a single 'first' face condition.

Behavioral Data Analysis

The log-transformed reaction times to inverted face stimuli receiving a correct response were
submitted to a linear mixed model using the *mixed()* function from the *afex* package (Singmann et al.,
2021) in R (R Core Team, 2021). The model included fixed effect terms age group (young versus
older), face age congruency (own age versus other age), and their interaction. Same age faces were
those faces from the same age group as the participant (e.g., young faces were coded as same age
faces for young adults) whereas different age faces were those face stimuli depicting individuals not

in the participants age group (e.g., for young adults, older faces were coded as different age faces).
The model included random effects of participant, specifically a random intercept and random slope
of face age congruency. Degrees of freedom were estimated using the Satterthwaite (1946)
approximation. Bayes factors for the alterative (BF_{10}) and null (BF_{01}) hypotheses were computed
using the *BayesFactor* package (Morey and Rouder, 2018).

MRI Data Acquisition

MRI data were acquired with a 3T Philips Achieva MRI scanner (Philips Medical Systems,
Andover, MA, USA) equipped with a 32-channel receiver head coil. Functional images were
acquired with a blood oxygenation level dependent (BOLD), T2*-weighted echoplanar imaging
(EPI) sequence (SENSE factor = 1.5, flip angle = 70°, 80 × 80 matrix, FOV = 240 mm x 240 mm,
TR = 2000 ms, TE = 30 ms, 34 ascending slices, slice thickness = 3 mm, slice gap = 1 mm), and
were oriented parallel to the AC-PC line. Five "dummy" scans were acquired at the start of each
fMRI session and discarded to allow for equilibration of tissue magnetization. A total of 264
functional volumes were acquired during each of the two task runs, for a total of 528 brain volumes.
T1-weighted images (MPRAGE sequence, 240 × 240 matrix, 1 mm isotropic voxels) were acquired
for anatomical reference. Note that this task was completed prior to the encoding phase of a
memory study that comprised object and scene images (Koen et al., 2019).

fMRI Data Preprocessing and Analysis

The functional data were preprocessed with Statistical Parametric Mapping (SPM12,
Wellcome Department of Cognitive Neurology, London, UK) implemented in Matlab 2017b (The
Mathworks, Inc., USA). The images were reoriented, subjected to a two-pass realignment procedure,
whereby images were initially realigned to the first image of a session and then realigned to the mean
EPI image, and then corrected for slice acquisition time differences using sinc interpolation with
reference to the middle slice. Finally, images were spatially normalized to a study specific EPI
template (de Chastelaine et al., 2011) and smoothed with an 8mm full-width at half-maximum
kernel.

The fMRI data were analyzed with a two-stage random effects model. First level GLMs
modeled neural activity as a delta function convolved with a canonical HRF for the 8 event types
formed by crossing face age and the four item types (first, lag0 repeat, lag11 repeat, inverted face).
The first level GLM also included the 6 realignment parameters and session specific means as
nuisance variables.

The second-level group analyses were performed by submitting the 8 first-level beta maps to
a 2 (age group) by 8 (condition) mixed ANOVA using the factorial ANOVA module in SPM12. The
8 levels of the condition factor were those described above for the first-level models, with one
exception. Like the analysis of the reaction time data, face age was coded as a face age congruency
factor with own-age and other-age faces. Planned contrasts were conducted to identify voxels
showing significant effects of repetition, face age congruency, and to examine whether these effects
interacted with participant age or lag. Additionally, effects of face inversion were examined. Effects
were deemed significant if they survived p < .001 with a cluster-wise correction (p < .05, FWE)
based on Gaussian Random Field theory or if the peak voxel of a cluster survived p < .05 FWE
voxel-level correction. The rationale for using both approaches was to avoid Type-II errors both for
large clusters with (relatively) weak mean activation and for more focal clusters with a strong peak
response.

As noted in the Results, we further probed the whole-brain findings by extracting the mean
beta response from 5mm spheres centered on the peak effects. These values were subjected to
mixed factor ANOVAs using the JASP software (JASP Team, 2020). Bayes factors for effects
favoring the alternative (BF_{10}) and null (BF_{01}) hypotheses are reported for these follow-up analyses.

169   **Results**

Behavioral Data

Participants were accurate in identifying inverted faces (≥ 98%) and very rarely false alarmed
to upright faces (≤ .3%). The analysis of reaction times revealed null effects of age group, $F(1,47.98)$
= 1.93, p = .171, partial-η^2 = .04, BF_{10} = 0.44, BF_{01} = 2.28, face age congruency, $F(1, 47.87)$ = 0.003,
p= .958, partial-η^2 = .00, BF_{10} = 0.05, BF_{01} = 21.53, and the interaction between the two variables,
$F(1, 47.87)$ = 1.36, p= .250, partial-η^2 = .03, BF_{10} = 0.19, BF_{01} = 5.37.

Face Inversion Effects

We first examined the effects of face inversion by contrasting the activity elicited by upright
(first presentation only) and inverted faces. There was an increase in the BOLD signal for inverted
relative to upright races across a wide swath of the cortex with peaks primarily in the frontal and
temporal cortices, as well as the cerebellum. No voxels survived our statistical thresholds when
looking for voxels showing elevated BOLD signal for upright compared to inverted faces. An
additional contrast examining age differences in the face inversion effect identified clusters in the
right pre- and post-central gyri showing smaller face inversion effects for older relative to younger
adults. No significant clusters demonstrated an interaction between face inversion and face age
congruency (own vs. other age faces).

Repetition Suppression

The first planned contrast identified voxels showing effects of face repetition. No clusters
demonstrated significant repetition enhancement effects. However, clusters showing significant
repetition suppression effects were identified in bilateral fusiform gyrus (Table 2 and Figure 2A-B).
The suppression effects did not differ significantly between young and older adults, between
repetition lags, or between own-age and other-age faces according to follow-up analyses that
exclusively masked the repetition suppression contrast with interaction terms involving repetition
and the other factors (exclusive mask threshold set at p < .10, uncorrected). The absence of age
group differences in repetition suppression effects in the fusiform gyrus is consistent with some
prior fMRI findings (Goh et al., 2010). However, the lack of an effect of repetition lag is
inconsistent with previous research on face repetition (Henson et al., 2000; for related findings, see
Nagy and Rugg, 1989); this inconsistency might be due to the relatively short lag in our 'long-lag'
condition (maximum of 11 intervening trials before a face repeat) compared to prior studies.

We next conducted planned contrasts aimed at identifying voxels where face repetition
effects differed in magnitude between young and older adults (i.e., regions showing an Age Group
by Repetition Interaction, collapsed across lag). This contrast identified a cluster in the left interior
temporal gyrus that showed repetition suppression effects in older but not young adults (Figure 2C-
D). Repetition suppression indices (i.e., first minus repeat) extracted from a 5mm sphere centered
on the peak voxel manifesting the interaction revealed a significant repetition suppression effect for
repeated faces in older adults, $t(25)$ = 3.24, p = .003, Cohen's d = 0.635, BF_{10} = 11.93, BF_{01} = 0.08,

but repetition *enhancement* for repeated faces in the younger adults , $t(23) = 2.85$, $p = .009$, Cohen's d
$= 0.582$, $BF_{10} = 5.27$, $BF_{01} = 0.19$.

A further planned contrast found no evidence that face repetition effects interacted with
repetition lag or face-age congruency.

Own-Age biases during novel face viewing

We next conducted a planned contrast between the first presentation trials of upright, own-
age and other-age faces to identify voxels showing an own-age bias. Repeated faces were excluded to
mitigate the potentially confounding effects of familiarity or other repetition-related processes on
own-age bias effects, allowing us essentially to mirror the contrast employed to identify own-race
effects reported by Brown et al. (2017). The above-described contrast identified a cluster of voxels
demonstrating a face age congruency effect in the right anterior hippocampus (Table 2 and Figure
3). The cluster demonstrated elevated BOLD activity for own-age relative to other-age faces. This
effect did not significantly differ in magnitude between young and older adults based on the
outcome of an exclusive mask (at $p < .10$) of the above contrast with the interaction contrast
between age group and face age congruency. Converging with this finding, a conjunction analysis
performed by inclusively masking the own-age versus other-age contrasts conducted separately for
young and older adults (each thresholded at $p < .01$) revealed a cluster that overlapped with that
identified in the initial analyses.

Follow-up analyses investigated whether face repetition moderated the own-age bias effect
observed in the right anterior hippocampus. We conducted a 2 (age group) by 3 (face condition) by
2 (face age congruency) mixed ANOVA on beta estimates for the three upright face trial types (first,
lag 0, and lag 11) from a 5mm sphere centered on the peak of right anterior hippocampus cluster (x
$= 22$, y $= -10$, z $= -24$). Note that because the main effect of face age congruency is not independent
of the contrast used to define the original face age effect, the effect is circular and therefore not
considered here. The ANOVA revealed null effects for the two-way interaction between face age
congruency and age group, $F(1, 47) = 1.55$, $p = .219$, partial-$\eta^2 = .143$, $BF_{10} = 0.308$, $BF_{01} = 3.246$,
for face condition, $F(1.98, 92.93) = 1.26$, $p = .288$, $BF_{10} = 0.163$, $BF_{01} = 6.147$, and for the three-way
interaction, $F(1.98, 92.93) = 1.26$, $p = .288$, $BF_{10} = 0.163$, $BF_{01} = 6.147$. It is noteworthy that the
Bayes factors corresponding to evidence for the null (BF_{01}) provided moderate to strong evidence
favoring null effects according to the definitions proposed by Jeffreys (Jeffreys, 1939; Kass and
Raftery, 1995). In short, these null findings suggest that none of factors included in the ANOVA
significantly moderated the own-age bias effect observed in the right anterior hippocampus.

Discussion

The present study examined whether face age moderated repetition suppression effects
elicited by unfamiliar faces and whether there were any own-age biases in neural activity when
viewing first presentations of unfamiliar faces. There were three main findings. First, although we
identified significant age-invariant repetition suppression effects in bilateral fusiform gyrus, we did
not identify any clusters where suppression effects demonstrated an own-age bias. These null
findings are arguably inconsistent with perceptual expertise accounts of the own-age bias (Tanaka
and Pierce, 2009; Valentine, 1991; Wiese et al., 2013). Such accounts propose that own-age faces are
more efficiently processed than other-age faces due to more extensive experience with own-age
peers. If face repetition suppression effects reflect modulation of processes contributing to the
identification of individual faces (Hermann et al., 2017), then we might expect that own-age faces

would elicit a greater suppression effect than other-age faces, which we did not observe. However,
one limitation of the present design is that we only examined repetition effects for identical repeats
and did not include a condition that could examine 'release from suppression' (e.g., Goh et al., 2010;
Reggev et al., 2020). Recently, Reggev et al. (2020) argued in favor of a perceptual expertise account
of own-race biases based on findings of greater release from suppression for own-race relative to
other-race faces. Future research employing designs that allow measurement of release from
suppression is needed to pursue this issue.

Second, we identified a cluster in left inferior temporal cortex that demonstrated repetition
suppression effects for older adults but repetition enhancement effects in older adults. This region
has been previously reported to demonstrate repetition suppression effects for familiar (e.g., yellow
banana) but not novel concepts (e.g., purple banana) (Reggev et al., 2016). Given that familiar
concepts are accrued through experience, it is possible that the present interaction reflects age group
differences in exposure to faces (Golarai et al., 2017; for a related discussion, see Koen and Rugg,
2019). By this argument, the left inferior temporal gyrus is a region that is especially sensitive to
cumulative lifetime experience with a perceptual category such as faces. Future research will be
needed to test the validity of this proposal.

Lastly, we observed an own-age bias for the first presentation of unfamiliar faces in the right
anterior hippocampus. Both young and older adults showed elevated BOLD signal for own-age
relative to other-age faces. We stress that this finding of an own-age bias in the hippocampus
requires replication within an experimental task that allows for a behavioral assay of own-age bias,
such as recognition memory. Nonetheless, given the well-established role of the hippocampus in
memory encoding (Eichenbaum et al., 2007; Kim, 2011), we conjecture that the present finding is
relevant to the own-age bias that has been reported for recognition memory performance (Rhodes
and Anastasi, 2012). There are several possible accounts of this finding, which are not mutually
exclusive. One possibility is that it is easier to bind previously acquired personal or semantic
knowledge about familiar individuals to unfamiliar own-age than to other-age faces. Alternately,
own-age faces may more readily attract attention to discriminating facial features than other-age
faces, in turn modulating encoding-related hippocampal activity (Aly and Turk-Browne, 2015, 2017;
Uncapher and Rugg, 2009). Another possibility is that the present hippocampal effect reflects the
formation of face representations with higher fidelity than representations of other-age faces (for
reviews, see Ekstrom and Yonelinas, 2020; Yonelinas, 2013). Future studies that link the own-age
bias in recognition memory to hippocampal effects promise to shed light on these and other
possibilities.

There are limitations to this study. First, we did not replicate prior findings of own-age
biases in regions such as the prefrontal cortex, insula, amygdala, and fusiform gyrus (Ebner et al.,
2013, 2011a; Golarai et al., 2017; Wright et al., 2008; Ziaei et al., 2019). The reasons for this
replication failure are unclear. One possibility is that task demands (e.g., passive viewing versus
emotion recognition) contributed to the differences between the present and prior findings. Second,
the own-age bias observed in the hippocampus did not have a behavioral correlate. While this is a
potential limitation, we note that prior studies have reported own-age biases in neural data in the
absence of analogous findings in behavioral data (e.g., Ebner et al., 2013; Ziaei et al., 2019).

In conclusion, we did not find evidence for own-age biases in face repetition suppression
effects, which is arguably inconsistent with perceptual expertise accounts of the own-age bias. Of
importance, a cluster in the right anterior hippocampus showed an own-age bias to the first

presentation of unfamiliar faces in both young and older adults, which we speculate to be related to
own-age biases in memory.
295

References

Aly M, Turk-Browne NB. 2017. How Hippocampal Memory Shapes, and Is Shaped by, Attention
In: Hannula DE, Duff MC, editors. The Hippocampus from Cells to Systems. Cham: Springer
International Publishing. pp. 369–403. doi:10.1007/978-3-319-50406-3_12

Aly M, Turk-Browne NB. 2015. Attention Stabilizes Representations in the Human Hippocampus.
Cerebral Cortex bhv041. doi:10.1093/cercor/bhv041

Bartlett JC, Fulton A. 1991. Familiarity and recognition of faces in old age. *Memory & Cognition*
19:229–238. doi:10.3758/BF03211147

Brown TI, Uncapher MR, Chow TE, Eberhardt JL, Wagner AD. 2017. Cognitive control, attention,
and the other race effect in memory. *PLOS ONE* **12**:e0173579.
doi:10.1371/journal.pone.0173579

de Chastelaine M, Wang TH, Minton B, Muftuler LT, Rugg MD. 2011. The Effects of Age, Memory
Performance, and Callosal Integrity on the Neural Correlates of Successful Associative
Encoding. *Cerebral Cortex* **21**:2166–2176. doi:10.1093/cercor/bhq294

Ebner NC. 2008. Age of face matters: Age-group differences in ratings of young and old faces. *Behav*
Res **40**:130–136. doi:10.3758/brm.40.1.130

Ebner NC, Gluth S, Johnson MR, Raye CL, Mitchell KJ, Johnson MK. 2011a. Medial prefrontal
cortex activity when thinking about others depends on their age. *Neurocase* **17**:260–269.
doi:10.1080/13554794.2010.536953

Ebner NC, He Y, Fichtenholtz HM, McCarthy G, Johnson MK. 2011b. Electrophysiological
correlates of processing faces of younger and older individuals. *Soc Cogn Affect Neurosci* **6**:526–
535. doi:10.1093/scan/nsq074

Ebner NC, Johnson MR, Rieckmann A, Durbin KA, Johnson MK, Fischer H. 2013. Processing
own-age vs. other-age faces: Neuro-behavioral correlates and effects of emotion. *NeuroImage*
78:363–371. doi:10/f2zvnz

Eichenbaum H, Yonelinas AP, Ranganath C. 2007. The Medial Temporal Lobe and Recognition
Memory. *Annual Review of Neuroscience* **30**:123–152.
doi:10.1146/annurev.neuro.30.051606.094328

Ekstrom AD, Yonelinas AP. 2020. Precision, binding, and the hippocampus: Precisely what are we
talking about? *Neuropsychologia* **138**:107341. doi:10.1016/j.neuropsychologia.2020.107341

Goh JO, Suzuki A, Park DC. 2010. Reduced neural selectivity increases fMRI adaptation with age
during face discrimination. *NeuroImage* **51**:336–344. doi:10.1016/j.neuroimage.2010.01.107

Golarai G, Liberman A, Grill-Spector K. 2017. Experience Shapes the Development of Neural
Substrates of Face Processing in Human Ventral Temporal Cortex. *Cerebral Cortex* **27**.
doi:10/gmr7x9

Henson R, Shallice T, Dolan R. 2000. Neuroimaging Evidence for Dissociable Forms of Repetition
Priming. *Science* **287**:1269–1272. doi:10.1126/science.287.5456.1269

Henson RN. 2016. Repetition suppression to faces in the fusiform face area: A personal and
dynamic journey. *Cortex*, Special Issue:Repetition suppression-an integrative view **80**:174–184.
doi:10.1016/j.cortex.2015.09.012

Henson RNA, Rugg MD. 2003. Neural response suppression, haemodynamic repetition effects, and
behavioural priming. *Neuropsychologia* **41**:263–270. doi:10.1016/s0028-3932(02)00159-8

Hermann P, Grotheer M, Kovács G, Vidnyánszky Z. 2017. The relationship between repetition
suppression and face perception. *Brain Imaging and Behavior* **11**:1018–1028. doi:10.1007/s11682-
016-9575-9

JASP Team. 2020. JASP (Version 0.14.1) [Computer software].

Jeffreys J. 1939. Theory of probability, 1st ed. Oxford, UK: Oxford University Press.

Kass RE, Raftery AE. 1995. Bayes Factors. *Journal of the American Statistical Association* **90**:773–795.
doi:10/gdnbw3

Kim H. 2011. Neural activity that predicts subsequent memory and forgetting: A meta-analysis of 74
fMRI studies. *NeuroImage* **54**:2446–2461. doi:10.1016/j.neuroimage.2010.09.045

Koen JD, Hauck N, Rugg MD. 2019. The relationship between age, neural differentiation, and
memory performance. *Journal of Neuroscience* **39**:149–162.
doi:https://doi.org/10.1523/JNEUROSCI.1498-18.2018

Koen JD, Rugg MD. 2019. Neural dedifferentiation in the aging brain. *Trends in Cognitive Sciences*
23:547–559. doi:10.1016/j.tics.2019.04.012

Minear M, Park DC. 2004. A lifespan database of adult facial stimuli. *Behavior Research Methods,*
Instruments, & Computers **36**:630–633. doi:10.3758/bf03206543

Morey RD, Rouder JN. 2018. BayesFactor: Computation of bayes factors for common designs
(manual).

Nagy ME, Rugg MD. 1989. Modulation of Event-Related Potentials by Word Repetition: The
Effects of Inter-Item Lag. *Psychophysiology* **26**:431–436. doi:https://doi.org/10.1111/j.1469-
8986.1989.tb01946.x

Neumann MF, End A, Luttmann S, Schweinberger SR, Wiese H. 2015. The own-age bias in face
memory is unrelated to differences in attention—Evidence from event-related potentials.
Cognitive, Affective, & Behavioral Neuroscience **15**:180–194. doi:10.3758/s13415-014-0306-7

R Core Team. 2021. R: A language and environment for statistical computing (manual). Vienna,
Austria.

Reggev N, Bein O, Maril A. 2016. Distinct Neural Suppression and Encoding Effects for
Conceptual Novelty and Familiarity. *Journal of Cognitive Neuroscience* **28**:1455–1470.
doi:10.1162/jocn_a_00994

Reggev N, Brodie K, Cikara M, Mitchell JP. 2020. Human face-selective cortex does not distinguish
between members of a racial outgroup. *eNeuro*. doi:10.1523/ENEURO.0431-19.2020

Rhodes MG, Anastasi JS. 2012. The own-age bias in face recognition: A meta-analytic and
theoretical review. *Psychological Bulletin* **138**:146–174. doi:10.1037/a0025750

Satterthwaite FE. 1946. An Approximate Distribution of Estimates of Variance Components.
Biometrics Bulletin **2**:110. doi:10.2307/3002019

Singmann H, Bolker B, Westfall J, Aust F, Ben-Shachar MS. 2021. afex: Analysis of factorial
experiments (manual).

Tanaka JW, Pierce LJ. 2009. The neural plasticity of other-race face recognition. *Cognitive, Affective, &*
Behavioral Neuroscience **9**:122–131. doi:10/dwn4dz

Uncapher MR, Rugg MD. 2009. Selecting for Memory? The Influence of Selective Attention on the
Mnemonic Binding of Contextual Information. *Journal of Neuroscience* **29**:8270–8279.
doi:10.1523/jneurosci.1043-09.2009

Valentine T. 1991. A Unified Account of the Effects of Distinctiveness, Inversion, and Race in Face
Recognition. *The Quarterly Journal of Experimental Psychology Section A* **43**:161–204. doi:10/c3mcjt

Wiese H, Komes J, Schweinberger SR. 2013. Ageing faces in ageing minds: A review on the own-age
bias in face recognition. *Visual Cognition* **21**:1337–1363. doi:10.1080/13506285.2013.823139

Wiese H, Schweinberger SR, Hansen K. 2008. The age of the beholder: ERP evidence of an own-
age bias in face memory. *Neuropsychologia* **46**:2973–2985.
doi:10.1016/j.neuropsychologia.2008.06.007

Wolff N, Wiese H, Schweinberger SR. 2012. Face recognition memory across the adult life span:
Event-related potential evidence from the own-age bias. *Psychology and Aging* **27**:1066–1081.
doi:10.1037/a0029112

Wright CI, Negreira A, Gold AL, Britton JC, Williams D, Feldman Barrett L. 2008. Neural
correlates of novelty and face–age effects in young and elderly adults. *NeuroImage* **42**:956–968.
doi:10.1016/j.neuroimage.2008.05.015

Yonelinas AP. 2013. The hippocampus supports high-resolution binding in the service of
perception, working memory and long-term memory. *Behavioural Brain Research* **254**:34–44.
doi:10.1016/j.bbr.2013.05.030

Ziaei M, Persson J, Bonyadi MR, Reutens DC, Ebner NC. 2019. Amygdala functional network
during recognition of own-age vs. other-age faces in younger and older adults. *Neuropsychologia*
129:10–20. doi:10/gh8qgd

Table 1. Demographic and neuropsychological test data for young and older adults.

	Young Adults	Older Adults	p-value
N	24	26	
Age	23.04 (3.46)	69.19 (3.27)	
Sex (M/F)	12/12	14/12	
Education	15.92 (2.22)	16.96 (2.24)	.104
MMSE	29.54 (0.59)	29.35 (0.94)	.378
CVLT Short Delay – Free	13.08 (1.79)	10.38 (3.16)	< .001
CVLT Short Delay – Cued	13.67 (1.81)	11.67 (3.57)	.016
CVLT Long Delay – Free	13.54 (2.06)	10.23 (3.27)	< .001
CVLT Long Delay – Cued	14.12 (1.62)	11.92 (2.78)	.001
CVLT Recognition – Hits	15.42 (0.83)	15.00 (0.98)	.110
CVLT Recognition – False Alarms	0.46 (0.66)	3.08 (2.53)	< .001
Logical Memory I	30.62 (4.95)	26.00 (5.50)	.003
Logical Memory II	28.12 (5.78)	22.62 (6.03)	.002
Digit Span Total[1]	21.04 (4.53)	17.77 (2.55)	.004
SDMT	65.38 (13.99)	46.42 (7.74)	< .001
Trails A (secs)	21.43 (7.97)	31.55 (10.76)	< .001
Trails B (secs)	47.54 (19.53)	70.52 (24.87)	.002
F-A-S Total	48.29 (10.97)	45.50 (11.42)	.382
Category Fluency (Animals)	24.58 (5.67)	21.00 (4.66)	.020
WTAR (Raw)	41.42 (3.44)	43.08 (4.78)	.163
Raven's (List 1)	11.08 (.97)	9.19 (2.50)	.001
Visual Acuity (logMar)[2]	-.11 (.10)	.06 (.11)	< .001

Note. Standard deviations are reported in parentheses. The *p*-values were obtained from unequal-variance independent *t*-tests comparing young and older adults. [1]Digit span total equals the sum of forward and backward span. [2]Lower logMAR scores indicate better visual acuity. MMSE = Mini-mental State Exam; CVLT = California Verbal Learning Test II; SDMT = Symbol-Digit Modalities Test; WTAR = Wechsler Test of Adult Reading

407 **Table 2.** Regions showing effects of face repetition and age congruency.

Region	MNI Coordinates			Cluster size	Peak z
	x	y	z		
Inverted > Novel (p < 1e^{-10}, FWE-corrected height threshold)					
Cerebellum	24	-55	-28	14458	18.75
	-24	-58	-28		16.49
	-33	-55	-34		15.78
R. Insula/Inferior Frontal Gyrus	42	14	-7	914	13.87
	36	20	-1		13.76
	57	11	8		12.61
L. Insula/Inferior Frontal Gyrus	-42	2	5	704	12.80
	-60	8	29		12.39
	-33	20	2		11.67
R. Superior Frontal Gyrus	33	47	17	228	11.16
L. Superior Frontal Gyrus	-33	47	20	151	10.58
R. Middle Temporal Gyrus	51	-25	-13	1	8.34
Age Group by Face Inversion Interaction (Older > Young Adults)					
R. Postcentral Gyrus[a,b]	60	-7	32	318	4.77
L. Precentral Gyrus[a,b]	-54	-7	41	136	4.24
Face Repetition Suppression (First > Repeat)					
R. Fusiform Gyrus[a,b]	39	-43	-28	493	5.82
L. Fusiform Gyrus[a]	-39	-46	-28	328	4.20
Age Group by Repetition Interaction (Older: Repetition Suppression, Young: Repetition Enhancement)					
L. Inferior Temporal Cortex[a,b]	-51	-43	-19	128	5.22
Face Age Congruency Effects (Own-Age > Other-Age Faces)					
R. Hippocampus[b]	21	-10	-22	71	4.99

Note. The peaks for the Inverted > Upright contrast were obtained using a height threshold of $p <$
$1e^{-10}$ (FWE-corrected) to better delineate and characterize the peaks in the data. Regions showing
repetition suppression effects were identified from a planned contrast of first and repeated face trials
(i.e., a main effect of face repetition) without regard for other conditions (i.e., repetition lag, Age
Group, and Face Age Congruency). The left interior temporal cortex was identified by a contrast
modeling the Age Group by Repetition (first vs. repeated) interaction regardless of Lag and Face
Age Congruency. Face Age Congruency effects were identified from an own-age versus other-age
contrast for first presentation, upright face trials. [a]Significant at $p < .001$ height threshold with $p <$
$.05$, FWE-corrected cluster threshold). [b]Significant at the voxel-level corrected threshold ($p < .05$,
FWE-corrected height threshold).

Figure 1. Schematic of the face repetition task. Participants saw a series of young and older adult faces in a face inversion task. Participants were instructed to make a button press if the face was inverted, and otherwise to simply view the faces. Each face was shown for 1 second and followed by an inter-trial interval (fixation cross) for 1.25 seconds. There was a total of 8 conditions formed by crossing Face Age (young, older) and Trial Type (first, lag 0 repeat, lag 11 repeat, inverted). Additional details are reported in the Materials and Methods section.

Figure 2. Age-group invariant (top panels) and age-group dependent (bottom panels) face repetition suppression effects. (A) A cluster in the right fusiform gyrus (x = 39, y = -43, z = -28) showing suppression of the BOLD signal for repeated faces compared to the first presentation of faces in both young and older adults. A similar pattern was observed in the left fusiform gyrus (not shown, see Table 1). (B) Repetition suppression estimates (first minus repeated presentation) extracted from a 5mm sphere centered on the right fusiform gyrus peak voxel. (C) A cluster in the left inferior temporal cortex (x = 39, y = -43, z = -28) showing larger repetition suppression effects for older relative to younger adults. (D) Repetition suppression estimates extracted from a 5mm sphere centered on the left inferior temporal cortex peak voxel. The interaction in this region is driven by the combination of repetition suppression in older adults and repetition enhancement in younger adults. (A) and (C) are shown at p < .001, uncorrected, for visualization purposes, and depicted in neurological orientation (right is right). In (B) and (D), the solid black circles represent condition means, and the green and orange points depict data from individual participants. Error bars reflect the 95% confidence interval.

Figure 3. BOLD signals in the right hippocampus were greater when young and older adults viewed
faces belonging to their own age group relative to other age faces. (A) Viewing the first repetition of
novel faces was associated with an own-age bias in the right anterior hippocampus (x = 22, y = -10,
z = -24). The image is shown at a threshold of at p < .001 uncorrected, for visualization purposes.
(B) The beta values extracted from a 5mm sphere centered on the peak coordinate in the right
hippocampus. The solid black circles represent condition means, and the green and orange points
depict data from individual participants. Error bars reflect the 95% confidence interval.