1

A glycan gate controls opening of the SARS-CoV-2 spike protein

Terra Sztain[1†], Surl-Hee Ahn[1†], Anthony T. Bogetti[2], Lorenzo Casalino[1], Jory A. Goldsmith[3],

Ryan S. McCool[3], Fiona L. Kearns[1], J. Andrew McCammon[1], Jason S. McLellan[3], Lillian T.

Chong[2*], Rommie E. Amaro[1*]

1. Department of Chemistry and Biochemistry, UC San Diego, La Jolla, CA 92093

2. Department of Chemistry, University of Pittsburgh, Pittsburgh, PA 15260

3. Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX 78712

† These authors contributed equally to this work.

* contact authors: ramaro@ucsd.edu, ltchong@pitt.edu

Abstract

SARS-CoV-2 infection is controlled by the opening of the spike protein receptor binding domain

(RBD), which transitions from a glycan-shielded (down) to an exposed (up) state in order to bind

the human ACE2 receptor and infect cells. While snapshots of the up and down states have been

obtained by cryoEM and cryoET, details of the RBD opening transition evade experimental

characterization. Here, over 200 µs of weighted ensemble (WE) simulations of the fully

glycosylated spike ectodomain allow us to characterize more than 300 continuous, kinetically

unbiased RBD opening pathways. Together with biolayer interferometry experiments, we reveal

a gating role for the N-glycan at position N343, which facilitates RBD opening. Residues D405,

R408, and D427 also participate. The atomic-level characterization of the glycosylated spike

activation mechanism provided herein achieves a new high-water mark for ensemble pathway

simulations and offers a foundation for understanding the fundamental mechanisms of SARS-

CoV-2 viral entry and infection.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is an enveloped RNA virus and

the causative agent of COVID-19, a disease that has caused significant morbidity and mortality

worldwide.[1,2] The main infection machinery of the virus is the spike (S) protein, which sits on

the outside of the virus, is the first point of contact that the virion makes with the host cell, and is

a major viral antigen.[3] A significant number of cryoEM structures of the spike protein have been

recently reported, collectively informing on structural states of the spike protein. The vast

majority of resolved structures fall into either 'down' or 'up' states, as defined by the position of

the receptor binding domain (RBD), which modulates interaction with the ACE2 receptor for cell

entry.[4,5,6]

The RBDs must transition from a down to an up state for the receptor binding motif to be

accessible for ACE2 binding (**Figure 1**), and therefore the activation mechanism is essential for

cell entry. Mothes et al.[7] used single-molecule fluorescence (Förster) resonance energy transfer

(smFRET) imaging to characterize spike dynamics in real-time. Their work showed that the

spike dynamically visits four distinct conformational states, the population of which are

modulated by the presence of the human ACE2 receptor and antibodies. However smFRET, as

well as conventional structural biology techniques, are unable to inform on the atomic-level

mechanisms underpinning such dynamical transitions. Recently, all-atom molecular dynamics

(MD) simulations of the spike protein with experimentally accurate glycosylation together with

corroborating experiments indicated the extensive shielding by spike glycans, as well as a

mechanical role for glycans at positions N165 and N234 in supporting the RBD in the open

conformation.[8] Conventional MD simulations as performed in Casalino et al.[8] also revealed

microsecond-timescale dynamics to better characterize the spike dynamics but were limited to

sampling configurations that were similar in energy to the cryoEM structures. Several enhanced

sampling MD simulations have been performed to study this pathway, however, these

simulations lacked glycosylation for the spike protein[9] or involve the addition of an external

force[10] or require a massive amount of sampling (millisecond timescale).[11]

56
57 **Figure 1** (A) The SARS-CoV-2 spike head (gray) with glycans (dark blue) as simulated, with the stalk
58 domain and membrane (not simulated here, but shown in transparent for completeness). RBD shown in
59 cyan, receptor binding motif (RBM) in pink. Side view of the down (shielded, B) and open (exposed, C)
60 RBD. Top view of the closed (shielded, D) and open (exposed, E) RBM. Composite image of glycans
61 (dark blue lines) shows many overlapping snapshots of the glycans over the microsecond simulations.

62
In this study, we characterized the spike RBD opening pathway for the fully glycosylated SARS-

CoV-2 spike protein, in order to gain a detailed understanding of the activation mechanism. We

used the weighted ensemble (WE) path sampling strategy[12,13] (**Figure S1**) to enable the

simulation of atomistic pathways for the spike opening process. As a path sampling strategy, WE

focuses computing power on the functional transitions between stable states rather than the stable

states themselves.[14] This is achieved by running multiple trajectories in parallel and periodically

replicating trajectories that have transitioned from previously visited to newly visited regions of

configurational space[15], thus minimizing the time spent waiting in the initial stable state for

"lucky" transitions over the free energy barrier. Given that these transitions are much faster than

the waiting times,[16,17] the WE strategy can be orders of magnitude more efficient than

conventional MD simulations in generating pathways for rare events such as protein folding and

protein binding.[18,19] This efficiency is even higher for slower processes, increasing *exponentially*

with the effective free energy barrier.[20] Not only are dynamics carried out without any biasing

force or modifications to the free energy landscape, but suitable assignment of statistical weights

to trajectories provides an *unbiased* characterization of the system's time-dependent ensemble

properties.[13] The WE strategy therefore generates continuous pathways with unbiased dynamics,

yielding the most direct, atomistic views for analyzing the mechanism of functional transitions,

including elucidation of transient states that are too fleeting to be captured by laboratory

experiments. Furthermore, while the strategy requires a progress coordinate toward the target

state, the definition of this target state need not be fixed in advance when applied under

equilibrium conditions,[21] enabling us to refine the definition of the target open state of the spike

protein based on the probability distribution of protein conformations sampled by the simulation.

Our work characterizes a series of transition pathways of the spike opening and identifies key

residues, including a glycan at position N343, that participate in the opening mechanism. Our

simulation findings are corroborated by biolayer interferometry experiments, which demonstrate

a reduction in the ability of the spike to interact with ACE2 after mutation of these key residues.

Results and Discussion

Weighted ensemble simulations of spike opening

We used the WE strategy to generate continuous, atomistic pathways of the spike opening

process with unbiased dynamics (**Figure 2A-E, Movie S1**); these pathways were hundreds of ns

long, excluding the waiting times in the initial "down" state. The protein model was based on the

head region (residues 16 to 1140) of the glycosylated SARS-CoV-2 spike from Casalino et al.[8]

(**Figure 1**), which in turn was built on the cryoEM structure of the three-RBD-down spike (PDB

ID: 6VXX[5]). The entire simulation system, including explicit water and salt ions, reaches almost

half a million atoms. We focused sampling along a two-dimensional progress coordinate to track

RBD opening: the difference in the center of mass of the spike core to the RBD and the root

mean squared deviation (RMSD) of the RBD (**Figure 2F-G**). On the SDSC Comet and TACC

Longhorn supercomputers, 100 GPUs ran the WE simulations in parallel for over a month,

generating over 200 μs of glycosylated spike trajectories and more than 200 TB of trajectory

data. We simulated a total of 310 independent pathways, including 204 pathways from the RBD

"down" conformation (PDB ID: 6VXX[5]) to the RBD "up" conformation (PDB ID: 6VSB[4]) and

106 pathways from the RBD "down" to RBD "open" state in which the RBD twists open beyond

the cryoEM structure (PDB ID: 6VSB[4]). Remarkably, the open state that we sampled includes

conformations that align closely with the ACE2-bound spike cryoEM structure (PDB ID: 7A95[6])

even though this structure was not a target state of our progress coordinate (**Figure 2F-G**, **Movie**

S1, Figures S2-3). This result underscores the value of using (i) equilibrium WE simulations that

do not require a fixed definition of the target state and (ii) a two-dimensional progress coordinate

that allows the simulations to sample unexpected conformational space along multiple degrees of

freedom. The ACE2-bound spike conformation has also been sampled by the Folding@home

distributed computing project,[11] and RBD rotation has been detected in cryoEM experiments.[6]

The N343 glycan gates RBD opening

In the down state, the RBD of the SARS-CoV-2 spike is shielded by glycans at positions N165,

N234, and N343.[22] Beyond shielding, a structural role for glycans at positions N165 and N234

has been recently reported, stabilizing the RBD in the up conformation.[8]

Our WE simulations reveal an even more specific, critical role of a glycan in the opening

mechanism of the spike: the N343 glycan acts as a "glycan gate" pushing up the RBD by

intercalating between residues F490, Y489, F456, and R457 of the ACE2 binding motif in a

"hand-jive" motion (**Figure 2A-E**, **Figure 3**, **Movie S2**). This gating mechanism was initially

visualized in several successful pathways of spike opening and then confirmed through analysis

of all 310 successful pathways in which the N343 glycan was found to form contacts (within 3.5

Å) with each of the aforementioned residues in every successful pathway (**Figure S4**). The same

mechanistic behavior of the N343 glycan was observed in two fully independent WE

simulations, suggesting the result is robust despite potentially incomplete sampling that can

challenge WE and other enhanced sampling simulation methods.[15]

To test the role of the N343 glycan as a key gating residue, we performed biolayer interferometry

(BLI) experiments. BLI experiments assess the binding level of the spike receptor binding motif

(RBM, residues 438 to 508) to ACE2, acting as a proxy for the relative proportion of RBDs in

the up position for each spike variant. No residues directly involved in the binding were mutated

(i.e., at the RBM-ACE2 interface), to ensure controlled detection of the impact of RBD opening

in response to mutations. Although previous results have shown reduced binding levels for

N165A and N234A variants in the SARS-CoV-2 S-2P protein,[8] the N343A variant displayed an

even greater decrease in ACE2 binding, reducing spike binding level by ~56 % (**Figure 4, Table**

S1). As a negative control, the S383C/D985C variant,[23] which is expected to be locked by

disulfides into the three-RBD-down conformation, showed no association with the ACE2

receptor. These results support the hypothesis that the RBD up conformation is significantly

affected by glycosylation at position N343.

147
148
Figure 2 Atomically detailed pathways of spike opening. (A-E) Snapshot configurations along the opening pathway
with chain A shown in cyan, chain B in gray, and chain C in pink and the glycan at position N343 is shown in
magenta. (F) Scatter plot of data from the 310 continuous pathways with the RMSD of RBD to the 6VSB "up" state
plotted against RBD — core distance. Data points are colored based on % RBD solvent accessible surface area
compared to the RBD "down" state 6VXX. Location of snapshots shown in A-E are labeled. (G) Primary regions of
spike defined for tracking progress of opening transition. The spike core is composed of three central helices per
trimer, colored according to chain as in (A-E). The RBD contains a structured pair of antiparallel beta sheets and an
overlay of snapshots from a continuous WE simulation are shown colored along a spectrum resembling the palette in
(F). Overlayed cryoEM structures are highlighted and labeled including the initial RBD "down" state, 6VXX, the
target RBD "up" state and the ACE2 bound "open" state, 7A95.

159
160 **Figure 3** Glycan gating by N343. (A-D) Snapshot configurations along the opening pathway with chain A shown in
161 cyan, chain B in gray, chain C in pink, and the glycan at position N343 is shown in magenta. (A) RBD A in the
162 down conformation is shielded by the glycan at position N343 of the adjacent RBD B. (B-D) The N343 glycan
163 intercalates between and underneath the residues F490, Y489, F456, F457 to push the RBD up and open (D).

Figure 4 ACE2 binding is reduced by mutation of N343 glycosylation site and key salt bridge residues. (A) Biolayer interferometry sensorgrams of HexaPro spike variants binding to ACE2. For clarity, only the traces from the first replicate are shown. (B) Graph of binding response for BLI data collected in triplicate with error bars representing the standard deviation from the mean.

Atomic details of the opening mechanism

The RBD down state features a hydrogen bond between T415 of the RBD (chain A) and K986 of

chain C, a salt bridge between R457 of RBD$_A$ and D364 of RBD$_B$, and a salt bridge between

K462 of RBD$_A$ and D198 of NTD$_C$ (**Figure 5A-C,E Figure S5**). The hydrogen bond T415$_A$ -

K986$_C$ spends an average of 12% of the successful pathways to the up state before K986$_C$ makes

a short lived (2% average duration to up state) salt bridge with RBD$_A$ D427. (**Figure 5B,E**

Figure S3) Next, K986$_C$ forms salt bridges with E990$_C$ and E748$_C$ as the RBD$_A$ continues to

open. These contacts are formed in all 310 successful pathways (**Figure S5**). Mutation of K986

to proline has been used to stabilize the pre-fusion spike,[24,25] including in vaccine development,[26]

and these simulations provide molecular context to an additional role of this residue in RBD

opening.

Subsequently, at an average of 16% of the way through the successful pathways to the up state,

the $R457_A$ - $D364_B$ salt bridge is broken, prompting the RBD_A to twist upward, away from RBD_B

towards RBD_C and forms a salt bridge between R408 of RBD_A and D405 of RBD_C (**Figure 5C**

and 5E, Figure S5). This salt bridge persists for 20% of the successful trajectories to the up

state, and is also present in all 310 successful pathways.

A salt bridge between R466 of RBD_A and E132 from NTD_B is present in 189 out of 204

successful pathways to the up state, and all 106 pathways to the open state. This contact is most

prevalent during the transition between the up and open state. Finally, the salt bridge between

D428 of RBD_A and R454 of RBD_C is present only in all 106 pathways from the up to the open

state and is the last salt bridge between the RBD and the spike in the open state before the S1

subunit begins to peel off (**Figure 5D and 5E, Figure S5**), at which point the last remaining

contact to the RBD_A is the glycan at position N165 of NTB_B.

Additional BLI experiments of the key identified spike residues R408A, D405A, and D427A

corroborate the pathways observed in our simulations. Each of these reduces the binding

interactions of the spike with ACE2 by ~13%, ~27%, and ~52%, respectively (**Figure 4, Table**

S1). We also note that identified residues D198, N343, D364, D405, R408, T415, D427, D428,

R454, R457, R466, E748, K986 and E990 are conserved between SARS-CoV and SARS-CoV-2

spikes, supporting their significance in coordinating the primary spike function of RBD opening.

The emerging mutant SARS-CoV-2 strains B.1: D614G; B.1.1.7: H69-V70 deletion, Y144-Y145

deletions, N501Y, A570D, D614G, P681H, T716I, S982A, D1118H; B.1.351: L18F, D80A,

D215G, R246I, K417N, E484K, N501Y, D614G, A701V; P1: L18F, T20N, P26S, D138Y,

R190S, K417T, E484K, N501Y ,D614G, H655Y, T1027I and CAL.20C: L452R, D614G[27] do

not contain mutants in the residues we identified here to facilitate RBD opening. Analysis of

neighboring residues and glycans to those mutated in the emerging strains along the opening

pathway is detailed in **Table S2**, and distances between each residue and glycan to RBD$_A$ is

summarized in **Movie S3.**

212
213 **Figure 5** Salt bridges and hydrogen bonds along the opening pathway. (A-D) salt bridge or hydrogen bond contacts
214 made between RBD A, shown in blue, with RBD B, shown in gray, or RBD C shown in pink within the down,
215 transient, up, and open conformations, respectively. (E) Histogram showing the frequency at which residues from
216 (A-D) are within 3.5 Å of each other relative to RBD — core distance. Frequencies are normalized to 1.
217
218

Conclusions

We report extensive weighted ensemble molecular dynamics simulations of the glycosylated

SARS-CoV-2 spike head characterizing the transition from the down to up conformation of the

RBD. Over 200 microseconds of simulation provide more than 300 independent RBD opening

transition pathways. Analysis of these pathways from independent WE simulations indicates a

clear gating role for the glycan at N343, which lifts and stabilizes the RBD throughout the

opening transition. We also characterize an "open" state of the spike RBD, in which the N165

glycan of chain B is the last remaining contact with the RBD *en route* to further opening of S1.

Biolayer interferometry experiments of residues identified as key in the opening transitions,

including N343, D405, R408, and D427, broadly supported our computational findings. Notably,

a 56% decrease in ACE2 binding of the N343A mutant, compared to a 40% decrease in N234A

mutant, and 10% decrease in the N165A mutant reported previously[8] evidenced the key role of

N343 in gating and assisting the RBD opening process, highlighting the importance of sampling

functional transitions to fully understand mechanistic detail. Our work indicates a critical gating

role of the N343 glycan in spike opening and provides new insights to mechanisms of viral

infection for this important pathogen.

Data accession

238 We endorse the community principles around open sharing of COVID19 simulation data.[28] All

239 simulation input files and data are available at the NSF MolSSI COVID19 Molecular Structure

240 and Therapeutics Hub at https://covid19.molssi.org and the Amaro Lab website

241 http://amarolab.ucsd.edu.

Author contributions

T.S. and S.A. contributed equally to this work. R.E.A. and L.T.C. oversaw the project. T.S. and L.C. prepared the simulation model. T.S. and S.A. performed WE simulations and A.B. provided WESTPA scripts. S.A., A.B., T.S., and L.T.C. carried out WE analysis. T.S., and F.L.K. performed simulation analyses. L.C., T.S., and F.L.K. created figures and movies. J.S.M. designed and oversaw biolayer interferometry experiments. R.M. and J.A.G. performed biolayer interferometry experiments and wrote the corresponding parts in the Results and Methods sections. T.S., S.A., L.T.C., and R.E.A. wrote the manuscript with contributions from all authors.

Funding

T.S. is funded by NSF GRFP DGE-1650112. S.A. is funded by NIH grant R01-GM31749. This work was supported by NIH GM132826, NSF RAPID MCB-2032054, an award from the RCSA Research Corp., and a UC San Diego Moores Cancer Center 2020 SARS-COV-2 seed grant to R.E.A.; NIH grant R01-GM31749 to J.A.M.; NIH grant R01-AI127521 to J.S.M; and NIH (1R01GM115805-01) and NSF (CHE-1807301) to L.T.C.

Acknowledgements

We are grateful for the efforts of the Texas Advanced Computing Center (TACC) Longhorn team and for the compute time made available through a Director's Discretionary Allocation (made possible by the National Science Foundation award OAC-1818253). We thank Dr. Zied Gaieb for helpful discussions around system construction. We thank Mahidhar Tatineni for help with computing on SDSC Comet, as well as a COVID19 HPC Consortium Award for compute time. We also thank Prof. Carlos Simmerling and his research group (SUNY Stony

Brook), and Prof. Adrian Mulholland and his research group (University of Bristol) for helpful

discussions related to the spike protein, as well as Prof. Daniel Zuckerman, Dr. Jeremy

Copperman, Dr. Matthew Zwier, and Dr. Sinam Saglam for helpful methodological discussions.

References

(1) Chan, J. F.-W.; Yuan, S.; Kok, K.-H.; To, K. K.-W.; Chu, H.; Yang, J.; Xing, F.; Liu, J.;

Yip, C. C.-Y.; Poon, R. W.-S.; Tsoi, H.-W.; Lo, S. K.-F.; Chan, K.-H.; Poon, V. K.-M.;

Chan, W.-M.; Ip, J. D.; Cai, J.-P.; Cheng, V. C.-C.; Chen, H.; Hui, C. K.-M.; Yuen, K.-Y.

A Familial Cluster of Pneumonia Associated with the 2019 Novel Coronavirus Indicating

Person-to-Person Transmission: A Study of a Family Cluster. *The Lancet* **2020**, *395*

(10223), 514–523. https://doi.org/10.1016/S0140-6736(20)30154-9.

(2) Lu, R.;Zhao, X.; Li, J.; Niu, P.; Yang, B.; Wu, H.; Wang, W.; Song, H.; Huang, B.; Zhu,

N.; Bi, Y.; Ma, X.; Zhan, F.; Wang, L.; Hu, T.; Zhou, H.; Hu, Z.; Zhou, W.; Zhao, L.;

Chen, J.; Meng, Y.; Wang, J.; Lin, Y.; Yuan, J.; Xie, Z.; Ma, J.; Liu, W. J.; Wang, D.; Xu,

W.; Holmes, E. C.; Gao, G. F.; Wu, G.; Chen, W.; Shi, W.; Tan. W. Genomic

Characterisation and Epidemiology of 2019 Novel Coronavirus: Implications for Virus

Origins and Receptor Binding. *Lancet* **2020**, *395* (10224), 565–574.

https://doi.org/10.1016/s0140-6736(20)30251-8.

(3) Li, F. Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol*

2016, *3* (1), 237–261. https://doi.org/10.1146/annurev-virology-110615-042301.

(4) Wrapp, D.; Wang, N.; Corbett, K. S.; Goldsmith, J. A.; Hsieh, C.-L.; Abiona, O.; Graham,

B. S.; McLellan, J. S. Cryo-EM Structure of the 2019-NCoV Spike in the Prefusion

Conformation. *Science* **2020**, *367* (6483), 1260–1263.

https://doi.org/10.1126/science.abb2507.

(5) Walls, A. C.; Park, Y.-J.; Tortorici, M. A.; Wall, A.; McGuire, A. T.; Veesler, D. Structure,

Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* **2020**, *181* (2),

281-292.e6. https://doi.org/10.1016/j.cell.2020.02.058.

(6) Benton, D. J.; Wrobel, A. G.; Xu, P.; Roustan, C.; Martin, S. R.; Rosenthal, P. B.; Skehel, J.

J.; Gamblin, S. J. Receptor Binding and Priming of the Spike Protein of SARS-CoV-2 for

Membrane Fusion. *Nature* **2020**, *588* (7837), 327–330. https://doi.org/10.1038/s41586-020-

2772-0.

(7) Lu, M.; Uchil, P. D.; Li, W.; Zheng, D.; Terry, D. S.; Gorman, J.; Shi, W.; Zhang, B.;

Zhou, T.; Ding, S.; Gasser, R.; Prévost, J.; Beaudoin-Bussières, G.; Anand, S. P.; Laumaea,

A.; Grover, J. R.; Liu, L.; Ho, D. D.; Mascola, J. R.; Finzi, A.; Kwong, P. D.; Blanchard, S.

C.; Mothes, W. Real-Time Conformational Dynamics of SARS-CoV-2 Spikes on Virus

Particles. *Cell Host Microbe* **2020**, *28* (6), 880-891.e8.

https://doi.org/10.1016/j.chom.2020.11.001.

(8) Casalino, L.; Gaieb, Z.; Goldsmith, J. A.; Hjorth, C. K.; Dommer, A. C.; Harbison, A. M.;

Fogarty, C. A.; Barros, E. P.; Taylor, B. C.; McLellan, J. S.; Fadda, E.; Amaro, R. E.

Beyond Shielding: The Roles of Glycans in the SARS-CoV-2 Spike Protein. *ACS Cent. Sci.*

2020. https://doi.org/10.1021/acscentsci.0c01056.

(9) Gur, M.; Taka, E.; Yilmaz, S. Z.; Kilinc, C.; Aktas, U.; Golcuk, M. Conformational

Transition of SARS-CoV-2 Spike Glycoprotein between Its Closed and Open States. *J.*

Chem. Phys. **2020**, *153* (7), 075101. https://doi.org/10.1063/5.0011141.

(10) Fallon, L.; Belfon, K.; Raguette, L.; Wang, Y.; Corbo, C.; Stepanenko, D.; Cuomo, A.;

Guerra, J.; Budhan, S.; Varghese, S.; Rizzo, R.; Simmerling, C. Free Energy Landscapes

for RBD Opening in SARS-CoV-2 Spike Glycoprotein Simulations Suggest Key

Interactions and a Potentially Druggable Allosteric Pocket. **2020**.

https://doi.org/10.26434/chemrxiv.13502646.v1.

(11) SARS-CoV-2 Simulations Go Exascale to Capture Spike Opening and Reveal Cryptic

Pockets Across the Proteome | bioRxiv

https://www.biorxiv.org/content/10.1101/2020.06.27.175430v3 (accessed Jan 13, 2021).

(12) Huber, G. A.; Kim, S. Weighted-Ensemble Brownian Dynamics Simulations for Protein

Association Reactions. *Biophys. J.* **1996**, *70* (1), 97–110. https://doi.org/10.1016/S0006-

3495(96)79552-8.

(13) Zhang, B. W.; Jasnow, D.; Zuckerman, D. M. The "Weighted Ensemble" Path Sampling

Method Is Statistically Exact for a Broad Class of Stochastic Processes and Binning

Procedures. *J. Chem. Phys.* **2010**, *132* (5), 054107. https://doi.org/10.1063/1.3306345.

(14) Chong, L. T.; Saglam, A. S.; Zuckerman, D. M. Path-Sampling Strategies for Simulating

Rare Events in Biomolecular Systems. *Curr Opin Struct Biol* **2017**, *43*, 88–94.

https://doi.org/10.1016/j.sbi.2016.11.019.

(15) Zuckerman, D. M.; Chong, L. T. Weighted Ensemble Simulation: Review of Methodology,

Applications, and Software. *Annu Rev Biophys* **2017**, *46*, 43–57.

https://doi.org/10.1146/annurev-biophys-070816-033834.

(16) Pratt, L. R. A Statistical Method for Identifying Transition States in High Dimensional

Problems. *J. Chem. Phys.* **1986**, *85* (9), 5045–5048. https://doi.org/10.1063/1.451695.

(17) Zuckerman, D. M.; Woolf, T. B. Transition Events in Butane Simulations: Similarities

across Models. *J. Chem. Phys.* **2002**, *116* (6), 2586–2591.

https://doi.org/10.1063/1.1433501.

(18) Adhikari, U.; Mostofian, B.; Copperman, J.; Subramanian, S. R.; Petersen, A. A.;

Zuckerman, D. M. Computational Estimation of Microsecond to Second Atomistic Folding

Times. *J Am Chem Soc* **2019**, *141* (16), 6519–6526. https://doi.org/10.1021/jacs.8b10735.

(19) Saglam, A. S.; Chong, L. T. Protein–Protein Binding Pathways and Calculations of Rate

Constants Using Fully-Continuous, Explicit-Solvent Simulations. *Chem. Sci.* **2019**, *10* (8),

2360–2372. https://doi.org/10.1039/C8SC04811H.

(20) DeGrave, A. J.; Ha, J.-H.; Loh, S. N.; Chong, L. T. Large Enhancement of Response Times

of a Protein Conformational Switch by Computational Design. *Nature Communications*

2018, *9* (1), 1013. https://doi.org/10.1038/s41467-018-03228-6.

(21) Suárez, E.; Lettieri, S.; Zwier, M. C.; Stringer, C. A.; Subramanian, S. R.; Chong, L. T.;

Zuckerman, D. M. Simultaneous Computation of Dynamical and Equilibrium Information

Using a Weighted Ensemble of Trajectories. *J. Chem. Theory Comput.* **2014**, *10* (7), 2658–

2667. https://doi.org/10.1021/ct401065r.

(22) Watanabe, Y.; Allen, J. D.; Wrapp, D.; McLellan, J. S.; Crispin, M. Site-Specific Glycan

Analysis of the SARS-CoV-2 Spike. *Science* **2020**.

https://doi.org/10.1126/science.abb9983.

(23) Henderson, R.; Edwards, R. J.; Mansouri, K.; Janowska, K.; Stalls, V.; Gobeil, S. M. C.;

Kopp, M.; Li, D.; Parks, R.; Hsu, A. L.; Borgnia, M. J.; Haynes, B. F.; Acharya, P.

Controlling the SARS-CoV-2 Spike Glycoprotein Conformation. *Nature Structural &*

Molecular Biology **2020**, *27* (10), 925–933. https://doi.org/10.1038/s41594-020-0479-4.

(24) Pallesen, J.; Wang, N.; Corbett, K. S.; Wrapp, D.; Kirchdoerfer, R. N.; Turner, H. L.;

Cottrell, C. A.; Becker, M. M.; Wang, L.; Shi, W.; Kong, W.-P.; Andres, E. L.; Kettenbach,

A. N.; Denison, M. R.; Chappell, J. D.; Graham, B. S.; Ward, A. B.; McLellan, J. S.

Immunogenicity and Structures of a Rationally Designed Prefusion MERS-CoV Spike

Antigen. *PNAS* **2017**, *114* (35), E7348–E7357. https://doi.org/10.1073/pnas.1707304114.

(25) Hsieh, C.-L.; Goldsmith, J. A.; Schaub, J. M.; DiVenere, A. M.; Kuo, H.-C.; Javanmardi,

K.; Le, K. C.; Wrapp, D.; Lee, A. G.; Liu, Y.; Chou, C.-W.; Byrne, P. O.; Hjorth, C. K.;

Johnson, N. V.; Ludes-Meyers, J.; Nguyen, A. W.; Park, J.; Wang, N.; Amengor, D.;

Lavinder, J. J.; Ippolito, G. C.; Maynard, J. A.; Finkelstein, I. J.; McLellan, J. S. Structure-

Based Design of Prefusion-Stabilized SARS-CoV-2 Spikes. *Science* **2020**, *369* (6510),

1501–1505. https://doi.org/10.1126/science.abd0826.

(26) Ryan Cross. The Tiny Tweak behind COVID-19 Vaccines. *Chemical & Engineering News*.

September 29, 2020.

(27) Corum, J.; Zimmer, C. Coronavirus Variants and Mutations. *The New York Times*. February

10, 2021.

(28) Amaro, R. E.; Mulholland, A. J. A Community Letter Regarding Sharing Biomolecular

Simulation Data for COVID-19. *J. Chem. Inf. Model.* **2020**, *60* (6), 2653–2656.

https://doi.org/10.1021/acs.jcim.0c00319.