TITLE

The genome of the CTG(Ser1) yeast S*cheffersomyces stipitis* is plastic

Samuel Vega Estevez[1], Andrew Armitage[2], Helen J. Bates[3], Richard J. Harrison[3] and

Alessia Buscaino[1#]

[1] University of Kent, School of Biosciences, Kent Fungal Group, Canterbury Kent,

CT2 7NJ, UK.

[2] Natural Resources Institute, University of Greenwich, Chatham Maritime, Kent,

ME4 4TB, UK.

[3] NIAB Cambridge Crop Research, 93 Lawrence Weaver Road , Cambridge

CB3 0LE

[#] Address correspondence to Alessia Buscaino

Email: A.Buscaino@kent.ac.uk

ABSTRACT

Microorganisms need to adapt to environmental changes, and genome plasticity can

lead to rapid adaptation to hostile environments by increasing genetic diversity.

Here, we investigate genome plasticity in the CTG(Ser1) yeast *Scheffersomyces*

stipitis, an organism with an enormous potential for second-generation biofuel

production. We demonstrate that *S. stipitis* has an intrinsically plastic genome and

that different *S. stipitis* isolates have genomes with distinct chromosome

organisation. Real-time evolution experiments show that *S. stipitis* genome plasticity

is common and rapid as extensive genomic changes with fitness benefits are

detected following *in vitro* evolution experiments. Hybrid MinION Nanopore and

Illumina genome sequencing identifies retrotransposons as major drivers of genome

diversity. Indeed, the number and position of retrotransposons is different in different

S. stipitis isolates, and retrotransposon-rich regions of the genome are sites of

chromosome rearrangements. Our findings provide important insights into the

adaptation strategies of the CTG (Ser1) yeast clade and have critical implications in

the development of second-generation biofuels. These data highlight that genome

plasticity is an essential factor to be considered for the development of sustainable

S. stipitis platforms for second-generation biofuels production.

INTRODUCTION

Eukaryotic genomes are often described as stable structures with well-preserved

chromosome organisation, and genome instability is viewed as harmful. However, an

increasing body of evidence demonstrates that eukaryotic microorganisms have a

plastic genome and genome instability is instrumental for rapid and reversible

adaptation to hostile environments (1–4). This is because genomic instability can

increase genetic diversity, allowing the selection of genotype(s) better adapted to a

new environment (5, 6). Repetitive DNA elements are major contributors to genome

plasticity as repeats can undergo inter and intra-locus recombination, resulting in

gene conversion, gross chromosomal rearrangements and segmental aneuploidies

(7). Transposable Elements (TE), a specific class of repetitive elements, alter

genome organisation by recombination-dependent mechanisms and by jumping to

new sites in the genome (8). TEs belong to two major classes: DNA transposons

(Class II) and retrotransposons (Class I). DNA transposons utilise a "cut and paste"

mechanism in which the parental element excises from its original location before

integrating elsewhere (9). In contrast, retrotransposons replicate through reverse

transcription of their RNA and integrate the resulting cDNA into another locus.

Retrotransposons can be further classified into Long Terminal Repeats (LTR)

retrotransposon and non-LTR retrotransposons (10). LTR retrotransposons are

characterised by two LTR sequences flanking an internal coding region containing

the genes encoding for the structural protein GAG and enzyme POL required for

reverse transcription and integration (11). While POL enzymes are conserved across

organisms, GAG proteins are poorly conserved (12). LINE elements are one of the

most abundant non-LTR retrotransposons and they are typically composed of a 5'

non-coding region, two ORFs (ORF1 and ORF2) and a 3' non-coding region that is

marked by a poly(A) tail (13). ORF1 proteins have a diverse amino acid sequence,

but they often contain a DNA-binding motif (14). ORF2 encodes endonuclease and

reverse transcriptase activity that are critical for transposition (15).

The CTG (Ser1) clade of fungi ,in which the CTG codon is translated as serine rather

than leucine, is an important group of ascomycetous yeasts featuring yeasts that

hold great promises in biotechnology, such as *Scheffersomyces stipitis*, and

dangerous human fungal pathogens, such as *Candida albicans* (16).

The CTG(Ser1) clade comprises several species with different lifestyles and

genomic organisations, including haploid and diploid species that colonise diverse

environments by reproducing sexually or para-sexually (16–19). One common

feature of CTG(Ser1) species is their ability to adapt remarkably well to extreme

environments (20). For example, CTG(Ser1) yeasts can grow on various carbon

sources and are highly tolerant to environmental changes such as changes in

osmolarity (16, 19, 20). It is well established that genome plasticity is a critical

adaptive mechanism in the human fungal pathogens *Candida albicans*, the most

studied CTG (Ser1)-clade member (4). In *C. albicans*, stress increases genome

instability by affecting the rate and type of genomic rearrangements (21). Different

classes of DNA repeats drive this genetic variation, including TEs, long repeats and

Major Repeat Sequences (MRS) (22–24). It is still unknown whether genome

plasticity is a general feature of the CTG(Ser1) clade and whether DNA repeats are

drivers for genome diversity across this yeast group.

This study investigates genome plasticity in *S. stipitis*, a CTG (Ser1)-clade yeast with

great potential for the eco-friendly and ethical production of second-generation

biofuels (25–27). Second-generation biofuels are generated by fermentation of

lignocellulose biomass, produced in large amounts (>1.3 billion tons produced

annually) as waste following agricultural and forestry processing operation (27).

Lignocellulose is a heteropolymer composed of fermentable hexose sugars, such a

glucose, and pentose sugars, such as xylose (28). The yeast *Saccharomyces*

cerevisiae, usually the organism of choice for industrial production of ethanol, is not

suitable for the production of second-generation ethanol because it cannot ferment

pentose sugars as it lacks specific transporters and enzymatic network important for

their metabolism (28). *S. stipitis* holds excellent potential for biofuel derived from

green waste because it is one of the few yeast species that can ferment both hexose

and pentose sugars (25–27). *S. stipitis* is a non-pathogenic haploid yeast that is

found in the gut of wood-ingesting beetles, in hardwood forests or areas high in

agricultural waste (29). Contrary to *C. albicans*, *S. stipitis* has a canonical sexual

cycle whereby mating of haploid cells generate diploid cells that undergo meiosis

and produce haploid spores (30). Although several *S. stipitis* natural isolates are

used for the optimisation of second-generation biofuels production, the genome of

only one strain (Y-11545) has been sequenced and assembled to the chromosomal

level (31). The Y-11545 genome has a size of 15.4 Million base pair (Mbp) organised

in 8 chromosomes and containing ~6000 protein-coding genes (31–33). *S. stipitis*

chromosomes are marked by regional centromeres composed of full-lenghts LTR

retrotrasposons (Tps5a, Tps5b and Tps5c) and non-coding, non-autonomous LARD

(large retrotransposon derivative) elements (31, 33).

To investigate the plasticity of the *S. stipitis* genome, we have taken several

complementary approaches. Firstly, we systematically identified *S. stipitis* DNA

repeats and investigated the genotypic diversity of 27 different *S. stipitis* natural

isolates collected from different environments. Secondly, we combined MinION

Nanopore with Illumina genome sequencing to generate a high-quality chromosome-

level sequence assembly of a second *S. stipitis* natural isolate (Y-7124) and

compared its genome structure to the reference Y-11545 genome. Lastly, we

performed *in vitro* evolution experiments and analysed *S. stipitis* genome

organisation changes following laboratory passaging under stress or unstressed

growth conditions. Thanks to this combined approach, we discovered that the *S.*

stipitis genome is plastic. Genome plasticity is not regulated by stress, however large

chromosome rearrangements are linked to adaptation to hostile environments. We

demonstrate that different *S. stipitis* natural isolates have distinct chromosomal

organisations and that transposable elements drive this extensive intra-species

genetic variation. Our findings have important implications for second-generation

biofuel production as genome plasticity is a paramount factor to be considered for

the successful development of superior biofuel-producer *S. stipitis* strains.

138   **MATERIAL AND METHODS**

Yeast strains and Growth Conditions

Strains were obtained from the Agricultural Research Service (ARS) Collection, run

by the Northern Regional Research Laboratory (NRRL) (Peoria, Illinois, USA), or the

National Collection of Yeast Cultures (NCYC) (Norwich, United Kingdom) (**Table S1)**

and confirmed by sequencing (primers AB798 and AB799 of the 26S rDNA (D1/D2

domain) (34) (**Table S2**). Routine culturing was performed at 30 ºC with 200 rpm

agitation on Yeast Extract-Peptone-D-Glucose (YPD) media. Phenotypic and *in vitro*

evolution analyses were conducted on Synthetic Complete (SC) media containing

glucose (SC-G), xylose (SC-X), or a mixture of 60% glucose and 40% xylose (SC-

G+X). SC-G was used as a reference media as glucose is the preferred carbon

source for both the model system *S. cerevisiae* and *S. stipitis*, SC-X was used

because of *S. stipitis* unique ability to utilise xylose as a carbon source and SC G+X

was used because this sugar combination resemble the ratio found in lignocellulose

(28). Uridine (0.08 g/L in YPD and SC) and adenine hemisulfate (0.05 g/L in YPD)

were added as growth supplements. Solid media were prepared by adding 2%

agar.

Contour-clamped homogeneous electric field (CHEF) electrophoresis

Intact yeast chromosomal DNA was prepared as previously described (35).

Briefly, cells were grown overnight and spheroplast were prepared in an agarose

plug by treating cells (~ OD_{600}=7) with 0.6 mg/ml Zymolyase 100T (Amsbio #120493-

1) in 1% Low Melt agarose (Biorad® # 1613112). Chromosomes were separated in a

1% Megabase agarose gel (Bio-Rad) in 0.5X TBE using a CHEF DRII apparatus.

Run conditions as follows: 60-120s switch at 6 V/cm for 12 hours followed by a 120-

300s switch at 4.5 V/cm for 12 hours, 14 °C. Chromosomes were visualised by

staining the gel 0.5x TBE with ethidium bromide (0.5 µg/ml) for 30 minute, followed

by destaining in water for 30 minutes. Images were capture using a Syngene GBox

Chemi XX6 gel imaging system.

Southern Blotting

DNA from CHEF gels were transferred overnight to a Zeta-Probe GT

Membrane (Biorad®, #162-0196) in 20x SSC and crosslinked using UV (150 mJ).

Probing and detection of the DNA were conducted as previously described (36).

Briefly, probes were generated by PCR incorporation of DIG-11-dUTP into target

sequences following manufacturer's instructions (Roche). Chromosome 5-

chromosome 7 translocation was detected using primers AB1028 and AB1029

amplifying a 180 bp region of chromosome 5 (Chr5 nt: 448,855-449,034) in Y-11545

and in chromosome 7 of Y-7124 (Chr7 nt: 494,698-494,877) (**Table S2**). The

membrane was hybridised overnight at 42 ºC with DIG Easy Hyb (Roche®,

11603558001). The DNA was detected with anti-digoxigenin-Alkaline Phosphatase

antibody (Roche®, #11093274910) and CDP Star ready to use (Roche®,

#12041677001) according to manufacturer instructions.

Phenotypic characterisation

Growth analyses were performed using a plate reader (SpectrostarNano,

BMG labtech) in 96 well plate format at 30 °C for 48 hours in SC-G, SC-X or SC-

G+X. The growth rate (μ, hours^{-1}) was calculated using: $\mu = (\ln(X_2)-\ln(X_1))/(t_2-t_1)$,

where: (i) X_1 is the biomass concentration (OD_{600}) at time point one (t_1, lag time) (ii)

X_2 is the biomass concentration (OD_{600}) at time point two (t_2, end of exponential

growth phase). The maximum OD (OD units) was determined with the MAX() from

Excel (Microsoft®). The lag time (minutes) was determined visually as the time in

which the exponential growth starts. Experiments were performed in 3 technical and

3 biological replicates. Heatmaps showing the average of 3 biological replicates were

generated by R using the library *pheatmap*. ANOVA test was performed to study

differences on growth rate, maximum OD and lag time between the strains. The

equality of variances presumption was tested using Levene's test, whereas the

normality of the data was tested by Shapiro-Wilk. When both assumptions were

satisfied, a Tukey's honest significant test was used to determine significant

differences between the natural isolates and the reference Y-11545 strain. When

assumption of equal variance were violated, one-way test was used to indicate

significance. In the case of equal variances, but a non-normal distribution of data, the

Kruskal-Wallis rank sum test was used to indicate statistical differences and

significance was determined by pairwise testing. A p-value lower than 0.05 was

considered significant for all these statistical tests.

Adaptive Laboratory Evolution

A single colony of the *S. stipitis* strain NRRL Y-7124 was grown overnight in 5

ml of YPD at 30 ºC, plated in YPD at a cell density of 100 and grown 48 hours at 30

ºC. 36 single colonies were streaked in two SC-G+X plates and grown at 30 ºC and

37 ºC, respectively and streaked daily for a total of 56 passages (8 weeks). The

karyotype variability of the colonies was assessed by CHEF electrophoresis.

Phenotypic differences were assessed by spotting assays. Strains with

rearrangements were grown overnight is SC-G+X and were diluted to an $OD_{600}=1$.

From this, five 1/10 serial dilutions were prepared and the cells were plated in SC-

G+X using a replica plater (Sigma Aldrich, R2383-1EA) and grown for 48 hours at

both 30 ºC and 37 ºC. Strains with no karyotypic modifications after evolution were

also used as control.

Identification of DNA repeats

Long sequences (>100 nucleotides) present more than once in the Y-11545

and Y-7124 genomes were identified by aligning each genome to itself using

BLASTN. Repetitive elements (E < 1e-04) were manually verified using

IGV/SNAPGene, and clustered repeats were combined. This repeats dataset was

manually examined to further classify it as (a) related to transposable elements (b)

telomeric repeats, (c) centromeres (d) belonging to protein coding gene families and

(e) MRS repeats. Transposons were classified using established guidelines (10).

Briefly, LTR-transposons were identified by detecting two Long-terminal Repeat

sequences (size 260-430 nt) flanking an internal coding region. These potential LTR-

transposons were further annotated for the presence of the following marks: LTR

flanked by a TG and CA di-nucleotides, presence of a Primer Binding Site (PBS) with

homology to *S. stipitis* tRNAs (GtRNAdb (http://gtrnadb.ucsc.edu/index.html),

presence of a coding region with homology to *pol* gene and containing an Integrase

(INT), Reverse Transcriptase (RT) and RNAse H (RH) domain. Non-LTR LINE

transposons were identified by detection of coding regions homologous to LINE

retrotransposons ORF1 (containing a Zn-finger), ORF2 (containing an Endonuclease

and a Reverse Transcriptase domain) and terminal Poly-A sequence.

Retrotransposons were classified into different families based on sequence similarity

with a 90% cut-off. Terminal telomeric tandem repeats were identified using Tandem

Repeats Finder (37) with default parameters. Regional centromeres were identified

based on them being the only regions of the genome with a large retrotransposon

Tps5 cluster (~ 20-40 kb) as previously described (33). Gene families were identified

by extracting coding-regions from our repeats datasets and performing Clustal

Omega sequence alignment and PFAMs domain identification using

SMART(http://smart.embl.de) (38). The identified gene families were compared to

published information (39). The presence of MRS repeats was explored using

BLASTN and by searching for clusters of non-coding tandem repeats, a hallmark of

C. albicans MRS, with no-homology to retrotransposons and not-located at

chromosome ends. Sequence alignments were visualised with Jalview v2.11.1.0

(40). Phylogenetic trees were generated with phyloT : a phylogenetic tree generator

(biobyte.de) using default parameters and visualised with Itol (https://itol.embl.de/).

Genome sequencing

The genome of *S. stipitis* isolate Y-7124 was sequenced by Illumina short-

read and MinION long-read technologies. To this end, DNA was extracted from an

overnight culture using the QIAGEN genomic tip 100/G kit (Qiagen®, #10243)

according to manufacturing protocol. For long-read sequencing, MinION (Oxford

Nanopore, Oxford UK) was performed on a DNA library prepared from size selected

gDNA. DNA fragments greater than 30 Kb were selected using a Blue Pippin (Sage

Science) and concentrated using Ampure beads. From this, a DNA library was

prepared using a Ligation Sequencing Kit 1D (SQK-LSK108) and run on the Oxford

Nanopore MinION flowcell FLOMIN 106D. The same gDNA extract was also used

for the preparation of Illumina libraries. In this case, the DNA was sheared using the

Covaris M220 with microTUBE-50 (Covaris 520166) and size selected using the

Blue Pippin (Sage Science). The library was constructed using a PCR-free kit with

NEBNext End Repair (E6050S), NEBNext dA-tailing (E6053S) and Blunt T/A ligase

(M0367S) New England Biolabs modules. Sequencing was performed on a MiSeq

Benchtop Analyzer (Illumina) using a 2x300bp PE (MS-102-3003) flow cell.

Genome assembly

Base-calling and demultiplexing were conducted with Albacore v2.3.3 (available at

https://community.nanoporetech.com). Adapters and low-quality data were trimmed

using the eautils package fastq-mcf 1.04.636

(https://expressionanalysis.github.io/ea-utils/). On nanopore sequence data, adapter

trimming was performed with Porechop v.0.1.0 (https://github.com/rrwick/Porechop).

Genome assembly was completed using long reads, with read correction performed

with Canu v1.8 (41) followed by assembly in SMARTdenovo github commit id

61cf13d (42)). The draft assembly was corrected using the corrected nanopore reads

through five rounds of Racon github commit 24e30a9 (43), and then by raw fast5

files using 10 rounds of Nanopolish v0.9.0 (44). Illumina sequencing reads were then

used to polish the resulting assembly through 10 rounds of Pilon v1.17 (45).

Following genome assembly, BUSCO v3 was run to assess evolutionary conserved

gene content (46), using the Saccharomycetales_odb9 gene database. The

Saccharomycetales database contains 1711 genes, which are therefore expected to

be present in *S. stipitis*. Of these, 1683 (98.36%) were identified in the Y-7124

assembly demonstrating a good level of completeness (>95%) (**Table S8**). Assembly

size and contiguity statistics were assessed using QUAST v4.5 (47). This initial

assembly of the nuclear genome contained 10 contigs. A chromosome level

assembly was produced by identification of overlapping regions between the contigs:

a 244 Kbp overlapping region between contig 7 and 2 led to the final assembly of

Chromosome 1, a 83 Kpb overlapping region between contig 9 and 10 led to the final

assembly of Chromosome 8.

Genome annotation

Genome annotation was performed using FUNGAP v1.0.1 (48) with fastq

reads from NCBI SRA accession SRR8420582 used as RNA-Seq training data and

protein sequences taken from NCBI assembly accession GCA_000209165.1 for *S.*

stipitis NRRL Y-11545 (CBS6054) used for example proteins. Protein fasta files were

extracted from predicted gene models using the yeast mitochondrial code (code 3)

and the alternative yeast nuclear code (code 12). Functional annotation of gene

models was performed through BLASTp searches vs all proteins from the NCBI

reference fungal genomes (downloaded 11th April 2020), retrieving the top-scoring

blast hit with an E-value < 1×10^{-30}. These annotations were supplemented with

domain annotations from Interproscan v5.42-78.0 (49). The annotated genome was

submitted to NCBI, with submission files prepared using GAG v2.0.1

(http://genomeannotation.github.io/GAG.), Annie github commit 4bb3980

(http://genomeannotation.github.io/annie) and table2asn_GFF v1.23.377 (available

from https://ftp.ncbi.nih.gov/toolbox/ncbi_tools/converters/by_program/tbl2asn/).

Comparative genomics

Whole-genome alignment between Y-7124 and Y-11545 was performed using

the nucmer tool from the MUMmer package v4.0 (50) with results visualised using

Circos v0.6 (51). Orthology analysis was performed between predicted proteins from

these isolates using OrthoFinder v2.3.11 (52), with results visualised using the

package VennDiagram in R (53).

Sequence variants were identified in Y-7124 through comparison to the Y-

11545 assembly. Short read sequence data for Y-7124 were aligned to the reference

genome using BWA v 0.7.15-r1140 (54), before filtering using using picardtools

v2.5.0 to remove optical duplicates (http://broadinstitute.github.io/picard/). SNP and

insertion/deletion (InDel) calling was performed using GATK4 (55). Low confidence

variants were then filtered using VCFtools v0.1.15 (56) using minimum mapping

quality of 40, phred quality of 30, read depth of 10 and genotype quality of 30. Effect

of variants on NRRL Y-11545 gene models was determined using SnpEff v4.2 (57).

312   **RESULTS**

Classification of *S. stipitis* DNA repeats

DNA repeats are drivers of genome variation. Understanding the repertoire of

repetitive elements associated with a genome is critical to gain insights into the

genome diversity of a specific organism. Comparative genomic analyses have

identified different repetitive elements in some CTG(Ser1) clade members, yet a

comprehensive survey of *S. stipitis* repetitive elements is lacking (18, 58). Therefore,

we sought to classify the major classes of repetitive elements associated with the Y-

11545 sequenced genome by aligning the genomic sequence of each strain to itself

and identifying long sequences (>100 nucleotides) present more than once in the

genome. The genomic position of these repeats was manually verified, and clustered

repeats were combined and categorised depending on their genomic position,

structure and sequence similarity. Our analyses identified known *S. stipitis* repeat-

rich loci such as centromeric transposon-clusters, the NUPAV sequence, an

integrated L-A ds-RNA virus, and several gene families (32, 33, 59). As observed in

other members of the CTG (Ser1)-clade, we did not detect any MRS repeats, a class

of repetitive elements found only in *C. albicans* and the closely related *C. dubliensis*

and *C. tropicalis* species (58, 60). Here we focus on intra-chromosomal or inter-

chromosomal repeats that have not been described to date: non-centromeric TEs,

subtelomeric regions and telomeric repeats (**Fig 1**).

We identified six novel retrotransposon families scattered along chromosome arms:

3 LTR retrotransposons (*Ava*, *Bea* and *Caia*) and 3 LINE retrotransposons (*Ace*, *Bri*

and *Can*) (**Fig 1A**, **Table S3**). *Ava*, *Bea* and *Caia* have a similar structure where two

identical LTR sequences flank an internal domain. The internal domain contains two

ORFs: one encoding for a putative POL and one encoding for an *S. stipitis*-specific

protein that we named LTR-Associated Protein (Lap1 in *Ava*, Lap2 in *Bea* and Lap3

in *Caia*). Homology search failed to identify any GAG gene associated with the *Ava*,

Bea and *Caia* retrotransposons. As Gag proteins are poorly conserved among

different organisms, we hypothesise that the Lap proteins are Gag proteins.

Ace, *Bri* and *Can* are LINE elements composed of the Non-Coding regions NC-1 and

NC-2 surrounding an internal coding region encoding for a Pol enzyme and an *S.*

stipitis-specific LINE Associated protein (Linea1 in *Ace*, Linea2 in *Bri* and Linea3 in

Can). Linea1 and Linea2, but not Linea 3, have a Zinc-Finger DNA binding motif

(**Table S3**). Comparison across the CTG (Ser1)-clade revealed that *S. stipits* TE

repertoire is typical of this clade. Indeed, retrotransposons are common in this yeast

group: the genome of all species analysed contains LTR elements, whereas LINE

elements are present in 6/8 species (**Fig 1B, Table S4**). Similarly to other CTG-

(Ser1) clade yeast, we did not detect any DNA transposons integrated into the *S.*

stipitis genome (**Fig 1B, Table S4**).

Our repeat analysis demonstrates that the terminal sequences of *S. stipitis*

chromosomes are repeat-rich and composed of two elements with different degree

of repetitiveness: telomere proximal-repeats and subtelomeric regions. The

telomeric repeats are non-canonical and composed of 24-nucleotide units repeated

in tandem. Each unit contains a TG motif reminiscent of typical telomeric repeats

(**Fig 1C**). *S. stipitis* subtelomeric regions (the ~30KB region adjacent to telomeric

repeats) are enriched in retrotransposon-derived elements. Indeed, DNA sequences

with homology to *Bea* LTR-retrotransposons and *Ace* LINE-elements are found in

5/16 subtelomeric regions (**Fig 1D, Table S3**). No full length-retrotransposons are

detected at these genomic locations. Subtelomeric regions contain several gene

families members, including gene encoding for ATP-dependent DNA helicases

(found in 7/16 subtelomeres), fungal-specific transcription factors (8/16

subtelomeres), MFS transporters (8/16 subtelomeres) and Agglutinine-like proteins

(11/16 subtelomeres) (**Fig 1D, Table S5**) (39). In summary, our analysis

demonstrates that the *S. stipitis* genome contains several classes of repetitive

elements that could be major

contributors of genome plasticity.

***S. stipitis* natural isolates have distinct genomic organisations**

Having identified *S. stipitis* DNA repeats, our next step was to examine *S. stipitis*

phenotypic and genotypic diversity across a geographically diverse set of strains

(n=27) that were collected in different habitats (**Table S1** source NRRL and NCYC

collection), and that includes the sequenced Y-11545 strain (31). rDNA fingerprinting

confirm that all isolates belong to the *S. stipitis* species (D1/D2 domain of the S26S

rDNA similarity >99 %) (**Table S6**). Phenotypic analyses established that the natural

isolates vary in their ability to utilise and grow on different carbon sources. Indeed,

when compared to the reference Y-11545 strain, different natural isolates cultured in

Synthetic Complete media containing the hexose sugar Glucose (SC-G), the

pentose sugar Xylose (SC-X) or a mixture of both sugars as found in lignocellulose

(SC-G+X) display distinct growth rate, maximum culture density and lag phase (**Fig**

2A and **Table S7**). To determine whether the natural isolates have distinct genomic

organisations, we analysed their karyotype by chromosomes Contour-clamped

Homogenous Electric Field (CHEF) gel electrophoresis, a technique allowing

chromosome separation according to size. The CHEF electrophoresis analysis

reveals clear differences in chromosome patterns demonstrating that *S. stipitis*

natural isolates have a genome organised in different-sized chromosomes (**Figure**

2B). We concluded that intra-species phenotypic and genotypic variation is a

common feature of *S. stipitis.*

Hybrid genomic sequencing identifies transposable elements as drivers of *S.*

***stipitis* genome plasticity**

To date, only one *S. stipitis* isolate (Y-11545) has been sequenced and assembled

at chromosome level (31). To gain insights into *S. stipitis* genetic diversity, we

generated a chromosome-level sequence assembly of a second *S. stipitis* natural

isolate (Y-7124) by combining MinION Nanopore with Illumina genome sequencing.

This isolate was chosen because *(i)* karyotypic analysis reveals that its genomic

organisation is distinct from the genomic organisation of the reference strain Y-

11545, and *(ii)* Y-7124 is widely used both for industrial applications and for basic

research (61).

The Y-7124 genome was sequenced to 186.88x coverage resulting in a 15.69 Mb

assembly arranged in 10 contigs (**Table S8**). High accuracy reads from Illumina-

sequencing enabled the correction of errors that are associated with the MinION

technology. A final chromosome-level assembly was produced by manually

identifying overlapping regions between contigs. Comparing the Y-7124 and Y-

11545 nucleotide sequences reveals that the two natural isolates overall share a

similar coding DNA sequence. The total number of SNPs between the two natural

isolates is 50,495 SNPs, equating to one variant every 306 bases. The majority of

these SNPs are synonymous changes (16,294 =74.25%), while ~25% (5,622) of

SNPs are missense and only (0.13% (28) are nonsense (**Table S9**). Despite this

high DNA sequence similarity, the Y-7124 genome is organised in eight

chromosomes with different sizes and organisations from that of Y-11545 (**Fig 3A**).

Comparison of the Y-7124 and Y-11545 genomes establishes that retrotransposons

are significant drivers of *S. stipitis* genome diversity as one of the most prominent

differences between the two genomes is the abundance and localisation of these

retrotransposons (**Fig 3B**). Indeed, the number of LTR and LINE non-centromeric

retrotransposons and transposons-derived repeats is greater in the Y-11545

reference genome compared to the Y-7124 genome: retrotransposons, solo LTR and

truncated LINE elements account for approximately 2% of the reference Y-11545

genome and only for ~1% of the Y-7124 genome (**Fig 3C**). We classified

retrotransposons loci present in both isolates (ancestral loci), those present in the

reference Y-11545 genome but absent in Y-7124 (deletion loci) and those not

present in the reference genome but present in a given strain (insertion loci). Out of

69 transposons loci, only ten ancestral loci (~15%) were detected in the two isolates.

These sites are likely to be inactive transposons or transposons that rarely

transpose. In addition, we detected 42 deletion loci (60 %) and 17 (24%) insertion

loci (**Fig 3D**). The presence of deletion and insertion loci suggests that *S. stipitis* LTR

transposons and LINE elements are active and competent of transposition. Although

active transposons can insert into genes to cause functional consequences (62), we

did not detect any TE-driven alteration in coding regions.

Transposable Elements are sites of chromosome rearrangements

Comparison of the Y-11545 and Y-7124 genome reveals that transposon-rich

regions are sites of complex chromosome rearrangements. Indeed, a transposon-

rich region is the breakpoint of a reciprocal translocation between chromosome 5

and chromosome 7. This translocation causes the size change of chromosome 5 [Y-

7124] and chromosome 7 [Y-7124] detected by CHEF karyotyping (**Fig 4A**). Southern

analyses with a probe specific for chromosome 5 [Y-11545] confirms this finding (**Fig**

S1). The evolutionary history of Y-11545 and Y-7124 is unknown, and therefore it is

difficult to predict the molecular events underlying these genomic changes. However,

sequence analysis of the rearrangement breakpoint reveals that this structural

variation occurs in a genomic region that *(i)* contains homologous sequences

between chromosome 5 and 7 and *(ii)* is transposon-rich and contains two inverted

repeats on chromosome 7 (**Fig 4B**). A second significant difference between the

genome organisation of Y-11545 and Y-7124 is found at subtelomeric regions: these

regions differ in the number and organisation of subtelomeric gene families and in

the number of transposon-associated repeats (**Fig 4C**). Lastly, we detected a distinct

centromeres organisation where the numbers of *Tps5* retrotransposons, LTRs and

LARD regions differ between the two isolates (**Fig 4D**). The presence of transposons

and transposon-derived repeats associated with all these genomic locations strongly

suggest that retrotransposons have mediated the chromosomal rearrangement by

recombination-mediated mechanisms. Therefore, changes in transposons

organisation are responsible for the bulk of genomic changes identified in two

different natural isolates.

***S. stipitis* real-time evolution leads to extensive genomic changes**

Our results demonstrate that intraspecies genetic diversity is common in *S. stipitis*.

However, as the evolutionary history of the analysed natural isolates are unknown, it

is difficult to predict whether the observed genomic changes are due to the selection

of rare genomic rearrangements events. To determine the time scale of *S. stipitis*

genome evolution, we investigated the genome organisation of 72 single colonies

passaged daily for 8 weeks (56 passages, ~672 divisions) in SC-G+X, as its sugar

composition resembles what found in lignocellulose (29) (**Fig 5A**). Strains were

grown at 30 °C, a temperature that does not lead to any growth defect, and 37 °C, a

stressful temperature that strongly inhibits *S. stipits* growth (**Fig 5B**). CHEF gel

electrophoresis was conducted to identify possible changes in the chromosome

organisation of the evolved strains. This analysis identifies genome rearrangements

in 19/36 strains evolved at 30 °C and 12/36 strains evolved at 37 °C (Blue and

Magenta- **Fig 5C**). Thus, changes in chromosome organisation were detected in the

presence (37 °C) or absence (30 °C) of stress. To test whether chromosome

rearrangements are associated with a fitness benefit, we tested the ability of the

parental and 37 °C-evolved strains to grow in SC-G+X media at permissive (30 C)

and restrictive (37 C) temperature (**Fig 5D**). This analysis demonstrates that 37 °C-

evolved strains with no chromosomal rearrangement grow poorly at 37 °C (**Fig 5D**).

In contrast, 5/12 37 °C-evolved strains with chromosome rearrangements grow

better than the parental strain at this restrictive temperature (**Fig 5D**). This result

suggests that changes in chromosome organisation have an adaptive value. Thus,

genome plasticity is a defining feature of the *S. stipitis* genome, and its genome can

rapidly change in mitotic cells propagated *in vitro*. Our results strongly suggest that

the extensive genomic changes can lead to adaptation to hostile environments.

DISCUSSION

Here we demonstrate that the yeast *S. stipitis* has a plastic genome and that

genome plasticity is linked to adaptation to hostile environments. We show that non-

centromeric retrotransposons are significant drivers of *S. stipitis* genome diversity.

These findings have important implications for developing economically viable

second-generation biofuels and better understanding the CTG (Ser1)-clade biology.

Retrotransposon are drivers of *S. stipitis* genome diversity

Our repetitive sequence analysis demonstrates that *S. stipitis* has a DNA repeats

content typical of the CTG (Ser1)-clade including TEs, non-canonical terminal

telomeric repeats and subtlomeric regions. As observed in other members of the

CTG (Ser1)-clade (60), we did not detect any DNA-transposons or MRS repeats.

One of our major findings is that that non-centromeric retrotransposons are

significant drivers of *S. stipitis* genome diversity. Our data support the hypothesis

that *S. stipitis* TEs generate genome diversity via two distinct mechanisms:

transposition into new genomic locations and recombination-mediated chromosome

rearrangements. Indeed, we demonstrated that the number and genomic position of

non-centromeric retrotransposons vary between the Y-11545 and Y-7124 *S. stipitis*

isolates. Significantly, we did not detect transposon insertions into coding regions.

However, transposons might alter *S. stipitis* gene expression by inserting into gene

regulatory regions (62). We propose that *S. stipitis* transposons are active and

generate genome diversity by jumping into different genomic locations. Our data also

indicate that TEs can generate further genome diversity though either homologous

recombination of nearly identical TE copies or by faulty repair of double-strand

breaks generated during transposable elements excision (62). Indeed, we find that

the translocation breakpoint between chromosome 5 and chromosome 7 is enriched

in retrotransposons. Furthermore, TE-rich subtelomeric regions and centromeres

have a distinct organisation in the two analysed isolates suggesting that the

transposons drive this genetic diversity. We hypothesise that transposons elements

cause the genetic variability observed during laboratory passaging. In the future, it

will be important to apply the hybrid genome sequencing approaches presented in

this study to dissect the nature of these rearrangements.

Genome plasticity and production of second-generation biofuels

One of our key findings is that the *S. stipitis* genome is intrinsically plastic and that

chromosome rearrangements are frequent events under stress or unstressed

conditions. Second-generation biofuels, generated by fermentation of agriculture and

forestry waste, have an enormous potential to meet future energy demands and

significantly reduce petroleum consumption. To meet the requirements for industrial

applications, second-generation biofuels need to be generated by microorganisms

that can efficiently utilise and ferment all the sugars found in lignobiomass (63).

Consequently, *S. stipitis* is one of the most promising yeast for producing second-

generation bioethanol as it can efficiently ferment both hexose and pentose sugars

(25, 26, 29). However, robust economically viable *S. stipitis* platforms still require

significant development as this organism struggles to survive under the harsh

environments generated during second-generation biofuel production. For example,

S. stipitis growth and fermentation is inhibited by the chemical pre-treatment required

to extract glucose and xylose from lignobiomass (61). Growth is also inhibited at high

ethanol concentration, and *S. stipitis* ferments xylose less efficiently than glucose.

Evolutionary engineering approaches under selective conditions (i.e. presence of

inhibitory compounds, high concentration of xylose or ethanol) have been applied to

isolate better performing *S. stipitis* strains (61).

Our data predict that the genetic make-up and associated improved phenotypes of

superior biofuel-producer strains are unstable and that the genetic drivers of

improved phenotypes might be lost over time. This hypothesis could explain why

short-read Illumina genome sequencing has failed to identify point mutations or

indels that could explain the superior performance of *S. stipitis* strains (64). It is also

possible that *S. stipitis* superior strains carry stable complex chromosomal

rearrangements with a breakpoint at DNA repeats. Such rearrangements could not

have been identified by Illumina sequencing as short sequenced fragments will not

resolve changes associated with long repetitive elements. Thus, economically viable

use of *S. stipitis* for second-generation biofuels production will require an in-depth

analysis of the genomic structures of superior strains.

Genome plasticity in the CTG (Ser1)-clade

The CTG-Ser1 clade is an incredibly diverse yeast group that includes many

important human pathogens and non-pathogenic species (17). Our data support the

hypothesis that genome plasticity is a general feature of the CTG (Ser1) yeast clade

as it has been observed in *C. albicans* and *S. stipitis* ((21, 22, 65) and this study),

two organisms with a very different lifestyle. Indeed, while *C. albicans* is a diploid

opportunistic human fungal pathogen that lives almost exclusively in the human host,

S. stipitis is a haploid non-pathogenic yeast found in the gut of wood-ingesting

beetles hardwood forests or areas high in agricultural waste (29, 66). Furthermore,

while *C. albicans* lacks a canonical sexual cycle and its associated meiosis, *S.*

stipitis has a canonical sexual cycle whereby mating of haploid cells generate diploid

cells that undergo meiosis and produce haploid spores (30).

Our results highlight that stress regulates genome plasticity differently in *C. albicans*

and *S. stipitis*. It has been demonstrated that stress exacerbates *C. albicans* genome

instability (21, 67). In contrast, we found that *S. stipitis* genome instability is not

regulated by stress as we detected a similar rate of chromosomal rearrangements

when cells are continuously passaged in unstress (30 ºC) or stress (37 ºC)

conditions. Importantly, we also demonstrated that the large genomic changes are

associated with fitness benefits suggesting that genome plasticity is instrumental for

adaptation to hostile environments.

In summary, our study demonstrates for the first time that *S. stipitis* genome is

plastic. Understanding the cause and effect of this extensive genome plasticity is of

paramount importance to understand the biology of the CTG(Ser1)-clade of fungi.

DATA AVAILABILITY

This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank

under the accession JADGGA000000000. The version described in this paper is

version JADGGA010000000. Illumina and nanopore sequence data associated with

this work have been deposited on the Sequence Read Archive (SRA) under

BioProject PRJNA609885.

FUNDING

This work was supported by the University of Kent Vice-Chancellor's Research

Scholarship (to SV), BBSRC grants (Grant number BB/L008041/1 to AB,

BB/P020364/1 to RJH) and an MRC grant (MR/M019713/1 to AB)

CONFLICT OF INTERESTS DISCLOSURE

None declared.

ACKNOWLEDGEMENTS

We thank Dr Patricia Slininger, members of the Buscaino Lab, the Kent Fungal

Group and Dr Jan Soetaert, for discussion and critical reading of the manuscript.

REFERENCES

1. Burns KH. 2017. Transposable elements in cancer. Nat Rev Cancer 17:415–
424.

2. Colnaghi R, Carpenter G, Volker M, O'Driscoll M. 2011. The consequences of
structural genomic alterations in humans: Genomic Disorders, genomic
instability and cancer. Semin Cell Dev Biol 22:875–885.

3. Galhardo RS, Hastings PJ, Rosenberg SM. 2007. Mutation as a Stress
Response and the Regulation of Evolvability. Crit Rev Biochem Mol Biol
42:399–435.

4. Buscaino. 2019. Chromatin-Mediated Regulation of Genome Plasticity in
Human Fungal Pathogens. Genes (Basel) 10:855.

5. Hirakawa MP, Martinez D a, Sakthikumar S, Anderson MZ, Berlin A, Gujja S,
Zeng Q, Zisson E, Wang JM, Greenberg JM, Berman J, Bennett RJ, Cuomo C
a, Aviv R. 2015. Genetic and phenotypic intra-species variation in Candida
albicans 1–13.

6. Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A,
Barre B, Freel K, Llored A, Cruaud C, Labadie K, Aury J-M, Istace B,
Lebrigand K, Barbry P, Engelen S, Lemainque A, Wincker P, Liti G,
Schacherer J. 2018. Genome evolution across 1,011 Saccharomyces
cerevisiae isolates Species-wide genetic and phenotypic diversity. Nature.

7. Aguilera A, García-Muse T. 2013. Causes of Genome Instability. Annu Rev
Genet 47:1–32.

8. Fedoroff N. 2000. Transposons and genome evolution in plants. Proc Natl
Acad Sci U S A. National Academy of Sciences.

9. Feschotte C, Pritham EJ. 2007. DNA Transposons and the Evolution of

Eukaryotic Genomes. Annu Rev Genet 41:331–368.

10. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A,

Leroy P, Morgante M, Panaud O, Paux E, SanMiguel P, Schulman AH. 2007.

A unified classification system for eukaryotic transposable elements. Nat Rev

Genet. Nature Publishing Group.

11. Havecker ER, Gao X, Voytas DF. 2004. The diversity of LTR retrotransposons.

Genome Biol 5.

12. Llorens C, Muñoz-Pomer A, Bernad L, Botella H, Moya A. 2009. Network

dynamics of eukaryotic LTR retroelements beyond phylogenetic trees. Biol

Direct 4:1–31.

13. Finnegan DJ. 1997. Transposable elements: How non-LTR retrotransposons

do it. Curr Biol 7:245–248.

14. Januszyk K, Li PWL, Villareal V, Branciforte D, Wu H, Xie Y, Feigon J, Loo JA,

Martin SL, Clubb RT. 2007. Identification and solution structure of a highly

conserved C-terminal domain within ORF1p required for retrotransposition of

long interspersed nuclear element-1. J Biol Chem 282:24893–24904.

15. Luan DD, Korman MH, Jakubczak JL, Eickbush TH. 1993. Reverse

transcription of R2Bm RNA is primed by a nick at the chromosomal target site:

A mechanism for non-LTR retrotransposition. Cell 72:595–605.

16. Papon N, Courdavault V, Clastre M. 2014. Biotechnological potential of the

fungal CTG clade species in the synthetic biology era. Trends Biotechnol

32:167–168.

17. Gabaldón T, Naranjo-Ortíz MA, Marcet-Houben M. 2016. Evolutionary

genomics of yeast pathogens in the Saccharomycotina. FEMS Yeast Res 16.

18. Butler G, Rasmussen MD, Lin MF, Santos MAS, Sakthikumar S, Munro CA,

Rheinbay E, Grabherr M, Forche A, Reedy JL, Agrafioti I, Arnaud MB, Bates S,

Brown AJP, Brunke S, Costanzo MC, Fitzpatrick DA, De Groot PWJ, Harris D,

Hoyer LL, Hube B, Klis FM, Kodira C, Lennard N, Logue ME, Martin R,

Neiman AM, Nikolaou E, Quail MA, Quinn J, Santos MC, Schmitzberger FF,

Sherlock G, Shah P, Silverstein KAT, Skrzypek MS, Soll D, Staggs R,

Stansfield I, Stumpf MPH, Sudbery PE, Srikantha T, Zeng Q, Berman J,

Berriman M, Heitman J, Gow NAR, Lorenz MC, Birren BW, Kellis M, Cuomo

CA. 2009. Evolution of pathogenicity and sexual reproduction in eight Candida

genomes. Nature 459:657–662.

19. Wohlbach DJ, Kuo A, Sato TK, Potts KM, Salamov AA, LaButti KM, Sun H,

Clum A, Pangilinan JL, Lindquist EA, Lucas S, Lapidus A, Jin M, Gunawan C,

Balan V, Dale BE, Jeffries TW, Zinkel R, Barry KW, Grigoriev I V., Gasch AP.

2011. Comparative genomics of xylose-fermenting fungi for enhanced biofuel

production. Proc Natl Acad Sci 108:13212–13217.

20. Krassowski T, Coughlan AY, Shen X-X, Zhou X, Kominek J, Opulente DA,

Riley R, Grigoriev I V., Maheshwari N, Shields DC, Kurtzman CP, Hittinger CT,

Rokas A, Wolfe KH. 2018. Evolutionary instability of CUG-Leu in the genetic

code of budding yeasts. Nat Commun 9:1887.

21. Forche A, Abbey D, Pisithkul T, Weinzierl MA, Ringstrom T, Bruck D, Petersen

K, Berman J. 2011. Stress alters rates and types of loss of heterozygosity in

Candida albicans. MBio 2.

22. Todd RT, Selmecki A. 2020. Expandable and reversible copy number

amplification drives rapid adaptation to antifungal drugs. Elife 9:1–33.

23. Todd RT, Wikoff TD, Forche A, Selmecki A. 2019. Genome plasticity in

Candida albicans is driven by long repeat sequences. Elife 8.

24. Hirakawa MP, Martinez DA, Sakthikumar S, Anderson MZ, Berlin A, Gujja S,

Zeng Q, Zisson E, Wang JM, Greenberg JM, Berman J, Bennett RJ, Cuomo

CA. 2015. Genetic and phenotypic intra-species variation in Candida albicans.

Genome Res 25:413–25.

25. du Preez JC, van Driessel B, Prior BA. 1989. D-xylose fermentation by

Candida shehatae and pichia stipitis at low dissolved oxygen levels in fed-

batch cultures. Biotechnol Lett 11:131–136.

26. du Preez JC, van Driessel B, Prior BA. 1989. Ethanol tolerance of Pichia

stipitis and Candida shehatae strains in fed-batch cultures at controlled low

dissolved oxygen levels. Appl Microbiol Biotechnol 30:53–58.

27. Sims REH, Mabee W, Saddler JN, Taylor M. 2010. An overview of second

generation biofuel technologies. Bioresour Technol 101:1570–1580.

28. Robak K, Balcerek M. 2018. Review of second generation bioethanol

production from residual biomass. Food Technol Biotechnol 56:174–187.

29. Suh SO, Marshall CJ, McHugh J V., Blackwell M. 2003. Wood ingestion by

passalid beetles in the presence of xylose-fermenting gut yeasts. Mol Ecol

12:3137–3145.

30. Melake T, Passoth V, Klinner U. 1996. Characterization of the genetic system

of the xylose-fermenting yeast Pichia stipitis. Curr Microbiol 33:237–242.

31. Jeffries TW, Grigoriev I V., Grimwood J, Laplaza JM, Aerts A, Salamov A,

Schmutz J, Lindquist E, Dehal P, Shapiro H, Jin YS, Passoth V, Richardson

PM. 2007. Genome sequence of the lignocellulose-bioconverting and xylose-

fermenting yeast Pichia stipitis. Nat Biotechnol 25:319–326.

32. Lynch DB, Logue ME, Butler G, Wolfe KH. 2010. Chromosomal G + C content

evolution in yeasts: Systematic interspecies differences, and GC-poor troughs

at centromeres. Genome Biol Evol 2:572–583.

33. Coughlan AY, Wolfe KH. 2019. The reported point centromeres of

Scheffersomyces stipitis are retrotransposon long terminal repeats. Yeast

36:275–283.

34. Villa-Carvajal M, Querol A, Belloch C. 2006. Identification of species in the

genus Pichia by restriction of the internal transcribed spacers (ITS1 and ITS2)

and the 5.8S ribosomal DNA gene. Antonie van Leeuwenhoek, Int J Gen Mol

Microbiol 90:171–181.

35. Schwartz DC, Cantor CR. 1984. Separation of yeast chromosome-sized DNAs

by pulsed field gradient gel electrophoresis. Cell 37:67–75.

36. Ketel C, Wang HSW, McClellan M, Bouchonville K, Selmecki A, Lahav T,

Gerami-Nejad M, Berman J. 2009. Neocentromeres form efficiently at multiple

possible loci in Candida albicans. PLoS Genet 5:e1000400.

37. Benson G. 1999. Tandem repeats finder: A program to analyze DNA

sequences. Nucleic Acids Res 27:573–580.

38. Letunic I, Bork P. 2018. 20 years of the SMART protein domain annotation

resource. Nucleic Acids Res 46:D493–D496.

39. Jeffries TW, Grigoriev I V, Grimwood J, Laplaza JM, Aerts A, Salamov A,

Schmutz J, Lindquist E, Dehal P, Shapiro H, Jin Y-S, Passoth V, Richardson

PM. 2007. Genome sequence of the lignocellulose-bioconverting and xylose-

fermenting yeast Pichia stipitis. Nat Biotechnol 25:319–326.

40. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview

Version 2--a multiple sequence alignment editor and analysis workbench.

Bioinformatics 25:1189–91.

41. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017.

Canu: Scalable and accurate long-read assembly via adaptive κ-mer weighting

and repeat separation. Genome Res 27:722–736.

42.   GitHub - ruanjue/smartdenovo: Ultra-fast de novo assembler using long noisy

reads.

43.   Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo

genome assembly from long uncorrected reads. Genome Res 27:737–746.

44.   Loman NJ, Quick J, Simpson JT. 2015. A complete bacterial genome

assembled de novo using only nanopore sequencing data. Nat Methods

12:733–735.

45.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo

CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: An Integrated Tool

for Comprehensive Microbial Variant Detection and Genome Assembly

Improvement. PLoS One 9:e112963.

46.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. 2015.

BUSCO: Assessing genome assembly and annotation completeness with

single-copy orthologs. Bioinformatics 31:3210–3212.

47.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: Quality

assessment tool for genome assemblies. Bioinformatics 29:1072–1075.

48.   Min B, Grigoriev I V, Choi I-G. 2017. FunGAP: Fungal Genome Annotation

Pipeline using evidence-based gene model evaluation. Bioinformatics

33:2936–2937.

49.   Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H,

Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A,

Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5:

Genome-scale protein function classification. Bioinformatics 30:1236–1240.

50.   Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018.

737   MUMmer4: A fast and versatile genome alignment system. PLOS Comput Biol

738   14:e1005944.

739   51.   Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones

740   SJ, Marra MA. 2009. Circos: An information aesthetic for comparative

741   genomics. Genome Res 19:1639–1645.

742   52.   Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole

743   genome comparisons dramatically improves orthogroup inference accuracy.

744   Genome Biol 16:157.

745   53.   Chen H, Boutros PC. 2011. VennDiagram: A package for the generation of

746   highly-customizable Venn and Euler diagrams in R. BMC Bioinformatics 12:35.

747   54.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-

748   Wheeler transform. Bioinformatics 25:1754–1760.

749   55.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A,

750   Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA. 2010. The genome

751   analysis toolkit: A MapReduce framework for analyzing next-generation DNA

752   sequencing data. Genome Res 20:1297–1303.

753   56.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA,

754   Handsaker RE, Lunter G, Marth GT, Sherry ST, McVean G, Durbin R. 2011.

755   The variant call format and VCFtools. Bioinformatics 27:2156–2158.

756   57.   Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X,

757   Ruden DM. 2012. A program for annotating and predicting the effects of single

758   nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila

759   melanogaster strain w1118; iso-2; iso-3. Fly (Austin) 6:80–92.

760   58.   Jackson AP, Gamble JA, Yeomans T, Moran GP, Saunders D, Harris D, Aslett

M, Barrell JF, Butler G, Citiulo F, Coleman DC, de Groot PWJ, Goodwin TJ, Quail MA, McQuillan J, Munro CA, Pain A, Poulter RT, Rajandream M-A, Renauld H, Spiering MJ, Tivey A, Gow NAR, Barrell B, Sullivan DJ, Berriman M. 2009. Comparative genomics of the fungal pathogens Candida dubliniensis and Candida albicans. Genome Res 19:2231–44.

59. Frank AC, Wolfe KH. 2009. Evolutionary capture of viral and plasmid DNA by yeast nuclear Chromosomes. Eukaryot Cell 8:1521–1531.

60. Butler G, Rasmussen MD, Lin MF, Santos M a S, Sakthikumar S, Munro C a, Rheinbay E, Grabherr M, Forche A, Reedy JL, Agrafioti I, Arnaud MB, Bates S, Brown AJP, Brunke S, Costanzo MC, Fitzpatrick D a, de Groot PWJ, Harris D, Hoyer LL, Hube B, Klis FM, Kodira C, Lennard N, Logue ME, Martin R, Neiman AM, Nikolaou E, Quail M a, Quinn J, Santos MC, Schmitzberger FF, Sherlock G, Shah P, Silverstein K a T, Skrzypek MS, Soll D, Staggs R, Stansfield I, Stumpf MPH, Sudbery PE, Srikantha T, Zeng Q, Berman J, Berriman M, Heitman J, Gow N a R, Lorenz MC, Birren BW, Kellis M, Cuomo C a. 2009. Evolution of pathogenicity and sexual reproduction in eight Candida genomes. Nature 459:657–62.

61. Slininger PJ, Shea-Andersh M a, Thompson SR, Dien BS, Kurtzman CP, Balan V, da Costa Sousa L, Uppugundla N, Dale BE, Cotta M a. 2015. Evolved strains of Scheffersomyces stipitis achieving high ethanol productivity on acid- and base-pretreated biomass hydrolyzate at high solids loading. Biotechnol Biofuels 8:1–27.

62. Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M, Izsvák Z, Levin HL, Macfarlan TS, Mager DL, Feschotte C. 2018. Ten things you should know about transposable elements 06 Biological

Sciences 0604 Genetics. Genome Biol 19:1–12.

63. Jansen MLA, Bracher JM, Papapetridis I, Verhoeven MD, de Bruijn H, de Waal

PP, van Maris AJA, Klaassen P, Pronk JT. 2017. Saccharomyces cerevisiae

strains for second-generation ethanol production: from academic exploration to

industrial implementation. FEMS Yeast Res. Oxford Academic.

64. Smith D, Quinlan A. 2008. Rapid whole-genome mutational profiling using

next-generation sequencing technologies. Genome Res 1638–1642.

65. Selmecki A, Forche A, Berman J. 2010. Genomic plasticity of the human

fungal pathogen Candida albicans. Eukaryot Cell 9:991–1008.

66. Fisher MC, Hawkins NJ, Sanglard D, Gurr SJ. 2018. Worldwide emergence of

resistance to antifungal drugs challenges human health and food security.

Science (80-) 360:739–742.

67. Freire-Benéitez V, Gourlay S, Berman J, Buscaino A. 2016. Sir2 regulates

stability of repetitive domains differentially in the human fungal pathogen

Candida albicans. Nucleic Acids Res https://doi.org/10.1093/nar/gkw594.

68. Jackson AP, Gamble J a, Yeomans T, Moran GP, Saunders D, Harris D, Aslett

M, Barrell JF, Butler G, Citiulo F, Coleman DC, de Groot PWJ, Goodwin TJ,

Quail M a, McQuillan J, Munro C a, Pain A, Poulter RT, Rajandream M-A,

Renauld H, Spiering MJ, Tivey A, Gow N a R, Barrell B, Sullivan DJ, Berriman

M. 2009. Comparative genomics of the fungal pathogens Candida dubliniensis

and Candida albicans. Genome Res 19:2231–44.

FIGURE LEGENDS

Figure 1. Classification of non-centromeric *S. stipitis* repeats

A) Schematics of non-centromeric retrotransposons identified in this study.

For each transposon, subclass, superfamily and family is indicated. The organisation

of coding and non-coding sequences of each transposon is displayed. **B)** Cladogram

showing CTG (Ser1)-clade species with known transposable elements (this study

and (60, 68). The presence (V) or absence (X) of a TE is indicated. **C)** Sequence

alignment of telomeric terminal repeats in members of the CTG (Ser1)-clade (*C.*

lusitaniae, *S. stipitis*, *L. elongisporus*, *C. albicans* and *C. tropicalis*) (this study and

(60, 68). Consensus sequence to the *S. cerevisiae* telomeric repeats is indicated

(Magenta box). **D)** Schematics of gene family members associated with *S. stipitis*

subtelomeres (30 Kb from chromosome end).

Figure 2. Phenotypic and Genotypic Diversity in *S. stipitis*

A) Heatmaps comparing growth rate (Left), maximum OD (Middle) and lag time

(Right) for each *S. stipitis* natural isolate in comparison to the reference Y-11545

strain (blue). Analyses were performed in Glucose (G), Xylose (X) and

Glucose/Xylose (G+X) media. The heatmap data are the average of 3 biological

replicates. **B)** Karyotyping of *S. stipits* natural isolates by CHEF electrophoresis. Y-

11545 strain is highlighted in blue and the size of its eight chromosome is indicated.

Figure 3. Differences in TE distribution and organisation

A) The genomic organisation of Y-11545 and Y-7124 is distinct .*Left:* Schematics of

Y-11545 chromosome organisation. Chromosome (Chr) number and size (Mbp) is

indicated. *Middle:* Karyotyping of *S. stipits* Y-11545 and Y-7124 strains by CHEF

electrophoresis. *Right:* Schematics of Y-7124 chromosome organisation.

Chromosome (Chr) number and size (Mbp) is indicated **B)** Schematics of non-

centromeric transposon family distribution in Y-11545 (*left*) and Y-7124 (*right*) **C)**

Copy Number of full-length transposons (Left) and transposon-associated repeats

associated with the Y-11545 (dark grey) and Y-7124 (light grey) genome. **D)**

Percentage (%) of Ancestral, Deletion and Insertion sites associated with the Y-

11545 and Y-7124 genomes

Figure 4. Chromosome rearrangements between *S. stipitis* natural isolates

A) Circos plot displaying macrosynteny between Y-11545 (Left) and Y-7124 (Right).

Chromosome (Chr) number and size is indicated. Recriprocal translocation between

the two genomes is highlighted in purple and pink. **B)** Schematics of repetitive

sequences associated with the translocation junction in the Y-11545 (Left) and Y-

7124 (Right) genomes. **C)** Subtelomeric gene families and TEs distribution in the Y-

11545 and Y-7124 genomes **D)** Schematics of centromere organisation in the Y-

11545 (Left) and Y-7124 (Right)

Figure 5 The *S. stipitis* genome is plastic following real-time evolution

A) Schematics of laboratory evolution strategy **B)** S. stipitis growth curve in SC G+X

liquid media at permissive (30 ºC) and restrictive (37 ºC) temperature **C)** Karyotype

organisation of *S. stipitis* colonies following 8 weeks of laboratory evolution at 30 ºC

and 37 ºC. Variation in the structure following laboratory evolution at 30 ºC (blue)

and 37 ºC (magenta) is indicated **D)** Serial dilution assay showing growth of parental

(P) and 37 ºC-evolved strains without (NO Rearrangements) and with

(Rearrangements) at 30 ºC and 37 ºC. The CHEF analysis strain number is indicated

(Magenta). * indicates colonies with a fitness advantage compared to the parental

strain.

Figure S1. Southern Blot analysis confirm the Chr5 / Chr7 translocation

Left: Schematics of chromosome 5 (Chr 5) and chromosome 7 (Chr 7) in Y-11545

(Left) and Y-7124 (Right). Reciprocal translocation is highlighted in purple and pink.

Southern Probe is indicated. Right: Southern Blot of Y-11545 and Y-7124

chromosomes separated by CHEF gel electrophoresis. Full chromosome profiling

(EtBr) and Southern Blot results (Southern) are indicated

Fig 1

Fig 2

A

B

Chromosome Organisation : CHEF

A

B

Key

Centromere   Subtelomere   *Ava*   *Bea*   *Caia*   *Ace*   *Bri*   *Can*

C

Full-length Retrotransposons

Transposon-derived Repeats

D

Fig 4

Fig 5