Poly (I:C)-induced maternal immune activation affects mouse visual discrimination performance and reversal learning in a sex-dependent manner.

Xin Zhao, Hieu Tran, Holly DeRosa, Ryland C. Roderick, Amanda C. Kentner*

School of Arts & Sciences, Health Psychology Program, Massachusetts College of Pharmacy and Health Sciences, Boston Massachusetts, United States 02115

Corresponding author:
Amanda Kentner
amanda.kentner@mcphs.edu
Office #617-274-3360
Fax # 617-732-2959

Keywords
Prenatal immune activation; Gestational; Poly (I:C); Poly I:C; Toll-like 3 receptors; Fetal programming; Touchscreen; Visual discrimination; Reversal learning; Hippocampus; Prefrontal cortex; Hypothalamus; Compensatory mechanisms; Acetylcholine; Environmental enrichment; Two-hit; Double hit; Behavioral flexibility; Cognitive flexibility; Prefrontal cortex; Hippocampus; Sex differences

Abstract

While there is a strong focus on the negative consequences of maternal immune activation (MIA) on the developing brain, very little attention is directed towards potential advantages of early life challenges. In this study we utilized a polyinosine-polycytidylic acid (poly(I:C)) MIA model to test visual discrimination (VD) and reversal learning (RL) in mice using touchscreen technology. Significant sex differences emerged in that MIA improved the latency for males to make a correct choice in the VD task while females reached criterion sooner, made fewer errors and utilized fewer correction trials in RL compared to saline-treated controls. These surprising improvements were accompanied by the sex-specific upregulation of several neural markers critical to cognitive functioning (e.g., Gabrg2, Grin1, Grin2b, Htr2a, Chrm1, Prkca, and Camk2a mRNA in prefrontal cortex (PFC)), indicative of compensatory plasticity in response to the MIA challenge. In contrast, when exposed to a "two-hit" stress model (MIA combined with loss of the social component of environmental enrichment (EE)), mice showed no evidence of anhedonia but required an increased number of PD and RL correction trials. These animals also had significant reductions of CamK2a mRNA in the PFC. Appropriate functioning of synaptic plasticity, via mediators such as this protein kinase and others, are critical for behavioral flexibility. Although EE has been implicated in delaying the appearance of symptoms associated with certain brain disorders, these findings are in line with evidence that it also makes individuals more vulnerable to its loss. Overall, with the right "dose", early life stress exposure can confer at least some functional advantages, which are lost when the number or magnitude of these exposures become too great.

Introduction

Numerous epidemiological studies have identified an association between inflammatory insults during pregnancy and the prevalence of neuropsychiatric disorders, such as autism and schizophrenia, in offspring (Babulas et al., 2006; Estes and McAllister, 2016; Sørensen et al., 2008). However, at the population level, only a subset of children exposed to prenatal infections develop disease-relevant behavioral abnormalities (Mahic et al., 2017; Meyer, 2019).This suggests the etiology of these neurodevelopmental disorders may involve a synergism of prenatal immune challenge and other environmental risk factors (e.g. stress) in later life. Adding to this complexity, the effects of prenatal immune insults on child health outcomes also vary depending biological sex (Gilman et al., 2016; Goldstein et al., 2014; Mac Giollabhui et al., 2019). Hence, it is essential to determine sources that contribute to the heterogeneous outcomes of MIA and to identify factors for building neurodevelopmental resilience and susceptibility to this early-life adversity.

In preclinical research, rodent maternal immune activation (MIA) models have been employed to mimic the effects of gestational infections and to investigate its underlying biological mechanisms (Estes and McAllister, 2016; Kentner et al., 2019a; Knuesel et al., 2014). Specifically, a mid-gestational injection of polyinosinic:polycytidylic acid (poly (I:C)), a viral mimetic toll-like receptor 3 agonist, can induce an extensive collection of innate immune responses (Mueller et al., 2019) and lead to a wide array of abnormalities in neurophysiology, behavior and cognitive abilities (see (Haddad et al., 2020) for review). Whereas the vast majority of previous studies have addressed the adverse effects of MIA, prior research has also revealed that MIA-treated offspring can exhibit improvements in cognitive functioning (Makinson et al., 2019; Nakamura et al., 2021), blunted responses to a second immune challenge in adolescence (Clark et al., 2019), and a higher level of resilience to the disruptive effects of isolation rearing on behavior and neurophysiology (Goh et al., 2020). Recently, it has been shown that poly (I:C)-induced MIA at a specific window of

parvalbumin interneuron development can lead to improvements in spatial working memory (Nakamura et al., 2021). These findings indicate that MIA may act like other mild early-life stressors to facilitate the development of protection against stressors in later life (Fujioka et al., 2001; Cannizzaro et al., 2006; Li et al., 2013). However, our understanding of the adaptations and potential priming effects of MIA on resiliency are relatively underexplored. Notably, the 'hidden talents' approach has provided a framework for investigating how harsh and unpredictable childhood environments may promote changes in behaviors and cognition that are adaptive for future harsh environment in humans (Ellis et al., 2020), which may explain some of these seemingly paradoxical observations in our animal studies.

The complexity of the living environment affects brain development at the functional, anatomical, and molecular level (Kempermann, 2019; Ohline and Abraham, 2019; Zhang et al., 2018). Increased complexity of the laboratory animal home cage, so-called environmental enrichment (EE), is characterized by exposure to environments with rich social, motor, cognitive and sensory stimulation. Although EE has been implicated in delaying the appearance of symptoms associated with certain brain disorders (Chourbaji et al., 2011; Herring et al., 2009; Van Dellen et al., 2000), it also makes individuals that are exposed to EE more vulnerable to its loss. Indeed, when the environment changes from being enriched to more impoverished (EE removal) behavioral alterations linked to depressive symptomology emerge (Morano et al., 2019; Smith et al., 2017). Moreover, these behavioral changes are accompanied by dysregulation of the hypothalamus-pituitary-adrenal axis activity in female rats (Morano et al., 2019). EE removal may therefore be considered as a second hit factor which could potentially reinforce the disruptions induced by the first hit (e.g., MIA) in early-life and ultimately lead to onset of a full clinical syndrome.

In the present study, we aim to examine how MIA interacts with the loss of the social component of life-long EE, influencing neuropathology and cognitive functioning of both male and female offspring. To examine the neuropathological alterations, we focused on the prefrontal cortex (PFC), an important brain region for cognitive control (Miller and Cohen, 2001). In this region, we measured neurochemical alterations in the PFC, analyzed as mRNA expression of neural markers associated with different neurotransmitter pathways (e.g., monoaminergic, glutamatergic, and cholinergic).

2. Materials and Methods

2.1. Animals

Male and female C57BL/6J mice arrived from the Jackson Laboratory (Bar Harbor, ME), and housed at 20°C on a 12 h light/dark cycle (0700–1900 light) with ad libitum access to food and water. **Figure 1** details the timeline of experimental procedures. Female mice were housed as pairs in either larger environmental enrichment cages (EE; N40HT mouse cage, Ancare, Bellmore, NY), with access to toys, tubes, a Nylabone, Nestlets® (Ancare, Bellmore, NY) and Bed-r' Nest® (ScottPharma Solutions, Marlborough MA), or standard sized cages (SD; N10HT mouse cage, Ancare, Bellmore, NY) with Nestlets® and a Nylabone only. Males were paired in SD conditions until breeding. At that point, male animals were housed with two EE or SD females. Dams received either a 20 mg/kg intraperitoneal injection of polyinosine-polycytidylic acid (poly (I:C); tlrl-picw, lot number PIW-41-03, InvivoGen), or vehicle (sterile pyrogen-free 0.9% NaCl) on the morning of gestational day (G)12. Offspring were weaned into same-sex groups on P21 and maintained in their housing assignments (SD = 2-3 animals/cage; EE = 4-5 animals/cage) until P90. Additional methodological details, including the validation of poly (I:C), can be found in the reporting table from Kentner et al. (2019), provided as **Supplementary Table 1**. The MCPHS University Institutional Care and Use Committee approved all procedures described, which were carried out in compliance

with the recommendations outlined by the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health.

2.3. Touchscreen Learning

Impairments in the cognitive functions were evaluated by leveraging a touchscreen-based operant task paradigm. Specifically, pair-wise discrimination (PD) and reversal learning (RL) tasks have been employed to assess impairments in motivation, memory, rule learning, and cognitive flexibility (Bryce and Howland, 2015; Bussey et al., 2012; Horner et al., 2013; Lins et al., 2018). Moreover, the cognitive tasks were conducted following social isolation, which has been implicated in mediating a range of behavioral deficits in adulthood including sensorimotor processing, social interaction abnormalities, heightened anxiety and cognitive dysfunction (Bakshi and Geyer, 1999; Fone and Porkess, 2008). By integrating this component, this study will shed light on the mechanisms underlying the variation in the neurodevelopmental resilience and susceptibility to a 'second-hit' stressor in later life. On P90, animals were individually housed in their SD or EE conditions and food restricted (at 85-90% of their free-feeding weight which was maintained until the end of the study) to facilitate touchscreen responding for a milkshake reward (Strawberry Nesquik®). Prior to touchscreen training, animals were habituated to the liquid food reward, placed on a plastic dish, in their home cages. Eight touchscreen sound-attenuated operant chambers (Campden Instruments Ltd, UK) were used to run the visual discrimination and reversal tasks. Details on the apparatus and dimensions have been reported previously (Horner et al., 2013). Training and testing were completed using the ABET II software.

Eight touchscreen-equipped operant conditioning chambers for mice (Bussey-Saksida Touch Systems, Lafayette Instrument Company) in sound attenuating cubicles were used. Mice were first habituated to the chambers then trained to a) recognize the sound and light cues, b) interact physically with the touchscreen, and c) collect the liquid rewards following the manufacturer's suggested protocol, and as previously outlined by Lins et al (2018). Chambers were thoroughly cleaned with Quat TB, and touchscreens with 10% ethanol alcohol (to maintain touch sensitivity), between each animal and session.

Visual discrimination: In this task, novel black and white images (the classic 'fan' and 'marbles') were simultaneously presented to mice (**Figure 1**). Left and right placement of the visual stimuli were presented in a pseudo randomized manner. For each animal, one of the two images was always correct (S+), regardless of its left vs right placement. When the animal responded to this S+ image, they received a continuously reinforced (100% probability of reward) delivery of ~7 ul of liquid milkshake. If the mouse responded to the incorrect (S-) image, no reward was produced and the animal had a 5-second 'time out' period, followed by a correction trial. Once the animal correctly responded to the correction trial (where the two images were presented in the same spatial configuration as the previous trial and the animal had responded incorrectly), the next trial of the session commenced. The designation of the correct stimulus (fan vs marble) was counterbalanced across mice. Criterion was reached when the animal completed 30 trials in 1 hour (excluding correction trials) per daily session with >85% correct for two consecutive days. If an animal could not meet criterion after 30 days, they did not progress onto the next task (Radke et al., 2019; Kenton et al., 2020).

Reversal learning: Reversal learning was evaluated after successful completion of the visual discrimination task. In reversal learning the previously correct image (S+) becomes the incorrect visual stimuli (S-) while the previously incorrect image (S-) becomes the correct choice (S+) that would result in a liquid reward delivery. Criterion was reached when animals reached 30 trials in 60 minutes with >85% correct for two consecutive days or were tested for 10 days.

2.4. Sucrose Preference Test

We utilized the 16-hr overnight sucrose preference test (Connors et al., 2014; Kentner et al., 2010). Here, male and female animals were given two bottles, each containing either a 1% sucrose solution or water. Sucrose preference was determined by calculating the ratio of sucrose intake (grams) to total fluid intake (grams) and converted into a percent score. The placement of the sucrose and water bottles was counterbalanced between trials to prevent side preferences. At no time were animals deprived of food or water.

2.5. Tissue collection and analysis

To circumvent the confound of touchscreen training modifying the brain, we collected tissue from littermates matched to our touchscreen animals on P85 (Group 1). The animals that underwent touchscreen training were Group 2. This allows for a better assessment of the role of poly (I:C) on the brain in our data interpretation. A mixture of Ketamine/Xylazine (150 mg/kg, i.p/15 mg/kg, i.p) was used to anesthetize animals. Animals were perfused intracardially with a chilled phosphate buffer solution. Prefrontal cortex (PFC) and ventral hippocampus were dissected from the left hemisphere, frozen on dry ice and stored at −75°C until processing. The right hemisphere was post-fixed in a 4% paraformaldehyde, phosphate buffer 0.1M solution (BM-698, Boston BioProducts) overnight at 4°C. Tissue was submerged in ice cold 10% sucrose in PBS (with 0.1% sodium azide) and incubated at 4°C overnight. The next day, solution was replaced with 30% sucrose in PBS for 3 days. Tissue was then flash frozen with 2-methylbutane (O3551-4, Fisher Scientific) and stored at −75°C until sectioning.

2.6. RT-PCR

Using the RNeasy Lipid Tissue Mini Kit (Qiagen, 74804), total RNA was extracted from frozen tissue and resuspended in RNase-free water. A NanoDrop 2000 spectrophotometer (ThermoFisher Scientific) was used to quantify isolated RNA. Total RNA was reverse transcribed to cDNA with the Transcriptor High Fidelity cDNA Synthesis Kit (#5081963001, Millipore Sigma) using the manufacturers protocol, and the final cDNA solution was stored at -20°C until analysis. Quantitative real-time PCR with Taqman™ Fast Advanced Master Mix (#4444963, Applied Biosystems) was used to measure the mRNA expression of Gamma-Aminobutyric Acid Type A Receptor Subunit Gamma2 (Gabrg2; Mm00433489_m1), glutamate receptor, ionotropic, NMDA1 (Grin1; Mm00433800_m1), glutamate receptor, ionotropic, NMDA2A (Grin2a; Mm00433802_m1), glutamate receptor, ionotropic, NMDA2B (Grin2b; Mm00433820_m1), solute carrier family 18 (vesicular monoamine), member 3/VAChT (Slc18a3; Mm00491465_s1), 5-hydroxytryptamine (serotonin) receptor 2A (Htr2a; Mm00555764_m1), dopamine receptor D1 (Drd1; Mm02620146_s1), dopamine receptor D2 (Drd2; Mm00438545_m1), cholinergic receptor, muscarinic 1, CNS (Chrm1; Mm00432509_s1), cholinergic receptor, muscarinic 4 (Chrm4; Mm00432514_s1), choline acetyltransferase (Chat; Mm01221880_m1), calcium/calmodulin dependent protein kinase II alpha (Camk2a, Mm00437967_m1), and protein kinase C alpha (Prkca, Mm00440858_m1). Reactions were analyzed in triplicate using optical 96-well plates (Applied Biosystems StepOnePlus™ Real-Time PCR System) and relative gene expression levels were evaluated using the $2^{-\Delta\Delta^{CT}}$ method with 18S (Hs99999901_s1). We selected this housekeeping gene as it was not affected by MIA or housing. Gene expression was normalized in relation to 18S and presented as mean expression relative to same sex SD-saline treated controls.

2.7. Statistical analyses

The statistical software package Statistical Software for the Social Sciences (SPSS) version 26.0 (IBM, Armonk, NY) was used for all analyses, except for the survival curves which were evaluated with the Log-rank (Mantel Cox test) using GraphPad Prism (Version 9.0). Two or three-way ANOVAs were used as appropriate. In cases of significantly skewed data (Shapiro-Wilks), Kruskal-Wallis tests (non-parametric equivalent to ANOVA, reported as X^2) were applied. Body weight was used as a covariate for sucrose preference data. LSD post hocs were applied, except where there were fewer than three levels in which case pairwise t-tests and Levene's were utilized. Multiple testing procedures for the gene expression data were analyzed using the False Discovery Rate (FDR). Partial eta-squared ($n_p{}^2$) is also reported as an index of effect size (the range of values being 0.02 = small effect, 0.13 = moderate effect, 0.26 = large effect; Miles and Shevlin, 2001).

3. Results

3.1. Behavioral tests

3.1.1 Touchscreen

The overall Log-rank (Mantel-Cox) test was significant across all groups, suggesting a difference in session-to-criterion survival curves (X^2 (7) = 19.82, p = 0.006; **Figure 2A**). Post hoc tests failed to identify differences across MIA, housing, or sex using adjusted alpha values that accounted for the number of comparisons made (p>0.01). However, ANOVA revealed a main effect of sex on number of sessions required to reach criterion in the discrimination task (F(1, 67) = 12.086, p = 0.001, np^2 = 0.153; male: 9.94±0.82 vs female: 15.9±1.38).

The total number of discrimination trials and errors did not differ across groups (p>0.05; **Figure 2B**). A MIA by housing interaction (F(1,67) = 4.028, p = 0.049, np^2 = 0.057) revealed that EE-poly (I:C) mice required more correction trials compared to their EE-saline (t(37) = -3.416, p = 0.002) and SD-poly IC counterparts (t(34) = -2.788, p = 0.009). This suggests that the combination of multiple early life stressors (MIA + companion loss) compounded impairments on this measure (**Figure 2B**). Latency to collect rewards and make incorrect responses was not different as a function of sex, MIA, or housing (p>0.05; **Figure 2C**). However, there were significant sex differences in the average latency to make a correct response in the visual discrimination task (X^2 (1) = 5.311, p = 0.021). Specifically, male treated MIA (X^2 (1) = 6.501, p = 0.011; poly (I:C): 4.89 ± 2.78 vs SD: 7.84 ± 0.92) mice were faster than females, and female EE ((X^2 (1) = 4.116, p = 0.042; SD: 7.14 ± 0.84 vs EE: 8.89 ± 0.704) mice were slower than males, to respond when making a correct choice (**Figure 2C**). In alignment with other research suggesting that females thrive better in social environments, this suggests that female EE mice were more sensitive to the loss of companionship in their homepage.

For reversal learning, there were no group differences in session-to-criterion survival curves (Long-rank test, X^2 (7) = 10.26, p = 0.17 (N.S); **Figure 3A**) or in the number of sessions to reach reversal criterion (p>0.05). The total number of trials completed were lower for female poly (I:C) animals relative to their saline counterparts (sex x MIA: F(1, 56) = 14.064, p = 0.001, np^2 = 0.201; female poly (I:C) vs female saline (p = 0.002); **Figure 3B**). Additionally, female poly (I:C) mice made fewer errors compared to female saline animals (sex x MIA: F(1, 56) = 11.24, p = 0.001, np^2 = 0.167; female poly (I:C) vs female saline (p = 0.005); **Figure 3B**) suggesting that early life challenges may not only lead to impairments, but may be associated with some performance benefits. A significant sex x housing x MIA interaction (F(1, 57) = 7.048, p = 0.010, np^2 = 0.110) showed that male (p = 0.027) and female (p = 0.0001) SD-poly (I:C) animals required fewer correction trials than same-sex EE-poly (I:C)

mice. This is in line with evidence that multiple hits can result in cognitive deficits that are not apparent with only one stressor exposure (Bilbo et al., 2005). Moreover, female SD-poly (I:C) mice had fewer correction trials than SD-saline animals (p = 0.0001; **Figure 3B**), again highlighting the potential for early life adversity to confer functional advantages. There were no differences in average latency to collect rewards or to make incorrect and correct responses on the reversal trials (p >0.05; **Figure 3C**).

3.1.2. Sucrose Preference Tests

To determine whether EE-poly (I:C) animals had motivational, rather than cognitive, impairments associated with the multiple stressor experience (MIA + social companion loss) we conducted a 16-hour sucrose preference test. We did not observe anhedonia using this measure (p>0.05). Moreover, animals were not different in their latency to collect the milkshake rewards (**Figure 2C; Figure 3C**). Therefore, we did not find motivational deficits in these animals that would interfere with their touchscreen responding.

3.2. Taq Man qPCR
3.2.1. GABA and Glutamatergic Receptor Expression

Changes in excitatory/inhibitory balance within the PFC, mediated by GABA and glutamate signaling, can impair behavioral flexibility (Bissonette & Powell, 2012; Enomoto et al., 2011). Here, we observed that levels of Gabrg2 in the PFC were elevated in poly (IC) treated ($F(1, 28) = 5.007$, $p = 0.033$, $np^2 = 0.152$; poly (I:C): 1.10 ± 1.09 vs saline: 0.85 ± 0.07) and standard housed mice (males: $F(1, 28) = 10.768$, $p = 0.003$, $np^2 = 0.278$; SD: 1.16 ± 0.09 vs EE: 0.79 ± 0.09; females: $F(1, 28) = 9.606$, $p = 0.004$, $np^2 = 0.255$; SD: 1.09 ± 0.05 vs EE: 0.82 ± 0.07; **Supplementary Table 1**). Grin1 expression was increased in female SD-poly (IC) animals compared to SD-saline (p = 0.004) and EE-poly (IC) mice (p = 0.003; housing x MIA interaction: $F(1, 28) = 13.250$, $p = 0.001$, $np^2 = 0.321$; **Figure 4A**). There were no housing or MIA effects on Grin2a (p>0.05; **Supplementary Table 1**), however PFC levels of Grin2b were elevated in male poly (IC) mice ($F(1, 28) = 21.745$, $p = 0.0001$, $np^2 = 0.437$; poly (IC): 1.28 ± 0.52 vs saline: 0.95 ± 0.04; **Figure 4B**). Grin2b expression in PFC could be an indication of region-specific compensatory plasticity following MIA, since decreased levels of this receptor are reported in other brain areas and with other immunogens (Connors et al., 2014). Hippocampal levels of each of these genes can be found in **Table 1**.

3.2.2. Markers of Monoamine and Cholinergic Functioning

A variety of receptor mechanisms involved in monoaminergic and cholinergic signaling are critical for PFC mediated cognitive functioning (Hupalo et al., 2019; Waltz, 2017). While male SD mice had higher Drd1 expression levels compared to EE animals (p = 0.017; main effect of housing: $F(1, 28) = 6.458$, $p = 0.017$, $np^2 = 0.187$; SD: 1.109 ± 0.07 vs EE: 0.85 ± 0.07; **Supplementary Table 1**) in the PFC, there were no significant group effects of Drd2 (p>0.05; this and hippocampal dopamine receptor genes are provided in **Supplementary Table 1**). In contrast, Htr2a levels in PFC were elevated in poly (I:C) males compared to saline (p = 0.007; main effect of MIA: $F(1, 28) = 8.484$, $p = 0.007$, $np^2 = 0.233$; poly (I:C): 1.304 ± 0.12 vs saline: 0.90 ± 0.07; **Figure 4C**). SD-poly (I:C) was also associated with increased Htr2a in the PFC of female mice compared to SD-saline (p = 0.001) and EE-poly (I:C) females (p = 0.001; MIA x housing interaction: $F(1, 28) = 14.615$, $p = 0.001$, $np^2 = 0.343$; **Figure 4C**). The contribution of the 5HT2a receptor to cognition is complicated by conflicting reports of its antagonism both enhancing (Amodeo et al., 2014, Baker et al., 2011) and inhibiting behavioral flexibility (Boulougouris et al., 2008), and of its activation impairing probabilistic reversal learning (Amodeo et al., 2020). This suggests that the MIA-induced upregulation of PFC Ht2a in the current study should be associated with a cognitive

impairment. Instead, we observed MIA mice to have a general improved performance in PD and RL touchscreen tasks. A critical difference in our work could be that the elevated receptor level is likely reflective of a chronic, rather than acute, change. It is also possible that pharmacological 5Ht2a receptor activation may improve behavioral flexibility at different doses and more targeted central administration routes. There were no group effects (p>0.05) for the vesicular acetylcholine transporter Slc18a3 (**Figure 4D**) or choline acetyltransferase mRNA (Chat; **Figure 4E**) in the PFC.

The muscarinic receptors Chrm1 and Chrm4 have both been implicated in the pathogenesis of schizophrenia (Gibbons & Dean, 2016). Given that poly (I:C) induced MIA in rodents is considered a model for this neurodevelopmental disorder (Meyer, 2019), and the role of these receptors in cognitive functioning, we were interested in their expression in this model. Chrm4 was significantly reduced in EE-saline vs EE-poly (I:C) male, but not female, mice in the PFC (p = 0.0001; MIA x housing interaction: $F(1, 28) = 4.909$, p = 0.035, $np^2 = 0.149$; **Supplementary Table 1**). While Chrm1 was elevated in female EE compared to SD mice (p = 0.001; main effect housing: $F(1, 28) = 14.893$, p = 0.001, $np^2 = 0.347$; EE:1.37±0.08 vs SD: 1.05±0.03; **Figure 4F**), in males there was a significant main effect of MIA ($F(1, 28) = 16.688$, p = 0.0001, $np^2 = 0.373$). Here, male poly (I:C) had elevated PFC levels of this muscarinic receptor compared to saline (p =0.0001; poly (I:C): 1.40±0.06 vs saline: 1.07±0.05; **Figure 4F**). Chrm1 is thought to be important for synaptic plasticity (Gibbons & Dean, 2016) and treatment with selective agonists for this receptor have resulted in improvements in performance in spatial memory tasks (Vanover., et al., 2008), reversal learning (Shirey et al., 2009), and novel object recognition (Bradley et al., 2010). Our data of elevated PFC Chrm1 expression, alongside improved behavioral flexibility, in MIA males are in line with these findings.

3.2.3. Synaptic Plasticity Markers

Appropriate functioning of synaptic plasticity in the PFC, via mediators such as CaMKII and other protein kinases, are critical for behavioral flexibility (**Ma et al., 2015; Natividad et al., 2015**). In males, both Camk2a (**Figure 4G**) and Prkca (**Figure 4H**) expression levels were elevated in poly (I:C) compared to saline treated mice (main effect of MIA; CamK2a: $F(1, 28) = 9.320$, p = 0.005, $np^2 = 0.250$; poly (I:C): 1.44±0.07 vs saline: 1.17±0.07; Prkca: $F(1, 28) = 27.233$, p = 0.0001, $np^2 = 0.493$; poly (I:C): 1.75±0.13vs saline: 0.99±0.06). SD-poly (I:C) females had heightened expression of Camk2a (MIA x housing interaction: $F(1, 28) = 43.633$, p = 0.001, $np^2 = 0.609$) and Prkca ($F(1, 28) = 14.416$, p = 0.005, $np^2 = 0.340$) compared to SD-saline (Camk2a: p= 0.001; Pkrca: p = p = 0.002) and EE-poly (I:C) animals (Camk2a: p = 0.0001; Prkca: p = 0.003). EE-saline animals also had increased levels of Camk2a (p = 0.001) compared to EE-poly (I:C) animals, suggesting that multiple stressor hits may inhibit some of the advantages conferred by stressful experiences.

Discussion

In the current study, we demonstrate that poly (I:C)-induced MIA imposes sex-specific enhancements on a variety of learning motifs. This runs in parallel with recent findings that gestationally treated poly (I:C) mice had enhanced spatial working memory (Nakamura et al., 2021) and LPS-induced MIA improved cognitive performance in the 5-choice serial reaction task (Makinson et al., 2019). At first glance, these findings stand in contrast to previous studies reporting impaired performance in working memory, executive function and cognitive flexibility following MIA (Amodeo et al., 2019; Lins et al., 2018; Meehan et al., 2017; Meyer et al., 2010; Murray et al., 2017; Wallace et al., 2014). However, it should be noted that several factors such as the immune stimulus intensity and the timing of challenge may contribute to the discrepancies between these findings. For example, compared to earlier poly

(I:C) exposure points, MIA applied at later gestational stages (e.g. GD 14.5 and 17.5) had a bigger influence on schizophrenia-related behaviors (Meehan et al., 2017). Our results suggest that MIA during mid-gestation may have an enhancement effect on certain aspects of cognitive functioning. This also echoes the notion of 'hidden talents' (Ellis et al., 2020) and suggests that improved cognitive abilities may act as an adaptive mechanism, preparing individuals to cope with future harsh environments.

Sex differences in the cognitive abilities have been well studied in both humans and animal models (review see (Hyde, 2016; Jonasson, 2005) and we found in the current study that males required fewer sessions than females to reach criterion in the PD task. Generally, touchscreen-based PD and RL tasks are used to assess impairments in motivation, memory, rule learning, and cognitive flexibility (Bryce and Howland, 2015; Bussey et al., 2012; Horner et al., 2013; Lins et al., 2018) and the sex difference we observed may reflect a small male advantage in some aspects of these abilities (Berger-Sweeney et al., 1995; Jonasson, 2005). That said, we found female MIA SD mice required fewer correction trials than their SD-saline controls in the RL task. This is consistent with a previous study showing that in the 5-choice serial reaction time task, LPS-treated females learned the task more quickly and displayed a higher level of motivation to earn a reinforcer (decreased time required to reach criterion) (Makinson et al., 2019). These sex difference could also be indicative of the different strategies used by male and female animals that lead to different performance outcomes (Chen et al., 2021).

Long-term EE exposure followed by the loss of social enrichment dampened the enhancing effects of MIA. For example, male and female SD-poly (I:C) animals required fewer correction trials than same-sex EE-poly (I:C) mice. In the current study, the cognitive tasks were conducted following social isolation, which mediated a range of behavioral deficits in adulthood including sensorimotor processing, social interaction abnormalities, heightened anxiety, and cognitive dysfunction (Bakshi and Geyer, 1999; Fone and Porkess, 2008). Although both SD and EE animals were subjected to the deprivation of the social component, social isolation impacted the cognitive performance in EE mice more negatively. This is consistent with data showing that loss of enrichment changes the behavioral and physiological phenotypes of animals that had been exposed to long-term EE (Morano et al., 2019). Specifically, single housing following prolonged EE exposure was associated with increased weight gain, elevated helplessness and passive coping behaviors, and blunted hypothalamic-pituitary-adrenocortical activity (Morano et al., 2019; Smith et al., 2017). In contrast, isolation rearing following MIA in SD-housed animals was protective against negative impacts on behavior and neurophysiology (Goh et al., 2020).

At the neurophysiological level, the mechanisms underlying the enhancing effects of MIA remain to be elucidated. Our current results suggest that protein kinase C (PKC) signaling in the PFC may be a potential pathway for mediating these effects. PKC isoforms are critically involved in many types of learning and memory (Alkon et al., 2007; Nelson et al., 2008; Sun and Alkon, 2010). Moreover, PKC can be activated by various synaptic inputs and intracellular signals that are involved in modulating cognitive functions, including glutamatergic (Hasham et al., 1997), cholinergic (Chen et al., 2005; Xiong et al., 2019) and serotonergic inputs (Carr et al., 2002; Carr et al., 2003). Therefore, the enhancing effects of MIA on cognition may be mediated through these inputs on the PKC signaling pathway. This is supported by our data showing upregulated *Prcka*, *Grin2b*, *Chrm1* and *Htr2a* in the poly (I:C)-treated male mice.

Another important enzyme, calcium/calmodulin-dependent protein kinase type II subunit alpha (CamK2a), interacts with NMDA subunits and together have a critical role in plasticity and learning (Ma et al., 2015; Takahashi et al., 2009). Given the parallel increase of these markers in our poly (I:C) animals (Group 1), we originally predicted that this upregulation

was a strategy enacted by the brain in an attempt to compensate for the early life inflammatory challenge, but that we would still observe functional deficits in touchscreen performance (Group 2 animals). Instead, the MIA animals surprised us by demonstrating sex-specific improvements in their PD and RL tasks.

As a complement to previous studies on MIA's impacts, the current study reveals, and characterizes, additional enhancing effects of MIA on cognitive functioning. We also demonstrate that long-term EE exposure may dampen the enhancing effects of MIA when there is a loss of the enrichment in later life, providing additional support for the detrimental effects of multiple hits. Our results shed light on the variable effects of MIA in the etiology of neurodevelopmental disorders and may help facilitate potential therapeutic or preventive strategies by taking these variables into consideration.

Funding and Disclosure
This project was funded by NIMH under Award Number R15MH114035 (to ACK), a MCPHS Center for Undergraduate Research Mini-Grant (to HT), and a Summer Undergraduate Research Fellowship (to R.C.R). The authors wish to thank Drs. Johnny Kenton and Jared Young for their helpful advice on the touchscreen tasks. The authors would also like to thank the MCPHS University School of Pharmacy and School of Arts & Sciences for their continual support. The content is solely the responsibility of the authors and does not necessarily represent the official views of any of the financial supporters.

Author Contributions
X.Z., H.T., H.D., & R.C.R., ran the experiments; X.Z., & A.C.K. analyzed and interpreted the data, and wrote the manuscript; A.C.K., H.T., & H.D made the figures. A.C.K. designed and supervised the study.

Declaration of Competing Interest
The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Figure Captions

Figure 1. Outline of experimental procedures.

Figure 2. Paired visual discrimination learning in male (left) and female (right) MIA offspring. A) Survival analysis of sessions to discrimination criterion. B) Total number of trials, errors, and correction trials completed during all discrimination sessions. C) Latency (seconds) to respond for a correct choice, incorrect choice, and reward during all discrimination sessions. Data expressed as mean $\pm$ SEM, n=8-14 litters represented per sex, MIA, and housing group. *$p < 0.05$, **$p < 0.01$, versus SD-saline; #$p < 0.05$, ##$p < 0.01$, versus EE-poly (I:C); [a]$p < 0.05$, [aa]$p < 0.01$, main effect of housing; [b]$p < 0.05$, [bb]$p < 0.01$, main effect of MIA.

Figure 3. Reversal learning in male (left) and female (right) MIA offspring. A) Survival analysis of sessions to reversal criterion. B) Total number of trials, errors, and correction trials completed during all reversal sessions. C) Latency (seconds) to respond for a correct choice, incorrect choice, and reward during all reversal sessions. Data expressed as mean $\pm$ SEM, n=7-11 litters represented per sex, MIA, and housing group. *$p < 0.05$, **$p < 0.01$,

versus SD-saline; #p < 0.05, ##p < 0.01, versus EE-poly (I:C); [a]p < 0.05, [aa]p < 0.01, main effect of housing; [b]p < 0.05, [bb]p < 0.01, main effect of MIA.

Figure 4. Prefrontal cortex gene expression in male (left) and female (right) MIA offspring on postnatal day 85. Levels of A) Grin1, B) Grin2b, C) Htr2a, D) Slc18a3 E) Chat, F) Chrm1, G) Chrm4, and H) Prkca mRNA. Gene expression data are expressed as mean ± SEM, n=8 litters represented per sex, MIA, and housing group. *p < 0.05, **p < 0.01, versus SD-saline; #p < 0.05, ##p < 0.01, versus EE-poly (I:C).

References

Alkon, D.L., Sun, M.-K., Nelson, T.J., 2007. PKC signaling deficits: a mechanistic hypothesis for the origins of Alzheimer's disease. Trends Pharmacol. Sci. 28, 51-60.

Amodeo, D.A., Hassan, O., Klein, L., Halberstadt, A.L., Powell, S.B., 2020. Acute serotonin 2A receptor activation impairs behavioral flexibility in mice. Behavioural Brain Research. 395, 112861.

Amodeo, D.A., Jones, J.H., Sweeney, J.A., Ragozzino, M.E., 2014. Risperidone and the 5□HT2A Receptor Antagonist M 100907 Improve Probabilistic Reversal Learning in BTBR T+ tf/J Mice, Autism Research 7, 555–567.

Babulas, V., Factor-Litvak, P., Goetz, R., Schaefer, C.A., Brown, A.S., 2006. Prenatal exposure to maternal genital and reproductive infections and adult schizophrenia. Am. J. Psychiatry 163, 927-929.

Baker, P.M., Thompson, J.L., Sweeney, J.A., Ragozzino, M.E., 2011. Differential effects of 5-HT2A and 5-HT2C receptor blockade on strategy-switching, Behavioural Brain Research 219, 123–131.

Bakshi, V.P., Geyer, M.A., 1999. Ontogeny of isolation rearing-induced deficits in sensorimotor gating in rats. Physiol. Behav. 67, 385-392.
Bayer, T.A., Falkai, P., Maier, W., 1999. Genetic and non-genetic vulnerability factors in schizophrenia: the basis of the" two hit hypothesis.". Journal of psychiatric research.

Berger-Sweeney, J., Arnold, A., Gabeau, D., Mills, J., 1995. Sex differences in learning and memory in mice: effects of sequence of testing and cholinergic blockade. Behav. Neurosci. 109, 859.

Bissonette, G.B., Powell, E.M., 2012. Reversal learning and attentional set-shifting in mice. Neurophamacology, 62, 1168-1174.

Bilbo, S.D., Levkoff, L.H., Mahoney, J.H., Watkins, L.R., Rudy, J.W., Maier, S.F., 2005b. Neonatal infection induces memory impairments following an immune challenge in adulthood. Behav. Neurosci. 119, 293–301.

Boulougouris, V., Glennon, J.C., Robbins, T.W., 2008. Dissociable effects of selective 5-HT 2A and 5-HT 2C receptor antagonists on serial spatial reversal learning in rats,

Neuropsychopharmacology, 33, 2007–2019.

Bradley, S.R., Lameh, J., Ohrmund, L., Son, T., Bajpai, A., Nguyen, D., Friberg, M., Burstein,
E.S., Spalding, T.A., Ott, T.R., Schiffer, H.H., Tabatabaei, A., McFarland, K., Davis, R.E., Bonhaus, D.W., 2010. AC-260584, an orally bioavailable M1 muscarinic receptor allosteric agonist, improves cognitive performance in an animal model. Neuropharmacology 58, 365–373.

Bryce, C.A., Howland, J.G., 2015. Stress facilitates late reversal learning using a touchscreen-based visual discrimination procedure in male Long Evans rats. Behav. Brain Res. 278, 21-28.

Bussey, T., Holmes, A., Lyon, L., Mar, A., McAllister, K., Nithianantharajah, J., Oomen, C., Saksida, L., 2012. New translational assays for preclinical modelling of cognition in schizophrenia: the touchscreen testing method for mice and rats. Neuropharmacology 62, 1191-1203.

Carr, D.B., Day, M., Cantrell, A.R., Held, J., Scheuer, T., Catterall, W.A., Surmeier, D.J., 2003. Transmitter modulation of slow, activity-dependent alterations in sodium channel availability endows neurons with a novel form of cellular plasticity. Neuron 39, 793-806.

Chen, C.S., Becket Ebitz, R., Bindas, S.R., Redish, A.D., Hayden, B.Y., Grissom, N.M. , 2021. Divergent strategies for learning in males and females. Current Biology, 31, 39-50.

Chen, Y., Cantrell, A.R., Messing, R.O., Scheuer, T., Catterall, W.A., 2005. Specific Modulation of Na+ Channels in Hippocampal Neurons by Protein Kinase C□. J. Neurosci. 25, 507-513.

Cho, K.K., Hoch, R., Lee, A.T., Patel, T., Rubenstein, J.L., Sohal, V.S., 2015. Gamma rhythms link prefrontal interneuron dysfunction with cognitive inflexibility in Dlx5/6+/– mice. Neuron 85, 1332-1343.

Chourbaji, S., Brandwein, C., Gass, P., 2011. Altering BDNF expression by genetics and/or environment: impact for emotional and depression-like behaviour in laboratory mice. Neurosci. Biobehav. Rev. 35, 599-611.

Clark, S.M., Notarangelo, F.M., Li, X., Chen, S., Schwarcz, R., Tonelli, L.H., 2019. Maternal immune activation in rats blunts brain cytokine and kynurenine pathway responses to a second immune challenge in early adulthood. Prog. Neuro-Psychopharmacol. Biol. Psychiatry 89, 286-294.

Connors, E.J., Shaik, A.N., Migliore, M.M., Kentner, A.C., 2014. Environmental enrichment mitigates the sex-specific effects of gestational inflammation on social engagement and the hypothalamic pituitary adrenal axis-feedback system. Brain, Behavior, and Immunity, 42, 178-190.

Deslauriers, J., Racine, W., Sarret, P., Grignon, S., 2014. Preventive effect of α-lipoic acid on prepulse inhibition deficits in a juvenile two-hit model of schizophrenia. Neuroscience 272,

261-270.

Ellis, B.J., Abrams, L.S., Masten, A.S., Sternberg, R.J., Tottenham, N., Frankenhuis, W.E., 2020. Hidden talents in harsh environments. Development and psychopathology 1, 19.

Enomoto T, Tse MT, Floresco SB. Reducing prefrontal gamma-aminobutyric acid activity induces cognitive, behavioral, and dopaminergic abnormalities that resemble schizophrenia. Biological Psychiatry. 2011; 69:432–441

Estes, M.L., McAllister, A.K., 2016. Maternal immune activation: Implications for neuropsychiatric disorders. Science 353, 772-777.

Fone, K.C., Porkess, M.V., 2008. Behavioural and neurochemical effects of post-weaning social isolation in rodents—relevance to developmental neuropsychiatric disorders. Neurosci. Biobehav. Rev. 32, 1087-1102.

Fuchs, E.C., Zivkovic, A.R., Cunningham, M.O., Middleton, S., LeBeau, F.E., Bannerman, D.M., Rozov, A., Whittington, M.A., Traub, R.D., Rawlins, J.N.P., 2007. Recruitment of parvalbumin-positive interneurons determines hippocampal function and associated behavior. Neuron 53, 591-604.

Fung, S.J., Webster, M.J., Sivagnanasundaram, S., Duncan, C., Elashoff, M., Weickert, C.S., 2010. Expression of interneuron markers in the dorsolateral prefrontal cortex of the developing human and in schizophrenia. Am. J. Psychiatry 167, 1479-1488.

Gilman, S., Cherkerzian, S., Buka, S., Hahn, J., Hornig, M., Goldstein, J., 2016. Prenatal immune programming of the sex-dependent risk for major depression. Transl. Psychiatry 6, e822-e822.

Giovanoli, S., Engler, H., Engler, A., Richetto, J., Feldon, J., Riva, M.A., Schedlowski, M., Meyer, U., 2016. Preventive effects of minocycline in a neurodevelopmental two-hit model with relevance to schizophrenia. Transl. Psychiatry 6, e772-e772.

Giovanoli, S., Engler, H., Engler, A., Richetto, J., Voget, M., Willi, R., Winter, C., Riva, M.A., Mortensen, P.B., Feldon, J., 2013. Stress in puberty unmasks latent neuropathological consequences of prenatal immune activation in mice. Science 339, 1095-1099.

Goh, J.-Y., O'Sullivan, S.E., Shortall, S.E., Zordan, N., Piccinini, A.M., Potter, H.G., Fone, K.C., King, M.V., 2020. Gestational poly (I:C) attenuates, not exacerbates, the behavioral, cytokine and mTOR changes caused by isolation rearing in a rat 'dual-hit' model for neurodevelopmental disorders. Brain Behav. Immun. 89, 100-117.

Goldstein, J., Cherkerzian, S., Seidman, L., Donatelli, J.-A., Remington, A., Tsuang, M., Hornig, M., Buka, S., 2014. Prenatal maternal immune disruption and sex-dependent risk for psychoses. Psychol. Med. 44, 3249.

Gibbons, A., Dean, B., 2016. The cholinergic system: an emerging drug target for schizophrenia. Current Pharmaceutical Design, 22, 2124-2133.

Haemisch, A., Voss, T., Gärtner, K., 1994. Effects of environmental enrichment on

aggressive behavior, dominance hierarchies, and endocrine states in male DBA/2J mice. Physiol. Behav. 56, 1041-1048.

Haddad, F., Patel, S., Schmid, S., 2020. Maternal immune activation by Poly I: C as a preclinical model for neurodevelopmental disorders: a focus on autism and schizophrenia. Neurosci. Biobehav. Rev.

Hasham, M.I., Pelech, S.L., Krieger, C., 1997. Glutamate-mediated activation of protein kinase C in hippocampal neurons. Neurosci. Lett. 228, 115-118.

Hashimoto, T., Volk, D.W., Eggan, S.M., Mirnics, K., Pierri, J.N., Sun, Z., Sampson, A.R., Lewis, D.A., 2003. Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. J. Neurosci. 23, 6315-6326.

Herring, A., Ambrée, O., Tomm, M., Habermann, H., Sachser, N., Paulus, W., Keyvani, K., 2009. Environmental enrichment enhances cellular plasticity in transgenic mice with Alzheimer-like pathology. Exp. Neurol. 216, 184-192.

Horner, A.E., Heath, C.J., Hvoslef-Eide, M., Kent, B.A., Kim, C.H., Nilsson, S.R., Alsiö, J., Oomen, C.A., Holmes, A., Saksida, L.M., 2013. The touchscreen operant platform for testing learning and memory in rats and mice. Nature protocols 8, 1961-1984.

Hupalo, S., Bryce, C.A., Bangasser, D.A., Berridge, C.W., Valentino, R. J., Floresco, S.B. 2019. Corticotropin-releasing factor (CRF) circuit modulation of cognition and motivation. Neuroscience and Biobehavioral Reviews, 103, 50-59.

Hyde, J.S., 2016. Sex and cognition: gender and cognitive functions. Curr. Opin. Neurobiol. 38, 53-56.

Jonasson, Z., 2005. Meta-analysis of sex differences in rodent models of learning and memory: a review of behavioral and biological data. Neurosci. Biobehav. Rev. 28, 811-825.

Kempermann, G., 2019. Environmental enrichment, new neurons and the neurobiology of individuality. Nat. Rev. Neurosci. 20, 235-245.

Kentner, A.C., McLeod, S.A., Field, E.F., Pittman, Q.J., 2010. Sex-dependent effects of neonatal inflammation on adult inflammatory markers and behavior. Endocrinology, 151, 2689-2699.

Kenton, J.A., Castillo, V.K., Kehrer, P., Brigman, J.L., 2020. Moderate prenatal alcohol exposure impairs visual-spatial discrimination in a sex-specific manner: effects of testing order and difficulty on learning performance. Alcoholism Clinical and Experimental Research 44(10) DOI: 10.1111/acer.14426.

Lin, E.-J.D., Choi, E., Liu, X., Martin, A., During, M.J., 2011. Environmental enrichment exerts sex-specific effects on emotionality in C57BL/6J mice. Behav. Brain Res. 216, 349-357.

Lins, B.R., Hurtubise, J.L., Roebuck, A.J., Marks, W.N., Zabder, N.K., Scott, G.A., Greba, Q., Dawicki, W., Zhang, X., Rudulier, C.D., 2018. Prospective analysis of the effects of

maternal immune activation on rat cytokines during pregnancy and behavior of the male offspring relevant to schizophrenia. ENeuro. 5: ENEURO.0249-18.2018.

Ma, J., Duan, Y., Qin, Z., Wang, J., Liu, W., Xu, M., Zhou, S., Cao, X., 2015. Overexpression of αCaMKII impairs behavioral flexibility and NMDAR-dependent long-term depression in the medial prefrontal cortex. Neuroscience, 310, 528-540.

Mac Giollabhui, N., Breen, E.C., Murphy, S.K., Maxwell, S.D., Cohn, B.A., Krigbaum, N.Y., Cirillo, P.M., Perez, C., Alloy, L.B., Drabick, D.A., 2019. Maternal inflammation during pregnancy and offspring psychiatric symptoms in childhood: timing and sex matter. Journal of psychiatric research 111, 96-103.

Mahic, M., Che, X., Susser, E., Levin, B., Reichborn-Kjennerud, T., Magnus, P., Stoltenberg, C., Chauhan, L., Briese, T., Bresnahan, M., 2017. Epidemiological and serological investigation into the role of gestational maternal influenza virus infection and autism spectrum disorders. Msphere 2.

Makinson, R., Lloyd, K., Grissom, N., Reyes, T.M., 2019. Exposure to in utero inflammation increases locomotor activity, alters cognitive performance and drives vulnerability to cognitive performance deficits after acute immune activation. Brain Behav. Immun. 80, 56-65.

Marashi, V., Barnekow, A., Ossendorf, E., Sachser, N., 2003. Effects of different forms of environmental enrichment on behavioral, endocrinological, and immunological parameters in male mice. Horm. Behav. 43, 281-292.

Meehan, C., Harms, L., Frost, J.D., Barreto, R., Todd, J., Schall, U., Weickert, C.S., Zavitsanou, K., Michie, P.T., Hodgson, D.M., 2017. Effects of immune activation during early or late gestation on schizophrenia-related behaviour in adult rat offspring. Brain Behav. Immun. 63, 8-20.

Meyer, U., 2019. Neurodevelopmental resilience and susceptibility to maternal immune activation. Trends Neurosci. 42, 793-806.

Meyer, U., Knuesel, I., Nyffeler, M., Feldon, J., 2010. Chronic clozapine treatment improves prenatal infection-induced working memory deficits without influencing adult hippocampal neurogenesis. Psychopharmacology 208, 531-543.

Miller, E.K., Cohen, J.D., 2001. An integrative theory of prefrontal cortex function. Annu. Rev. Neurosci. 24, 167-202.

Miles, J., Shevlin, M., 2001. Applying Regression and Correlation: A Guide for Students and Researchers, first ed. (London).

Moncek, F., Duncko, R., Johansson, B., Jezova, D., 2004. Effect of environmental enrichment on stress related systems in rats. J. Neuroendocrinol. 16, 423-431.

Morano, R., Hoskins, O., Smith, B.L., Herman, J.P., 2019. Loss of environmental enrichment elicits behavioral and physiological dysregulation in female rats. Front. Behav. Neurosci. 12, 287.

Morano, R., Hoskins, O., Smith, B.L., Herman, J.P., 2019. Loss of environmental enrichment elicits behavioral and physiological dysregulation in female rats. Front. Behav. Neurosci. 12, 287.

Mueller, F.S., Richetto, J., Hayes, L.N., Zambon, A., Pollak, D.D., Sawa, A., Meyer, U., Weber-Stadlbauer, U., 2019. Influence of poly (I: C) variability on thermoregulation, immune responses and pregnancy outcomes in mouse models of maternal immune activation. Brain Behav. Immun. 80, 406-418.

Murray, B.G., Davies, D.A., Molder, J.J., Howland, J.G., 2017. Maternal immune activation during pregnancy in rats impairs working memory capacity of the offspring. Neurobiol. Learn. Mem. 141, 150-156.

Nakamura, J.P., Gillespie, B., Gibbons, A., Jaehne, E.J., Du, X., Chan, A., Schroeder, A., van den Buuse, M., Sundram, S., Hill, R.A., 2021. Maternal immune activation targeted to a window of parvalbumin interneuron development improves spatial working memory: Implications for autism. Brain Behav. Immun. 91, 339-349.

Natividad, L.A., Steinman, M.Q., Laredo S.A., Irimia, C., Polis. I.Y., Lintz, R., Buczynski, M.W., Martin-Fardon, R., Roberto, M., Parsons, L.H., 2015. Phosphoylation of calcium/calmodulin-dependent protein kinase II in the rat dorsal medial prefrontal cortex is associated with alcohol-induced cognitive inflexibility. Addict. Biol., 23, 1117-1129.

Nelson, T.J., Sun, M.-K., Hongpaisan, J., Alkon, D.L., 2008. Insulin, PKC signaling pathways and synaptic remodeling during memory storage and neuronal repair. Eur. J. Pharmacol. 585, 76-87.

Ohline, S., Abraham, W., 2019. Environmental enrichment effects on synaptic and cellular physiology of hippocampal neurons. Neuropharmacology 145, 3-12.

Paxinos, G., Franklin, K.B.J., 2019. The mouse brain in stereotaxic coordinates. 5th Edition. Academic Press, Elsevier.

Peña, Y., Prunell, M., Dimitsantos, V., Nadal, R., Escorihuela, R.M., 2006. Environmental enrichment effects in social investigation in rats are gender dependent. Behav. Brain Res. 174, 181-187.

Radke, A.K., Zweifel, L.S., Holmes, A., 2019. NMDA receptor deletion on dopamine neurons disrupts visual discrimination and reversal learning. Neuroscience Letters, 699, 109-114.

Radke, A.K., Piantadosi, P.T., Uhl, G.R., Hall, F.S., Holmes, A., 2020. Improved visual discrimination learning in mice with partial 5-HT2B gene deletion. Neuroscience Letters, 738, 135378.

Shirey, J.K., Brady, A.E., Jones, P.J., Davis, A.A., Bridges, T.M., Kennedy, J.P., Jadhav, S.B., Menon, U.N., Xiang, Z., Watson, M.L., Christian, E.P., Doherty, J.J., Quirk, M.C.,

Snyder, D.H., Lah, J.J., Levey, A.I., Nicolle, M.M., Lindsley, C.W., Conn, P.J., 2009. A selective allosteric potentiator of the M1 muscarinic acetylcholine receptor increases activity of medial prefrontal cortical neurons and restores impairments in reversal learning. J. Neurosci. 29, 14271–14286.

Smith, B.L., Lyons, C.E., Correa, F.G., Benoit, S.C., Myers, B., Solomon, M.B., Herman, J.P., 2017. Behavioral and physiological consequences of enrichment loss in rats. Psychoneuroendocrinology 77, 37-46.

Sohal, V.S., Zhang, F., Yizhar, O., Deisseroth, K., 2009. Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. Nature 459, 698-702.

Sørensen, H.J., Mortensen, E.L., Reinisch, J.M., Mednick, S.A., 2008. Association between prenatal exposure to bacterial infection and risk of schizophrenia. Schizophr. Bull. 35, 631-637.

Sun, M.-K., Alkon, D.L., 2010. Pharmacology of protein kinase C activators: cognition-enhancing and antidementic therapeutics. Pharmacol. Ther. 127, 66-77.

Takahashi, E., Niimi, K., Itakura, C., 2009. Enhanced CaMKII activity and spatial cognitive function in SAMP6 Mice. Behavioral Neuroscience, 123, 527-532.

Testa, D., Prochiantz, A., Di Nardo, A.A., 2019. Perineuronal nets in brain physiology and disease, Semin. Cell Dev. Biol. Elsevier, pp. 125-135.

Van Dellen, A., Blakemore, C., Deacon, R., York, D., Hannan, A.J., 2000. Delaying the onset of Huntington's in mice. Nature 404, 721-722.

Vanover, K.E., Veinbergs, I., Davis, R.E., 2008. Antipsychotic-like behavioral effects and cognitive enhancement by a potent and selective muscarinic M-sub-1 receptor agonist, AC-260584. Behav. Neurosci. 122, 570–575.

Wallace, J., Marston, H., McQuade, R., Gartside, S., 2014. Evidence that aetiological risk factors for psychiatric disorders cause distinct patterns of cognitive deficits. European Neuropsychopharmacology 24, 879-889.

Waltz, J.A., 2017. The neural underpinnings of cognitive flexibility and their disruption in psychiatric illness. Neuroscience, 345, 203-217.

Xiong, C.H., Liu, M.G., Zhao, L.X., Chen, M.W., Tang, L., Yan, Y.H., Chen, H.Z., Qiu, Y., 2019. M1 muscarinic receptors facilitate hippocampus-dependent cognitive flexibility via modulating GluA2 subunit of AMPA receptors. Neuropharmacology, 146, 242-251.

Zhang, T.-Y., Keown, C.L., Wen, X., Li, J., Vousden, D.A., Anacker, C., Bhattacharyya, U., Ryan, R., Diorio, J., O'Toole, N., 2018. Environmental enrichment increases transcriptional and epigenetic differentiation between mouse dorsal and ventral dentate gyrus. Nature communications 9, 1-11.

Table 1. Prefrontal cortex mRNA expression in offspring exposed to maternal immune activation and housed in SD or EE

Brain region	Sex	Neural marker	SD Saline	SD Poly (I:C)	EE Saline	EE Poly (I:C)
Prefrontal Cortex	Males	Gabgr2 [*, ##]	1.00 ± 0.09	1.32 ± 0.13	0.69 ± 0.09	0.88 ± 0.14
		Grin1	1.00 ± 0.06	1.42 ± 0.11	1.45 ± 0.16	1.38 ± 0.14
		Grin2a	1.00 ± 0.07	1.18 ± 0.08	0.82 ± 0.08	1.06 ± 0.07
		Grin2b [***]	1.00 ± 0.08	1.25 ± 0.08	0.91 ± 0.06	1.30 ± 0.08
		Drd1 [#]	1.00 ± 0.10	1.22 ± 0.10	0.83 ± 0.07	0.88 ± 0.13
		Drd2	1.14 ± 0.24	1.44 ± 0.33	0.96 ± 0.14	1.20 ± 0.10
		5htr2a [**]	1.00 ± 0.10	1.44 ± 0.15	0.81 ± 0.09	1.17 ± 0.19
		Slc18a [*]	1.14 ± 0.33	2.26 ± 0.50	1.99 ± 0.44	2.97 ± 0.60
		Chrm1 [***]	1.00 ± 0.06	1.32 ± 0.07	1.14 ± 0.09	1.48 ± 0.10
		Chrm4 [†]	1.00 ± 0.11	1.18 ± 0.12	0.83 ± 0.07 [bbb]	1.51 ± 0.12
		Chat [†]	1.00 ± 0.09	0.40 ± 0.17 [a]	0.76 ± 0.08	0.93 ± 0.22
		Camk2a [**, ##]	1.00 ± 0.06	1.35 ± 0.08	1.35 ± 0.09	1.52 ± 0.11
		Prkca [***]	1.00 ± 0.09	1.85 ± 0.18	0.98 ± 0.09	1.64 ± 0.19
	Females	Gabgr2 [*, ##]	1.00 ± 0.08	1.17 ± 0.06	0.89 ± 0.09	0.75 ± 0.10
		Grin1 [††]	1.00 ± 0.06	1.40 ± 0.10 [aa,bb]	1.82 ± 0.22	1.28 ± 0.05
		Grin2a	1.00 ± 0.72	0.78 ± 0.03	0.66 ± 0.06	0.70 ± 0.06
		Grin2b	1.00 ± 0.05	1.09 ± 0.06	0.92 ± 0.05	0.97 ± 0.07
		Drd1	1.00 ± 0.13	1.02 ± 0.08	0.92 ± 0.06	0.80 ± 0.15
		Drd2	1.00 ± 0.13	1.57 ± 0.37	1.53 ± 0.34	1.26 ± 0.26
		5htr2a [††]	1.00 ± 0.09	1.50 ± 0.08 [aaa,bbb]	1.07 ± 0.07	0.91 ± 0.11
		Slc18a	1.00 ± 0.19	0.99 ± 0.21	1.49 ± 0.34	1.25 ± 0.32
		Chrm1 [##]	1.00 ± 0.04	1.10 ± 0.07	1.49 ± 0.12	1.26 ± 0.10
		Chrm4	1.00 ± 0.16	1.21 ± 0.09	1.32 ± 0.07	1.01 ± 0.13
		Chat	1.00 ± 0.24	1.11 ± 0.47	1.43 ± 0.19	1.19 ± 0.50
		Camk2a [††]	1.00 ± 0.07	1.67 ± 0.11 [aaa,bbb]	1.86 ± 0.21 [bbb]	1.32 ± 0.10
		Prkca [††]	1.00 ± 0.07	1.59 ± 0.15 [aa,bb]	1.43 ± 0.10 [bb]	1.10 ± 0.15
Hippocampus	Males	Gabgr2 [†††]	1.00 ± 0.07	2.19 ± 0.16 [aaa,bb]	1.86 ± 0.07 [aaa]	1.29 ± 0.30
		Grin1 [*]	1.00 ± 0.10	1.45 ± 0.14	0.89 ± 0.07	1.19 ± 0.16
		Grin2a [#]	1.00 ± 0.08	1.04 ± 0.14	1.14 ± 0.09	0.91 ± 0.07
		Grin2b [†††]	1.00 ± 0.06	1.62 ± 0.10 [aaa,bb]	1.43 ± 0.09 [aaa,b]	0.96 ± 0.17
		Drd1	1.00 ± 0.10	1.62 ± 0.35	1.49 ± 0.25	1.56 ± 0.26
		Drd2	1.00 ± 0.10	2.07 ± 0.19	1.59 ± 0.12	1.57 ± 0.33
	Females	Gabgr2 [††]	1.00 ± 0.15	1.49 ± 0.08 [aa,bb]	0.84 ± 0.05	0.77 ± 0.12
		Grin1 [##]	1.00 ± 0.10	0.84 ± 0.06	1.32 ± 0.09	1.45 ± 0.17
		Grin2a	1.00 ± 0.12	0.82 ± 0.07	0.81 ± 0.05	0.69 ± 0.06
		Grin2b [†]	1.00 ± 0.11	1.15 ± 0.06 [bbb]	0.86 ± 0.05	0.72 ± 0.04
		Drd1 [#]	1.00 ± 0.19	0.96 ± 0.11	0.68 ± 0.08	0.76 ± 0.9
		Drd2 [†††]	1.00 ± 0.12	1.60 ± 0.16 [aa,b]	1.61 ± 0.19 [a,b]	1.10 ± 0.08

Data are mean ± SEM.

† P<0.05, significant interaction (MIA x Housing)

* P<0.05, ** P<0.01, significant main effect of MIA (saline vs. poly (I:C))

# P<0.05, ## P<0.01, significant main effect of housing (SD vs. EE)

Lower case letters indicate significant differences between conditions ([a]different from SD-Saline, [b]different from EE-Poly (I:C))