Demands of visual processing hierarchy shape laminar compartmentalization of attention modulation of luminance contrast in area V4

Xiang Wang[1], Anirvan S. Nandy[1,3†] and Monika P. Jadi[1,2,3,*,†]

[1]Interdepartmental Neuroscience Program, Yale University, New Haven, CT 06511

[2]Department of Psychiatry, Yale University, New Haven, CT 06511

[3]Department of Neuroscience, Yale University, New Haven, CT 06511

[†]senior author

[*]corresponding author (monika.jadi@yale.edu)

ABSTRACT

Contrast is a key feature of the visual scene that aids object recognition. Attention has been shown to selectively enhance the responses to low contrast stimuli in visual area V4, a critical hub that sends projections both up and down the visual hierarchy. Veridical encoding of contrast information is a key computation in early visual areas, while later stages encode higher level features that benefit from improved sensitivity to low contrast. How area V4 meets these distinct information processing demands in the attentive state is not known. We found that attentional modulation of contrast responses in area V4 is cortical layer and cell-class specific. Putative excitatory neurons in the superficial output layers that project to higher areas show enhanced boosting of low contrast information. On the other hand, putative excitatory neurons of deep output layers that project to early visual areas exhibit contrast-independent scaling. Computational modeling revealed that such layer-wise differences may result from variations in spatial integration extent of inhibitory neurons. These findings reveal that the nature of interactions between attention and contrast in V4 is highly compartmentalized, in alignment with the demands of the visual processing hierarchy.

INTRODUCTION

Voluntary attention is essential for sensory guided behavior and memory formation (Petersen and Posner, 2012). Failures in sensory processing and selective attention are aspects of many mental illnesses, including schizophrenia and mood disorders (Fioravanti et al., 2005; McIntyre et al., 2010; Neuchterlein et al., 1991). Visual spatial attention plays a critical role in visual sensory processing: It allows improved perception of behaviorally relevant target stimuli among competing distractors by boosting the apparent visibility of the target (Carrasco et al., 2004). At the neuronal level, attention modulates the activity of cortical neurons that encode an attended visual stimulus at various stages of visual processing (Bisley and Goldberg, 2003; Ghose and Maunsell, 2008; Moran and Desimone, 1985; Motter, 1993; Reynolds et al., 1999; Treue and Martinez Trujillo, 1999; Treue and Maunsell, 1996). In visual areas such as V4 and MT, attention modulates neuronal mean firing rates, increases their firing reliability, and reduces the co-variability among pairs of neurons (Cohen and Maunsell, 2009; Mitchell et al., 2007, 2009; Reynolds and Chelazzi, 2004; Treue and Martinez Trujillo, 1999). However, the computational principles that underlie the activity of neuronal populations that represent both sensory information and the attentional state remain poorly understood (Moore and Zirnsak, 2017; Reynolds and Chelazzi, 2004).

Object recognition is mediated by a hierarchy of cortical visual processing areas that form the ventral visual stream. Contrast is a key feature of the visual scene that aids object recognition, and the encoding of contrast information is one of the most important computations performed by early visual areas. On the other hand, visual features represented in higher areas such as the inferotemporal (IT) cortex benefit from improved sensitivity to low contrast stimuli (Avidan et al., 2002; Rolls and Baylis, 1986). Visual area V4 is a critical hub in the ventral stream that sends feedforward projections to areas such as IT and feedback projections to early visual processing areas (Anderson and Martin, 2006; Douglas and Martin, 1991; Van Essen and Maunsell, 1983). Attention has been shown to selectively enhance the responses to low contrast stimuli (Martinez-Trujillo and Treue, 2002; Reynolds et al., 2000). Attention mediated selective enhancement of low

contrast features is thought to aid invariant representations in higher object recognition
areas downstream of V4 (Roe et al., 2012). However, such a bias in the attention-
modulated feedback from V4 to upstream visual areas can disrupt the contrast-based
feature extraction functions of these stages. How area V4 meets these distinct information
processing demands of the visual processing hierarchy is not known. While attention can
enhance V4 responses in a contrast-independent manner (response gain) under certain
experimental conditions (Williford and Maunsell, 2006), an understanding of robust
mechanisms of feedback from V4 that does not interfere with the contrast landscape of
scene representations in early visual areas remains elusive.
69
One possibility is that distinct subpopulations in V4 mediate these functional demands.
Indeed, the sensory cortical sheet, including area V4, is not a homogeneous piece of
tissue along its depth; rather, it has a six-layered or laminar structure made up of multiple
cell classes, of both excitatory and local inhibitory kind, with largely stereotypical
anatomical connectivity between and within layers (Douglas and Martin, 2004). Layer 4
(the *input* layer) is the primary target of projections carrying visual information from early
areas, such as V1, V2, and V3 (Felleman and Van Essen, 1991; Ungerleider et al., 2008).
Visual information is then processed by local neural subpopulations as it is sent to layers
2/3 (the *superficial* layer) and layers 5/6 (the *deep* layer), which serve as output nodes in
the laminar circuit (Hirsch and Martinez, 2006; Rockland and Pandya, 1979). The
superficial layers feed information forward to downstream visual areas, such as IT (Borra
et al., 2010; Distler et al., 1993), whereas the deep layers send feedback information to
upstream early visual areas (Callaway, 1998; Gattass et al., 2014; Mehta et al., 2000;
Ungerleider et al., 2008). This anatomical organization suggests distinct functional roles
(D'Souza and Burkhalter, 2017), and differential attentional modulation of sensory
representation among cell-class and layers-specific neural subpopulations. In support of
this idea, a recent study of simultaneous depth recordings in visual area V4 has shown
layer-specific attentional modulation of average neuronal responses, reliability of
responses, and correlations between responses of pairs of neurons (Nandy et al., 2017).
Therefore, to fully understand the attentional modulation of sensory computations, it is

essential to investigate the modulation of sensory representation in these subpopulations.

Our broad hypothesis is that the attentional modulation of contrast computations in area

V4 is not homogeneous, but rather is layer- and cell-class specific and that these

differences reflect the different computational demands on these subpopulations.

Considering their key contribution to feedback projections to early visual areas, we

specifically expect that projection neurons in the deep layers show uniform attentional

modulation across all contrasts in order to minimally impact the faithful representation of

contrast landscape in their target areas.

In this study, we characterized layer- and cell-class specific neural subpopulations from

extracellular simultaneous laminar recordings of single neurons within area V4 of

macaque monkeys performing an attention-demanding task. Using unsupervised

clustering techniques on spiking properties, we distinguished five functional clusters of

neurons. We distinguished layer identities – superficial, input or deep – of these neurons

using features of local field potentials. To test our hypothesis, we characterized the

attentional modulation of contrast response functions in these sub-populations. We

interpreted our findings within a computational framework of attentional modulation of

contrast responses (Reynolds and Heeger, 2009), which yielded predictions for distinct

mechanistic roles of these neural subpopulations in attentive perception.

RESULTS

In the primate visual system, cortical sensitivity to features such as luminance contrast

varies with the locus of spatial attention; contrast response functions (CRF) of cortical

neurons are measured to quantify this dependence (Kastner and Ungerleider, 2000;

Reynolds and Chelazzi, 2004; Reynolds et al., 2000). However, the laminar- and cell-

class specific dependence of CRF on the attentive state is not known. Using linear array

electrodes, we recorded neuronal activity from well-isolated single units, multi-unit

clusters, and local field potentials (LFPs) in visual area V4 of two rhesus macaques (right

hemisphere in monkey A, left hemisphere in monkey C) during an attention demanding

orientation change detection task (Figure 1A, B; see Methods). We used current source

density (CSD) analysis to identify different laminar compartments (*superficial, input*, and

deep), and assigned isolated single units to one of the three layers (see Methods). In the

main experiment, we presented a sequence of paired Gabor stimuli with different

contrasts (Figure 1B); one stimulus was presented inside the receptive fields (RFs) of the

recorded neurons and the other at an equally eccentric location across the vertical

meridian. Attention was cued either to the stimuli within the neurons' RFs ("attend-in") or

to the stimuli in the contralateral visual hemifield ("attend-away").

Attentional Modulation of Contrast Response Function

To examine the effects of attention on individual neurons, we used the method of ordinary

least squares to fit each neuron's contrast responses from both attentional states to a

hyperbolic ratio function (Figure 1C). This function is described by four parameters: r_{max},

c_{50}, m, and n, where r_{max} is the attainable maximum response, c_{50} is the contrast at

which neuronal response is half-maximal, m is the baseline activity, and n describes the

nonlinearity of the function. Attention effects differ considerably for individual neurons.

Attention either enhances or suppresses neuronal responses at different contrast levels

(Figure 1D). We quantified the effect of attention on every recorded neuron by computing

the attentional modulation index (AMI) using contrast responses from both attention

conditions (see Methods). We saw a significant variance of AMI values at each contrast

level (Figure 1E). We also examined how attention impacts the values of best-fitting

parameters (Figure 1F). The mean AMIs for r_{max} and m are significantly higher than zero

(Mann-Whitney U test, $p < 0.01$ for both distributions), which is consistent with previous

observations in V4 (Williford and Maunsell, 2006). The same percentage change in r_{max}

and m (15% increase) supports an effect of contrast independent scaling by attention.

The average modulations of c_{50} and n are significantly smaller than zero (Mann-Whitney

U test, $p < 0.01$ for c_{50} and $p \ll 0.01$ for n), suggesting an increased sensitivity to low

contrast stimuli and a reduction in the sensitivity to contrast change, respectively. The

bootstrap sampling distributions of the mean difference from 0 support the average

attention effects on r_{max}, n and m (Figure 1G). These results indicate that the overall

effect of attention on V4 neuron responses cannot be simply explained as selective

boosting of low contrast. It is a combination of modulations in multiple parameters of the

contrast response function (Figure 1F, G).

Classification of Single Units Using Electrophysiological Features

To investigate whether attention modulates different classes of neurons uniformly or

differentially, we characterized classes of single units based on two electrophysiological

properties extracted from extracellular recordings: the peak-to-trough duration (PTD) and

the local variation (Lv). Properties of the action potential waveform, especially the PTD,

have been extensively used to classify neurons into narrow- (putative inhibitory) and

broad-spiking (putative excitatory) cells (Constantinidis and Goldman-Rakic, 2002;

Diester and Nieder, 2008; Hussar and Pasternak, 2009; Johnston et al., 2009; Kaufman

et al., 2010; Mitchell et al., 2007; Wilson et al., 1994). The shapes of average spiking

waveform for all single units in our data were also highly variable (Figure 2A). We

exploited the information structure in the entire waveforms by applying principal

component analysis (PCA). The correlation pattern between the first two components of

the PCA (cumulative percentages of explained variance: 59.62%, 83.10%) supported the

idea that neurons can be separated into meaningful clusters by waveform shape

measures (Figure 2B). The clusters generated by neurons' PTDs in the PCA component

space were minimally overlapped (Supp. Figure 2E). Therefore, we chose PTD instead

of PCA components as one of the classification features for further analysis since the

PTD is more interpretable.

Firing variability measures have been previously used as an additional electrophysiology-

based dimension along which neurons have been found to be separable (Anderson et al.,

2011; Ardid et al., 2015; Degenetais et al., 2002). We used Lv, a measure that effectively

characterizes neurons' intrinsic spiking, and controls the effect of transient variations in

firing rates (Shinomoto et al., 2003) (see Methods). To achieve stable classification of

single units across attention conditions, we verified that Lv was not significantly

modulated by attentional states (Figure 2C).

We used a meta-clustering analysis based on the *k*-means clustering algorithm (see

Methods) in the two-dimensional space of PTD and Lv, and identified five clusters of

isolated single units (Figure 2D) (Ardid et al., 2015; Hartigan and Wong, 1979). The five-

cluster result was picked because it was the largest set of distinct cell classes that

characterized a majority (99.7%) of single units in the dataset (Supp. Figure 2A). Narrow-

spiking cells become a cluster by themselves, while those classified as broad-spiking cells

(Mitchell et al., 2007; Nandy et al., 2017) are split into four clusters. Based on the average

PTD and Lv of each cluster, we termed these five clusters as Narrow, Medium Regular,

Medium Bursty, Broad Regular, and Broad Bursty.

We validated our classification results using several methods (see Methods). First, we

gathered additional support for the meta-clustering based number of clusters by applying

a data-driven approach based on a novel form of cross-validation (Fu and Perry, 2020).

The method incorporates clustering results from the unsupervised algorithm into its

supervised training of linear classifiers to produce cross-validation errors (see Methods).

The five-cluster result showed the lowest cross-validation error (Supp. Figure 2B).

Second, we validated the stability of the clustering result by bootstrap subsampling

analysis (Hennig, 2007). The Jaccard similarity, averaged across subsamples, is a

measure of each cluster's robustness regarding its sensitivity to the amount of data. All

clusters in the five-cluster result had average Jaccard similarities greater than 0.5,

implying that clusters remained stable under subsampling (Supp. Figure 2C). A cell-wise

co-clustering matrix showing the probability that each pair of neurons belongs to a same

cluster across all subsamples also supported the number of clusters we chose (Supp.

Figure 2D). Third, we visualized our dataset by applying nonlinear transformations: t-SNE

(Hinton and Roweis, 2003) and UMAP (McInnes et al., 2018). Although these techniques

are generally suited for embedding high-dimensional data for visualization in a low-

dimensional space, their algorithms that enlarge the distance differences in the original

dataset also make them useful for recovering well-separated clusters. When we explored

the hyperparameters of both algorithms, we found that most of the five clusters were still

separable in both t-SNE and UMAP space (Figure 2E; Supp. Figure 2G, H). Notably, all

four non-Narrow clusters were separable, including the Medium Regular and the Medium

Bursty which occupied distinct locations in the t-SNE and UMAP space (Supp. Figure 2G,

H).

One of the assumptions we made to use the PTD as a clustering feature was that it

captures a significant amount of the variations of neurons' spiking waveforms. We tested

this assumption by clustering neurons in the principal component space of the AP

waveform and comparing them with neuronal groups defined by their PTD. We divided

neurons into narrow- (0-250 μs), medium- (250-350 μs), and broad-spiking (350-550 μs)

groups, and found that the 3 clusters generated from the *k*-means clustering were

consistent with the 3 neuronal groups defined by the spike width (Supp. Figure 2F).

The clusters differ in terms of their firing rates (Supp. Figure 2I). Notably, Narrow class

neurons exhibited higher firing rates than the Broad Regular cluster when averaged

across layers (mean 10.2 Hz compared to 5.6 Hz, Mann-Whitney U test, $p < 0.05$). It is in

agreement with previous findings that narrow-spiking neurons, considered putative

inhibitory interneurons, show higher firing rates than broad-spiking neurons, thought to

be putative excitatory pyramidal cells (Connors and Gutnick, 1990; McCormick et al.,

1985; Mitchell et al., 2007; Nowak et al., 2003; Povysheva et al., 2006).

Cell-Class and Layer-Specific Attentional Modulation

We next examined how attention modulates contrast responses for each cell class. We

first computed the AMIs of best-fitting CRF parameters for every cell class. The pattern

of modulations of CRF parameters was distinct for individual cell classes (Figure 2F).

Narrow and Medium Regular cell classes showed significant positive modulations of

r_{max} only, implying a contrast-independent effect by attention. On the other hand, both

Broad Regular and Broad Bursty classes showed significant negative modulations of c_{50}

(Figure 2F), suggesting a selective enhancement of responses to low contrast stimuli.

This effect was novel to these classes and not revealed in the analysis of unclassified

neurons (Figure 1G). None of the remaining cell classes – Narrow, Medium Bursty and

Medium Regular – showed a significant modulation of c_{50} by attention, an effect that matched the analysis of unclassified neurons (Figure 1G). Medium Bursty neurons showed a modulation pattern that was distinct from the ones for any of the other four cell classes: significant positive modulations of r_{max} and baseline activity, implying a pure response gain effect by attention.

To further investigate the cell-class specific attentional modulation at each contrast level, we computed the AMI as a function of contrast using CRFs from both attentional states for every single unit and then averaged AMIs across single units within a cluster (Figure 3A, left panel). We found that the AMIs of Narrow and Medium Regular classes were relatively less dependent on contrast, whereas the remaining clusters appeared to be modulated by attention in a contrast-dependent manner (Figure 3A, left panel). When averaged across all contrasts, attention positively modulated firing rates for all cell classes except the Medium Regular class (Mann-Whitney U test, $p < 0.01$ except for MR). Further, attentional modulation differed in significant ways among the non-Narrow clusters (Figure 3A, right panel). To quantify the contrast dependence of attentional modulation for each single unit, we first averaged the AMIs within the low-contrast and the high-contrast ranges with the contrast boundary set at each unit's best-fitting c_{50} parameter. We then defined the contrast dependence index (CDI) of a single unit as the difference between the two average AMIs normalized by the AMI averaged across all contrasts (see Methods). Contrast independent modulation would then result in CDI = 0, reflecting a pure scaling effect of attention on the CRF. A positive CDI would indicate a more robust attentional modulation at the low-contrast range. A negative CDI would suggest a stronger attention effect on neural responses at the high-contrast range (Figure 3B). We examined the CDI distribution within each cell class and found that the Narrow and Medium Regular classes showed small mean CDIs, and their distributions were not significantly different from zero. However, the other 3 clusters (Medium Bursty, Broad Regular, Broad Bursty) exhibited more positive CDIs (Figure 3C). These results are consistent with our findings of AMIs of CRF parameters for each cell class (Figure 2F), confirming that attention modulated Narrow and Medium Regular cell classes' responses regardless of the stimulus contrast.

On the other hand, the modulations for Medium Bursty, Broad Regular, and Broad Bursty

classes were dependent on contrast and were more robust in the low-contrast range.

We further inspected the laminar profile of the attention effect and its contrast

dependence for every cell class (Figure 3D, E). We excluded from our analysis clusters

that contained an insufficient number of units (n < 10) in a layer. When averaged across

contrasts, (Figure 3D, right panels), Narrow class neurons showed significant attentional

modulations in the input layer, but not in the superficial or deep layer (Figure 3D, right

panels, Mann-Whitney U test, $p_{superficial} = 0.79$, $p_{input} < 0.01$, $p_{deep} = 0.06$). On the other

hand, Broad Regular neurons were robustly modulated by attention across all cortical

layers (Figure 3D, right panels, Mann-Whitney U test, $p_{superficial} \ll 0.01$, $p_{input} \ll 0.01$, p_{deep}

$\ll 0.01$). The AMI difference between these two cell classes is in agreement with the

differences between narrow- and "broad"-spiking cells previously reported in these

cortical layer (Nandy et al., 2017); it is important to note that the AMI patterns across

layers were distinct for the other three cell classes (Figure 3D). Two key laminar patterns

of contrast dependence emerged from these 5 clusters. First, the attentional modulation

of the Narrow cell class was independent of contrast across all cortical layers. Second,

the Broad Regular cell class exhibited a strong contrast dependence and, specifically, a

significant modulation in the low-contrast range in the superficial and input layers; but its

dependence on contrast was not significant in the deep layer (Figure 3E). It is important

to note that at least one non-Narrow class (Medium Regular) was functionally similar to

Narrow neurons in superficial and input layers. Also notably, the laminar differences did

not emerge when all units in a layer were analyzed as either a single class or more

conventionally as narrow vs. "broad" classes.

Laminar network mechanisms of contrast dependence of AMI across layers

We next used computational modeling to gain insights into the possible neural

mechanisms underlying the layer- and cell-class specific AMI dependency on stimulus

contrast. Variation in CDI across experimental paradigms has been previous observed

(Martinez-Trujillo and Treue, 2002; Reynolds et al., 2000; Williford and Maunsell, 2006),

and explained by paradigm-specific normalization due to attention (Reynolds and Heeger,

2009). We hypothesized that normalization mechanisms can also explain the layer-

specific differences in CDI in our empirical findings (Figure 3D, E). To test this, we first

interpreted our results in the context of the normalization model of attention (Reynolds

and Heeger, 2009) to generate predictions about layer-specific cortical connectivity that

might underlie the variations in CDI. The normalization model of attention proposes a

computational principle that accounts for various attention effects on neurons' contrast

response functions (Reynolds and Heeger, 2009). Normalization model assumes that the

relative sizes of excitatory receptive field and suppressive field of neurons, and the

'attention field' of the experimental paradigm shape the net suppressive drive to individual

neurons. The suppressive drive ultimately determines the CDI of individual neurons in a

population. We thus investigated the consequences of varying the relative sizes of

excitatory receptive field and suppressive field of individual neurons on attentional

modulations of CRFs (see Methods). This inquiry was motivated by the observation that

neuronal receptive field sizes change along the cortical depth in sensory areas (Gilbert,

1977; Sur et al., 1985; Vaiceliunaite et al., 2013), and based on the assumption that

'attention field' sizes are constant for an experimental paradigm.

We simulated the normalization model with different sizes of excitatory receptive field and

suppressive field of neurons, and generated neuronal responses to different stimulus

contrasts in "attend in" and "attend away" conditions (Figure 4A, top panel). We computed

the AMI and the CDI for each combination of size parameters (see Methods). We find

that the CDI depends both on the excitatory receptive field size and on the suppressive

field size. Holding the attention field size and the stimulus size fixed, a smaller

suppressive field or a smaller excitatory receptive field leads to a greater CDI of the

attentional modulation (Figure 4A, middle panel). On the other hand, a larger suppressive

field or a larger excitatory receptive field results in a smaller CDI (Figure 4A, middle panel).

These results hold for a wide range of values of the stimulus size and the attention field

size. The pattern is robust when the attention field and the stimulus are both small or large

(Supp. Figure 4B, i). The results are also stable for both a linear and saturating transfer

function assumption between the stimulus contrast and excitatory drive in the normalization model (Supp. Figure 4B, ii). We also computed the AMI of suppressive drive of neurons for each combination of size parameters. The CDI of model neurons is roughly proportional to the AMI of suppressive drive (Figure 4A, bottom panel). Greater the AMI of suppressive drive, stronger is the CDI of model neurons, and vice versa. Since Broad Regular neurons are putative excitatory pyramidal cells, these results suggest two possible neural mechanisms that explain the laminar profile of CDIs of Broad Regular neurons: the suppressive field size increases along the depth of V4 (Figure 4A, middle panel) or the excitatory receptive field is more extensive in the deeper layer of V4 (Supp. Figure 4C).

The normalization model predicts the AMI of the suppressive drive (Figure 4A, bottom panel) to be correlated with the CDI of neuronal responses (Figure 4A, middle panel) (Reynolds and Heeger, 2009). However, the suppressive field in the model can be implemented by various biophysical mechanisms (Carandini, 2004). One possible mechanism is shunting inhibition via lateral connections from other neurons in the cortical neighborhood (Carandini and Heeger, 1994; Carandini et al., 1997; Kouh and Poggio, 2008), in which case the receptive field of local inhibitory neurons can approximate the suppressive field. Since the average AMI of the putative inhibitory (Narrow) cluster and CDI of putative excitatory (Broad) clusters in the input and deep layers in our empirical data (Figure 3D right panels, Figure 3E) is also correlated, we further explored this mechanism mediated by local inhibitory neurons. Under this assumption, the prediction about the changes in suppressive field size down the cortical depth from the normalization model transforms into one about changes in the excitatory (E) - inhibitory (I) connectivity along the cortical depth. Similarly, the prediction about the changes in excitatory receptive field sizes down the cortical depth can also transforms into one about the changes in the E-E connectivity along the cortical depth (Gilbert and Wiesel, 1985; Hirsch and Gilbert, 1991). The layer-specificity of cortical connectivity implies different temporal signatures of neural activity across layers.

We next used a spiking network model to examine the effects of excitatory and inhibitory

receptive field sizes on spike-time correlation between populations of local excitatory (E)

and inhibitory neurons (I). Our spiking network model focuses on connectivity

mechanisms for generating variable sizes of suppressive and excitatory receptive fields

in a cortical network. The amplitude of the spike-time correlation between neurons has

been shown to depend on both the connection strength and the background synaptic

noise (Ostojic et al., 2009). Therefore, the spike-time correlation between neurons can be

a proxy for the size of the postsynaptic neuron's receptive field. We hypothesized that a

smaller receptive field of the postsynaptic neuron would make the local connections more

dominant against background inputs and lead to a higher spike-time correlation between

the locally connected neurons. We examined how spike-time correlations change as a

function of the inhibitory or excitatory receptive field size in a conductance-based model

of spiking neurons (see Methods). We set up 10 local networks or "columns" of E and I

units that were interconnected in a ring formation (Figure 4B, Supp. Figure 4C). Neurons

within the same column were mutually coupled, while interactions between columns were

confined to excitatory connections to local E and I neurons whose strengths decayed with

distance between columns. All connections occurred with a probability of 0.5. We

modeled the receptive field size as the standard deviation (σ_I or σ_E) of the connection

strength between columns (Figure 4B, Supp. Figure 4C). We performed simulations that

generated spiking activity in response to a step input (Figure 4B, bottom panel). The

spike-time correlation between local E and I populations was calculated using pooled

spike trains within the same column; the resulting spike-time correlation was averaged

across columns. We found that the inhibitory receptive field size has a critical impact on

the spike-time correlation amplitude in such a network (Figure 4C), while the excitatory

receptive field size has little effect (Supp. Figure 4C). A larger inhibitory receptive field

(larger values of σ_I) leads to a lower spike-time correlation between the local E and I

populations in the network (Figure 4C). This result suggests that the prediction about

inhibitory receptive field sizes down the cortical depth as the basis of CDI variation of

Broad Regular neurons can be tested by examining the spike-time correlation between

local E and I populations within each layer.

To test this prediction in our dataset, we computed the session-averaged spike-time

correlation between Narrow (putative inhibitory neurons) and Broad Regular (putative

excitatory neurons) single units within each layer (see Methods). We found that the spike-

time correlation amplitudes were higher in the superficial layer and the input layer than

that in the deep layer (Figure 4D). We compared the spike-time correlations in the deep

layer with those in either superficial or input layers, averaged within 3 different 50ms time

windows. The 95% confidence interval of the mean difference between layers in either

comparison was greater than 0 for the center window (Supp. Figure 4D). In accordance

with our findings from the E-I network models (Figure 4C), this suggests that inhibitory

neurons in the deep layer exhibit relatively broader receptive fields, which supports the

prediction by the normalization model of attention (Figure 4A, middle panel). Our findings

thus provide a parsimonious explanation for the layer- and cell-class specific contrast

dependence of attentional modulation observed in area V4 (Figure 4E).

DISCUSSION

Spatial attention plays a critical role in sensory guided behavior. It is thought to achieve

this by enhancing the responses to low contrast stimuli in mid-tier visual cortical areas

such as V4. While later stages of the visual processing hierarchy are thought to benefit

from this manipulation, V4 also sends feedback projections to early visual areas that use

veridical representation of contrast to aid object recognition. How area V4 meets these

distinct information processing demands is not known. Contrary to the simplifying

assumptions of prior empirical studies, we tested the hypothesis that V4 customizes its

output to different stages of the visual processing hierarchy through layer- and cell-class

specific attentional modulation of contrast computations. Recent advances in

experimental techniques have shown layer- and cell-class specific functional specificity

of computations in the cortical circuit (Adesnik and Naka, 2018; Adesnik and Scanziani,

2010; Naka and Adesnik, 2016; Olsen et al., 2012). However, these studies have been

limited to species in which higher cognitive functions, such as attention, are challenging

to study. Using computational approaches on laminar neural data in area V4 of the

macaque, we find that the attentional modulation of neural responses to visual luminance

contrast is indeed layer- and cell-class specific. We classified neurons into five functional

cell classes defined by their action potential widths and the statistics of firing variability

(Figure 2D); these classes show specificity in attention effects on their contrast response

functions (Figure 2F) and the contrast dependence of attentional modulation (Figure 3C).

Specifically, Narrow neurons show contrast-independent response modulation across

layers; Broad Regular neurons, the putative projection neurons, exhibit significant

contrast dependence of attentional modulation in the superficial layers, that project to

higher level visual areas, but not in the deep layers, that project to earlier visual areas

(Figure 3D, E). Notably, this highly significant laminar difference was not observable

without cell-class identification. These results provide the first evidence for our broad

hypothesis that attentional modulation of contrast computations in the visual cortex is

heterogeneous across those cell classes and layers that project to distinct stages of the

visual processing hierarchy. The qualitative nature of the attention modulation of contrast

in our data is not only distinct but suggests optimization for the computational demands

of the target stages. Selective boosting of responses to low contrast stimuli is

compartmentalized to the superficial output layers that project representations such as

extended contours and object surfaces to higher areas (see Roe et al., 2012 for a review).

Contrast-independent scaling of neural responses is confined to the deep output layers.

Neurons in these layers project back to early visual areas that are reliant on faithful

representation of luminance contrast for low-level feature extraction. We speculate that

the contrast-independent attentive feedback provides a spatial boost signal to early visual

areas that do not receive direct inputs from attention control centers such as the frontal

eye fields (Ungerleider et al., 2008). This also aligns with the predictive coding model of

object recognition, wherein V4 is a higher-level area in the object recognition hierarchy

that generates predictions of lower-level activity, without corrupting the sensory

landscape that is needed for error correction (Rao and Ballard, 1999).

When interpreted within the framework of the normalization model of attention (Figure

4A), the layer-specific attention modulation predicts differences in the spatial pooling of

local inhibitory populations across layers. Such differences further predict a layer-specific

signature of correlations between the activities of local inhibitory and putative excitatory

neurons when explored in a spiking E-I network model (Figure 4B, C). We find robust

evidence for differences in inhibitory spatial pooling across layers through our analyses

of correlations between putative inhibitory and putative excitatory neurons in the

superficial, input, and deep layers of the cortex (Figure 4D, E).

Classification of cell-types

The duration of the extracellular spike waveform has been used to distinguish putative

inhibitory interneurons from putative excitatory pyramidal cells in a wide range of species

and across various brain regions (Ardid et al., 2015; Bruno and Simons, 2002;

Constantinidis and Goldman-Rakic, 2002; Csicsvari et al., 1999; Fox and Ranck, 1981;

Frank et al., 2001; Mitchell et al., 2007; Nandy et al., 2017; Rao et al., 1999; Simons,

1978; Swadlow, 2003; Wilson et al., 1994). In terms of attention effects, narrow-spiking

neurons show stronger attention-dependent increases in absolute firing rates and firing

reliability than broad-spiking cells (Mitchell et al., 2007). Statistics of the firing pattern and

unsupervised clustering algorithms are also effective in identifying subpopulations of

neurons with distinct functional properties (Ardid et al., 2015; Compte et al., 2003;

Gouwens et al., 2019; Hawken et al., 2020; Shinomoto et al., 2009). It is important to note

that the clusters we distinguished based on spike width and firing variability may not

correspond to neuronal classes differentiated based on morphology or protein expression

patterns (Migliore and Shepherd, 2005; Tasic et al., 2018; Zeng and Sanes, 2017). Two

possible correspondences exist between the Narrow neurons and interneurons, and

between the Broad Regular neurons and pyramidal cells (Connors and Gutnick, 1990;

McCormick et al., 1985; Nowak et al., 2003). We find significant differences in both the

firing rate (Supp. Figure 2I) and the attentional modulation of firing rates (Figure 3A, D)

between clusters, suggesting their different functional roles in attention-mediated visual

processing. Crucially, these distinct functional roles are reflected by the differences in

contrast dependence of attentional modulation.

Relation to prior studies of spatial attention in V4

Prior studies evaluating attention effects on neuronal contrast responses proposed either contrast-independent scaling of responses, termed as response gain (McAdams and Maunsell, 1999a; Morrone et al., 2002; Pestilli et al., 2009; Treue and Martinez Trujillo, 1999) or boosting of responses to low contrast stimuli, termed as contrast gain (Li and Basso, 2008; Li et al., 2008; Martinez-Trujillo and Treue, 2002; Reynolds et al., 2000) or an intermediate effect between the two (Huang and Dobkins, 2005; Williford and Maunsell, 2006). Although the overall attentional modulation of best-fitting CRF parameters in our dataset is consistent with the intermediate effect (Figure 1F, G), attention effects on individual clusters are highly variable: a mixture effect of response gain and contrast gain is observed for Broad Regular and Broad Bursty units; Medium Bursty cluster shows a response gain change; Medium Regular and Narrow neurons are only modulated in their maximum responses (Figure 2F). Furthermore, some clusters, such as Broad Regular and Broad Bursty neurons, exhibit larger attention-dependent increases in response than the population mean, especially within the low-contrast range (Figure 3A). These observations suggest that attentional modulation of firing rate for certain cell classes may be more robust than that gleaned from previous studies that averaged across the whole recorded population. These cell-class specific increases in firing rate may significantly improve the signal-to-noise ratios of individual cell classes, and therefore, act as another important contributor to the improvement of psychophysical performance due to attention in addition to reductions in correlations (Cohen and Maunsell, 2009; Mitchell et al., 2009).

Our interpretation of the normalization model

The predictions from the normalization model (NM) of attention provide one possible explanation for the diverse contrast modulation patterns across layers. NM assumes both stimulus parameters and attention condition to contribute to the normalization input to local excitatory neurons. The stimuli presented in our experiments were optimized for the recording site and did not change with attention condition, and hence are not assumed to contribute differentially to the normalization mechanism. NM also assumes the sizes of attention field of the population to contribute to the normalization input to individual

neurons. The attention field in NM describes the attention gain for each neuron in the

population and depends on the animal's attentional strategy employed during the

experiment (Herrmann et al., 2010). The neural substrate for the attention field is

unspecified in the NM, but we assumed the attention field to be constant across the

cortical depth since the data was collected using a fixed experimental paradigm. However,

given a lack of the biophysical mechanism underlying attentional modulation, our

understanding of the attention field may be subject to future revision. The extent of

excitatory receptive field, also termed as the stimulation field, in the NM can be mediated

by various cortical connectivity patterns. While we explored a lateral pooling mechanism

as the determinant of the receptive field extent of neurons, innervation specificity of

feedforward synaptic input could be an alternative mechanism (Bruno and Simons, 2002;

Hubel and Wiesel, 1962).

The variation in contrast dependence of attentional modulation observed across layers

and cell classes (Figure 3D, E) in our data is explained by the NM in a most parsimonious

way via the variability of the suppressive field size (Figure 4). However, the NM is agnostic

to the neural machinery dedicated to the formation of neuronal tunings or the

implementation of attentional modulation. To explore the implications of its field size

predictions on spike-time correlations in a biophysical model, we considered the model's

stimulation field as the receptive field of putative excitatory projection neurons in a column,

and its suppressive field as the receptive field of local inhibitory interneurons.

We implemented a spiking network model to relate the NM's predictions of variable

suppressive field sizes to variations in spike-time correlations in our data. It is important

to note that our model is not a spiking network implementation of the entirety of attention

computations described by the NM. The suppressive field in NM, which mediates divisive

normalization, is a computation that can be can be implemented through a variety of

mechanisms (see Reynolds and Heeger, 2009 for review). We chose one of the candidate

suppression mechanisms – pooling of lateral inputs by local inhibitory interneurons

(Carandini and Heeger, 1994; Carandini et al., 1997; Troyer et al., 1998). A feedforward

mechanism of variable suppressive fields would yield a very similar prediction for spike-
time correlations between local E and I populations. Our choice was guided by excellent
agreement between the NM model AMI predictions and modulation patterns of related
clusters in the input and deep layers. It is, however, important to note that in the superficial
layers, putative inhibitory neurons (Narrow cluster) lack significant attention modulation
in spite of robust boosting of responses to low contrast stimuli in putative excitatory
neurons (Broad clusters). This does not agree with the predictions of the normalization
model. There are three possible explanations for this observation: 1. Suppressive drive
to broad-spiking neurons in superficial layer is not provided by local inhibitory neurons
within that layer. 2. Superficial layer broad-spiking neurons inherit their contrast
dependent attention modulation from the input layer. 3. Suppressive drive to broad-
spiking neurons in the superficial layer is provided by non-PV local inhibitory neurons
within the layer. Since PV neurons are a majority of the local interneuron population which
itself occupies roughly 20% of the total neural population in the cortex, it is highly possible
that our recordings did not sample the other inhibitory neuronal types. Indeed, studies
from the mouse visual cortex suggest that SOM+ neurons play a key role in mediating
lateral inhibition to layer 2/3 pyramidal neurons (Adesnik et al, 2012). Further studies are
needed to distinguish the contributions of local vs feedforward computations to the
attention effects in superficial layers.

When testing the model's predictions in our dataset, we ascribed the stimulation field to
any of the non-Narrow clusters, including the Broad Regular cluster identified in our layer-
specific CDI analysis (Figure 3E). We ascribed the suppressive field to the receptive field
of the Narrow cluster (putative interneurons). While the experimental data for the Broad
Regular cluster robustly validates the model predictions (Figure 4D), the Broad Bursty
and Medium Regular classes show a comparable trend (Supp. Figure 4D). We could not
perform a robust analysis for the remaining non-Narrow cell classes in a subset of layers
due to a lack of sufficient experimental data (Figure 3E).

Conclusion

Attention increases the signal detection abilities of individual neurons. Whether the

attention mediated firing rate variability is unchanged (McAdams and Maunsell, 1999b)

or reduced (Mitchell et al., 2007), the response gain alone results in improved signal-to-

noise ratio of individual neurons, and enhances the discriminability of the attended signal

(McAdams and Maunsell, 1999b; Verghese, 2001). Attention mediated increases in

neural responses to low- and intermediate-contrast stimuli can extend the separation

between the neuron's stimulus-evoked responses and its spontaneous activity, thereby

improving the neuron's sensitivity to low-contrast stimuli. There has, however, been a

long-standing debate regarding the nature of interactions between attention and visual

scene contrast that mediate object recognition. Previous theoretical studies have sought

to resolve this based on the nature of differences in experimental paradigms (Reynolds

and Heeger, 2009). Our work has exploited advanced experimental techniques to bring

novel understanding of these interactions. Superficial cortical layers in area V4 that

project to higher object recognition stages exhibit enhancement of low contrast stimuli.

Deep layers that project to earlier visual areas exhibit contrast independent attentional

scaling of neuronal responses. By identifying the compartmentalization of attention

modulation among cortical layers, our study has uncovered a new dimension: the nature

of interactions between attention and contrast is aligned with the demands of the visual

processing hierarchy. A previous study has suggested that encoding of scene contrast

and spatial attention by distinct neural populations in area V1 could fulfill its visual

processing demands in the face of contrast dependent attentional feedback (Pooresmaeili

et al., 2010). Our work has revealed an elegant mechanism of meeting these needs via

laminar compartmentalization of attention modulation in area V4 that contributes to this

feedback. Low-frequency synchrony between the thalamus and visual cortex has been

suggested to guide the higher-frequency synchronization of inter-area activity that is

critical to the communication of attention signals between brain areas (Saalmann et al.,

2012). A contrast-independent effect of attention in the deep layer of V4 may also drive

alpha rhythms of pulvino-cortical loops irrespective of stimulus conditions and maintain

the transmission of attentional priorities across the cortex. Future studies are needed to

test these and related hypotheses about the different functional roles of contrast-attention

interactions in different cortical layers.

FIGURE CAPTIONS

Figure 1. Attentional modulation of Contrast Response Function

(A) Orientation change detection task. While the monkey maintained fixation, two oriented

Gabor stimuli were flashed on and off at two locations: one within the RF overlap region of the

recorded V4 column and the other at a location of equal eccentricity across the vertical

meridian. The covert attention of the monkey was cued to one of the two locations. One of the

two stimuli changed its orientation at an unpredictable time. The monkey was rewarded for

making a saccade to the location of orientation change (95% probability of change at the cued

location; 5% probability at uncued location [foil trials]). If no change happened (catch trials), the

monkey was rewarded for maintaining fixation.

(B) An example trial showing the single-unit signals in the attend-in condition. The time axis is

referenced to the appearance of the fixation spot. Spikes (vertical ticks) in each channel come

from the single unit with the highest spike rate in this trial. The gray boxes depict stimulus

presentation epochs. In this particular trial, 8 sample stimuli with different contrasts were

presented, followed by a target stimulus flash with an orientation change that the monkey

responded to correctly. Two different waveforms were shown for two single units.

(C) The mathematical function we used to fit neuronal contrast response functions is shown on

the top. Schematics at the bottom show the effect of positive attentional modulation of each

parameter on the contrast response functions.

(D) The best-fitting contrast response functions of three example neurons in "attend in" and

"attend away" conditions. Mean ± SEM. Insets show the attentional modulation indices

calculated as a function of contrast.

(E) The AMI as a function of contrast for each of the 255 visually responsive single units, with

the three example units in (C) highlighted.

(F) Attention effects on the best-fitting parameters of the contrast response function. Each

histogram plots the AMI distribution of a particular parameter across the population, with the

dashed line marking the 0 modulation and the arrow with a number depicting the median AMI

value. The median AMI is significantly different from zero for all 4 parameters (Mann-Whitney U

test, $p < 0.01$).

(G) The mean difference of AMI from 0 for the 4 parameters are shown in the Cumming

estimation plot. Mean differences are plotted as bootstrap sampling distributions. Each mean

difference is depicted as a dot. Each 95% confidence interval (CI) is indicated by the ends of the

vertical error bars. The faded color represents that the 95% CI include 0.

Figure 2. Classification of Single Units Using Electrophysiological Features.

(A) Mean waveforms for all 410 single units recorded. Waveforms were smoothed using spline

interpolation and their heights were normalized to help compare spike widths.

(B) Distribution of all single units in the space of the first two principal components (PCs) of the

waveforms. The non-Gaussian structure implies that spike shape is a viable feature for

classifying single units.

(C) Histogram of the local variation AMI for all units with available local variation (n=341). The

dashed line marks the 0 AMI value. The arrow depicts the median value of the distribution. The

average local variation of the population is not significantly modulated by attention (Mann-

Whitney U test, $p = 0.37$).

(D) *k*-means clustering of 341 single units based on PTD and spiking variability. Cell classes are

named after their spiking widths (narrow, medium, broad) and their spiking patterns (regular,

bursty). Single units within each range of spike width are highlighted in the component space on

the top. Unclassified units are displayed as black crosses in the feature space.

(E) t-SNE embedding of the same data in (D) in a 2-dimensional space. The number at the left

bottom corner of each panel represents the perplexity parameter of the t-SNE embedding.

(F) The Cumming estimation plot shows the bootstrap sampling distributions of AMIs of CRF

parameters for each cell class. Distributions with CIs including 0 are displayed in faded colors.

The CRF parameters were only available for visually responsive single units.

Figure 3. Contrast Dependency of AMI is Cell-Class and Layer-Specific

(A) The left panel shows the AMI of contrast responses as a function of contrast averaged

across visually responsive single units in each cluster. Mean ± SEM. The black line indicates

the population mean. The right panel shows the mean AMI averaged across contrast for each

cluster. Asterisk indicates either the distribution is significantly different from zero or two

distributions are significantly different (Mann-Whitney U test, $p < 0.05$).

(B) To quantify the contrast dependence of attentional modulation, we averaged the AMI for a

single unit separately within the low-contrast range and the high-contrast range (using the c_{50} as

the low- to high-contrast threshold). We then defined the contrast dependence index (CDI) as

the difference between the average AMI within the low-contrast range and that within the high-

contrast range, normalized by the mean AMI across the whole contrast range. The schematic

shows the interpretation of different ranges of CDI in terms of the AMI.

(C) The Cumming estimation plot shows the raw data of CDIs (left) and the bootstrap sampling

distribution of the mean (right) for each cell class. The plus signs are the outliers within the axis

range, and the arrows depict the outliers outside the axis limit. The number of valid units for

each cell class is shown on the top of the swarm plot. Distributions for cell classes with CIs

inclusive of 0 are shown in faded colors.

(D) Layer-wise AMI (mean ± SEM) of contrast responses for each cell class as a function of

contrast (left) or averaged across contrast (right). Asterisk indicates either the distribution is

significantly different from zero or two distributions are significantly different (Mann-Whitney U

test, p < 0.05). Cell classes that contain fewer than 10 units (including outliers) are excluded.

(E) Layer-wise CDI for five clusters of units, all units, and non-narrow units. The Cumming

estimation plot shows the bootstrap sampling distribution of the mean CDI. Distributions with CIs

inclusive of 0 are illustrated in faded colors. The number of units excluding outliers is shown on

the top of each plot. Distributions for cell classes with CIs inclusive of 0 are shown in faded

colors. For the raw data of the layer-wise CDIs, see Supp. Figure 3B.

Figure 4. Computational Models Provide A Parsimonious Explanation for the Laminar

Profile of AMI Contrast Dependence

(A) Predictions from the normalization model of attention with different suppressive field sizes or

different excitatory (E) receptive field sizes. The top panel shows contrast response functions for

a simulated neuron in the normalization model, when attending to a stimulus within the neuron's

receptive field (black curve) and when attending toward the opposite hemifield (gray curve). The

orange curve represents the AMI. The black dot shows the inflection point of "attend away"

responses that was used to delimit the low- and high-contrast ranges. The middle panel shows

CDIs for simulated neurons as a function of the E receptive field size and the suppressive field

size while holding the stimulus size and the attention field size fixed. The white rectangles depict

a potential mechanism that leads to the observed variation of CDIs across layers (change in

suppressive field size). The black asterisk corresponds to the model parameters used for the

simulation above. The bottom panel shows AMIs of suppressive drive as a function of the E

receptive field size and the suppressive field size. For both simulations, the attention field size is

30 and the stimulus size is 5. The normalization model predicts the AMI of the suppressive drive

to be correlated with the CDI of neuronal responses.

(B) Simulations of a conductance-based E-I network with columnized connections. Schematics

of the E-I networks corresponding to the possible mechanism in (A) are shown on top. 800 E

and 200 I units were evenly distributed in 10 columns around a ring. We interpret the

normalization model's suppressive field as the receptive field of inhibitory neurons in the E-I

network model. E and I units from the same column are mutually coupled. We modeled I

receptive field size as the standard deviations (σ_I) of E-I connections (W_{ie}) across columns

(middle panel, -5 and 5 are the same column). We changed the range of E-I connections across

columns (W_{ie}, shades of green) while keeping other connections the same (gray, including W_{ee},

W_{ii}, W_{ei}). At the bottom, the raster plot shows the spiking activity for all units organized by their

column IDs (blue, I; red, E) in response to a step input. The box depicts a 200 ms window used

for computing cross-correlations between E and I populations.

(C) Cross-correlograms between E and I populations in the same column with different I

receptive field sizes. Cross-correlations were calculated using the pooled spike trains of E units

and I units from the same column across 500 repeats of identical simulation and averaged

across 10 columns. A larger I receptive field reduces the cross-correlation between local E and I

populations. Mean ± SEM.

(D) Cross-correlograms (mean ± SEM) between Narrow and Broad Regular cell classes in the

superficial, input, and deep layer. Cross-correlations were calculated using the pooled spike

trains of Narrow class (putative inhibitory) neurons and Broad Regular class (putative excitatory)

neurons, and were averaged across sessions. The arrows mark 3 time intervals during which

we averaged the cross-correlations and compared the mean differences between the superficial

(or input) and the deep layers. Asterisk: The mean difference of cross-correlations in the center

interval (-75 ms to 75 ms) has a 95% CI above 0. For the estimation plot, see Supp. Figure 4B.

(E) Proposed E-I networks in V4 accounting for the layer-wise CDI variations. The empirical

data and the model simulations imply a larger inhibitory pooling size in the deep layer than

those in the superficial and input layer. The arrows depict the canonical information flow

pathways in a columnar circuit.

Supp. Figure 2. Validations for the Single-unit Classification and the AMI of CRF

Parameters for Each Cell Class

(A) Percentage of classified neurons in the total sample as a function of the number of clusters

(k) input to the k-means clustering algorithm. The 5-cluster result was able to identify the largest

set of distinct clusters while classifying most of the units.

(B) Cross-validation (CV) errors for different numbers of clusters. The 5-cluster result shows the

lowest CV error.

(C) The minimum Jaccard index across clusters for each k from the subsampling analysis. The

analysis was applied to neuronal data from either of the two attention conditions or to combined

data. Clusters that have Jaccard indices above 0.5 are considered as stable.

(D) The cell-wise co-clustering matrix showing the probability of single units belonging to the

same cluster in the subsampling analysis.

(E) In the principal component space of spike shape, we colored single units based on their

spike width range (open circles; narrow, medium, broad). The clusters generated from the peak-

to-trough duration were minimally overlapped.

(F) In the principal component space of spike shape, we colored single units either based on

their spike width range (open circles; narrow, medium, broad) or by running the k-means

clustering algorithm with the first 2 PCs (closed circles). The clusters generated from the k-

means clustering match the ones grouped by the peak-to-trough duration, suggesting that peak-

to-trough duration is an efficient measure to capture the variance in spike shapes.

(G and H) Embedding the data used for the k-means clustering in a 2-dimensional space using

t-SNE (G) or UMAP (H).

(I) Mean firing rate for visually responsive single units split by cell class or by layer. Neuronal

firing rates were calculated from stimulus flashes with the highest common contrast across two

monkey experiments in the "attend away" condition. The number of single units within each

cluster is shown. In each layer, we only analyzed clusters containing more than 10 single units.

Asterisk indicates either the distribution is significantly different from zero or two distributions are

significantly different (Mann-Whitney U test, $p < 0.05$). Mean ± SEM.

(J) The Cumming estimation plot shows the raw data (left) of AMIs of best-fitting CRF

parameters and the bootstrap sampling distribution of each cell class's mean (right).

Supp. Figure 3. Raw Data of AMIs and CDIs of AMI for Each cell class

(A) The AMI of firing rate as a function of contrast for single units within each cell class.

(B) The raw data of cluster-wise CDIs of AMI within each layer. The plus signs are the outliers

within the axis range, and the arrows depict the outliers outside the axis limit. The number of

valid units for each cell class is shown on the top of the swarm plot.

(C) Layer-wise AMI (mean ± SEM) for all units, Narrow unit, and non-narrow units as a function

of contrast (left) or averaged across contrast (right). Asterisk indicates either the distribution is

significantly different from zero or two distributions are significantly different (Mann-Whitney U

test, $p < 0.05$).

Supp. Figure 4. Normalization Model of Attention and CCG Analyses Between Cell

Classes

(A) The structure of the normalization model of attention. The left panel shows a pair of

orientated grating stimuli with identical contrasts, acting as input to the model. The central black

dot indicates the fixation point. The dashed red circle indicates the receptive field of the model

neuron centered on the grating stimulus. The stimulus drive shown in the middle panel is a

collection of neural activity driven by the stimuli. Neurons are arranged based on their receptive

field center (horizontal position) and orientation preference (vertical position). The values of the

stimulus drive are shown by brightness. The top panel shows the attention field as a function of

the receptive field center and the orientation preference. In this case, the attention is guided to

the right stimulus position and does not vary with orientation. Gray areas indicate values of 1,

and white areas indicate values greater than 1. The suppressive drive at the bottom is

calculated from the point-by-point product of the stimulus drive and the attention field and then

pooled over space and orientation according to the suppressive field size. The stimulus drive is

multiplied by the attention field and then divided by the suppressive field to generate the output

firing rates of model neurons (right panel).

(B) i, CDIs for simulated neurons in the normalization model with different stimulus sizes and

attention field sizes. In each panel, we vary the E receptive field size relative to the attention

field size (x-axis), and the suppressive field size relative to the E receptive field size (y-axis).

The pattern of CDI holds for a range of values of stimulus size and attention field size. ii, CDIs

for simulated neurons in the normalization model with different types of inputs. We changed the

stimulus drive input to the normalization model to have either a nonlinear or an attention-

modulated contrast response function. We tested both the response gain (10% increase in

overall response) and the contrast gain (1% of increase in detected contrast) effects. For these

simulations, the attention field size is 30 and the stimulus size is 5. The pattern of CDI holds for

different types of inputs.

(C) Changes in E receptive field size (white box) can also lead to the variation of CDIs across

layers (left panel). We tested this hypothesis in the E-I network by adjusting the standard

deviation of between-column E-E connections (W_{ee}) from narrow (green) to broad (orange) while

keeping other connections the same (gray, including W_{ee}, W_{ii}, W_{ie}) (middle panel). Cross-

correlograms between E and I populations in the same column suggest that different E

receptive field sizes have little impact on the spike-time correlations of local neural activity

across layers (right panel).

(D) Cross-correlograms (mean ± SEM) between Narrow and 3 other cell classes in the

superficial, input, and deep layer. Cross-correlations were calculated using the pooled spike

trains of Narrow class and the other cell class (Board Bursty, Medium Regular, or Medium

Bursty) and were averaged across sessions.

(E) The Cumming estimation plot shows the mean difference for cell-class specific comparisons

of average cross-correlations between the superficial (*Super.*) and deep layers or between the

input and deep layers. We picked 3 time intervals to compute the average cross-correlations

(rows). The raw data of average cross-correlations is plotted on the left in each panel. Each

mean difference between layers is plotted on the right as a bootstrap sampling distribution.

METHODS

Attention Task and Electrophysiological Recording:

Well-isolated single units were recorded from area V4 of two rhesus macaques during an

attention-demanding orientation change detection task (Figure 1A). The task design and

the experimental procedures are described in detail in previous studies (Nandy et al.,

2019; Nandy et al., 2017). While the monkey maintained fixation, two oriented Gabor

stimuli were flashed on for 200 ms and off for variable intervals (randomly chosen

between 200 and 400 ms). The contrast of each stimulus was randomly chosen from a

uniform distribution of 6 contrasts (c = [10%, 18%, 26%, 34%, 42%, and 50%]). One of

the stimuli was located at the receptive field overlap region of the recorded neurons and

the other at an equally eccentric location across the vertical meridian. At the beginning of

a block of trials, the monkey was spatially cued to covertly attend to one of the two spatial

locations using instruction trials in which only one stimulus was presented. One of the two

stimuli changed in orientation at an unpredictable time (minimum 1s, maximum 5s, mean

3s). The monkey was rewarded for making a saccade to the location of orientation change.

95% of the orientation changes occur at the cued location, and 5% occur at the uncued

location (foil trials). We observed impaired performance and slower reaction times for the

foil trials, suggesting that the monkey was indeed using the spatial cue to perform the

task. The difficulty of the task was controlled by changing the degree of orientation change

(randomly chosen from the following: 1°, 2°, 3°, 4°, 6°, 8°, 10°, and 12°). If no change

occurred before 5 s, the monkey was rewarded for holding fixation (catch trial, 13% of

trials).

While the monkey was performing the attention task, we used artificial dura chambers to

facilitate the insertion of 16-channel linear array electrodes ("laminar probes", Plexon,

Plexon V-probe) or single tungsten microelectrodes (FHC Inc) into cortical sites near the

center of the prelunate gyrus. Neuronal signals were recorded, filtered, and stored using

the Multichannel Acquisition Processor system (Plexon). Neuronal signals were classified

as either isolated single units or multiunit clusters by the Plexon Offline Sorter program.

For the data collected from linear array electrodes, we used current source density

analysis (Mitzdorf, 1985) to identify the superficial (Layers 1-3), input (Layer 4), and deep

(Layers 5 and 6) layers of the cortex based on the second derivative of the flash-triggered

LFPs (Bollimunta et al., 2008; Schroeder and Lakatos, 2009; Schroeder et al., 1998;

Nandy et al., 2019; Nandy et al., 2017). Cell bodies of single units with bi-phasic action

potential waveforms were assigned to the same layer in which the electrode channel was

situated during recordings. Units that had tri-phasic waveforms or other shapes were

excluded from analyses. Extracellular data were collected over 32 sessions (23 sessions

in monkey A, 9 in monkey C) using linear array electrodes and 42 sessions (24 sessions

in monkey A, 18 in monkey C) using single tungsten electrodes, yielding 410 single units

in total (337 units using linear array electrodes and 73 units using single tungsten

electrodes). Unit yield per session was considerably higher in monkey C than monkey A,

resulting in a roughly equal contribution of both monkeys toward the population data.

Contrast Response Function (CRF):

Neuronal responses were analyzed only for correctly performed trials, excluding instruction trials. We restricted all data analysis to non-target stimuli because neuronal responses to target stimuli were generally affected by the behavioral response or the reward delivery, which occurs on correct trials after the target's appearance. Moreover, the larger number of non-target stimuli compared to target stimuli provided a more reliable response strength measure. For both attention conditions, the firing rate of a single unit in response to a particular contrast was measured by counting the number of spikes within a period of 60-260 ms after stimulus onset. Its baseline firing rate in each attention condition was extracted from a 200 ms window before a stimulus flash. The mean firing rates and the standard deviations (SDs) were generated across all stimulus flashes. We considered a neuron as visually responsive if any contrast responses exceeded its baseline firing rate by 4 SDs for both attention conditions. We found that 255 of 410 single units were significantly driven by the task stimuli and had valid Lv measures (See *Analysis of Spiking Activity*).

We drew 1000 random samples of contrast responses from a normal distribution with the same mean and standard deviation as the experimental data for each visually responsive single unit. For each attention condition, we computed the CRF for each random sample by applying an ordinary least square fit to a hyperbolic ratio function:

$$r = r_{max} \cdot \frac{c^n}{c^n + c_{50}^n} + m \tag{1}$$

where r is the neuronal response, r_{max} is the maximum attainable response, c is the contrast, c_{50} is the contrast at which response is half-maximal, m is the baseline activity, and n describes the steepness of the response function and represents the neuron's sensitivity to contrast. This function has been shown to provide a good fit to contrast response functions from visual cortices in cat and macaque monkey (Albrecht and Hamilton, 1982; Williford and Maunsell, 2006). We then averaged the best-fitting CRFs

882   across random samples to generate the mean CRF for each visually responsive single

883   unit (Figure 1D).

885   *Analysis of Spiking Activity*:

886   For every single unit, the spiking variability was measured by the local variation (Lv),

887   which quantifies the average differences between consecutive inter-spike intervals (ISIs).

889
$$Lv = \frac{3}{N-2} \sum_{i=1}^{N-1} \frac{(\Delta t_i - \Delta t_{i+1})^2}{(\Delta t_i + \Delta t_{i+1})^2} \qquad (2)$$

where Δt_i is a given ISI and N represents the total number of spikes within the time

window. The advantage of Lv over other spiking measures such as the Fano factor and

coefficient of variation is that it is more robust to changes in firing rate (Shinomoto et al.,

2003). We computed each unit's Lv using its spike train during a stimulus flash and

averaged across all flashes (restricted to non-target stimuli).

For completely Poisson processes (where neuronal firing rates are fixed and spike times

are random) the Lv is 1, whereas more regular activity takes values significantly lower

than 1, and bursty spiking takes values significantly larger than 1.

Of 410 single units, we included 341 neurons with enough spikes to compute Lv for

further clustering analysis.

Clustering Analysis:

We used the *k*-means clustering algorithm (Hartigan and Wong, 1979) to characterize

cell classes upon the space of peak-to-trough duration (PTD) and Lv. To estimate a range

of the number of clusters, we used a set of indices that evaluate the quality of clustering

(Halkidi et al., 2001; Jain and Dubes, 1988; Milligan and Cooper, 1985; Vendramin et al.,

2010): Rand, Mirkin, Hubert, Silhouette, Davies–Bouldin, Calinski–Harabasz, Hartigan,

Homogeneity and Separation indices. We ran 50 replicates of the *k*-means clustering for

different numbers of clusters, from *k* = 1 to *k* = 40. For each *k*, we selected the best

replicate according to the minimum squared Euclidean distance from all cluster elements

to their respective centroids. We also ran 10 identical realizations, each with a random

set of initial centroids to exclude the initialization issues. We evaluated validation indices

for each realization, and due to random initializations, most validation indices showed

increased variability after saturation, suggesting excessive partitions in the clustering

process. Based on this method, a range of 2 to 10 clusters was shown to be proper for

our dataset.

We then used a meta-clustering analysis (Ardid et al., 2015) to select the most appropriate

number of clusters: we ran 500 realizations of the k-means for each k and selected the

best replicate from 50 replicates for each realization. After 500 realizations of each k, we

computed the probability that pairs of neurons belonged to a same cluster. Valid clusters

were identified by setting a probability threshold ($p \geq 0.9$). We considered clusters with at

least five single units as reliable. We identified the most appropriate number of clusters

($k = 5$) as the largest number of reliable cell classes that classified the most neurons in

the dataset (Supp. Figure 2A).

Clustering Validation:

We validated our clustering analysis in three ways. First, we applied a data-driven

approach based on a form of cross-validation (Fu and Perry, 2020). We organized our

data into a matrix with each row representing a single unit and each column representing

a feature for clustering. We then randomly partition the rows and columns into K and L

subsets, respectively. Each fold is represented by a pair (r, s) of integers, with $r \in$

$\{1, ..., K\}$ and $s \in \{1, ..., L\}$. Fold (r, s) treats the rth row subset as "test" observations,

and the sth column subset as "responses". The remaining $(K - 1)$ row subsets are "train"

observations, and the $(L - 1)$ column subsets are "predictors". For our dataset, we take

$K = 5$ and $L = 2$. We applied the same clustering procedures described above to the

"responses" data of "train" observations to generate the cluster labels and cluster means

for "train" observations. Then, we trained a linear discriminant analysis classifier with

equal class priors to predict those cluster labels from the "predictors" data of "train"

observations. The classifier was then applied to the "predictors" data of "test"

observations to generate their predicted cluster labels as well as predicted cluster means.

The cross-validation error was then computed by averaging the squared differences
between the "responses" of "test" observations and their predicted cluster means. Using
such a method, we calculated the cross-validation error for each k (from $k = 2$ to $k = 10$)
in the k-means clustering results (Supp. Figure 2B), and $k = 5$ showed the lowest cross-
validation error.

Second, we validated the stability of our clustering analysis by subsampling analysis
(Hennig, 2007). We generated 100 random subsamples containing 90% of the trials from
"attend in" or "attend away" or both conditions. We computed the Lv for every single unit
in subsamples. Random subsamples were then clustered by the k-means algorithm with
k from 3 to 10. The Jaccard similarities were calculated between original clusters and
clusters from the subsample, and the maximum was found for each original cluster. These
Jaccard similarities were averaged across all subsample runs. Clusters with average
Jaccard similarities below 0.5 were thought to be unstable. We reported the minimum
Jaccard similarity across original clusters for each k (Supp. Figure 2C), and all clusters
when $k = 5$ were stable. A cell-wise co-clustering matrix was also generated during this
procedure (Supp. Figure 2D), and it also supported our estimation of cluster stability.

Third, we used dimensionality reduction techniques to deal with the concern that cell
classes in our dataset may not be separable by the linear combinations of the two features
we used as input to perform k-means clustering. We applied both the t-distributed
stochastic neighbor embedding (t-SNE; Hinton and Roweis, 2003) algorithm and the
uniform manifold approximation and projection (UMAP; McInnes et al., 2018) algorithm
to our singe-unit data. Within a range of both algorithms' critical parameters, we find that
the clusters from k-means clustering were still well separated (Supp. Figure 2G, H).

Attentional Modulation Index and its Contrast Dependency:

The attentional modulation index (AMI) of a neuron during the stimulus presentation with
a specific contrast c was calculated using the best-fitting contrast response functions (r)
from both attention conditions:

$$AMI(c) = \frac{r(c)^{IN} - r(c)^{AWAY}}{r(c)^{IN} + r(c)^{AWAY}} \tag{3}$$

The contrast dependence of the AMI was measured by the contrast dependence index (CDI):

$$CDI = \frac{\overline{AMI_{low}} - \overline{AMI_{high}}}{|\overline{AMI_{all}}|} \tag{4}$$

where $\overline{AMI_{low}}$ and $\overline{AMI_{high}}$ are the average AMIs within the low-contrast range and the high-contrast range, respectively. $\overline{AMI_{all}}$ is the average AMI across all contrasts. c_{50} from the best-fitting CRF during "attend away" condition delimited the range of low contrast ($c < c_{50}$) and the range of high contrast ($c \geq c_{50}$). CDI measures how the AMI of a neuron fluctuates with the contrast of the stimulus. A zero CDI indicates that the AMI is independent of the contrast of the stimulus. More robust attentional modulation at the low-contrast range leads to positive CDIs, and more potent attention effects at the high-contrast range result in negative CDIs (Figure 3B). AMI and CDI were only calculated for those visually responsive neurons whose laminar locations were identified (n = 255).

Normalization Model Simulations:

We used the normalization model of attention (Reynolds and Heeger, 2009) to explore the neural mechanisms behind the variety of attentional modulation across layers (Supp. Figure 4A). The normalization model posits that the resulting firing rate (R) of the population of simulated neurons can be produced from a function of the stimulus drive (E), the attention field (A), and the suppressive drive (S):

$$R(c; x, \theta) = \frac{A(x, \theta)E(x, \theta; c)}{S(x, \theta; c) + \sigma} \tag{5}$$

where x and θ represent the receptive field center and orientation preference of each neuron in the population. c is stimulus contrast and σ is a constant that controls the contrast gain of the neurons' response. The stimulus drive is derived from the stimulus

and the stimulation field of the model neuron, which is its receptive field in the spatial and

orientational space. The attention field describes the strength of attentional modulation

as a function of receptive field center and orientation preference. The attentional

modulation is 1 for unattended space and is greater than 1 for a small range of locations

around the attended stimulus. We computed the suppressive drive by pooling the product

of the stimulus drive and the attention field over spatial positions and orientations:

1004

1005
$$S(x, \theta; c) = s(x, \theta) * [A(x, \theta)E(x, \theta; c)] \qquad (6)$$

1006

1007 where $s(x, \theta)$ is the suppressive field and $*$ represents convolution. The stimulus,

1008 stimulation field (excitatory receptive field), attention field, and suppressive field all had

1009 Gaussian profile in space and orientation.

1010

For simulations in Figure 4A, the stimulus size was 5 and the attention field size was 30.

The CDI pattern holds for a range of stimulus sizes and attention field sizes (Supp. Figure

4B). The excitatory receptive field size and the suppressive field size were varied

according to their ratios relative to the attention field size. For a given pair of stimulus size

and attention field size, we changed the ratio of the attention field size to the excitatory

receptive field size from 0.5 to 3 and the ratio of the suppressive field size to the excitatory

receptive field size from 1 to 6. The orientation tuning width of the excitatory receptive

field was 60°, and the suppressive field was nonspecific. A baseline activity of 0.5 was

added after the normalization. For each combination of parameters, the AMIs were

calculated using the model neuron responses from two attention conditions. The CDIs of

AMIs were computed from the average AMIs within the low-contrast range and the high-

contrast range delimited by the CRF's inflection point from the "attend away" condition.

For simulations in Supp. Figure 4C, we further modified the stimulus drive of the model

to have either a nonlinear or an attention-modulated contrast response function. The

nonlinear function was implemented as

1026
$$r(c) = \frac{c}{c + \sigma}$$

1027 where σ is 0.26, matching the average c_{50} of our data. We also applied either a
1028 multiplicative response gain (10% of increase in overall response) or a contrast gain (1%
1029 of increase in perceived contrast) to test the effects of different attention modulation of
1030 inputs on the model neurons' responses.

1031

1032 *Computational Model*:

1033 We set up a conductance-based model of N_E excitatory (E) and N_I inhibitory (I) neurons
1034 with a connection probability of 0.5 (Figure 4B). Neurons were evenly divided into 10
1035 columns or local E-I sub-networks around a ring with the following within-column synaptic
1036 weights:

1037

$$1038 \quad \text{E to E}: w_{EE} = \frac{W_{EE}}{N_E}; \text{ I to I}: w_{II} = \frac{W_{II}}{N_I}; \text{ E to I}: w_{IE} = \frac{W_{IE}}{N_E}; \text{ I to E}: w_{EI} = \frac{W_{EI}}{N_I}$$

1039

1040 We only modeled E to I connections and E to E connections between different columns.
1041 The synaptic weights fell off with column distance following a Gaussian profile:

1042

$$w^{ij} = \frac{W}{N_E} \times \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2}\left(\frac{d_{ij}}{\sigma}\right)^2\right) \tag{7}$$

1044

where w^{ij} is the synaptic weight between two columns (w_{IE}^{ij} or w_{EE}^{ij}) and d_{ij} represents
the distance from column j to column i. σ controls the pooling size of the postsynaptic
inhibitory (σ_I) or excitatory (σ_E) neuron.

We simulated models of $N_E = 800$ excitatory and $N_I = 200$ inhibitory spiking units. The
spiking units were modeled as Izhikevich neurons (Izhikevich, 2003) with the following
dynamics:

1051

$$\frac{dv}{dt} = 0.04v^2 + 5v + 140 - u + I \tag{8}$$

1053

$$\frac{du}{dt} = a(bv - a) \tag{9}$$

$$if \; v \geq 30 \; mV, then \begin{cases} v \leftarrow c \\ u \leftarrow u + d \end{cases} \tag{10}$$

v represents the membrane potential of the neuron and u is a membrane recovery variable. I is the current input to the neuron (synaptic and injected DC currents). The parameters a, b, c, and d determine intrinsic firing patterns and were chosen as follows:

Regular spiking excitatory units: $a = 0.02, b = 0.2, c = -65, d = 8$

Fast spiking inhibitory units: $a = 0.1, b = 0.2, c = -65, d = 2$

Presynaptic excitatory neurons generate fast (AMPA) and slow (NMDA) synaptic currents, while presynaptic inhibitory neurons generate fast GABA currents:

$$I_{syn} = \sum_i g_{AMPA}(t)(v(t) - V_{AMPA}) + \sum_j g_{NMDA}(t)(v(t) - V_{NMDA})$$
$$+ \sum_k g_{GABA}(t)(v(t) - V_{GABA}) \tag{11}$$

where $V_{AMPA} = 0$, $V_{NMDA} = 0$, $V_{GABA} = -70$ are the respective reversal potentials (mV). The synaptic time course g(t) was modeled as a difference between exponentials:

$$g(t) = \frac{1}{\tau_d - \tau_r}\left[exp\left(-\frac{t - \tau_l}{\tau_d}\right) - exp\left(-\frac{t - \tau_l}{\tau_r}\right) \right] \tag{12}$$

where the parameters τ_d, τ_r, and τ_l are the decay, rise, and latency time constants with the following values (Brunel and Wang, 2003): AMPA: $\tau_d = 2$ ms, $\tau_r = 0.5$ ms, $\tau_l = 1$ ms; NMDA: $\tau_d = 80$ ms, $\tau_r = 2$ ms, $\tau_l = 1$ ms; GABA: $\tau_d = 5$ ms, $\tau_r = 0.5$ ms, $\tau_l = 1$ ms; The AMPA to NMDA ratio is 0.45 (Myme et al., 2003).

We simulated the network with a DC step current ($I_{DC} = 4$) of duration 1.2 s. Synaptic

noise was sampled from a normal distribution ($I_{syn-noise} \sim \mathcal{N}(\mu = 0, \sigma = 3)$). We pooled

over spike trains of excitatory units and inhibitory units in each column separately and

calculated the shuffled-corrected jittered cross-correlations from the two population spike

trains binned at 1 ms within the 200 ms time window (800-1000 ms) after the initial

transient response across 500 repeats of the simulation. Cross-correlations for different

choices of σ_I or σ_E were reported as the average across columns (Figure 4C) (Harrison

et al., 2007; Harrison and Geman, 2009).

1088

Spike Train Cross-correlations:

The population cross-correlograms in Figure 4 report shuffled-corrected jittered cross-

correlations (Harrison et al., 2007; Harrison and Geman, 2009). We computed the jittered

cross-correlations by resampling two spike trains within a specific time window such that

for each spike in the original data, a spike is chosen at random with replacement from

within the same time window across trials, thus preserving the PSTH at the resolution of

the jitter window. We computed the jittered cross-correlations with 4, 8, and 16 jitter

windows, and the results of 8 jitter windows were shown. Shuffled cross-correlations were

calculated by cross-correlating the first population spike train with the randomly permuted

second population spike train. Both types of cross-correlations were averaged across

trials and were further normalized by the geometric mean of the two spike trains' firing

rates and a triangular function that corrects for the amount of overlap for the different lags.

The normalized shuffled cross-correlation was then subtracted from the normalized

jittered cross-correlation to produce the shuffled-corrected jittered cross-correlation.

1103

1104   **AUTHOR CONTRIBUTIONS**

1105   MPJ & ASN conceptualized the project. XW analyzed the data, previously collected by

1106   ASN, and performed the computational modeling. MPJ supervised the project. XW, MPJ

1107   and ASN wrote the manuscript.

ACKNOWLEDGEMENTS

This research was supported by NIH R00 EY025026 to MPJ, NARSAD Young Investigator Grant, Ziegler Foundation Grant and Yale Orthwein Scholar Funds to ASN, and by NEI core grant for vision research P30 EY026878 to Yale University.

SUPPLEMENTARY MATERIAL

Figures S2-S4

REFERENCES

Adesnik, H., and Naka, A. (2018). Cracking the Function of Layers in the Sensory Cortex. Neuron *100*, 1028-1043.

Adesnik, H., and Scanziani, M. (2010). Lateral competition for cortical space by layer-specific horizontal circuits. Nature *464*, 1155-1160.

Albrecht, D.G., and Hamilton, D.B. (1982). Striate cortex of monkey and cat: contrast response function. J Neurophysiol *48*, 217-237.

Anderson, E.B., Mitchell, J.F., and Reynolds, J.H. (2011). Attentional modulation of firing rate varies with burstiness across putative pyramidal neurons in macaque visual area V4. J Neurosci *31*, 10983-10992.

Anderson, J.C., and Martin, K.A. (2006). Synaptic connection from cortical area V4 to V2 in macaque monkey. J Comp Neurol *495*, 709-721.

Ardid, S., Vinck, M., Kaping, D., Marquez, S., Everling, S., and Womelsdorf, T. (2015). Mapping of Functionally Characterized Cell Classes onto Canonical Circuit Operations in Primate Prefrontal Cortex. The Journal of Neuroscience *35*, 2975.

Avidan, G., Harel, M., Hendler, T., Ben-Bashat, D., Zohary, E., and Malach, R. (2002). Contrast sensitivity in human visual areas and its relationship to object recognition. J Neurophysiol *87*, 3102-3116.

Bisley, J.W., and Goldberg, M.E. (2003). Neuronal activity in the lateral intraparietal area and spatial attention. Science *299*, 81-86.

Bollimunta, A., Chen, Y., Schroeder, C.E., and Ding, M. (2008). Neuronal Mechanisms of Cortical Alpha Oscillations in Awake-Behaving Macaques. The Journal of Neuroscience *28*, 9976.

Borra, E., Ichinohe, N., Sato, T., Tanifuji, M., and Rockland, K.S. (2010). Cortical connections to area TE in monkey: hybrid modular and distributed organization. Cereb Cortex *20*, 257-270.

Brunel, N., and Wang, X.J. (2003). What determines the frequency of fast network oscillations with irregular neural discharges? I. Synaptic dynamics and excitation-inhibition balance. J Neurophysiol *90*, 415-430.

Bruno, R.M., and Simons, D.J. (2002). Feedforward mechanisms of excitatory and inhibitory cortical receptive fields. J Neurosci *22*, 10966-10975.

Callaway, E.M. (1998). Local circuits in primary visual cortex of the macaque monkey.
Annu Rev Neurosci *21*, 47-74.
Carandini, M. (2004). Receptive Fields and Suppressive Fields in the Early Visual System.
Cognitive Neurosciences Iii, Third Edition *3*, 313-326.
Carandini, M., and Heeger, D.J. (1994). Summation and division by neurons in primate
visual cortex. Science *264*, 1333-1336.
Carandini, M., Heeger, D.J., and Movshon, J.A. (1997). Linearity and normalization in
simple cells of the macaque primary visual cortex. J Neurosci *17*, 8621-8644.
Carrasco, M., Ling, S., and Read, S. (2004). Attention alters appearance. Nat Neurosci
7, 308-313.
Cohen, M.R., and Maunsell, J.H. (2009). Attention improves performance primarily by
reducing interneuronal correlations. Nat Neurosci *12*, 1594-1600.
Compte, A., Constantinidis, C., Tegner, J., Raghavachari, S., Chafee, M.V., Goldman-
Rakic, P.S., and Wang, X.J. (2003). Temporally irregular mnemonic persistent activity in
prefrontal neurons of monkeys during a delayed response task. J Neurophysiol *90*, 3441-
3454.
Connors, B.W., and Gutnick, M.J. (1990). Intrinsic firing patterns of diverse neocortical
neurons. Trends Neurosci *13*, 99-104.
Constantinidis, C., and Goldman-Rakic, P.S. (2002). Correlated discharges among
putative pyramidal neurons and interneurons in the primate prefrontal cortex. J
Neurophysiol *88*, 3487-3497.
Csicsvari, J., Hirase, H., Czurko, A., Mamiya, A., and Buzsaki, G. (1999). Oscillatory
coupling of hippocampal pyramidal cells and interneurons in the behaving Rat. J Neurosci
19, 274-287.
D'Souza, R.D., and Burkhalter, A. (2017). A Laminar Organization for Selective Cortico-
Cortical Communication. Front Neuroanat *11*, 71.
Degenetais, E., Thierry, A.M., Glowinski, J., and Gioanni, Y. (2002). Electrophysiological
properties of pyramidal neurons in the rat prefrontal cortex: an in vivo intracellular
recording study. Cereb Cortex *12*, 1-16.
Diester, I., and Nieder, A. (2008). Complementary contributions of prefrontal neuron
classes in abstract numerical categorization. J Neurosci *28*, 7737-7747.
Distler, C., Boussaoud, D., Desimone, R., and Ungerleider, L.G. (1993). Cortical
connections of inferior temporal area TEO in macaque monkeys. J Comp Neurol *334*,
125-150.
Douglas, R.J., and Martin, K.A. (1991). A functional microcircuit for cat visual cortex. J
Physiol *440*, 735-769.
Douglas, R.J., and Martin, K.A. (2004). Neuronal circuits of the neocortex. Annu Rev
Neurosci *27*, 419-451.
Felleman, D.J., and Van Essen, D.C. (1991). Distributed hierarchical processing in the
primate cerebral cortex. Paper presented at: Cereb cortex (Citeseer).
Fioravanti, M., Carlone, O., Vitale, B., Cinti, M.E., and Clare, L. (2005). A meta-analysis
of cognitive deficits in adults with a diagnosis of schizophrenia. Neuropsychol Rev *15*, 73-
95.

Fox, S.E., and Ranck, J.B., Jr. (1981). Electrophysiological characteristics of hippocampal
complex-spike cells and theta cells. Exp Brain Res *41*, 399-410.
Frank, L.M., Brown, E.N., and Wilson, M.A. (2001). A comparison of the firing properties
of putative excitatory and inhibitory neurons from CA1 and the entorhinal cortex. J
Neurophysiol *86*, 2029-2040.
Fu, W., and Perry, P.O. (2020). Estimating the Number of Clusters Using Cross-Validation.
Journal of Computational and Graphical Statistics *29*, 162-173.
Gattass, R., Galkin, T.W., Desimone, R., and Ungerleider, L.G. (2014). Subcortical
connections of area V4 in the macaque. J Comp Neurol *522*, 1941-1965.
Ghose, G.M., and Maunsell, J.H. (2008). Spatial summation can explain the attentional
modulation of neuronal responses to multiple stimuli in area V4. J Neurosci *28*, 5115-
5126.
Gilbert, C.D. (1977). Laminar differences in receptive field properties of cells in cat
primary visual cortex. J Physiol *268*, 391-421.
Gilbert, C.D., and Wiesel, T.N. (1985). Intrinsic connectivity and receptive field properties
in visual cortex. Vision Res *25*, 365-374.
Gouwens, N.W., Sorensen, S.A., Berg, J., Lee, C., Jarsky, T., Ting, J., Sunkin, S.M., Feng,
D., Anastassiou, C.A., Barkan, E.*, et al.* (2019). Classification of electrophysiological and
morphological neuron types in the mouse visual cortex. Nat Neurosci *22*, 1182-1195.
Halkidi, M., Batistakis, Y., and Vazirgiannis, M. (2001). On Clustering Validation
Techniques. Journal of Intelligent Information Systems *17*, 107-145.
Harrison, M., Amarasingham, A., and Geman, S. (2007). Jitter methods for investigating
spike train dependencies. Computational and Systems Neuroscience Abstracts III-17.
Harrison, M.T., and Geman, S. (2009). A rate and history-preserving resampling algorithm
for neural spike trains. Neural Comput *21*, 1244-1258.
Hartigan, J.A., and Wong, M.A. (1979). Algorithm AS 136: A K-Means Clustering
Algorithm. Journal of the Royal Statistical Society Series C (Applied Statistics) *28*, 100-
108.
Hawken, M.J., Shapley, R.M., Disney, A.A., Garcia-Marin, V., Henrie, A., Henry, C.A.,
Johnson, E.N., Joshi, S., Kelly, J.G., Ringach, D.L.*, et al.* (2020). Functional Clusters of
Neurons in Layer 6 of Macaque V1. J Neurosci *40*, 2445-2457.
Hennig, C. (2007). Cluster-wise assessment of cluster stability. Computational Statistics
& Data Analysis *52*, 258-271.
Herrmann, K., Montaser-Kouhsari, L., Carrasco, M., and Heeger, D.J. (2010). When size
matters: attention affects performance by contrast or response gain. Nat Neurosci *13*,
1554-1559.
Hinton, G.E., and Roweis, S.T. (2003). Stochastic neighbor embedding. Paper presented
at: Advances in neural information processing systems.
Hirsch, J.A., and Gilbert, C.D. (1991). Synaptic physiology of horizontal connections in
the cat's visual cortex. J Neurosci *11*, 1800-1809.
Hirsch, J.A., and Martinez, L.M. (2006). Laminar processing in the visual cortical column.
Curr Opin Neurobiol *16*, 377-384.
Huang, L., and Dobkins, K.R. (2005). Attentional effects on contrast discrimination in
humans: evidence for both contrast gain and response gain. Vision Res *45*, 1201-1212.

Hubel, D.H., and Wiesel, T.N. (1962). Receptive fields, binocular interaction and
functional architecture in the cat's visual cortex. J Physiol *160*, 106-154.

Hussar, C.R., and Pasternak, T. (2009). Flexibility of sensory representations in prefrontal
cortex depends on cell type. Neuron *64*, 730-743.

Izhikevich, E.M. (2003). Simple model of spiking neurons. IEEE Trans Neural Netw *14*,
1569-1572.

Jain, A.K., and Dubes, R.C. (1988). Algorithms for clustering data (Prentice-Hall, Inc.).

Johnston, K., DeSouza, J.F., and Everling, S. (2009). Monkey prefrontal cortical
pyramidal and putative interneurons exhibit differential patterns of activity between
prosaccade and antisaccade tasks. J Neurosci *29*, 5516-5524.

Kastner, S., and Ungerleider, L.G. (2000). Mechanisms of visual attention in the human
cortex. Annu Rev Neurosci *23*, 315-341.

Kaufman, M.T., Churchland, M.M., Santhanam, G., Yu, B.M., Afshar, A., Ryu, S.I., and
Shenoy, K.V. (2010). Roles of monkey premotor neuron classes in movement preparation
and execution. J Neurophysiol *104*, 799-810.

Kouh, M., and Poggio, T. (2008). A canonical neural circuit for cortical nonlinear
operations. Neural Comput *20*, 1427-1451.

Li, X., and Basso, M.A. (2008). Preparing to move increases the sensitivity of superior
colliculus neurons. J Neurosci *28*, 4561-4577.

Li, X., Lu, Z.L., Tjan, B.S., Dosher, B.A., and Chu, W. (2008). Blood oxygenation level-
dependent contrast response functions identify mechanisms of covert attention in early
visual areas. Proc Natl Acad Sci U S A *105*, 6202-6207.

Martinez-Trujillo, J., and Treue, S. (2002). Attentional modulation strength in cortical area
MT depends on stimulus contrast. Neuron *35*, 365-370.

McAdams, C.J., and Maunsell, J.H. (1999a). Effects of attention on orientation-tuning
functions of single neurons in macaque cortical area V4. J Neurosci *19*, 431-441.

McAdams, C.J., and Maunsell, J.H. (1999b). Effects of attention on the reliability of
individual neurons in monkey visual cortex. Neuron *23*, 765-773.

McCormick, D.A., Connors, B.W., Lighthall, J.W., and Prince, D.A. (1985). Comparative
electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. J
Neurophysiol *54*, 782-806.

McInnes, L., Healy, J., and Melville, J. (2018). Umap: Uniform manifold approximation
and projection for dimension reduction. arXiv preprint arXiv:180203426.

McIntyre, R.S., Kennedy, S.H., Soczynska, J.K., Nguyen, H.T., Bilkey, T.S.,
Woldeyohannes, H.O., Nathanson, J.A., Joshi, S., Cheng, J.S., Benson, K.M.*, et al.*
(2010). Attention-deficit/hyperactivity disorder in adults with bipolar disorder or major
depressive disorder: results from the international mood disorders collaborative project.
Prim Care Companion J Clin Psychiatry *12*, PCC.09m00861.

Mehta, A.D., Ulbert, I., and Schroeder, C.E. (2000). Intermodal selective attention in
monkeys. II: physiological mechanisms of modulation. Cereb Cortex *10*, 359-370.

Migliore, M., and Shepherd, G.M. (2005). Opinion: an integrated approach to classifying
neuronal phenotypes. Nat Rev Neurosci *6*, 810-818.

Milligan, G.W., and Cooper, M.C. (1985). An examination of procedures for determining
the number of clusters in a data set. Psychometrika *50*, 159-179.

Mitchell, J.F., Sundberg, K.A., and Reynolds, J.H. (2007). Differential attention-
dependent response modulation across cell classes in macaque visual area V4. Neuron
55, 131-141.
Mitchell, J.F., Sundberg, K.A., and Reynolds, J.H. (2009). Spatial attention decorrelates
intrinsic activity fluctuations in macaque area V4. Neuron *63*, 879-888.
Mitzdorf, U. (1985). Current source-density method and application in cat cerebral cortex:
investigation of evoked potentials and EEG phenomena. Physiological Reviews *65*, 37-
100.
Moore, T., and Zirnsak, M. (2017). Neural Mechanisms of Selective Visual Attention.
Annu Rev Psychol *68*, 47-72.
Moran, J., and Desimone, R. (1985). Selective attention gates visual processing in the
extrastriate cortex. Science *229*, 782-784.
Morrone, M.C., Denti, V., and Spinelli, D. (2002). Color and luminance contrasts attract
independent attention. Curr Biol *12*, 1134-1137.
Motter, B.C. (1993). Focal attention produces spatially selective processing in visual
cortical areas V1, V2, and V4 in the presence of competing stimuli. J Neurophysiol *70*,
909-919.
Myme, C.I., Sugino, K., Turrigiano, G.G., and Nelson, S.B. (2003). The NMDA-to-AMPA
ratio at synapses onto layer 2/3 pyramidal neurons is conserved across prefrontal and
visual cortices. J Neurophysiol *90*, 771-779.
Naka, A., and Adesnik, H. (2016). Inhibitory Circuits in Cortical Layer 5. Front Neural
Circuits *10*, 35.
Nandy, A., Nassi, J.J., Jadi, M.P., and Reynolds, J. (2019). Optogenetically induced low-
frequency correlations impair perception. Elife *8*, e35123.
Nandy, A.S., Nassi, J.J., and Reynolds, J.H. (2017). Laminar Organization of Attentional
Modulation in Macaque Visual Area V4. Neuron *93*, 235-246.
Neuchterlein, K.H., Dawson, M.E., Ventura, J., Miklowitz, D., and Konishi, G. (1991).
Information-processing anomalies in the early course of schizophrenia and bipolar
disorder. Schizophrenia Research *5*, 195-196.
Nowak, L.G., Azouz, R., Sanchez-Vives, M.V., Gray, C.M., and McCormick, D.A. (2003).
Electrophysiological classes of cat primary visual cortical neurons in vivo as revealed by
quantitative analyses. J Neurophysiol *89*, 1541-1566.
Olsen, S.R., Bortone, D.S., Adesnik, H., and Scanziani, M. (2012). Gain control by layer
six in cortical circuits of vision. Nature *483*, 47-52.
Ostojic, S., Brunel, N., and Hakim, V. (2009). How connectivity, background activity, and
synaptic properties shape the cross-correlation between spike trains. J Neurosci *29*,
10234-10253.
Pestilli, F., Ling, S., and Carrasco, M. (2009). A population-coding model of attention's
influence on contrast response: Estimating neural effects from psychophysical data.
Vision Res *49*, 1144-1153.
Petersen, S.E., and Posner, M.I. (2012). The attention system of the human brain: 20
years after. Annu Rev Neurosci *35*, 73-89.
Pooresmaeili, A., Poort, J., Thiele, A., and Roelfsema, P.R. (2010). Separable codes for
attention and luminance contrast in the primary visual cortex. J Neurosci *30*, 12701-12711.

Povysheva, N.V., Gonzalez-Burgos, G., Zaitsev, A.V., Kroner, S., Barrionuevo, G., Lewis,
D.A., and Krimer, L.S. (2006). Properties of excitatory synaptic responses in fast-spiking
interneurons and pyramidal cells from monkey and rat prefrontal cortex. Cereb Cortex *16*,
541-552.
Rao, R.P., and Ballard, D.H. (1999). Predictive coding in the visual cortex: a functional
interpretation of some extra-classical receptive-field effects. Nat Neurosci *2*, 79-87.
Rao, S.G., Williams, G.V., and Goldman-Rakic, P.S. (1999). Isodirectional tuning of
adjacent interneurons and pyramidal cells during working memory: evidence for
microcolumnar organization in PFC. J Neurophysiol *81*, 1903-1916.
Reynolds, J.H., and Chelazzi, L. (2004). Attentional modulation of visual processing.
Annu Rev Neurosci *27*, 611-647.
Reynolds, J.H., Chelazzi, L., and Desimone, R. (1999). Competitive mechanisms
subserve attention in macaque areas V2 and V4. J Neurosci *19*, 1736-1753.
Reynolds, J.H., and Heeger, D.J. (2009). The normalization model of attention. Neuron
61, 168-185.
Reynolds, J.H., Pasternak, T., and Desimone, R. (2000). Attention increases sensitivity
of V4 neurons. Neuron *26*, 703-714.
Rockland, K.S., and Pandya, D.N. (1979). Laminar origins and terminations of cortical
connections of the occipital lobe in the rhesus monkey. Brain Res *179*, 3-20.
Roe, A.W., Chelazzi, L., Connor, C.E., Conway, B.R., Fujita, I., Gallant, J.L., Lu, H., and
Vanduffel, W. (2012). Toward a unified theory of visual area V4. Neuron *74*, 12-29.
Rolls, E.T., and Baylis, G.C. (1986). Size and contrast have only small effects on the
responses to faces of neurons in the cortex of the superior temporal sulcus of the monkey.
Exp Brain Res *65*, 38-48.
Saalmann, Y.B., Pinsk, M.A., Wang, L., Li, X., and Kastner, S. (2012). The pulvinar
regulates information transmission between cortical areas based on attention demands.
Science *337*, 753-756.
Schroeder, C.E., and Lakatos, P. (2009). Low-frequency neuronal oscillations as
instruments of sensory selection. Trends in Neurosciences *32*, 9-18.
Schroeder, C.E., Mehta, A.D., and Givre, S.J. (1998). A spatiotemporal profile of visual
system activation revealed by current source density analysis in the awake macaque.
Cerebral Cortex *8*, 575-592.
Shinomoto, S., Kim, H., Shimokawa, T., Matsuno, N., Funahashi, S., Shima, K., Fujita, I.,
Tamura, H., Doi, T., Kawano, K.*, et al.* (2009). Relating neuronal firing patterns to
functional differentiation of cerebral cortex. PLoS Comput Biol *5*, e1000433.
Shinomoto, S., Shima, K., and Tanji, J. (2003). Differences in Spiking Patterns Among
Cortical Neurons. Neural Computation *15*, 2823-2842.
Simons, D.J. (1978). Response properties of vibrissa units in rat SI somatosensory
neocortex. J Neurophysiol *41*, 798-820.
Sur, M., Garraghty, P.E., and Bruce, C.J. (1985). Somatosensory cortex in macaque
monkeys: laminar differences in receptive field size in areas 3b and 1. Brain Res *342*,
391-395.
Swadlow, H.A. (2003). Fast-spike interneurons and feedforward inhibition in awake
sensory neocortex. Cereb Cortex *13*, 25-32.

Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J.,
Garren, E., Economo, M.N., Viswanathan, S., *et al.* (2018). Shared and distinct
transcriptomic cell types across neocortical areas. Nature *563*, 72-78.
Treue, S., and Martinez Trujillo, J.C. (1999). Feature-based attention influences motion
processing gain in macaque visual cortex. Nature *399*, 575-579.
Treue, S., and Maunsell, J.H. (1996). Attentional modulation of visual motion processing
in cortical areas MT and MST. Nature *382*, 539-541.
Troyer, T.W., Krukowski, A.E., Priebe, N.J., and Miller, K.D. (1998). Contrast-invariant
orientation tuning in cat visual cortex: thalamocortical input tuning and correlation-based
intracortical connectivity. J Neurosci *18*, 5908-5927.
Ungerleider, L.G., Galkin, T.W., Desimone, R., and Gattass, R. (2008). Cortical
connections of area V4 in the macaque. Cereb Cortex *18*, 477-499.
Vaiceliunaite, A., Erisken, S., Franzen, F., Katzner, S., and Busse, L. (2013). Spatial
integration in mouse primary visual cortex. J Neurophysiol *110*, 964-972.
Van Essen, D.C., and Maunsell, J.H.R. (1983). Hierarchical organization and functional
streams in the visual cortex. Trends in Neurosciences *6*, 370-375.
Vendramin, L., Campello, R.J.G.B., and Hruschka, E.R. (2010). Relative clustering
validity criteria: A comparative overview. Statistical Analysis and Data Mining: The ASA
Data Science Journal *3*, 209-235.
Verghese, P. (2001). Visual search and attention: a signal detection theory approach.
Neuron *31*, 523-535.
Williford, T., and Maunsell, J.H.R. (2006). Effects of Spatial Attention on Contrast
Response Functions in Macaque Area V4. J Neurophysiol *96*, 40-54.
Wilson, F.A., O'Scalaidhe, S.P., and Goldman-Rakic, P.S. (1994). Functional synergism
between putative gamma-aminobutyrate-containing neurons and pyramidal neurons in
prefrontal cortex. Proc Natl Acad Sci U S A *91*, 4009-4013.
Zeng, H., and Sanes, J.R. (2017). Neuronal cell-type classification: challenges,
opportunities and the path forward. Nat Rev Neurosci *18*, 530-546.
1393

Figure 1 - Wang

Figure 1. Attentional Modulation of Contrast Response Function

(A) Orientation change detection task. While the monkey maintained fixation, two oriented Gabor stimuli were flashed on and off at two locations: one within the RF overlap region of the recorded V4 column and the other at a location of equal eccentricity across the vertical meridian. The covert attention of the monkey was cued to one of the two locations. One of the two stimuli changed its orientation at an unpredictable time. The monkey was rewarded for making a saccade to the location of orientation change (95% probability of change at the cued location; 5% probability at uncued location [foil trials]). If no change happened (catch trials), the monkey was rewarded for maintaining fixation. Two different waveforms were shown for two single units.

(B) An example trial showing the single-unit signals in the attend-in condition. The time axis is referenced to the appearance of the fixation spot. Spikes (vertical ticks) in each channel come from the single unit with the highest spike rate in this trial. The gray boxes depict stimulus presentation epochs. In this particular trial, 8 sample stimuli with different contrasts were presented, followed by a target stimulus flash with an orientation change that the monkey responded to correctly.

(C) The mathematical function we used to fit neuronal contrast response functions is shown on the top. Schematics at the bottom show the effect of positive attentional modulation of each parameter on the contrast response function.

(D) The best-fitting contrast response functions of three example neurons in "attend in" and "attend away" conditions. Mean ± SEM. Insets show the attentional modulation indices calculated as a function of contrast.

(E) The AMI as a function of contrast for each of the 255 visually responsive single units, with the three example units in (C) highlighted.

(F) Attention effects on the best-fitting parameters of the contrast response function. Each histogram plots the AMI distribution of a particular parameter across the population, with the dashed line marking the 0 modulation and the arrow with a number depicting the median AMI value. The median AMI is significantly different from zero for all 4 parameters (Mann-Whitney U test, p < 0.01).

(G) The mean difference of AMI from 0 for the 4 parameters are shown in the Cumming estimation plot. Mean differences are plotted as bootstrap sampling distributions. Each mean difference is depicted as a dot. Each 95% confidence interval (CI) is indicated by the ends of the vertical error bars. The shaded color represents that the 95% CI does not include 0.

Figure 2 - Wang

Figure 2. Classification of Single Units Using Electrophysiological Features.
(A) Mean waveforms for all 410 single units recorded. Waveform heights have been normalized to help compare spike widths.
(B) Distribution of all single units in the space of the first two principal components (PCs) of the waveforms. The non-Gaussian structure implies that spike shape is a viable feature for classifying single units.
(C) Histogram of the local variation AMI for all units with available local variation (n=341). The dashed line marks the 0 AMI value. The arrow depicts the median value of the distribution. The average local variation of the population is not significantly modulated by attention (Mann-Whitney U test, p = 0.37).
(D) k-means clustering of 341 single units based on PTD and spiking variability. Cell classes are named after their spiking widths (narrow, medium, broad) and their spiking patterns (regular, bursty). Single units within each range of spike width are highlighted in the component space on the top. Unclassified units are displayed as black crosses in the feature space.
(E) t-SNE embedding of the same data in (D) in a 2-dimensional space. The number at the left bottom corner of each panel represents the perplexity paramter of the t-SNE embedding.
(F) The Cumming estimation plot shows the bootstrap sampling distributions of AMIs of CRF parameters for each cell class. The CRF parameters were only available for visually responsive single units.

Figure 3. Contrast Dependency of AMI is Cell-Class and Layer-Specific

(A) The left panel shows the AMI of contrast responses as a function of contrast averaged across visually responsive single units in each cluster. Mean ± SEM. The black line indicates the population mean. The right panel shows the mean AMI averaged across contrast for each cluster. Asterisk indicates either the distribution is significantly different from zero or two distributions are significantly different (Mann-Whitney U test, $p < 0.05$).

(B) To quantify the contrast dependence of attentional modulation, we averaged the AMI for a single unit separately within the low-contrast range and the high-contrast range (using the c_{50} as the low- to high-contrast threshold). We then defined the contrast dependence index (CDI) as the difference between the average AMI within the low-contrast range and that within the high-contrast range, normalized by the mean AMI across the whole contrast range. The schematic shows the interpretation of different ranges of CDI in terms of the AMI.

(C) The Cumming estimation plot shows the raw data of CDIs (left) and the bootstrap sampling distribution of the mean (right) for each cell class. The plus signs are the outliers within the axis range, and the arrows depict the outliers outside the axis limit. The number of valid units for each cell class is shown on the top of the swarm plot. Distributions for cell classes with CIs inclusive of 0 are shown in faded colors.

(D) Layer-wise AMI (mean ± SEM) of contrast responses for each cell class as a function of contrast (left) or averaged across contrast (right). Asterisk indicates either the distribution is significantly different from zero or two distributions are significantly different (Mann-Whitney U test, $p < 0.05$). Cell classes that contain fewer than 10 units (including outliers) are excluded.

(E) Layer-wise CDI for five clusters, all units, and non-narrow units. The Cumming estimation plot shows the bootstrap sampling distribution of the mean CDI. Distributions with CIs inclusive of 0 are illustrated in faded colors. The number of units excluding outliers is shown on the top of each plot. For the raw data of the layer-wise CDIs, see Supp. Figure 3B.

Figure 4 - Wang

Figure 4. Computational Models Provide A Parsimonious Explanation for the Laminar Profile of AMI Contrast Dependence
(A) Predictions from the normalization model of attention with different suppressive field sizes or different excitatory (E) receptive field sizes. The top panel shows contrast response functions for a simulated neuron in the normalization model, when attending to a stimulus within the neuron's receptive field (black curve) and when attending toward the opposite hemifield (gray curve). The aorange curve represents the AMI. The black dot shows the inflection point of "attend away" responses that was used to delimit the low- and high-contrast ranges. The middle panel shows CDIs for simulated neurons as a function of the E receptive field size and the suppressive field size while holding the stimulus size and the attention field size fixed. The white rectangles depict a potential mechanism that leads to the observed variation of CDIs across layers (change in suppressive field size). The black asterisk corresponds to the model parameters used for the simulation above. The bottom panel shows AMIs of suppressive drive as a function of the E receptive field size and the suppressive field size. For both simulations, the attention field size is 30 and the stimulus size is 5. The normalization model predicts the AMI of the suppressive drive to be correlated with the CDI of neuronal responses.
(B) Simulations of a conductance-based E-I network with columnized connections. Schematics of the E-I networks corresponding to the possible mechanism in (A) are shown on top. 800 E and 200 I units were evenly distributed in 10 columns around a ring. We interpret the normalization model's suppressive field as the receptive field of inhibitory neurons in the E-I network model. E and I units from the same column are mutually coupled. We modeled I receptive field size as the standard deviations (σ_I) of E-I connections (W_{ie}) across columns (middle panel, -5 and 5 are the same column). We increased the range of E-I connections across columns (W_{ie}) (shades of green) while keeping other connections the same (gray, including W_{ee}, W_{ii}, W_{ei}). At the bottom, the raster plot shows the spiking activity for all units organized by their column IDs (blue, I; red, E) in response to a step input. The box depicts a 200 ms window used for computing cross-correlations between E and I populations.
(C) Cross-correlograms between E and I populations in the same column with different I receptive field sizes. Cross-correlations were calculated using the pooled spike trains of E units and I units from the same column across 500 repeats of identical simulation and averaged across 10 columns. A larger I receptive field reduces the cross-correlation between local E and I populations. Mean ± SEM.
(D) Cross-correlograms (mean ± SEM) between Narrow and Broad Regular cell classes in the superficial, input, and deep layer. Cross-correlations were calculated using the pooled spike trains of Narrow class (putative inhibitory) neurons and Broad Regular class (putative excitatory) neurons, and were averaged across sessions. The arrows mark 3 time intervals during which we averaged the cross-correlations and compared the mean differences between the superficial (or input) and the deep layers. Asterisk: The mean difference of cross-correlations in the center interval (-75 ms to 75 ms) has a 95% CI above 0. For the estimation plot, see Supp. Figure 4B.
(E) Proposed E-I networks in V4 accounting for the layer-wise CDI variations. The empirical data and the model simulations imply a larger inhibitory pooling size in the deep layer than those in the superficial and input layer. The arrows depict the canonical information flow pathways in a columnar circuit.

Supp. Figure 2. Validations for the Single-unit Classification and the AMI of CRF Parameters for Each Cell Class

(A) Percentage of classified neurons in the total sample as a function of the number of clusters (k) input to the k-means clustering algorithm. The 5-cluster result was able to identify the largest set of distinct clusters while classifying most of the units.

(B) Cross-validation (CV) errors for different numbers of clusters. The 5-cluster result shows the lowest CV error.

(C) The minimum Jaccard index across clusters for each k from the subsampling analysis. The analysis was applied to neuronal data from either of the two attention conditions or to combined data. Clusters that have Jaccard indices above 0.5 are considered as stable.

(D) The cell-wise co-clustering matrix showing the probability of single units belonging to the same cluster in the subsampling analysis.

(E) In the principal component space of spike shape, we colored single units based on their spike width range (open circles; narrow, medium, broad). The clusters generated from the peak-to-trough duration were minimally overlapped.

(F) In the principal component space of spike shape, we colored single units either based on their spike width range (open circles; narrow, medium, broad) or by running the k-means clustering algorithm with the first 2 PCs (closed circles). The clusters generated from the k-means clustering match the ones grouped by the peak-to-trough duration, suggesting that peak-to-trough duration is an efficient measure to capture the variance in spike shapes.

(G and H) Embedding the data used for the k-means clustering in a 2-dimensional space using t-SNE (G) or UMAP (H).

(I) Mean firing rate for visually responsive single units split by cell class or by layer. Neuronal firing rates were calculated from stimulus flashes with the highest common contrast across two monkey experiments in the "attend away" condition. The number of single units within each cluster is shown. In each layer, we only analyzed clusters containing more than 10 single units. Asterisk indicates either the distribution is significantly different from zero or two distributions are significantly different (Mann-Whitney U test, $p < 0.05$). Mean ± SEM.

(J) The Cumming estimation plot shows the raw data (left) of AMIs of best-fitting CRF parameters and the bootstrap sampling distribution of each cell class's mean (right).

Supp. Figure 2 - Wang

Supp. Figure 2. Validations for the Single-unit Classification and the AMI of CRF Parameters for Each Cell Class

(A) Percentage of classified neurons in the total sample as a function of the number of clusters (k) input to the k-means clustering algorithm. The 5-cluster result was able to identify the largest set of distinct clusters while classifying most of the units.

(B) Cross-validation (CV) errors for different numbers of clusters. The 5-cluster result shows the lowest CV error.

(C) The minimum Jaccard index across clusters for each k from the subsampling analysis. The analysis was applied to neuronal data from either of the two attention conditions or to combined data. Clusters that have Jaccard indices above 0.5 are considered as stable.

(D) The cell-wise co-clustering matrix showing the probability of single units belonging to the same cluster in the subsampling analysis.

(E) In the principal component space of spike shape, we colored single units based on their spike width range (open circles; narrow, medium, broad). The clusters generated from the peak-to-trough duration were minimally overlapped.

(F) In the principal component space of spike shape, we colored single units either based on their spike width range (open circles; narrow, medium, broad) or by running the k-means clustering algorithm with the first 2 PCs (closed circles). The clusters generated from the k-means clustering match the ones grouped by the peak-to-trough duration, suggesting that peak-to-trough duration is an efficient measure to capture the variance in spike shapes.

(G and H) Embedding the data used for the k-means clustering in a 2-dimensional space using t-SNE (G) or UMAP (H).

(I) Mean firing rate for visually responsive single units split by cell class or by layer. Neuronal firing rates were calculated from stimulus flashes with the highest common contrast across two monkey experiments in the "attend away" condition. The number of single units within each cluster is shown. In each layer, we only analyzed clusters containing more than 10 single units. Asterisk indicates either the distribution is significantly different from zero or two distributions are significantly different (Mann-Whitney U test, $p < 0.05$). Mean ± SEM.

(J) The Cumming estimation plot shows the raw data (left) of AMIs of best-fitting CRF parameters and the bootstrap sampling distribution of each cell class's mean (right).

Supp. Figure 3. Raw Data of AMIs and CDIs of AMI for Each cell class
(A) The AMI of firing rate as a function of contrast for single units within each cell class.
(B) The raw data of cluster-wise CDIs of AMI within each layer. The plus signs are the outliers within the axis range, and the arrows depict the outliers outside the axis limit. The number of valid units for each cell class is shown on the top of the swarm plot.
(C) Layer-wise AMI (mean ± SEM) for all units, Narrow unit, and non-narrow units as a function of contrast (left) or averaged across contrast (right). Asterisk indicates either the distribution is significantly different from zero or two distributions are significantly different (Mann-Whitney U test, $p < 0.05$).

Supp. Figure 4 - Wang

A

B

Supp. Figure 4. Normalization Model of Attention and CCG Analyses Between Cell Classes

(A) The structure of the normalization model of attention. The left panel shows a pair of orientated grating stimuli with identical contrasts, acting as input to the model. The central black dot indicates the fixation point. The dashed red circle indicates the receptive field of the model neuron centered on the grating stimulus. The stimulus drive shown in the middle panel is a collection of neural activity driven by the stimuli. Neurons are arranged based on their receptive field center (horizontal position) and orientation preference (vertical position). The values of the stimulus drive are shown by brightness. The top panel shows the attention field as a function of the receptive field center and the orientation preference. In this case, the attention is guided to the right stimulus position and does not vary with orientation. Gray areas indicate values of 1, and white areas indicate values greater than 1. The suppressive drive at the bottom is calculated from the point-by-point product of the stimulus drive and the attention field and then pooled over space and orientation according to the suppressive field size. The stimulus drive is multiplied by the attention field and then divided by the suppressive field to generate the output firing rates of model neurons (right panel).

(B) i, CDIs for simulated neurons in the normalization model with different stimulus sizes and attention field sizes. In each panel, we vary the E receptive field size relative to the attention field size (x-axis), and the suppressive field size relative to the E receptive field size (y-axis). The pattern of CDI holds for a range of values of stimulus size and attention field size. ii, CDIs for simulated neurons in the normalization model with different types of inputs. We changed the stimulus drive input to the normalization model to have either a nonlinear or an attention-modulated contrast response function. We tested both the response gain (10% increase in overall response) and the contrast gain (1% of increase in detected contrast) effects. For these simulations, the attention field size is 30 and the stimulus size is 5. The pattern of CDI holds for different types of inputs.

(C) Changes in E receptive field size (white box) can also lead to the variation of CDIs across layers (left panel). We tested this hypothesis in the E-I network by adjusting the standard deviation of between-column E-E connections (W_{ee}) from narrow (green) to broad (orange) while keeping other connections the same (gray, including W_{ee}, W_{II}, W_{Ie}) (middle panel). Cross-correlograms between E and I populations in the same column suggest that different E receptive field sizes have little impact on the spike-time correlations of local neural activity across layers (right panel).

(D) Cross-correlograms (mean ± SEM) between Narrow and 3 other cell classes in the superficial, input, and deep layer. Cross-correlations were calculated using the pooled spike trains of Narrow class and the other cell class (Board Bursty, Medium Regular, or Medium Bursty) and were averaged across sessions.

(E) The Cumming estimation plot shows the mean difference for cell-class specific comparisons of average cross-correlations between the superficial (*Super.*) and deep layers or between the input and deep layers. We picked 3 time intervals to compute the average cross-correlations (rows). The raw data of average cross-correlations is plotted on the left in each panel. Each mean difference between layers is plotted on the right as a bootstrap sampling distribution.

C

Supp. Figure 4. Normalization Model of Attention and CCG Analyses Between Cell Classes

(A) The structure of the normalization model of attention. The left panel shows a pair of orientated grating stimuli with identical contrasts, acting as input to the model. The central black dot indicates the fixation point. The dashed red circle indicates the receptive field of the model neuron centered on the grating stimulus. The stimulus drive shown in the middle panel is a collection of neural activity driven by the stimuli. Neurons are arranged based on their receptive field center (horizontal position) and orientation preference (vertical position). The values of the stimulus drive are shown by brightness. The top panel shows the attention field as a function of the receptive field center and the orientation preference. In this case, the attention is guided to the right stimulus position and does not vary with orientation. Gray areas indicate values of 1, and white areas indicate values greater than 1. The suppressive drive at the bottom is calculated from the point-by-point product of the stimulus drive and the attention field and then pooled over space and orientation according to the suppressive field size. The stimulus drive is multiplied by the attention field and then divided by the suppressive field to generate the output firing rates of model neurons (right panel).

(B) i, CDIs for simulated neurons in the normalization model with different stimulus sizes and attention field sizes. In each panel, we vary the E receptive field size relative to the attention field size (x-axis), and the suppressive field size relative to the E receptive field size (y-axis). The pattern of CDI holds for a range of values of stimulus size and attention field size. ii, CDIs for simulated neurons in the normalization model with different types of inputs. We changed the stimulus drive input to the normalization model to have either a nonlinear or an attention-modulated contrast response function. We tested both the response gain (10% increase in overall response) and the contrast gain (1% of increase in detected contrast) effects. For these simulations, the attention field size is 30 and the stimulus size is 5. The pattern of CDI holds for different types of inputs.

(C) Changes in E receptive field size (white box) can also lead to the variation of CDIs across layers (left panel). We tested this hypothesis in the E-I network by adjusting the standard deviation of between-column E-E connections (W_{ee}) from narrow (green) to broad (orange) while keeping other connections the same (gray, including W_{ee}, W_{II}, W_{Ie}) (middle panel). Cross-correlograms between E and I populations in the same column suggest that different E receptive field sizes have little impact on the spike-time correlations of local neural activity across layers (right panel).

(D) Cross-correlograms (mean ± SEM) between Narrow and 3 other cell classes in the superficial, input, and deep layer. Cross-correlations were calculated using the pooled spike trains of Narrow class and the other cell class (Board Bursty, Medium Regular, or Medium Bursty) and were averaged across sessions.

(E) The Cumming estimation plot shows the mean difference for cell-class specific comparisons of average cross-correlations between the superficial (*Super.*) and deep layers or between the input and deep layers. We picked 3 time intervals to compute the average cross-correlations (rows). The raw data of average cross-correlations is plotted on the left in each panel. Each mean difference between layers is plotted on the right as a bootstrap sampling distribution.

Supp. Figure 4. Normalization Model of Attention and CCG Analyses Between Cell Classes

(A) The structure of the normalization model of attention. The left panel shows a pair of orientated grating stimuli with identical contrasts, acting as input to the model. The central black dot indicates the fixation point. The dashed red circle indicates the receptive field of the model neuron centered on the grating stimulus. The stimulus drive shown in the middle panel is a collection of neural activity driven by the stimuli. Neurons are arranged based on their receptive field center (horizontal position) and orientation preference (vertical position). The values of the stimulus drive are shown by brightness. The top panel shows the attention field as a function of the receptive field center and the orientation preference. In this case, the attention is guided to the right stimulus position and does not vary with orientation. Gray areas indicate values of 1, and white areas indicate values greater than 1. The suppressive drive at the bottom is calculated from the point-by-point product of the stimulus drive and the attention field and then pooled over space and orientation according to the suppressive field size. The stimulus drive is multiplied by the attention field and then divided by the suppressive field to generate the output firing rates of model neurons (right panel).

(B) i, CDIs for simulated neurons in the normalization model with different stimulus sizes and attention field sizes. In each panel, we vary the E receptive field size relative to the attention field size (x-axis), and the suppressive field size relative to the E receptive field size (y-axis). The pattern of CDI holds for a range of values of stimulus size and attention field size. ii, CDIs for simulated neurons in the normalization model with different types of inputs. We changed the stimulus drive input to the normalization model to have either a nonlinear or an attention-modulated contrast response function. We tested both the response gain (10% increase in overall response) and the contrast gain (1% of increase in detected contrast) effects. For these simulations, the attention field size is 30 and the stimulus size is 5. The pattern of CDI holds for different types of inputs.

(C) Changes in E receptive field size (white box) can also lead to the variation of CDIs across layers (left panel). We tested this hypothesis in the E-I network by adjusting the standard deviation of between-column E-E connections (W_{ee}) from narrow (green) to broad (orange) while keeping other connections the same (gray, including W_{ee}, W_{II}, W_{Ie}) (middle panel). Cross-correlograms between E and I populations in the same column suggest that different E receptive field sizes have little impact on the spike-time correlations of local neural activity across layers (right panel).

(D) Cross-correlograms (mean ± SEM) between Narrow and 3 other cell classes in the superficial, input, and deep layer. Cross-correlations were calculated using the pooled spike trains of Narrow class and the other cell class (Board Bursty, Medium Regular, or Medium Bursty) and were averaged across sessions.

(E) The Cumming estimation plot shows the mean difference for cell-class specific comparisons of average cross-correlations between the superficial (*Super.*) and deep layers or between the input and deep layers. We picked 3 time intervals to compute the average cross-correlations (rows). The raw data of average cross-correlations is plotted on the left in each panel. Each mean difference between layers is plotted on the right as a bootstrap sampling distribution.