Virus relatedness predicts susceptibility in novel host species

Ryan M. Imrie*, Katherine E. Roberts, Ben Longdon

Centre for Ecology & Conservation, Biosciences, College of Life and Environmental Sciences,

University of Exeter, Penryn Campus, Penryn, Cornwall

*corresponding author: ri237@exeter.ac.uk

Abstract

As a major source of outbreaks and emerging infectious diseases, virus host shifts cause significant

health, social and economic damage. Predicting the outcome of infection with novel combinations of

virus and host remains a key challenge in virus research. Host evolutionary relatedness can explain

variation in transmission rates, virulence, and virus community composition between host species,

but there is much to learn about the potential for virus evolutionary relatedness to explain variation

in the ability of viruses to infect novel hosts. Here, we measure correlations in the outcomes of

infection across 45 *Drosophilidae* host species with four *Cripavirus* isolates that vary in their

evolutionary relatedness. We found positive correlations between every pair of viruses tested, with

the strength of correlation tending to decrease with greater evolutionary distance between viruses.

These results suggest that virus evolutionary relatedness can explain variation in the outcome of

host shifts and may be a useful proxy for determining the likelihood of novel virus emergence.

Author Summary

Many outbreaks and new infectious diseases are caused by viruses jumping into new host species. Predicting when jumps will occur, what the characteristics of viruses will be, and how they are likely to evolve in their new hosts are major challenges, requiring an understanding of the interactions between virus and host that is not often available for new viruses or variants. Instead, while studies of the new virus are ongoing, we rely on comparisons to closely related viruses to fill the gaps in our knowledge. How much we can trust these comparisons to be informative, and how their usefulness decreases when comparing viruses that are more distantly related, has not been well studied. Here we used four viruses of fruit flies with different levels of relatedness to begin to answer these questions. We found that, across 45 different fly species, infections were more similar in closely related viruses. The similarity of infections between any two viruses closely matched how similar the genomes of each virus were, suggesting this may be a good rule of thumb to assess how reliable comparisons between viruses are, and how well we can estimate the likelihood of new viruses jumping between host species.

Introduction

Virus host shifts, where viruses jump to and establish onward transmission in novel host species, are

a major source of outbreaks and emerging infectious diseases [1–3]. Many human infections,

including HIV, Ebola, and recently SARS-CoV-2, have shifted into humans from other species, and

continue to cause significant health, social and economic damage [4–7]. Predicting and preventing

virus host shifts have consequently become major goals of virus research [8]. Many challenges

remain in achieving these goals, including improving our understanding of the host, virus, and

ecological factors that influence the outcome of host shifts [9,10], and deciding on the best strategy

for surveillance [11,12]. A better grasp of the evolutionary and epidemiological trajectories of viruses

after host shifts is also required to infer which pathogens may become established in novel host

species [13].

Several studies have investigated how host evolutionary relatedness can explain variation in the

outcome of infection across host species. Greater phylogenetic distance between the natural

(donor) and recipient hosts is associated with decreased likelihood of cross-species transmission

[14,15] and reduced onward transmission within the novel host species [16]. Phylogenetic distance

between hosts also explains variation in virulence after cross-species transmission, which increases

when viruses jump between more distantly related hosts [16–18]. Groups of closely related hosts

have also been shown to share similar levels of susceptibility to novel viruses, independent of the

distance to the natural host [19,20], and harbour similar virus communities [21–23].

Fewer studies have investigated how the evolutionary relatedness and genetics of viruses influence

the outcome of host shifts. Within species, genotype-by-genotype interactions between hosts and

viruses can be important in determining the outcomes of infection [24–26], but little is known about

the importance of genetic differences between viruses in determining infection across host species

[20]. Following host shifts, virus genomes accumulate mutations as they adapt to the novel host

[27,28], and higher pre-standing genetic variation may make some viruses more likely to emerge

than others [29,30]. Despite high rates of geographical overlap between potential hosts [31], closely

related viruses tend to infect the same broad host taxa [32], suggesting they share similar

constraints on their host ranges. However, shifts between divergent host species are also common

across every virus family [33]. A study of fungal pathogens of plants found that, in addition to the

host phylogeny being an important determinant of a pathogen's ability to infect novel host species,

related pathogens share similar host ranges [34]. It has yet to be determined if a similar pathogen

phylogenetic effect exists in viruses.

Although we do not know the extent to which closely related viruses share traits, inferring the

characteristics of viruses from better studied relatives is sometimes necessary. This is often the case

during the early stages of outbreaks, where primary research on new viruses or variants is not yet

available. When SARS-CoV-2 first emerged, its characteristics and epidemiological trajectory were

inferred from the closely related SARS-CoV, MERS-CoV, endemic coronaviruses responsible for

common colds [35], and from other pandemic respiratory viruses such as influenza [36].

Comparisons to previous outbreaks were also used to parameterise disease models in the 2009

H1N1 pandemic [37,38], the 2014 Ebolavirus outbreak [39], and in forecast models of seasonal

influenza [40]. Even for viruses that are not newly emerged, many experimental models of infection

rely on surrogates when the virus of interest is unavailable, non-permissive in cell culture or animal

models, or requires considerable adaptation to experimental hosts [41,42].

In these approaches, the evolutionary scale at which comparisons are being made is typically not

quantified, relying instead on the classification of different virus strains and species [43]. As such, it

is difficult to assess their ability to predict the characteristics of newly emerging viruses.

Comparisons between more distantly related viruses, such as bat and canine rabies viruses [44] and

diverged lineages of influenza viruses [45–47], found stark differences across larger evolutionary

scales. Given the huge diversity of viruses in wildlife [48], and our inability to exhaustively test every

feasible virus variant [8], a greater understanding of the correlations that exist between viruses, and

the evolutionary distances at which these correlations break down, is necessary to understand the

likelihood of novel virus emergence.

In this study, we have investigated how patterns of host susceptibility are correlated between

closely related viruses, using experimental infections with four *Cripavirus* isolates (family

Dicistroviridae) across a panel of 45 host species of *Drosophilidae*. Three of the viruses are isolates of

Drosophila C virus (DCV-C, DCV-EB and DCV-M), a well-studied virus isolated from *Drosophila*

melanogaster [49]. The fourth virus is the closely related Cricket Paralysis virus (CrPV) which was

isolated from Australian field crickets (*Teleogryllus commodus*) and is a widely used model insect

pathogen [50–52]. Both DCV and CrPV infect a broad range of host species [53], cause virulent

infections in adult flies [19,54], and share similar mechanisms for co-opting the host translation

machinery [55]. A major-effect resistance gene called *pastrel* increases resistance to DCV in *D.*

melanogaster [56–58] and has also been shown to provide cross-resistance to CrPV along with

another gene, *Ubc-E2H* [59]. Both DCV and CrPV are targeted by the host antiviral RNAi pathway and

each encodes a potent suppressor of antiviral RNAi. However, these suppressors have different

functions and target different components of the RNAi pathway [60,61]. DCV and CrPV also differ in

their tissue pathology; DCV has been shown to infect gut tissues, causing intestinal obstruction

following septic inoculation in *D. melanogaster*, which was not observed in CrPV infection [62].

Although little is known about the differences between DCV isolates, they have been shown to cause

similar levels of virulence in *D. melanogaster* and interact differently with the endosymbiont

Wolbachia [63].

Previous work in this host system has shown that susceptibility to DCV-C varies across host species,

with the host phylogeny explaining a large proportion of the variance in both viral load and virulence

[19]. The host phylogeny is also an important determinant of the evolution of DCV-C in novel hosts,

with evidence that mutations that adapt the virus to one host may also adapt it to closely related

host species. This suggests virus genotype could alter the likelihood of host shifts in *Drosophila* [28].

Here we measured the ability of four viruses to replicate and persist across host species and

examined correlations between these viruses in the context of their evolutionary relatedness.

Materials & Methods

Fly Stocks

Flies were taken from laboratory stocks of 45 different species of *Drosophila* (S1 Text Table A).

Before experiments began all included stocks were confirmed to be negative for infection with DCV

and CrPV by quantitative reverse transcription PCR (qRT-PCR, described below). Stocks were

maintained in multi-generation Drosophila stock bottles (Fisherbrand) at 22˚C, in a 12-hour light-

dark cycle. Each bottle contained 50ml of one of four varieties of food media (S1 Text

Supplementary Methods).

Host Phylogeny

The method used to infer the host phylogeny has been described in detail elsewhere [19]. Briefly,

publicly available sequences of the *28S*, *Adh*, *Amyrel*, *COI*, *COII*, *RpL32*, and *SOD* genes were collected

from Genbank (see https://doi.org/10.6084/m9.figshare.13079366.v1 for a full breakdown of genes

and accessions by species). Gene sequences were aligned in Geneious v9.1.8

(https://www.geneious.com) using a progressive pairwise global alignment algorithm with free end

gaps and a 70% similarity IUB cost matrix. Gap open penalties, gap extension penalties, and

refinement iterations were kept as default.

Phylogenetic reconstruction was performed using BEAST v1.10.4 [64] with genes partitioned into

separate ribosomal (*28S*), mitochondrial (*COI*, *COII*), and nuclear (*Adh*, *Amyrel*, *RpL32*, *SOD*) groups.

The mitochondrial and nuclear groups were further partitioned into groups for codon position 1+2

and codon position 3, with unlinked substitution rates and base frequencies across codon positions.

Each group was fitted to separate relaxed uncorrelated lognormal molecular clock models using

random starting trees and 4-category gamma-distributed HKY substitution models. The BEAST

analysis was run twice, with 1 billion MCMCglmm generations sampled every 100,000 iterations,

using a birth-death process tree-shape prior. Model trace files were evaluated for chain

convergence, sampling, and autocorrelation using Tracer v1.7.1 [65]. The reconstructed tree was

visualised using ggtree v2.0.4 [66].

Virus Isolates

Virus stocks were kindly provided by Julien Martinez (DCV isolates) [63], and Valérie Dorey and

Maria Carla Saleh (CrPV) [61]. DCV-C, DCV-EB and DCV-M were originally isolated from fly stocks

with origins in three separate continents; DCV-C and DCV-EB were isolated from lab stocks

established by wild capture in Charolles, France and Ellis Beach, Australia respectively, while DCV-M

was isolated directly from wild flies in Marrakesh, Morocco [49]. The CrPV isolate was collected from

Teleogryllus commodus in Victoria, Australia [67]. Virus stocks were diluted in Ringers solution [68]

to equalise the relative amounts of viral RNA and checked for contamination with CrPV (DCV

isolates) and DCV (CrPV isolate) by qRT-PCR as described below.

Virus phylogeny

Full genome sequences for DCV-C (*MK645242*), DCV-EB (*MK645239*), DCV-M (*MK645243*), and CrPV

(*NC_003924*) were retrieved from the NCBI Nucleotide database. Annotations of ORFs for the

replicase polyprotein (CrPV: *Q9IJX4*, DCV: *O36966*) and structural polyprotein (CrPV: *P13418*, DCV:

O36967) were collected from the UniProtKB database and used to separate the coding and non-

coding regions of each virus. ORF sequences were concatenated and aligned using the Geneious

progressive pairwise translation alignment algorithm with a Blosum50 cost matrix and default

parameters. Alignments were manually checked for quality and sequences aligning to CrPV ORF1

nucleotides 1-387 and 2704-2728 were removed due to the presence of large indels.

Phylogenetic reconstruction was performed using BEAST v1.10.4 with translated ORF sequences

fitted to an uncorrelated relaxed lognormal molecular clock model using a speciation birth-death

process tree-shape prior. A Blosum62 substitution model [69] with a gamma distribution of rate

variation with four categories and a proportion of invariable sites was used. The model was run for

10 million MCMC generations sampled every 1,000 iterations and evaluated in Tracer v1.7.1 as

above.

Inoculation

Before inoculation, 0-1 day old male flies were kept in vials containing cornmeal media (S1 Text

Supplementary Methods), and were transferred to fresh media every 2 days for one week. Vials

contained between 5 and 20 flies (mean = 14.5) and were kept at 22°C at 70% relative humidity in a

12-hour light-dark cycle. Flies were inoculated at 7-8 days old under CO_2 anaesthesia via septic pin

prick with 12.5µm diameter stainless steel needles (Fine Science Tools, CA, USA). These needles

were bent approximately 250µm from the end to provide a depth stop and dipped in virus solution

before being pricked into the pleural suture of each fly. Inoculated flies were then snap frozen

immediately in liquid nitrogen, providing a 0 days-post-infection (dpi) timepoint, or maintained in

cornmeal vials for a further 2 days ± 3 hours before freezing, providing a 2 dpi time point. Within

replicate blocks the 0 and 2 dpi vials for each virus were inoculated on the same day, and together

constituted one biological replicate. We aimed to collect three biological replicates for each species

and virus combination, with the order of species, vial (0 or 2 dpi), and virus randomised for each

replicate block.

Measuring Change in Viral Load

To measure the change in viral load between 0 and 2 dpi, total RNA was extracted from flies

homogenised in Trizol (Invitrogen, supplied by ThermoFisher) using chloroform-isoproponyl

extraction, and reverse transcribed using Promega GoScript reverse transcriptase (Sigma) with

random hexamer primers. qRT-PCR was carried out on 1:10 diluted cDNA on an Applied Biosystems

StepOnePlus system using Sensifast Hi-Rox Sybr kit (Bioline). Cycle conditions were as follows: initial

denaturation at 95˚C for 120 seconds, then 40 cycles of 95˚C for 5 seconds, and 60˚C for 30 seconds.

DCV isolates were measured using the same primer pair (Forward: 5'-GACACTGCCTTTGATTAG-3',

Reverse: 5'-CCCTCTGGGAACTAAATG-3') which targeted a conserved location and had similarly high

efficiencies across all isolates. For CrPV the following primers were used: Forward: 5'-

TTGGCGTGGTAGTATGCGTAT-3', Reverse: 5'-TGTTCCGTCCTGCGTCTC-3'. *RpL32* housekeeping gene

primers varied by species (S1 Text Tables B & C). For each sample, two technical replicates were

performed for each amplicon (viral and *RpL32*).

Between-plate variation in C_T values was estimated and corrected for using a linear model with plate

ID and biological replicate ID as parameters, as described elsewhere [70,71]. Mean viral C_T values

from technical replicate pairs were normalised to *RpL32* and converted to fold-change in viral load

using the $2^{-\Delta\Delta CT}$ method, where $\Delta C_T = C_{T:Virus} - C_{T:Rpl32}$, and $\Delta\Delta C_T = \Delta C_{T:day0} - \Delta C_{T:day2}$.

Amplification of the correct products was verified by melt curve analysis. Repeated failure to amplify

product, the presence of melt curve contaminants, or departures from the melt curve peaks of

positive samples (± 1.5˚C for viral amplicons, ± 3˚C for *Rpl32*) in either the 0 or 2 dpi samples were

used as exclusion criteria for biological replicates. In total, of the 180 unique combinations of host

species and virus measured, 3 biological replicates were obtained for 161 combinations, 2 replicates

for 18 combinations, and 1 replicate for 1 combination (*Drosophila virilis,* CrPV).

Statistical Analysis

233 Phylogenetic generalised linear mixed models were used to investigate the effects of host and virus

234 relatedness on viral load. Multivariate models were fitted using the R package MCMCglmm [72] with

235 the viral load of each of the four virus isolates as the response variable. The structures of the models

236 were as follows:

$$y_{hiv} = \beta_{1:v} + \mu_{p:hv} + \mu_{s:hv} + e_{hiv} \; (1)$$

$$y_{hiv} = \beta_{1:v} + \mu_{p:hv} + e_{hiv} \; (2)$$

In these models, y_{hiv} is the change in viral load for virus v in the i^{th} biological replicate of host species

h. The fixed effect β_1 represents the intercepts for each virus isolate, the random effect μ_p represents

the effects of the host phylogeny assuming a Brownian motion model of evolution, and e represents

the model residuals. Model (1) also includes a random species-specific effect that is independent of

the host phylogeny ($\mu_{s:hv}$). This explicitly estimates the non-phylogenetic component of between-

species variance and allows the proportion of variance explained by the host phylogeny to be

calculated. $\mu_{s:hv}$ was removed from model (2) as model (1) struggled to separate the phylogenetic

and species-specific traits. Wing size, measured as the length of the IV longitudinal vein from the tip

of the proximal segment to the join of the distal segment with vein V [73], provided a proxy for body

size [74] and was included in a further model as a fixed effect ($wingsize_h.\beta_2$).

Within each of these models the random effects and residuals were assumed to follow a

multivariate normal distribution with a centred mean of zero and a covariance structure of $V_p \otimes A$ for

the phylogenetic effects, $V_s \otimes I$ for species-specific effects, and $V_e \otimes I$ residuals, where $\otimes$ represents

the Kronecker product. A represents the host phylogenetic relatedness matrix, I an identity matrix

and V represents 4x4 covariance matrices describing the between-species variances and covariances

of changes in viral load for the different viruses. Specifically, the matrices V_p and V_s describe the

phylogenetic and non-phylogenetic between-species variances in viral load for each virus and the

covariances between them, while the residual covariance matrix V_e describes within-species

variance that includes both true within-species effects and measurement errors. Since each

biological replicate was tested with a single virus isolate, the covariances of V_e cannot be estimated

and were set to zero.

Models were run for 13 million MCMC generations, sampled every 5,000 iterations with a burn-in of

3 million generations. Parameter expanded priors were placed on the covariance matrices, resulting

in multivariate F distributions with marginal variance distributions scaled by 1,000. Inverse-gamma

priors were placed on the residual variances, with a shape and scale equal to 0.002. To ensure the

model outputs were robust to changes in prior distribution, models were also fitted with flat and

inverse-Wishart priors, which gave qualitatively similar results.

The proportion of the between species variance that can be explained by the phylogeny was

calculated from model (1) using the equation $v_p /(v_p + v_s)$, where v_p and v_s represent the phylogenetic

and species-specific components of between-species variance [75], and is equivalent to phylogenetic

heritability or Pagel's lambda [76,77]. The repeatability of viral load measurements was calculated

from model (2) as $v_p/(v_p + v_e)$, where V_e is the residual variance of the model [78]. Inter-specific

correlations in viral load were calculated from model (2) V_p matrix as $cov_{x,y}/\sqrt{(var_x + var_y)}$. Parameter

estimates reported are means of the posterior density, and 95% credible intervals (95% CI) were

taken to be the 95% highest posterior density intervals.

The data files and R scripts used in this study are available in an online repository:

https://doi.org/10.6084/m9.figshare.13750711.v1.

Results

Change in viral load is a repeatable trait among host species

To investigate how patterns of host susceptibility differ between related viruses, we experimentally

infected 45 species of *Drosophila* with four virus isolates: DCV-C, DCV-EB, DCV-M, and CrPV. The DCV

isolates were all highly similar (>93% genome and ORF amino acid identity), with the closest

relationship between DCV-C and DCV-EB. CrPV formed an outgroup to the DCV isolates (57-59%

identity, Fig. 1, Table 1).

In total, 15,657 flies were inoculated, and the change in viral load after two days of infection was

determined by qRT-PCR (Fig. 2). The mean viral load within host species ranged from an

approximately 2.7-billion-fold increase in *D. persimilis* infected with DCV-M to a 2.5-fold decrease in

Z. tuberculatus infected with DCV-C. Viral loads across host species tended to be higher for the DCV

isolates, with a mean fold-increase of roughly 11,000 - 19,000, and lower for CrPV, with a mean fold-

increase of roughly 1,600.

295
296   **Figure 1: Phylogeny of virus isolates.** Evolutionary relationships estimated from open reading frame (ORF)
297   amino acid sequences presented in a midpoint-rooted tree. Node labels represent the posterior probabilities
298   of each clade, and the scale bar represents amino acid substitutions per site.

	DCV-C	DCV-EB	DCV-M	CrPV
DCV-C		97.10%	94.00%	57.40%
DCV-EB	98.81%		93.90%	57.30%
DCV-M	98.30%	98.20%		57.40%
CrPV	59.00%	58.70%	58.70%	

Table 1: Virus isolate sequence similarity. Percentage sequence identity was calculated from multiple-

alignment of whole genome nucleotides (white) or concatenated amino acid sequences of ORFs 1 & 2 (grey).

Phylogenetic generalised linear mixed models were fitted to the data to determine the proportion of

variation in viral load explained by the host phylogeny and the repeatability of viral load as a host

trait (Table 2). The phylogeny explained an estimated 79% of the variation in viral load for CrPV but

only 9-21% of the variation for the DCV isolates, with wide credible intervals on all the DCV

estimates. This was due to the model struggling to separate the phylogenetic and species-specific

effects for these viruses. The repeatability of viral load was high for both CrPV (0.66) and the DCV

isolates (0.92-0.96), with the between-species phylogenetic component (v_p) explaining a high

proportion of the variation in viral load with little within-species variation or measurement error (v_r).

We found no significant effect of wing length (a proxy for host body size) on viral load for any of the

included viruses, with all estimates having wide credible intervals overlapping zero.

Figure 2: Change in viral load across a diverse panel of *Drosophila* host species for different virus isolates.
Bar height and colour show the mean change in RNA viral load by 2 dpi on a log2 scale, with error bars
representing the standard error of the mean. The phylogeny of *Drosophila* hosts is presented on the left. The
scale bar represents number of nucleotide substitutions per site and the scale axis represents the approximate
age since divergence in millions of years (my) based on estimates from [79,80]. The virus cladogram is based
on the evolutionary relationships shown in Figure 1.

Virus	Mean change in viral load (log$_2$)	Repeatability	Variance explained by phylogeny
DCV-C	13.50 (95% CI: 11.17, 15.89)	0.96 (95% CI: 0.93, 0.98)	0.11 (95% CI: 0, 0.35)
DCV-EB	14.22 (95% CI: 11.42, 16.76)	0.96 (95% CI: 0.93, 0.98)	0.09 (95% CI: 0, 0.32)
DCV-M	13.63 (95% CI: 10.52, 16.59)	0.92 (95% CI: 0.87, 0.96)	0.23 (95% CI: 0, 0.51)
CrPV	10.66 (95% CI: 8.59, 12.66)	0.66 (95% CI: 0.46, 0.83)	0.79 (95% CI: 0.50, 1.00)

323 **Table 2: Estimates of mean change in viral load, repeatability, and the proportion of variation explained by**
324 **the host phylogeny.** Estimates of the mean change in viral load and repeatability are taken from model (2),
325 while estimates of the variation explained by the host phylogeny are taken from model (1).
326

Correlations between viruses deteriorate with evolutionary distance

We next estimated the inter-specific correlations in patterns of susceptibility between the different

viruses (Fig. 3). We found strong positive correlations between the DCV isolates (r > 0.93), with the

strongest correlation between DCV-C and DCV-EB (r = 0.97). Correlations between DCV isolates and

the more distantly related CrPV were positive (r = 0.52 to 0.59) but weaker than the correlations

between the DCV isolates. DCV-C was more strongly correlated to DCV-EB than to DCV-M (Δr = 0.04,

95% CI = 0.00, 0.09), and more strongly correlated to DCV-M than to CrPV (Δr = 0.40, 95% CI = 0.18,

0.82), reflecting a deterioration in correlation strength with evolutionary distance. Point estimates

imply a similar pattern for DCV-EB, but the evidence for a stronger correlation with DCV-C than DCV-

M was not well supported (S1 Text Table D). Together, this suggests that correlations in patterns of

susceptibility across hosts deteriorate with virus genome divergence, although differences between

the closely related DCV isolates were not always detectable.

340
341

Fig 3: Correlation in viral load between virus isolates. Individual points represent the mean change in viral load after 2 dpi for each host species on a log2 scale, and trendlines have been added from a univariate least-squares linear model for illustrative purposes. Correlations (r) are the total inter-specific correlations and 95% CIs from the output of model (2).

Discussion

Closely related host species present similar environments to novel viruses [75,81], and so often

share similar levels of susceptibility to a given virus [14–20]. Likewise, closely related viruses are

often assumed to share characteristics that make their host interactions, transmission, and

evolutionary trajectories comparable [35–42]. Here, we measured the strength of correlations in the

outcomes of infection with four Cripavirus isolates across 45 host species of *Drosophilidae*, to

investigate whether the susceptibility of novel hosts is similar for closely related viruses. We found

positive correlations between every pair of viruses tested, and the strength of correlation tended to

decrease with greater evolutionary distance between viruses. Between DCV isolates, differences in

correlation strength were detected but not always well supported. However, DCV isolates were

consistently more strongly correlated to each other than to the more distantly related CrPV. This

suggests that between virus variation in the outcome of host shifts may be explained by the

evolutionary relatedness of the viruses.

The strong positive correlations between DCV isolates are likely due to high levels of sequence

conservation resulting in only small differences in their ability to infect different host species.

However, in other instances a small number of mutations have been shown to allow some viruses to

successfully infect a novel host [82–85]. We find some evidence of such effects here. For example, in

Zaprionus davidi DCV-EB shows a decline in viral load, suggesting it is failing to replicate and persist

in this host species, whereas the other DCV isolates and CrPV show an increase in viral load in the

same host. The weaker correlations between DCV and CrPV may be due to interactions with

different host traits that vary in their patterns across the host phylogeny. CrPV and DCV are known

to have distinct methods of suppression of the host antiviral RNAi pathway [60,61] and cause

pathology in different tissues [62]. Additionally, their relatively high levels of sequence divergence

(57-59% identity) may have resulted in changes in the ability of each virus to bind to host cell

receptors, utilise host replication machinery, or subvert host immune defences [86].

Our findings suggest a general rule of thumb: that the ability of a virus to infect a novel host can be

inferred based on its relatedness to other viruses. Many examples exist of one or a few virus

mutations causing large phenotypic changes that would be exceptions to this rule [87–90]. We do

not know the rate at which exceptions like these occur in natural systems and predicting them would

require a mechanistic understanding of the underlying host-virus interactions. Our data also suggest

that host susceptibility is not specific to individual viruses and that certain host clades may be

broadly susceptible to infection. Susceptible host clades may act as reservoirs and donors of

emerging infectious diseases and identifying them in relevant systems could inform control and

prevention strategies [91]. Further work is now needed to expand these findings to broader groups

of viruses, to provide a true comparative framework for understanding virus traits, and to test the

importance of the virus phylogeny in understanding the potential outcomes of virus host shifts.

Acknowledgements

We would like to thank Julien Martinez, Valérie Dorey, and Maria Carla Saleh for kindly providing us

with DCV and CrPV virus isolates, and Jarrod Hadfield for his advice on MCMCglmm model fitting.

Ben Longdon and Katherine E. Roberts are supported by a Sir Henry Dale Fellowship jointly funded

by the Wellcome Trust and the Royal Society (Grant Number 109356/Z/15/Z). Ryan M. Imrie is

supported by a studentship funded by the Natural Environment Research Council (NERC) GW4+

Doctoral Training Partnership and the College of Life and Environmental Sciences, University of

Exeter.

References

1. Woolhouse MEJ, Gowtage-Sequeria S. Host Range and Emerging and Reemerging Pathogens. Emerg Infect Dis.
2005;11(12):1842–7. doi: 10.3201/eid1112.050997

2. Cleaveland S, Laurenson MK, Taylor LH. Diseases of humans and their domestic mammals: pathogen characteristics, host
range and the risk of emergence. Philosophical Transactions Royal Soc Lond Ser B Biological Sci. 2001;356(1411):991–9.
doi: 10.1098/rstb.2001.0889

3. Taylor LH, Latham SM, woolhouse MEJ. Risk factors for human disease emergence. Philosophical Transactions Royal Soc
Lond Ser B Biological Sci. 2001;356(1411):983–9. doi: 10.1098/rstb.2001.0888

4. Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of
human coronaviruses. J Adv Res. 2020;24:91–8. doi: 10.1016/j.jare.2020.03.005

5. Miller IF, Becker AD, Grenfell BT, Metcalf CJE. Disease and healthcare burden of COVID-19 in the United States. Nat Med.
2020;26(8):1212–7. doi: 10.1038/s41591-020-0952-y

6. Frank TD, Carter A, Jahagirdar D, Biehl MH, Douwes-Schultz D, Larson SL et al. Global, regional, and national incidence,
prevalence, and mortality of HIV, 1980–2017, and forecasts to 2030, for 195 countries and territories: a systematic analysis
for the Global Burden of Diseases, Injuries, and Risk Factors Study 2017. Lancet Hiv. 2019;6(12):e831–59. doi:
10.1016/s2352-3018(19)30196-1

7. Huber C, Finelli L, Stevens W. The Economic and Social Burden of the 2014 Ebola Outbreak in West Africa. J Infect Dis.
2018;218(Supplement_5):S698–704. doi: 10.1093/infdis/jiy213

8. Holmes EC. What can we predict about viral evolution and emergence? Curr Opin Virol. 2013;3(2):180–4. doi:
10.1016/j.coviro.2012.12.003

9. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL et al. Pathways to zoonotic spillover. Nat Rev
Microbiol. 2017;15(8):502–10. doi: 10.1038/nrmicro.2017.45

10. Olival KJ, Hosseini PR, Zambrana-Torrelio C, Ross N, Bogich TL, Daszak P. Host and viral traits predict zoonotic spillover
from mammals. Nature. 2017;546(7660):646–50. doi: 10.1038/nature22975

11. Holmes EC, Rambaut A, Andersen KG. Pandemics: spend on surveillance, not prediction. Nature. 2018;558(7709):180–
2. doi: 10.1038/d41586-018-05373-w

12. Levinson J, Bogich TL, Olival KJ, Epstein JH, Johnson CK, Karesh W et al. Targeting Surveillance for Zoonotic Virus
Discovery - Volume 19, Number 5—May 2013 - Emerging Infectious Diseases journal - CDC. Emerg Infect Dis.
2013;19(5):743–7. doi: 10.3201/eid1905.121042

13. Geoghegan JL, Holmes EC. Predicting virus emergence amid evolutionary noise. Open Biol. 2017;7(10):170189. doi:
10.1098/rsob.170189

14. Gilbert GS, Webb CO. Phylogenetic signal in plant pathogen–host range. Proc National Acad Sci. 2007;104(12):4979–83.
doi: 10.1073/pnas.0607968104

15. Streicker DG, Turmelle AS, Vonhof MJ, Kuzmin IV, McCracken GF, Rupprecht CE. Host Phylogeny Constrains Cross-
Species Emergence and Establishment of Rabies Virus in Bats. Science. 2010;329(5992):676–9. doi:
10.1126/science.1188836

16. Guth S, Visher E, Boots M, Brook CE. Host phylogenetic distance drives trends in virus virulence and transmissibility
across the animal–human interface. Philosophical Transactions Royal Soc B Biological Sci. 2019;374(1782):20190296. doi:
10.1098/rstb.2019.0296

17. Farrell MJ, Davies TJ. Disease mortality in domesticated animals is predicted by host evolutionary relationships. Proc
National Acad Sci. 2019;116(16):201817323. doi: 10.1073/pnas.1817323116

18. Mollentze N, Streicker DG, Murcia PR, Hampson K, Biek R. Virulence mismatches in index hosts shape the outcomes of
cross-species transmission. Proc National Acad Sci. 2020;117(46):28859–66. doi: 10.1073/pnas.2006778117

19. Longdon B, Hadfield JD, Day JP, Smith SCL, McGonigle JE, Cogni R et al. The Causes and Consequences of Changes in
Virulence following Pathogen Host Shifts. Plos Pathog. 2015;11(3):e1004728. doi: 10.1371/journal.ppat.1004728

20. Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM. Host Phylogeny Determines Viral Persistence and
Replication in Novel Hosts. Plos Pathog. 2011;7(9):e1002260. doi: 10.1371/journal.ppat.1002260

21. Davies TJ, Pedersen AB. Phylogeny and geography predict pathogen community similarity in wild primates and humans.
Proc Royal Soc B Biological Sci. 2008;275(1643):1695–701. doi: 10.1098/rspb.2008.0284

22. Shaw LP, Wang AD, Dylus D, Meier M, Pogacnik G, Dessimoz C et al. The phylogenetic range of bacterial and viral
pathogens of vertebrates. Mol Ecol. 2020;29(17):3361–79. doi: 10.1111/mec.15463

23. Albery GF, Eskew EA, Ross N, Olival KJ. Predicting the global mammalian viral sharing network using phylogeography.
Nat Commun. 2020;11(1):2260. doi: 10.1038/s41467-020-16153-4

24. Hudson AI, Fleming-Davies AE, Páez DJ, Dwyer G. Genotype-by-genotype interactions between an insect and its
pathogen. J Evolution Biol. 2016;29(12):2480–90. doi: 10.1111/jeb.12977

25. Lambrechts L, Fellous S, Koella JC. Coevolutionary interactions between host and parasite genotypes. Trends Parasitol.
2006;22(1):12–6. doi: 10.1016/j.pt.2005.11.008

26. Carpenter JA, Hadfield JD, Bangham J, Jiggins FM. Specific interactions between host and parasite genotypes do not act
as a constraint on the evolution of antiviral resistance in Drosophila. Evolution. 2012;66(4):1114–25. doi: 10.1111/j.1558-
5646.2011.01501.x

27. Longdon B, Brockhurst MA, Russell CA, Welch JJ, Jiggins FM. The Evolution and Genetics of Virus Host Shifts. Plos
Pathog. 2014;10(11):e1004395. doi: 10.1371/journal.ppat.1004395

28. Longdon B, Day JP, Alves JM, Smith SCL, Houslay TM, McGonigle JE et al. Host shifts result in parallel genetic changes
when viruses evolve in closely related species. Plos Pathog. 2018;14(4):e1006951. doi: 10.1371/journal.ppat.1006951

29. Woolhouse M, Scott F, Hudson Z, Howey R, Chase-Topping M. Human viruses: discovery and emergence. Philosophical
Transactions Royal Soc B Biological Sci. 2012;367(1604):2864–71. doi: 10.1098/rstb.2011.0354

30. Woolhouse MEJ, Adair K, Brierley L. RNA Viruses: A Case Study of the Biology of Emerging Infectious Diseases. Microbiol
Spectr. 2014;1(1):83–97. doi: 10.1128/microbiolspec.oh-0001-2012

31. Jenkins CN, Pimm SL, Joppa LN. Global patterns of terrestrial vertebrate diversity and conservation. Proc National Acad
Sci. 2013;110(28):E2602–10. doi: 10.1073/pnas.1302251110

32. Kitchen A, Shackelton LA, Holmes EC. Family level phylogenies reveal modes of macroevolution in RNA viruses. Proc
National Acad Sci. 2011;108(1):238–43. doi: 10.1073/pnas.1011090108

33. Geoghegan JL, Duchêne S, Holmes EC. Comparative analysis estimates the relative frequencies of co-divergence and
cross-species transmission within viral families. Plos Pathog. 2017;13(2):e1006215. doi: 10.1371/journal.ppat.1006215

34. Vienne DMD, Hood ME, Giraud T. Phylogenetic determinants of potential host shifts in fungal pathogens. J Evolution
Biol. 2009;22(12):2532–41. doi: 10.1111/j.1420-9101.2009.01878.x

35. Zhu Z, Lian X, Su X, Wu W, Marraro GA, Zeng Y. From SARS and MERS to COVID-19: a brief summary and comparison of
severe acute respiratory infections caused by three highly pathogenic human coronaviruses. Respir Res. 2020;21(1):224.
doi: 10.1186/s12931-020-01479-w

36. Petersen E, Koopmans M, Go U, Hamer DH, Petrosillo N, Castelli F et al. Comparing SARS-CoV-2 with SARS-CoV and
influenza pandemics. Lancet Infect Dis. 2020;20(9):e238–44. doi: 10.1016/s1473-3099(20)30484-9

37. Hsieh Y-H. Age groups and spread of influenza: implications for vaccination strategy. Bmc Infect Dis. 2010;10(1):106.
doi: 10.1186/1471-2334-10-106

38. Chao DL, Halloran ME, Obenchain VJ, Longini IM. FluTE, a Publicly Available Stochastic Influenza Epidemic Simulation
Model. Plos Comput Biol. 2010;6(1):e1000656. doi: 10.1371/journal.pcbi.1000656

39. Gomes MFC, Piontti AP y, Rossi L, Chao D, Longini I, Halloran ME et al. Assessing the International Spreading Risk
Associated with the 2014 West African Ebola Outbreak. Plos Curr Outbreaks. 2014;6. doi:
10.1371/currents.outbreaks.cd818f63d40e24aef769dda7df9e0da5

40. Du X, King AA, Woods RJ, Pascual M. Evolution-informed forecasting of seasonal influenza A (H3N2). Sci Transl Med.
2017;9(413):eaan5325. doi: 10.1126/scitranslmed.aan5325

41. Ruiz SI, Zumbrun EE, Nalca A. Animal Models for the Study of Human Disease (Second Edition). Part K Viral Dis.
2017;853–901. doi: 10.1016/b978-0-12-809468-6.00033-4

42. Cann JA, Jahrling PB, Hensley LE, Wahl-Jensen V. Comparative Pathology of Smallpox and Monkeypox in Man and
Macaques. J Comp Pathol. 2013;148(1):6–21. doi: 10.1016/j.jcpa.2012.06.007

43. Simmonds P, Aiewsakun P. Virus classification – where do you draw the line? Arch Virol. 2018;163(8):2037–46. doi:
10.1007/s00705-018-3938-z

44. Troupin C, Dacheux L, Tanguy M, Sabeta C, Blanc H, Bouchier C et al. Large-Scale Phylogenomic Analysis Reveals the
Complex Evolutionary History of Rabies Virus in Multiple Carnivore Hosts. Plos Pathog. 2016;12(12):e1006041. doi:
10.1371/journal.ppat.1006041

45. Tan Y, Guan W, Lam TT-Y, Pan S, Wu S, Zhan Y et al. Differing Epidemiological Dynamics of Influenza B Virus Lineages in
Guangzhou, Southern China, 2009-2010. J Virol. 2013;87(22):12447–56. doi: 10.1128/jvi.01039-13

46. Vijaykrishna D, Holmes EC, Joseph U, Fourment M, Su YC, Halpin R et al. The contrasting phylodynamics of human
influenza B viruses. Elife. 2015;4:e05055. doi: 10.7554/elife.05055

47. Chen R, Holmes EC. The Evolutionary Dynamics of Human Influenza B Virus. J Mol Evol. 2008;66(6):655. doi:
10.1007/s00239-008-9119-z

48. Shi M, Lin X-D, Chen X, Tian J-H, Chen L-J, Li K et al. The evolutionary history of vertebrate RNA viruses. Nature.
2018;556(7700):197–202. doi: 10.1038/s41586-018-0012-7

49. Johnson KN, Christian PD. Molecular Characterization of Drosophila C Virus Isolates. J Invertebr Pathol. 1999;73(3):248–
54. doi: 10.1006/jipa.1998.4830

50. Cherry S, Silverman N. Host-pathogen interactions in drosophila: new tricks from an old friend. Nat Immunol.
2006;7(9):911–7. doi: 10.1038/ni1388

51. Bonning BC, Miller WA. Dicistroviruses. Entomology. 2010;55(1):129–50. doi: 10.1146/annurev-ento-112408-085457

52. Reinganum C, O'Loughlin GT, Hogan TW. A nonoccluded virus of the field crickets Teleogryllus oceanicus and T.
commodus (Orthoptera: Gryllidae). J Invertebr Pathol. 1970;16(2):214–20. doi: 10.1016/0022-2011(70)90062-5

53. Scotti PD, Longworth JF, Plus N, Croizier G, Reinganum C. The Biology and Ecology of Strains of an Insect Small RNA
Virus Complex. Adv Virus Res. 1981;26:117–43. doi: 10.1016/s0065-3527(08)60422-4

54. Nayak A, Kim DY, Trnka MJ, Kerr CH, Lidsky PV, Stanley DJ et al. A Viral Protein Restricts Drosophila RNAi Immunity by
Regulating Argonaute Activity and Stability. Cell Host Microbe. 2018;24(4):542-557.e9. doi: 10.1016/j.chom.2018.09.006

55. Majzoub K, Hafirassou ML, Meignin C, Goto A, Marzi S, Fedorova A et al. RACK1 Controls IRES-Mediated Translation of
Viruses. Cell. 2014;159(5):1086–95. doi: 10.1016/j.cell.2014.10.041

56. Magwire MM, Fabian DK, Schweyen H, Cao C, Longdon B, Bayer F et al. Genome-Wide Association Studies Reveal a
Simple Genetic Basis of Resistance to Naturally Coevolving Viruses in Drosophila melanogaster. Plos Genet.
2012;8(11):e1003057. doi: 10.1371/journal.pgen.1003057

57. Cao C, Cogni R, Barbier V, Jiggins FM. Complex Coding and Regulatory Polymorphisms in a Restriction Factor Determine
the Susceptibility of Drosophila to Viral Infection. Genetics. 2017;206(4):2159–73. doi: 10.1534/genetics.117.201970

58. Cogni R, Cao C, Day JP, Bridson C, Jiggins FM. The genetic architecture of resistance to virus infection in Drosophila. Mol
Ecol. 2016;25(20):5228–41. doi: 10.1111/mec.13769

59. Martins NE, Faria VG, Nolte V, Schlötterer C, Teixeira L, Sucena É et al. Host adaptation to viruses relies on few genes
with different cross-resistance properties. Proc National Acad Sci. 2014;111(16):5938–43. doi: 10.1073/pnas.1400378111

60. Nayak A, Berry B, Tassetto M, Kunitomi M, Acevedo A, Deng C et al. Cricket paralysis virus antagonizes Argonaute 2 to
modulate antiviral defense in Drosophila. Nat Struct Mol Biol. 2010;17(5):547–54. doi: 10.1038/nsmb.1810

61. Rij RP van, Saleh M-C, Berry B, Foo C, Houk A, Antoniewski C et al. The RNA silencing endonuclease Argonaute 2
mediates specific antiviral immunity in Drosophila melanogaster. Gene Dev. 2006;20(21):2985–95. doi:
10.1101/gad.1482006

62. Chtarbanova S, Lamiable O, Lee K-Z, Galiana D, Troxler L, Meignin C et al. Drosophila C Virus Systemic Infection Leads to
Intestinal Obstruction. J Virol. 2014;88(24):14057–69. doi: 10.1128/jvi.02320-14

63. Martinez J, Bruner-Montero G, Arunkumar R, Smith SCL, Day JP, Longdon B et al. Virus evolution in Wolbachia-infected
Drosophila. Proc Royal Soc B. 2019;286(1914):20192117. doi: 10.1098/rspb.2019.2117

64. Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian Phylogenetics with BEAUti and the BEAST 1.7. Mol Biol Evol.
2012;29(8):1969–73. doi: 10.1093/molbev/mss075

65. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior Summarization in Bayesian Phylogenetics Using
Tracer 1.7. Systematic Biol. 2018;67(5):901–4. doi: 10.1093/sysbio/syy032

66. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc Bioinform. 2020;69(1):e96. doi:
10.1002/cpbi.96

67. Johnson KN, Christian PD. A molecular taxonomy for cricket paralysis virus including two new isolates from Australian
populations ofDrosophila (Diptera: Drosophilidae). Arch Virol. 1996;141(8):1509–22. doi: 10.1007/bf01718251

68. Drosophila Ringer's solution. Cold Spring Harbor Protocols. 2007; doi: 10.1101/pdb.rec10919

69. Henikoff S, Henikoff JG. Amino acid substitution matrices from protein blocks. Proc National Acad Sci.
1992;89(22):10915–9. doi: 10.1073/pnas.89.22.10915

70. Ruijter JM, Thygesen HH, Schoneveld OJ, Das AT, Berkhout B, Lamers WH. Factor correction as a tool to eliminate
between-session variation in replicate experiments: application to molecular biology and retrovirology. Retrovirology.
2006;3(1):2. doi: 10.1186/1742-4690-3-2

71. Ruijter JM, Villalba AR, Hellemans J, Untergasser A, Hoff MJB van den. Removal of between-run variation in a multi-
plate qPCR experiment. Biomol Detect Quantification. 2015;5:10–4. doi: 10.1016/j.bdq.2015.07.001

72. Hadfield JD. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package.
Journal of Statistical Software [Internet]. 2010;33(2):1–22. Available from: http://www.jstatsoft.org/v33/i02/

73. Gilchrist GW, Huey RB, Serra L. Rapid evolution of wing size clines in Drosophila subobscura. Genetica. 2001;112–
113(1):273–86. doi: 10.1023/a:1013358931816

74. Huey RB, Moreteau B, Moreteau J-C, Gibert P, Gilchrist GW, Ives AR et al. Sexual size dimorphism in a Drosophila clade,
the D. obscura group. Zoology. 2006;109(4):318–30. doi: 10.1016/j.zool.2006.04.003

75. Freckleton RP, Harvey PH, Pagel M. Phylogenetic Analysis and Comparative Data: A Test and Review of Evidence. Am
Nat. 2002;160(6):712–26. doi: 10.1086/343873

76. Pagel M. Inferring the historical patterns of biological evolution. Nature. 1999;401(6756):877–84. doi: 10.1038/44766

77. Housworth EA, Martins EP, Lynch M. The Phylogenetic Mixed Model. Am Nat. 2004;163(1):84–96. doi: 10.1086/380570

78. Falconer D. Introduction to quantitative genetics. 4th ed. Pearson Education India; 1996.

79. Russo CA, Takezaki N, Nei M. Molecular phylogeny and divergence times of drosophilid species. Mol Biol Evol.
1995;12(3):391–404. doi: 10.1093/oxfordjournals.molbev.a040214

80. Obbard DJ, Maclennan J, Kim K-W, Rambaut A, O'Grady PM, Jiggins FM. Estimating Divergence Dates and Substitution
Rates in the Drosophila Phylogeny. Mol Biol Evol. 2012;29(11):3459–73. doi: 10.1093/molbev/mss150

81. Poulin R, Krasnov BR, Mouillot D, Thieltges DW. The comparative ecology and biogeography of parasites. Philosophical
Transactions Royal Soc B Biological Sci. 2011;366(1576):2379–90. doi: 10.1098/rstb.2011.0048

82. Boon M, Holtappels D, Lood C, Noort V van, Lavigne R. Host Range Expansion of Pseudomonas Virus LUZ7 Is Driven by a
Conserved Tail Fiber Mutation. Phage. 2020;1(2):87–90. doi: 10.1089/phage.2020.0006

83. Lounková A, Kosla J, Přikryl D, Štafl K, Kučerová D, Svoboda J. Retroviral host range extension is coupled with Env-
activating mutations resulting in receptor-independent entry. Proc National Acad Sci. 2017;114(26):E5148–57. doi:
10.1073/pnas.1704750114

84. Allison AB, Organtini LJ, Zhang S, Hafenstein SL, Holmes EC, Parrish CR. Single Mutations in the VP2 300 Loop Region of
the Three-Fold Spike of the Carnivore Parvovirus Capsid Can Determine Host Range. J Virol. 2016;90(2):753–67. doi:
10.1128/jvi.02636-15

85. Vries RP de, Peng W, Grant OC, Thompson AJ, Zhu X, Bouwman KM et al. Three mutations switch H7N9 influenza to
human-type receptor specificity. Plos Pathog. 2017;13(6):e1006390. doi: 10.1371/journal.ppat.1006390

86. Rothenburg S, Brennan G. Species-Specific Host–Virus Interactions: Implications for Viral Host Range and Virulence.
Trends Microbiol. 2019;28(1):46–56. doi: 10.1016/j.tim.2019.08.007

87. Plante JA, Liu Y, Liu J, Xia H, Johnson BA, Lokugamage KG et al. Spike mutation D614G alters SARS-CoV-2 fitness. Nature.
2020;1–6. doi: 10.1038/s41586-020-2895-3

88. Wong G, He S, Leung A, Cao W, Bi Y, Zhang Z et al. Naturally-occurring single mutations in Ebola observably impact
infectivity. J Virol. 2018;93(1):JVI.01098-18. doi: 10.1128/jvi.01098-18

89. Tsetsarkin KA, Vanlandingham DL, McGee CE, Higgs S. A Single Mutation in Chikungunya Virus Affects Vector Specificity
and Epidemic Potential. Plos Pathog. 2007;3(12):e201. doi: 10.1371/journal.ppat.0030201

90. Shan C, Xia H, Haller SL, Azar SR, Liu Y, Liu J et al. A Zika virus envelope mutation preceding the 2015 epidemic enhances
virulence and fitness for transmission. Proc National Acad Sci. 2020;117(33):20190–7. doi: 10.1073/pnas.2005722117

91. Streicker DG, Gilbert AT. Contextualizing bats as viral reservoirs. Science. 2020;370(6513):172–3. doi:
10.1126/science.abd4559

Figure 1

Figure 2

Figure 3