From GWAS Variant to Function: a Study of ~148,000 Variants for Blood Cell Traits.

Quan Sun*,[1] Cheynna A. Crowley*,[1] Le Huang,[2] Jia Wen,[3] Jiawen Chen,[1] Erik L. Bao, [9,10] Paul L. Auer,[4] Guillaume Lettre,[5,6] Alexander P. Reiner,[7,8] Vijay G. Sankaran,[9,10] Laura M. Raffield,[3] Yun Li[1,3,11, #]

1. Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

2. Curriculum in Bioinformatics and Computational Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

3. Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

4. Zilber School of Public Health, University of Wisconsin-Milwaukee, Milwaukee, WI, USA

5. Montreal Heart Institute, Montreal, Quebec, Canada

6. Department of Medicine, Faculty of Medicine, Université de Montréal, Montreal, Quebec, Canada

7. Department of Epidemiology, University of Washington, Seattle, WA, USA

8. Fred Hutchinson Cancer Research Center, University of Washington, Seattle, WA, USA

9. Division of Hematology/Oncology, Boston Children's Hospital and Department of Pediatric Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA, USA

10. Broad Institute of Harvard and MIT, Cambridge, MA, USA

11. Department of Computer Science, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

*These authors contributed equally to this work.

Corresponding author: Yun Li. Department of Genetics, Biostatistics and Computer Science, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA. Fax: (919) 843-4682; E-mail: yunli@med.unc.edu

Abstract

Genome-wide association studies (GWAS) have identified hundreds of thousands of genetic variants associated with complex diseases and traits. However, most variants are noncoding and not clearly linked to genes, making it challenging to interpret these GWAS signals. We present a systematic variant-to-function study, prioritizing the most likely functional elements of the genome for experimental follow-up, for >148,000 variants identified for hematological traits. Specifically, we developed VAMPIRE: Variant Annotation Method Pointing to Interesting Regulatory Effects, an interactive web application implemented in R Shiny (http://shiny.bios.unc.edu/vampire/). This tool efficiently integrates and displays information from multiple complementary sources, including epigenomic signatures from blood cell relevant tissues or cells, functional and conservation summary scores, variant impact on protein and gene expression, chromatin conformation information, as well as publicly available GWAS and phenome-wide association study (PheWAS) results. Leveraging data generated from independently performed functional validation experiments, we demonstrate that our prioritized variants, genes, or variant-gene links are significantly more likely to be experimentally validated. This study not only has important implications for systematic and efficient revelation of functional mechanisms underlying GWAS variants for hematological traits, but also provides a prototype that can be adapted to many other complex traits, paving the path for efficient variant to function (V2F) analyses.

Keywords: Genome-wide association studies, variant to function, functional annotations, experimental validations, blood cell traits

Introduction

Genome-wide association studies (GWAS) have identified thousands of genetic loci and hundreds of thousands of genetic variants associated with various complex human diseases and traits, but the underlying genetic mechanism for the vast majority of these GWAS signals remains elusive. With extensive sequencing and GWAS efforts, there is a pressing need to convert the large and ever growing number of significant GWAS variant-trait pairs into human-interpretable functional or mechanistic knowledge[1]. Most variants identified through GWAS reside in the noncoding regions (e.g., >95% for blood cell traits[2]), and most signals include multiple highly correlated variants or variants in strong linkage disequilibrium (LD). Pinpointing the most likely causal variants within GWAS signals, and linking these variants to their target genes, is challenging, particularly as the number of GWAS loci and variants increases. For hematological traits, for instance, our recent GWAS meta-analyses[3; 4] have revealed over seven thousand loci, with >148,000 variants associated with at least one blood cell index at stringent genome-wide significance threshold. Comprehensive and computationally efficient annotation and prioritization of such GWAS findings are of ever-increasing interest.

Understanding how genetic variants contribute to a phenotype is often referred to as the variant-to-function (V2F) problem. Responding to this problem requires us to determine causal genetic variants, relative cell types/states, their target genes and cellular/physiological functions[5]. Functional experiments are needed to fully reveal molecular mechanisms, but we cannot yet afford to perform time-, money- and labor-consuming experimental validations of thousands of loci involving hundreds of thousands of potentially functional variants or regulatory elements controlling their nearby genes, since each gene is likely regulated by multiple variants and each

variant may regulate multiple genes. Thus, computational methods are needed to screen potential

variants and their effector genes for further experiments.

In this study, we focus on hematological traits. Hematological phenotypes (red blood cell, white

blood cell, and platelet counts and indices) are critical physiological intermediaries in oxygen

transport, immunity, infection, thrombosis, and hemostasis and are associated with autoimmune,

allergic, infectious, and cardiovascular diseases. Hematological traits are highly heritable [6], and

recent large GWAS for hematological traits (including nearly 750,000 participants) identified

thousands of variant-trait associations [2; 4]. In addition, there are multiple large-scale functional

experiments already available[2; 7; 8] for hematological traits, as well as fairly comprehensive

functional annotation resources relevant to blood tissues. This makes hematological traits an

ideal model for this type of V2F computational solution.

We have developed VAMPIRE: Variant Annotation Method Pointing to Interesting Regulatory

Effects, a tool for the user to explore annotations encompassing epigenomic signatures, variant

impact on protein and gene expression, chromatin conformation information from Hi-C and

similar technologies, as well as publicly available GWAS and PheWAS results, creating a

comprehensive annotation profile for variants from recent trans-ethnic blood cell trait

publications[3; 4] with a flexible interface for adding additional future GWAS results. This

interactive web application implemented in R Shiny provides a model display mechanism for

annotating GWAS variants from diverse complex traits, allowing selection of most likely causal

variants and their effector genes for experimental follow-up. Importantly, we show the value of

how variants and genes nominated by VAMPIRE can highlight key regulators of blood cell traits

using independent functional assessment, confirming the value of this annotation tool. While

blood cell traits are the focus for VAMPIRE, this framework (including our R Shiny application)

is adaptable for annotation of other complex trait GWAS results and will facilitate the connection

between variant and function.

Methods

Variant Annotations

The current version of VAMPIRE includes GWAS results from two studies (as detailed in

Supplemental Methods), including all variants in 95% credible sets for fine-mapped

hematological trait associated loci from Chen et al. (N=148,019 variants) [4] and lead variants

(N=2) from a TOPMed imputed GWAS meta-analysis in African American and Hispanic/Latino

populations[3]. We plan to extend VAMPIRE as new trans-ethnic blood cell trait genetic analyses

are released.

The sources of the annotation used are stated clearly in the VAMPIRE online application, with

links or references to the original data sources. As a brief summary, the annotation categories are

trivially split into six types ("variant level", "1D","2D","3D", "PheWAS","GWAS"). First,

"variant level" contains data on phenotypic association from the original publication or preprint

(such as the p-value for association with a given hematological trait, effect size, and posterior

probability of inclusion for fine-mapping credible sets). Second, "1D" refers to epigenomic or

sequence constraints features. This displays selected output from WGSA [9] including functional

prediction scores, conservation scores, and epigenetic information gathered from GeneHancer [10],

FANTOM5 [11; 12], Roadmap [13], and ENCODE [14]. ATAC-seq peaks from recent studies for blood

cell traits [15; 16] and key histone ChIP-seq peaks such as H3K9me3, H3K36me3, H3K4me1,

H3K4me3, and H3K27Ac generated across blood cell related tissues from Roadmap

Epigenomics are also included [13; 17]. We further include information regarding whether each

variant resides in any selective sweep region detected from multiple populations in the 1000

Genomes Project [18] using the S/HIC method [19; 20]. Information is displayed based on the tissue

relevance to the blood cell phenotype (see Supplemental Methods). All variants have 1D

annotation, but for prioritization purposes as described below in the five categories for

noncoding variant annotation, we define 1D annotation as FANTOM5_enhancer_robust =Y

(yes), or Genehancer_feature="Promoter" or "Enhancer" or "Promoter/Enhancer", or coreMarks

(for any relevant roadmap epigenomic category) = "Enhancers" or "Active TSS." Users can then

additionally filter by criteria such as functional prediction and conservation scores.

For the "2D" annotations, we included impact on gene expression and splicing ratios (eQTL and

sQTL information) and impact on protein abundance (pQTL information [21]) from public sources

relevant to blood cell traits. This includes both bulk and cell type specific sources from the

public domain (eQTLGen [22], CAGE [23], BIOS [24] for whole blood, and Raj et al for purified CD4+

T cells and monocytes [25]). Information available in these sources varies, but generally we at a

minimum display the effect size estimate, p-value, the allele assessed, and the gene or protein

involved. Variants were matched across sources based on chromosome, position, and alleles of

each variant. Only significant results (based on FDR or other publication specific thresholds)

from the respective sources are displayed in VAMPIRE; we do note that formal co-localization

analyses would still need to be performed to determine if blood cell related and gene/protein

expression QTL signals truly coincide.

For the "3D" annotations, we include information on 3D genome conformation, linking blood

lineage specific regulatory elements to target genes from various sources. More specifically,

using Hi-C data we incorporated statistically significant long-range chromatin interactions

(LRCI) [17; 26; 27] calculated from Fit-Hi-C [28], loops using the HiCCUPs methodology [26], and super-

FIREs for related tissues [17]. Two Promoter-Capture Hi-C (PCHi-C) data sources [29; 30] were also

incorporated and matched with the 2D results to highlight consistent evidence regarding the

affected gene(s) across "2D" and "3D" annotations. VAMPIRE displays information on the

number of loops, LRCI, PCHi-C interactions, FIREs, or super-FIREs, as well as significance

measures such as p-values, FDR, or CHICAGO scores where applicable. This "3D" annotation

information can also be visualized via our HUGIn browser [31].

The last two data groups present results from two PheWAS sources [4; 32] and GWAS results of

blood cell traits from GWAS catalog [33], allowing the user to evaluate if hematological trait

associated variants may also influence other complex traits.

To visualize and leverage these multiple annotation categories for further analysis or

prioritization of experimental validations, VAMPIRE efficiently displays and integrates relevant

variant information, allowing the user to investigate either all the variants annotated or subsets

based on annotation category groupings, searching either by variant or by gene name. The

comprehensive annotation for the variants is summarized using a five category grouping created

for highlighting the most promising variants as they have various types of annotation.

Specifically, the five categories for noncoding variants are (1) the most restrictive category,

containing variants that have 1D, 2D, and 3D annotation and the genes implicated by 2D and 3D

evidence are consistent; (2) containing variants with 1D, 2D, and 3D evidence, but the genes

implicated from different resources are not consistent; (3) 2D and 3D with consistent gene

evidence between the 2D and 3D annotations; (4) variants with 2D and 3D information and no

consistent gene implied; (5) variants with 1D and 3D evidence. We also have a predicted high

impact coding variant category displayed, including high confidence loss of function (LoF)

variants and likely influential missense, in frame indels, and synonymous variants. Variants

without strongly compelling variant annotation are still displayed, but are not listed in these high

priority categories. The user can further subset results by hematological trait, hematological trait

category, or (for the Chen et al paper [4]) the ancestry specific grouping in which a given credible

set was derived (trans-ethnic, European, East Asian, South Asian, Hispanic/Latino, or African

ancestry). In addition, the user can restrict the amount of information presented by selecting

which tables to be displayed. All tables can be exported in a csv or tab delimited format.

Enrichment analysis

To assess whether the variants prioritized by VAMPIRE are more likely to be functionally

impactful, we performed enrichment analysis at three different levels: variant level, gene level,

and variant-gene pair level, leveraging data generated from previously published functional

experiments [2; 7; 8]. For each set of analysis, we conducted Fisher's exact test and calculated odds

ratios (OR) and one-sided p-values.

At the variant level, we assessed the enrichment of variants that modify transcription factor (TF)

binding motif[2] among our annotation category 1 variants. Recently, Vuckovic et al. [2]

characterized variants that affect erythropoiesis or hematopoiesis by modifying related TF

motifs, such as for KLF1, KLF6, MAFB, and GATA1. We chose these four erythroid TFs as

positive control TFs and two non-erythroid TFs (IRF1 and IRF8) as negative controls.

At the gene level, we evaluated the genes interrogated by Nandakumar et al. [8] with a pooled

short hairpin RNA (shRNA) based loss-of-function approach. Specifically, Nandakumar et al.

studied 389 candidate genes in the neighborhood of 75 loci associated with red blood cell traits

[34], to identify potential causal genes underlying these GWAS signals. We assessed the

enrichment of genes validated by shRNA experiments among those prioritized in VAMPIRE's

category 1. Note that the categories were previously defined at variant level. Here we extent

variant category to gene category as the strongest category where a genome-wide significant

variant linked to this gene falls in.

At the variant-gene pair level, we employed the enhancer-gene connections validated via

CRISPRi-FlowFISH experiments by Fulco et al. [7] in their activity-by-contact (ABC) paper.

Specifically, Fulco et al. tested pairs of candidate *cis* regulatory elements (CREs, ~500bp regions)

and their potential effector genes via CRISPRi perturbations of the CREs, in multiple cell lines

including the K562 cells. Fulco et al. tested 4,124 CRE-gene pairs in total, of which 175 were

significant from their experiments. We overlapped their tested CREs with variants in our

VAMPIRE annotation database. We define a VAMPIRE variant-gene pair confirmed if the

variant overlaps an ABC validated CRE *and* the linked genes in VAMPIRE (from QTL and

chromatin capture conformation evidence) overlaps the corresponding effector gene for that CRE

via ABC's CRISPRi-FlowFISH experiment. We focused on ABC experiments performed on the

K562 cells (instead of GM12878 cells, where a very small number of CREs were tested) as the

number of tested CRE-gene pairs was not too small for robust statistical inference. Matching the

K562 cell line, we focused only on variants associated with red blood cell traits. Similar to the

above two sets of enrichment analyses, we focused on annotations in VAMPIRE's prioritization

category 1. Specifically, we tested whether variant-gene pairs prioritized in VAMPIRE's

category 1 are enriched within ABC's validated enhancer-gene connections. Given the CREs

tested in the ABC paper are rather short (~500bp), we also performed sensitivity analysis by first

extending the CRE regions by +/- 1kb and +/- 5kb and then overlapping variants with these

extended CREs, to ensure robust conclusions.

Results

Overview of VAMPIRE annotations

The overall framework of VAMPIRE is illustrated in **Figure 1**. We started with all variants in

95% credible sets from our recent trans-ethnic study for hematological traits (total 148,019

variants) [4] and lead variants (2 variants) from Kowalski et al.[3]. We incorporated six types of

annotations (detailed in **Methods**): GWAS summary statistics and posterior probability of

inclusion from our previous fine-mapping analyses [4]; epigenomic or sequence constraints

features (1D); eQTL, sQTL and pQTL information (2D); information on 3D genome

conformation (3D); results from two PheWAS sources [4; 32] (PheWAS); and GWAS results from

blood cell traits from GWAS catalog [33] (GWAS).

To visualize and prioritize variants, their corresponding candidate regulatory regions, and their

potential effector genes, we leverage the aforementioned six types of annotation to group these

~148,000 variants into various prioritization categories. Specifically, for non-coding variants,

we classified them into five categories (detailed in Methods). Among them, category 1 is the

most restrictive category, containing variants that have 1D, 2D, and 3D annotation and the genes

implicated by 2D and 3D evidence are consistent. Variants not falling into any of the five

categories are classified as uncategorized. In addition, each gene is categorized according to the

prioritization categories of its linked variant(s). When its linked variants fall in multiple

categories, the gene is assigned to the most highly prioritized category.

Enrichment analysis

Our enrichment analyses employing multiple previously published functional validation

experiments encompassing variant-level, gene-level, and variant-gene pair levels all showed

promising results. Specifically, at the variant level, we found significant enrichment of variants

affecting TF binding motifs among variants prioritized in category 1 of VAMPIRE (**Figure 2**),

for all the erythroid TFs ($p < 8.1E-4$) but GATA1 ($p = 0.18$) (**Table 1**), likely due a smaller

sample size of variants. In contrast, neither of the two negative control TFs (IRF1 and IRF8)

showed any significant enrichment ($p = 0.22$ and 0.62). At the gene level, we focused on two

statistics: (1) number of genes selected for shRNA experiments, since genes were more likely to

be selected for experiments when they demonstrated some prior evidence of potential causality,

and (2) number of genes validated ($p < 0.05$) by shRNA experiments. We compared the number

of genes in our annotation category 1 and all other categories, and found that both shRNA

candidate genes ($p = 3.5E-13$) and significant genes ($p = 3.1E-8$) show strong enrichment among

those in our annotation category 1 (**Table 2**), and the estimated enrichment score for significant

genes (OR $= 4.65$) is almost double of that for candidate genes (OR $= 2.37$). These results

suggest the genes prioritized by VAMPIRE's category 1 annotations are more likely to be

functional.

Finally, at the variant-gene pair level, we also observed enrichment among variants selected into

VAMPIRE's category 1 (**Table 3**). When restricting only to variants in category 1 and associated

with red blood cell traits and without extending the CRE regions, only 7 of VAMPIRE's variant-

gene pairs can be found in ABC's CRISPRi-FlowFISH experiments, of which 6 are not

significant and 1 is significant. While not significant (p = 0.26), the direction of enrichment is

nevertheless encouraging (one of seven, or 14.3%, confirmed by CRISPRi-FlowFISH

experiments) and 3-fold greater than that among all/background pairs from Fulco et al. [7], where

175 out of 4124 variant-gene pairs (4.2%) were confirmed. Note that all the confirmed pairs

were linked with variants associated with red blood cell traits. Further generalizing to all

VAMPIRE annotation categories and to variants associated with any blood cell trait, the

enrichment OR increases to 8.30 with p-value 9.0E-5, indicating that variant-gene pairs

prioritized by VAMPIRE's five categories have much higher odds of being functional. To further

accommodate causal variants tagged by GWAS variants not falling into the short 500bp CREs,

we extended the CREs by +/- 1kb or +/- 5kb, and performed similar enrichment analysis. Our

conclusions remained qualitatively similar (**Table 3**), but the enrichments increased in

significance, thanks to larger sample size (in this context, the larger number of variant-gene pairs

contributing to the analysis) and suggesting that more liberal windows of *cis*-regulatory regions

can capture a higher rate of functional variant-gene pairs. For example, the enrichment for

category 1 variants associated with red blood cell (RBC) traits reached an OR of 15.77 (p=3.8E-

6) and 16.68 (p=3.1E-15) for 1kb and 5kb extension, respectively. We thus conclude that such

enrichment is significant and robust to the extension of CREs.

Application example

Figure 3 shows one example at the *CALR* locus associated with red blood cell traits. Fulco et al.

confirmed by CRISPRi-FlowFISH experiment that CRE chr19:12,996,905-12,998,745 (hg19)

regulates gene *CALR* (adjusted p-value 1.9E-7)[7]. Annotations compiled by VAMPIRE suggest,

consistently, that *CALR* is linked to rs8110787 (chr19:12,999,458, hg19) in category 1.

rs8110787 is associated with several RBC traits [4], including hematocrit (HCT), mean corpuscular

hemoglobin (MCH), mean corpuscular volume (MCV) and red blood cell counts (RBC). Based

on genomic distance alone, *CALR* is not the nearest gene to rs8110787, with several other closer

genes. However, based on H3K27ac HiChIP data in K562 cells [35], rs8110787 significantly

interacts with *CALR* promoter region ($p < 1E-120$), suggesting that *CALR* is a potential target

gene regulated by the CRE around rs8110787. This variant is also an eQTL of *CALR* from

CAGE [23] ($p = 9.4E-16$) and BIOS [24] ($p = 1.0E-25$), and is an enhancer in K562 Leukemia cells

(E123) from Roadmap [13], adding additional evidence. Our VAMPIRE successfully highlights

this rs8110787-*CALR* pair in its category 1.

As a further example of the utility of the VAMPIRE application, we present the annotation

results for one of the lead genome-wide significant variants from recent trans-ethnic GWAS

analyses from Chen et al. [4] For our analysis, we were particularly interested in exploring low

frequency variants, and those more common in those of non-European ancestry. We were able to

quickly rank and prioritize variants for further examination using the annotation categories

described above, including the low frequency variant rs112097551 associated with mean

corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and red blood cell count.

This low frequency intergenic variant rs112097551 (*GATA2-RPN1* locus, 0.15% minor allele

frequency in Chen et al. trans-ethnic analysis [4]) has no close linkage disequilibrium proxies in

African or European populations, and thus was not compared to other highly correlated variants.

Based on variant frequency, particularly in European ancestry populations, we had no

expectation this variant would have eQTL or pQTL evidence (2D annotation), given currently

available sample sizes for eQTL and pQTL analysis. For low frequency variants, 1D and 3D

annotation would be the highest annotation category likely for a variant of interest like

rs112097551. The variant is ~5x more common among African versus non-African samples in

gnomAD version 2.1.1. It is the only variant in the credible set in fine-mapping analyses from

Chen et al. 1D annotation suggests this variant is highly conserved (CADD Phred score of 20.4,

meaning the variant is amongst the top 1% of deleterious variants in the human genome), and it

is rated as deleterious by FATHMM-XF (rank score 0.99169, close to the maximum score of 1).

It is also in open chromatin in megakaryocyte–erythroid progenitor cells, based on hematopoietic

ATAC-seq data [36]. 3D annotation from PCHi-C data in erythroblasts from Javierre et al. [29] links

this variant to the gene *RUVBL1* ~500Kb away, as well as noncoding transcripts *RNU2-37P* and

RUVBL1-AS1. Based on this data, which can be quickly displayed using the VAMPIRE

application, we are currently working on *in vitro* follow-up of this candidate functional enhancer

variant [37].

Discussion

As genotyped sample sizes increase and meta-analysis efforts grow ever larger, more variant-

trait pairs are identified for complex traits than can be easily annotated on a variant by variant

basis. New, user-friendly applications are needed for rapid display of functional annotation

information and prioritization of variants for further functional follow-up to pave the V2F path.

Our VAMPIRE tool provides an example of how the publicly available code can be adapted to

accommodate other sources of annotation specific to other complex trait GWAS results or to

accommodate future blood cell trait GWAS and annotation resources. In addition to *a priori*

providing one category of coding variants and 5 categories of non-coding variants that warrant

prioritization consideration, VAMPIRE allows users to decide their own categories based on

arbitrary combinations of the annotations at adjustable thresholds (for example, prioritizing high

CADD score variants, or variants in open chromatin in blood cells based on ATAC-seq). Along

with the addition of more blood cell trait genetics papers published in the future, VAMPIRE

could also be used as written to annotate GWAS results for other blood related phenotypes, such

as recent GWAS of risk of myeloproliferative neoplasm or clonal hematopoiesis [38; 39].

As we accumulate additional functional validation data, including high-throughput massively

parallel reporter assays (MPRA), medium-throughput CRISPRi/CRISPRa and low throughput

mouse xenotransplant experiments, we will provide statistics summarizing experimental

validation results (e.g., number of variants in the category followed-up, proportion that show

evidence of functional impact in their experiments) for each of the 6 VAMPIRE categories and

for user defined categories. Importantly, we illustrate the value of VAMPIRE using existing

independent functional validation and therefore illuminate the value of this type of annotation

tool in enabling one to go from variant to function for blood cell traits and other complex

phenotypes.

We also note that there are some limitations of VAMPIRE. First, comprehensive annotations

specific to various cell types and cell states would further enhance classification and

prioritization accuracy of functional variants or regulatory elements and their target genes.

Although data is increasingly being generated by us [15; 16] and others [29; 35], and has been

incorporated into VAMPIRE where available, interrogations in a cell-type- or state- specific

manner are still much needed. For instance, our recent work has demonstrated cell-type or tissue

specific FIREs [17; 40] and super interactive promoters (SIP)[41] play key regulatory role and aid the

identification and prioritization of functional regulatory elements and their corresponding genes.

As more experimental data are generated, we will update VAMPIRE accordingly. Second, our

list of 148,019 variants derives primarily from fine-mapping studies, which may be inaccurate in

loci where more than one independent or partially independent signals exist. However, this

limitation cannot be resolved before more powerful methods are developed for fine-mapping

analysis for trans-ethnic GWAS. Finally, most of the annotations are based on analyses in

European ancestry individuals (e.g. eQTL, pQTL, chromatin conformation etc.). Many ongoing

efforts including ours are generating resources for non-European ancestry samples. For example,

we are involved in several recently funded efforts to generate RNA-sequencing data in non-

European ancestry individuals in hematopoietic cell types and anticipate relevant eQTL and

sQTL annotations being added to VAMPIRE in upcoming years.

In conclusion, we have built a comprehensive annotation tool, VAMPIRE, which provides

characterization and prioritization of blood cell trait related GWAS signals. Our results using

existing functional experiments demonstrate that variants and genes prioritized by VAMPIRE

are significantly more likely to be functionally validated at either the variant, gene, or variant-

gene pair level. Annotation tools like VAMPIRE, which could be easily modified to apply to

additional complex traits and diseases, are necessary to translate knowledge of GWAS

significant variants to target genes and biological insights, and to guide our decisions to prioritize

experimental validations of most likely functional regulatory variants/elements and their effector

genes.

Appendix

A1. Supplementary methods.

Declaration of Interests

The authors declare no competing interests.

Acknowledgement

This work was supported by the National Center for Advancing Translational Sciences, National Institutes of Health [R01HL146500 to A.P.R., R01HL129132 to Y.L., KL2TR002490 to L.M.R., R01DK103794 to V.G.S.], and the New York Stem Cell Foundation. C.A.C. and Y.L. are also partially supported by R01GM105785 and U01DA052713. V.G.S. is a New York Stem Cell Foundation-Robertson Investigator. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

We would like to thank the Blood Cell Consortium (BCX) and the HemeNet investigators for comments on earlier versions of VAMPIRE. We want to thank Hanling Wang for visualization assistance. We also want to thank the Li lab members for feedback on the R Shiny app.

Web Resources

VAMPIRE: http://shiny.bios.unc.edu/vampire/

GWAS summary statistics from Chen et al.[4]: http://www.mhi-humangenetics.org/en/resources/

GWAS Catalog: https://www.ebi.ac.uk/gwas/

PheWAS website: http://pheweb.sph.umich.edu

Data Availability

The data underlying this article are available in the article and in its online supplementary material.

Figures

Figure 1. Overall framework of this study. VAMPIRE starts with GWAS variants in the 95%

credible sets, integrates different annotations and assigns them into different prioritization

categories. We further demonstrated that our top prioritized category is enriched with variants

that were experimentally validated. VAMPIRE provides a prototype that can be adapted to many

other complex traits, paving the path for efficient variant to function (V2F) analyses.

393   **Figure 2. Variant level TF motif enrichment analysis.** Each dot represents an enrichment

394   score with the line depicting 95% confidence interval (CI). All the upper bounds of these CIs are

395   infinity. The p-values of the enrichment are reflected by the dot size at the OR point estimate

396   with a larger dot indicating more significant the enrichment.

Figure 3. Variant-gene pair example (rs8110787-*CALR*) visualization from HUGIn2 [31].

Fulco et al. confirmed via CRISPRi experiments that chr19:12996905-12998745 (hg19)

regulates gene *CALR* (adjusted p-value 1.9E-7) which is highly expressed in Erythroblasts[7].

Based on annotations in VAMPIRE, *CALR* is linked to rs8110787 (chr19:12999458, hg19) in

prioritization category 1, including higher than expected physical interactions with the *CALR*

locus from erythroblasts pcHiC data[29], eQTL of *CALR* in CAGE[23] and BIOS[24], erythroid ATAC-

seq peak[16] and H3K27ac peak in K562 leukemia cell[13]. rs8110787 is associated with several

RBC traits (namely hematocrit, mean corpuscular hemoglobin, mean corpuscular volume, and

red blood cell count) as reported in Chen et al. [4].

Tables

	Category 1	Uncategorized	p-value	Odds ratio
All RBCT variants	5,687	21,947		
KLF1	34	34	7.10E-08	3.86
KLF6	21	14	4.30E-07	5.79
MAFB	13	13	8.10E-04	3.86
GATA1	8	19	0.18	1.63
IRF1	12	49	0.62	0.95
IRF8	19	58	0.22	1.26

Table 1. Variant level transcription factor (TF) motif enrichment analysis. Four erythroid TFs and two non-erythroid TFs were selected. Fisher's exact test was applied to test for enrichment. Three erythroid TFs show enrichment for our VAMPIRE annotation category 1 (MAFB, KLF6, KLF1, $p<0.05$). GATA1 motif variants also have some evidence of enrichment (odds ratio = 1.625) but this enrichment is not significant ($p=0.18$), likely due to smaller sample size of variants. Two non-hematopoiesis transcription factors selected as controls don't show significant enrichment with VAMPIRE functional annotation category 1. RBCT, red blood cell trait associated.

	Category 1	Other categories	p-value	Odds ratio
All category genes	9,857	7,408		
shRNA Candidate genes	262	83	3.50E-13	2.37
shRNA Validated genes	68	11	3.10E-08	4.65

Table 2. Gene level enrichment analysis. Fisher's exact test was applied to test for enrichment. Both shRNA experiment candidate genes and validated genes show significant enrichment in our most restrictive VAMPIRE annotation category (category 1).

	Not significant	Significant	Significant %	p-value	Odds ratio
All pairs from Fulco et al.	3,949	175	4.24		
Confirmed pairs in category 1 for RBC traits	6	1	14.29	0.26	3.76
Confirmed pairs in category 1 for all traits	6	1	14.29	0.26	3.76
Confirmed pairs in	19	7	26.92	9.00E-05	8.3

all categories for all

traits

Confirmed pairs in category 1 for RBC traits (+/- 1kb)	10	7	41.18	3.80E-06	15.77
Confirmed pairs in category 1 for all traits (+/- 1kb)	21	9	30	3.50E-06	9.66
Confirmed pairs in all categories for all traits (+/- 1kb)	70	21	23.08	4.60E-10	6.76
Confirmed pairs in category 1 for RBC traits (+/- 5kb)	27	20	42.55	3.10E-15	16.68
Confirmed pairs in category 1 for all traits (+/- 5kb)	64	23	26.44	3.80E-12	8.1
Confirmed pairs in all categories for all traits (+/- 5kb)	160	37	18.78	3.10E-13	5.21

Table 3. Variant-Gene pair level enrichment analysis. We performed analysis for three

variant annotation pools (category 1, red blood cell (RBC) trait associated; category 1, any blood

cell trait associated; any annotation priority category (1-5), any blood cell trait associated) and

three CRE lengths. Fisher's exact test was applied to test for enrichment. We found enrichment

for all three variant annotation pools. These enrichments are also robust to the extension of

CREs.

References

1. Liggett, L.A., and Sankaran, V.G. (2020). Unraveling Hematopoiesis through the Lens of
Genomics. Cell 182, 1384-1400.
2. Vuckovic, D., Bao, E.L., Akbari, P., Lareau, C.A., Mousas, A., Jiang, T., Chen, M.H., Raffield,
L.M., Tardaguila, M., Huffman, J.E., et al. (2020). The Polygenic and Monogenic Basis of
Blood Traits and Diseases. Cell 182, 1214-1231.e1211.
3. Kowalski, M.H., Qian, H., Hou, Z., Rosen, J.D., Tapia, A.L., Shan, Y., Jain, D., Argos, M.,
Arnett, D.K., Avery, C., et al. (2019). Use of >100,000 NHLBI Trans-Omics for Precision
Medicine (TOPMed) Consortium whole genome sequences improves imputation quality
and detection of rare variant associations in admixed African and Hispanic/Latino
populations. PLoS genetics 15, e1008500.
4. Chen, M.H., Raffield, L.M., Mousas, A., Sakaue, S., Huffman, J.E., Moscati, A., Trivedi, B.,
Jiang, T., Akbari, P., Vuckovic, D., et al. (2020). Trans-ethnic and Ancestry-Specific
Blood-Cell Genetics in 746,667 Individuals from 5 Global Populations. Cell 182, 1198-
1213.e1114.
5. Nandakumar, S.K., Liao, X., and Sankaran, V.G. (2020). In The Blood: Connecting Variant to
Function In Human Hematopoiesis. Trends in genetics : TIG 36, 563-576.
6. Garner, C., Tatu, T., Reittie, J.E., Littlewood, T., Darley, J., Cervino, S., Farrall, M., Kelly, P.,
Spector, T.D., and Thein, S.L. (2000). Genetic influences on F cells and other
hematologic variables: a twin heritability study. Blood 95, 342-346.
7. Fulco, C.P., Nasser, J., Jones, T.R., Munson, G., Bergman, D.T., Subramanian, V.,
Grossman, S.R., Anyoha, R., Doughty, B.R., Patwardhan, T.A., et al. (2019). Activity-by-
contact model of enhancer-promoter regulation from thousands of CRISPR
perturbations. Nat Genet 51, 1664-1669.
8. Nandakumar, S.K., McFarland, S.K., Mateyka, L.M., Lareau, C.A., Ulirsch, J.C., Ludwig, L.S.,
Agarwal, G., Engreitz, J.M., Przychodzen, B., McConkey, M., et al. (2019). Gene-centric
functional dissection of human genetic variation uncovers regulators of hematopoiesis.
eLife 8.
9. Liu, X., White, S., Peng, B., Johnson, A.D., Brody, J.A., Li, A.H., Huang, Z., Carroll, A., Wei,
P., Gibbs, R., et al. (2016). WGSA: an annotation pipeline for human genome
sequencing studies. Journal of medical genetics 53, 111-112.
10. Fishilevich, S., Nudel, R., Rappaport, N., Hadar, R., Plaschkes, I., Iny Stein, T., Rosen, N.,
Kohn, A., Twik, M., Safran, M., et al. (2017). GeneHancer: genome-wide integration of
enhancers and target genes in GeneCards. Database : the journal of biological
databases and curation 2017.
11. Lizio, M., Harshbarger, J., Shimoji, H., Severin, J., Kasukawa, T., Sahin, S., Abugessaisa, I.,
Fukuda, S., Hori, F., Ishikawa-Kato, S., et al. (2015). Gateways to the FANTOM5
promoter level mammalian expression atlas. Genome Biol 16, 22.
12. Lizio, M., Abugessaisa, I., Noguchi, S., Kondo, A., Hasegawa, A., Hon, C.C., de Hoon, M.,
Severin, J., Oki, S., Hayashizaki, Y., et al. (2019). Update of the FANTOM web resource:

expansion to provide additional transcriptome atlases. Nucleic Acids Res 47, D752-
D758.
13. Bernstein, B.E., Stamatoyannopoulos, J.A., Costello, J.F., Ren, B., Milosavljevic, A.,
Meissner, A., Kellis, M., Marra, M.A., Beaudet, A.L., Ecker, J.R., et al. (2010). The NIH
Roadmap Epigenomics Mapping Consortium. Nat Biotechnol 28, 1045-1048.
14. Consortium, E.P. (2004). The ENCODE (ENCyclopedia Of DNA Elements) Project. Science
306, 636-640.
15. Ludwig, L.S., Lareau, C.A., Bao, E.L., Nandakumar, S.K., Muus, C., Ulirsch, J.C.,
Chowdhary, K., Buenrostro, J.D., Mohandas, N., An, X., et al. (2019). Transcriptional
States and Chromatin Accessibility Underlying Human Erythropoiesis. Cell Rep 27,
3228-3240.e3227.
16. Ulirsch, J.C., Lareau, C.A., Bao, E.L., Ludwig, L.S., Guo, M.H., Benner, C., Satpathy, A.T.,
Kartha, V.K., Salem, R.M., Hirschhorn, J.N., et al. (2019). Interrogation of human
hematopoiesis at single-cell and single-variant resolution. Nature Genetics 51, 683-693.
17. Schmitt, A.D., Hu, M., Jung, I., Xu, Z., Qiu, Y., Tan, C.L., Li, Y., Lin, S., Lin, Y., Barr, C.L., et
al. (2016). A Compendium of Chromatin Contact Maps Reveal Spatially Active Regions
in the Human Genome. Cell reports 17, 2042-2059.
18. Genomes Project, C., Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M.,
Korbel, J.O., Marchini, J.L., McCarthy, S., McVean, G.A., et al. (2015). A global
reference for human genetic variation. Nature 526, 68-74.
19. Schrider, D.R., and Kern, A.D. (2016). S/HIC: Robust Identification of Soft and Hard Sweeps
Using Machine Learning. PLoS Genet 12, e1005928.
20. Schrider, D.R., and Kern, A.D. (2017). Soft Sweeps Are the Dominant Mode of Adaptation in
the Human Genome. Mol Biol Evol 34, 1863-1877.
21. Emilsson, V., Ilkov, M., Lamb, J.R., Finkel, N., Gudmundsson, E.F., Pitts, R., Hoover, H.,
Gudmundsdottir, V., Horman, S.R., Aspelund, T., et al. (2018). Co-regulatory networks
of human serum proteins link genetics to disease. Science (New York, NY) 361, 769-
773.
22. Võsa, U., Claringbould, A., Westra, H.-J., Bonder, M.J., Deelen, P., Zeng, B., Kirsten, H.,
Saha, A., Kreuzhuber, R., Kasela, S., et al. (2018). Unraveling the polygenic architecture
of complex traits using blood eQTL metaanalysis. bioRxiv, 447367.
23. Lloyd-Jones, L.R., Holloway, A., McRae, A., Yang, J., Small, K., Zhao, J., Zeng, B., Bakshi,
A., Metspalu, A., Dermitzakis, M., et al. (2017). The Genetic Architecture of Gene
Expression in Peripheral Blood. Am J Hum Genet 100, 228-237.
24. Zhernakova, D.V., Deelen, P., Vermaat, M., van Iterson, M., van Galen, M., Arindrarto, W.,
van 't Hof, P., Mei, H., van Dijk, F., Westra, H.J., et al. (2017). Identification of context-
dependent expression quantitative trait loci in whole blood. Nat Genet 49, 139-145.
25. Raj, T., Rothamel, K., Mostafavi, S., Ye, C., Lee, M.N., Replogle, J.M., Feng, T., Lee, M.,
Asinovski, N., Frohlich, I., et al. (2014). Polarization of the effects of autoimmune and
neurodegenerative risk alleles in leukocytes. Science 344, 519-523.
26. Rao, S.S.P., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D., Robinson, J.T.,
Sanborn, A.L., Machol, I., Omer, A.D., Lander, E.S., et al. (2014). A 3D map of the
human genome at kilobase resolution reveals principles of chromatin looping. Cell 159,
1665-1680.
27. Gorkin, D.U., Qiu, Y., Hu, M., Fletez-Brant, K., Liu, T., Schmitt, A.D., Noor, A., Chiou, J.,
Gaulton, K.J., Sebat, J., et al. (2019). Common DNA sequence variation influences 3-
dimensional conformation of the human genome. Genome Biol 20, 255.
28. Ay, F., Bailey, T.L., and Noble, W.S. (2014). Statistical confidence estimation for Hi-C data
reveals regulatory chromatin contacts. Genome Research 24, 999-1011.
29. Javierre, B.M., Burren, O.S., Wilder, S.P., Kreuzhuber, R., Hill, S.M., Sewitz, S., Cairns, J.,
Wingett, S.W., Varnai, C., Thiecke, M.J., et al. (2016). Lineage-Specific Genome

Architecture Links Enhancers and Non-coding Disease Variants to Target Gene
Promoters. Cell 167, 1369-1384 e1319.
30. Jung, I., Schmitt, A., Diao, Y., Lee, A.J., Liu, T., Yang, D., Tan, C., Eom, J., Chan, M., Chee,
S., et al. (2019). A compendium of promoter-centered long-range chromatin interactions
in the human genome. Nat Genet 51, 1442-1449.
31. Martin, J.S., Xu, Z., Reiner, A.P., Mohlke, K.L., Sullivan, P., Ren, B., Hu, M., and Li, Y.
(2017). HUGIn: Hi-C Unifying Genomic Interrogator. Bioinformatics 33, 3793-3795.
32. Denny, J.C., Bastarache, L., Ritchie, M.D., Carroll, R.J., Zink, R., Mosley, J.D., Field, J.R.,
Pulley, J.M., Ramirez, A.H., Bowton, E., et al. (2013). Systematic comparison of
phenome-wide association study of electronic medical record data and genome-wide
association study data. Nature Biotechnology 31, 1102-1111.
33. Buniello, A., MacArthur, J.A.L., Cerezo, M., Harris, L.W., Hayhurst, J., Malangone, C.,
McMahon, A., Morales, J., Mountjoy, E., Sollis, E., et al. (2019). The NHGRI-EBI GWAS
Catalog of published genome-wide association studies, targeted arrays and summary
statistics 2019. Nucleic Acids Res 47, D1005-d1012.
34. van der Harst, P., Zhang, W., Mateo Leach, I., Rendon, A., Verweij, N., Sehmi, J., Paul,
D.S., Elling, U., Allayee, H., Li, X., et al. (2012). Seventy-five genetic loci influencing the
human red blood cell. Nature 492, 369-375.
35. Mumbach, M.R., Satpathy, A.T., Boyle, E.A., Dai, C., Gowen, B.G., Cho, S.W., Nguyen,
M.L., Rubin, A.J., Granja, J.M., Kazane, K.R., et al. (2017). Enhancer connectome in
primary human cells identifies target genes of disease-associated DNA elements. Nat
Genet 49, 1602-1612.
36. Corces, M.R., Buenrostro, J.D., Wu, B., Greenside, P.G., Chan, S.M., Koenig, J.L., Snyder,
M.P., Pritchard, J.K., Kundaje, A., Greenleaf, W.J., et al. (2016). Lineage-specific and
single-cell chromatin accessibility charts human hematopoiesis and leukemia evolution.
Nat Genet 48, 1193-1203.
37. Hu, Y., Stilp, A.M., McHugh, C.P., Jain, D., Zheng, X., Lane, J., Rao, S., de Bellefon, S.M.,
Raffield, L.M., Chen, M.-H., et al. (2020). Whole genome sequencing association
analysis of quantitative red blood cell phenotypes: the NHLBI TOPMed program.
medRxiv, 2020.2012.2009.20246736.
38. Bick, A.G., Weinstock, J.S., Nandakumar, S.K., Fulco, C.P., Bao, E.L., Zekavat, S.M.,
Szeto, M.D., Liao, X., Leventhal, M.J., Nasser, J., et al. (2020). Inherited causes of
clonal haematopoiesis in 97,691 whole genomes. Nature 586, 763-768.
39. Bao, E.L., Nandakumar, S.K., Liao, X., Bick, A.G., Karjalainen, J., Tabaka, M., Gan, O.I.,
Havulinna, A.S., Kiiskinen, T.T.J., Lareau, C.A., et al. (2020). Inherited myeloproliferative
neoplasm risk affects haematopoietic stem cells. Nature 586, 769-775.
40. Crowley, C., Yang, Y., Qiu, Y., Hu, B., Abnousi, A., Lipiński, J., Plewczyński, D., Wu, D.,
Won, H., Ren, B., et al. (2020). FIREcaller: Detecting frequently interacting regions from
Hi-C data. Comput Struct Biotechnol J 19, 355-362.
41. Song, M., Pebworth, M.P., Yang, X., Abnousi, A., Fan, C., Wen, J., Rosen, J.D., Choudhary,
M.N.K., Cui, X., Jones, I.R., et al. (2020). Cell-type-specific 3D epigenomes in the
developing human cortex. Nature 587, 644-649.
575