Frequent origins of traumatic insemination involve convergent shifts in sperm and genital morphology

Jeremias N. Brand[1*], Gudrun Viktorin[1], R. Axel W. Wiberg[1], Christian Beisel[2], Luke J. Harmon[3] and Lukas Schärer[1]

[1] University of Basel, Department of Environmental Sciences, Zoological Institute, Vesalgasse 1, 4051 Basel, Switzerland

[2] Department of Biosystems Science and Engineering, ETH Zürich, Basel, Switzerland

[3] Department of Biological Sciences, University of Idaho, Moscow, USA

Short title:

Frequent origins of traumatic insemination

*Corresponding Author

University of Basel,

Department of Environmental Sciences, Zoological Institute,

Vesalgasse 1, 4051 Basel, Switzerland

Email: jeremias.br@gmail.com

Abbreviations

PC: principal component

pPCA: phylogenetically corrected principal component analysis

Abstract

Traumatic insemination is a mating behaviour during which the (sperm) donor uses a traumatic intromittent organ to inject an ejaculate through the epidermis of the (sperm) recipient, thereby frequently circumventing the female genitalia. It likely evolves due to sexual selection and sexual conflict, since it allows the donor to bypass pre- and postcopulatory choice and resistance mechanisms of the recipient. Several striking cases of traumatic insemination have been studied to date, but the frequency of its evolution, the intermediate stages via which it originates, and the morphological changes that such shifts involve remain poorly understood. Based on observations of reproductive traits in 145 species of the free-living flatworm genus *Macrostomum*—in combination with a robust phylogenomic phylogeny—we present comparative work on the evolution of traumatic insemination. We identify at least nine, but up to 13, independent origins of traumatic insemination, which involve convergent shifts in both sperm design and male and female genital morphology. We further find that traumatic insemination may originate via internal wounding. Finally, we highlight the large diversity of reproductive traits across the genus and show evidence for male-female coevolution. Our findings indicate that sexual selection and sexual conflict repeatedly favour the evolution of behaviours—like traumatic insemination—that allow donors to bypass control mechanisms of recipients, leading to predictable shifts in both male and female reproductive traits.

Keywords: traumatic mating, hypodermic insemination, copulatory wounding, phylogenetics, evolution, female genitalia, correlated evolution, parallel evolution, sexually antagonistic coevolution

Introduction

The sexes frequently show a difference in mating propensity because male fertility (i.e. fertilized egg production) is often limited by the number of matings a male can achieve, while female fertility is often limited by the amount of resources a female can invest into eggs and offspring [1–3]. The resulting conflict over mating rate has far-reaching consequences, often resulting in "Darwinian sex roles" with choosy females and eager males [4]. Choice may be beneficial for females, since it allows them to preferentially reproduce with males based on genetic compatibility, genetic quality [5] and/or direct benefits (e.g. nuptial gifts [6]). Indeed, evidence for female choice is abundant and there are many species where females mate multiply, suggesting that polyandry may indeed result in such benefits [7]. However, females may also simply mate multiply as a result of male harassment, and while multiple mating could be costly to females, resisting male harassment might be even costlier [7, 8]. Costly harassment is expected to arise frequently, since female choice necessarily goes against the rejected males' interests [9], potentially leading to sexually antagonistic coevolution between male persistence and female resistance traits [8, 10, 11].

In polyandrous species, sexual selection and sexual conflict will continue after copulation through intricate interactions of the female genital tract with the male intromittent organs and the received ejaculate [12–14]. Female genitalia might exert postcopulatory control mechanisms through differential sperm storage, sperm ejection or sperm digestion, thus applying a selective filter on male genital and ejaculate traits. In analogy to the precopulatory conflict, it is then possible for traits in males to arise that attempt to bypass or influence the female choice and resistance mechanisms, again resulting in sexually antagonistic coevolution.

Sexually antagonistic coevolution can drive the emergence of male traits that inflict considerable harm on females [11, 15, 16]. A striking example that implicates such harm is traumatic insemination, which occurs in some internally fertilising species and involves the infliction of a wound to the female's integument through which the male then transfers its ejaculate [17]. Since traumatic insemination frequently occurs in both gonochoristic (separate-sexed) and hermaphroditic species

[17], we in the following use the more general terms (sperm) donor and (sperm) recipient to refer to

the two sexual roles, with no loss of generality [18].

Although traumatic insemination often results in costs to the recipients [15, 17, 19–22], it has evolved

repeatedly across various animal groups [17]. While natural selection might play a role in its evolution

in some taxa—especially the endoparasitic Strepsiptera [23, 24]—it likely often evolves because it

allows the donor to bypass pre- and postcopulatory choice and resistance mechanisms of the recipient.

Specifically, traumatic insemination can enable donors to force copulation and thus minimise the

control that the recipient could otherwise exert over mating [15]. Traumatic insemination usually also

allows the donor to bypass the recipient's genitalia, by depositing sperm either closer to the site of

fertilisation [24, 25] or even directly within the relevant tissue [15, 26], thus likely reducing the

recipient's ability to control the fate of the received ejaculate [12, 17]. In this view, traumatic

insemination allows the donor to bypass the influence of the recipient's sexually antagonistic choice

and resistance mechanisms, temporarily gaining an advantage in the coevolutionary chase.

However, since conflicts persist under traumatic insemination, we expect selection to then act on

traits that allow the recipient to regain control over mating and/or the fate of the received ejaculate.

For example, some species of bed bugs have evolved what could be considered a secondary vagina,

a structure that has been shown to reduce the costs incurred due to traumatic insemination [19, 27].

But even without the emergence of new organs, the recipients could evolve behavioural or

physiological responses to avoid traumatic insemination (such as parrying strikes during penis

fencing in polyclad flatworms [28]) or to manipulate and control the hypodermically received

ejaculate (e.g. similar to sperm digestion in copulating species [29–31]).

Besides bypassing recipient choice and resistance mechanisms, traumatic insemination could also

evolve due to sperm competition, since in many internally fertilising species sperm of unrelated

donors compete within the female genital tract for fertilisation of the recipient's eggs [32]. In this

context, traumatic insemination might allow donors to bias sperm competition in their favour and

prevent competing donors from removing their previously donated sperm, potentially resulting in

paternity benefits [17]. Indeed, traumatic insemination seems to affect sperm competition in a family

of spiders, where sperm precedence is biased towards the first male in a species with traumatic

insemination, while it is biased towards the second male in its non-traumatically mating relatives [17,

23, 33]. In contrast, traumatic insemination is associated with last male precedence in one species of

bed bug [26], so the effects of it on sperm competition might depend on a species' morphology and

ecology.

Interestingly, since traumatic insemination likely affects not only how sperm interact, e.g. concerning

sperm mixing, sperm removal and sperm density, but also the degree to which they are accessible to

recipient control, it could influence the nature of both sperm competition and cryptic female choice.

Cryptic female choice, sperm removal, and sperm displacement can introduce skews in the

distribution of sperm competing in the female storage organ, which may reduce the intensity of sperm

competition [34–38]. These same postcopulatory mechanisms can also favour sperm designs that

increase the chance of being retained in storage. This could include adaptations to better anchor or

position sperm, making them able to resist removal or displacement by either rival donors or the

recipient [39]. Traumatic insemination could lead to less removal and more mixing, resulting in a

more fair-raffle like sperm competition [40, 41], in which paternity success is more determined by

the relative number of sperm donated, leading to strong selection on sperm number. Since sperm

number is generally assumed to trade-off with sperm size [42–45], it is expected that traumatic

insemination is associated with smaller (and potentially also more numerous) sperm [40, 41].

Traumatic insemination might evolve more frequently in hermaphrodites due to sexual conflict over

mating roles [12, 18, 46–48]. In general, and in analogy to the situation outlined above for

gonochorists [1], a hermaphrodite that already carries enough received sperm to fertilise its own eggs

might gain little from additional matings as a recipient, while it might still gain additional

fertilisations by acting as a donor [12]. It is thus likely that, on average, individual hermaphrodites

show a preference for sperm donation [12, 18, 46–48] and this prediction is supported in several

laboratory studies [47, 49, 50]. Traumatic insemination then potentially allows unilateral enforcement

of donation while avoiding receipt. Additionally, harmful traits are expected to evolve more readily

in simultaneous hermaphrodites because fitness costs incurred by a recipient may be partially

compensated by fitness benefits from the same individual acting as a donor. A hermaphrodite may

thus also engage in matings that involve harmful effects, as long as the net fitness benefits of mating

are positive [46, 51]. Therefore, hermaphrodites could engage in traumatic insemination even if this

results in considerable harm to the recipient.

Indeed, traumatic insemination appears to arise more commonly in simultaneous hermaphrodites

compared to gonochorists, since they comprise 11 out of the 23 well-supported independent origins

listed in [17], which is a lot considering that hermaphrodites make up ~6% of animals [52].

Hermaphroditic animals are thus ideal study organisms for investigations of traumatic insemination,

since—while it has been studied in some charismatic systems [15, 25, 28, 53–56]—we know little

about how frequently it evolves or via which intermediate states a copulating species can transition

to traumatic insemination [17, 20, 22].

Here we present comparative work on the evolution of traumatic insemination across the genus

Macrostomum, a species-rich taxon of hermaphroditic free-living flatworms. In *Macrostomum*,

traumatic insemination is called hypodermic insemination, since in several species the donor uses a

needle-like stylet (**Fig 1**) to inject sperm through the epidermis of the mating partner and sperm then

move through the body of the recipient to the site of fertilisation [41, 57, 58]. Injected sperm can

often be observed inside the parenchymal tissues of these highly transparent animals [41, 57–59],

making it feasible to screen a large number of species for the convergent evolution of hypodermic

insemination. And while we here present evidence that not all traumatically mating *Macrostomum*

species may inject sperm through the external epidermis, we nevertheless use the term hypodermic

insemination for consistency with the earlier literature.

The genus consists of two phylogenetically well-separated clades, with one clade currently thought

to only mate through hypodermic insemination ("hypodermic clade", referred to as Clade 1 in [41])

and a second clade primarily containing reciprocally mating species ("reciprocal clade", referred to

as Clade 2 in [41]). Within the reciprocal clade, hypodermic insemination has previously been shown

to have convergently evolved in *M. hystrix* [41]. Reciprocally copulating *Macrostomum* species

engage in a copulatory handshake, where two individuals insert their often relatively blunt stylet (Fig

1) via their partner's female genital opening into the female sperm storage organ, the female antrum

(further only called antrum), so that both can donate and receive sperm in the same mating [60]. Many

reciprocally copulating species then perform a postcopulatory behaviour, where worms place their

mouth over their own female genital opening and suck [41, 61], presumably in an attempt to remove

components of the received ejaculate from the antrum (called the suck behaviour, [39, 60]). This

removal of ejaculate could target manipulative seminal fluids, since the ejaculate of the main

Macrostomum model species, *M. lignano*, contains substances that influence the mating partners

behaviour, including the propensity to perform the suck behaviour [62, 63]. The suck behaviour could

also be aimed at reducing the number of stored sperm (e.g. to lower the risk of polyspermy), constitute

a form of cryptic female choice (e.g. to favour donors of higher quality), and/or represent a resistance

trait in sexual conflict over mating roles (i.e. to undo unwanted sperm receipt) [41].

If the suck behaviour is a recipient resistance trait, we might expect the evolution of donor persistence

traits, potentially leading to antagonistic coevolution [8]. Indeed, the sperm of reciprocally copulating

species generally have a thin anterior feeler and two stiff lateral bristles that could represent such

persistence traits (**Fig 1**), serving to anchor the sperm in the antrum to prevent removal during the

suck behaviour [39, 41]. In contrast, sperm of species with hypodermic insemination (i.e. the

hypodermic clade and *M. hystrix*) lack lateral bristles and have a simplified morphology, presumably

because they no longer need to resist the suck behaviour [39, 41] which has so far never been observed

in species with hypodermic insemination. These sperm may instead be adapted to efficiently move

through the partner's tissues (**Fig 1**), and one such adaptation could include a size reduction, as

outlined above. Moreover, while species with reciprocal copulation have an antrum with a thickened

epithelium, the species with hypodermic insemination have a simple antrum, presumably because it

no longer interacts with the donor's stylet and sperm, and instead is used for egg-laying only [41].

Based on these findings, the observed adaptations to reciprocal copulation and hypodermic

insemination have been described as the reciprocal and hypodermic mating syndrome, respectively,

since they each constitute specific combinations of morphological (sperm, stylet and antrum) and

behavioural traits [41].

If hypodermic insemination evolved as a resolution of sexual conflict over the mating roles, then we

would expect it to occur frequently, but it is currently unclear whether hypodermic insemination has

convergently arisen more than once within the reciprocal clade. It is also unclear if the transitions are

reversible or if the emergence of hypodermic insemination alters the coevolutionary dynamics

between donor and recipient, so that they cannot readily revert back to reciprocal copulation. Little

molecular data is currently available for most species in the genus *Macrostomum* and the description

of important structures, such as the sperm morphology and aspects of the mating behaviours, are often

neglected in the taxonomic literature.

Here we collected molecular and morphological information on 145 *Macrostomum* species to identify

additional independent origins of hypodermic insemination, to highlight species that represent

possible transitional states in the evolution of hypodermic insemination, and to quantitatively assess

convergent changes in both sperm design and in male and female genital morphology that accompany

its evolution. Using ancestral state reconstruction, we further ask whether species can revert back to

reciprocal copulation once hypodermic insemination has arisen. Moreover, if hypodermic

insemination evolves in the context of sexually antagonistic coevolution, then signatures of male-

female coevolution in the form of a diversity of resistance and persistence traits should be evident

across the genus. Indeed it is predicted that donors might carry multiple persistence traits even if

recipients resistance renders them ineffective [64]. We thus test for covariation between male and

female genital morphology and survey the genus for novel resistance and persistence traits.

Results

Species collected and phylogenetics

We included 145 *Macrostomum* species in our analysis, and established a phylogeny inferred from a combination of newly generated transcriptome data (98 species, see Material and Methods) and a partial *28S rRNA* gene sequence (permitting to add 47 additional species). Except for 14 species, we collected and systematically documented all species ourselves. Based on an integrative approach— using detailed documentation of *in vivo* morphology and *28S rRNA* sequences—we identified 51 species as previously described and a striking 94 species as likely new to science (see Material and Methods for details on species delimitation; **Tab 1** for a summary of all collected species; **Tab S1** for the sample sizes of morphological measures across species; and also SI Taxonomy). Since our focus here was not on taxonomy, we delimited new species as operational taxonomic units. We deposited extensive image, video [65], geographic and molecular data for all specimens collected and their operational species assignment (see **Tab S2**), facilitating their future description (see http://macrostomorpha.info and [65]).

We inferred phylogenies based on two protein alignments generated from transcriptome data (see Material and Methods), one with many genes while having lower occupancy (L alignment with 8218 genes) and one with high occupancy but fewer genes (H alignment with 385 genes, **Tab 2**). For both alignments, we calculated phylogenies using maximum likelihood with IQ-TREE [66] (referred to as L-IQ-TREE and H-IQ-TREE) and summary methods with ASTRAL III [67] (L-ASTRAL and H-ASTRAL). We applied summary methods to account for incomplete lineage sorting and the gene tree – species tree conflict it can cause. Second, to include species for which we lacked transcriptomes, we added information from *28S rRNA* to the H alignment to construct a combined (C) maximum likelihood phylogeny (C-IQ-TREE, see Material and Methods). Third, we performed two types of Bayesian analyses on the H alignment, one that like the maximum-likelihood analysis operates on partitioned amino acids [68] (H-ExaBayes) and a second performed on the unpartitioned DNA coding

sequence alignment to better account for rate heterogeneity across sites [69] (H-PhyloBayes, see

Material and Methods).

Across these phylogenies, all species represented by more than one transcriptome (i.e. 15 x 2, 3 x 3,

and 1 x 9 transcriptomes) were monophyletic, with the notable exception of *M. lignano* (**Fig S1**). One

of the *M. lignano* transcriptomes was from the DV1 inbred line from the type locality near Bibione,

Italy [70] and the other from an outbred population from the Sithonia Peninsula, Greece [61, 71].

M. lignano was monophyletic in the L-IQ-TREE, H-PhyloBayes, and both ASTRAL phylogenies,

but the Greek population was sister to *M. janickei* in the H-IQ-TREE and H-ExaBayes phylogenies,

although node support for this split was low for H-IQ-TREE (**Fig S1**), and removal of cDNA synthesis

primer sequences from the alignment rendered *M. lignano* monophyletic in H-IQ-TREE (see SI

Primer removal and Material and Methods). A closer investigation of the morphology of the Greek

vs. Italian population also revealed that the former had a considerably larger stylet (96.7 vs. 60.7 µm)

and longer sperm (76.5 vs. 62.7 µm). We still consider *M. lignano* a proper species, but a closer

comparison of the Greek and Italian populations would be interesting.

Next, we discuss other agreements and discrepancies between the different inferred phylogenies. The

grouping of the major clades was mostly consistent across all phylogenetic approaches (**Fig 2** and

Fig S1). All phylogenies recovered six large species groups (further called the hypodermic, spirale,

lignano, finlandense, tuba, and tanganyika clade), two smaller species groups (the minutum and

hamatum clade), and two consistent species pairs (*M*. sp. 45 + 46 and *M*. sp. 4 + 89). However, the

backbone, and the positions of some species with long branches (*M*. sp. 37, *M*. sp. 39, *M*. sp. 90 and

M. curvituba), differed depending on the alignment and method used. Despite these discrepancies,

the Robinson-Foulds distances [72] between the phylogenies were low (**Tab S3**), indicating good

agreement between methods.

In all phylogenies, the hypodermic clade was deeply split from the reciprocal clade, but with large

phylogenetic distances also appearing within the hypodermic clade (see also below). Also consistent

was the grouping of the tanganyika, tuba and finlandense clades (**Fig 2** and **Fig S1**). The hamatum

and minutum clades, together with *M.* sp. 4 + 89, were always the closest relatives to the former three

clades. However, the exact relatedness patterns were uncertain. In the L-IQ-TREE, H-IQ-TREE and

H-ExaBayes phylogenies, *M.* sp. 4 + 89 were the closest relatives to these three clades, followed by

the hamatum and minutum clades. In contrast, in the ASTRAL and the H-PhyloBayes phylogenies,

the hamatum clade was more closely related to the minutum clade (with the latter nested within the

former in case of H-PhyloBayes) and both were sister to the grouping of tanganyika, tuba and

finlandense. Moreover, the exact branching order at the base of the reciprocal clade was not clearly

resolved. The spirale clade split off first in the H-IQ-TREE and H-PhyloBayes phylogenies, while

the lignano clade split off first in the other phylogenies. Consistent with the conflict between methods,

the quartet support from the ASTRAL analysis indicated gene tree – species tree conflict at most

nodes in the phylogeny's backbone (**Fig S2**). These internal nodes were separated by short branches

suggestive of rapid speciation events, such as during adaptive radiation [73], where substantial

incomplete lineage sorting is expected. This pattern is also consistent with ancient hybridisation,

which is a distinct possibility, since there is evidence for hybridisation within the genus under

laboratory conditions [74].

The topology of C-IQ-TREE was identical to H-IQ-TREE when we removed all the species added

based on *28S rRNA* (**Fig 3**) and thus adding these species did not negatively influence the overall

topology of the tree. Node support in C-IQ-TREE was somewhat lower, as could be expected given

the placement of the additional species based solely on *28S rRNA* sequences. Nevertheless, the added

inferential power obtained from a ~50% increase in species representation is highly worthwhile, and

we therefore focus on results based on this combined phylogeny in the main text.

As already mentioned above, many species in the hypodermic clade were highly molecularly

diverged, even though they are difficult to distinguish morphologically. Most of these species had

stylets that consisted of a short proximal funnel that tapered to a curved and drawn-out asymmetrical

needle-like thickening (**Fig 3**A, top). It was possible to distinguish some species based on general

habitus. These differences are reflected by the four deeply split clades containing, *M. rubrocinctum*,

M. hystricinum, *M. gabriellae*, and *M. pusillum*, respectively (although the latter two clades were also

quite similar in habitus). We have indicated substantial morphological similarity by appending a letter

to the species name (e.g. *M. hystricinum a*, *c*, *d*). Investigations of species within the hypodermic

clade without support from molecular data thus require considerable caution. Moreover, given these

striking morphological similarities, it may often not be clear to which of these species the name-

bearing type specimens belong to [41], so the species names in this clade should be considered

tentative. Eventually, one may need to name these species afresh, applying more extensive molecular

species delimitation, and either suppress the original names of species without detailed enough

morphological descriptions and/or lacking adequate type material, or to define neotypes, including

molecular voucher specimens [75]. While such a detailed taxonomical revision of the genus

Macrostomum goes well beyond the scope of the current study, we feel that readers need to be aware

of these caveats (see also SI Taxonomy).

Morphological diversity

Before we report on our formal analysis of convergent evolution and coevolution, we highlight

several exciting morphologies, which considerably expand the known morphospace of *Macrostomum*

(**Fig 3**). As mentioned, the hypodermic clade shows little variation in terms of stylet and sperm

morphology. However, the stylet of *M*. sp. 93 differed clearly from that stereotypical form, by having

a small proximal funnel extending into a straight and obliquely-cut tube (**Fig 3**A). This shape is

similar to the stylets of several species in the reciprocal clade, namely *M. shenda*, *M*. sp. 34 and

M. sp. 64 (as well as *M. orthostylum*, for which we have no phylogenetic placement). Since we

observed hypodermic received sperm in both *M*. sp. 93 and *M*. sp. 64 (**Fig 3**), this shape appears

adapted for hypodermic insemination. And while we did not observe hypodermic received sperm in

M. orthostylum and *M*. sp. 34 (nor was such sperm reported for *M. shenda* by [76]), these species are

nevertheless likely to also mate through hypodermic insemination, not least since stylets with similar

shapes are also used for hypodermic insemination in related macrostomid flatworms [77].

In contrast to the largely canalized stylet and sperm of the hypodermic clade, we found remarkable

variation in these structures within the reciprocal clade (**Fig 3**). For example, we documented five

species in the spirale clade that had stylets with lateral protrusions close to the distal opening. The

protrusion is shaped like a rod or spike in *M. evelinae*, *M.* sp. 29 and *M.* sp. 42, while it consists of

two thin filaments in *M.* sp. 30 and *M.* sp. 43 (see also SI Taxonomy). These highly modified stylets

may have coevolved with the complex antra in these species, which have two separate chambers

connected via a ciliated muscular sphincter. The sperm of these species is also remarkable because

they carry only a single bristle instead of the typical two. The sperm of *M.* sp. 30 carry only a single

curved bristle, while the bristle of *M. evelinae*, *M.* sp. 29, and *M.* sp. 42 is additionally modified,

being thicker and appearing flexible, frequently curving back towards the sperm body. A second

bristle might also be absent in the closely related *M.* sp. 13 (indicated with a shaded second bristle in

Fig 3), but the available material currently does not allow an unambiguous assessment.

Sperm are also highly variable across the entire reciprocal clade. Particularly striking are the sperm

modifications of *M.* sp. 82, which give the anterior part of the sperm a translucent appearance under

phase contrast (discussed in more detail below). We also document extraordinarily long sperm in

several species in the tanganyika clade. While it is not entirely clear which part of the sperm is

modified here, it appears that they have very long sperm feelers, with the bristles thus being located

unusually far posterior. Finally, we observed numerous species that either had reduced (length <3.2

µm) or no sperm bristles. Such reduction and losses repeatedly occur across the whole reciprocal

clade and appear to coincide with changes in the stylet and antrum morphology (see our formal tests

below). Finally, a striking modification of sperm design occurs in *M. distinguendum* (finlandense

clade), which appears to lack sperm bristles, but instead carries novel club-shaped lateral appendages,

which in light microscopic images bear no resemblance to the usual sperm bristles.

The antrum morphology was also highly variable within the reciprocal clade, with several species

having complex female antra with multiple chambers. Particularly striking are the two female genital

openings present in four quite distantly related taxa (see SI Taxonomy). A second opening is present

in *M. spiriger*, *M. gieysztori* (and its three close relatives, *M.* sp. 16, *M.* sp. 17 and *M.* sp. 18),

M. paradoxum, and *M.* sp. 82, suggesting multiple independent origins across the genus (see SI

Female opening). The observed variation in male and female genital morphology, and sperm design

could be important for sexual selection, as we outline in the discussion.

Frequent origins of hypodermic insemination

We inferred the number of convergent transitions to hypodermic insemination using ancestral state

reconstruction (ASR) of three reproductive traits: received sperm location, sperm bristle state, and

antrum state (see SI Morphology and Tab 3). Because received sperm location involves the direct

observation of sperm within the recipient's tissue, it provides direct evidence for hypodermic

insemination. However, observation of injected sperm in field-collected specimens can be

challenging, especially in species with low investment into sperm production, thus lowering the

sample size. Since tests of correlated evolution revealed strong associations of hypodermic sperm

with absent/reduced sperm bristles and a thin antrum (see the next section), we also performed ASR

using these states as proxies for hypodermic insemination. Finally, we performed ASR on the inferred

mating syndrome, which represents a synthesis of all available information (see Material and Methods

and Tab 3). We performed ASR, with all traits scored as binary and, where appropriate, also as trinary,

to test if hypodermic insemination could evolve via an intermediate state (see Material and Methods).

All reconstructions indicated frequent origins of hypodermic insemination (Tab 4 and Fig S3). In all

analyses with trinary states, an ordered transition model without gains once traits have been lost

(ORD-Dollo) was preferred, and in all analyses with binary states, a model without gains (Dollo) was

preferred. However, other models, including some permitting gains, also received at least some

support (Tab 4). ASR of trinary states inferred frequent transitions to the intermediate state, which

were driven by the ordered model's requirements to transition through it. These transitions were often

placed along internal branches of the phylogeny, primarily within the finlandense clade, which

contains several species with reduced or absent states and, nested within them, two species with

present states (*M*. sp. 12 and *M*. sp. 44, with received sperm in the antrum, long bristles, and assigned

to the reciprocal mating syndrome; Fig S3 A, C, F).

We estimated a lower bound for the number of transitions by requiring an origin of the derived state

to be separated by other such origins via nodes with a >95% posterior probability of having the

ancestral state. Applying this rule to traits scored as binary, we find nine transitions to hypodermic

received sperm, 17 losses/reductions of sperm bristles, 13 simplifications of the antrum, and 13

transitions to the hypodermic or intermediate mating syndrome (see red stars and numbers in Fig S3).

Moreover, these lower-bound estimates were slightly lower for trinary states. Finally, we also

performed these analyses for the H-IQ-TREE and H-ExaBayes phylogenies, and found qualitatively

very similar results to those of the C-IQ-TREE phylogeny reported above, albeit, since they contain

fewer species, showing somewhat lower numbers of transitions (Tab S4).

Correlated evolution

We performed tests of correlated evolution to ask if the numerous convergent changes in received

sperm location, sperm bristle state and antrum state are evolutionarily dependent. We found strong

support for correlated evolution of received sperm location with both sperm bristle state and antrum

state (Fig 4A+B). This supports previous findings that hypodermic insemination is associated with

changes in sperm design and antrum simplification [41]. Therefore, when observations of received

sperm are missing, both sperm bristle state and antrum state are likely good proxies for the mating

syndrome. We expand on the earlier analyses by also providing evidence for the correlated evolution

between the sperm bristle state and antrum state (Fig 4C), which was implied in [41], but not formally

tested. We find substantially stronger support for correlated evolution than [41] across the board, with

Bayes factors that are 7.7 and 7 times larger for the correlated evolution of sperm bristle state or

antrum state with the inferred mating syndrome, respectively. Moreover, the analyses were robust,

with respect to the phylogeny and the priors used (see SI Correlated evolution).

Hypodermic insemination and convergence in morphospace

Next, we used phylogenetically corrected principal component analysis (pPCA) to investigate if these convergent transitions to hypodermic insemination also coincide with changes in a greater variety of reproductive traits (see SI Morphology). The first two principal components, PC1 and PC2, together captured nearly half of the variation in the analysed reproductive traits (Fig 5), followed by additional principal components with relatively small contributions (Tab S5). Specifically, PC1 captured a change in stylet phenotype, with larger values indicating species with longer, more curved stylets, that are distally more symmetric and less sharp (Fig 5). Larger values of PC1 also indicated both longer sperm and bristles, and an increased probability for the sperm to carry a brush. Finally, high values of PC1 indicated a thickened antrum with a more pronounced cellular valve, and a more complex internal structure. In comparison, PC2 had a less clear interpretation, with high values indicating larger species with larger proximal and distal stylet openings.

Species in the hypodermic clade (stippled outlines) had similar values in PC1 and mainly differed in PC2. Interestingly, species from the reciprocal clade that we had assigned to the hypodermic mating syndrome (solid and left yellow) grouped closely with the species in the hypodermic clade, indicating striking convergence in morphospace concerning stylet, sperm and antrum morphology (see also Fig S4). PC1 further separated species based on the received sperm location, with hypodermic received sperm (right yellow) only found in species with low PC1 values, indicating that PC1 captures a morphology necessary for hypodermic insemination. Almost all species with reduced (triangles) or absent (circles) sperm bristles grouped closely together in PC1, with the notable exception of *M*. sp. 68 and *M*. sp. 82 (black arrowheads), which cluster together with other species that we assigned to the reciprocal mating syndrome. We observed sperm in the antrum of both species (i.e. in 2 of 7 specimens in *M*. sp. 68 and 16 of 21 specimens in *M*. sp. 82) and the antrum is similar in both, with a long muscular duct that performs a 90° turn towards the anterior before it enters a second chamber that is strongly muscular (Fig 6). Moreover, both species have a similar L-shaped stylet with

402 a blunt tip, which makes it unlikely that they mate through hypodermic insemination (see also

403 Discussion).

Pathways to hypodermic insemination

While 117 of the 145 studied species could be assigned to either the hypodermic or reciprocal mating

syndrome, some were not easily identified as performing either type of mating behaviour (Tab 3),

which may either have been due to lack of detailed observations, or more interestingly, due to

potentially transitional patterns that deviated from these syndromes. This included two species that

we categorised as intermediate because we observed sperm both in the antrum and embedded inside

the recipient's tissues (light green triangles in Fig 5). Both species (*M*. sp. 3 and *M*. sp. 101) have a

sharp stylet and sperm with reduced bristles, fitting with the hypodermic mating syndrome. But they

also have a thickened antrum wall, which indicates the reciprocal mating syndrome, and these species

indeed have intermediate values of PC1. Multiple observations of hypodermic sperm in *M*. sp. 3 show

them to be embedded deeply in the anterior wall of the antrum and more deeply in the tissue lateral

to the body axis, extending up to the ovaries, and in the tail plate (Fig 7). And while in *M.* sp. 101 we

did not observe sperm as deeply in the recipient's tissues, some were fully embedded in the cellular

valve and just anterior to it, and thus close to the developing eggs (Fig 8). One explanation for these

findings would be that during mating, the stylet of both species pierces the recipient's antrum wall

and sperm is traumatically injected into the body internally. Unfortunately, no copulations were

observed in mating observations of *M*. sp. 101, but this species has been seen performing the suck

behaviour, as expected if ejaculate is, at least partially, deposited in the antrum (pers. comm. P.

Singh). We lack mating observations for *M*. sp. 3 and further investigations of the mating behaviour

and the antrum histology of both species would be highly desirable.

Three additional species (*M*. sp. 14, *M*. sp. 51 and *M*. sp. 89) were also difficult to classify because,

although their morphology indicates hypodermic insemination, we clearly observed received sperm

in the antrum. *M*. sp. 51 and *M*. sp. 89 grouped with the hypodermically mating species in PC1 (red

arrowheads in Fig 5), while we did not include *M*. sp. 14 in this analysis due to missing data for sperm

bristle length. We found sperm within the antrum in only 1 of 4 specimens in *M*. sp. 51 and 1 of 12

specimens in *M*. sp. 89, and it is thus possible that sperm is hypodermically injected and may later

enter the antrum when an egg passes through the cellular valve into the antrum before egg laying. But

these species could also represent an intermediate state between the mating syndromes. We observed

sperm in the antrum in 3 of 5 specimens in *M*. sp. 14 and it therefore seems less likely that sperm

entering during the transition of the egg into the antrum is the cause of its presence here as well.

Instead, sperm is probably deposited in the antrum by the mating partner during copulation. However,

based on its general morphology, we predict that closer investigations of this species will reveal

hypodermic received sperm in a similar location as found in the other intermediate species (*M*. sp. 3

and *M*. sp. 101).

Finally, we were not able to assign *M*. sp. 10 to a mating syndrome, because—although we found

received sperm in the antrum and its sperm carry long bristles—it also has a sharp stylet and a simple

antrum. From our previous findings, we would expect this species to have a thickened antrum due to

its interaction with sperm and the mating partner's stylet. This discrepancy could possibly be

attributed to misclassification of the antrum morphology, since *M*. sp. 10 has very pronounced shell

glands, making it difficult to see the anterior part of the antrum, possibly obscuring a thickening or

cellular valve (see specimen IDs MTP LS 788 and MTP LS 801 for a possibly thin cellular valve).

Hypodermic insemination and sperm morphology

In addition to the changes in sperm design mentioned above, we tested whether hypodermic

insemination is associated with a change in sperm length using phylogenetic least squares (PGLS)

regression. We used received sperm location, sperm bristle state, antrum state, and the inferred mating

syndrome as predictors and the $\log_{10}$ transformed sperm length as the response variable. In all cases,

the states that indicate the reciprocal mating syndrome were associated with longer sperm, with the

largest effect for antrum state, followed by the inferred mating syndrome (Fig 9). This seems

reasonable, since the bristle type falsely classified *M*. sp. 68 and *M*. sp. 82 as hypodermically mating,

while the received sperm location and inferred mating syndrome analyses had lower samples size

than that with antrum state. The predictive value of the PGLS models was high, indicating that a large

proportion of the variation in sperm length is explained by the phylogeny and these indicators of the

reciprocal mating syndrome (Fig 9, Tab S6). Note that despite these strong associations, there is

considerable overlap in sperm length between the species exhibiting the different states, with some

species with the reciprocal mating syndrome having short sperm (Fig 9, Tab S6) and an overall 6.7-

fold variation in sperm length across all species (mean length ranging from 25.6 to 173.1 µm).

Coevolution of male and female genitalia

To investigate coevolution between male and female genital traits, we independently summarised five

male and four female genital traits using pPCA. Stylet PC1 was positively loaded with stylet length

and the width of the distal opening and negatively loaded with distal asymmetry. Therefore, high

values of Stylet PC1 represent a more elongate stylet with a wider and less sharp distal opening (Fig

10A, Tab S7). Antrum PC1 was positively loaded with all input variables, meaning that large values

represent more complex female genitalia (Fig 10B). A PGLS regression of Stylet PC1 on Antrum

PC1 across all species revealed a significant positive relationship (Fig 10C). This relationship closely

matches the loadings on PC1 in the pPCA analysis of all reproductive traits (see Fig 5) and could

therefore be mainly driven by the simplification of the antrum in hypodermically mating species.

Therefore, we repeated the analysis on the subset of species assigned to the reciprocal mating

syndrome. We again found a positive relationship between Stylet PC1 and Antrum PC1, indicative

of male-female coevolution of genital morphology among the reciprocally mating species (Fig 10C).

Discussion

Across the genus *Macrostomum*, hypodermic insemination has evolved independently at least 9 times when assessed based on the location of received sperm, and at least 13 times when our more inclusive inferred mating syndrome is considered. The frequent origins of this type of traumatic insemination is remarkable considering that according to Lange et al. [17] merely 12 and 11 such origins have to date been documented in gonochorists and simultaneous hermaphrodites, respectively (including the two cases previously documented in *Macrostomum*). Our detailed analysis of a single genus of free-living flatworms thus approximately doubles the number of documented origins of hypodermic insemination among hermaphrodites. Moreover, three additional origins of traumatic insemination have also recently been documented in the Macrostomorpha [77], the parent group of the genus *Macrostomum*, suggesting that the origin of traumatic insemination occurs commonly in the Macrostomorpha, and maybe other groups of free-living flatworms.

The majority of the collected species are likely undescribed, and a large proportion of the diversity in this genus is yet to be discovered. Since free-living flatworms remain understudied in many regions of the world [78], we have likely not yet documented all convergent origins of hypodermic insemination within *Macrostomum*. As a case and point, the deeply split placement of *M.* sp. 15 (from South Africa) and *M.* sp. 118 (from Southern France) suggests the existence of an additional clade, possibly hinting at large amounts of additional molecular diversity. While we have no observations on the sperm morphology of *M.* sp. 15 or *M.* sp. 118, the stylet morphology of *M.* sp. 15 is suggestive of hypodermic insemination, and closer investigations of this clade would be interesting since this could facilitate inference about the ancestral state leading to the reciprocal clade.

Frequent convergent evolution of this extreme type of mating bolsters the interpretation that it represents an adaptive resolution to sexual conflict over mating rate, mating role or both [12, 18, 28, 39, 41, 46]. Hypodermic insemination likely is an alternative strategy in an ongoing evolutionary chase between donor and recipient, with donor persistence traits, such as complex sperm with bristles

and manipulative seminal fluids, and recipient resistance traits, such as the suck behaviour and

complex female genitalia, engaged in constant antagonistic coevolution [12, 17, 20, 46]. Moreover,

this coevolution does not only seem to drive the frequent origin of hypodermic insemination, but also

appears to result in the emergence of multiple morphological innovations, such as modifications of

stylet, sperm and antrum morphology (see also below).

Interestingly, we find no clear evidence for reversals back to reciprocal mating once hypodermic

insemination has arisen. However, while a Dollo model was preferred in all our ancestral state

reconstructions, this evidence was not completely decisive, as alternative models also received at

least some support. Reciprocal copulation clearly is the ancestral state of the reciprocal clade, but the

state of the most recent common ancestor of the genus is less clear, allowing for either a gain or a

loss (Fig S3). Similarly, in the finlandense clade we either have two independent losses with *M*. sp. 12

and *M*. sp. 44 retaining the ancestral state, or a single loss with a gain in these two sibling species

(Fig S3).

Since hypodermic insemination fundamentally alters the nature of the mating interaction, it might be

difficult for a species to revert to copulation. Specifically, once copulation no longer involves

inserting the stylet into the antrum, a reversal back to copulation would presumably require both

mating partners to again engage in a reciprocal behaviour. Also, a simplification of the antrum could

further hinder reversals, since reciprocally copulating species have traits that presumably reduce the

risk of injury (e.g. a thickened antrum epithelium). If these traits have been lost and are missing in

hypodermically mating species, then occasional reciprocal copulations could result in high fitness

costs for both partners.

In contrast, hypodermic insemination can presumably be performed unilaterally and might not require

the cooperation of both partners, making an evolutionary transition in that direction more likely.

Consequently, the origin of hypodermic insemination likely is a one-way process that canalises taxa

into this mating behaviour. However, across macroevolutionary time recipients may evolve

secondary female genitalia to avoid costs and regain control over the received ejaculate [17]. Why

this has not occurred in *Macrostomum* is unclear, but it might imply that costs of hypodermic

insemination are generally low (possibly due to the striking regeneration ability of these flatworms

[79]) or that the location of insemination is too variable for the evolution of a localised novel organ.

The analysis of correlated evolution clearly shows a strong association between both the sperm bristle

and antrum state with the received sperm location. While on first sight this might simply appear to

represent a confirmation of previous findings [41], albeit with more species and consequently a larger

Bayes factors, it actually represents a very important qualitative improvement of the evidence for

these associations. This is due to how tests of correlated evolution generally, and the Pagel test

implemented in the BayesTraits analysis specifically, are set up. Tests of correlated evolution evaluate

evidence for convergence, by testing whether a model, in which the evolution of the state of one trait

is dependent of the state of the other trait, is more likely than a model where such shifts occur

independently. While these tests supposedly correct for phylogenetic dependencies, they, somewhat

counterintuitively, can support the dependent model of evolution even with only a single

(unreplicated) origin of the trait states in question, provided that these origins map to appropriate

nodes in the phylogeny (as recently outlined: [80, 81]). However, descendants of a clade often share

multiple traits, many of which may not be functionally linked (as in the example given by [80] of the

joint presence of fur and middle ear bones in mammals). Naturally, one aim of such a correlation

analysis is to explore whether there might be a causal relationship between the traits in question.

Thus, while the previous findings with two independent origins of hypodermic insemination in

Macrostomum could be considered evidence for correlated evolution [41], that evidence is not as

decisive as the large Bayes factors may have suggested. By sampling many more convergent events,

we here could remedy the limitations of these earlier results, giving us substantially more confidence

that our findings indeed reveal a causal link between these traits and allowing us to evaluate the

consequences of shifts to hypodermic insemination in a more quantitative context.

In their earlier study [41] posited that specific combinations of adaptations were necessary for

efficient hypodermic insemination, consisting of a needle-like stylet and sperm adapted for movement

through tissue. As a consequence of hypodermic insemination, the antrum should then become

simpler, since it no longer interacts with the sperm and stylet of the partner, instead being only used

for egg-laying. While the data of [41] was suggestive, the issue was that the second hypodermic origin

consisted only of *M. hystrix*, which made quantitative analysis difficult. The principal component

analysis we performed here shows that species with hypodermic insemination indeed have similar

values of PC1, with such values corresponding tightly to the mating syndromes described by [41]

(but note that we here slightly adjusted the definitions of the syndromes, since we lacked behavioural

observations for most species). We show that hypodermic insemination is indeed associated with a

distinct syndrome of reproductive traits that have convergently evolved many times in this genus.

Therefore, these traits likely constitute predictable adaptations to this type of mating.

We observed several species with an unclear or intermediate mating syndrome, and these species

suggest a possible route to hypodermic insemination via the initial evolution of a traumatic stylet in

reciprocally mating species. A sharp stylet could provide anchorage during copulation, as potentially

occurs in *M. spirale* (pers. obs.) and *M. hamatum* (pers. comm. P. Singh), but it may also serve to

stimulate the partner during copulation, or aid in the destruction or removal of rival sperm already

present in the partner's antrum. Finally, internal wounding by the stylet may help to embed sperm in

the antrum wall and/or cellular valve to prevent their removal, either by rival mating partners or by

the recipient during the suck behaviour. However, the fact that we observe many *Macrostomum*

species with stylets with blunt distal thickenings (Fig 3) suggests that such internal wounding may

not always be advantageous for the donor and that these structures may have evolved to avoid harm

to the recipient [41]. Irrespective of the initial selective advantage that internal wounding may confer,

it could then evolve further to complete internal traumatic insemination, and eventually complete

avoidance of the female genitalia and hypodermic insemination via the epidermis.

Accidental sperm transfer due to copulatory wounding has generally been suggested as a possible

route from copulation to traumatic insemination (e.g. in traumatically mating bedbugs) [17, 20]. In

bedbugs, the attachment during mating is a two-step process, indicating that traumatic insemination

was evolutionarily preceded by traumatic penetration for attachment [17]. Similar transitions have

also been proposed for *Drosophila* species in the melanogaster group, where extragenital wounding

structures are typically used for anchorage during copulation. In traumatically inseminating

Drosophila, these structures are modified and pierce the mating partner's integument to inject sperm

into the genital tract [55, 82]. In this light our findings are remarkable, because previous examples

only compared species with and without traumatic insemination, whereas we potentially observe

species "in the act" of transitioning to traumatic insemination. These intermediate species should be

excellent targets for future studies of the costs and benefits of traumatic insemination, especially

because the two candidates represent independent evolutionary transitions.

The effects of traumatic insemination on sperm morphology are mostly unknown, and we are only

aware of the previous study on hypodermic insemination in *Macrostomum* [41]. We here present

quantitative evidence for a shift in sperm length under hypodermic insemination. The PGLS analysis

showed that, on average, the sperm of species with the hypodermic mating syndrome are significantly

shorter. There are several possible explanations for these findings. First, hypodermic insemination

avoids the recipient's genitalia, and these potentially allow both cryptic female choice (e.g. via the

suck behaviour), and sperm displacement or removal by competing donors. Hypodermic insemination

could therefore alter the mode of postcopulatory sexual selection and particularly sperm competition.

This is because postcopulatory sexual selection can introduce skews in sperm representation [35, 37,

38, 83], which likely results in lower sperm competition compared to a "fair-raffle" type sperm

competition, where sperm are expected to mix more freely [32, 43, 44]. Such skews seem less likely

to occur under hypodermic insemination. If sperm size trades-off with sperm number [42–44] then

more intense sperm competition under hypodermic insemination will likely favour the evolution of

smaller sperm [40, 41].

Second, sperm in *Macrostomum* is quite large compared to the size of the antrum, and it intimately

interacts with its epithelium, often being partially embedded in the cellular valve with the feeler [39,

41, 60, 84]. Sperm is also in close contact with rival sperm when animals mate multiply [85, 86].

Under such conditions of high sperm density, i.e. when sperm displacement is likely (e.g. [87–89]),

sperm are predicted to be bigger compared to species in which the sperm storage organ is substantially

larger than the sperm [45, 90]. While in species with hypodermic insemination the sperm will still

intimately interact with the partner's tissue, the "storage organ" could now potentially include the

whole body of the recipient and would then be considerably larger compared to the antrum. Third,

sperm size could decrease due to natural selection, if small sperm are better able to move through the

dense parenchymal tissue of the mating partner [41]. We know little about sperm movement within

the recipient's tissues, but it seems analogous to the movement one can observe within the antrum,

namely via undulation of the sperm body using cortical microtubules [91]. Presumably, smaller sperm

need less energy to overcome tissue resistance and thus might move more efficiently.

Note, that these three explanations are not mutually exclusive, and their relative importance might

depend on the physiology, morphology, and ecology of each species. Even though sperm morphology

is exceptionally diverse, little is known about its functional significance [92]. Because traumatic

insemination originates frequently, it offers an exciting opportunity to disentangle how natural and

sexual selection shape sperm morphology. Such investigations can contribute to an integrative view,

considering not only the sexual selection perspective, but also the direct functional constraints a

sperm's ecology imposes (e.g. survival in sperm storage vs. movement through tissue) [93].

To disentangle mechanisms shaping sperm length evolution, we should ideally investigate the sperm

morphology of other species with traumatic insemination and make use of natural variation in the

location of sperm injection. For example, in bedbugs, the elaboration of the sperm receiving organ

varies considerably from just being a slightly thickened epithelium to a complex spermalege [27, 94].

If movement efficiency is a crucial constraint, we might expect a negative correlation between sperm

length and tissue transit time. Also of interest are comparative investigations of sperm length in taxa

with species with traumatic insemination directly into the recipient's reproductive tract (e.g. the fly

Drosophila parabipectinata [55] or the spider *Harpactea sadistica* [33]), because here movement

through tissue is absent and presumably factors related to sexual selection dominate.

Besides changes in sperm length, we strengthen earlier findings that hypodermic insemination is

associated with the loss of sperm bristles [41]. Furthermore, we document hypodermic received sperm

in species with reduced bristles, indicating that hypodermic insemination likely precedes the complete

loss of bristles. The ancestral state reconstruction using three sperm bristle states found support for

an ordered evolutionary model, suggesting a transition via an intermediate state may be the rule. If

bristles only confer a selective advantage in reciprocally mating species, we could expect them to be

lost through drift once hypodermic insemination evolves, since these structures are no longer needed

to persist against the suck behaviour. Moreover, sperm bristles might also be selected against in

hypodermically mating species if they result in costs for the donor, such as a reduced rate of

spermatogenesis of complex sperm. Indeed, spermatogenesis of the complex sperm with bristles of

M. lignano takes about six days, which is two days longer than the development of the simpler sperm

in *M. pusillum* [95–97]. But note that this could also be due to sperm length differences between these

species, since longer sperm have been associated with longer development times [98, 99]. Moreover,

hypodermic sperm that carry bristles could also have reduced mobility [41].

The fact that there are hypodermically mating species with reduced bristles suggests that selection

might only be efficient at reducing bristle size until they no longer result in strong costs. Short bristles

might not hinder movement much, and once the bristles are reduced to such an extent that they do not

protrude much from the sperm, they could potentially be retained across longer evolutionary times.

The sperm of a member of the *M. pusillum* species-complex in the hypodermic clade contains

electron-dense bodies [100] that are similar to the bristle anchor structures identified in the

reciprocally mating *M. tuba* and *M. lignano* [91, 101]. If these structures are indeed remnants of

bristles, this would support the hypothesis (in agreement with our ancestral state reconstruction) that

bristles are symplesiomorphic in *Macrostomum*—with bristle loss as the derived condition—

suggesting that such greatly reduced bristles are not costly or detrimental to sperm function.

Moreover, sperm bristles have not been observed in three species of *Psammomacrostomum* (pers.

obs.), the sister taxon of *Macrostomum* in the Macrostomidae [77], nor in a presumably closely

associated genus (e.g. *Dunwichia* [102]). Sperm bristles thus appear to be a novel trait that is restricted

to the genus *Macrostomum* (and some closely associated and taxonomically problematic genera that

should be synonymised with *Macrostomum*; see SI Taxonomy). However, more detailed

investigations of sperm ultrastructure of species in the hypodermic clade and other Macrostomorpha

are needed to evaluate this hypothesis.

Our data further suggest that the correlation between reduced/absent bristles and the hypodermic

mating syndrome is not perfect, because we document two species without sperm bristles, that very

likely mate through reciprocal copulation (*M*. sp. 68 and *M*. sp. 82; black arrows in Fig 5). In both

species, we have observed sperm in the antrum that in both is very muscular and complex (**Fig 6**).

We speculate that sperm is deposited deeply inside the antrum, so that sperm bristles may no longer

serve an anchoring function in these species. Additionally, the sperm of *M*. sp. 82 has a peculiar feeler

(Fig 6), which could represent an adaptation to this complex antrum morphology. It is also not clear

if these species perform the suck behaviour, as this was not observed in mating observations of

M. sp. 82 (pers. comm. P. Singh), while we currently have no data at all on the mating behaviour of

M. sp. 68. The loss of sperm bristles in reciprocally mating species should caution us against a too

hasty conclusion that bristle loss is adaptive. Instead, bristles could be lost simply due to drift because

they no longer serve a function.

We also find clear evidence for male-female genital coevolution, both across all species and when

the analysis is restricted to species assigned to the reciprocal mating syndrome. This evidence is in

line with previous findings of such coevolution in hermaphrodites (e.g. [53, 103]) and contributes to

a growing body of evidence that male-female genital coevolution frequently occurs in both

hermaphrodites and gonochorists [104–107]. Genital coevolution is expected both under scenarios of

sexual selection and sexual conflict. Under the sexual selection perspective, we expect coevolution

due to cryptic female choice, where the recipient will choose based on genital traits of the donor

[108]. Donors are therefore selected to closely match their genital morphology to the selection criteria

of the recipient. Similarly, we expect coevolution to result from sexual conflict, since persistence and

resistance traits of the donor and recipient will coevolve due to sexually antagonistic selection [8].

The number and nature of the traits involved in this coevolution, as well as their respective selective

optima, might potentially differ between species, driving trait diversification and speciation [109,

110].

We see signatures of evolutionary diversification across the genus, with the emergence of several

novel and clade-specific traits. Five related species in the spirale clade carry lateral stylet protrusions,

which might serve to anchor the donor during copulation or allow sperm removal, and these species

have complex antra and an unusual sperm design with only a single bristle. Moreover, several species

in the tanganyika clade have long sperm, which could constitute a novel persistence trait serving a

similar function as is hypothesised for the sperm bristles. It appears possible that these long structures,

which are probably greatly elongated sperm feelers, can embed themselves in the recipient's antrum.

They may thereby anchor the sperm and help to prevent removal, making bristles less important to

resist the suck behaviour. Indeed, several of these species have quite short bristles (including the

above-mentioned *M*. sp. 68 that lacks bristles entirely), in spite of showing the reciprocal mating

syndrome, thus supporting this interpretation. Other novel traits include the peculiar sperm feeler of

M. sp. 82, the novel sperm appendages in *M. distinguendum*, and the frequent emergence of a sperm

velum, the function of which is currently completely unknown.

While coevolution can drive trait diversification, it is also interesting to consider cases of

convergence. Repeated origins of a trait indicate that it is a readily available target of selection.

Besides hypodermic insemination, another example of convergent evolution in *Macrostomum* is the

origin of a second female genital opening. The phylogeny suggests that a second female genital

opening has evolved at least four times independently within the genus (see SI Female openings for

a more detailed discussion). In all species, the novel second opening is associated with a muscular

bursa that could allow cryptic female choice, since it might eject sperm through muscular contraction.

Such contraction is also at play during the suck behaviour, where, at least in *M. hamatum*, sperm can

be seen to be ejected from the antrum even before the worm places its mouth on the female genital

opening (pers. comm. P. Singh).

In summary, our findings document a dynamic evolutionary history of male-female coevolution

driving frequent innovations. Such coevolution does not only lead to trait diversification, but likely

also drives the frequent origin of traumatic insemination, because it allows donors to overcome pre-

and postcopulatory choice and/or resistance mechanisms of the recipient. The drastic and repeatable

morphological changes associated with traumatic insemination indicate a shift in functional

constraints on sperm morphology and function, which are presumably not unique to these flatworms,

and which should therefore also be studied in gonochoristic models for traumatic insemination. Such

studies will allow us to disentangle specific and general factors advancing our general understanding

of sperm form and function.

Material and Methods

Field collections and documentation

Almost all specimens were collected from the field in freshwater, brackish or marine habitats, either from sediment samples or from water plants (details on all sampled specimens and their measurements in **Tab S2**). We documented specimens extensively with digital photomicrography, as previously described [41, 77, 84], using light microscopes (Leica DM2500, Olympus BH2, Leitz Diaplan, Zeiss Axioscope 5) with differential interference contrast (DIC) and equipped with digital cameras (Sony DFW-X700, Sony DFK41 and Ximea xiQ). We collected both images and videos at various magnifications (40x to 1000x), documenting the general habitus and details of internal organs, which is possible due to the small size and high transparency of these organisms. To document sperm morphology we amputated a worm's tail slightly anterior of the seminal vesicle, ruptured the seminal vesicle (as described in [111]), and documented the released sperm in a smash preparation using DIC (and sometimes also phase-contrast microscopy). When possible, we prepared whole-mount permanent preparations of these amputated tails to preserve a physical specimen of the male intromittent organ, the stylet [41, 77]. Finally, we preserved the entire animal, or its anterior portion when amputated, for molecular analysis, in either RNAlater solution (Sigma, stored at 4°C up to a few weeks and then at -80°C) or in absolute ethanol (stored cool for up to a few weeks and then at -20°C).

Morphological data

To characterise morphology, we collected both quantitative (Q) and categorical (C) data from the detailed images and videos of the collected specimens (or from the taxonomical descriptions of the few species we did not collect ourselves). Categorical data were determined on a per species basis, while quantitative data were taken per individual. We measured body size (Q) as the total body area and either measured or scored various aspects of the stylet (Q: length, curviness, width of the proximal

opening, width of the distal opening, and asymmetry of the distal thickening; C: sharpness of the

distal thickening), the sperm (Q: total length, bristle length; C: sperm bristle state, presence of a brush,

and presence of a velum) and antrum (Q: number of genital openings; C: antrum thickness, presence

and thickness of anterior cellular valve, antrum chamber complexity, and an overall compound

measure of antrum complexity). We refer the reader to the SI Morphology section for detailed

explanations of these measures. Morphometric analyses were performed using the software ImageJ

(version 1.51w) and the plugin ObjectJ (version 1.04r), which allows marking structures in the

original images in a non-destructive manner. The pixel length of structures was converted into μm,

by calibrating the different microscope setups using a stage micrometre. For comparative analysis we

transformed body area ($\log_{10}$ of the square-root) and $\log_{10}$ transformed all linear measures (stylet

length, width of the proximal opening, width of the distal opening, sperm length, and bristle length).

The sample sizes for all quantitative measurements are given in **Tab S1** and species averages in Tab

S9.

Sequence data generation

We extracted both DNA and RNA from the RNAlater samples using the Nucleospin XS Kit in

combination with the Nucleospin RNA/DNA Buffer Set (Macherey-Nagel), and we extracted DNA

from the ethanol samples using the DNeasy Blood and Tissue kit (Qiagen, Germany). Extracted DNA

and RNA was stored at -80°C. We amplified a partial *28S rRNA* sequence from DNA samples using

PCR primers ZX-1 and 1500R and, for some fragments, additional nested PCR using primers ZX-1

and 1200R, or 300F and 1500R, with polymerases and cycling conditions as previously described

[41, 61]. We sequenced the resulting fragments from both sides using the PCR primers (Microsynth,

Switzerland), assembled them in Geneious (v 11.1.5) with the built-in assembler (using the Highest

sensitivity setting), with minor manual trimming and correction. For some sequences, we obtained

additional sequences using internal primers 1090F, ECD2 (both [41]).

To generate the RNA-Seq library for *M. clavituba* we extracted RNA from 40 pooled animals using

Tri^{TM} reagent (Sigma) and then prepared the library using the TruSeq® Stranded mRNA kit

(Illumina). And to generate the RNA-Seq libraries for all other selected RNA samples, we used

SMART-Seq v4 (Clontech Laboratories, Inc.) in combination with the Nextera XT DNA library

preparation kit (Illumina). When at least 5ng total RNA was available, we performed the SMART-

Seq v4 protocol with 12 preamplification steps and otherwise we used 1ng and performed 15

preamplification steps. Libraries were checked for quality using a Fragment Analyzer (Agilent) and

then sequenced as 101 paired-end reads on the HiSeq2500 platform (using the HiSeq® SBS Kit v4,

Illumina) at the Genomics Facility Basel of the University of Basel and the Department of Biosystems

Science and Engineering of the ETH Zürich.

Species delimitation

Most collected specimens could not be assigned to a taxonomically described *Macrostomum* species

(http://turbellaria.umaine.edu) [112] and are therefore likely new to science. We present

transcriptome-level information for most species but could not conduct RNA-Seq on all >1600

collected specimens. We, therefore, relied on morphology and the partial *28S rRNA* sequences for

species assignment since this fragment is widely used as a DNA barcode for flatworms (e.g. [77, 113–

115]. We constructed haplotype networks from the partial *28S rRNA* sequences using the TCS

algorithm [116] implemented in the TCS software [117] and chose to delimit species with >3

mutational differences in the network. Such a difference in this *28S rRNA* fragment indicates distinct

species among microturbellarians [115], but it is insufficient to clearly distinguish recently diverged

species within *Macrostomum* [61]. Additional markers like COI are frequently used to detect recent

divergence (e.g. [41, 77]), but despite considerable efforts, we were unable to develop universal

primers (a common issue in flatworms, see e.g. [118]) and individual primer optimisation, as in [61],

was not feasible here. Instead, we chose to also delimit species with ≤3 differences in this *28S rRNA*

fragment if they showed clear diagnostic differences in morphology. We thus err on the side of

lumping specimens, with species with shallow molecular and no diagnostic morphological divergence

being assigned to the same operational species. We provide haplotype networks accompanied by

drawings of the diagnostic features for all species (see SI Haplotype networks).

When the morphology of a species was diagnostic, we sequenced several specimens, when possible

from different sample locations, to confirm that they were indeed molecularly similar and then

assigned additional specimens based on morphology only. When no unambiguous diagnostic traits

could be defined (as was the case for many species of the hypodermic clade) we sequenced all

specimens collected for molecular assignment. We used 668 sequences generated for this study (604

sequences, Accessions: MT428556-MT429159) or from public databases (64 sequences),

representing the available diversity across the genus. We also included 15 sequences from seven

species of our chosen outgroup genus *Psammomacrostomum*. We aligned sequences using MAFFT

(„—genafpair –maxiterate 1000") and generated haplotype networks for the recovered clades. We

removed all columns that contained at least one gap before running TCS, leaving us with an alignment

with 787 sites and 385 variable bases. We thus ignore indels to avoid the generation of cryptic species

solely based on these, since scoring indels is non-trivial, and they are frequently treated as missing

data [119].

Phylogenetics

We used 134 transcriptomes representing 105 species, including four distant outgroups,

Haplopharynx rostratum, *Microstomum lineare*, *Myozonaria bistylifera*, and *Karlingia* sp. 1, three

species from the sister genus, *Psammomacrostomum* (see also [77]), and 98 *Macrostomum* species in

the phylogenomic analysis. This included the four publicly available high-quality transcriptomes of

M. hystrix, *M. spirale*, *M. pusillum* [120], and *M. lignano* [121, 122], as well as 130 *de novo*

assembled transcriptomes. Transcriptomes were assembled as previously described [120] and were

mostly derived from whole single specimens (and a few from anterior fragments only, which may

reduce the transcriptome repertoire), while nine were generated by combining RNA-Seq data sets

from several animals or pooling animals that were collected at the same location and assigned to the

species based on our taxonomic expertise into one sample (for details on the transcriptomes used see

Tab S8). We assessed transcriptome quality using TransRate (version 1.0.2, [123]), which maps the

reads back to the assembly and calculates mapping metrics, and BUSCO (version 2.0, [124]), which

searches for the presence of a curated set of core conserved genes. Specifically, we ran the BUSCO

analysis with the metazoan dataset consisting of 978 genes (version uploaded 2016-11-01). These

BUSCO scores were also used to select one representative transcriptome when multiple

transcriptomes were available for a species (see below). We determined the empirical insert size of

our libraries by mapping the reads to the assemblies using SNAP (version 1.0, [125]) and then

extracting the mean insert size using Picard (version 2.20.2).

To infer a set of orthologous genes (orthologs) we predicted open reading frames (ORFs) for each

transcriptome using TransDecoder (version 5.3.0 [126]) with Pfam searches (version 32.0) to retain

transcripts with predicted proteins and kept only one ORF per transcript using the "--

single_best_only" option. We then clustered predicted proteins with at least 99.5% sequence identity

using the CD-HIT clustering algorithm (version 4.7, [127]). The amino acid sequences where then

processed with OrthoFinder (version 2.2.6, [128]) using the „-os" option to perform a length-adjusted

reciprocal BLAST searches followed by MCL clustering. We processed the resulting set of

homologous genes (homogroups) using modified scripts from the phylogenomic dataset construction

workflow [129]. We aligned all homogroups that contained at least 10 species using MAFFT (version

7.310, „–genafpair –maxiterate 1000" [130]), inferred the best substitution model with ModelFinder

[131], and the gene tree using IQ-TREE (version 1.5.5, [66], command: "-mset DCMut, JTTDCMut,

LG, mtZOA, PMB, VT, WAG –mfreq FU,F –mrate G"). Then we trimmed the gene trees using

"trim_tips.py" to remove tip branches that were longer than two substitutions per site and

"mask_tips_by_taxonID_transcripts.py" to remove monophyletic paralogs by choosing the sequence

with the best representation in the alignment. We split off subtrees with long internal branches using

"cut_long_internal_branches.py" and inferred orthologs using the rooted outgroup method in

"prune_paralogs_RT.py". This method uses known outgroup taxa to root the phylogenies, which then

allows to infer the history of speciation and duplication and extract the most inclusive set of orthologs.

We defined *Haplopharynx rostratum*, *Microstomum lineare*, *Myozonaria bistylifera*, and

Karlingia sp. 1 as the outgroup and all *Macrostomum* and *Psammomacrostomum* as an ingroup

(following the phylogeny of [77]), since the algorithm does not include the defined outgroup in the

output, and we then used *Psammomacrostomum* to root our final ortholog trees. Next, we realigned

the orthogroups using MAFFT and trimmed the alignment with ZORRO [132], discarding any

columns in the alignment with a score of less than five and filtering alignments that were shorter than

50 amino acids after trimming. Finally, we inferred ortholog gene trees with 100 non-parametric

bootstraps with IQ-TREE, by inferring the best fitting model „-mset DCMut, JTTDCMut, LG,

mtZOA, PMB, VT, WAG -mfreq FU,F -mrate E,I,G,I+G". These best fitting substitution models

were later also used for the partitioned maximum-likelihood analysis.

We generated two gene matrices, one matrix containing many genes but a relatively moderate

occupancy (called L for low occupancy) and one with a lower gene number but a higher occupancy

(called H for high occupancy, **Tab 2**). We conducted most analyses on the H alignment for

computational tractability and since missing data can have a negative influence on tree inference,

particularly when using Bayesian methods ([133], but see [134]), and only the less computationally

demanding summary and maximum-likelihood methods were also applied to the L alignment. So for

both alignments, we calculated a maximum likelihood species phylogeny using IQ-TREE ("L-IQ-

TREE" and "H-IQ-TREE") with a partition for each gene and using the best substitution model for

each gene (see above). We calculated 1000 ultrafast bootstraps and conducted an approximate-

likelihood ratio test to assess branch support. We inferred Bayesian phylogenies using ExaBayes

(version 1.5, [68]) for the H alignment ("H-ExaBayes") with partitions for each gene, equal prior

probability on all available amino acid substitution models, and with gamma models for all partitions.

We ran four independent chains retaining every 500th iteration and discarding the first 365,000

iterations as burn-in. We terminated the analysis after 1.46 million generations since the average

deviation of split frequencies between all the chains was below 1%, indicating convergence. We

further assessed convergence using Gelman's R of the likelihoods with the R package coda [136],

which showed three chains had converged, while one appeared stuck on a local peak. Since all chains

quickly converged on the same topology, the local peak probably occurs due to differences in the

substitution models applied, with the HIVB model being present at a low rate in the three converged

chains, but being absent in the fourth. We combined the three converged chains, discarded the fourth

and calculated a consensus tree using quadratic path distance optimisation (using the ls.consensus

function in the R package phytools).

To account for potential issues caused by model misspecification, we also performed an unpartitioned

analysis using the CAT model implemented in PhyloBayes (version 1.5, [69]). Because a full GTR

model of the amino acids was too parameter rich, we ran the tool on the DNA coding sequence of the

H alignment and fit the CAT+GTR model ("H-PhyloBayes"). Two chains were run in parallel on 400

CPUs for two weeks. Due to the high cost of the analysis, we terminated the run after 23,311

iterations, at which point the chains showed an identical topology (after removal of 10,000 iterations

as a burn-in). At this point the realised difference in the likelihoods between the two H-PhyloBayes

chains was high (0.21 with an ESS of 858) suggesting the chains had not converged. However,

Gelman diagnostics for the likelihoods suggested convergence. Our chains could potentially be at a

local peak where the hypodermic clade is paraphyletic, possibly due to poor mixing, potentially

biasing the results. Therefore, we discuss the H-PhyloBayes phylogeny, but exclude it from the

comparative analysis.

To construct phylogenies while accounting for potential incomplete lineage sorting, we ran the quartet

based method implemented in ASTRAL-III (version: 5.6.1, [67]) on both alignments ("L-ASTRAL"

and "H-ASTRAL"). We assessed the level of gene tree – species tree conflict at each node of the

phylogeny using the quartet support score, which gives the proportion of quartet trees induced by the

gene trees that are supporting the species tree topology as opposed to the two possible alternatives

[67]. Strong support for the species tree partition can be interpreted as little disagreement between

the gene trees, while support for the alternate topologies indicates strong gene tree – species tree

conflict.

Furthermore, we chose representative *28S rRNA* sequences from the haplotype network analysis to

infer phylogenetic placement for species without a transcriptome. For species with a transcriptome,

we mostly chose the *28S rRNA* sequence derived from the same specimen (for exceptions see

Tab S8). We aligned the sequences using MAFFT ("--maxiterate 1000 –globalpair"), trimmed the

start and end using trimAl ("–nogaps –terminalonly" [137]) and determined the best substitution

model using ModelFinder with the BIC criterion. We then combined this alignment with the H amino

acid alignment, and analysed it with the best fitting substitution model using IQ-TREE as described

above (referred to as "C-IQ-TREE", called C for combined). For comparative analysis, we

transformed the phylogenies (C-IQ-TREE, H-IQ-TREE and H-ExaBayes) to be ultrametric and with

a root depth of 1 using the penalized marginal likelihood approach [138] implemented in the software

TreePL [139].

We performed all follow-up analyses on the H-IQ-TREE and the H-ExaBayes phylogenies to explore

possible effects of phylogenetic uncertainty and included the C-IQ-TREE phylogeny to incorporate

more species. We pruned the phylogenies to include one representative tip per species, choosing the

transcriptome with the best BUSCO score. We present the results obtained with all three phylogenies,

but primarily discuss the ones obtained with the C-IQ-TREE phylogeny, since all results are

quantitatively similar and qualitatively identical.

After the above analyses were completed, we discovered that some transcripts of the RNA-Seq

libraries constructed with the SMART-Seq v4 cDNA kit contained cDNA synthesis primer

sequences. We did not remove these primers before transcriptome assembly since i) the manufacturer

specifically states they should not occur in RNA-Seq reads if used in combination with the Nextera

XT DNA library preparation kit (Appendix C in SMART-Seq v4 Ultra Low Input RNA Kit for

Sequencing User Manual, Takara Bio Inc. available at: https://www.takarabio.com/assets/a/114825)

and ii) because the relevant primer sequence is proprietary (both confirmed by Takara EU tech

support in November 2020), so that we initially did not have access to its sequence. We only

discovered the offending primer sequence due to ongoing work on two *Macrostomum* genome

projects. Because the analyses presented here had at this point used approximately 650,000 CPU

hours of computation, and since the effect of the primer was likely small, we elected to perform some

follow-up analyses to determine how robust our results were to its removal, rather than opting for a

complete re-assembly and re-analysis of all the data. As we outline in SI Primer removal, the removal

of the cDNA synthesis primer sequences had little effect on the topology and branch lengths of L-IQ-

TREE, H-IQ-TREE and C-IQ-TREE and, therefore, we present the results based on the original

alignments.

Frequent origins of hypodermic insemination

The original definitions of the mating syndromes integrated both morphological and behavioural traits

[41]. Because we here lacked behavioural data for most species, we adapted the definitions, relying

instead on several morphological traits and the observed received sperm location only to derive the

inferred mating syndrome (Tab 3). We assigned species to the hypodermic mating syndrome if we

exclusively found hypodermic received sperm, since this represents strong evidence for hypodermic

insemination, as opposed to species where we observed both hypodermic sperm and received sperm

in the female antrum, which we classify as intermediate (Tab 3). Moreover, because hypodermic

sperm can be difficult to observe, especially in species with low investment into sperm production,

we also assigned species that lacked received sperm observations to the hypodermic mating syndrome

based on their morphology alone, namely when they had a simple antrum, a sharp stylet, and absent

or reduced sperm bristles (**Tab 3**). And while observing received sperm in the female antrum may

not exclude occasional hypodermic insemination, it is a strong indication of the reciprocal mating

syndrome, especially when it occurs in a species with a blunt stylet. We, therefore, assigned all species

with received sperm in the antrum and a blunt stylet to the reciprocal mating syndrome (**Tab 3**). And

since some reciprocally mating species also have a sharp stylet (e.g. *M. spirale*), which could possibly

wound the partner internally during mating (pers. obs.), we also assigned these species to the

reciprocal mating syndrome, provided that we observed received sperm in the antrum, and that they

had sperm with bristles (**Tab 3**). These assignments based on morphology alone are supported by our

analysis of correlated evolution, showing a strong association between the received sperm location

and both sperm bristle state and antrum type, respectively (see Results). The inferred mating

syndrome is therefore a more inclusive classification of hypodermic insemination compared to an

assignment based on received sperm location alone.

We estimated ancestral states of the mating syndrome and three reproductive traits linked to it, namely

received sperm location, sperm bristle state, and antrum state. First, we used binary scorings (see SI

Morphology), equivalent to how they were used in the tests for correlated evolution (see below).

However, since we predicted that losses/reductions of the trait would transition via an intermediate

state, we also reconstructed ancestral states with the inferred mating syndrome, received sperm

location and sperm bristle state scored as trinary states. We estimate the history of transitions using

stochastic character mapping [140] with the R package phytools [141]. We determined the

appropriate transition matrix for reconstruction by fitting MK-models with either all state transitions

with equal rates (ER), with symmetric rates (SYM), with all rates different (ARD), and with a model

without the possibility of gains once the trait is lost (Dollo). For traits with trinary states, we

additionally fit an ordered model, where transitions have to go via the intermediate state (ORD) and

an ordered Dollo model without the possibility of gains once the trait is completely lost, but allowing

transitions back from the intermediate state (ORD-Dollo). We reconstructed ancestral states for all

models with a corrected AIC weight >0.15 (Table 3). We used the fully Bayesian implementation of

stochastic character mapping with a gamma prior on each transition ($\alpha = 1$, $\beta = 1$, this results in a low

prior on the number of transitions) and reconstructed 1000 character histories (10.000 iterations burn-

in followed by 10,000 iterations and retaining every 10^{th} character history). We summarised the

number of transitions as the average number of changes as well as the 95% credible interval.

Correlated evolution

We performed a number of tests of correlated evolution to ask if the numerous convergent changes in received sperm location, sperm bristle state and antrum state were evolutionarily dependent. But since we do not have direct observations of received sperm in all species, we first conducted a correlation test between sperm bristle state and received sperm location, and then tested for correlated evolution between both of these variables and the antrum type. We scored all traits as binary and applied Pagel's correlation test [142] as implemented in BayesTraits3 (available at http://www.evolution.rdg.ac.uk/BayesTraitsV3.0.2/BayesTraitsV3.0.2.html). This test compares the marginal likelihood of a model where the transition probability of the first trait depends on the state of the second trait with a model where the two traits evolve independently. For all analysis we ran four independent MCMC chains for 510 million iterations, discarding the first 10 million iterations as burn-in and retaining every 1000^{th} iteration. We calculated the marginal likelihood of the models using the stepping stone sampler of BayesTraits3 with 1000 power posteriors estimated with 10,000 iterations each. We assessed convergence of the chains by calculating Gelman's R using the coda R package [136] and upon confirming convergence merged the chains for further analysis. To evaluate sensitivity to the prior, we ran all analyses with a uniform prior (U 0 100), an exponential prior (exp 10) and a reversible-jump hyperprior with a gamma distribution between 0 and 1 for both the rate prior and the hyperprior (rjhp gamma 0 1 0 1). Models were compared with Bayes factors using the marginal likelihoods calculated by the stepping stone sampler (i.e. BF=2(logLH$_{dependent}$ − logLH$_{independent}$)). To assess the influence of the phylogeny we conducted these tests on three different phylogenies. This setup resulted in 54 model runs consisting of 216 MCMC chains (three trait pairs x three priors x three topologies x four chains for the independent and the dependent model; see also the Correlated evolution section in the Supporting Information). The number of species included was largest for the correlation between sperm bristle state and antrum type (C-IQ-TREE: 124; H-IQ-TREE & H-ExaBayes: 95), and similar for received sperm location and antrum type (C-IQ-TREE: 100; H-

993 IQ-TREE & H-ExaBayes: 77) and received sperm location and sperm bristle state (C-IQ-TREE: 101;

994 H-IQ-TREE & H-ExaBayes: 76).

Hypodermic insemination and convergence in morphospace

We conducted a multivariate analysis to investigate whether the convergent evolution of hypodermic

insemination is associated with changes in a variety of reproductive traits (see SI Morphology, and

Tables A1-2 and Figures A1-3 therein). For this we summarized all our data on stylet, sperm and

antrum morphology (including both quantitative and categorical data) using principal component

analysis. Since regular principal component analysis assumes independence of observations, an

assumption violated by the phylogenetic relationships of species [143], we calculated a

phylogenetically corrected principal component (pPCA), using the phyPCA function in phytools with

the lambda model. Since we combined data with different scales, we used the correlation matrix for

all calculations. When discussing loadings of principal components we apply an aggressive threshold

of ±0.5, since although this results in erosion of power, it keeps false-positive rate within expectations

[144].

Hypodermic insemination and sperm morphology

To test the influence of hypodermic insemination on sperm length, we performed phylogenetically

corrected ordinary least squared regression (PGLS) with the *gls* function in the R package *nlme*

(version 3.1). We used *gls* because it allowed us to simultaneously incorporate phylogenetic signal in

the residuals and account for variation in the number of measured specimens by using the sample size

of the response as weights. We determined the best fitting evolutionary model for the covariance in

the residuals by comparing corrected AIC of PGLS fitted with Brownian motion, lambda or Ornstein-

Uhlenbeck models. We assessed if the assumptions of the PGLS were met by checking the

distributions of the phylogeny-corrected residuals for normality and profiled the likelihood of the

parameter of the correlation structure (i.e. lambda or alpha). Since R-squared values are problematic

for PGLS models [145] we calculated R_{pred} [146] to show model fits. As predictors, we used the

binary traits included in the test of correlated evolution (received sperm location, sperm bristle state

and antrum state) since they all are strong indicators of hypodermic insemination. Moreover, we also

included the inferred mating syndrome as a predictor, but coded it as binary (hypodermic and

reciprocal), and excluding the intermediate syndrome due to the low sample size of this group.

Analysing all four of these predictors is somewhat redundant, but while received sperm location is

the most direct evidence for hypodermic insemination it results in a reduced sample size (see above)

and the other predictors allow the inclusion of more species.

Coevolution of male and female genitalia

If male and female genitalia coevolve, we expect correlations in aspects of their morphology across

species. Ideally, this is detected by identifying homologous male and female structures in all species

and correlating their shape evolution (e.g. [107]). However, the stylets within the reciprocal clade are

highly variable, making the identification and placement of consistent landmarks challenging.

Similarly, capturing the morphology of the antrum is challenging because these soft internal

structures are highly transparent in live specimens. And while the antrum can be studied in

histological sections, it can appear distorted and contracted due to the fixation, making it difficult to

accurately measure length or volume. With these caveats in mind, we independently summarised

male and female genital complexity using pPCAs (see above) and then performed a PGLS regression

(see above) between the first principal components of each pPCA. We used five stylet traits (stylet

length, curviness, width of proximal opening, width of distal opening, and distal asymmetry) and four

antrum traits (thickness, cellular valve, chamber complexity and number of genital openings). We

performed the PGLS regression both across all species and as well as only including species

categorised as showing the reciprocal mating syndrome, since the species with the hypodermic mating

syndrome were nearly invariant.

Acknowledgment

1041

We thank the numerous people that have helped with field work. Especially, we are grateful for the

help of, in no particular order, Werner Armonies, Benny Glasgow, Mohamed Charni, Edith Zemp,

Bernhard Egger, Peter Ladurner, Gregor Schulte, Floriano Papi, Kazuya Kobayashi, Christopher

Laumer, Wim Willems, Tom Artois, Christian Lott, Miriam Weber, Ana-Maria Leal-Zanchet, Kaja

Wasik, Mariana Adami, Walter Salzburger, Adrian Indermaur, Bernd Egger, Fabrizia Ronco, Heinz

Büscher, Victoria Huwiler, Philipp Kaufmann, Michaela Zwyer, Stefanie von Fumetti, Joe Ryan,

Mark Q. Martindale, Marta Chiodin, John Evans, Leigh Simmons, Mauro Tognon, Piero Tognon,

Cristiano Tognon, Pragya Singh, Nikolas Vellnow, Christian Felber, Ulf Jondelius, Sarah Atherton,

Tim Janicke, Georgina Rivera-Ingraham, Ben Byrne, Yvonne Gilbert, Rod Watson, Jochen Rink,

Miquel Vila-Fare, Helena Bilandžija, and Sasho Trajanovski. We thank Katja Eschbach of the

Genomics Facility Basel for preparing and sequencing RNA-Seq libraries. We thank Peter Fields and

Lukas Zimmermann for IT advice. We thank Jürgen Hottinger, Daniel Lüscher and Yasmin Picton

for administrative and technical support. We thank Dita Vizoso for the use of sperm and stylet

drawings for some of the previously described species and for providing inspiration for the new

drawings. We thank Yu Zhang and his collaborators for kindly providing specimens of

M. baoanensis. We thank Nick Goldman and Ziheng Yang for organising the summer school on

Computational Molecular Evolution, which has helped and inspired the first author to explore

phylogenetics. Calculations were performed, in part, at sciCORE (http://scicore.unibas.ch/) scientific

computing centre at the University of Basel.

Funding

1061

1062 This work was supported by Swiss National Science Foundation (SNSF) research grants

1063 31003A_162543 and 310030_184916 to LS.

Competing Interests

All Authors declare that they have no competing interests.

Data availability

The raw sequencing data generated for this study are available in the NCBI

Sequence Read Archive repository with the following accession: PRJNA635941. The partial *28S*

rRNA sequence are available on NCBI with the following accessions: MT428556-MT429159.

Extensive image and video material of all documented specimens are deposited on Zenodo at:

10.5281/zenodo.4482135. Transcriptome assemblies, gene alignments and phylogenetic trees are

deposited on Zenodo at: 10.5281/zenodo.4543289.

References

1. Bateman AJ. Intra-sexual selection in *Drosophila*. Heredity. 1948;2:349–68.

2. Arnold SJ. Bateman's principles and the measurement of sexual selection in plants and animals. Am Nat. 1994;144:S126–49.

3. Janicke T, Häderer IK, Lajeunesse MJ, Anthes N. Darwinian sex roles confirmed across the animal kingdom. Sci Adv. 2016;2:e1500983–e1500983.

4. Parker GA. The sexual cascade and the rise of pre-ejaculatory (Darwinian) sexual selection, sex roles, and sexual conflict. Cold Spring Harb Perspect Biol. 2014;6:a017509–a017509.

5. Puurtinen M, Ketola T, Kotiaho JS. The Good-Genes and Compatible-Genes Benefits of Mate Choice. Am Nat. 2009;174:741–52.

6. Arnqvist G, Nilsson T. The evolution of polyandry: multiple mating and female fitness in insects. Anim Behav. 2000;60:145–64.

7. Hosken DJ, Stockley P. Benefits of Polyandry: A Life History Perspective. In: Macintyre RJ, Clegg MT, editors. Evolutionary Biology. Boston, MA: Springer US; 2003. p. 173–94. doi:10.1007/978-1-4757-5190-1_4.

8. Arnqvist G, Rowe L. Sexual conflict. Princeton, N.J: Princeton University Press; 2005.

9. Parker GA. Sexual conflict over mating and fertilization: An overview. Philos Trans R Soc B Biol Sci. 2006;361:235–59.

10. Rice WR. Sexually antagonistic male adaptation triggered by experimental arrest of female evolution. Nature. 1996;381:232–4.

11. Arnqvist G, Rowe L. Antagonistic coevolution between the sexes in a group of insects. Nature. 2002;415:787–9.

12. Charnov EL. Simultaneous hermaphroditism and sexual selection. Proc Natl Acad Sci U S A. 1979;76:2480–4.

13. Birkhead TR, Pizzari T. Evolution of sex: postcopulatory sexual selection. Nat Rev Genet. 2002;3:262–73.

14. Wedell N, Hosken DJ. The evolution of male and female internal reproductive organs in insects. In: Leonard JL, Córdoba-Aguilar A, editors. The evolution of primary sexual characters in animals. Oxford ; New York: Oxford University Press; 2010.

15. Morrow EH, Arnqvist G. Costly traumatic insemination and a female counter-adaptation in bed bugs. Proc R Soc B Biol Sci. 2003;270:2377–81.

16. Morrow EH, Arnqvist G, Pitnick S. Adaptation versus pleiotropy: why do males harm their mates? Behav Ecol. 2003;14:802–6.

17. Lange R, Reinhardt K, Michiels NK, Anthes N. Functions, diversity, and evolution of traumatic mating: function and evolution of traumatic mating. Biol Rev. 2013;88:585–601.

18. Schärer L, Janicke T, Ramm SA. Sexual conflict in hermaphrodites. Cold Spring Harb Perspect
Biol. 2015;7:a017673.

19. Reinhardt K, Naylor R, Siva-Jothy MT. Reducing a cost of traumatic insemination: female
bedbugs evolve a unique organ. Proc R Soc B Biol Sci. 2003;270:2371–5.

20. Reinhardt K, Anthes N, Lange R. Copulatory wounding and traumatic insemination. Cold Spring
Harb Perspect Biol. 2015;7:a017582.

21. Benoit JB, Jajack AJ, Yoder JA. Multiple traumatic insemination events reduce the ability of bed
bug females to maintain water balance. J Comp Physiol B. 2012;182:189–98.

22. Tatarnic NJ. Traumatic Insemination and Copulatory Wounding. In: Reference Module in Life
Sciences. Elsevier; 2018.

23. Tatarnic NJ, Cassis G, Siva-Jothy MT. Traumatic insemination in terrestrial arthropods. Annu
Rev Entomol. 2014;59:245–61.

24. Kathirithamby J, Hrabar M, Delgado JA, Collantes F, Dötterl S, Windsor D, et al. We do not
select, nor are we choosy: reproductive biology of Strepsiptera (Insecta). Biol J Linn Soc.
2015;116:221–38.

25. Peinert M, Wipfler B, Jetschke G, Kleinteich T, Gorb SN, Beutel RG, et al. Traumatic
insemination and female counter-adaptation in Strepsiptera (Insecta). Sci Rep. 2016;6:25052.

26. Stutt AD, Siva-Jothy MT. Traumatic insemination and sexual conflict in the bed bug *Cimex
lectularius*. Proc Natl Acad Sci. 2001;98:5683–7.

27. Siva-Jothy M t. Trauma, disease and collateral damage: conflict in cimicids. Philos Trans R Soc
B Biol Sci. 2006;361:269–75.

28. Michiels NK, Newman LJ. Sex and violence in hermaphrodites. Nature. 1998;391:647–647.

29. Sluys R. Sperm resorption in triclads (Platyhelminthes, Tricladida). Invertebr Reprod Dev.
1989;15:89–95.

30. Koene JM. Tales of two snails: sexual selection and sexual conflict in Lymnaea stagnalis and
Helix aspersa. Integr Comp Biol. 2006;46:419–29.

31. Koene JM, Montagne-Wajer K, Roelofs D, Ter Maat A. The fate of received sperm in the
reproductive tract of a hermaphroditic snail and its implications for fertilisation. Evol Ecol.
2009;23:533–43.

32. Parker GA. Sperm competition and the evolution of ejaculates: towards a theory base. In: Sperm
competition and sexual selection. T.R. Birkhead and A.P. Møller. San Diego: Academic Press; 1998.
p. 3–54.

33. Řezáč Milan. The spider *Harpactea sadistica*: Co-evolution of traumatic insemination and
complex female genital morphology in spiders. Proc R Soc B Biol Sci. 2009;276:2697–701.

34. Parker GA. Sperm competition games: raffles and roles. Proc R Soc Lond B Biol Sci.
1990;242:120–6.

35. Charnov EL. Sperm competition and sex allocation in simultaneous hermaphrodites. Evol Ecol.
1996;10:457–62.

36. Greeff JM, Parker GA. Spermicide by females: what should males do? Proc R Soc B Biol Sci.
2000;267:1759–63.

37. van Velzen E, Scharer L, Pen I. The effect of cryptic female choice on sex allocation in
simultaneous hermaphrodites. Proc R Soc B Biol Sci. 2009;276:3123–31.

38. Schärer L, Pen I. Sex allocation and investment into pre- and post-copulatory traits in
simultaneous hermaphrodites: the role of polyandry and local sperm competition. Philos Trans R Soc
B Biol Sci. 2013;368:20120052–20120052.

39. Vizoso DB, Gunde Rieger, Lukas Schärer. Goings-on inside a worm: functional hypotheses
derived from sexual conflict thinking. Biol J Linn Soc. 2010;99:370–83.

40. Schärer L, Janicke T. Sex allocation and sexual conflict in simultaneously hermaphroditic
animals. Biol Lett. 2009;5:705–8.

41. Schärer L, Littlewood DTJ, Waeschenbach A, Yoshida W, Vizoso DB. Mating behavior and the
evolution of sperm design. Proc Natl Acad Sci. 2011;108:1490–5.

42. Parker GA. Selection on non-random fusion of gametes during the evolution of anisogamy. J
Theor Biol. 1978;73:1–28.

43. Parker GA. Why are there so many tiny sperm? Sperm competition and the maintenance of two
sexes. J Theor Biol. 1982;:281–94.

44. Parker GA. Sperm competition games: sperm size and sperm number under adult control. Proc R
Soc Lond B Biol Sci. 1993;253:245–54.

45. Immler S, Pitnick S, Parker GA, Durrant KL, Lüpold S, Calhim S, et al. Resolving variation in
the reproductive tradeoff between sperm size and number. Proc Natl Acad Sci. 2011;108:5325–30.

46. Michiels NK. Mating conflicts and sperm competition in simultaneous hermaphrodites. In: Sperm
Competition and Sexual Selection. T.R. Birkhead and A.P. Møller. Elsevier; 1998. p. 219–54.

47. Anthes N, Putz A, Michiels NK. Sex role preferences, gender conflict and sperm trading in
simultaneous hermaphrodites: a new framework. Anim Behav. 2006;72:1–12.

48. Anthes N. Mate choice and reproductive conflict in simultaneous hermaphrodites. In: Kappeler
P, editor. Animal Behaviour: Evolution and Mechanisms. Berlin, Heidelberg: Springer Berlin
Heidelberg; 2010. p. 329–57.

49. Anthes N, David P, Auld JR, Hoffer JNA, Jarne P, Koene JM, et al. Bateman gradients in
hermaphrodites: an extended approach to quantify sexual selection. Am Nat. 2010;176:249–63.

50. Pélissié B, Jarne P, David P. Sexual selection without sexual dimorphism: Bateman gradients in
a simultaneous hermaphrodite. Evolution. 2012;66:66–81.

51. Michiels NK, Koene JM. Sexual selection favors harmful mating in hermaphrodites more than in
gonochorists. Integr Comp Biol. 2006;46:473–80.

52. Jarne P, Auld JR. Animals mix it up too: the distribution of self-fertilization among
hermaphroditic animals. Evolution. 2006;60:1816–24.

53. Koene JM, Schulenburg H. Shooting darts: co-evolution and counter-adaptation in
hermaphroditic snails. BMC Evol Biol. 2005;5:25.

54. Koene JM, Pförtner T, Michiels NK. Piercing the partner's skin influences sperm uptake in the
earthworm Lumbricus terrestris. Behav Ecol Sociobiol. 2005;59:243.

55. Kamimura Y. Twin intromittent organs of *Drosophila* for traumatic insemination. Biol Lett.
2007;3:401–4.

56. Tatarnic NJ, Cassis G. Surviving in sympatry: Paragenital divergence and sexual mimicry
between a pair of traumatically inseminating plant bugs. Am Nat. 2013;182:542–51.

57. Ramm SA, Vizoso DB, Schärer L. Occurrence, costs and heritability of delayed selfing in a free-
living flatworm. J Evol Biol. 2012;25:2559–68.

58. Ramm SA, Schlatter A, Poirier M, Schärer L. Hypodermic self-insemination as a reproductive
assurance strategy. Proc R Soc B Biol Sci. 2015;282:20150660.

59. Winkler L, Ramm SA. Experimental evidence for reduced male allocation under selfing in a
simultaneously hermaphroditic animal. Biol Lett. 2018;14:20180570.

60. Schärer L, Joss G, Sandner P. Mating behaviour of the marine turbellarian *Macrostomum* sp.:
these worms suck. Mar Biol. 2004;145:373–80.

61. Schärer L, Brand JN, Singh P, Zadesenets KS, Stelzer C-P, Viktorin G. A phylogenetically
informed search for an alternative *Macrostomum* model species, with notes on taxonomy, mating
behavior, karyology, and genome size. J Zool Syst Evol Res. 2020;58:41–65.

62. Patlar B, Weber M, Temizyürek T, Ramm SA. Seminal fluid-mediated manipulation of post-
mating behavior in a simultaneous hermaphrodite. Curr Biol. 2020;30:143–9.

63. Weber M, Patlar B, Ramm SA. Effects of two seminal fluid transcripts on post-mating behaviour
in the simultaneously hermaphroditic flatworm *Macrostomum lignano*. J Evol Biol. 2020;:jeb.13606.

64. Holland B, Rice WR. Perspective: chase-away sexual selection: antagonistic seduction versus
resistance. Evolution. 1998;52:1–7.

65. Brand JN. Specimens from Brand et al. 2020, "Frequent origins of traumatic insemination involve
convergent shifts in sperm and genital morphology". Zenodo.
http://doi.org/10.5281/zenodo.XXXXX. 2020.

66. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015;32:268–74.

67. Zhang C, Rabiee M, Sayyari E, Mirarab S. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics. 2018;19 Suppl 6.

68. Aberer AJ, Kobert K, Stamatakis A. Exabayes: Massively parallel bayesian tree inference for the
whole-genome era. Mol Biol Evol. 2014;31:2553–6.

69. Lartillot N, Rodrigue N, Stubbs D, Richer J. PhyloBayes MPI: Phylogenetic Reconstruction with
Infinite Mixtures of Profiles in a Parallel Environment. Syst Biol. 2013;62:611–5.

70. Vellnow N, Vizoso DB, Viktorin G, Schärer L. No evidence for strong cytonuclear conflict over
sex allocation in a simultaneously hermaphroditic flatworm. BMC Evol Biol. 2017;17:103.

71. Zadesenets KS, Vizoso DB, Schlatter A, Konopatskaia ID, Berezikov E, Schärer L, et al. Evidence
for karyotype polymorphism in the free-living flatworm, *Macrostomum lignano*, a model organism
for evolutionary and developmental biology. Plos One. 2016;11:e0164915.

72. Robinson DF, Foulds LR. Comparison of phylogenetic trees. Math Biosci. 1981;53:131–47.

73. Irisarri I, Singh P, Koblmüller S, Torres-Dowdall J, Henning F, Franchini P, et al. Phylogenomics
uncovers early hybridization and adaptive loci shaping the radiation of Lake Tanganyika cichlid
fishes. Nat Commun. 2018;9:3159.

74. Singh P, Ballmer DN, Laubscher M, Schärer L. Successful mating and hybridisation in two
closely related flatworm species despite significant differences in reproductive morphology and
behaviour. Sci Rep. 2020;10:12830.

75. Pleijel F, Jondelius U, Norlinder E, Nygren A, Oxelman B, Schander C, et al. Phylogenies without
roots? A plea for the use of vouchers in molecular phylogenetic studies. Mol Phylogenet Evol.
2008;48:369–71.

76. Xin F, Zhang S-Y, Shi Y-S, Wang L, Zhang Y, Wang A-T. *Macrostomum shenda* and *M. spiriger*,
two new brackish-water species of *Macrostomum* (Platyhelminthes: Macrostomorpha) from China.
Zootaxa. 2019;4603:105.

77. Janssen T, Vizoso DB, Schulte G, Littlewood DTJ, Waeschenbach A, Schärer L. The first multi-
gene phylogeny of the Macrostomorpha sheds light on the evolution of sexual and asexual
reproduction in basal Platyhelminthes. Mol Phylogenet Evol. 2015;92:82–107.

78. Leasi F, Sevigny JL, Laflamme EM, Artois T, Curini-Galletti M, de Jesus Navarrete A, et al.
Biodiversity estimates and ecological interpretations of meiofaunal communities are biased by the
taxonomic approach. Commun Biol. 2018;1:1–12.

79. Egger B, Ladurner P, Nimeth K, Gschwentner R, Rieger R. The regeneration capacity of the
flatworm *Macrostomum lignano*—on repeated regeneration, rejuvenation, and the minimal size
needed for regeneration. Dev Genes Evol. 2006;216:565–77.

80. Maddison WP, FitzJohn RG. The unsolved challenge to phylogenetic correlation tests for
categorical characters. Syst Biol. 2015;64:127–36.

81. Uyeda JC, Zenil-Ferguson R, Pennell MW. Rethinking phylogenetic comparative methods. Syst
Biol. 2018;67:1091–109.

82. Kamimura Y. Copulation anatomy of *Drosophila melanogaster* (Diptera: Drosophilidae): wound-
making organs and their possible roles. Zoomorphology. 2010;129:163–74.

83. Schärer L. Tests of sex allocation theory in simultaneously hermaphroditic animals. Evolution.
2009;63:1377–405.

84. Ladurner P, Schärer L, Salvenmoser W, Rieger RM. A new model organism among the lower
Bilateria and the use of digital microscopy in taxonomy of meiobenthic Platyhelminthes:

Macrostomum lignano, n. sp. (Rhabditophora, Macrostomorpha). J Zool Syst Evol Res. 2005;43:114–26.

85. Janicke T, Marie-Orleach L, De Mulder K, Berezikov E, Ladurner P, Vizoso DB, et al. Sex allocation adjustment to mating group size in a simultaneous hermaphrodite. Evolution. 2013;67:3233–42.

86. Marie-Orleach L, Janicke T, Vizoso DB, David P, Schärer L. Quantifying episodes of sexual selection: insights from a transparent worm with fluorescent sperm. Evolution. 2016;70:314–28.

87. Miller GT, Pitnick S. Sperm-female coevolution in *Drosophila*. Science. 2002;298:1230–3.

88. Lüpold S, Manier MK, Berben KS, Smith KJ, Daley BD, Buckley SH, et al. How multivariate ejaculate traits determine competitive fertilization success in *Drosophila melanogaster*. Curr Biol. 2012;22:1667–72.

89. Manier MK, Lüpold S, Belote JM, Starmer WT, Berben KS, Ala-Honkola O, et al. Postcopulatory sexual selection generates speciation phenotypes in *Drosophila*. Curr Biol. 2013;23:1853–62.

90. Parker GA, Immler S, Pitnick S, Birkhead TR. Sperm competition games: Sperm size (mass) and number under raffle and displacement, and the evolution of P2. J Theor Biol. 2010;264:1003–23.

91. Willems M, Leroux F, Claeys M, Boone M, Mouton S, Artois T, et al. Ontogeny of the complex sperm in the macrostomid flatworm *Macrostomum lignano* (Macrostomorpha, Rhabditophora). J Morphol. 2009;270:162–74.

92. Birkhead TR, Hosken DJ, Pitnick S. Sperm Biology. 2009.

93. Reinhardt K, Dobler R, Abbott J. An Ecology of Sperm: Sperm Diversification by Natural Selection. Annu Rev Ecol Evol Syst. 2015;46:435–59.

94. Hoogstraal H, Usinger RL. Monograph of Cimicidae (Hemiptera-Heteroptera). J Parasitol. 1967;53:222.

95. Schärer L, Ladurner P, Seifarth C, Salvenmoser W, Zaubzer J. Tracking sperm of a donor in a recipient: an immunocytochemical approach. Anim Biol. 2007;57:121–36.

96. Giannakara A, Schärer L, Ramm SA. Sperm competition-induced plasticity in the speed of spermatogenesis. BMC Evol Biol. 2016;16. doi:10.1186/s12862-016-0629-9.

97. Giannakara A, Ramm SA. Self-fertilization, sex allocation and spermatogenesis kinetics in the hypodermically inseminating flatworm *Macrostomum pusillum*. J Exp Biol. 2017;220:1568–77.

98. Pitnick S, Markow TA, Spicer GS. Delayed male maturity is a cost of producing large sperm in Drosophila. Proc Natl Acad Sci. 1995;92:10614–8.

99. Pitnick S. Investment in Testes and the Cost of Making Long Sperm in Drosophila. Am Nat. 1996;148:57–80.

100. Rohde K, Faubel A. Spermatogenesis of *Macrostomum pusillum* (Platyhelminthes, Macrostomida). Invertebr Reprod Dev. 1997;32:209–15.

101. Rohde K, Watson N. Ultrastructure of spermatogenesis and sperm of *Macrostomum tuba*. J Submicrosc Cytol Pathol. 1991;23:23–32.

102. Faubel A, Blome D, Cannon LRG. Sandy beach meiofauna of eastern Australia (southern
Queensland and New South Wales). I. Introduction and macrostomida (Platyhelminthes). Invertebr
Syst. 1994;8:899–1007.

103. Anthes N, Schulenburg H, Michiels NK. Evolutionary links between reproductive morphology,
ecology and mating behavior in opisthobranch gastropods. Evolution. 2008;62:900–16.

104. Brennan PLR, Prum RO, McCracken KG, Sorenson MD, Wilson RE, Birkhead TR. Coevolution
of Male and Female Genital Morphology in Waterfowl. PLOS ONE. 2007;2:e418.

105. Arnqvist G, Rowe L. Correlated evolution of male and female morphologles in water striders.
Evol Int J Org Evol. 2002;56:936–47.

106. McPeek MA, Shen L, Farid H. The Correlated Evolution of Three-Dimensional Reproductive
Structures Between Male and Female Damselflies. Evolution. 2009;63:73–83.

107. Simmons LW, Fitzpatrick JL. Female genitalia can evolve more rapidly and divergently than
male genitalia. Nat Commun. 2019;10:1312.

108. Eberhard W. Female control: sexual selection by cryptic female choice. Princeton: Princeton
University   Press;   1996.
http://www.degruyter.com/view/books/9780691207209/9780691207209/9780691207209.xml.
Accessed 16 May 2020.

109. Arnqvist G, Edvardsson M, Friberg U, Nilsson T. Sexual conflict promotes speciation in insects.
Proc Natl Acad Sci. 2000;97:10460–4.

110. Ritchie MG. Sexual Selection and Speciation. Annu Rev Ecol Evol Syst. 2007;38:79–102.

111. Janicke T, Schärer L. Sperm competition affects sex allocation but not sperm morphology in a
flatworm. Behav Ecol Sociobiol. 2010;64:1367–75.

112. Tyler S, Schilling S, Hooge M, Bush L F. Turbellarian taxonomic database. 2006.

113. Chambrier A, Zehnder M, Vaucher C, Mariaux J. The evolution of the Proteocephalidea
(Platyhelminthes, Eucestoda) based on an enlarged molecular phylogeny, with comments on their
uterine development. Syst Parasitol. 2004;57:159–71.

114. Chambrier A de, Waeschenbach A, Fisseha M, Scholz T, Mariaux J. A large 28S rDNA-based
phylogeny confirms the limitations of established morphological characters for classification of
proteocephalidean tapeworms (Platyhelminthes, Cestoda). ZooKeys. 2015;500:25–59.

115. Scarpa F, Cossu P, Sanna D, Lai T, Norenburg JL, Curini-Galletti M, et al. An 18S and 28S-
based clock calibration for marine Proseriata (Platyhelminthes). J Exp Mar Biol Ecol. 2015;463:22–
31.

116. Templeton AR, Crandall KA, Sing CF. A cladistic analysis of phenotypic associations with
haplotypes inferred from restriction endonuclease mapping and DNA sequence data. III. Cladogram
estimation. Genetics. 1992;132:619–33.

117. Clement M, Posada D, Crandall KA. TCS: a computer program to estimate gene genealogies.
Mol Ecol. 2000;9:1657–9.

118. Vanhove MPM, Tessens B, Schoelinck C, Jondelius U, Littlewood DTJ, Artois T, et al.
Problematic barcoding in flatworms: A case-study on monogeneans and rhabdocoels
(Platyhelminthes). ZooKeys. 2013;:355–79.

119. Simmons MP, Müller K, Norton AP. The relative performance of indel-coding methods in
simulations. Mol Phylogenet Evol. 2007;44:724–40.

120. Brand JN, Wiberg RAW, Pjeta R, Bertemes P, Beisel C, Ladurner P, et al. RNA-Seq of three
free-living flatworm species suggests rapid evolution of reproduction-related genes. BMC Genomics.
2020;21:462.

121. Wudarski J, Simanov D, Ustyantsev K, de Mulder K, Grelling M, Grudniewska M, et al.
Efficient transgenesis and annotated genome sequence of the regenerative flatworm model
Macrostomum lignano. Nat Commun. 2017;8:2120.

122. Grudniewska M, Mouton S, Grelling M, Wolters AHG, Kuipers J, Giepmans BNG, et al. A
novel flatworm-specific gene implicated in reproduction in *Macrostomum lignano*. Sci Rep.
2018;8:3192.

123. Smith-Unna R, Boursnell C, Patro R, Hibberd JM, Kelly S. TransRate: reference-free quality
assessment of de novo transcriptome assemblies. Genome Res. 2016;26:1134–44.

124. Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, et al. BUSCO
applications from quality assessments to gene prediction and phylogenomics. Mol Biol Evol. 2017.
https://academic.oup.com/mbe/advance-article/doi/10.1093/molbev/msx319/4705839. Accessed 8
Dec 2017.

125. Zaharia M, Bolosky WJ, Curtis K, Fox A, Patterson D, Shenker S, et al. Faster and more accurate
sequence alignment with SNAP. ArXiv Prepr ArXiv11115572. 2011.

126. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. *De novo*
transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation
and analysis. Nat Protoc. 2013;8:1494–512.

127. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 2012;28:3150–2.

128. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. Genome Biol. 2015;16:157.

129. Yang Y, Smith SA. Orthology inference in nonmodel organisms using transcriptomes and low-
coverage genomes: improving accuracy and matrix occupancy for phylogenomics. Mol Biol Evol.
2014;31:3081–92.

130. Nakamura T, Yamada KD, Tomii K, Katoh K, Hancock J. Parallelization of MAFFT for large-
scale multiple sequence alignments. Bioinformatics. 2018;34:2490–2.

131. Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler A von, Jermiin LS. ModelFinder: fast
model selection for accurate phylogenetic estimates. Nat Methods. 2017;14:587–9.

132. Wu M, Chatterji S, Eisen JA. Accounting for alignment uncertainty in phylogenomics. PLOS
ONE. 2012;7:e30288.

133. Roure B, Baurain D, Philippe H. Impact of missing data on phylogenies inferred from empirical
phylogenomic data sets. Mol Biol Evol. 2013;30:197–214.

134. Tan G, Muffato M, Ledergerber C, Herrero J, Goldman N, Gil M, et al. Current methods for
automated filtering of multiple sequence alignments frequently worsen single-gene phylogenetic
inference. Syst Biol. 2015;64:778–91.

135. Wang H-C, Minh BQ, Susko E, Roger AJ. Modeling site heterogeneity with posterior mean site
frequency profiles accelerates accurate phylogenomic estimation. Syst Biol. 2018;67:216–35.

136. Plummer M, Best N, Cowles K, Vines K. CODA: convergence diagnosis and output analysis
for MCMC. R News. 2006;6:7–11.

137. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25:1972–3.

138. Sanderson MJ. Estimating absolute rates of molecular evolution and divergence times: A
penalized likelihood approach. Mol Biol Evol. 2002;19:101–9.

139. Smith SA, O'Meara BC. treePL: divergence time estimation using penalized likelihood for large
phylogenies. Bioinformatics. 2012;28:2689–90.

140. Bollback JP. SIMMAP: Stochastic character mapping of discrete traits on phylogenies. BMC
Bioinformatics. 2006;7:88.

141. Revell LJ. phytools: an R package for phylogenetic comparative biology (and other things):
phytools: R package. Methods Ecol Evol. 2012;3:217–23.

142. Pagel M. Detecting correlated evolution on phylogenies: a general method for the comparative
analysis of discrete characters. Proc R Soc Lond B. 1994;255:37–45.

143. Revell LJ. Size-correction and principal components for interspecific comparative studies.
Evolution. 2009;63:3258–68.

144. Peres-Neto PR, Jackson DA, Somers KM. Giving meaningful interpretation to ordination axes:
assessing loading significance in principal component analysis. Ecology. 2003;84:2347–63.

145. Ives AR. R^2s for Correlated Data: Phylogenetic Models, LMMs, and GLMMs. Syst Biol.
2019;68:234–51.

146. Ives A, Li D. rr2: An R package to calculate R^2s for regression models. J Open Source Softw.
2018;3:1028.

147. Wang A-T. Three new species of the genus *Macrostomum* from China (Platyhelminthes,
Macrostomida, Macrostomidae). Acta Zootaxonomica Sin. 2005;30:714–20.

148. Sun T, Zhang L, Wang A-T, Zhang Y. Three new species of freshwater *Macrostomum*
(Platyhelminthes, Macrostomida) from southern China. Zootaxa. 2015;4012:120–34.

149. Lin Y, Zhou W, Xiao P, Zheng Y, Lu J, Li J, et al. Two new species of freshwater *Macrostomum*
(Rhabditophora: Macrostomorpha) found in China. Zootaxa. 2017;4329:267.

150. Lin Y-T, Feng W-T, Xin F, Zhang L, Zhang Y, Wang A-T. Two new species and the molecular
phylogeny of eight species of *Macrostomum* (Platyhelminthes: Macrostomorpha) from southern
China. Zootaxa. 2017;4337:423.

151. Wang L, Xin F, Fang C-Y, Zhang Y, Wang A-T. Two new brackish-water species of
Macrostomum (Platyhelminthes, Macrostomida) from mangrove wetland in southern China. Zootaxa.
2017;4276:107.

1408

Figures

1410

Fig 1. Representative drawings of the morphology of the stylet (male intromittent organ) (left) and the sperm
(right) of three *Macrostomum* species.
Top: *M.* sp. 92, a hypodermically mating species from the hypodermic clade, with a typical needle-like stylet and a
simple sperm morphology. Middle: The well-studied model organism *M. lignano* with the typical morphology for
reciprocally mating species, showing a stylet with blunt distal thickenings and a complex sperm with an anterior feeler,
two stiff lateral bristles, and a terminal brush. Bottom: *M.* sp. 9 representing one of the convergent origins of
hypodermic insemination in the reciprocal clade, showing a stylet with a highly asymmetric and sharp distal thickening
and sperm with reduced sperm bristles, no brush, but a thin velum along the shaft. Note that, given the striking diversity
across the *Macrostomum* genus, it is not possible to clearly delimit all the sperm traits originally defined in *M. lignano*
in some of the species.

1421

Fig 2. Simplified phylogenies of the genus *Macrostomum*.
Consistently recovered clades are collapsed into named triangles, with the height of the base proportional to the number
of species, and single species names preceded by the genus abbreviation (in italics). Not shown here is the variable
placement of *M.* sp. 39 within the hypodermic clade (see Figure S1). (A) Maximum-likelihood phylogeny (H-IQ-
TREE) (B) Bayesian phylogeny (H-ExaBayes). (C) Summary method phylogeny (H-ASTRAL). Branch length
represents substitutions per site in A & B and coalescent units in C. Support values for these groupings are generally
high (for details see **Fig S1**).

1429

sp. 47
sp. 1
sp. 38
sp. 100
rubrocinctum
sp. 80
sp. 39
hystricinum c
sp. 105
hystricinum a
hystricinum d
sp. 6
hystricinum
sp. 32
sp. 81
sp. 93
sp. 48
sp. 49
gabriellae
sp. 78
pusillum
pusillum a
sp. 22
sp. 97
sp. 95
sp. 92
sp. 99
sp. 104
sp. 15
sp. 118
curvituba
sp. 98
sp. 96
spirale
balticum
axi
meridionalis
tenuicauda
longituba
sp. 2
clavituba
sp. 11
ermini
sp. 23
sp. 102
sp. 13
sp. 30
sp. 43
sp. 29
evelinae
sp. 42
spiriger
sp. 5
mediterraneum
kepneri
sp. 46
baoanensis
sp. 45
sp. 85
cliftonensis
lignano
janickei
sp. 14
sp. 103
sp. 108
mirumnovem
sp. 101
shenzhenensis
mystrophorum
sp. 3
shenda
hystrix
sp. 34
zhujiangensis
sp. 41
qiaochengensis
sp. 36

received sperm
sperm bristle
antrum
sharpness
inferred syndrome

0.2

to Fig. 3B

20µm
stylet

20µm
sperm

1430

from
Fig. 3A

sp. 90
minutum
sp. 54
sp. 112
hamatum
sp. 86
sp. 83
sp. 106
sp. 82
sp. 89
sp. 4
bicaudatum
sp. 37
heyuanensis
rostratum
retortum
paradoxum
sp. 55
lankouensis
inductum
zhaoqingensis
chongqingensis
shiyanensis
sinensis
karlingi
distinguendum
sp. 7
platensis
sp. 51
finlandense
sp. 12
sp. 44
sp. 50
sp. 24
sp. 35
sp. 9
quiritium
tuba
acus
obtusa
sp. 75
sp. 10
sp. 76
sp. 66
poznaniense
sp. 63
sp. 62
sp. 16
sp. 17
sp. 18
gieysztori
sp. 64
sp. 65
sp. 74
sp. 60
sp. 117
sp. 67
sp. 70
sp. 77
sp. 56
sp. 71
sp. 58
sp. 68
sp. 72
sp. 61
sp. 69
sp. 57
sp. 59

received sperm
sperm bristle
antrum
sharpness
inferred syndrome

0.2

no obs.
no obs.
no obs.
stylet ½
no obs.
no obs.
no obs.
no obs.
no obs.
no obs.
no obs.

20μm
stylet

20μm
sperm

1431

Fig 3. Phylogeny of the genus *Macrostomum*, showing the striking diversity in stylet and sperm morphology across the genus.
The ultrametric phylogeny (C-IQ-TREE) includes all 145 studied species (with 77 species depicted in Fig 3A and 68 species in Fig 3B). Branch supports are ultrafast bootstraps (top, black if 100) and approximate likelihood ratio tests (bottom, grey if 100). Species without available transcriptomes that were added based on a *28S rRNA* fragment are indicated with red branches. Columns indicate the states of five reproductive traits from light to dark (i.e. yellow, light green and dark green for trinary states; or yellow and dark green for binary states; grey indicates missing data): received sperm location (hypodermic, both, in antrum), sperm bristle state (absent, reduced, present), antrum state (simple, thickened), sharpness of stylet (sharp, neutral, blunt), inferred mating syndrome (hypodermic, intermediate, reciprocal). Stylet and sperm are drawn based on our live observations, except for species with underlined names, which were redrawn based on the species description (*M. acus*, *M. obtusa* and *M. sinensis* from Wang 2005; *M. heyuanensis* and *M. bicaudatum* from Sun et al. 2015; *M. chongqingensis* and *M. zhaoqingensis* from Lin et al. 2017a; *M. shiyanensis* and *M. lankouensis* from Lin et al. 2017b; *M. shenzhenensis* and *M. qiaochengensis* from Wang et al. 2017; and *M. spiriger* and *M. shenda* from Xin et al. 2019). The stylet of *M*. sp. 15 is not drawn to scale, the stylets of some species are drawn at half size (stylet ½), and the stylet of *M*. sp. 23 is not drawn since it was incomplete (specimen ID MTP LS 913). Unobserved structures are marked as no observation (no obs.).

Fig 4. Results of correlated evolution analysis between (A) received sperm location and sperm bristle state, (B) received sperm location and antrum state, and (C) sperm bristle state and antrum state.
Shown are transition matrices for the dependent model from BayesTraits analysis, which was always preferred over the independent model. Transition rates are scaled so that the largest is unity (and arrow sizes are proportional). Also given are the likelihoods of the independent (Lh_{ind}) and dependent (Lh_{dep}) models, and the resulting BayesFactors (BF). An exponential prior and the C-IQ-TREE phylogeny was used for the results shown here. See SI Correlated evolution for runs with other priors (uniform and reversible-jump hyperprior) and other phylogenies (H-IQ-TREE and H-ExaBayes), which show qualitatively similar results.

1458

Fig 5. Results of a phylogenetically corrected principal component analysis of the measured quantitative and categorical (italics) reproductive traits.
Left: Loadings of PC1 and PC2, with the percentage of variance explained at the bottom. Right: 2D morphospace defined by PC1 and PC2. As indicated by the legend, the shape represents the sperm bristle state, while the colours represent the inferred mating syndrome (left side) and the received sperm location (right side). Species from the hypodermic clade are outlined with stippled lines. Red arrowheads indicate two species (*Macrostomum* sp. 51 and *M*. sp. 89) that cluster closely with species assigned to the hypodermic mating syndrome, but in which we observed received sperm in the antrum. Black arrowheads indicate two species (*M*. sp. 68 and *M*. sp. 82) assigned to the reciprocal syndrome, which have no discernible sperm bristles (see also Fig 6). The phylogenetic relationships of these species are represented as a phylomorphospace animation in Fig S4. Results shown here based on C-IQ-TREE, while detailed results including analyses with other phylogenies (H-IQ-TREE and H-ExaBayes) are in Tab S5**Fig S3**.

1470

1471

Fig 6. Details on the reproductive morphology of *Macrostomum* sp. 68 (A-E) and *M.* sp. 82 (F-J).
M. sp. 68 (A) Stylet drawing showing the blunt distal thickenings; (B) distal stylet tip in a smash preparation (specimen ID MTP LS 2611). (C) Sperm image (MTP LS 2686) and drawing showing what seems to be a long feeler, but no apparent sperm bristles. (D-E). Details of the antrum (MTP LS 2562) indicating the muscular connection between the female genital opening and the antrum (arrowhead in D) and the anterior second chamber containing at least one received sperm (arrowhead in E). *M.* sp. 82 (F) Drawing of the stylet showing the slight blunt distal thickenings. (G) Distal stylet tip *in situ* (top, MTP LS 2845) and in a smash preparation (bottom, MTP LS 2846). (H) Sperm image (MTP LS 2877) and drawing indicating the modified anterior part of the sperm (shaded grey) and a less dense area approximately 1/3 along the sperm, which could be a vestigial bristle anchor location (arrowhead). (I-J) Details of the antrum (MTP LS 2848) indicating the anterior genital opening, the bursa pore (I, left arrowhead) next to the posterior genital opening, the gonopore (I, right arrowhead), both connecting into a large chamber containing many received sperm (J, arrowhead). Scale bars represent 100 µm in the antrum images and 20 µm otherwise.

1484

Fig 7. Detailed observations of received sperm location in *Macrostomum* sp. 3.
(A) Drawing of the sexual organs (te: testis, vd: vas deferens, ov: ovary, eg: developing egg, cv: cellular valve, an:
antrum, fs: false seminal vesicle, ts: true seminal vesicle, st: stylet), with sperm drawn at locations where they were
observed. (B-E) show sperm *in situ*. (B) sperm in the antrum (specimen ID MTP LS 3286) embedded in the cellular
valve (anterior); (C) sperm in cellular valve with loose tissue (MTP LS 3314); (D) sperm close to developing egg (MTP
LS 3314); (E) sperm close to the ovary (MTP LS 3317). Scale bars represent 100µm in (A) and 50 µm otherwise.

1491

1492

Fig 8. Detailed observations of received sperm location in *Macrostomum* sp. 101.
(A) Drawing of antrum region (ov: ovary, eg: developing egg, cv: cellular valve, an: antrum), with sperm drawn at locations where they were observed. (B-D) show sperm *in situ* with anterior on the top. (B) sperm (arrow) embedded in cellular valve (specimen ID MTP LS 3206); (C) sperm (arrow) in cellular valve and antrum (MTP LS 3127); (D) same specimen as C with dislodged sperm (arrow) surrounded by what appears to be cellular valve derived tissue. Scale bars represent 100µm in (A) and 50µm otherwise.

1499

1500

1501

Fig 9. Sperm length of species dependent on (A) sperm bristle state, (B) received sperm location (C) antrum state or (D) inferred mating syndrome.
Values are slightly jittered in the x direction, and the y-axis is on a log-scale. Within each panel the main results of PGLS analysis are given and in all tests the slopes were significant at p <0.001. Detailed results including analyses with different phylogenies (H-IQ-TREE and H-ExaBayes) are given in Tab S6.

1507

Fig 10. Phylogenetically corrected principal component analyses of stylet (A) and antrum (B) traits, and evidence for male-female coevolution (C).
Loadings of Stylet and Antrum PC1, with the percentage of variance explained at the bottom, of the stylet (A) and antrum (B) traits, respectively. Categorical reproductive traits are in italics. (C) Shows the results from PGLS regression between Stylet PC1 and Antrum PC1 from (A-B). Regression was performed across all species (solid line, upper statistics) and restricted to species of the reciprocal mating syndrome (dashed line, lower statistics). The inferred mating syndrome of species is indicated by the colour of the dots. Results based on C-IQ-TREE, detailed results including analysis with other phylogenies (H-IQ-TREE and H-ExaBayes) are in Tab S7.

1518   # Tables

1519

1520   **Tab 1. Summary of the taxonomic status of all the included *Macrostomum* species.**
1521   Species inferred based on single specimens (N=1) are listed separately.

Species in this study	N = 1	N > 1	Total
All	18	127	145
Described	5	46	51
Likely undescribed & immature	3	2	5
Likely undescribed, mature & no transcriptome	5	17	22
Likely undescribed, mature & transcriptome	5	62	67

1522

1523

1524   **Tab 2. Characteristics of the used protein alignments.**
1525   We used one alignment aimed at a high number of genes (L alignment) and one alignment aimed at high occupancy (H
1526   alignment). The alignments used were trimmed to only include regions with a high probability of being homologous
1527   using ZORRO, and the statistics are given for the alignments before and after trimming.

Alignment metrics	L untrimmed	L	H untrimmed	H
No. genes	8128	8128	385	385
No. amino acids AA	5,057,157	1,687,014	200,729	94,625
No. variable sites	3,263,955	1,157,689	135,887	74,175
No. parsimony informative sites	2,287,246	934,803	103,425	63,066
Missing data (%)	78.1	59.3	55.7	22.9

1528

1529

1530

Tab 3. Assignment of the inferred mating syndrome based on different reproductive traits.

Species were assigned to an inferred mating syndrome based on the location of received sperm in the body (antrum, in the antrum only; hypodermic, hypodermic only; both, in the antrum and hypodermic; NA, no observation), the sperm bristle state (absent, reduced or present), the antrum state (simple or thickened), and the shape of the distal thickening of the stylet (sharp or blunt). 26 species with either not enough (22 species) or contradictory (four species) information were not assigned to a syndrome. Note, that all 24 species with only hypodermic sperm had the same morphological states, but this was not a condition for their assignment (hence the brackets). Similarly, all 69 species assigned to the reciprocal mating syndrome had a thickened antrum, but this was not a condition for their assignment.

	Received sperm location				Morphology			
Syndrome	**Antrum**	**Hypodermic**	**Both**	**NA**	**Sperm bristle**	**Antrum**	**Stylet**	**N**
Hypodermic		24			(Reduced/absent)	(Simple)	(Sharp)	24
Hypodermic				18	Reduced/absent	Simple	Sharp	18
Intermediate			2		Reduced	Thickened	Sharp	2
Reciprocal	61			6	Any state	(Thickened)	Blunt	67
Reciprocal	8				Present	(Thickened)	Sharp	8
Unclear	7			19	Other combinations			26

1539

1540

1541

Tab 4. Ancestral state reconstructions of reproductive traits, including received sperm location, sperm bristle state, antrum state, and inferred mating syndrome.
A range of MK-models were compared based on their AIC weights (ER: equal rate, SYM: symmetrical rate, ORD-Dollo: ordered model without gains once the trait is in state 0, Dollo: model without gains, ORD: ordered model, ARD: all rates different). For each trait the model with the highest AICc weight is shown in bold type, but we estimated the number of transitions between the states using stochastic character mapping with 1000 posterior samples for all models with an AICc weight >0.15. Given are the average number of transitions and the 2.5% and 97.5% quantiles in brackets. Results are based on the C-IQ-TREE phylogeny. For the quantitatively similar results with the H-IQ-TREE and H-ExaBayes phylogenies see Table S4.

Reproductive trait	model	dfs	log lik	AICc	ΔAICc	AICc weight	$N_{Species}$	$N_{Changes}$	0→1	1→0	1→2	2→1	0→2	2→0
Received sperm location 0: hypodermic only 1: hypodermic & in antrum 2: in antrum only	ER	1	-43.3	88.6	10.1	0.004								
	SYM	3	-40.9	88.1	9.6	0.005								
	ORD-Dollo	**3**	**-36.1**	**78.5**	**0**	**0.579**	**102**	**21.2 (16, 32)**	**-**	**8.0 (7, 13)**	**4.0 (0, 11)**	**9.3 (7, 13)**	**-**	**-**
	Dollo	4	-36.1	80.6	2.2	0.196	102	10.8 (9, 17)	-	0.8 (0, 4)	0.9 (0, 4)	2.6 (2, 5)	-	6.5 (4, 8)
	ORD	4	-36.1	80.6	2.2	0.196	102	23.4 (16, 38)	0.7 (0, 4)	8.4 (6, 13)	4.8 (0, 15)	9.6 (7, 15)	-	-
	ARD	6	-36.1	85.1	6.6	0.021								
Received sperm location 0: hypodermic 1: in antrum only	ER	1	-36.8	75.6	3.1	0.135								
	Dollo	**1**	**-35.2**	**72.5**	**0**	**0.638**	**102**	**9.3 (9, 11)**	**-**	**9.3 (9, 11)**				
	ARD	2	-35.2	74.5	2.1	0.227	102	9.9 (9, 14)	0.9 (0, 4)	9.0 (8, 12)				
Sperm bristle state 0: absent 1: reduced 2: present	ER	1	-82.9	167.8	27.2	0								
	SYM	3	-81.7	169.5	28.9	0								
	ORD-Dollo	**3**	**-67.2**	**140.6**	**0**	**0.578**	**131**	**36.3 (29, 48)**	**-**	**12.2 (11, 17)**	**6.7 (1, 16)**	**17.5 (14, 23)**	**-**	**-**
	Dollo	4	-67.2	142.7	2.1	0.2	131	29.3 (22, 41)	-	7.1 (3, 13)	5.3 (1, 13)	12.4 (8, 18)	-	4.6 (0, 8)
	ORD	4	-67.2	142.7	2.1	0.199	131	37.1 (29, 50)	0.8 (0, 5)	13.0 (10, 20)	6.0 (1, 14)	17.3 (14, 22)	-	-
	ARD	6	-67.2	147.1	6.5	0.023								
Sperm bristle state 0: absent or reduced 1: present	ER	1	-57.6	117.3	5.7	0.038								
	Dollo	**1**	**-54.8**	**111.6**	**0**	**0.635**	**131**	**18.8 (18, 22)**	**-**	**18.8 (18, 22)**				
	ARD	2	-54.4	112.9	1.3	0.327	131	20.5 (17, 29)	2.7 (0, 9)	17.8 (15, 23)				
Antrum state 0: simple 1: thickened	ER	1	-48.1	98.2	3.8	0.086								
	Dollo	**1**	**-46.2**	**94.4**	**0**	**0.577**	**127**	**14.7 (14, 17)**	**-**	**14.7 (14, 17)**				
	ARD	2	-45.7	95.5	1.1	0.337	127	15.2 (13, 21)	2.1 (0, 6)	13.1 (11, 16)				
Inferred mating syndrome 0: hypodermic 1: intermediate 2: reciprocal	ER	1	-59.3	120.6	16.2	0								
	SYM	3	-54.4	114.9	10.4	0.003								
	ORD-Dollo	**3**	**-49.1**	**104.5**	**0**	**0.578**	**119**	**34.1 (27, 47)**	**-**	**13.7 (12, 18)**	**7.1 (2, 16)**	**13.2 (10, 19)**	**-**	**-**
	Dollo	4	-49.1	106.6	2.1	0.199	119	16.6 (14, 25)	-	0.7 (0, 5)	1.1 (0, 6)	2.8 (2, 6)	-	11.9 (9, 15)
	ORD	4	-49.1	106.6	2.1	0.198	119	35.4 (27, 50)	1.0 (0, 5)	14.3 (11, 20)	7.2 (2, 16)	12.9 (9, 18)	-	-
	ARD	6	-49.1	111	6.5	0.022								
Inferred mating syndrome 0: hypodermic & intermediate 1: reciprocal	ER	1	-49.4	100.9	14.675	0.001								
	Dollo	**1**	**-42.1**	**86.26**	**0**	**0.997**	**119**	**14.7 (14, 17)**	**-**	**14.7 (14, 17)**				
	ARD	2	-47.3	98.68	12.421	0.002								

Supp. Figures

Fig S1. Phylogenetic trees inferred using various methods on the L and H alignments.
Clades are coloured consistently between the panels, with six large species groups (hypodermic, dark grey; spirale, orange; lignano, dark green; finlandense, purple; tuba, pink; tanganyika, brown), two smaller species groups (minutum, yellow; hamatum, light green), and two consistent species pairs (*M*. sp. 45 + 46, light grey; *M*. sp. 4 + 89, dark blue). Transcriptomes that are placed differently between approaches are highlighted in red. Species assignments are given as a short six letter abbreviation in the name of the transcriptomes (three letters each for the genus and species, respectively). See

Tab S8 for details.

see the file Fig_S1.pdf

Fig S2. The two ASTRAL phylogenies with quartet support shown for each node.
(A) L-ASTRAL phylogeny, and (B) H-ASTRAL phylogeny. The pie charts at the nodes represent the quartet support for the three possible partitions at that node. Blue represents support for the species tree topology and orange and red represent support for one or the other of the two alternative partitions. Mostly blue pie charts indicate little conflict. This analysis indicates that there is substantial gene tree – species tree conflict throughout the genus.

see the file Fig_S2.pdf

Fig S3. Ancestral state reconstructions of reproductive traits using the C-IQ-TREE phylogeny.
The trait and type of scoring (binary/trinary) is indicated at the bottom of each panel. Stochastic character mapping is summarised with pie charts representing the proportion of stochastic maps with the respective state. Shown is the reconstruction of the best-fitting ordered model without losses. The average number of transitions is given in Tab 4, while the red stars and numbers indicate the lower-bound number of transitions that have likely occurred (i.e. separated by nodes with >95% posterior probability of the ancestral state), while acknowledging that the ancestral state of the genus is often unclear (hence the brackets). The colours indicate the six large species groups (hypodermic, dark grey; spirale, orange; lignano, dark green; finlandense, purple; tuba, pink; tanganyika, brown), two smaller species groups (minutum, yellow; hamatum, light green), and two consistent species pairs (*M*. sp. 45 + 46, light grey; *M*. sp. 4 + 89, dark blue).

see the file Fig_S3.pdf

Fig S4. Animation of the phylomorphospace represented by PC1 and PC2 of the species in the C-IQ-TREE phylogeny.
The animation initially shows a cladogram that then gradually transforms into the phylomorphospace, which was calculated using the phylomorphospace function in phytools.

see the file Fig_S4.gif

Supp. Tables

Tab S1. The number of specimens analysed per *Macrostomum* species for all the included quantitative traits.

see the file Tab_S1.xlsx

Tab S2. Details on all specimens included in this study.

see the file Tab_S2.xlsx

Tab S3. Robinson-Foulds distance matrix between the various phylogenetic inference methods.
The first row gives the alignment used for inference (L or H) and the second row the inference software used.

see file Tab_S3.xlsx

Tab S4. Ancestral state reconstruction using stochastic character mapping.

see file Tab_S4.xlsx

Tab S5. Scores and loadings from the phylogenetically corrected principal component analysis.

see file Tab_S5.xlsx

Tab S6. Results of PGLS analysis of states indicating reciprocal copulation versus hypodermic insemination on sperm length.
All predictors were binary, with the reference level being the state indicating hypodermic insemination.

see file Tab_S6.xlsx

Tab S7. Results from PGLS correlating the first principal components of a phylogenetically corrected principal component analysis (pPCA) analysis including five stylet traits with the first principal component of a pPCA analysis including four antrum traits.
Analysis was performed across all species and restricted to the reciprocal mating syndrome. Also given are pPCA loadings and results for all three phylogenies.

see file Tab_S7.xlsx

Tab S8. Details on the transcriptomes used in this study.

see file Tab_S8.xlsx

Tab S9. Mean species values for all morphological variables.

see file Tab_S9.xlsx