The spatial signal in area LIP is not an obligatory correlate of perceptual evidence during informed saccadic choices

Joshua A Seideman, Terrence R Stanford[1], and Emilio Salinas[1*]

Department of Neurobiology and Anatomy, Wake Forest School of Medicine, 1 Medical Center Blvd., Winston-Salem, NC 27157-1010, USA

[1]Co-senior authors

[*]Correspondence should be addressed to:

> Emilio Salinas
> Department of Neurobiology and Anatomy
> Wake Forest School of Medicine
> 1 Medical Center Blvd.
> Winston-Salem, NC 27157-1010, USA
> Phone: (336) 713-5176
> e-mail: esalinas@wakehealth.edu

Abbreviations: CRDM, compelled random dot motion; RDM, random dot motion; RF, response field; rPT, raw processing time; RT, reaction time; SD, standard deviation; SE, standard error

Running title: LIP differentiation dissociated from evidence accumulation

Grants: Research was supported by the NIH through grants R01EY025172, R01EY021228 and F31EY029154 from the NEI, and by the NIH-NINDS Training Grant T32NS073553.

Disclosures: The authors declare no conflicts of interest, financial or otherwise.

Author contributions: JAS, ES and TRS designed the research; JAS collected data; JAS, ES and TRS analyzed data; JAS, ES and TRS wrote the manuscript.

Keywords: attention, accumulation of evidence, decision making, eye movement, parietal cortex, oculomotor

draft February 16, 2021, 11:48

The lateral intraparietal area (LIP) contains spatially selective neurons that are partly responsible for determining where to look next, and are thought to serve a variety of sensory, motor planning, and cognitive control functions within this role[1,2,3]. Notably, according to numerous studies in monkeys[4,5,6,7,8,9,10,11,12], area LIP implements a fundamental perceptual process, the gradual accumulation of sensory evidence in favor of one choice (e.g., look left) over another (look right), which manifests as a slowly developing spatial signal during a motion discrimination task. However, according to recent inactivation experiments[13,14], this signal is unnecessary for accurate task performance. Here we reconcile these contradictory findings. We designed an urgent version of the motion discrimination task in which there is no systematic lag between the perceptual evaluation and the motor action reporting it, and such that the evolution of the subject's choice can be tracked millisecond by millisecond[15,16,17,18]. We found that while choice accuracy increased steeply with increasing sensory evidence, at the same time, the spatial signal became progressively weaker, as if it hindered performance. In contrast, in a similarly urgent task in which the discriminated stimuli and the choice targets were spatially coincident, the neural signal seemed to facilitate performance. The data suggest that the ramping activity in area LIP traditionally interpreted as evidence accumulation likely corresponds to a slow, post-decision shift of spatial attention from one location (where the motion occurs) to another (where the eyes land).

The lateral intraparietal area (**LIP**) combines sensory and cognitive information to highlight behaviorally relevant locations or visual features to look at. Although this may involve many sophisticated perceptual operations[3,19,20,21], the accumulation of sensory evidence (or, more generally, temporal integration) is one of major theoretical importance. First, by some accounts[22,23], it is an obligatory antecedent to perceptually guided choices regardless of task details, sensory modality, or effector. And second, its manifestation in LIP provides key experimental justification for sequential sampling models, which comprise the most widespread computational framework for reproducing reaction time (**RT**) and accuracy data in deterministic choice tasks[24,25,26,27]. In this framework, the gradual differentiation between spatial locations signaled by LIP corresponds directly to the gradual formation of the perceptual decision[28,29].

The random-dot motion (**RDM**) discrimination task (Fig. 1a) has been pivotal to this functional interpretation. In it, the subject must look at one of two choice targets to indicate the net direction of motion of a cloud of flickering dots, and in numerous variants of the task[4,5,6,7,8,9,10,11,12], LIP neurons gradually signal the chosen location while simultaneously reflecting the particulars of the perceptual discrimination. However, in recent inactivation experiments[13,14], the LIP spatial signal was disrupted with minimal consequence to performance (effects were seen on RT but not on accuracy), consistent with a more indirect relationship between LIP activity and decision formation[29,30].

We propose a simple yet potentially far-reaching explanation for this puzzling combination of findings: the perceptual evaluation of the motion stimulus occurs more rapidly (∼200 ms) than is generally assumed and may *precede* the LIP differentiation in many instances. So, what appears to be a gradual accumulation of sensory evidence is likely the byproduct of task designs that promote a slow, post-decision shift of attention from one spatial location (where the dots are) to another (where the chosen target is).

This hypothesis makes a stark prediction. Consider the RDM task performed with high urgency, such that perceptual and motor planning processes run concurrently (Fig. 1b, c). This will produce correct trials that are rapid (low RT) but still informed by the motion stimulus. If LIP neurons accumulate evidence, then in those trials they must still differentiate and indicate the impending choice, with stronger evidence yielding stronger differentiation. Alternatively, if the

Figure 1. Urgent and non-urgent motion discrimination tasks. Subjects had to report the direction of motion (left or right) of a cloud of flickering dots by looking at one of two peripheral targets. **a**, RDM task (non-urgent). The motion stimulus is presented and evaluated (Cue, 600–1000 ms) well before the go signal (fixation point offset; Go). **b**, CRDM task (urgent). The motion stimulus is presented (Cue) after the go signal (Go), with an unpredictable delay between them (Gap, 0–250) and a limited RT time window for responding (350–425 ms). The perceptual evaluation must occur during the cue-viewing interval (rPT = RT − gap), as the motor plan develops. **c**, Percentage of correct responses as a function of rPT, or tachometric curve. Results are from CRDM behavioral sessions for monkeys C (left) and T (right) for 100% (red; C: 9544, T: 33974 trials) and a lower coherence (black; C: 7909, T: 12066 trials). Shades indicate ± 1 SE from binomial statistics.

spatial differentiation in LIP occurs after the motion stimulus has been evaluated, its development on such rapid trials will be curtailed, and stronger evidence will not prevent its attenuation or abolition altogether.

Urgent versus non-urgent choices

To test this prediction, we recorded single-neuron activity in area LIP during two variants of the RDM discrimination task. In the standard, non-urgent version (Fig. 1a), the motion stimulus is presented first (Cue, 600–1000 ms) and is followed by the offset of the fixation point (Go), which means "respond now!" In the urgent or compelled random-dot motion (**CRDM**) discrimination task (Fig. 1b), the order of events is reversed; the go signal is given first, before the stimulus is shown, and the subject must respond within a short time window after the go (350–425 ms). Although the required perceptual judgment is the same, the tasks differ critically in the order in which perceptual and motor processes are engaged: in the former the saccade can be prepared with relative leisure, after the perceptual evaluation is completed, whereas in the latter the motor plan is initiated early and the perceptual evaluation must occur while the developing motor plan advances. Under time pressure, saccades can be triggered before, during, or shortly after the perceptual evaluation, and may result in guesses, partially informed, or fully informed choices (Fig. 1c). Perceptual and motor performance are effectively decoupled[15,16,17,18] (Fig. S1).

Our two monkey subjects performed the two choice tasks in interleaved blocks of trials (in addition to single-target tasks traditionally used to characterize LIP activity; Fig. 2a, b). In the standard, non-urgent RDM task, most choices were correct (93% and 84% correct for monkeys C and T at 100% coherence; Fig. S2), and the recorded LIP activity evolved as reported previously[4,5,8,11] (Fig. 2c). The neurons responded briskly upon presentation of a choice target in the response field (**RF**), continued firing at an elevated rate, and began signaling the choice about 200 ms after the onset of the motion stimulus (Fig. 2d, red arrow), at which point their activity increased for saccades into the RF and decreased for saccades away.

Figure 2. LIP activity in urgent versus non-urgent random-dot motion discrimination. **a**, Responses during visually guided saccades. Traces show normalized firing rate (mean $\pm$ 1 SE across cells; $n = 50$) as a function of time for correct trials into (magenta) or away from the cell's RF (green). Same scales for other panels. The gray bar indicates the go signal range for 90% of the trials. **b**, Responses during memory guided saccades ($n = 49$). **c**, Responses in the non-urgent RDM task ($n = 51$). **d**, Spatial signal magnitude as a function of time. S_{ROC} measures the statistical separation between the traces in **c** (same time axis; bin width = 50 ms). Red arrow marks approximate onset of differentiation. **e**, **f**, Responses in the CRDM task ($n = 51$) during guesses (**e**, rPT $\leq$ 150 ms) and fully informed choices (**f**, rPT $\geq$ 200 ms). **g**, Presaccadic S_{ROC} for individual neurons ($n = 51$) during guesses (x-axis) and fully informed choices in the CRDM task (right y-axis), and in the non-urgent RDM task (left y-axis). Spike counts for computing S_{ROC} are from shaded windows in **c**–**f**. Side plot shows bootstrapped distributions of mean values. **h**, Behavioral (black) and neuronal (brown) performance curves from the same CRDM sessions (mean $\pm$ 1 SE across trials). S_{ROC} is from presaccadic spikes pooled across neurons ($n = 51$) and sorted by rPT (bin width = 51 ms). All motion data are for 100% coherence.

To interpret this growing differential signal (with magnitude quantified by S_{ROC}, Fig. 2d) as an immediate correlate of the perceptual evaluation — one that is causal to the choice — one must assume that the evaluation begins about 200–250 ms after cue onset. And indeed, many experiments are consistent with such a protracted time scale[6,7,8,10,11,24]. However, none of these studies tracked the timecourse of performance explicitly, moment by moment. By doing this, we find that after 250 ms of stimulus viewing time the motion discrimination is essentially over.

Perceptual and neural discrimination under time pressure

In the CRDM task, the key variable is the amount of time during which the stimulus can be seen and analyzed before movement onset, which we call the raw processing time (**rPT**, computed as RT − gap in each trial; Fig. 1b). Plotting choice accuracy as a function of rPT yields a detailed, high-resolution account of the temporal evolution of the perceptual judgment (Figs. 1c, 2h). According to this 'tachometric curve,' in trials with rPT $\lesssim$ 140 ms the stimulus is seen so briefly that the motion direction cannot be resolved, which results in uninformed choices, or guesses (∼50%

correct). Choice accuracy then rises rapidly after the 150 ms mark, reaching asymptotic performance for rPTs of 200–250 ms. This amount of viewing time is sufficient for evaluating the RDM stimulus and reliably determining its motion direction.

As in other urgent tasks with similar designs[15,16,17,18], the rPT measured in each trial quantifies the degree to which sensory evidence guided the corresponding choice (or the probability that the choice was guided). Thus, if the differential signal in LIP reflects the amount of evidence accumulated in each trial, then it should be larger for fully informed discriminations (at long rPTs) than for guesses (at short rPTs), and its evolution should parallel the rise of the tachometric curve.

Contrary to this expectation, the recorded LIP activity showed quite the opposite. During performance of the CRDM task, the neural responses favoring each of the two possible eye movements were clearly separated just prior to saccade onset (Fig. 2e, f). Quantitatively, the presaccadic separation was less definitive than that in the non-urgent condition (Fig. 2g, red data), but the urgent differential signal still pointed reliably to the eventual choice. Crucially, however, across the sample of individual neurons recorded in the CRDM task ($n = 51$), the differential signal measured during fully informed, correct choices (rPT $\geq$ 200 ms) was considerably weaker than that during guesses (rPT $\leq$ 150 ms; Fig. 2g, blue data, $p = 0.001$, permutation test). More evidence yielded less differentiation. Furthermore, when the presaccadic responses were pooled across neurons and binned by rPT to calculate the magnitude of the spatial signal as a continuous function of rPT (Methods), the resulting neurometric curve *decreased* steadily for rPT $>$ 100 ms (Fig. 2h, brown curve) — in sharp contrast to choice accuracy (Fig. 2h, black curve). In the CRDM task, the stronger the influence of perception on the choice, the weaker the observed LIP differentiation.

Relationship between LIP differentiation and trial outcome

Everything else being equal, the neural encoding of perceptual information upon which choices are made is typically more robust for correct than for incorrect outcomes[31,32,33,34]. This is true across tasks, circuits, and modalities, and should apply to urgent choices too. During short-rPT trials (rPT $\leq$ 150 ms), the differential response in LIP was identical for correct and incorrect choices (Fig. 3a, gray bars), as anticipated given that those were all guesses. During informed discriminations (rPT $>$ 150 ms), however, the differentiation was greater for errors than for correct choices (Fig. 3a, b, blue vs. purple data, $p = 0.0006$, resampling test) — again, opposite to the trend expected from an evidence accumulation process.

In urgent tasks, the relationship between behavioral performance and single-neuron activity is revealed most effectively by conditioning the former on the latter. First, for a given experimental condition (saccade into or away from the RF), the responses collected from a neuron are sorted according to their magnitude (above vs. below the median), and then performance is compared across the corresponding groups of trials (Fig. S3; Methods). The resulting tachometric curves conditioned on evoked activity reveal if, when, and how the subject's behavior changes when the recorded neurons fire more or less than average. According to this analysis, performance was consistently worse ($p < 10^{-5}$, resampling test) in trials that were congruent with stronger spatial differentiation (Fig. 3c), as if a more robust spatial signal interfered with the urgent motion discrimination.

Spatial conflict within LIP

Why is the LIP differentiation suppressed in the CRDM task, and more so for informed choices? There are two likely reasons, both brought about by urgency. First, the differential signal is curtailed when it has less time to develop (Fig. 2g, red data), a general effect[18] consistent with

Figure 3. LIP differentiation may help or hinder performance. **a**, LIP activity in the CRDM task during guesses (rPT ≤ 150 ms, gray) and informed choices (rPT > 150 ms, blue, purple) sorted by outcome (x-axis). Activity indicates presaccadic spike counts normalized and pooled across neurons ($n = 51$). Data are mean and 95% CIs across trials. **b**, Differential signal magnitudes for the three conditions in **a** indicated by color. Curves are bootstrapped distributions. **c**, Performance in the CRDM task conditioned on neuronal activity. Trials were classified according to their presaccadic spike counts as either congruent (red) or incongruent (black) with strong differentiation. Inset shows bootstrapped distribution for the mean difference in percent correct between curves for rPTs of 140–280 ms. **d–f**, As in **a–c**, but for the urgent color discrimination task ($n = 56$; in **d**, **e**, rPT ≤ 125 ms for guesses and rPT > 125 ms for informed choices).

our initial hypothesis. And second, given LIP's participation in attentional deployment[2,35,36], the particular geometry of the task must create a spatial conflict: the early motor plan, initiated shortly after the go signal[15,18], automatically allocates attentional resources to the planned saccade endpoint(s)[37,38,39,40], but attention should be directed to the RDM stimulus, which defines the perceptually relevant location[13,14]. A spatial competition ensues[35,41]. Evidence of this is plainly manifest in the behavioral CRDM data (Fig. S4).

To investigate the contributions of these two factors, limited time and attentional conflict, we recorded LIP activity from the same monkeys during two versions, urgent and non-urgent, of a discrimination task in which the subject must make an eye movement to the peripheral stimulus that matches the color of the fixation point[42] (Fig. S5). The key difference here is that the conflict described above is eliminated: the relevant color cues are found at the choice targets, and deploying attention/perceptual resources to them should be of benefit, if not a necessity, to the required discrimination (Fig. S6).

During the non-urgent color-matching task, the sampled neurons (which again exhibited characteristic LIP response features; Fig. 4a, b) differentiated saccades into versus away from the RF (Fig. 4c, d) slightly earlier than during the standard RDM task (Figs. 2d, 4d, arrows). But, overall, under relaxed, non-urgent conditions, the evoked spatial signal developed with comparable timecourse and strength in the motion- and color-based tasks, in spite of their distinct spatial and feature requirements. Under time pressure, though, the comparison across tasks was striking. During the urgent color-matching task, the differential response in LIP was larger for informed than uninformed discriminations (Fig. 4e–g); its magnitude increased over time in parallel with the monkeys' choice accuracy (Fig. 4h); it was weaker for errors than correct choices during informed trials (Fig. 3d, e); and it acted as if to improve the monkeys' performance (Fig. 3f). In this case, the greater the influence of perception on the choice, the stronger the spatial signal observed in LIP.

Figure 4. LIP activity in urgent versus non-urgent color discrimination. Same format and conventions as in Fig. 2. Data are from $n = 56$ sampled neurons, except in **c**, **d**, and **g** (red data), for which $n = 43$. In **e**, rPT ≤ 125 ms. In **f**, rPT ≥ 175 ms.

These results in the color-matching task experiment confirm that an informed spatial signal can emerge very rapidly in LIP[43,44]. They show that time pressure alone does not necessarily abolish or reverse the expected correlation between evidence and LIP differentiation, and so cannot explain the CRDM results. Rather, the data suggest that the anti-correlation between CRDM performance and LIP spatial signal strength results from urgency together with spatial conflict between the perceptually relevant location and the saccade endpoint. In this case, spatial selection of the saccade target corresponds to attention being diverted away from the location of the dots during a brief but critical period of time when the motion stimulus is being evaluated.

In both the motion- and color-based tasks the results were robust with respect to the subjects' performance level (Fig. S7), the criteria used for including/excluding neurons (Figs. S8, S9, S10), and how the effects were quantified (Fig. S11).

Conclusions

The highly robust target selection seen during non-urgent conditions (RDM task) would lead one to conclude, as have countless past studies, that LIP differentiation is an obligatory, causal antecedent to perceptually informed choices, and that greater differentiation implies more or stronger perceptual evidence. Yet, for equally informed choices made urgently (CRDM task), the spatial signal was markedly diminished, and it decreased with increasing evidence. While counterintuitive, this result remains consistent with a representation of spatial priority[2,35,41].

In general, the differential activation of oculomotor neurons simply points to locations that are important; the saccade endpoint, the place where sensory evidence is found, or locations that contain reward information, a symbolic instruction, or a salient distracter[2,35,36,45,46,47]. Such versatility is the hallmark of attention-related activity. During motion discrimination, the location

that is relevant to the perceptual evaluation is where the dots are, and indeed, inactivation exper-
iments indicate that the LIP neurons with RFs covering the dots are precisely the ones that matter
the most for performance[13,14]. Under urgent conditions (CRDM task), stronger differentiation be-
tween the two perceptually irrelevant choice-target locations would denote a firmer attentional
commitment, and likely a stronger conflict with the RFs covering the dots. However, when the
urgency requirement is relaxed (standard RDM task), attention can be deployed to the location
of the dots even before motion onset, and can remain there as long as necessary to then shift to
the chosen saccade target. The focus on the dots need not be long (∼100 ms, considering the time
to transition from chance to asymptotic performance; Figs. 1c, 2h), and the timecourse and mag-
nitude of the shift may still depend on the strength of the sensory evidence. If so, the resulting
post-perceptual differentiation may appear causal to the choice.

Methods

Subjects and setup

All experimental procedures were conducted in accordance with NIH guidelines, USDA regula-
tions, and the policies set forth by the Institutional Animal Care and Use Committee (IACUC) of
Wake Forest School of Medicine. The subjects in this experiment were two adult male rhesus mon-
keys (Macaca mulatta) weighing between 8.5 and 11 kg. For each animal, an MRI-compatible post
(Crist Instruments, MD, USA) was implanted on the skull while under general anesthesia. The
post served to fix the position of the head during all experimental sessions. Following head-post
implantation, both subjects were trained to perform oculomotor response tasks in exchange for
water reward. After reaching a criterion level (> 75% accuracy for each task), craniotomies were
made and recording cylinders (Crist Instruments, MD, USA) were placed over the left LIP of each
monkey (monkey C: left hemisphere; monkey T: left and right hemispheres; stereotactic coordi-
nates: 5 posterior, 12 lateral[48,49]) while under general anesthesia. Neural recordings commenced
after a 1-2 week recovery period following cylinder placement.

Behavioral and neurophysiological recording systems

Eye position was monitored using an EyeLink 1000 Plus infrared tracking system (SR Research;
Ottawa, Canada) at a sampling rate of 500 or 1000 Hz. For sessions in which dot-motion tasks
were performed, all gaze-contingent stimulus presentation and reward delivery were controlled
using Psychtoolbox[50,51] version 2.0; for all other sessions, gaze-contingent stimulus presentation
and reward delivery were controlled via a custom-designed PC-based software package (Ryklin
Software). Visual stimuli were presented on a Viewpixx/3D display (Vpixx Technologies, Quebec,
Canada; 1920 × 1080 screen resolution, 120 Hz refresh rate, 12 bit color) placed 57 cm away from
the subject. Viewing was binocular. Neural activity was recorded using single tungsten micro-
electrodes (FHC, Bowdoin, ME; 2–4 MΩ impedance at 1 kHz) driven by a hydraulic microdrive
(FHC). A Cereplex M headstage (Blackrock Microsystems, Utah, USA) filtered (0.03 Hz – 7.5 kHz),
amplified, and digitized electrical signals, which were then sent to a Cereplex Direct (Blackrock
Microsystems) data acquisition system. Single neurons were isolated online based on amplitude
criteria and/or waveform characteristics.

Behavioral tasks

218 Three design elements are the same for all the tasks. (1) Each trial begins with presentation of a
219 central spot and the monkey fixating it for 300–800 ms. (2) The offset of the fixation spot is the
220 go signal that instructs the monkey to make a saccade. (3) To yield a reward (drop of liquid), the
221 saccade must be to the correct location and must be initiated within an allotted RT window. The

RT is always measured as the time elapsed between fixation offset and saccade onset (equal to the time point following the go signal at which the eye velocity first exceeds a criterion of 25 deg/s). In non-urgent tasks the monkey is allowed to initiate an eye movement within 600 ms of the go signal, whereas in urgent tasks this must happen within 350–425 ms.

Visually- and memory-guided saccade tasks: Two standard single-target tasks were used to characterize the visuomotor properties of LIP neurons. In both tasks, after the monkey fixates, a peripheral target is presented (Target on) either within or diametrically opposed to the RF of the recorded neuron. For the delayed visually-guided saccade task, after a variable delay (500–1000 ms), the fixation spot disappears (Go) and the monkey is required to make a saccade to the peripheral target. For the memory-guided saccade task, after being displayed for 250 ms, the peripheral target is extinguished (Target off) and the monkey is required to maintain fixation throughout a subsequent delay interval (500–1000 ms). After this memory interval, the fixation spot disappears (Go) and the monkey is required to make a saccade to the remembered target location.

Non-urgent RDM motion discrimination task: This two-alternative task (Fig. 1a) is similar to previous implementations of the RDM discrimination task[4,5,8,11]. Upon fixation and after a short delay (300–500 ms), two gray stimuli, the potential targets, are presented (Targets on), one in the RF and one diametrically opposed. After a delay (250–750 ms), a cloud of randomly moving dots appears in the center of the screen or just above the fixation point for 600–1000 ms (Cue). Then, after another delay period (300–500 ms; Delay), the fixation spot is extinguished (Go), which instructs the monkey to make a choice. If the saccade is to the stimulus in the direction of the dot motion (and is made within 600 ms), the monkey obtains a liquid reward. The direction of motion, toward one choice target or the other, is assigned randomly from trial to trial. The difficulty of the task varies with stimulus coherence, which is the percentage of dots that move in a consistent direction across video frames. Monkeys worked with coherence values of 100%, 50%, 25%, 6% and 3%, but the neural data were recorded at 100% (Fig. S2).

Compelled random-dot motion discrimination task: The CRDM task (Fig. 1b) is an urgent version of the RDM discrimination task just described. The geometry, reward size, and stimuli are the same; only the temporal requirements are different. In this case, the monkey fixates, the two peripheral gray stimuli are shown (Targets on), and after a delay (250–750 ms), the go signal is given (Go), urging the subject to respond as quickly as possible (within 350–425 ms). At this point in the trial, however, no information is available yet to guide the choice. That information, conveyed by the cloud of flickering dots, is revealed later (Cue), after an unpredictable amount of time following the go (Gap; 0–250 ms). Subjects are tasked with looking to the peripheral choice alternative that is congruent with the net direction of motion of the dots (Saccade).

On each trial, the raw processing time, or rPT, is the maximum amount of time that is potentially available for seeing and evaluating the motion stimulus. It is the time interval between cue onset and saccade onset (rPT = RT − gap). We refer to it as 'raw' because it includes any afferent or efferent delays in the circuitry[15]. Gap values (0–250 ms) varied randomly from trial to trial and were chosen to yield rPTs covering the full range between guesses and informed choices.

Non-urgent color discrimination task: In this task (Fig. S5a), the color of the central fixation spot (red or green) defines the identity of the eventual target. Upon fixation and after a short delay (300–800 ms), two gray stimuli, the potential targets, are presented (Targets on), one in the RF and one diametrically opposed. After a delay (250–750 ms), one of the gray stimuli changes to red and the other to green (Cue). After a cue viewing period (500–1000 ms), the fixation spot is extinguished (Go), which instructs the monkey to make a choice. If the ensuing saccade is to the stimulus that matches the color of the prior fixation spot and is made within 600 ms, the monkey obtains a reward. Colors and locations for target and distracter are randomly assigned in each trial.

Urgent color discrimination task: This task (Fig. S5b), also referred to as the compelled-saccade task[15,16,18,42], requires the same red-green discrimination as in the easier non-urgent version. In this case, after the monkey fixates (300–800 ms) and the two gray stimuli in the periphery are displayed (Targets on; 250–750 ms), the fixation spot disappears (Go). This instructs the monkey to make a choice, although the visual cue that informs the choice (one gray spot turning red and the other green; Cue) is revealed later, after an unpredictable period of time following the go signal (Gap; 0–250 ms). To obtain a reward, the monkey must look to the peripheral stimulus that matches the color of the initial fixation spot (Saccade) within the allowed RT window (350–425 ms). As with the CRDM task, the key variable that determines performance is the rPT.

Tachometric curves and rPT intervals

All data analyses were performed in Matlab (The MathWorks, Natick MA). To compute the tachometric curve and rPT distributions, trials were grouped into rPT bins of 51 ms, with bins shifting every 1 ms. Numbers of correct and incorrect trials were then counted within each bin.

When parsing trials into short and long rPT time bins (Figs. 2e–g, 3a, b, d, e, 4e–g), we considered the distributions of processing times from all the recording sessions in each task. The threshold for guesses (rPT $\leq$ 150 for the CRDM task; rPT $\leq$ 125 ms for the color task) corresponded to the point at which the fractions of correct and incorrect trials started diverging steadily with rPT. Trials above this cutoff were considered informed, and trials above this cutoff plus 50 ms, which brought the fraction correct about 75% of the way from chance to asymptotic, were considered fully informed. The results depended minimally on the exact cutoffs used.

Tachometric curves conditioned on neuronal activity (Fig. 3c, f) were computed as follows. First, for each neuron, spike counts from a presaccadic window ($-50:0$ ms, aligned on saccade) were collected and sorted into two conditions, saccade-in (S_{in}) and saccade-out (S_{out}) choices. The trials in each condition were then split into two groups, with spike counts below the median for the condition, or with spike counts at or above it. Four groups of trials resulted: S_{in} high firing, S_{in} low firing, S_{out} high firing, and S_{out} low firing. Data from all the neurons in a sample were aggregated, and a tachometric curve was generated for each group (Fig. S3). The first and last groups are congruent with a strong spatial signal, whereas the other two are incongruent. Because the results were consistent for S_{in} and S_{out} conditions (Fig. S3), congruent and incongruent trials were combined across conditions.

Characterization of neural activity

On line, RF location was determined from activity levels measured around the time of saccade onset during performance of the visually- or memory-guided saccade task. All neurons included in the current study ($n =$51 for CRDM task, $n =$56 for urgent color discrimination task) were significantly activated during performance of the urgent tasks, both in response to visual stimuli presented in their RF (window: 20:150 ms, aligned on targets on) as well as prior to saccades executed into their RF (window: $-100:0$ ms, aligned on saccade) relative to respective baseline measures. In addition, all neurons included exhibited significant delay period activity in the visually- and/or memory-guided saccade tasks. For all such determinations, significance ($p < 0.01$) was calculated numerically via permutation tests[52] in which the two group labels (e.g., 'baseline' and 'response period') were randomly permuted. These physiological response properties (i.e., visual, delay period, and presaccadic activation) are characteristic of LIP neurons that project directly to saccade production centers[53] (i.e., the superior colliculus).

Some additional neurons that were recorded and fully characterized (15 in the CRDM experiment, 26 in the color-based) were excluded from the studied samples for any of the following

reasons: they had no significant visual or memory activity in the single-target tasks; they were
not significantly activated presaccadically; or their spatial preference for contralateral/ipsilateral
stimuli either was ambiguous or clearly flipped between different tasks. Importantly, though, ex-
cept for small quantitative variations, all results were essentially identical with inclusion of all
such neurons (Fig. S8).

For each neuron, continuous firing rate traces, or spike density functions, were generated by
aligning the recorded spike trains to relevant task events (e.g., cue onset, saccade onset), convolv-
ing them with a gaussian kernel ($\sigma =$15 ms), and averaging across trials. Normalized population
traces (as in panels a–c, e, f of Figs. 2, 4) were generated by dividing each cell's response curve by
its maximum firing rate value and then averaging across cells. For each cell, this maximum rate
was calculated from the recorded urgent trials (motion- or color-based) and was used to normalize
the population traces for all other tasks.

ROC analyses and neurometric curves

The magnitude of spatial differentiation, or S_{ROC}, was used to quantify the degree to which LIP
neurons were differentially activated in S_{in} versus S_{out} choices. This measure corresponds to
the accuracy with which an ideal observer can classify data samples from two distributions (of
responses in S_{in} and S_{out} trials, in this case), and is equivalent to the area under the receiver
operating characteristic, or ROC, curve[54,55]. Values of 0.5 correspond to distributions that are in-
distinguishable (chance performance, full overlap), whereas values of 0 or 1 correspond to fully
distinguishable distributions (perfect performance, no overlap). Here, $S_{ROC} > 0.5$ always indi-
cates higher activity for saccades into the RF than away from the RF. Presaccadic S_{ROC} values
(Figs. 2g, h, 3b, e, 4g, h) were computed using spike counts measured prior to choice onset (-50:0
ms, relative to saccade onset) and sorted according to trial outcome.

For the urgent tasks, continuous neurometric functions comparable to the behavioral tacho-
metric curves (Figs. 2h, 4h) were generated by first pooling the data across neurons and then cal-
culating S_{ROC} as a function of rPT (bin width $=$ 51 ms, shifted every 1 ms). The pooling involved
two steps. First, the presaccadic spike counts of each neuron were normalized by subtracting a
constant, θ, that was cell-specific, and then the normalized spike counts from all the neurons were
sorted into two groups, for S_{in} and S_{out} trials. The pooled S_{ROC} compared responses from these
two pooled distributions within each rPT bin (see Fig. S12 for an example). For each neuron, the
constant θ was equal to $(m_{in} + m_{out})/2$, where m_{in} and m_{out} are the mean spike counts for S_{in} and
S_{out} trials. Other normalization schemes produced qualitatively similar trends. This procedure,
pooling the data first and then computing S_{ROC}, generated more precise results than the reverse,
i.e., first computing S_{ROC} for each cell and then averaging across cells. However, the latter al-
ternative produced qualitatively consistent results (Fig. S11). We stress that, although the pooled
S_{ROC} values that make up the neurometric curve vary with rPT, they were always based on spike
counts measured just prior to the saccade.

For the non-urgent tasks (Figs. 2d, 4d), continuous S_{ROC} values were again computed by di-
viding time into sliding bins (bin width $=$ 50 ms, shifted every 1 ms). For each bin, the spikes
counted for each neuron in each condition (S_{in} and S_{out} trials) were used to calculate that cell's
S_{ROC}, and then values were averaged across cells. Pooling was unnecessary in this case because
more data were available in each time bin, but the results with pooling were very similar.

Effect sizes for mean S_{ROC} values were computed by bootstrapping[56,57]; that is, by repeatedly
resampling the underlying data with replacement ($10^4 - 10^5$ iterations) and recomputing the mean
S_{ROC} each time. In Figs. 2g, 4g (insets), the resampling was over neurons; in Fig. 3b, e, it was over
trials in the two pooled distributions (for S_{in} and S_{out} conditions). Effect sizes for other quantities

(e.g., Δc in Fig. 3c, f) were also calculated through bootstrapping. Having generated these effect-size distributions for any two conditions (e.g., correct vs. incorrect choices, or long vs. short rPTs), we could calculate from them a significance value for the mean difference. Instead, however, for any relevant comparison between two conditions, the p-value of the difference was calculated separately using a permutation test[52] for paired data or an equivalent resampling test for non-paired data, as these tests provide slightly more accurate and specific comparisons against the null hypothesis (of no difference between the distributions from which the two data sets originated). For example, to compare the mean S_{ROC} for short- versus long-rPT trials (Figs. 2g, 4g, insets), we randomly permuted the 'short' and 'long' labels for each neuron and recomputed the difference between S_{ROC} means 10^5 times. All reported significance values were calculated similarly, via permutation or resampling tests (one-sided).

References

1. Snyder, L.H., Batista, A.P., and Andersen, R.A. (2000). Intention-related activity in the posterior parietal cortex: a review. Vision Res. 40, 1433–1442.

2. Bisley JW, Goldberg ME (2010) Attention, intention, and priority in the parietal lobe. Annu Rev Neurosci 33:1–21.

3. Freedman DJ, Ibos G (2018) An integrative framework for sensory, motor, and cognitive functions of the posterior parietal cortex. Neuron 97:1219–1234.

4. Shadlen MN, Newsome WT (2001) Neural basis of a perceptual decision in the parietal cortex (area LIP) of the rhesus monkey. J Neurophysiol 86:1916–1936.

5. Roitman JD, Shadlen MN (2002) Response of neurons in the lateral intraparietal area during a combined visual discrimination reaction time task. J Neurosci 22: 9475–9489.

6. Huk AC, Shadlen MN (2005) Neural activity in macaque parietal cortex reflects temporal integration of visual motion signals during perceptual decision making. J Neurosci 25:10420–1036.

7. Hanks TD, Ditterich J, Shadlen MN (2006) Microstimulation of macaque area LIP affects decision-making in a motion discrimination task. Nat Neurosci 9:682–689.

8. Kiani R, Hanks TD, Shadlen MN (2008) Bounded integration in parietal cortex underlies decisions even when viewing duration is dictated by the environment. J Neurosci 28:3017–3029.

9. Rorie AE, Gao J, McClelland JL, Newsome WT (2010) Integration of sensory and reward information during perceptual decision-making in lateral intraparietal cortex (LIP) of the macaque monkey. PLoS One 5:e9308.

10. Churchland AK, Kiani R, Chaudhuri R, Wang XJ, Pouget A, Shadlen MN (2011) Variance as a signature of neural computations during decision making. Neuron 69:818–831.

11. de Lafuente V, Jazayeri M, Shadlen MN (2015) Representation of accumulating evidence for a decision in two parietal areas. J Neurosci 35:4306–4318.

12. Kira S, Yang T, Shadlen MN (2015) A neural implementation of Wald's sequential probability ratio test. Neuron 85:861–873.

13. Katz LN, Yates JL, Pillow JW, Huk AC (2016) Dissociated functional significance of decision-related activity in the primate dorsal stream. Nature 535:285–288.

14. Zhou Y, Freedman DJ (2019) Posterior parietal cortex plays a causal role in perceptual and

categorical decisions. Science 365:180–185.

15. Stanford TR, Shankar S, Massoglia DP, Costello MG, Salinas E (2010) Perceptual decision
making in less than 30 milliseconds. Nat Neurosci 13:379–385.

16. Costello MG, Zhu D, Salinas E, Stanford TR (2013) Perceptual modulation of motor — but
not visual — responses in the frontal eye field during an urgent-decision task. J Neurosci
33:16394–16408.

17. Salinas E, Steinberg BR, Sussman LA, Fry SM, Hauser CK, Anderson DD, Stanford TR (2019)
Voluntary and involuntary contributions to perceptually guided saccadic choices resolved
with millisecond precision. eLife 8:e46359.

18. Scerra VE, Costello MG, Salinas E, Stanford TR (2019) All-or-none context dependence de-
lineates limits of FEF visual target selection. Curr Biol 29:294–305.

19. Gottlieb J (2007) From thought to action: the parietal cortex as a bridge between perception,
action, and cognition. Neuron 53:9–16.

20. Bennur S, Gold JI (2011) Distinct representations of a perceptual decision and the associated
oculomotor plan in the monkey lateral intraparietal area. J Neurosci 31:913–921.

21. Freedman DJ, Assad JA (2016) Neuronal mechanisms of visual categorization: an abstract
view on decision making. Annu Rev Neurosci 39:129–147.

22. O'Connell RG, Dockree PM, Kelly SP (2012) A supramodal accumulation-to-bound signal
that determines perceptual decisions in humans. Nat Neurosci 15:1729–1735.

23. Twomey DM, Kelly SP, O'Connell RG (2016) Abstract and effector-selective decision sig-
nals exhibit qualitatively distinct dynamics before delayed perceptual reports. J Neurosci
36:7346–7352.

24. Palmer J, Huk AC, Shadlen MN (2005) The effect of stimulus strength on the speed and
accuracy of a perceptual decision. J Vis 5:376–404.

25. Bogacz R, Brown E, Moehlis J, Holmes P, Cohen JD (2006) The physics of optimal decision
making: a formal analysis of models of performance in two-alternative forced-choice tasks.
Psychol Rev 113:700–765.

26. Ratcliff R, McKoon G (2008) The diffusion decision model: theory and data for two-choice
decision tasks. Neural Comput 20:873–922.

27. Forstmann BU, Ratcliff R, Wagenmakers EJ (2016) Sequential sampling models in cognitive
neuroscience: advantages, applications, and extensions. Annu Rev Psychol 67:641–666.

28. Gold JI, Shadlen MN (2007) The neural basis of decision making. Annu Rev Neurosci 30:535–
574.

29. Huk AC, Katz LN, Yates JL (2017) The role of the lateral intraparietal area in (the study of)
decision making. Annu Rev Neurosci 40:349–372.

30. Yates JL, Park IM, Katz LN, Pillow JW, Huk AC (2017) Functional dissection of signal and
noise in MT and LIP during decision-making. Nat Neurosci 20:1285–1292.

31. Salinas E, Hernández H, Zainos A, Romo R (2000) Periodicity and firing rate as candidate
neural codes for the frequency of vibrotactile stimuli. J Neurosci 20:5503–5515.

32. Cook EP, Maunsell JHR (2002) Dynamics of neuronal responses in macaque MT and VIP
during motion detection. Nat Neurosci 10: 985–994.

33. Matsumora T, Koida K, Komatsu H (2008) Relationship between color discrimination and

neural responses in the inferior temporal cortex of the monkey. J Neurophysiol 100:3361–
3374.

34. Nienborg H, Cohen MR, Cumming BG (2012) Decision-related activity in sensory neurons: correlations among neurons and with behavior. Annu Rev Neurosci 35:463–483.

35. Bisley JW, Goldberg ME (2003) Neuronal activity in the lateral intraparietal area and spatial attention. Science 299: 81–86.

36. Herrington TM, Assad JA (2010) Temporal sequence of attentional modulation in the lateral intraparietal area and middle temporal area during rapid covert shifts of attention. J Neurosci 30:3287–3296.

37. Kowler E, Anderson E, Dosher B, Blaser E (1995) The role of attention in the programming of saccades. Vision Res 35:1897–1916.

38. Deubel H, Schneider WX (1996) Saccade target selection and object recognition: evidence for a common attentional mechanism. Vision Res 36:1827–1837.

39. Moore T, Fallah M (2004) Microstimulation of the frontal eye field and its effects on covert spatial attention. J Neurophysiol 91: 152–162.

40. Steinmetz NA, Moore T (2014) Eye movement preparation modulates neuronal responses in area V4 when dissociated from attentional demands. Neuron 83:496–506.

41. Falkner AL, Krishna BS, Goldberg ME (2010) Surround suppression sharpens the priority map in the lateral intraparietal area. J Neurosci 30:12787–12797.

42. Seideman JA, Salinas E, Stanford TS (2019) Perceptual modulation of parietal activity during urgent saccadic choices. bioRxiv 2019.12.12.874313 [Preprint].

43. Ipata AE, Gee AL, Goldberg ME, Bisley JW (2006) Activity in the lateral intraparietal area predicts the goal and latency of saccades in a free-viewing visual search task. J Neurosci 26:3656–3661.

44. Sapountzis P, Paneri S, Gregoriou GG (2018) Distinct roles of prefrontal and parietal areas in the encoding of attentional priority. Proc Natl Acad Sci USA 115:E8755–E8764.

45. Ignashchenkova A, Dicke PW, Haarmeier T, Thier P (2004) Neuron-specific contribution of the superior colliculus to overt and covert shifts of attention. Nat Neurosci 7:56–64.

46. Peck CJ, Jangraw DC, Suzuki M, Efem R, Gottlieb J (2009) Reward modulates attention independently of action value in posterior parietal cortex. J Neurosci 29:11182–11191.

47. Zhou HH, Thompson KG (2009) Cognitively directed spatial selection in the frontal eye field in anticipation of visual stimuli to be discriminated. Vision Res 49:1205–1215.

48. Colby CL, Duhamel JR, Goldberg ME (1996) Visual, presaccadic, and cognitive activation of single neurons in monkey lateral intraparietal area. J Neurophysiol 76:2841–2852.

49. Snyder LH, Batista AP, Andersen RA (1998) Change in motor plan, without a change in the spatial locus of attention, modulates activity in posterior parietal cortex. J Neurophysiol 79:2814–2819.

50. Brainard DH (1997). The psychophysics toolbox. Spatial Vision 10:433–436.

51. Kleiner M, Brainard D, Pelli D (2007) Whats new in Psychtoolbox-3? Perception 36 (ECVP Abstract Supplement).

52. Siegel S, Castellan NJ (1988) Nonparametric statistics for the behavioral sciences. Boston, MA: McGraw-Hill.

53. Paré M, Wurtz RH (1997) Monkey posterior parietal cortex neurons antidromically activated from superior colliculus. J Neurophysiol 78:3493–3497.

54. Green DM, Swets JA (1966) Signal Detection Theory and Psychophysics. New York: Wiley.

55. Fawcett T (2006) An introduction to ROC analysis. Pattern Recognit Lett 27:861–874. doi:10.1016/j.patrec.2005.10.010.

56. Efron B (1982) The jacknife, the bootstrap and other resampling plans. (Society for Industrial and Applied Mathematics: Philadelphia).

57. Hesterberg T (2014) What teachers should know about the bootstrap: resampling in the undergraduate statistics curriculum. arXiv: 1411.5279.

516
Figure S3. Behavioral performance in the urgent tasks conditioned on LIP neuronal activity. **a**,
Tachometric curves from all the CRDM trials in which the outcome was a saccade into the recorded
cell's RF (S_{in}). For each neuron ($n = 51$), S_{in} trials were split according to whether the response
was at or above the median (brown curve, high firing), or below the median (gray curve, low
firing). The response was the spike count elicited in the 50 ms immediately preceding the onset
of the saccade. High firing in S_{in} trials is congruent with a strong spatial signal, whereas low
firing is incongruent. **b**, Tachometric curves from all the CRDM trials in which the outcome was
a saccade away from the recorded cell's RF (S_{out}). For each neuron, S_{out} trials were split accord-
ing to whether the response was at or above the median (blue curve, high firing), or below the
median (purple curve, low firing). Low firing in S_{out} trials is congruent with a strong spatial sig-
nal, whereas high firing is incongruent. **c**, Results combining all congruent and incongruent trials
across conditions. **d–f**, As in **a–c**, but for the urgent color discrimination task ($n = 56$). Significance
values shown are from resampling tests on the mean difference between each pair of curves for
$140 \leq$ rPT ≤ 280 ms. N indicates number of trials. Data in **c**, **f** are the same as in Fig. 3c, f. Note
that the results are consistent between S_{in} and S_{out} conditions.

Figure S4. Evidence of spatial conflict in the CRDM task. a, Percentage of correct choices as a function of raw processing time. The data comprise all trials from the behavioral sessions of monkey C (left column) and monkey T (right column), and include all coherences. Trials ($n = 30473$ for monkey C, $n = 52945$ for monkey T) were sorted into 50 ms bins (red curve, bins shifted every 1 ms), as done for tachometric curves shown in other figures, or into 1 ms bins (black dots). b, Processing time distributions for correct (cyan) and incorrect (blue) choices from the same trials in a, sorted into 1 ms bins. c, Enlarged view of the data in b between 40 and 140 ms of rPT. Note the prominent dip in the number of events at 80–100 ms. We interpret the lack of saccades during this narrow interval as an interruption of the ongoing motor plans due to the onset of the moving-dot stimulus. This is entirely consistent both with the capture of exogenous attention by a salient stimulus[17,35,101] and with the related phenomenology of saccadic inhibition[102,103,104]. The timing of this dip (∼90 ms after cue onset) is also consistent with a slight decrease in LIP activity often observed[4,5,8,11] in the non-urgent RDM task (Fig. S6a). For a brief moment, the cue-driven activation at the location of the random dots is in intense conflict with the oculomotor activity that generates saccades to the choice targets.

548
Figure S5. Urgent and non-urgent color-discrimination tasks. Subjects had to look to the periph-
eral target that matched the color of the fixation point. **a**, Easy, non-urgent task. The color stimuli
are presented (Cue, 500–1000 ms) and evaluated well before the go signal (fixation point offset;
Go). **b**, Urgent task. The color stimuli are presented (Cue) after the go signal (Go), with an unpre-
dictable delay between them (Gap, 0–250) and a limited RT time window for responding (350–425
ms). The perceptual evaluation must occur during the cue-viewing interval (rPT = RT − gap), as
the motor plan develops. **c**, Percentage of correct responses as a function of rPT, or tachometric
curve. Results are from all the recording sessions during which the urgent color-discrimination
task was performed by monkeys C (left, $n = 7330$ trials) and T (right, $n = 10745$ trials). Shades
indicate ± 1 SE from binomial statistics.

559
Figure S6. Cue-driven LIP responses in the non-urgent motion- and color-based discrimination
tasks are indicative of the deployment of spatial attention. **a**, Neuronal activity into (magenta)
and away from the RF (green) in the non-urgent RDM task ($n = 51$). Same data as in Fig. 2c, but
restricted to the period following cue onset. **b**, Spatial signal magnitude, S_{ROC}, as a function of
time, for the data in **c**. Arrow marks approximate onset of differentiation. Same data as in Fig. 2d,
but over a restricted time period. **c**, **d**, As in **a**, **b**, but for the non-urgent color discrimination task
($n = 43$). Same data as in Fig. 4c, d, but over a restricted time period. Note the change in activity
in the 50–150 ms following cue onset (gray, shaded areas): after the motion stimulus appears near
fixation (panel **a**), the LIP activity associated with the choice-target locations decreases slightly, as
found in previous studies[4,5,8,11]; in contrast, after the color change of the choice targets (panel **c**),
activity clearly increases. The decrease in **a** is maximal ∼90 ms after cue onset, at the same time
that (uninformed) saccades to the choice targets are completely suppressed in the urgent CRDM
task (Fig. S4). Interpreting the LIP activity as an attention signal, the motion stimulus near fixation
acts as if to suppress attention at the choice targets, whereas the color change at those targets acts
as if to increase the intensity of the existing signal.

 Figure S7. Modulation of LIP activity during high- versus low-performance sessions. For both the
motion- and color-based urgent tasks, recording sessions were ranked according to overall percent
correct. The 20 sessions with the best performance (gray traces) and the 20 sessions with the poor-
est performance (brown traces) were selected, and analyses were run for each group separately.
a, Tachometric curves, i.e., percent correct as a function of processing time, for the best and worst
CRDM sessions. **b**, Neurometric curves in the CRDM task, i.e., magnitude of presaccadic differ-
entiation as a function of processing time (as in Fig. 2h). In both groups of sessions, stronger LIP
differentiation was associated with less evidence (shorter rPTs). **c**, Tachometric curves conditioned
on neuronal activity in the CRDM task (as in Fig. 3c). In both groups of sessions, stronger LIP dif-
ferentiation (congruent condition) was associated with worse perceptual performance. **d–f**, Same
as **a–c**, but for the urgent color discrimination task (as in Figs. 4h and 3f). In this task, stronger
LIP differentiation was associated with more evidence (longer rPTs; **e**) and improved perceptual
performance (**f**). In each task, the relationship between neuronal activity in LIP and behavior was
similar for the two groups of sessions, in spite of the dramatically different performance levels.

590
Figure S8. Results with extended neuronal populations. The results in the main figures were
based on neurons ($n = 51$ for the CRDM task, $n = 56$ for the urgent color discrimination task) that
were fully characterized and satisfied several criteria: they had both visually-driven and saccade-
related responses, and their RFs were well defined and consistent across tasks (Methods). This
resulted in the exclusion of 15 additional neurons recorded in the CRDM experiment and 26 in
the color-based. Here we show the results of analyses that included all the neurons recorded in
the CRDM (left column, $n = 66$) and the urgent color discrimination task (right column, $n = 82$),
with no exclusions. **a**, Mean, normalized neuronal activity pooled across neurons and sorted by
experimental condition (as in Fig. 3a, d). **b**, Distributions for mean presaccadic differentiation
values (S_{ROC}), obtained by bootstrapping, for each of the three conditions above, as indicated by
the corresponding colors (as in Fig. 3b, e). **c**, Neurometric curves, i.e., magnitude of presaccadic
differentiation as a function of processing time (as in Figs. 2h, 4h). **d**, Behavioral performance
conditioned on neuronal activity (as in Fig. 3c, e). Inclusion of the additional populations did not
alter the results substantially.

Figure S11. Alternate procedure for combining the data across neurons. In the main figures (Figs. 2h, 4h, 3a, d), we first pooled all the spike counts from all the neurons, separately for saccades into (S_{in}) and away from the RF (S_{out}), and then quantified the separation between the two resulting distributions by computing the S_{ROC} (Methods; Fig. S12). Here we present an alternate analysis in which S_{in} and S_{out} conditions are first contrasted for each neuron and then the results are averaged across neurons. Results are for the CRDM (left column, $n = 51$) and the urgent color discrimination task (right column, $n = 56$). **a**, LIP activity during guesses and informed choices sorted by outcome (same short- and long-rPT intervals as in Fig. 3a, d). Activity corresponds to presaccadic spike counts (same as in other analyses) that were z-scored for each neuron and then averaged across neurons. Data are mean $\pm$ 1 SE across cells. **b**, Differential signal magnitudes for the three conditions in **a** indicated by color. Here, the differential signal of each cell is the mean difference between the z-scored responses in S_{in} and S_{out} conditions. Thus, the mean Δz is equal to $\langle z_{IN} - z_{OUT} \rangle$, where the brackets indicate an average is over neurons. Curves are bootstrapped distributions. **c**, Mean neurometric curve, i.e., magnitude of presaccadic differentiation as a function of processing time. In this case, a neurometric curve was first computed for each individual neuron (bin width = 81 ms) and then the results were averaged over neurons. Shades indicate $\pm$ 1 SE across cells. The results of these analyses are more variable than those in the main figures, but show the same qualitative trends for how spatial discriminability depends on processing time and trial outcome in each task.

Supplementary references

101. Busse L, Katzner S, Treue S (2008) Temporal dynamics of neuronal modulation during exogenous and endogenous shifts of visual attention in macaque area MT. Proc Natl Acad Sci USA 105:16380–16385.

102. Bompas A, Sumner P (2011) Saccadic inhibition reveals the timing of automatic and voluntary signals in the human brain. J Neurosci 31:12501–12512.

103. Buonocore A, Purokayastha S, McIntosh RD (2017) Saccade reorienting is facilitated by pausing the oculomotor program. J Cogn Neurosci 18:1–13.

104. Salinas E, Stanford TR (2018) Saccadic inhibition interrupts ongoing oculomotor activity to enable the rapid deployment of alternate movement plans. Sci Rep 8:14163.