Perceptual restoration fails to recover unconscious processing for smooth eye movements after occipital stroke

Sunwoo Kwon[1,2,3], Berkeley K. Fahrenthold[3,4], Matthew R. Cavanaugh[3,4], **Krystel R. Huxlin[2,3,4]* and Jude F. Mitchell[1,2]***

*Equal senior authors

[1]Herbert Wertheim School of Optometry & Vision Science, University of California, Berkeley, CA, USA

[2]Department of Brain and Cognitive Sciences, [3]Center for Visual Science,[4]Flaum Eye Institute, University of Rochester, NY, USA

Corresponding author: Jude F. Mitchell, Dept. Brain and Cognitive Sciences, 360 Meliora Hall, 775 Library Road, University of Rochester, Rochester NY 14627, USA. Email: jfmitch27@ur.rochester.edu

Keywords: Pre-saccadic attention, occipital stroke, hemianopia, vision restoration, eye movements, motion processing

Abstract

The visual pathways that guide actions do not necessarily mediate conscious perception. Patients

with primary visual cortex (V1) damage lose conscious perception but often retain unconscious

abilities (e.g. blindsight). Here, we asked if saccade accuracy and post-saccadic following

responses (PFRs) that automatically track target motion upon saccade landing are retained when

conscious perception is lost. We contrasted these behaviors in the blind and intact-fields of 11

chronic V1-stroke patients, and in 8 visually-intact controls. Saccade accuracy was relatively

normal in all cases. Stroke patients also had normal PFR in their intact-fields, but no PFR in their

blind-fields. Thus, V1 damage did not spare the unconscious visual processing necessary for

automatic, post-saccadic smooth eye movements. Importantly, visual training that recovered

motion perception in the blind-field did not restore the PFR, suggesting a clear dissociation

between pathways mediating perceptual restoration and automatic actions in the V1-damaged

visual system.

Introduction

Human observers use eye movements to bring targets of interest to central vision for detailed inspection. For moving targets, they do so effortlessly, with a combination of rapid saccades and smooth eye movements. When observers acquire a moving target via a saccade, they can continue to track it with smooth eye movements that match the eye's velocity to the motion of the target (Buizza & Schmid, 1986; Lisberger & Westbrook, 1985; Lisberger et al., 1987; Rashbass, 1961; Tychsen & Lisberger, 1986). Those pursuit movements can show accurate velocity tracking, matched to the target from the moment of saccade landing, indicating motion integration and predictive planning for the target prior to the saccade (Gardner & Lisberger, 2001). Other smooth eye movements can occur involuntarily, as in ocular following, where the eyes drift after a saccade in response to the onset of wide-field motion (Gellman et al., 1990; Miles et al., 1986).

Recently, we found that saccade planning to peripheral, static apertures containing motion involuntarily generates predictive, smooth eye movements at the saccade target, even when there are no task demands to follow the target's motion (Kwon et al., 2019). These involuntary smooth eye movements anticipate the post-saccadic motion in the target aperture, generating a low-gain, following response along the target's motion direction, which we named the "post-saccadic following response" (PFR). The PFR appears to reflect automatic, unconscious visual processing that occurs during a saccade target's selection - i.e., during pre-saccadic planning (Kwon et al., 2019). Previous studies of pre-saccadic attention have shown that perceptual enhancements for the saccade target are automatic and obligatory (Deubel & Schneider, 1996; Kowler et al., 1995; Rolfs & Carrasco, 2012; Rolfs et al., 2011), and involve selection of target features such as spatial frequency and orientation (Li et al., 2016), as well as

motion features (White et al., 2013), which can drive smooth eye movements. Thus, the PFR

may represent an automatic consequence of attentional selection for the motion target prior to the

saccade; alternatively, it may play a role in priming the motor system for subsequent tracking of

that target. Critically for the current study, the automatic selection of stimulus motion during

pre-saccadic attention results in PFR, which we will use as a read-out of the motion processing

for smooth eye movements at specific peripheral locations. Both voluntary and involuntary

smooth eye movements are thought to rely on processing of stimulus motion mediated through

neural pathways in the middle temporal (MT) area (Bakst et al., 2017; Mustari et al., 2009;

Newsome et al., 1985; Nuding et al., 2008). Area MT receives strong cortical inputs, routed

through primary visual cortex (area V1), but it also receives direct input from sub-cortical

centers, which bypass V1 (Glickstein et al., 1980; Maunsell & van Essen, 1983; Rodman et al.,

1989; Sincich et al., 2004; Tamietto & Morrone, 2016; Ungerleider et al., 1984; Van Essen et al.,

1981). To what extent these different routes of information that are transferred to MT contribute

to voluntary and involuntary, smooth eye movements, and to perception, remains to be fully

elucidated. Notably, prior studies have suggested that motion pathways driving involuntary

smooth eye movements differ from those mediating perception (Glasser & Tadin, 2014; Price &

Blum, 2014; Simoncini et al., 2012; Spering & Carrasco, 2012; Spering et al., 2011). As such,

the fact that MT receives inputs from sub-cortical centers and from other cortical areas (via V1)

prompted the hypothesis that sub-cortical pathways to MT may support smooth eye movements

while conscious visual motion perception relies predominantly on input routed via V1 (Spering

et al., 2011). Damage to V1 as a result of unilateral occipital stroke offers a unique opportunity

to test this hypothesis in humans. Indeed, unilateral V1-strokes cause a loss of conscious visual

perception in the contralateral visual hemifield (Smith, 1962; Teuber et al., 1960), but sub-

cortical projections to MT are generally spared. Importantly, these projections are thought to

underlie the preservation of unconscious residual abilities such as blindsight (Mazzi et al., 2019;

Sanchez-Lopez et al., 2019; Tamietto & Morrone, 2016; Weiskrantz et al., 1974). A key role of

MT in blindsight has also been inferred from the particular stimulus properties needed to elicit

blindsight: visual targets presented in the blind-field have to be relatively large, coarse, moving

or flickering (Weiskrantz et al., 1995), containing high luminance contrasts, low spatial

frequencies and high temporal frequencies (Sahraie et al., 2008) – stimulus properties that elicit

strong responses from MT neurons (Born & Bradley, 2005; Movshon & Newsome, 1996).

However, the impact of V1 damage on unconscious, motion-dependent visual processes used to

guide smooth eye movements, such as the PFR, has not been investigated.

Moreover, although visual training can restore conscious visual motion perception in

parts of the blind-field of V1-stroke patients (Cavanaugh et al., 2019; Cavanaugh et al., 2015;

Das et al., 2014; Elshout et al., 2016; Huxlin et al., 2009; Saionz et al., 2020; Vaina et al., 2014),

we do not know how effectively patients can use such restored percepts to guide actions. Here,

we used a cued, saccade task to demonstrate remarkably preserved ability of trained, V1-stroke

patients to correctly target motion-containing peripheral stimuli presented in both their intact and

blind-fields. However, by continuously tracking eye movements, we were also – for the first time

– able to capture the impact of V1 damage on the PFR. Our unique data reveal a key role for V1

in this unconscious, automatic, oculomotor behavior. By the same token, they provide new

insights into the likely neural pathways mediating restored, conscious, motion perception after

V1 damage *versus* those involved in the predictive processing necessary for a normal PFR.

Results

To investigate how V1 damage impacts unconscious motion processing for smooth eye

movements, we contrasted the PFR for saccades made to motion stimuli placed in the intact and

the blind-fields of eleven cortically-blind (CB) V1-stroke patients (**Fig. 1**) after visual restoration

training at some of these blind-field locations.

Figure. 1 | Occipital lesions, Humphrey visual field maps and PFR testing locations. Single

radiographic images through the brains of each V1-stroke participant, illustrating region(s) of occipital

damage (red arrows), shown with right brain hemispheres on image right. The location, size and shape of

visual stimuli presented during the PFR testing protocol are indicated by colored circles superimposed on

initial 24-2 Humphrey visual field maps acquired for the tested eye in each case. Red circles: blind-field

testing locations; blue circles: intact-field locations; OS: left eye; OD: right eye; DWI: diffusion-weighted

imaging; FLAIR: T2-weighted fluid-attenuated inversion recovery; CT: computed tomography.

We identified four optimal equi-eccentric testing locations in the intact and blind-fields of each

participant, assessed from Humphrey perimetry **(Fig. 1)**, and measured both the PFR and motion

integration thresholds using random dot stimuli **(Fig. 2A)** at these locations. Our main goal was

to assess how PFR tracked with perception in blind- and intact regions of the visual field, rather

than specifically at trained, blind-field locations. As such, because PFR testing locations had to

be within a certain range of eccentricities, and mirror symmetric across the 4 visual field

quadrants, they did not always fall directly on all trained, blind-field location. The tight location

and task specificity of training-induced global motion recovery in the blind-field (Huxlin et al.,

2009; Das et al., 2014; Saionz et al., 2020) allowed us to attain a large range of perceptual

thresholds at blind-field PFR test locations, against which to plot PFR gain.

Global motion perception at PFR-tested locations was assessed by measuring Normalized

Direction Range (NDR) thresholds **(Table 1;** additional details in Methods). In brief, at each

location of interest, patients were asked to perform a two-alternative, direction discrimination

(left vs. right) task using random dot stimuli in which the range of dot directions was varied in a

3:1 staircase procedure, from 0° (coherent motion) to 360° (random motion). The direction range

for which a criterion performance of 75% correct was attained over 100 trials was then computed,

then normalized to 360° to generate NDR threshold (see Methods) ranging from 1 (poor

performance/unmeasurable threshold) to $\leq$ 0.35 (excellent performance, within the normal range).

Figure. 2 | Experimental paradigms for measuring motion perception and oculomotor functions. A:
Trial sequence for assessing global motion perception: trials started with a fixation period of 1000 ms,
followed by appearance of a random dot stimulus either in the blind or intact-field for 500 ms. Dots
moved globally to the right or left, with a range of directions defined an adaptive staircase. On each trial,
subjects were asked to report the stimulus' global direction of motion by pressing the left or right arrow
keys on a keyboard. They received auditory feedback on the correctness of each response. **B:** Trial
sequence for assessing oculomotor behavior: each trial started with a variable fixation period after which,
participants were presented with 4 equi-eccentric motion apertures and a spatial cue at fixation. Dot
motion apertures were Gaussian-enveloped and contained 100 % coherent motion along a randomly
assigned direction (clockwise or counter-clockwise) in each aperture, that was tangential to the center-out
saccade to the aperture. The spatial cue (50 ms) indicated a peripheral target aperture towards which the

participant was instructed to initiate a saccade as fast as possible. In half the trials, the dot motion stimuli

persisted for 500 ms, and were thus present upon saccade offset. In remaining trials, stimuli disappeared

during saccade flight, such that no stimulus motion was present at the fovea upon saccade landing.

In the blind-fields, depending on whether the tested locations overlapped with a trained location,

performance was either at chance or had improved to measurable and sometimes near-normal

NDR thresholds, assessed in each patient's own intact visual field **(Table 1)**. The end result was

a set of 14 blind-field locations across 11 patients, where perceptual thresholds ranged from

unmeasurable (reflecting inability to do the task) to near-normal.

Table 1. Demographics and global motion integration thresholds in retrained, V1-stroke

participants. M, male; F, female; NDR, normalized direction range (low value=good performance,

1=bad performance/no measurable threshold). Each NDR threshold denotes performance measured at a

single blind- or intact-field location in each patient (see **Fig. 1** for positioning of these locations relative to

the pre-training Humphrey visual field). Pre-training NDR thresholds were always 1 in blind-fields.

Subject	Sex	Age (years)	Time post-stroke (months)	Blind-field NDR thresholds	Intact-field NDR thresholds
CB1	F	27	65.0	1	0.3
CB2	F	68	24.8	0.2	0.2
CB3	F	57	48.6	0.7, 0.8	0.3
CB4	M	66	32.2	0.5	0.2, 0.2, 0.2
CB5	M	54	52.2	1	0.3, 0.3
CB6	M	79	22.7	1, 0.1	0.1, 0.2
CB7	M	53	36.8	1	0.3
CB8	M	52	65.5	1, 0.2	0.1, 0.2
CB9	F	65	56.5	0.3	0.1, 0.1, 0.02

| CB10 | F | 31 | 66.8 | 0.3 | 0.2 |
| CB11 | F | 43 | 6.0 | 0.1 | 0.3, 0.2, 0.2 |

Oculomotor behavior was measured using a cued-saccade task for motion stimuli at 4 peripheral

locations (colored circles in **Fig. 1; Fig. 2B**). Of particular note, the motion direction inside the

apertures was irrelevant to the task and subjects were not asked to track it or report it.

Additionally, in half the trials, the motion stimulus disappeared during saccade flight to

disambiguate the contributions of pre-saccadic motion processing to the post-saccadic following

response.

Basic saccade behavior of V1-stroke participants – accuracy and latency

Stroke participants were generally able to use central spatial cues at fixation to plan saccades to

peripheral aperture locations. We first quantified saccade accuracy in a binary fashion by

whether saccade end-points fell within the diameter of the stimulus aperture. We then also

assessed spatial accuracy by the mean absolute error in position of the saccade end-point from

the aperture center. A saccade was labelled "correct" when it fell at least 3.5˚ from the saccade

target center within 90 ms of the eye leaving the fixation window. Stroke participants made

slightly more correct saccades to targets in their intact-field (mean 96.8 ± 3.7 %) than to those in

their blind-field (mean 89.4 ± 15.5 %), likely reflecting more reliable target identification in the

intact-field (t_{42}=2.5078, p=0.016, BF=3.4414). However, when correctly selecting targets in their

blind-fields, there was high spatial accuracy of saccades to those targets, as measured by the

location of the end-points relative to the aperture center. Specifically, stroke participants had a

mean absolute landing error relative to the stimulus center of 1.59 ± 0.07 deg (SEM) in the

intact-field and 1.59 ± 0.08 deg (SEM) in the blind-field, which was not significantly different

(t_{10}=0.1946, p=0.8496, BF=0.3024). The latency of saccades was also similar for blind- and

intact-field targets (intact-field latencies: 360 ± 45 ms; blind-field latencies: 382 ± 57 ms; t_{42}=-

1.3593, p=0.181, BF=0.6472).

Finally, we compared saccade accuracy and latency of stroke patients to those of

visually-intact controls from a previous study (Kwon et al., 2019). Saccade accuracies in the

intact-fields of stroke patients were slightly impaired compared to those of visually-intact

controls (t_{36}=1.5634, p=0.1267, BF=0.8863) who exhibited saccade accuracy of 98.7 ± 0.8 % as

compared to 96.8 ± 3.7 % for stroke patients. However, stroke patients had significantly longer

(t_{36}=-5.4737, p<0.0001, BF>1000) saccade latencies of 360 ± 45 ms in their intact-fields,

compared to visually-intact controls, whose latencies averaged 260 ± 10 ms (Kwon et al., 2019).

We consider several possible causes for these differences in the Discussion, including age, and

challenges specific to saccade planning in the presence of a blind-field.

Predictive oculomotor behavior in the intact-field of V1-stroke participants

In intact portions of their visual fields, stroke patients' post-saccadic smooth eye movements

reflected the direction of target motion of the pre-saccadic stimulus immediately upon saccade

offset. In a typical trial, a saccade made to a target aperture in the intact-field exhibited a smooth

drift in eye position from the saccade end-point along the direction of target motion (see example

for a single stroke patient in **Fig. 3A**). We quantified the time course of the drift in eye position

by computing the eye velocity projected along the direction of target motion, where positive

values reflect following of motion. We term this the PFR velocity. Across patients, we observed

a net positive PFR velocity (**Fig. 3C**). By including a stimulus manipulation in which the motion

target disappeared during saccade flight, we were able to confirm that the PFR velocity was

driven exclusively by pre-saccadic motion in the peripheral aperture. Within the first 100 ms

after saccade offset (*the "open-loop" period*) the PFR velocity did not differ depending on

whether the stimulus remained present after the saccade (red trace in **Fig. 3C**) or if it was

removed during saccade flight (blue trace in **Fig. 3C**). By including trials where we removed the

stimulus in-flight, we could eliminate direct post-saccadic stimulation of motion at the fovea, and

thus isolate the predictive, "open-loop" component of the PFR velocity. After 100 ms from

saccade offset, the presence or absence of foveal motion did influence post-saccadic eye

movements. Specifically, the PFR velocity continued to increase along the target motion

direction in the foveal-motion-present condition (red trace in **Fig. 3C**), whereas it decreased

when no stimulus was present upon saccade landing (blue trace in **Fig. 3C**).

Figure. 3 | Oculomotor behavior in perceptually-trained, V1-stroke participants. A: Eye-movement

traces to a cued target in the intact-field (white background) of a single V1-stroke patient in a stimulus

absent condition (i.e. with no foveal motion upon saccade landing at the cued target). Small, connected

black dots denote the raw eye movement sampled from our eye tracker; the green circle represents the

electronic window around the fixation spot; random dot stimuli were presented inside the 4 dashed circles

and their global motion direction is indicated by large arrows inside each circle. Note the accurate saccade

to the target center, and how the eye follows the pre-saccadic target motion direction (red arrow) upon

saccade landing. **B**: Raw eye movement traces to a cued target in the blind-field (grey background) of a

single V1-stroke patient in a stimulus absent condition. Note successful saccade landing onto the cued

target but how the eye fails to follow the pre-saccadic target motion direction. Labeling conventions as in

A. **C**: Eye velocity traces for saccades to intact portions of the visual field, averaged across intact-field

locations per patient then averaged across all 11 stroke patients. In half the trials, stimuli were present

upon saccade landing, resulting in foveal motion (red trace). In the remaining trials, stimuli were absent -

i.e. there was no foveal motion upon saccade landing on the target (blue trace). Error bars = 2 SEM across

subjects. Average eye velocities were projected along the target motion direction time-locked prior to the

saccade onset (-100 to 0 ms) and offset (0 to 200 ms), such that positive values reflected motion

consistent with the stimulus, while negative values reflected motion opposite. **D***:* Eye velocity traces for

saccades to blind portions of the visual field, averaged across blind-field locations per patient and then

averaged across all 11 stroke patients (same conventions as in A). Error bars = 2 SEM across subjects.

Note the near-zero eye movement velocity during the "open-loop" period, reflecting the lack of PFR.

For subsequent analyses, we focused on the "open-loop" period (i.e., within 100 ms after saccade

offset), as the PFR velocity in this period depends on pre-saccadic motion information

accumulated from the peripheral target aperture (Kwon et al., 2019). The average velocity of

these open-loop following responses, both for stimulus-present and -absent trials, was small in

magnitude relative to the stimulus speed (10 deg/sec), ranging from 5-15 % relative velocity

gain. These lower gain responses are consistent with an involuntary following response, such as

ocular following (Gellman et al., 1990; Miles et al., 1986), rather than voluntary pursuit of a

target. Overall, the pattern of oculomotor behavior in the intact-field of stroke patients was

highly similar in its time-course and magnitude to that previously measured in visually-intact

controls (Kwon et al., 2019).

Figure. 4 | PFR gain in the intact and blind-field of V1-stroke participants. A*: Plot of mean PFR gain

in visually-intact controls (Kwon et al., 2019), and in V1-stroke patients' intact and blind-fields during

the open-loop period. The PFR gain is here represented as a proportion of pursuit gain along the target

motion direction (1 = perfect following, 0 = no following, -1 = following in the opposite direction).

Individual dots represent the mean PFR for each participant. There was no significant difference in PFR

gain between the intact-fields of stroke patients and visually-intact controls. PFR gain approximated 0 for

saccades to targets in stroke patients' blind-fields. Error bars represent 2 SEM across subjects. **B***: PFR

gain for individual intact-field locations (open circles) and **C:** blind-field locations (grey circles) as a

function of the normalized direction range (NDR) threshold measured at each location in stroke

participants. Within intact regions of the visual field, there was an inverse correlation between this global

motion threshold and PFR gain: the lower the threshold (i.e. the better the motion perception), the higher

the PFR gain. This relationship was lost in the blind-field of stroke patients, with no significant

correlation between NDR thresholds and PFR gain – the latter remaining abnormally low.

Visually-intact controls in our prior study had a net positive PFR gain in the open-loop epoch

that differed significantly from zero (t_7=4.31, p=0.004, BF=14.2833 – dark grey bar in **Fig. 4A**).

PFR gain in the intact-field of stroke patients (white bar in **Fig. 4A,** 15.7% gain, CI_{95} = [0.1564,

0.1576]) was not significantly different from PFR gain in our prior, visually-intact controls (t_{17}=-

0.577, p=0.5715, BF=0.4578), and also showed a net positive effect that differed significantly

from zero (t_{10}=5.4058, p<0.0001, BF>100). Finally, we noted a significant correlation (R^2=0.351,

F_{18}=9.,73, p=0.0059, BF=7.3164) between perceptual performance measured by NDR thresholds,

and the magnitude of the PFR (white circles, **Fig. 4B**) in the intact-field of stroke participants.

Predictive oculomotor behavior in the blind-field of V1-stroke participants

Although saccades landed correctly on target stimuli in the blind-field of our stroke participants

~89 % of the time, post-saccadic eye movements differed dramatically from those in the same

participants' intact-field. Specifically, in the open-loop period, the eyes no longer moved along

the direction of motion in the target (see example in **Fig. 3B**) – in other words, there was no

positive PFR velocity. This pattern was reflected in the average PFR velocity across all 11 stroke

patients' blind-fields (**Fig. 3D**), irrespective of visual rehabilitation training. In contrast to

oculomotor behavior when making a saccade to targets in their intact-fields, stroke patients did

not show any positive PFR velocity within the first 20-100 ms after saccade offset ("open-loop"

period), whether the motion stimulus was present (red trace) or absent (blue trace) post-

saccadically (**Fig. 3D**).

Beyond 100 ms from the saccade offset, post-saccadic foveal motion – when present –

was sufficient to drive an ocular following response (red trace, **Fig. 3D**). The PFR during this

later "closed-loop" period (100-200 ms) was driven by foveal motion for blind-fields as much as

the intact-fields (**Supplementary Fig. S1**). PFR gain was positive and significant both for the

intact-fields (18.6% gain, CI_{95} = [18.5%, 18.6%], t_{10}=4.2314, p=0.0017, BF=25.4627) and the

blind-fields (15.0% gain, CI_{95} = [14.9%, 15.0%], t_{10}=8.4130, p<0.0001, BF>1000) with no

significant difference between them (t_{10}=1.2041, p=0.2563, BF=0.5355). This is an important

observation, as it confirms that post-saccadic ocular following remained functional in these

patients. Only the predictive component during the open-loop period was abnormal, reflected by

the absence of the PFR velocity in stimulus-absent trials (blue trace, **Fig. 3D**). Consistent with

these observations, PFR gain in the blind-field of our stroke patients (light grey bar in **Fig. 4A,**

0.45% gain, CI_{95} = [0.43%, 0.47%]) was significantly lower than PFR gain in their intact-field

(t_{10}=5.6451, p=0.0001, BF=150), and was not significantly different from zero (t_{10}= 0.3796,

p=0.7122, BF=0.3166).

Since portions of the blind-fields of stroke patients underwent training that restored

global motion perception, we next asked if such training, when it was coincident with locations

tested on the PFR task, restored PFR gains. Perceptual recovery did not restore PFR: there was

no significant correlation between NDR and PFR gains (R^2=0.0655, F_{12}=0.841, p=0.377,

BF=0.3020) in the blind-field of stroke patients (grey circles, **Fig. 4C**). At blind-field locations

where post-training stroke patients attained NDR thresholds <1, the PFR gain was not

significantly different from 0 (mean PFR gain = 0.0177 ± 0.07, t_8=0.7524, p=0.4734,

BF=0.4066). Even when we isolated blind-field locations where training restored "normal" NDR

thresholds ($\leq$ 0.35), contrasting them against blind-field locations where NDR thresholds were

>0.35, PFR gain was not significantly different between those groups (t_{12}=-0.4140, p=0.6862,

BF=0.4749). Since many blind-field locations failed to recover perception to the same range as

the intact-fields (NDR $\leq$ 0.35), we examined a subset of blind-fields (N=6) where discrimination

was matched to performance of intact-fields. Still we found there was no significant PFR for

those blind-fields (t_5=0.5411, p=0.6117, BF=0.4206) and that subset differed significantly from

the intact-fields (t_{24}=2.7424, p=0.0057, BF=9.0755), and further the correlation remains non-

significant for blind-fields in that range (R^2=0.362, F_4=2.267, p=0.2066, BF=0.6993). In an

additional analysis, we fit an exponential curve to describe how PFR varied with NDR for the

intact-fields, and asked if PFR at blind-field location that recovered NDR (i.e., NDR $\leq$ 0.35) fell

within its confidence intervals. Only 1 of 6 such blind-field locations overlapped the intact-field

confidence intervals (**Supplementary Fig. S2**), providing strong evidence against those

distributions being matched across NDR (p $<=$ 1.79x10^{-6}, binomial test).

Finally, we should note that the above analyses (**Fig. 4B-C; Supplementary Fig. S2**)

treated individual visual field locations from the same participant as independent samples. Prior

work in CB patients suggests that both baseline/pre-training performance and the ability of a

location to improve are not uniform across blind-fields, even at identical eccentricities; training-

induced changes in discrimination performance are tightly restricted to the trained, blind-field

locations and do not spread by more than 1˚ outside the boundaries of each trained location (Das

et al., 2014; Huxlin et al., 2009; Sahraie et al., 2006; Saionz et al., 2020). Nonetheless, we

repeated our analyses by pooling the intact and blind-fields within each participant and

confirmed that our results remained consistent (**Supplementary Fig. S3**). In conclusion, over

several tests, we found no significant recovery of PFR gain with recovery in perceptual motion

performance.

Discussion

For the first time, the present study measured the impact of V1 damage and subsequent, training-

induced, visual restoration on both voluntary saccadic behavior and a class of unconscious,

automatic, post-saccadic smooth eye movements: the PFR. Our findings revealed an unexpected,

critical reliance of pre-saccadic visual motion processing on visual pathways that include V1. We

first confirmed that V1-stroke patients exhibit normal saccade accuracy and normal PFR when

saccading to motion targets in intact regions of their visual fields, where vision is mediated by

intact V1. However, while the same patients exhibited normal saccade accuracies for targets

presented in their blind-fields, they had no measurable PFR, even after visual discrimination

training recovered global motion perception at those blind-field locations. Thus, restoration of

motion perception did not automatically restore the unconscious visual motion processing

necessary for the PFR. This is surprising because traditionally, patients with V1 damage are

well-known for having preserved, unconscious visual processing in their blind-fields – under the

umbrella of blindsight phenomena (reviewed in Weiskrantz, 1996, 2009). That visual

discrimination training can recover the ability to perform the relatively complex computations

needed to integrate motion direction into a global percept available to consciousness – even to

the extent of attaining normal NDR thresholds - inside chronic blind-fields is remarkable. That

this could occur without automatically restoring the unconscious global motion processing

necessary for predictive smooth eye movements was unexpected. Our findings suggest that

primary visual cortex (V1) may be key for both conscious visual perception (Tong, 2003) and for

unconscious visual processes that influence smooth eye movements induced by peripherally-

presented motion targets. They also suggest that visual restoration, after V1 damage, recruits

different neural circuits than are normally used for these processes in the intact visual system;

finally, it suggests that these newly-engaged circuits now dissociate conscious and unconscious

visual motion processing.

The extrastriate visual area critical for many aspects of visual motion processing - area

MT - receives strong inputs from V1 as well as from sub-cortical projections that bypass V1

(Glickstein et al., 1980; Hagan et al., 2019; Maunsell & van Essen, 1983; Rodman et al., 1989;

Sincich et al., 2004; Ungerleider et al., 1984; Van Essen et al., 1981). This diversity of inputs to

MT likely explains why, after V1 damage, residual visual motion processing persists inside the

resulting blind-fields (reviewed in Das & Huxlin, 2010; Melnick et al., 2016; Tamietto &

Morrone, 2016). Key for rehabilitation efforts, this residual processing can be leveraged by

intensive visual training to recover both simple and complex motion perception (Cavanaugh et

al., 2019; Das et al., 2014; Huxlin et al., 2009; Saionz et al., 2020).

In non-human primates, activity in area MT has been causally linked to perceptual reports

of motion in discrimination and detection tasks (Britten et al., 1992; Newsome et al., 1985;

Salzman et al., 1990; Siegel & Andersen, 1986), and to accuracy in pursuit eye movements

(Huang & Lisberger, 2009; Newsome et al., 1985; Osborne et al., 2005; Salzman et al., 1990;

Siegel & Andersen, 1986). The perception of velocity is also well correlated with velocity gain

in voluntary pursuit, supporting the notion that pursuit and perception share common motion

processing at the neural level (Gegenfurtner et al., 2003; Spering et al., 2005; Stone & Krauzlis,

2003). Voluntary pursuit and involuntary ocular following responses, such as the PFR, are also

thought to rely on motion processing by the dLGN, V1, as well as area MT and the medial

superior temporal (MST) area (Bakst et al., 2017; Mustari et al., 2009; Nuding et al., 2008;

Takemura et al., 2007). Just as pursuit is modulated by stimulus contrast (Spering &

Gegenfurtner, 2007; Spering et al., 2005), the PFR also exhibits a dependence on stimulus

contrast; in visually-intact humans, we saw a steep rise of the PFR contrast response function

starting right below 10 % luminance contrast, quickly reaching saturation at or above 15 %

contrast (**Supplementary Fig. S4**). This is comparable to the contrast response function of

neurons in macaque area MT (Heuer & Britten, 2002; Kohn & Movshon, 2003; Sclar et al.,

1990). However, pathways for ocular following are thought to be at least partly non-overlapping

with those involved in motion perception (Glasser & Tadin, 2014; Price & Blum, 2014;

Simoncini et al., 2012). Other studies in humans also support a distinction between neural

circuits underlying smooth eye movements and conscious motion percepts (Spering & Carrasco,

2012; Spering & Gegenfurtner, 2007; Spering et al., 2011). Consistent with those studies, we

now find that despite recovery of processing used for accurate, global motion perception in the

blind-field of V1-damaged humans, the PFR remains absent at trained, blind-field locations.

One possible explanation for this outcome [disassociation between perception and PFR]

is that training post-stroke lowered the thresholds used for reading out motion information in

area MT for perception but not smooth eye movements. If motion perception in the blind-field

was consistent with weak residual sensory signals, and the PFR response has a higher contrast

threshold than perception, then we could observe a lack of PFR while still retaining motion

discrimination at blind-field locations. Contrast thresholds for global motion stimuli in normal

participants range between 5-10% contrast (Fine et al., 2004). We examined PFR as a function of

contrast in normal participants to determine if it might have a higher threshold than that for

perception, but found thresholds were in a comparable range (Supplementary methods,

Supplementary Figure S4). Thus, we consider it unlikely that the lack of PFR reflects a

difference in threshold present in normal populations. However, there remains a possibility that

perceptual training in patients may cause lower the threshold used for perception to

accommodate weaker motion signals in area MT following V1 damage, but that a similar change

in threshold did not transfer to the read-out used for oculomotor signals. This would reflect a

disassociation between perception and eye movements at the level of read-out from area MT,

which would still have important implications for visual recovery and understanding resulting

differences in action and perception.

Another possible explanation for this disassociation is that while training post-stroke

improved processing for perception, it did not correct problems with pre-saccadic attention

and/or other aspects of saccade planning. It is well established that sensory processing among

neurons in MT and MST can be strongly influenced by attention (Treue & Maunsell, 1996;

Treue & Trujillo, 1999) and also by target selection immediately prior to saccades (Ferrera &

Lisberger, 1997; Recanzone & Wurtz, 2000). Recent studies suggest that like target selection in

voluntary pursuit, selective attention can modulate ocular following responses (Souto & Kerzel,

2014), and our findings here and previously (Kwon et al., 2019) support this notion. Pre-saccadic

attention is thought to operate through feedback from oculomotor planning areas to visual cortex

(Moore & Armstrong, 2003; Moore & Fallah, 2004), and while its impact has been studied

mainly in visual area V4, it is also thought to occur in MT/MST. Indeed, electrical micro-

stimulation in an oculomotor area, the frontal eye fields (FEF), influences selection of motion

signals prior to saccades and can alter subsequent saccade trajectories to favor stimulus motion

(Schafer & Moore, 2007). That the FEF and its projections to area MT are intact in V1-stroke

patients suggests preservation of pre-saccadic planning and attention selection for the saccade

target even when visual input is weak or abnormal in a blind-field.

Although the effects of attention have not been studied extensively in V1-damaged

patients, work to date suggests that some attentional mechanisms remain functional within

cortical blind-fields; as such, they could modulate motion signals at the level of MT/MST in the

current behavioral paradigm. For instance, covert spatial attention was reported to improve

stimulus detection in the blind-field (Poggel et al., 2006) and in a separate study, it was shown to

significantly decrease reaction times in V1-stroke patients performing an orientation

discrimination task without any speed-accuracy trade-off (Kentridge et al., 2004). Feature-based

attention was also able to improve fine direction discrimination training in cortically-blinded

fields (Cavanaugh et al., 2019). One piece of evidence suggesting that pre-saccadic attention

remains functional in the current experiments is that other aspects of saccade pre-planning

related to perceptual shifts in the position of motion targets, remain in the blind-field. Previous

studies reported a motion-induced perceptual shift for stimulus location along the direction of

stimulus motion (De Valois & De Valois, 1991; Nishida & Johnston, 1999; Ramachandran &

Anstis, 1990; Whitney & Cavanagh, 2000), which for saccades is reflected by a shift in their

end-points along the direction of target motion (Kosovicheva et al., 2014; Kwon et al., 2019;

Schafer & Moore, 2007). For stroke patients, we confirmed similar shifts in saccade end-points

in their intact-fields, as well as significant, albeit reduced shifts along the target motion in their

blind-fields (**Supplementary Fig. S5**). We also considered if the lack of PFR in the blind-field

may have resulted from inability to allocate attention for the saccade into the blind-field, against

the competing influence of distractor apertures within intact portions of the visual field. First, we

ran a control experiment in two patients to compare PFR when competing distractors were absent

(single aperture task) against the original design with four apertures (**Supplementary Fig. S6A**).

A central line cue directed patients to the saccade target, and as in the main experiment, their

saccades remained accurate for both intact (98.6 ± 1.2 %) and blind-field (98.7 ± 0.24 %)

locations. Importantly, we did not observe a measurable PFR in the blind-fields when distractor

motion was removed, and PFR remained robust in the intact-fields comparable to the original

design (**Supplementary Fig. S6B**). There also remained a correlation between PFR and NDR

thresholds for intact-fields, but not for the blind-fields (**Supplementary Fig. S6C**). Thus,

removing distractors had little impact on the PFR, suggesting the deficit was sensory in nature

rather than involving a misallocation of attention. As a final control analysis, we also measured

to what extent PFR was driven by the motion in distractor apertures. If the motion in the other

three apertures produced any measurable influence on PFR, then one might expect that influence

to be stronger when saccades were made into the blind-field because attention could not be

properly allocated to its location, resulting in higher attention to distracter locations. While we

did find that motion in distractor apertures contributed to a weak PFR gain that was significant

and measurable, contrary to that hypothesis, the gain was stronger for saccades made into intact

visual fields as compared to blind-fields (**Supplementary Fig. S7**). Thus, it seems unlikely that

the lack of PFR reflects an impairment to engage pre-saccadic attention into the blind-field.

Instead, we posit that perceptual recovery through repetitive discrimination training did not

entrain the specific motion processing pathways that support post-saccadic following.

An important consideration for the present experiments was whether failures to elicit

PFRs in cortically-blinded portions of the visual field might simply reflect a motor deficit for

accurately targeting peripherally-presented motion apertures. Because the PFR requires pre-

saccadic attention to select the target motion, any loss in target localization accuracy could

impair selection. Previous studies in monkeys with V1 lesions have found reduced spatial

accuracy for saccades made into the blind-field with larger end-point errors from 0.2 to 0.6

degrees at matched eccentricities for intact and blind-fields (estimated from data in Fig. 2 of

Yoshida et al., 2008). In the present study, we observed that stroke patients were less likely to

correctly *select* a target aperture in their blind *versus* intact-fields when given a central spatial

cue (87 % vs. 96 %). However, when the target was correctly selected, spatial accuracy of the

saccade end-points was normal (1.31° *versus* 1.38° absolute error relative to the target's center).

A key difference in the prior study is that Yoshida *et al.* (2008) used smaller stimuli, measuring

only 0.45 degrees in diameter, while we used large, dot-motion fields (5.5 degrees in diameter,

Gaussian enveloped - ***see Methods***). Because spatial accuracy of saccade landings was similar

for blind and intact-fields of our stroke participants, we conclude that a motor deficit for

targeting peripheral motion apertures in cortically-blind regions of the visual field was not a

likely explanation for the absence of a PFR.

The slightly larger number of errors made by stroke patients when selecting cued targets

in their blind-fields could also be consistent with a reduction in their relative, perceived salience.

Although all 4 motion apertures appeared simultaneously, iso-eccentrically and had equal

[~100 %] luminance contrast, it is possible that stroke participants required extra effort to ignore

blind-field-related perceptual inhomogeneities between the four apertures. Indeed, a prior study

in chronic V1-stroke patients showed depressed luminance contrast sensitivity for motion and

orientation discriminations at trained, blind-field locations, in spite of normal NDR thresholds at

these locations (Das et al., 2014). Thus, when patients are cued to saccade to a presumably less

salient target in their blind-field, this may require additional effort to suppress a reflexive

saccade to the more salient targets. In an anti-saccade task where a salient target is ignored in

order to plan a cued movement to the opposite [but empty] visual field, there is typically a

reduction in saccadic reaction time (Hallett, 1978; Munoz & Everling, 2004). If saccadic reaction

times slow down by 140 ms or more due to task difficulty, this impacts saccades to both visible

and anti-saccade locations due to the extra volitional demands (Hallett & Adams, 1980). In line

with this observation, we found that saccade reaction times were slower for stroke patients by

roughly 100-130 ms, both for the intact and blind-fields, relative to visually-intact controls.

Controls did differ in age from stroke patients (~20 years old *versus* ~57 years old), and prior

work showed a correlation between slower saccade latency and growing age; however, the

typical reduction from 20 to 80 years of age was 40-45 ms (Abel et al., 1983; Pirozzolo &

Hansch, 1981; Spooner et al., 1980). Therefore, it appears unlikely that age alone would account

for the >100 ms reduction in saccadic reaction times in stroke patients. Rather, we posit that

patients had to exert greater volitional control to select the cued target inside blind regions of

their visual field.

In summary, V1 damage in humans, such as occurs from occipital stroke, causes a

dramatic loss of conscious visual perception across large regions of the visual field, impairing

most aspects of daily living. Paradoxically, this condition was famously known for its relative

preservation of unconscious visual processes, such as those mediating blindsight. With the

advent of visual restoration training for this patient population, an important question in the field

has been to ascertain what aspects of visual processing can recover, which cannot, and why.

Peripheral visual motion processing is key to many aspects of daily living. Not only is it critical

for accurate perception and identification of targets, it is also essential for our motor actions and

reactions to these targets. Here we show that visual training that can restore perceptual

discrimination of peripheral motion does not automatically recover the PFR [or normal saccade

targeting to peripheral motion stimuli]. Our findings support a dissociation between smooth eye

movements, saccade targeting and perception following V1 damage, and suggest that V1 is

critical for driving smooth eye movements such as the PFR. A key realization emerging from

these results is that *alternative* pathways, which convey motion information from subcortical

centers directly to area MT, are insufficient to support predictive oculomotor behaviors when V1

is damaged, even if they are sufficient to mediate recovery of conscious motion perception.

A second insight attained presently is that repetitive motion discrimination training in CB

fields might only influence circuits and processes that support conscious perception, without

transfer to those driving motion-dependent, unconscious behaviors, such as the smooth eye

movements involved in the PFR. Importantly, prior to global motion training, CB patients are

typically unable to describe or even correctly guess global motion direction of random dot

stimuli in their blind-fields – as such, they have no demonstrable blindsight for the direction

integration task used here (Das et al., 2014; Huxlin et al., 2009; Saionz et al., 2020), and for

other global motion tasks (Azzopardi & Cowey, 2001; Vaina et al., 2014). By the end of training,

CB patients in the present study could correctly describe random dot stimuli and attain relatively

normal NDR threshold levels of performance – implying recovery of conscious perception and

not simply more effective blindsight. A dissociation between processing for conscious vision and

action in the context of restoring vision in CB naturally leads to parallels with the work of Milner

and Goodale (1992), which suggested that ventral stream projections from striate cortex to the

inferotemporal cortex play a major role in perception, while dorsal stream projections from

striate cortex to posterior parietal regions mediate the required sensorimotor transformations for

visually guided actions. Our current results suggest that after V1 damage, psychophysical

training enables residual motion processing, likely in area MT and related dorsal pathway areas,

to be used to recover perception without recovering the necessary processing for smooth eye

movements. This suggests a dissociation between perception and visually-guided actions in this

rehabilitation paradigm.

It remains to be determined if deliberate training on tasks that focus on saccade planning

to motion targets might recover predictive motor behaviors. Rehabilitation of predictive ocular

behaviors remains an uncharted area of research for V1-stroke patients, even though saccade

training is one of the few forms of rehabilitation more readily available to these patients

(Kerkhoff, 1999, 2000; Kerkhoff et al., 1992; Mannan et al., 2010; Nelles et al., 2001; Ong et al.,

2015; Pambakian et al., 2004; Roth et al., 2009; Sahraie et al., 2016; Spitzyna et al., 2007;

Trauzettel-Klosinski, 2010; Weinberg et al., 1977; Zihl, 1980). Of relevance to our observation

of an apparent dissociation between perception and eye movements after V1 damage, training

patients to saccade to targets in their blind-field does not induce perceptual recovery (Campion et

al., 1983; Pollock et al., 2019). Nonetheless, it is conceivable that an approach combining

perceptual training with training of ocular behaviors could improve the efficiency with which

patients use information from their blind-fields in everyday life.

Materials and Methods

Participants: eleven participants with long-standing cerebral blindness (CB) were recruited 2 to

5 years after a stroke that damaged their V1 unilaterally or in one case, bilaterally (see **Table 1**

for details). The location and nature of V1 damage was verified from clinical brain imaging

performed as part of each patient's standard of care. Homonymous visual field defects were

confirmed using monocular, Humphrey automated perimetry performed at the Flaum Eye

Institute of the University of Rochester (**Fig. 1**). Participants suffering from neglect, cognitive

impairments or ocular diseases were excluded from enrollment, as were those using psychoactive

drugs. All participants had their visual acuity corrected to normal (with glasses or contact lenses)

during testing.

Testing of V1-stroke participants occurred following completion of separate, visual

training studies whereby they underwent visual training at one or more blind-field locations.

Some of these trained, blind-field locations overlapped with PFR testing locations - some did not.

The end result was a set of 14 blind-field locations from 11 patients, where pre-training

performance was initially at chance – i.e. participants were unable to reliably discriminate left

from right coherent, global motion. Post-training however, depending on whether the tested

locations overlapped with a trained location, performance either remained at chance or improved,

generating measurable and occasionally near-normal direction integration (NDR) thresholds

(**Table 1**). For comparison we also measured performance at a set of 13, iso-eccentric, intact-

field locations (**Table 1;** see below for details of global motion assessment methods).

PFR data from stroke patients were contrasted with a previously published data set

obtained from eight, visually-intact controls (18 to 22 years old; 4 females and 4 males) who had

normal or corrected-to-normal vision (Kwon et al., 2019).

All procedures were approved by the Institutional Review Board of the University of

Rochester and adhered to the tenets of the Declaration of Helsinki. Written, informed consent

was obtained from each participant, and participation was at all times completely voluntary.

Testing and visual training to recover perception: In each participant, visual field deficits

were first estimated from Humphrey visual perimetry (Cavanaugh & Huxlin, 2017). This served

as a starting point to map the position of the blind-field border and select locations for PFR

testing. It is important to note however, that spared Humphrey visual field performance (i.e.

luminance detection) was not necessarily representative of spared motion discrimination at a

given blind-field location – some patients were unable to attain normal global motion thresholds

at locations partially spared on Humphrey perimetry (**i.e. CB6 blind-field location 2, Fig. 1,**

Table 1, Supplementary Fig. S2). Blind-field training was performed as previously-described

(Cavanaugh et al., 2022; Cavanaugh et al., 2015; Das et al., 2014; Huxlin et al., 2009; Saionz et

al., 2020). In brief, participants were assigned to train daily at home using a custom MATLAB-

based software on their personal computers and displays. They were supplied with chin/forehead

rests and instructed to position them such that the eyes were 42 cm away from their displays

during training. They performed 300 trials per training location per day, at least 5 days per week,

and emailed their auto-generated data log files back to the laboratory for analysis weekly.

Session thresholds were calculated by fitting a Weibull function with a threshold criterion of

75% correct performance. Once performance reached levels comparable to equivalent intact-

field locations, training moved 1 degree deeper into the blind-field along the x-axis (Cartesian

coordinate space). When subjects returned to the lab, discrimination performance at all home-

trained locations was verified with on-line fixation control using the Eyelink 1000 eye tracker. In

addition, we measured NDR thresholds at blind and intact-field locations chosen to undergo

testing on the PFR task (see below). Intact-field measurements were used to obtain a normative

range of NDR thresholds that was specific to each participant (internal control for blind-field

performance). Ultimately, this enabled us to assess whether PFR was tied to perceptual motion

integration performance in both intact and impaired regions of the visual field.

The locations used for PFR testing were mirror symmetric across the vertical and

horizontal meridians, and ranged in eccentricity from 5.8 to 10.4 degrees with a mean of 7.7

degrees. This was done to maintain consistency with the original study (Kwon et al., 2019) that

measured PFR in visually-intact participants, whose stimuli uniformly sampled ordinal and

cardinal directions at 7.1 degrees eccentricity. Partly because of this experimental requirement,

PFR-testing locations did not always coincide with all trained locations in CB patients' blind-

fields. However, this was not a problem, as our main goal here was to ascertain if and how PFR

gain tracked with NDR performance. Thus, we sampled regions of the blind-field that underwent

training and recovered normal NDR performance [CB2, CB6 location 2, CB8 location 2, CB9,

CB10, CB11], as well as locations that were not directly trained or were trained on a non-global

motion task, causing NDR thresholds to remain at 1 [CB1, CB5, CB6 location 1, CB7, CB8

location 1] or improve only partially [CB3 locations 1 and 2, CB4]. In sum, the specificity of

recovery to trained locations and tasks in CB fields (Das et al., 2014; Huxlin et al., 2009; Sahraie

et al., 2006; Saionz et al., 2020) allowed us to attain a wide range of NDR threshold outcomes,

against which to map PFR gain.

Apparatus & eye tracking for assessing global motion perception: participants were asked to

perform 100 trials of a 2-alternative, forced-choice, left-*versus*-right, global direction

discrimination task at 2 to 4, equi-eccentric, peripheral visual field locations chosen for testing of

predictive oculomotor behavior (circles superimposed on Humphrey visual fields in **Fig. 1**; red:

blind-field locations, blue: intact-field locations). All blind-field locations were tested in each

patient (red circles in **Fig. 1**). Time limitations restricted our ability to measure performance at

every intact-field location (blue circles in **Fig. 1**), but at least one intact-field location was

assessed in each participant. Across intact-field locations tested, we saw normal NDR thresholds

that varied from 0.1-0.3 (**Table 1**). Percent correct and direction range thresholds were measured

during in-lab testing, with central fixation enforced using an Eyelink 1000 eye tracker (SR

Research, Mississagua, Ontario, Canada). Tracking was binocular for all participants except for

CB3, who was tested monocularly because she exhibited convergence issues. As such, she had

her dominant (right) eye tracked and the non-dominant eye patched both for motion perception

and PFR testing. Stimuli were presented in a gaze-contingent manner in either intact or blind

regions of the visual field. Viewing distance to a luminance-calibrated CRT monitor (HP 7217

A, 48.5 x 31.5 cm, 1024x640p, refresh rate 120 Hz) was 42 cm, enforced by a chin/forehead rest.

Experiments were conducted using MATLAB (The MathWorks, Natick, MA, USA) and the

Psychophysics toolbox (Brainard, 1997; Kleiner et al., 2007; Pelli, 1997). At the start of each

trial, subjects were asked to fixate a small target at the center of the CRT monitor. The Eyelink

1000 eye tracker was accurate to within 0.25^{o}, with a sampling frequency of 1000 Hz (nominal,

standard values as reported by SR Research). Subjects were allowed a fixation window of only $\pm$

1^{o} around the fixation spot. If gaze moved outside this window during stimulus presentation, the

trial was aborted, reshuffled and patients received a noxious auditory tone as feedback,

reminding them to improve their fixation accuracy.

Following accurate fixation of the central spot for 1000 ms, a random dot stimulus

appeared in a 5^o diameter circular aperture, at one of the pre-determined locations in the

peripheral visual field (see colored circles in **Fig. 1**; NDR thresholds in **Table 1**). Black dots

moved on a mid-grey background with a 250 ms lifetime, a speed of 10 deg/s, and with a density

of 3 dots/deg^2. Stimuli were presented for 500 ms, accompanied by a tone to indicate stimulus

onset. Dots moved globally with a variable range of directions, uniformly distributed around the

left- or rightward vectors (Das et al., 2014; Huxlin et al., 2009; Saionz et al., 2020). On each

trial, subjects were asked to report the stimulus' global direction of motion by pressing the left or

right arrow keys on a keyboard (**Fig. 2A**). Task difficulty was adjusted using an adaptive

staircase (Levitt, 1971), which increased the range of dot directions from 0^o to 360^o in 40^o steps

after each set of 3 consecutive, correct responses; direction range was decreased by one 40^o step

for every incorrect response (Das et al., 2014; Huxlin et al., 2009; Saionz et al., 2020). Auditory

feedback was provided on each trial, indicating the correctness of each response. Participants

completed 100 trials at each of the four visual field locations. For each session, we fit a Weibull

function to the data to extract the direction range for which performance reached 75 % correct.

This value was then normalized to the maximum possible range of dot directions (360^o),

generating a normalized direction range (NDR) threshold, defined as:

NDR threshold = (360^o-Weibull-fitted direction range threshold)/360^o

Apparatus & eye tracking for PFR measurements: stimuli were generated using the

Psychophysics toolbox in MATLAB 2015b on a PC computer (Intel i7 CPU, Windows 7, 8 GB

RAM, GeForce Ti graphics card). They were presented on a gamma corrected display (BenQ

X2411z LED Monitor, resolution: 1920x1080p, refresh rate: 120 Hz, gamma correction: 2.2)

which had a dynamic luminance range from 0.5 to 230 cd/m^2, at a distance of 95.25 cm in a dark

room. Brightness on the display was set to 100 % and contrast to 50 %, and additional visual

features of the monitor such as blur reduction and low blue light were turned off. Gamma

corrections were verified with measurement by a photometer. Position of the left eye was

recorded continuously in all participants except for CB3, who had her right eye tracked (see

above). Eye position was recorded at 220 Hz using an infrared eye tracker (USB-220, Arrington

Research, Scottsdale, AZ, USA). The accuracy of the Arrington Eye Tracking system was 0.25˚,

with a precision of 0.15˚ (nominal, standard values as reported by Arrington Research). To

minimize potential head movements, participants performed the task using a bite bar.

PFR stimulus and task: CB patients performed a centrally-cued saccade task towards peripheral

motion apertures (**Fig. 2B**) as previously described in visually-intact controls (Kwon et al.,

2019). In brief, and as schematically illustrated in **Fig. 2B,** trials were initiated by fixation of a

small, dark, fixation spot presented on a gray background. After a variable fixation period of

150-200 ms, a saccade cue appeared at fixation together with four dot-motion apertures in a

square configuration (colored circles, **Fig. 1**). The cue (dark bar, 1˚ in length, extending from

fixation) was used to indicate the target aperture to which the participant should saccade. The

target aperture was 5.5˚ in diameter and was consistent for all the participants. For CB1, the

target apertures were centered at (±3˚, ±5˚). For CB2-8, the target apertures were centered at

(±5˚, ±5˚). For CB9, the target apertures were centered at (±5˚, ±10˚). For CB10, the target

apertures were centered at (±8˚, ±5˚). For CB11, the target apertures were centered at (±3˚,

±10˚). There were 180 dots total in each aperture, with dot luminance set to 0.5 cd/m^2 (100 %

contrast) and dot velocity fixed at 10 deg/s. Following parameters from our previous study

(Kwon et al., 2019), a Gaussian envelope was applied to each dot-motion aperture to create a

gradient in dot contrast from the center of the aperture (sigma= 1˚).

To avoid stereotyped eye movements, we varied saccade directions across trials. Thus,

the spatially cued motion aperture could appear in the intact or blind-field of a given participant,

on any given trial. Of particular note, *the motion itself or its direction were irrelevant to the task*.

The motion within the aperture was 100 % coherent and ran along a direction that was tangential

to an imaginary line from the fixation point to the aperture. For each aperture, the motion was

selected independent of the other apertures in one of the two tangential directions relative to the

center out saccade, either clockwise or counter-clockwise relative to the screen center.

We first compared eye movements in which the peripheral motion aperture was either

present or absent upon saccade offset. Participants were instructed to make a saccade to the

peripheral aperture as quickly as possible following the movement cue. A saccadic grace period

(i.e., a maximum latency) was allowed for participants to initiate the saccade. In half the trials,

selected at random, the stimulus motion remained present in all four apertures for 300 ms

following detection of the eye landing within 3.5˚ from the center of an aperture. In the other half

of trials, the stimulus was removed as soon as the eye had been detected leaving the fixation

window, thus leaving a blank screen through the post-saccadic period. A saccade was labelled

"correct" when it fell at least 3.5˚ from the saccade target center within 90 ms of the eye leaving

the fixation window. Participants completed as many trials as possible during a 1-1.5 hour

session that sampled equally from the four locations and different stimulus conditions. During a

typical session, stroke participants correctly completed on average 75.5 ± 32.0 (1 SD) trials in

each blind-field and 75.0 ± 31.8 (1 SD) trials in each intact-field. For comparison against a set of

eight normal controls, we analyzed eye movement data from a previous study (Kwon et al.,

2019). The previous study employed a task with a subset of trials identical to the current study,

but also in another half of trials tested an alternative stimulus configuration with the four

apertures along the cardinal axes rather than the quadrants, thus a total of eight locations. During

a typical session, normal controls completed on average 59.6 ± 13.8 (1 SD) trials in each of the

eight fields.

Eye movement recordings and PFR analysis: eye position data were collected as participants

performed saccades from fixation to the peripheral target. Eye tracking and saccade detection

procedures were identical to those previously published (Kwon et al., 2019). We sub-sampled

eye position using the ViewPoint Matlab toolbox (Arrington Research) at the display refresh rate

(120 Hz) to initiate gaze-contingent task events. For offline detection of saccadic eye

movements, we used the full eye position data recorded at 220 hz and applied an automatic

procedure that detected deviations in 2D horizontal and vertical eye velocity space (Engbert &

Mergenthaler, 2006; Kwon et al., 2019). Only the trials where the saccade was labelled "correct"

were included in the PFR analysis. We then focused our analysis by time locking eye velocity

traces on intervals 200 ms prior to saccade onset and 200 ms following saccade offset. Details

for eye position filtering, smoothing, and saccade detection were as previously described (Kwon

et al., 2019). In brief, the 2D eye velocity was computed from smoothed eye position traces and

then projected onto the motion vector in the target aperture on each trial. These projected

velocity traces were then aligned to saccade onset or offset, and averaged across trials for each

participant. Notably, the stimulus motion in our task was selected such that it was tangent to the

radial direction of the saccade from fixation out to the aperture (thus either clockwise or counter-

clockwise directions of motion), as in the original study (Kwon et al, 2019). This minimizes the

influence of velocity transients from the saccade itself in measuring the PFR. We have focused

analyses on the eye velocity projected onto stimulus motion for this reason, as it provides a

measure of pre-saccadic motion integration that is less contaminated by saccade induced

velocity. To quantify the net target-related eye velocity in each trial, we used a second measure

of eye velocity that did not involve any filtering or smoothing of eye position. We computed a

vector for the PFR in units of velocity (deg/sec) as the 2D vector difference in the raw (non-

smoothed) eye position from 20 to 100 ms after saccade offset normalized by that time interval.

Excluding the first 20 ms after saccade offset from this analysis interval reduced the influence of

saccade related effects to instead focus on post-saccadic smooth movements. Like velocity

traces, we projected this 2D vector onto the vector of the target's motion to produce a single

velocity value along the axis of stimulus motion, which we term the 'open-loop' PFR (Kwon et

al., 2019). To assess the average PFR across trials, we computed each CB patients' eye

movements relative to the target motion direction so that positive average eye velocities meant

that the eye was moving along the target motion direction, and negative average eye velocities

meant that the eye was moving opposite to the target motion direction.

Finally, we considered to what extent the post-saccadic following response tracked target

velocity by quantifying the PFR gain: the eye velocity computed from the open-loop PFR

normalized to the target velocity, with +1 indicating a perfect match of eye velocity to the target

motion, and negative values indicating eye velocity in the opposite direction.

Statistics: to evaluate the significance of PFR gain we computed the one-sample t-test to verify

it was greater than zero and we also computed the two-sample t-test to compare whether the PFR

gains differed between conditions comparing either intact *versus* blind-fields for CB participants,

or intact-fields for CB participants *versus* normal controls. We used the Pearson correlation to

assess the relationship between the PFR gains and NDR thresholds within each stroke patient for

intact and blind-field visual locations. Confidence intervals of 95% were computed by

bootstrapping the respective data at 10,000 times. Further statistical significance was computed

by calculating the Bayes factor for the respective t-tests and ANOVA.

References

Abel, L. A., Troost, B. T., & Dell'osso, L. F. (1983, 1983/01/01/). The effects of age on normal saccadic characteristics and their variability. *Vision research, 23*(1), 33-37. https://doi.org/https://doi.org/10.1016/0042-6989(83)90038-X

Azzopardi, P., & Cowey, A. (2001). Motion discrimination in cortically blind patients. *Brain, 124*(1), 30-46.

Bakst, L., Fleuriet, J., & Mustari, M. J. (2017). FEFsem neuronal response during combined volitional and reflexive pursuit. *Journal of vision, 17*(5), 13, 1-14.

Born, R. T., & Bradley, D. C. (2005). Structure and function of visual area MT. *Annu. Rev. Neurosci., 28*, 157-189.

Brainard, D. H. (1997, 01 Jan. 1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433. https://doi.org/https://doi.org/10.1163/156856897X00357

Britten, K. H., Shadlen, M. N., Newsome, W. T., & Movshon, J. A. (1992). The analysis of visual motion: a comparison of neuronal and psychophysical performance. *Journal of Neuroscience, 12*(12), 4745-4765.

Buizza, A., & Schmid, R. (1986). Velocity characteristics of smooth pursuit eye movements to different patterns of target motion. *Experimental brain research, 63*(2), 395-401.

Campion, J., Latto, R., & Smith, Y. M. (1983). Is blindsight an effect of scattered light, spared cortex, and near-threshold vision? *Behavioral and Brain Sciences, 6*(3), 423-486. https://doi.org/10.1017/S0140525X00016861

Cavanaugh, M. R., Barbot, A., Carrasco, M., & Huxlin, K. R. (2019, 2019/05/01/). Feature-based attention potentiates recovery of fine direction discrimination in cortically blind patients. *Neuropsychologia, 128*, 315-324. https://doi.org/https://doi.org/10.1016/j.neuropsychologia.2017.12.010

Cavanaugh, M. R., & Huxlin, K. R. (2017). Visual discrimination training improves Humphrey
     perimetry in chronic cortically induced blindness. *Neurology, 88*(19), 1856-1864.
     https://doi.org/10.1212/wnl.0000000000003921

Cavanaugh, M. R., Tadin, D., Carrasco, M., & Huxlin, K. R. (2022). Benefits of Endogenous
     Spatial Attention During Visual Double-Training in Cortically-Blinded Fields. *Frontiers
     in Neuroscience*, 421.

Cavanaugh, M. R., Zhang, R., Melnick, M. D., Das, A., Roberts, M., Tadin, D., Carrasco, M., &
     Huxlin, K. R. (2015). Visual recovery in cortical blindness is limited by high internal
     noise. *Journal of vision, 15*(10), 9-9. https://doi.org/10.1167/15.10.9

Das, A., & Huxlin, K. R. (2010). New Approaches to Visual Rehabilitation for Cortical
     Blindness: Outcomes and Putative Mechanisms. *The Neuroscientist, 16*(4), 374-387.
     https://doi.org/10.1177/1073858409356112

Das, A., Tadin, D., & Huxlin, K. R. (2014). Beyond Blindsight: Properties of Visual Relearning
     in Cortically Blind Fields. *The Journal of Neuroscience, 34*(35), 11652-11664.
     https://doi.org/10.1523/jneurosci.1076-14.2014

De Valois, R. L., & De Valois, K. K. (1991). Vernier acuity with stationary moving Gabors.
     Vision research, 31(9), 1619-1626.

Deubel, H., & Schneider, W. X. (1996). Saccade target selection and object recognition:
     Evidence for a common attentional mechanism. *Vision research, 36*(12), 1827-1838.

Elshout, J. A., van Asten, F., Hoyng, C. B., Bergsma, D. P., & van den Berg, A. V. (2016).
     Visual rehabilitation in chronic cerebral blindness: a randomized controlled crossover
     study. *Frontiers in neurology, 7*, 92.

Engbert, R., & Mergenthaler, K. (2006). Microsaccades are triggered by low retinal image slip.
Proceedings of the National Academy of Sciences, 103(18), 7192-7197.
https://doi.org/10.1073/pnas.0509557103

Ferrera, V. P., & Lisberger, S. G. (1997). Neuronal Responses in Visual Areas MT and MST
During Smooth Pursuit Target Selection. *Journal of Neurophysiology, 78*(3), 1433-1446.
https://doi.org/10.1152/jn.1997.78.3.1433

Fine, I., Anderson, C. M., Boynton, G. M., & Dobkins, K. R. (2004, 2004/04/01/). The
invariance of directional tuning with contrast and coherence. *Vision research, 44*(9), 903-
913. https://doi.org/https://doi.org/10.1016/j.visres.2003.11.022

Gardner, J. L., & Lisberger, S. G. (2001). Linked target selection for saccadic and smooth pursuit
eye movements. *Journal of Neuroscience, 21*(6), 2075-2084.

Gegenfurtner, K. R., Xing, D., Scott, B. H., & Hawken, M. J. (2003). A comparison of pursuit
eye movement and perceptual performance in speed discrimination. *Journal of vision,*
3(11), 865-876. https://doi.org/10.1167/3.11.19

Gellman, R., Carl, J., & Miles, F. (1990). Short latency ocular-following responses in man. *Vis*
Neurosci, 5(2), 107-122.

Glasser, D. M., & Tadin, D. (2014). Modularity in the motion system: Independent oculomotor
and perceptual processing of brief moving stimuli. *Journal of vision, 14*(3):28, 1-13.
https://doi.org/10.1167/14.3.28

Glickstein, M., Cohen, J. L., Dixon, B., Gibson, A., Hollins, M., Labossiere, E., & Robinson, F.
(1980). Corticopontine visual projections in macaque monkeys. *Journal of Comparative*
Neurology, 190(2), 209-229.

Goodale, M. A., & Milner, A. D. (1992). Separate visual pathways for perception and action.
Trends in neurosciences, 15(1), 20-25.

Hagan, M. A., Chaplin, T. A., Huxlin, K. R., Rosa, M. G. P., & Lui, L. L. (2019). Altered
Sensitivity to Motion of Area MT Neurons Following Long-Term V1 Lesions. *Cerebral*
Cortex, 30(2), 451-464. https://doi.org/10.1093/cercor/bhz096

Hallett, P. E. (1978, 1978/01/01/). Primary and secondary saccades to goals defined by
instructions. *Vision research, 18*(10), 1279-1296.
https://doi.org/https://doi.org/10.1016/0042-6989(78)90218-3

Hallett, P. E., & Adams, B. D. (1980, 1980/01/01/). The predictability of saccadic latency in a
novel voluntary oculomotor task. *Vision research, 20*(4), 329-339.
https://doi.org/https://doi.org/10.1016/0042-6989(80)90019-X

Heuer, H. W., & Britten, K. H. (2002). Contrast Dependence of Response Normalization in Area
MT of the Rhesus Macaque. *Journal of Neurophysiology, 88*(6), 3398-3408.
https://doi.org/10.1152/jn.00255.2002

Huang, X., & Lisberger, S. G. (2009). Noise Correlations in Cortical Area MT and Their
Potential Impact on Trial-by-Trial Variation in the Direction and Speed of Smooth-
Pursuit Eye Movements. *Journal of Neurophysiology, 101*(6), 3012-3030.
https://doi.org/10.1152/jn.00010.2009

Huxlin, K. R., Martin, T., Kelly, K., Riley, M., Friedman, D. I., Burgin, W. S., & Hayhoe, M.
(2009). Perceptual Relearning of Complex Visual Motion after V1 Damage in Humans.
The Journal of Neuroscience, 29(13), 3981-3991. https://doi.org/10.1523/jneurosci.4882-
08.2009

Kentridge, R. W., Heywood, C. A., & Weiskrantz, L. (2004, 2004/01/01/). Spatial attention
speeds discrimination without awareness in blindsight. *Neuropsychologia, 42*(6), 831-
835. https://doi.org/https://doi.org/10.1016/j.neuropsychologia.2003.11.001

Kerkhoff, G. (1999). Restorative and compensatory therapy approaches in cerebral blindness a
review. *Restorative neurology and neuroscience, 15*, 255-271.

Kerkhoff, G. (2000). Neurovisual rehabilitation: recent developments and future directions. *Journal of Neurology, Neurosurgery & Psychiatry, 68*(6), 691-706. https://doi.org/10.1136/jnnp.68.6.691

Kerkhoff, G., Münßinger, U., Haaf, E., Eberle-Strauss, G., & Stögerer, E. (1992). Rehabilitation of homonymous scotomata in patients with postgeniculate damage of the visual system: saccadic compensation training. *Restorative neurology and neuroscience, 4*, 245-254. https://doi.org/10.3233/RNN-1992-4402

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3?

Kohn, A., & Movshon, J. A. (2003, 2003/08/14/). Neuronal Adaptation to Visual Motion in Area MT of the Macaque. *Neuron, 39*(4), 681-691. https://doi.org/https://doi.org/10.1016/S0896-6273(03)00438-0

Kosovicheva, A. A., Wolfe, B. A., & Whitney, D. (2014). Visual motion shifts saccade targets. *Attention, Perception, & Psychophysics, 76*(6), 1778-1788.

Kowler, E., Anderson, E., Dosher, B., & Blaser, E. (1995). The role of attention in the programming of saccades. *Vision research, 35*(13), 1897-1916.

Kwon, S., Rolfs, M., & Mitchell, J. F. (2019). Presaccadic motion integration drives a predictive postsaccadic following response. *Journal of vision, 19*(11), 12, 11-19.

Levitt, H. (1971). Transformed Up□Down Methods in Psychoacoustics. *The Journal of the Acoustical Society of America, 49*(2B), 467-477. https://doi.org/10.1121/1.1912375

Li, H.-H., Barbot, A., & Carrasco, M. (2016). Saccade preparation reshapes sensory tuning. *Current Biology, 26*(12), 1564-1570.

Lisberger, S., & Westbrook, L. (1985). Properties of visual inputs that initiate horizontal smooth
pursuit eye movements in monkeys. *Journal of Neuroscience, 5*(6), 1662-1673.

Lisberger, S. G., Morris, E., & Tychsen, L. (1987). Visual motion processing and sensory-motor
integration for smooth pursuit eye movements. *Annual review of neuroscience*. 10:1, 97-
129

Mannan, S. K., Pambakian, A. L. M., & Kennard, C. (2010, 2010/11/01). Compensatory
strategies following visual search training in patients with homonymous hemianopia: an
eye movement study. *Journal of Neurology, 257*(11), 1812-1821.
https://doi.org/10.1007/s00415-010-5615-3

Maunsell, J., & van Essen, D. C. (1983). The connections of the middle temporal visual area
(MT) and their relationship to a cortical hierarchy in the macaque monkey. *Journal of
Neuroscience, 3*(12), 2563-2586.

Mazzi, C., Savazzi, S., & Silvanto, J. (2019). On the "blindness" of blindsight: What is the
evidence for phenomenal awareness in the absence of primary visual cortex (V1)?
Neuropsychologia, 128, 103-108.

Melnick, M. D., Tadin, D., & Huxlin, K. R. (2016). Relearning to See in Cortical Blindness. *The
Neuroscientist, 22*(2), 199-212. https://doi.org/10.1177/1073858415621035

Miles, F., Kawano, K., & Optican, L. (1986). Short-latency ocular following responses of
monkey. I. Dependence on temporospatial properties of visual input. *Journal of
Neurophysiology, 56*(5), 1321-1354.

Moore, T., & Armstrong, K. M. (2003, 2003/01/01). Selective gating of visual signals by
microstimulation of frontal cortex. *Nature, 421*(6921), 370-373.
https://doi.org/10.1038/nature01341

Moore, T., & Fallah, M. (2004). Microstimulation of the Frontal Eye Field and Its Effects on
Covert Spatial Attention. *Journal of Neurophysiology, 91*(1), 152-162.
https://doi.org/10.1152/jn.00741.2002

Movshon, J. A., & Newsome, W. T. (1996). Visual response properties of striate cortical neurons
projecting to area MT in macaque monkeys. *Journal of Neuroscience, 16*(23), 7733-
7741.

Munoz, D. P., & Everling, S. (2004, 2004/03/01). Look away: the anti-saccade task and the
voluntary control of eye movement. *Nature Reviews Neuroscience, 5*(3), 218-228.
https://doi.org/10.1038/nrn1345

Mustari, M. J., Ono, S., & Das, V. E. (2009). Signal processing and distribution in cortical-
brainstem pathways for smooth pursuit eye movements. *Annals of the New York Academy*
of Sciences, 1164, 147-154.

Nelles, G., Esser, J., Eckstein, A., Tiede, A., Gerhard, H., & Diener, H. C. (2001, 2001/06/29/).
Compensatory visual field training for patients with hemianopia after stroke.
Neuroscience Letters, 306(3), 189-192. https://doi.org/https://doi.org/10.1016/S0304-
3940(01)01907-3

Newsome, W., Wurtz, R., Dursteler, M., & Mikami, A. (1985). Deficits in visual motion
processing following ibotenic acid lesions of the middle temporal visual area of the
macaque monkey. *The Journal of Neuroscience, 5*(3), 825-840.
https://doi.org/10.1523/jneurosci.05-03-00825.1985

Nishida, S. y., & Johnston, A. (1999). Influence of motion signals on the perceived position of
spatial pattern. *Nature, 397*(6720), 610-612.

Nuding, U., Ono, S., Mustari, M. J., Buttner, U., & Glasauer, S. (2008). A theory of the dual
pathways for smooth pursuit based on dynamic gain control. *Journal of Neurophysiology,*
99(6), 2798-2808.

Ong, Y.-H., Jacquin-Courtois, S., Gorgoraptis, N., Bays, P. M., Husain, M., & Leff, A. P. (2015).
Eye-Search: A web-based therapy that improves visual search in hemianopia. *Annals of*
Clinical and Translational Neurology, 2(1), 74-78. https://doi.org/10.1002/acn3.154

Osborne, L. C., Lisberger, S. G., & Bialek, W. (2005, 2005/09/01). A sensory source for motor
variation. *Nature, 437*(7057), 412-416. https://doi.org/10.1038/nature03961

Pambakian, A. L. M., Mannan, S. K., Hodgson, T. L., & Kennard, C. (2004). Saccadic visual
search training: a treatment for patients with homonymous hemianopia. *Journal of*
Neurology, Neurosurgery & Psychiatry, 75(10), 1443-1448.
https://doi.org/10.1136/jnnp.2003.025957

Pelli, D. G. (1997, 01 Jan. 1997). The VideoToolbox software for visual psychophysics:
transforming numbers into movies. *Spatial Vision, 10*(4), 437.
https://doi.org/https://doi.org/10.1163/156856897X00366

1003

1004 Pirozzolo, F., & Hansch, E. (1981). Oculomotor reaction time in dementia reflects degree of
1005   cerebral dysfunction. *Science, 214*(4518), 349-351.
1006   https://doi.org/10.1126/science.7280699

1007

1008 Poggel, D. A., Kasten, E., Müller-Oehring, E. M., Bunzenthal, U., & Sabel, B. A. (2006,
1009   2006/06/30/). Improving residual vision by attentional cueing in patients with brain
1010   lesions. *Brain Research, 1097*(1), 142-148.
1011   https://doi.org/https://doi.org/10.1016/j.brainres.2006.04.011

1012

1013 Pollock, A., Hazelton, C., Rowe, F. J., Jonuscheit, S., Kernohan, A., Angilley, J., Henderson, C.
1014   A., Langhorne, P., & Campbell, P. (2019). Interventions for visual field defects in people
1015   with stroke. *Cochrane Database of Systematic Reviews*(5).
1016   https://doi.org/10.1002/14651858.CD008388.pub3

1017

1018 Price, N. S. C., & Blum, J. (2014, 2014/09/26/). Motion perception correlates with volitional but
1019   not reflexive eye movements. *Neuroscience, 277*, 435-445.
1020   https://doi.org/https://doi.org/10.1016/j.neuroscience.2014.07.028

1021

1022 Ramachandran, V. S., & Anstis, S. M. (1990). Illusory displacement of equiluminous kinetic
1023   edges. *Perception, 19*(5), 611-616.

1024

1025 Rashbass, C. (1961). The relationship between saccadic and smooth tracking eye movements.
1026   *The Journal of Physiology, 159*(2), 326-338.

1027

1028 Recanzone, G. H., & Wurtz, R. H. (2000). Effects of Attention on MT and MST Neuronal
1029   Activity During Pursuit Initiation. *Journal of Neurophysiology, 83*(2), 777-790.
1030   https://doi.org/10.1152/jn.2000.83.2.777

1031

1032 Rodman, H. R., Gross, C. G., & Albright, T. D. (1989). Afferent basis of visual response
1033   properties in area MT of the macaque. I. Effects of striate cortex removal. *Journal of
1034   Neuroscience, 9*(6), 2033-2050.

1035

1036 Rolfs, M., & Carrasco, M. (2012). Rapid simultaneous enhancement of visual sensitivity and
1037   perceived contrast during saccade preparation. *Journal of Neuroscience, 32*(40), 13744-
1038   13752a.

1039

1040 Rolfs, M., Jonikaitis, D., Deubel, H., & Cavanagh, P. (2011). Predictive remapping of attention
1041   across eye movements. *Nature neuroscience, 14*(2), 252-256.

1042

1043 Roth, T., Sokolov, A. N., Messias, A., Roth, P., Weller, M., & Trauzettel-Klosinski, S. (2009).
1044   Comparing explorative saccade and flicker training in hemianopia. *A randomized
1045   controlled study, 72*(4), 324-331. https://doi.org/10.1212/01.wnl.0000341276.65721.f2

1046

1047 Sahraie, A., Smania, N., & Zihl, J. (2016, 2016/09/15). Use of NeuroEyeCoach™ to Improve
1048   Eye Movement Efficacy in Patients with Homonymous Visual Field Loss. *BioMed
1049   Research International, 2016*, 5186461. https://doi.org/10.1155/2016/5186461

1050

1051 Sahraie, A., Trevethan, C. T., & MacLeod, M. J. (2008). Temporal properties of spatial channel
1052   of processing in hemianopia. *Neuropsychologia, 46*(3), 879-885.
1053   https://doi.org/10.1016/j.neuropsychologia.2007.11.008

1054

1055 Sahraie, A., Trevethan, C. T., MacLeod, M. J., Murray, A. D., Olson, J. A., & Weiskrantz, L.
1056   (2006). Increased sensitivity after repeated stimulation of residual spatial channels in
1057   blindsight. *Proceedings of the National Academy of Sciences, 103*(40), 14971-14976.

1058

1059 Saionz, E. L., Tadin, D., Melnick, M. D., & Huxlin, K. R. (2020). Functional preservation and
1060   enhanced capacity for visual restoration in subacute occipital stroke. *Brain, 143*(6), 1857-
1061   1872. https://doi.org/10.1093/brain/awaa128

1062

1063 Salzman, C. D., Britten, K. H., & Newsome, W. T. (1990, 1990/07/01). Cortical
1064   microstimulation influences perceptual judgements of motion direction. *Nature,*
1065   *346*(6280), 174-177. https://doi.org/10.1038/346174a0

1066

1067 Sanchez-Lopez, J., Pedersini, C. A., Di Russo, F., Cardobi, N., Fonte, C., Varalta, V., Prior, M.,
1068   Smania, N., Savazzi, S., & Marzi, C. A. (2019). Visually evoked responses from the blind
1069   field of hemianopic patients. *Neuropsychologia, 128*, 127-139.

1070

1071 Schafer, R. J., & Moore, T. (2007, 2007/11/08/). Attention Governs Action in the Primate
1072   Frontal Eye Field. *Neuron, 56*(3), 541-551.
1073   https://doi.org/https://doi.org/10.1016/j.neuron.2007.09.029

1074

1075 Sclar, G., Maunsell, J. H. R., & Lennie, P. (1990, 1990/01/01/). Coding of image contrast in
1076   central visual pathways of the macaque monkey. *Vision research, 30*(1), 1-10.
1077   https://doi.org/https://doi.org/10.1016/0042-6989(90)90123-3

1078

1079 Siegel, R., & Andersen, R. (1986). Motion perceptual deficits following ibotenic acid lesions of
1080   the middle temporal area (MT) in the behaving rhesus monkey. Soc Neurosci Abstr, Vol.
1081   12, 1183.

1082

1083 Simoncini, C., Perrinet, L. U., Montagnini, A., Mamassian, P., & Masson, G. S. (2012,
1084   2012/11/01). More is not always better: adaptive gain control explains dissociation
1085   between perception and action. *Nature neuroscience, 15*(11), 1596-1603.
1086   https://doi.org/10.1038/nn.3229

1087

1088   Sincich, L. C., Park, K. F., Wohlgemuth, M. J., & Horton, J. C. (2004). Bypassing V1: a direct
1089   geniculate input to area MT. *Nature neuroscience, 7*(10), 1123-1128.

1090

1091   Smith, J. L. (1962). Homonymous Hemianopia*: A review of one hundred cases. *American
1092   journal of ophthalmology, 54*(4), 616-623.

1093

1094   Souto, D., & Kerzel, D. (2014). Ocular tracking responses to background motion gated by
1095   feature-based attention. *Journal of Neurophysiology, 112*(5), 1074-1081.
1096   https://doi.org/10.1152/jn.00810.2013

1097

1098   Spering, M., & Carrasco, M. (2012). Similar effects of feature-based attention on motion
1099   perception and pursuit eye movements at different levels of awareness. *Journal of
1100   Neuroscience, 32*(22), 7594-7601.

1101

1102   Spering, M., & Gegenfurtner, K. R. (2007). Contrast and assimilation in motion perception and
1103   smooth pursuit eye movements. *Journal of Neurophysiology, 98*(3), 1355-1363.

1104

1105   Spering, M., Kerzel, D., Braun, D. I., Hawken, M. J., & Gegenfurtner, K. R. (2005). Effects of
1106   contrast on smooth pursuit eye movements. *Journal of vision, 5*(5), 455-465
1107   https://doi.org/10.1167/5.5.6

1108

1109   Spering, M., Pomplun, M., & Carrasco, M. (2011). Tracking without perceiving: a dissociation
1110   between eye movements and motion perception. *Psychological science, 22*(2), 216-225.

1111

1112   Spitzyna, G. A., Wise, R. J. S., McDonald, S. A., Plant, G. T., Kidd, D., Crewes, H., & Leff, A.
1113   P. (2007). Optokinetic therapy improves text reading in patients with hemianopic alexia.
1114   *A controlled trial, 68*(22), 1922-1930.
1115   https://doi.org/10.1212/01.wnl.0000264002.30134.2a

1116

1117   Spooner, J. W., Sakala, S. M., & Baloh, R. W. (1980). Effect of Aging on Eye Tracking.
1118   *Archives of Neurology, 37*(9), 575-576.
1119   https://doi.org/10.1001/archneur.1980.00500580071012

1120

1121   Stone, L. S., & Krauzlis, R. J. (2003). Shared motion signals for human perceptual decisions and
1122   oculomotor actions. *Journal of vision, 3*(11), 725-736. https://doi.org/10.1167/3.11.7

1123

1124   Takemura, A., Murata, Y., Kawano, K., & Miles, F. A. (2007). Deficits in Short-Latency
1125   Tracking Eye Movements after Chemical Lesions in Monkey Cortical Areas MT and
1126   MST. *The Journal of Neuroscience, 27*(3), 529-541.
1127   https://doi.org/10.1523/jneurosci.3455-06.2007

1128

1129   Tamietto, M., & Morrone, M. C. (2016). Visual plasticity: blindsight bridges anatomy and
1130   function in the visual system. *Current Biology, 26*(2), R70-R73.

1131

1132   Teuber, H.-L., Battersby, W. S., & Bender, M. B. (1960). Visual field defects after penetrating
1133   missile wounds of the brain. *Harvard University Press.*

1134

1135   Tong, F. (2003, 2003/03/01). Primary visual cortex and visual awareness. *Nature Reviews
1136   Neuroscience, 4*(3), 219-229. https://doi.org/10.1038/nrn1055

1137

1138   Trauzettel-Klosinski, S. (2010). Rehabilitation for Visual Disorders. *Journal of Neuro-
1139   Ophthalmology, 30*(1), 73-84. https://doi.org/10.1097/WNO.0b013e3181ce7e8f

1140

1141   Treue, S., & Maunsell, J. H. R. (1996, 1996/08/01). Attentional modulation of visual motion
1142   processing in cortical areas MT and MST. *Nature, 382*(6591), 539-541.
1143   https://doi.org/10.1038/382539a0

1144

1145   Treue, S., & Trujillo, J. C. M. (1999, 1999/06/01). Feature-based attention influences motion
1146   processing gain in macaque visual cortex. *Nature, 399*(6736), 575-579.
1147   https://doi.org/10.1038/21176

1148

1149   Tychsen, L., & Lisberger, S. G. (1986). Visual motion processing for the initiation of smooth-
1150   pursuit eye movements in humans. *Journal of Neurophysiology, 56*(4), 953-968.

1151

1152   Ungerleider, L. G., Desimone, R., Galkin, T. W., & Mishkin, M. (1984). Subcortical projections
1153   of area MT in the macaque. *Journal of Comparative Neurology, 223*(3), 368-386.

1154

1155   Vaina, L. M., Soloviev, S., Calabro, F. J., Buonanno, F., Passingham, R., & Cowey, A. (2014,
1156   Jun). Reorganization of retinotopic maps after occipital lobe infarction. *J Cogn Neurosci,*
1157   *26*(6), 1266-1282. https://doi.org/10.1162/jocn_a_00538

1158

1159   Van Essen, D., Maunsell, J., & Bixby, J. (1981). The middle temporal visual area in the
1160   macaque: myeloarchitecture, connections, functional properties and topographic
1161   organization. *Journal of Comparative Neurology, 199*(3), 293-326.

1162

1163   Weinberg, J., Diller, L., Gordon, W. A., Gerstman, L. J., Lieberman, A., Lakin, P., Hodges, G.,
1164   & Ezrachi, O. (1977, Nov). Visual scanning training effect on reading-related tasks in
1165   acquired right brain damage. *Arch Phys Med Rehabil, 58*(11), 479-486.

1166

1167   Weiskrantz, L. (1996). Blindsight revisited. *Current opinion in neurobiology, 6*(2), 215-220.

1168

1169   Weiskrantz, L. (2009). Blindsight: A case study spanning 35 years and new developments.
1170   *Oxford University Press.*

1171

1172   Weiskrantz, L., Barbur, J. L., & Sahraie, A. (1995). Parameters affecting conscious versus
1173   unconscious visual discrimination with damage to the visual cortex (V1). *Proceedings of*
1174   *the National Academy of Sciences of the United States of America, 92*(13), 6122-6126.
1175   https://doi.org/10.1073/pnas.92.13.6122

1176

1177   Weiskrantz, L., Warrington, E. K., Sanders, M., & Marshall, J. (1974). Visual capacity in the
1178   hemianopic field following a restricted occipital ablation. *Brain, 97*(1), 709-728.

1179

1180 White, A. L., Rolfs, M., & Carrasco, M. (2013). Adaptive deployment of spatial and feature-
1181   based attention before saccades. *Vision research, 85*, 26-35.

1182

1183 Whitney, D., & Cavanagh, P. (2000). Motion distorts visual space: shifting the perceived
1184   position of remote stationary objects. *Nature neuroscience, 3*(9), 954-959.

1185

1186 Yoshida, M., Takaura, K., Kato, R., Ikeda, T., & Isa, T. (2008). Striate Cortical Lesions Affect
1187   Deliberate Decision and Control of Saccade: Implication for Blindsight. *The Journal of*
1188   *Neuroscience, 28*(42), 10517-10530. https://doi.org/10.1523/jneurosci.1973-08.2008

1189

1190 Zihl, J. (1980, 1980/01/01/). "Blindsight": Improvement of visually guided eye movements by
1191   systematic practice in patients with cerebral blindness. *Neuropsychologia, 18*(1), 71-78.
1192   https://doi.org/https://doi.org/10.1016/0028-3932(80)90085-8

1193

1194

1195

Acknowledgments

The authors wish to thank Martin Rolfs, Duje Tadin, Ralf Haefner, and Gregory DeAngelis for enlightening discussions and comments on the manuscript. The authors also wish to thank Terrance Schaefer, who performed the Humphrey visual field tests presented here and Christine Callan for her excellent work as research coordinator.

Author Contributions

SK, KRH, and JFM designed the study; SK collected and analyzed the data with inputs from KRH and JFM; BKF and MRC collected perceptual performance data with inputs from KRH; SK, KRH and JFM wrote the manuscript and all authors commented on it.

Funding

The present study was funded by NIH (EY027314 and EY021209 to KRH; EY030998 to JFM; P30 EY001319 to the Center for Visual Science), by start-up funds to JFM from the University of Rochester, and by an unrestricted grant from the Research to Prevent Blindness (RPB) Foundation to the Flaum Eye Institute.

Competing interests

KRH is co-inventor on US Patent No. 7,549,743. The remaining authors have no competing interests.