Delayed induction of type I and III interferons mediates nasal epithelial cell permissiveness

to SARS-CoV-2

Nasal interferon responses to SARS-CoV-2

Authors

Catherine F Hatton[1a], Rachel A Botting[2a], Maria Emilia Dueñas[2a], Iram J Haq[1,3a], Bernard Verdon[2a],

Benjamin J Thompson[1], Jarmila Stremenova Spegarova[1], Florian Gothe[1,4], Emily Stephenson[2], Aaron I

Gardner[1], Sandra Murphy[2], Jonathan Scott[1], James P Garnett[1], Sean Carrie[5], Jason Powell[1], C M Anjam

Khan[2], Lei Huang[1], Rafiqul Hussain[6], Jonathan Coxhead[6], Tracey Davey[7], A John Simpson[1], Muzlifah

Haniffa[2,8,9,10], Sophie Hambleton[1,11], Malcolm Brodlie[1,3b], Chris Ward[1b], Matthias Trost[2b], Gary

Reynolds[2b], Christopher J A Duncan[1,12 b]∗

[a] equal contribution

[b] equal contribution

* corresponding author: christopher.duncan@ncl.ac.uk

Affiliations

1. Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University,

Newcastle upon Tyne, UK.

2. Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK.

3. Paediatric Respiratory Medicine, Great North Children's Hospital, Newcastle upon Tyne Hospitals

NHS Foundation Trust, Newcastle upon Tyne, UK.

4. Department of Pediatrics, Dr. von Hauner Children's Hospital, University Hospital, Ludwig-

Maximilians-Universität Munich, Munich, Germany.

5. Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK.

6. Genomics Core Facility, Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK.

7. Electron Microscopy Research Services, Newcastle University, Newcastle upon Tyne, UK.

8. NIHR Newcastle Biomedical Research Centre, Newcastle Hospitals NHS Foundation Trust, Newcastle

upon Tyne, UK

9. Department of Dermatology, Newcastle Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

10. Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, UK

11. Great North Children's Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, UK.

12. Department of Infection and Tropical Medicine, Newcastle upon Tyne Hospitals NHS Foundation

Trust, UK.

Abstract

The nasal epithelium is a plausible entry point for SARS-CoV-2, a site of pathogenesis and transmission,

and may initiate the host response to SARS-CoV-2. Antiviral interferon (IFN) responses are critical to

outcome of SARS-CoV-2. Yet little is known about the interaction between SARS-CoV-2 and innate

immunity in this tissue. Here we applied single-cell RNA sequencing and proteomics to a primary cell

model of human nasal epithelium differentiated at air-liquid interface. SARS-CoV-2 demonstrated

widespread tropism for nasal epithelial cell types. The host response was dominated by type I and III

IFNs and interferon-stimulated gene products. This response was notably delayed in onset relative to

viral gene expression and compared to other respiratory viruses. Nevertheless, once established, the

paracrine IFN response began to impact on SARS-CoV-2 replication. When provided prior to infection,

recombinant IFNβ or IFNλ1 induced an efficient antiviral state that potently restricted SARS-CoV-2

viral replication, preserving epithelial barrier integrity. These data suggest that the IFN-I/III response

to SARS-CoV-2 initiates in the nasal airway and suggest nasal delivery of recombinant IFNs to be a

potential chemoprophylactic strategy.

INTRODUCTION

SARS-CoV-2 is an emergent betacoronavirus responsible for coronavirus disease-19 (COVID-19)[1]. Since its identification in late 2019, global pandemic transmission of SARS-CoV-2 has resulted in over 192 million confirmed infections and approximately 4.1 million deaths. SARS-CoV-2 infects target cells via the entry receptor ACE2[2] leading to a spectrum of clinical outcomes, ranging from asymptomatic infection to death[3]. Although multiple host factors (e.g. age, male sex, obesity) contribute to adverse clinical outcome[4], the immune response also plays a decisive role, evidenced by the therapeutic benefit of immunomodulatory agents including corticosteroids[5] or IL6 inhibition[6]. Yet much remains to be understood about the immunopathogenesis of COVID-19. Identification of the cells hosting viral entry and characterisation of their response to infection is essential to understanding pathogenesis and improving therapy.

The nasal epithelium is believed to be a key entry point of SARS-CoV-2. Nasal epithelial tropism and efficient viral shedding from the nasopharynx apparently contributes to the high transmissibility of SARS-CoV-2[7], as well as to pathologic features such as anosmia[8]. As an early viral target cell, nasal epithelial cells may also set the tone for the systemic immune response, potentially influencing disease outcome[9]. These factors emphasise the need to study host-virus interaction in human nasal cells. *Ex vivo* single-cell transcriptomic studies indicate that ciliated and/or goblet cells in the nasal mucosa express *ACE2* and *TMPRSS2,* implicating them as probable SARS-CoV-2 target cells[10,11]. This has been confirmed by *in vitro* and *in vivo* studies demonstrating SARS-CoV-2 infection of human nasal epithelial cells[12-15]. Single-cell studies also revealed that nasal cells exhibit basal expression of an antiviral expression programme, characterised by induction of several interferon-stimulated genes (ISGs), suggesting that they may be primed to respond to viral infection[10,11]. Interestingly, *ACE2* is also regulated by interferons (IFNs)[11,16], implying a complex relationship between IFN signalling and tropism. Type I and type III IFN (IFN-I/III) systems play a critical role in human antiviral innate immunity[17] and have been recently implicated in defence against SARS-CoV-2 susceptibility to severe or life-threatening COVID-19 is associated with deleterious variants in *IFNAR* genes[18,19] and IFN-I

blocking autoantibodies[20]. *In vitro*, SARS-CoV-2 appears sensitive to the antiviral properties of IFN-I,

at least in cell lines[21,22], and this activity extends to *in vivo* model systems[9]. These findings motivate

studies to improve understanding of the interaction between SARS-CoV-2 and the IFN-I system in

primary human target cells, providing impetus to clinical trials of recombinant IFNs in treatment or

prophylaxis of COVID-19[23].

Organotypic cultures of primary human nasal epithelium differentiated at air-liquid interface (ALI) are

a translationally relevant primary cell model for studies of SARS-CoV-2 host-virus interaction[12], with

considerable potential to accelerate our understanding of pathogenesis. A small number of studies

using this model demonstrate that SARS-CoV-2 replicates efficiently in human nasal cells[12-14], yet

important questions concerning cellular tropism and their innate immune response remain

unresolved. Hou and colleagues report that only ciliated cells were permissive to SARS-CoV-2, despite

expression of ACE2 and TMPRSS2 by all cell types[13]. They hypothesised that post-entry factors, such

as innate immunity, might govern tropism. By contrast, Pizzorno and colleagues reported infection in

all major cell types (ciliated, secretory and basal cells)[14], consistent with prior indications from single-

cell RNA sequencing (scRNA-seq) data and studies in lower airway models[24,25]. While an IFN response

to SARS-CoV-2 can be detected in nasal cells[12,14], in apparent contrast to bronchial or alveolar

epithelial cells[26-28], the kinetics of induction and the antiviral function of IFNs in nasal epithelium has

not been systematically characterised.

Here we employed a comprehensive range of techniques, including scRNA-seq and proteomics, in

primary human nasal ALI cultures to define: (i) cellular tropism; (ii) the innate immune response to

SARS-CoV-2; and (iii) the antiviral activity of IFN-I/III. We observed broad cellular tropism of SARS-CoV-

2 for nasal epithelial cells, although secretory and ciliated cells were the most permissive. Nasal cells

mounted a delayed IFN response that began to exert control over viral replication at later times post-

infection. However, SARS-CoV-2 remained highly sensitive to IFN-restriction if exogenous IFN-I/III was

applied prior to infection. These data enrich our understanding of the interaction of SARS-CoV-2 and

the human IFN system at the earliest point of infection, with immediate therapeutic implications.

RESULTS

SARS-CoV-2 robustly infects primary differentiated nasal epithelial cultures

Primary nasal epithelial cultures were established from cryopreserved stocks from six adult donors, obtained prior to the SARS-CoV-2 pandemic. Cells were expanded, differentiated and then matured at ALI for 28 days, according to an established protocol[29]. We first sought to address their suitability as a model of SARS-CoV-2 infection. Single-cell RNA-seq libraries were generated from two representative donors, yielding 28,346 individual transcriptomes for analysis following quality control (Fig. S1, Table S1). Following dimensionality reduction and Leiden clustering, eight populations were discerned by their expression of established marker genes[11,30] (Fig. 1A, supplementary dataset 1). This annotation was further validated using Seurat label transfer from a published scRNA-seq dataset from nasopharyngeal swabs[15] (Fig. 1B). The major populations identified were ciliated, secretory, goblet and basal cells, alongside two rarer populations of FOXN4+ deuterosomal cells[31] and ionocytes. Cells expressed characteristic markers (Fig. 1C and supplementary dataset 1), corresponding closely to *ex vivo* single-cell data from nasal brusings[15] (Fig. 1B). Immunostaining verified the presence of major cell types in these cultures using well-established protein markers[13] - including acetylated alpha-tubulin-positive (AAT) ciliated cells, mucin 5B-positive (MUC5B) secretory cells, mucin 5AC (MUC5AC)-positive goblet cells, and tumour protein 63-positive (TP63) basal cells (Fig. S2) - with ciliated cells the most abundant cell population. Consistent with published scRNA-seq data[10,11,15], mRNA for key SARS-CoV-2 entry receptors, *ACE2* and *TMPRSS2*, was expressed, albeit at relatively low levels, alongside other genes implicated in SARS-CoV-2 entry such as *FURIN* and *CTSL* (Fig. S3)[32]. Robust expression of ACE2 and TMPRSS2 at the protein level was confirmed by immunoblotting of whole-cell lysates prepared from mature ALI cultures (Fig. 1D). To establish their permissiveness to infection, nasal ALI cultures were inoculated at the apical surface with a clinical SARS-CoV-2 isolate (BetaCoV/England/2/20) at an approximate multiplicity of infection (MOI) 0.1 - consistent with other studies (0.1-0.5)[12-14] - and monitored for infection over the next 72 h. Expression of SARS-CoV-2 *nucleocapsid (N)* gene and spike

(S) protein increased significantly over time, indicative of viral replication (Fig. 1E-F). This was

accompanied by the release of infectious viral particles, as determined by plaque assay of apical

washes on vero E6 cells, confirming productive infection (Fig. 1G). SARS-CoV-2 replication was

accompanied by a progressive decline in epithelial barrier integrity starting from 48 hours post-

infection (hpi), reflecting virus-induced epithelial dysfunction and/or potential cytopathic effect (Fig.

1H). These data established the suitability of the human nasal ALI system for modelling SARS-CoV-2

infection.

Figure 1. **Robust SARS-CoV-2 infection in a primary differentiated nasal epithelial ALI culture model.**

(A) UMAP visualisation of scRNA-seq data from nasal ALI cultures (28,346 single-cell transcriptomes

from two representative donors) revealed six major cell types. (B) Correlation between the annotation

from an external dataset of nasopharyngeal swabs and the assigned annotation of our scRNA-seq from

nasal ALI culture following label transfer. (C) Dot plot demonstrating expression of key markers

distinguishing cell types in annotated clusters, with intensity demonstrated by colour and size of the

dot representing the proportion of cells expressing the marker. (D) Immunoblot demonstrating ACE2

and TMPRSS2 expression by donor, representative of n=3 experiments. Nasal ALI cultures were

infected with SARS-CoV-2 (MOI 0.1) and subjected to various modalities to analyse infection. Whole-

cell lysates were prepared at the indicated times for RT-PCR analysis of expression of (E) SARS-CoV-2

nucleocapsid (*N*) gene expression normalised to the housekeeper *RNASEP2* (average of n=2 repeat

experiments in n=4 donors, mean $\pm$ SEM; * P < 0.05, **** P < 0.0001, ANOVA with Dunnett's post-test

correction compared to 0h). (F) Whole-cell lysates were prepared at the indicated times for

immunoblot analysis of expression of SARS-CoV-2 spike (S) and cleaved S2 protein expression

(representative of repeat experiments in n=4 donors). (G) Release of infectious viral particles was

determined by plaque assay of apical washings on permissive vero E6 cells (average of repeat

experiments in n=6 donors, mean $\pm$ SEM; **** P < 0.0001, ANOVA with Dunnett's post-test correction

compared to 24h). Dotted line represents lower limit of detection. (H) Transepithelial resistance

(TEER) measurements upon infection (expressed as % of mock-infected wells, n=6 donors, mean $\pm$

SEM; *** P < 0.001, **** P < 0.0001, ANOVA with Dunnett's post-test correction compared to 24h).

Evidence of broad cellular tropism of SARS-CoV-2

To revisit the question of whether individual cell types are more or less permissive to SARS-CoV-2[13,14],

we first examined viral gene expression by scRNA-seq analysis at 24 hpi, selected to represent an early

stage in the progress of infection. While all cell types expressed viral transcripts, there were notable

differences both in the proportion of cells infected, and the relative abundance of different viral

transcripts within these cells (Fig. 2A). Based on differential gene expression analysis between cell

types (Wilcoxon rank sum test, one vs rest, $P < 0.05$), secretory and ciliated cells expressed higher

levels of viral transcripts than other cell types, with viral transcripts most abundant in secretory cells

(Fig. 2A, supplementary dataset 2). Deuterosomal cells also expressed abundant viral transcripts but

were a rare population within these cultures, possibly limiting the power of this analysis. Basal cells

are also located away from the apical surface; physical inaccessibility to apically-applied virus at this

time point might at least partially account for this observation. To investigate tropism further, we

undertook immunofluorescence analysis of viral spike (S) protein expression at 48 hpi (Fig. 2C-D, see

Fig. S4 for background spike immunoreactivity in uninfected cells). This analysis revealed broadly

similar proportions of ciliated, secretory and basal cells expressing S protein, with significantly lower

spike immunoreactivity in MUC5AC positive (goblet) cells (Fig. 2D). However, the mean pixel intensity

of S protein was significantly greater in ciliated cells than in other cell types, and significantly increased

in secretory cells compared to basal cells (Fig. 2D). To corroborate these findings, we undertook

analysis of intracellular virion-like structures (VLSs) at 48 hpi by transmission electron microscopy

(TEM, Fig 2E), focusing on ciliated and secretory/goblet cells (the latter cell types were grouped for

analysis as they could not be reliably distinguished based on morphology). Intracellular VLSs were

observed in both ciliated and secretory/goblet cells, predominantly towards the apical surface (Fig.

2E). Consistent with immunofluorescence analysis of S protein intensity, there was a significant

increase in the number of VLSs per cell in ciliated compared with secretory/goblet cells (Fig. 2E).

Collectively, these data suggested that the virus is capable of entering, and replicating in, all major

nasal cell types, but with quantitative differences in efficiency.

Figure 2. Broad tropism of SARS-CoV-2 for nasal cells. Nasal ALI cultures were infected with SARS-

CoV-2 (MOI 0.1) and analysed using different modalities to explore tropism. At 24 hours post-infection

(hpi), cell suspensions were prepared from two representative donors for single-cell RNA sequencing

(scRNA-seq) and 28,346 individual transcriptomes passing QC were analysed. (A) Dot plot of scRNA-

seq data showing magnitude (colour) and proportion (size) of cell types expressing viral transcripts. E

= envelope; M = matrix; N = nucleocapsid; S = spike. (B) Relative proportion of infected cell types based

on expression of any viral transcript. Separately, nasal ALI cultures were fixed at 48 hpi and subjected

to immunofluorescence analysis. (C) Expression of viral S protein expression in ciliated (AAT), basal

cells (TP63), secretory (MUC5B) and goblet (MUC5AC) cells (arrowed) shown in (C). Scale bars = 10 μm

(representative of experiments in n=5 donors). (D) Quantification of cell-type specific expression of

viral S protein and S protein intensity at 48 hpi (n=5 donors, mean ± SEM; * P < 0.05, ** P < 0.01, ***

P < 0.001, **** P < 0.0001, ns = non-significant, ANOVA with Sidak's post-test correction for multiple

comparisons, indicated by lines). (E) Nasal ALI cultures were infected as above, fixed at 48 hpi for

transmission electron micrograph (TEM) analysis of SARS-CoV-2 infected ciliated and secretory/goblet

cells. Inserts (a) and (b) display virion-like structures in ciliated and secretory/goblet cells respectively.

Scale bars = 1 μm. Image analysis was undertaken to quantify virion-like structures as displayed in the

bar plot (n=3 donors, mean ± SEM ** P = 0.003, Mann-Whitney test).

Characterisation of individual nasal cell responses to SARS-CoV-2

Published *ex vivo* single-cell transcriptomic analyses report that nasal cell types exhibit the basal

expression of an innate antiviral gene signature, in the absence of viral infection, characterised by

several IFN-stimulated genes (ISGs)[10,11]. This signature correlated with *ACE2* expression, suggesting

conditioned expression to reduce susceptibility. Based on these reports, we examined scRNA-seq data

to characterise the innate antiviral response of nasal cells to SARS-CoV-2 infection at 24 hpi. In

unexposed cells, ISG signature scores were generated using context-specific ISGs from a published

IFN-treated nasal cell dataset[11] and compared using the Wilcoxon rank sum test. Gene set scores

greater than zero suggested expression levels higher than background gene expression, and was the

case for basal 1, secretory, goblet cells and ionocytes (Fig. 3A). The basal ISG signature was highest in

secretory cells despite abundant expression of viral RNA in these cells upon exposure to SARS-CoV-2

(Fig. 2A-B), suggesting basal ISG expression may not be sufficient to protect against infection. Next we

distinguished cells in three experimental conditions: unexposed (mock-infected); SARS-CoV-2-

exposed but uninfected (these 'bystander' cells would theoretically be exposed to IFNs and other

paracrine signals, but not infected); and SARS-CoV-2-infected (as defined by detectable expression of

SARS-CoV-2 transcripts). We undertook differential expression (DE) analysis between mock and

bystander or infected cells, labelling ISGs derived from the same list of context-specific ISGs (Fig. 3B,

supplementary datasets S3-4). There was minimal transcriptional response to infection in bystander

cells, including the absence of ISG induction, suggesting a lack of substantial paracrine IFN signalling

at this timepoint in keeping with reports in other airway models[25,27,28]. Interestingly, *IFITM* genes (ISGs

which have been paradoxically implicated in SARS-CoV-2 entry[33]) were downregulated in some

bystander cell populations. There was also minimal evidence of ISG induction in SARS-CoV-2 infected

cells, especially secretory, deuterosomal and goblet cells (Fig. 3B). In secretory and ciliated several

ISGs relating to antigen processing or presentation were downregulated upon infection (Fig. 3B). Basal

cells expressed a modest number of ISGs upon infection, specifically genes of the *IFITM* family, *IFI27*

and *IFI6* and the negative regulator of IFN-I signalling, *ISG15.* Consistent with this finding, gene-set

enrichment analysis (GSEA) revealed upregulation of IFN alpha/gamma responses in infected basal

cell populations but not in other cell types (Fig. 3C). The transcriptional response of infected secretory

and ciliated cells was characterised by widespread downregulation of expression, which may reflect

viral co-optation of transcriptional machinery of host cells, but this effect was not uniform. In

agreement with previous reports[34,35], genes related to oxidative phosphorylation were prominent

amongst downregulated genes, as were antigen presentation pathways. Pathway analysis also

predicted upregulation of NF-KB signalling in basal, secretory, goblet and ciliated cells, consistent with

previous findings[27,36]. Using DoRoTHea to explore regulon activity in these populations revealed higher

predicted *NFKB2* activity but limited evidence of widespread activation of IFN-mediated signalling (Fig.

S5). Transcripts for IFN-I (*IFNB, IFNK, IFNA5)* and IFN-III (*IFNL1)* were not significantly differentially

expressed and were detectable in only a small minority (~ 0.4%) of infected secretory cells (Fig. S6).

Whilst potentially consistent with the lack of paracrine signalling at this timepoint, this might also

reflect transient expression and/or insensitivity of detection by scRNA-seq; a similar pattern was

observed for other cytokines and chemokines (Fig. S6). Overall, this analysis showed that despite

evidence of NF-KB activation at 24 hpi, there was a minimal IFN response to SARS-CoV-2 in the cell

types with the highest levels of infection, consistent with previous reports in non-nasal epithelial

cells[25,27,28].

Figure 3. Characterisation of individual nasal cell transcriptional responses to SARS-CoV-2. Nasal ALI

cultures were infected with SARS-CoV-2 (MOI 0.1). At 24 hours post-infection (hpi), cell suspensions

were prepared from two representative donors for single-cell RNA sequencing (scRNA-seq) and

28,346 individual transcriptomes passing QC were analysed. (A) A composite interferon-stimulated

gene (ISG) signature score was defined based on a published nasal cell dataset from cells treated with

IFN alpha and IFN gamma. Gene set scores greater than zero suggest expression levels higher than

background gene expression. Wilcoxon rank sum testing was performed for each cell type vs all with

Benjamini-Hochberg correction (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ **** $P < 0.0001$). (B) Differential

expression (DE) analysis by Wilcoxon rank sum test was undertaken to compare mock-infected cell

transcriptomes with those from bystander cells (without detectable viral transcripts) and infected cells

(with detectable viral transcripts) from the virus-exposed cultures. Volcano plots were generated with

vertical lines marking -/+ 1.5 fold change cut-offs (note $\log_2$ scale) and the horizontal line marking an

adjusted P value cut-off of 0.05 (< 0.05 was considered statistically significant). Individual genes

coloured as non-significant (light blue) and significant (red). Labels indicate viral transcripts (dark blue)

and epithelial-cell specific ISGs (black). (C) Gene-set enrichment analysis was undertaken by ordering

genes by fold change difference between mock-infected and infected cells by cluster. Vertical lines

indicated adjusted P value cut-off of 0.05. NES = normalised enrichment score.

Kinetics of innate IFN-I/IFN-IIIs response to SARS-CoV-2

To investigate the kinetics of the IFN-I/III response, expression of IFN-I (*IFNA1* and *IFNB*) and IFN-III

(*IFNL1*) was examined by RT-PCR at 6, 24, 48 and 72 hpi at the same MOI (0.1) as previous experiments.

Induction of *IFNL1* and *IFNB* was low at 24 hpi, consistent with scRNA-seq findings, but increased

significantly by 48 and 72 hpi (Fig. 4A). *IFNA1* was not induced, as observed in our scRNA-seq data.

Compared to the timing of initiation of viral gene expression - which was detectable at 6 hpi and

approached its maximum level by 24 hpi (Fig. 1E) - the induction of IFNs appeared delayed, as

suggested by previous studies[12,14,36]. Infection was accompanied by progressive upregulation of

proinflammatory cytokines such as *IL6, IL1B* and *TNF*, consistent with initiation of an NF-KB-dependent

inflammatory response (Fig. 4B). To look for evidence of a paracrine response to IFN-I/III, we analysed

expression of the ISGs *RSAD2* and *USP18* by RT-PCR, as well as the expression of RSAD2, USP18, ISG15

and MX1 proteins by immunoblotting. There was an increase in ISG mRNA and protein expression at

later times following the onset of *IFN* gene expression (Fig. 4C-D), potentially suggestive of paracrine

JAK-STAT signalling. To explore early induction of IFNs in more detail, we compared the response to

SARS-CoV-2 with other RNA respiratory viruses, influenza A virus (IAV) and parainfluenza virus 3

(PIV3). In this experiment, significant induction of *IFNB* and *IFNL1* occurred in response to both PIV3

and IAV, but not SARS-CoV-2, at 24 hpi (Fig. S7), and was accompanied by upregulation of ISGs *USP18*

and *RSAD2*. Infection of cell lines at high MOI are reported to enhance the relatively inefficient IFN-I

induction to SARS-CoV-2[25]. To confirm that the attenuated production of *IFNL1* and *IFNB* at early times

was not dependent on MOI, we repeated SARS-CoV-2 infections at 20-fold higher MOI (2), alongside

IAV (Fig. 4E), or a preparation of Sendai virus (SeV) containing a high proportion of immunostimulatory

defective viral genomes[32] as a positive control (Fig. S8). At 6 hpi, a time point at which *IFNB* and *IFNL1*

were significantly induced by IAV, there was no detectable response to SARS-CoV-2 (Fig. 4E).

Compatible observations were made with SeV (Fig. S8). At 24 hpi, SARS-CoV-2 exposure led to no

detectable induction of *IFNB* and significantly less *IFNL1* than IAV (Fig. 4E). This differential response

was reflected in the robust expression of ISGs *RSAD2, USP18* and *ISG15* at 24 h post-inoculation with

IAV but not SARS-CoV-2 (Fig. 4F). These observations recapitulated our previous RT-PCR and scRNA-

seq data with a lower MOI, and are consistent with other reports[12,27,37]. Collectively, the results

indicate that nasal epithelial cells express IFN-I/IIIs during SARS-CoV-2 infection, but that the response

is delayed relative to viral replication.

Figure 4. Delayed induction of IFN-I/III signalling in SARS-CoV-2-infected nasal ALI cultures. Nasal ALI

cultures were infected with SARS-CoV-2 (MOI 0.1). Whole-cell lysates were prepared at the indicated

times for RT-PCR analysis of expression of (A) *IFNB*, *IFNL1* and *IFNA1* (B) *IL6*, *TNF* and *IL1B* and (C)

USP18 and *RSAD2* (average of n=2 repeat experiments in n=4 donors, mean ± SEM; * P < 0.05, ** P <

0.01, *** P < 0.001 **** P < 0.0001, ANOVA with Dunnett's post-test correction compared to 6h [B,

IL6], 24h [A, *IFNL1*] or 0h [all others]). ND, Not detected. (D) Whole-cell lysates were prepared at the

indicated times for immunoblot analysis of S/S2, MX1, USP18, RSAD2 and ISG15 expression

(representative of experiments in n=4 donors). Nasal ALI cultures were infected with SARS-CoV-2 or

influenza A virus (IAV H1N1, purple bars) at MOI 2. Whole-cell lysates were prepared at the indicated

times for RT-PCR analysis of expression of (E) *IFNB*, *IFNL1* and (F) the ISGs *USP18*, *RSAD2* and *ISG15*

(n=3 donors, mean ± SEM; * P < 0.05, ** P < 0.01, *** P < 0.001 **** P < 0.0001, ANOVA with Dunnett's

post-test correction compared to 0h).

IFN-signalling dominates the nasal host response to SARS-CoV-2 at the protein level

To validate and extend these findings, we undertook an unbiased assessment of the host response to

SARS-CoV-2 infection by proteomics analysis. Whole-cell lysates were prepared from SARS-CoV-2 and

mock-infected nasal ALI cultures from six donors at 72 hpi. Lysates were analysed by quantitative mass

spectrometry (quality control data in Fig. S9). Overall, this analysis detected the differential expression

(DE) of 180 proteins including viral proteins such as S, M, N, ORF1AB, ORF3A and ORF8 (Fig. 5A

supplementary dataset S4). The most highly increased host protein was Sorting Nexin 33 (SNX33), an

endosomal protein that has not yet been implicated in the life cycle of SARS-CoV-2. Notably, other

SNX proteins (e.g. SNX17 and SNX2) are involved in viral trafficking[38,39]. Infected and uninfected cells

clustered together by principal component analysis (Fig. 5B). Inspection of the DE proteins confirmed

a robust host innate immune response, dominated by ISG products (Fig. 5A). Functional annotation

revealed an enrichment of antiviral response and especially IFN-I signalling pathways (Fig. 5C, Table

S2). These data are consistent with our earlier findings and contrary to prior reports in cell lines or

human bronchial/tracheal epithelial cultures, where a robust endogenous IFN-I/III response to SARS-

CoV-2 was not detected[26-28]. Key antiviral ISG proteins identified included IFIT1-3, MX1-2, and the OAS

cluster (OAS1-3), the latter associated with genetic susceptibility to severe COVID-19[19] (Fig. 5C).

Significantly downregulated pathways were also identified, including TRIF-dependent toll-like

receptor signalling, as well as RNA polymerase II transcription and endosomal transport (Table S3).

This implied viral subversion of critical host functions, including host gene transcription, protein

trafficking and viral sensing. Proteins involved in the maintenance of epithelial tight junctions were

also downregulated, consistent with the loss of barrier integrity observed in earlier experiments (Fig.

1H).

340

Figure 5. An ISG response dominates the proteome of SARS-CoV-2 infected nasal ALI cultures.

Differential proteomic profiling of SARS-CoV-2-infected nasal ALI cultures. Mass spectrometry-based

proteomics was carried out on whole-cell lysates prepared at 72 hpi (n=6 donors per condition). (A)

Volcano plot illustrating 180 differentially expressed proteins with increased (orange points) and

decreased (purple points) expression in infected as compared to mock-infected samples. Dotted red

lines indicate those proteins with a fold change of > 1.5 and adjusted p values <0.05. (B) Principal

component analysis of the whole proteome data set. (C) Functional annotation network of

differentially expressed proteins.

Antiviral activity of IFN-I/III towards SARS-CoV-2 infection

Given the prominence of the IFN-I/III response in the proteome of SARS-CoV-2-infected cells at later

times post-infection, a key question was whether this IFN-I/III response had any impact on SARS-CoV-

2 replication. To address this question, nasal ALI cultures were treated with the JAK inhibitor

ruxolitinib (RUX). RUX antagonises signalling downstream of IFNAR and IFNLR, owing to the

involvement of JAK1 in both signalling pathways. We reasoned that blocking paracrine IFN-I/III

signalling would reveal its impact, if any, on SARS-CoV-2 replication. Cells were treated with 10 μM

RUX (a dose optimised in prior experiments[40]) or vehicle control (DMSO) in the basal medium for 24

hours prior to infection. Nasal cultures were infected at the apical surface (MOI 0.1), inhibitors were

refreshed every 24h and infection was monitored up to 96 hpi. Lysates were prepared and analysed

by RT-PCR and immunoblot. RUX treatment abolished expression of ISGs *USP18, RSAD2 and ISG15* at

the mRNA and protein level (Fig. 6A-B), indicating that ISG induction was dependent on paracrine IFN-

I/III signalling, as previously suggested (Fig. 4D). By 96 hpi, approximately 24 hours after ISGs were

reliably detected at the protein level (Fig. 4D, Fig. 5A-C), blockade of this endogenous IFN response by

RUX led to a significant increase in both viral gene expression, assessed by RT-PCR (*N* gene) and

immunoblot (S/S2 protein, Figs 6B-D), and apical release of infectious virus measured by plaque assay

(Fig. 6E). These data provided further evidence that SARS-CoV-2 triggered an endogenous paracrine

IFN-I/III response in nasal cells, which once established began to impact SARS-CoV-2 replication.

An important follow-up question was whether nasal cells could mount an antiviral state to SARS-CoV-

2, providing IFN-I/III was delivered in a timely fashion. To address this, nasal ALI cultures were pre-

treated with exogenous IFNβ (1000 IU/mL) or IFNλ1 (100 ng/mL) for 16h to induce an antiviral state,

subsequently infected with SARS-CoV-2 at MOI 0.01 and examined at 48 hpi. Analysis of infection by

immunoblotting of whole-cell lysates for spike (S/S2) protein expression or plaque assay of apical

washes demonstrated a significant reduction in infection with either IFNβ or IFNλ1 pre-treatment (Fig.

7A-B). This was accompanied by robust induction of antiviral ISG products (Fig. 7A), and preservation

of barrier integrity (Fig. S10). It is worth noting that the ISG expression induced in response to

recombinant IFN-I/III at 48 hpi was substantially greater than that induced by endogenous IFN-I/III

production (Fig. 7A). Thus exogenous IFN-I/III was capable of inducing in the nasal epithelium an

antiviral state that potently inhibited SARS-CoV-2 infection, providing it was delivered (a) prior to

infection, and (b) at sufficient concentration. This IFN-sensitivity of SARS-CoV-2 contrasts with the

relative resistance of SARS-CoV[21]. These data suggest that mucosal delivery of IFNβ or IFNλ1 is a

potential therapeutic strategy for SARS-CoV-2. In clinical practice, IFNs are unlikely to be used prior to

infection, unless this is part of a prophylactic regimen. To determine the effectiveness of exogenously

applied IFNs once SARS-CoV-2 infection is underway, infected cells were treated with IFNβ or IFNλ1

at 6 or 24 hpi and examined for S/S2 protein expression by immunoblot (Fig. 7C-D) and release of

infectious virus by plaque assay (Fig. 7E). In this experiment, IFNβ and IFNλ1 treatment at 6 hpi

continued to impact on SARS-CoV-2 infection, whereas addition after 24 hpi had minimal effect.

Interestingly, ISG induction was still observed in response to IFN treatment at 24 hpi, albeit at reduced

magnitude in the case of IFNβ (Fig. 7C-D). These data suggest that SARS-CoV-2 may impair, but does

not abolish, JAK-STAT signalling in infected cells, implying that recombinant IFNs may have a

therapeutic role in established SARS-CoV-2 infection, as recently shown in animal models[9] and in early

phase clinical trials[41].

Figure 6. Impact of endogenous IFN-I/III signalling on SARS-CoV-2 infection. Nasal ALI cultures

treated with ruxolitinib (RUX, 10 µM) or DMSO vehicle for 24 h prior to infection (MOI 0.1). Whole-

cell lysates were prepared at the indicated times for RT-PCR analysis of expression of (A) the ISGs

USP18, RSAD2 and *ISG15* (n=3 donors, mean ± SEM; *** P < 0.001, **** P < 0.0001, ANOVA with

Dunnett's post-test correction compared to mock-infected cells) or (B) viral *N* mRNA (n=3 donors,

mean ± SEM; * P = 0.035 ANOVA with Sidak's post-test correction compared to DMSO control). (C)

Whole-cell lysates were prepared at 96 hpi for immunoblot analysis of viral S/S2 protein and host

RSAD2, USP18 and ISG15 protein expression (representative blot shown of experiments in n=4

donors). (D) Densitometry analysis of S+S2 protein intensity relative to GAPDH, normalised to the

DMSO control (data from C, n=4 donors, mean ± SEM; ** P = 0.003, one-sample t test). (E) Plaque

assay of apical washes collected at the times indicated showing a significant increase in infectious

particle release at 96 hpi (same experimental conditions as C-D; n=4 donors, mean ± SEM; * P = 0.015,

ANOVA with Sidak's post-test correction compared to DMSO control). Dotted line indicates lower limit

of assay detection.

Figure 7. Exogenous IFN-I/III treatment controls SARS-CoV-2 replication. Nasal ALI cultures were pre-

treated for 16 h with IFNβ (1000 IU/mL) or IFNλ1 (100 ng/mL) prior to infection (MOI 0.01). (A)

Immunoblot of whole-cell lysates prepared from nasal ALI cultures at 48 hpi (representative of

experiments in n=4 donors). (B) Plaque assay of apical washes showing significant reduction in

infectious particle release at 48 hpi if pre-treated with IFNβ (1000 IU/mL) or IFNλ1 (100 ng/mL) (same

experimental conditions as A; n=5 donors, mean ± SEM; **** P < 0.0001, ANOVA with Dunnett's post-

test correction compared to untreated control). (C-D) Immunoblot of whole-cell lysates prepared at

48 hpi. Nasal ALI cultures were either pre-treated with IFNβ (1000 IU/mL, C) or IFNλ1 (100 ng/mL, D)

for 16h prior to infection with SARS-CoV-2 or IFN treatment was applied at 6 or 24 hpi. Results

representative of experiments in n=3 donors. (E) Plaque assay on apical washes collected at 48 hpi

from experiments in C-D (n=3 donors, mean $\pm$ SEM; * P < 0.05, ** P < 0.01, *** P < 0.001, ANOVA with

Dunnett's post-test correction compared to untreated control). Dotted line indicates lower limit of

assay detection.

DISCUSSION

We report the most comprehensive characterisation of the human nasal epithelial response to experimental SARS-CoV-2 infection to date, revealing a response dominated at later stages by IFN-I/IIIs and their downstream ISG products. This response partially contained SARS-CoV-2 at later times post-infection, while recombinant IFN-I/III treatment potently blocked SARS-CoV-2 replication, suggesting that mucosal delivery of IFNs could be a promising strategy for post-exposure prophylaxis. The nasal mucosa is likely to be a main point of entry of SARS-CoV-2. Prior single-cell transcriptomic studies implied an abundance of target cells in the nasal mucosa and further suggested that they may be poised to mount an antiviral response[10]. Yet few studies to date have characterised SARS-CoV-2 replication in primary human differentiated nasal cells[12-14], while we analyse the host-virus interaction comprehensively, at single-cell resolution and utilising proteomics. Our findings indicate that the host response to SARS-CoV-2 in nasal epithelium is dominated by paracrine IFN-I/III signalling, albeit this response is kinetically delayed. These data contrast with initial reports that SARS-CoV-2 did not induce a robust IFN response in airway epithelial cells[26-28], but are consistent with emerging evidence of IFN-I/III induction in nasal swabs from patients with COVID-19[15,42-44] and with more recent findings in lung airway models[36,43,45-47]. Blockade of the endogenous IFN response had an impact on SARS-CoV-2 infection at later stages post-infection, once the IFN response was established, underscoring the delayed kinetic but also emphasising its functional relevance. While the impact of endogenous IFN-I/III signalling upon SARS-CoV-2 replication has not to our knowledge been investigated in nasal cell models, our data are consistent with recent findings in some[43,47], but not all epithelial model systems[27,45]. Our experiments with IAV, PIV3 and SeV - viruses which induced the robust early expression of IFN-I/III, in line with previous studies[12,27,37] - confirm that this delay was not due to an intrinsic property of nasal epithelial cells. The expression of IFN evasion proteins[37,48], the sequestration of viral replication machinery within cytosolic vesicles[49], as well as global reductions in host mRNA content[50] and translational shutdown[51-53] induced by SARS-CoV-2 presumably underlie its capacity to subvert early IFN induction in infected cells. Consistent with this, there was evidence of

downregulation of immune pathways including TLR signalling in the proteome of infected cells.

However, an important question is what molecular patterns are responsible for IFN-I/III induction at

later times. Recent evidence implicates MDA5 as a major sensor of SARS-CoV-2 RNA in epithelial

cells[45-47], while other findings suggest a contribution from virus-mediated damage occurring after

several days of infection[54]. It will be important to address the relative contribution of host damage-

associated molecular patterns versus viral pathogen-associated molecular patterns (e.g. defective

viral genomes) accumulating during replication.

IFN-I/III signalling is plausibly implicated in protection against life-threatening COVID-19[18-20].

Consistent with this, circulating immune cells of patients with severe COVID-19 exhibit impaired ISG

responses[55-57]. However, whether the local airway IFN response in the early stages of infection has a

decisive role in shaping the subsequent clinical outcome of COVID-19 remains to be conclusively

determined. A compelling recent scRNA-seq study reported that patients going on to develop severe

disease exhibited a muted ISG response in the nasal airway, in contrast to those with milder disease[15],

and is supported by independent findings of attenuated nasal ISG induction in patients with

autoantibodies to IFN-I[58], who are prone to more severe disease[20]. Additional strands of evidence

suggest a potential link between airway IFN-I/III competence and clinical outcome in COVID-19. Age

remains the strongest risk factor for poor outcome in COVID-19, and the efficiency of IFN-I/III

induction is known to decline with advancing age[59], and appears to be greater in the nasal airways of

children than adults infected with SARS-CoV-2[60]. Other relevant environmental influences, such as

exposure to cigarette smoke or other viral infections, are also reported to perturb IFN-I/III responses

of airway cells in ways that may be relevant to COVID-19 pathogenesis[43,61].

The main limitation of our data in this nasal epithelial culture system is that it did not account for

professional immune cells present in the nasal mucosa, for example plasmacytoid dendritic cells[62],

which are capable of more rapidly mounting an IFN-I/III response to SARS-CoV-2[63], potentially tipping

the scales in favour of the host[64]. We studied cells derived from adult donors, however it is possible

that nasal cells from paediatric donors, who are naturally less susceptible to severe COVID-19, may

behave differently in terms of their reduced permissiveness to SARS-CoV-2 and/or the greater

efficiency of their innate IFN response[60,65]. Furthermore, SARS-CoV-2 variants with mutations in the

spike gene have emerged worldwide whilst we were undertaking the experiments described here;

these variants may impact viral replication and/or host immunity, and should be included in future

studies.

Nevertheless, our data, employing a variety of complementary methods, indicate that SARS-CoV-2 has

a relatively broad tropism for nasal epithelial cells, confirming suggestions from prior scRNA-seq

studies[8,10], other *in vitro* studies of primary nasal[14] and tracheobronchial cells[24,36], and importantly

recent scRNA-seq studies of nasal samples from COVID-19 patients[15]. We also identify tropism for the

rare deuterosomal cell, marked by expression of FOXN4, as recently reported[36,61,66]. Our findings

contrast with the results of Hou and colleagues, who reported exclusive tropism of SARS-CoV-2 for

ciliated cells in the nasal airway[13]. It is not immediately clear how to reconcile these findings, given

that secretory cells express relevant entry receptors[10,13] and have been identified as a major infected

cell type in infected patients[66]. Hou and colleagues used a fluorescent reporter virus, the tropism of

which might have been slightly narrower than clinical isolates. It is also worth noting that while we

show that all cell types contained SARS-CoV-2 protein, there was a significant reduction in the

proportion of goblet cells expressing spike protein, and the intensity of spike immunodetection was

significantly greater in ciliated and secretory cells than basal or goblet cells. Ciliated cells also

contained more virion-like structures per cell. Collectively, this implies that although all cell types are

permissive to SARS-CoV-2 entry, there may also be quantitative differences in the overall efficiency of

viral replication in different cell types. Hou and colleagues previously hypothesised that post-entry

factors such as intrinsic antiviral immunity might dictate permissiveness. As discussed, we found

limited evidence to support such a correlation, since while virtually all nasal epithelial cells

demonstrated a basal ISG signature - consistent with *ex vivo* nasal biopsy data[10] - this was apparently

insufficient to mediate resistance to SARS-CoV-2, at least at the time point analysed. However, it

remains possible that cell-type specific differences in the efficiency of induction of the IFN response

(for example in basal cells) might contribute to more subtle variation in permissiveness.

The differential response of basal cell types to SARS-CoV-2 at 24 hpi identified by our scRNA-seq

analysis appears a novel observation. Basal cells are the stem/progenitor cell population of the

airway[67]. Recent data indicate an emerging role for these cells as sentinels of the airway inflammatory

response[68]. For example, basal cells detect apoptotic cells in the context of viral inflammation[69],

retaining memory of prior immune exposure[70]. More generally, stem/progenitor cell types exhibit

enhanced intrinsic antiviral immunity[71]. Future studies should consider mechanism(s) governing the

seemingly distinct early antiviral response of nasal airway basal cells to SARS-CoV-2, and its functional

relevance.

Importantly, from a clinical perspective, the observation that IFN-I/III treatment prevented SARS-CoV-

2 infection *in vitro* indicates that chemoprophylaxis with IFN-I/III may have therapeutic value. This

approach has already been tested in a small clinical trial in China (although the absence of a control

group makes it impossible to judge the efficacy of this approach[72]). Immunisation is the most tractable

approach for large-scale primary prevention of COVID-19. However, owing to incomplete vaccine

coverage, and reduced vaccine effectiveness in immunocompromised populations or against mildly

symptomatic or asymptomatic infection, allied to the emergence of variants that may compromise

vaccine efficacy, there will likely continue to be a need for targeted chemoprophylactic therapies to

prevent transmission in specific circumstances. These include post-exposure prophylaxis of contacts -

to avoid the need for self-isolation - as well as pre-exposure prophylaxis for certain high-risk

encounters (e.g. in healthcare settings or prior to long-distance travel). Our data suggest that nasal

application of IFNβ or IFNλ1 might have an important role to play in this setting and argue for urgent

clinical assessment of this approach. In terms of the therapeutic efficacy of mucosally-administered

IFNβ in patients with established COVID-19[41], our findings suggest that early administration may be a

key factor determining clinical efficacy. Furthermore, studies in animal models indicate that

administration of IFNβ or IFNλ1 later in the disease course may have deleterious effects on viral

inflammation and/or airway cell regeneration[73-75], suggesting the existence of a relatively narrow

therapeutic window of opportunity.

In summary, we have shown that SARS-CoV-2 exhibits broad tropism for nasal epithelial cells, but with

preferential infection of ciliated and secretory cell types. Nasal cells mount a robust innate antiviral

response to SARS-CoV-2 dominated by paracrine IFN-I/III signalling, which is delayed in onset relative

to viral replication, but which is nevertheless capable of exerting partial control at later times post-

infection. Upon exposure to exogenous IFN-I/III treatment, these cells adopt a profound antiviral

state, highlighting a potential clinical role for recombinant IFNβ or IFNλ1 in chemoprophylaxis and/or

therapy of COVID-19.

METHODS

Adult nasal airway epithelial cell culture at air-liquid interface (ALI)

Adult primary human nasal airway epithelial cells were derived from excess clinical material obtained

during routine nasal surgical procedures[29]. Ethical approval for sample collection was provided

(Research Ethics Committee Reference 17/NE/0361) and written approved consent was provided

prior to sample collection. Tissue shaved from the superficial surface of the sample was chopped into

~2 mm^2 pieces and added to RPMI-1640 basal medium containing 0.1% protease (Sigma-Aldrich, UK)

and incubated overnight with gentle agitation at 4°C. All large pieces of tissue were discarded, and

residual protease was neutralized with 5% FCS. The preparation was centrifuged (200 g; 7 min) and

the pellet resuspended in PneumaCult-Ex Plus expansion medium (Stemcell Technologies), then

seeded onto 25 cm^2 tissue culture flasks pre-coated with 30 μg/mL Type I collagen (PureCol, Advanced

BioMatrix). Flasks were incubated in a humidified atmosphere containing 5% CO2 at 37°C, with

medium replaced every 48 hours. Cells were trypsinised at 60-80% confluence and cryopreserved for

future use. Upon thawing, cells were grown through an additional expansion phase, then transferred

in Ex Plus medium onto collagen-coated 6.5 mm polyester transwell membranes with 0.4 μm pore

size (Corning) at a density of 150,000 cells/cm^2. When cells were fully confluent, apical medium was

removed and basolateral medium was switched to PneumaCult-ALI-S (Stemcell Technologies). Cells

were maintained at air-liquid interface until fully differentiated. Barrier integrity of ALI cultures was

monitored by measuring trans-epithelial electrical resistance (TEER, EVOM 2, World Precision

Instruments). ALI cultures were validated for use in experiments based on microscopic appearance of

appropriate ciliated morphology and TEER > 500 Ω*cm^2. The sex and age of donors are included in

Table S4.

Viruses, cytokines and inhibitors

A clinical isolate of SARS-CoV-2 (BetaCoV/England/2/2020) was obtained from Public Health England

(PHE). This was isolated from a patient in January 2020 and thus represents an early strain of SARS-

CoV-2, not known to be affected by variants of concern. The initial stock was propagated once in vero

E6 cells. The same viral stock was used for all experiments. As SARS-CoV-2 is a Hazard Group 3

pathogen (Advisory Committee on Dangerous Pathogens, UK), all infection experiments were

performed in a dedicated Containment Level 3 (CL3) facility by trained personnel. Sendai virus (Cantell

strain) and parainfluenza virus 3 (PIV3) was obtained from Richard Randall (St Andrew's University).

Influenza A virus (A/PR8/1934/H1N1) was propagated and titred on MDCK cells. For nasal ALI

infections, apical poles were gently washed once with warm Dulbecco's modified Eagle's medium

(DMEM; Gibco, USA) and then infected with 60 µL dilution of virus in DMEM, at a MOI between 2 and

0.01 plaque-forming units per cell for 2 hours, when the virus-containing medium was removed.

DMEM was used as inoculum for mock infection. Apical washes (in warm phosphate-buffered saline)

were collected at different time points and stored at -80°C for plaque assays. Plaque assays were

undertaken in vero E6 cells using a 1.2% (w/v) microcrystalline cellulose overlay (Sigma-Aldrich).

Cytokines/inhibitors were used at the following concentrations: human recombinant IFNβ1 (1000

ng/mL; Avonex, NDC 59627-002-06, Biogen Inc, USA); IFNλ1 (100 ng/mL; 1598-IL-025, R&D Systems,

USA); and Ruxolitinib (10 µM; S1378, Calbiochem, USA) alongside the appropriate dilution of DMSO

vehicle. Treatment was applied through basolateral poles.

Single cell RNA sequencing (scRNA-seq) sample processing

For the droplet-encapsulation scRNA-seq experiments, ALI cultures were washed with PBS and then incubated with 1x Trypsin-EDTA (ThermoFisher Scientific, USA) for 10 min before the cells were diluted with DMEM and counted using a haemocytometer. 20,000 single cells were loaded onto each channel of a Chromium chip before encapsulation on the Chromium Controller (10x Genomics, USA). The single-cell sequencing libraries were generated using the Single Cell 5' V.1, as per the manufacturer's protocol. Libraries were sequenced using NovoSeq 6000 to achieve a minimum depth of 50,000 raw reads per cell. The libraries were sequenced using the following parameters: Read1: 26 cycles, i7: 8 cycles, i5: 0 cycles; Read2: 98 cycles to generate 75 bp paired-end reads.

Single-cell RNA sequencing (scRNA-seq) data generation and annotation

Sequencing data were demultiplexed and quantified using the Cellranger tool (version 4.0.0, 10x Genomics) and aligned to the combined human (official Cell Ranger reference, GRCh38-2020-A) and SARS-CoV-2 reference transcriptomes (Ensembl reference Sars_cov_2.ASM985889v3). CellBender (version 0.2.0)[76] was applied to the output from Cell Ranger software after alignment to remove background effect from ambient mRNA released during processing. Doublet detection and exclusion was performed using Scrublet (version 0.2.1) with thresholding of cells with a doublet score above two median absolute deviations from the median. Low-quality cells were removed using thresholds of < 200 genes and > 20% mitochondrial content. The analysis was performed using Seurat (version 4.0.1). Data were normalised and log-transformed using NormalizeData and the top 2000 variable genes identified using the FindVariableFeatures tool. The first 20 principal components were batch-adjusted using Harmony (by sample ID) and used to generate the nearest-neighbour graph. Dimensionality reduction and embedding was performed using Uniform Manifold Approximation and Projection (UMAP), with the neighbourhood graph clustered using the Leiden algorithm. The Wilcoxon rank sum test ($\log_2$ fold change threshold of 0.25, adjusted P value of 0.05) was used to identify

differentially expressed genes between clusters, and these were annotated based on expression of

markers from literature. This annotation was validated by comparing to a recently published scRNA-

seq dataset from ex vivo primary nasal cells[15]. The Seurat label transfer tool was used to assign

predicted identities to our data using the external published data as reference. The robustness of our

annotation was then assessed by the strength of its correlation with this prediction.

Gene set scoring and gene set enrichment analysis

A published gene set derived from IFN alpha or IFN gamma-treated human nasal basal cells[11] was used

to generate a list of epithelial-specific IFN-stimulated genes (ISGs). To calculate basal expression of

ISGs within the unexposed cells, over-expression of this gene list was assessed using the Seurat

AddModuleScore tool. Differences between clusters was compared by a Wilcoxon rank sum test

(adjusted alpha 0.05) with Benjamini-Hochburg multiple test correction applied.

Gene set enrichment analysis was performed using the fgsea tool[77]. Genes were ordered between

mock-infected and infected cells by fold-change in expression with Wilcoxon rank sum testing using

the FindAllMarkers function in Seurat, but without thresholds. Gene sets from Hallmark, Reactome

and Biocarta were used as reference after filtering to exclude those with fewer than 50 and greater

than 200 genes. Output from fgsea was further filtered to remove pathways that were not significantly

enriched in any cell type (adjusted P value <0.05) followed by further manual curation of the resultant

pathways.

Regulon scoring and analysis

The DoRoTHea/Viper package in R was used to score regulon activity by cell[78]. Human regulons from

DoRoTHea were filtered for those with a confidence score A-C. Normalised enrichment scores (NES)

for each transcription factor (TF) were calculated using run_viper with a minimum regulon size of 4 on

the complete gene expression matrix. To estimate TF activity over baseline within each cell type in the

infected cells, the median NES from the mock-infected clusters was subtracted from the equivalent in

the infected cluster. These scores were then scaled within each TF to give comparative estimate of TF

activity between clusters.

Quantitative RT-PCR

Total RNA was isolated using TRIzol[TM] reagent (Invitrogen, Carlsbad, CA, USA) according to the

manufacturer's instructions. For qPCR analyses of transcripts, 250 ng of RNA isolated from the nasal

epithelial cells was reverse transcribed with Superscript III (ThermoFisher Scientific), and the resulting

cDNA templates were subjected to qPCR with a TaqMan[TM] Gene Expression Master Mix (Applied

Biosystems, MA, USA) and AriaMx real-time PCR system (Agilent Technologies, CA, USA) according to

the manufacturer's instructions. The following TaqMan gene expression assays (Thermo Fischer) were

used: *IFNA1* (Hs03044218_g1), *TNF* (Hs00174128_m1). The primers were designed using the Roche

Universal Probe Library (UPL) Assay Design tool (Roche, Basel, Switzerland) with the indicated UPL

probes. All other primer and probe information is described in Table S5. Cycling conditions were as

follows: reverse transcription at 50°C for 15 min, followed by initial polymerase activation at 95°C for

10 min, then 40 cycles of denaturation at 95°C for 15 sec and annealing/extension at 60°C for 1 min.

The $2^{-\Delta\Delta Ct}$ method was used to calculate the relative expression of genes. Each sample was run in

duplicate. Samples were normalised to the endogenous housekeeping gene expression, either

RNASEP for *N* gene expression, or *18S* for all other genes.

Immunofluorescence

Immunofluorescence analysis was performed in accordance with published methods[79]. Infected and

mock-infected membranes were fixed *in situ* with 4% (w/v) paraformaldehyde overnight at 4°C, before

removal from transwells and sectioning. Membranes were washed twice for 5 min in PBS plus 0.1%

(v/v) Triton X-100 (Sigma-Aldrich) before being blocked with a 1% (w/v) BSA solution in PBS with 0.5%

(v/v) Tween 20 (Sigma-Aldrich; PBST) for one hour at room temperature (RT). Membranes were

incubated with primary antibodies for 2 hours at RT. Antibodies used are listed in Table S6.

Membranes were washed three times for 5 min in PBST before incubation with appropriate

fluorescence-conjugated secondary antibodies for 2 hours at RT. This process was repeated as part of

a sequential staining process where required. Membranes were washed three times for 5 min in PBST

before being incubated with DAPI (50 nM; Sigma-Aldrich) as a nuclear counterstain, and phalloidin,

DyLight 650 (1 unit/mL; ThermoFisher) where required, for 10 min. Membranes were then mounted

in MOWIOL mounting media (Sigma-Aldrich) and coverslips applied. Appropriate secondary only

controls were performed as required. Images were captured using a Nikon A1 confocal microscope

(Nikon, Japan), with all capture settings standardised. For analysis three random fields per sample

were captured at x20 magnification and analysed via Fiji (Version 2.0) using the Cell Counter plugin.

Total cell count, and number of spike protein-positive cells, were determined by segmenting images

using ZO-1 and DAPI to identify individual cells. Mean pixel intensity of the spike protein in positive

cells was also determined using the Plot Profile plugin.

Immunoblot

Immunoblotting was carried out as previously described[40]. Briefly, proteins from cell lysates were

separated by 10% sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) using

MOPS running buffer (Thermo Fisher, USA), and transferred to a nitrocellulose membrane (Millipore,

USA) using NuPage Tris-Bis Transfer Buffer (Thermo Fisher) for immunoblotting (for details of

antibodies see Table S6). Blots were developed with Pierce ECL Western blotting substrate (Thermo

Fisher) and imaged on a LI-COR Odyssey Fc (LI-COR, USA). Densitometry was undertaken using

ImageStudio software (LI-COR).

Transmission electron microscopy (TEM)

Cultures were fixed with 2% glutaraldehyde (Sigma-Aldrich, MO, USA) in 0.1 M sodium cacodylate (pH

7.4) buffer in the apical and basal compartment and then kept at 4°C overnight. For TEM resin

processing, the monolayer membranes were removed from the insert frame and placed in microwave

684 sample holders. The Pelco Biowave Pro+ microwave (Pelco, CA, USA), incorporating the Pelco ColdSpot

685 Pro system, was used for the following steps of the processing. The ColdSpot system

686 improves inconsistent wattage supply to the microwave compartment, therefore protecting samples

687 from excess microwave energy. The range temperature was set at 23–27°C. Following buffer rinses

688 (three pulses at 150 watts (W) for 40 secs) the samples were post-fixed in 1% osmium tetroxide for 8

689 min [pulse microwaved (MW), 100 W] and rinsed in distilled H_2O (three times at 150 W for 40 secs,

690 per step). Samples were dehydrated in a graded series of acetone (25%; 50%; 75%; three times with

691 100% (v/v); 150 W, 40 secs per step) before being impregnated with increasing concentrations of

692 epoxy resin (medium resin; TAAB, UK) in acetone (25%; 50%; 75%; three times at 100% (v/v); 300 W,

693 3 min per step). The samples were then embedded in 100% fresh resin and left to polymerise at 60°C

694 in a conventional oven for a minimum of 24 hours. All resin blocks were trimmed using a razor blade

695 to form a trapezoid block face. Sections were cut on an ultramicrotome using a diamond knife. Semi

696 thin sections (0.5 μm) were stained with toluidine blue and viewed on a light microscope to verify

697 presence of cell monolayers. Ultrathin sections (70 nm) were then cut and picked up onto pioloform-

698 coated copper grids. Grids were stained with 1% (w/v) uranyl acetate (30 min) and 3% (w/v) lead

699 citrate (7 min) to improve contrast. All sections were examined using a HT7800 120 kV TEM (Hitachi,

700 Japan). Digital micrographs were captured using an EMSIS Xarosa CMOS Camera with Radius software

701 (EMSIS, Germany). ImageJ software (Version 2.0) was used to enhance the contrast by increasing the

702 percentage of saturated pixels to 1% to aid virus-like particle identification. Virus-like particles

703 (∼70 nm in diameter) were counted using the cell counter plugin in at least 6 goblet cells and 16

704 ciliated cells per donor. Data were presented as number of virus-like particles per cell.

706 **Proteome sample preparation**

707 The protein concentration was determined by EZQ® protein quantification assay. Protein digestion was

708 performed using the S-Trap™ sample preparation method and TMT-16 plex labelling was carried out

709 as per the manufacturer's instructions. Samples were cleaned using MacroSpin columns, and dried

down prior to offline high-performance liquid chromatography fractionation. Peptides were

fractionated on a Basic Reverse Phase column on a Dionex Ultimate 3000 off-line LC system. A total of

18 fractions were collected, and each fraction was acidified and dried. Peptides were dissolved in 5%

formic acid, and each sample was independently analysed on an Orbitrap Fusion Lumos Tribrid mass

spectrometer, connected to an UltiMate 3000 RSLCnano System. All spectra were analysed using

MaxQuant 1.6.10.43 and searched against SwissProt *Homo sapiens* and Trembl SARS-CoV-2 FASTA

files. Reporter ion MS3 was used for quantification and the additional parameter of quantitation labels

with 16 plex TMT on N-terminus or lysine was included. A protein and peptide false discovery rate

(FDR) of less than 1% was employed in MaxQuant. Moderated t-tests, with patient accounted for in

the linear model, was performed using Limma, where proteins with an adjusted $P < 0.05$ were

considered as statistically significant. All analysis was performed using R. Raw data are present in

supplementary dataset 5. A comprehensive description of the methods can be found in the

supplementary methods.

Statistical analysis

Statistical analysis was performed and figures assembled using GraphPad Prism V9 (GraphPad

Software, USA). Data are presented as mean ± SEM of individual donor values (derived typically from

2-3 independent repeat experiments per donor). The donor was used as the unit of experiment for

statistical analysis purposes. Continuous data were normalised or log-transformed prior to analysis

using parametric significance tests, or if this was not possible, were analysed using nonparametric

significance tests. Differences between two groups were compared using an unpaired, two-tailed

Student's t-test (or Mann-Whitney test for TEM image analysis), whereas differences between more

than two groups used ANOVA, with Dunnett's post-test correction for multiple comparisons when

comparing to a single reference point (e.g. mock-infected or time zero) or with Sidak's post-test

correction for other multiple comparisons (e.g. between differently treated donors at the same time

point). In some cases, data were normalised to a reference point, where a one-sample t-test was used.

Unless stated otherwise, a two-tailed alpha of < 0.05 was the threshold for statistical significance.

Statistical analysis of proteomics and transcriptomics data sets is described in the relevant sections

above.

Data availability

Source data are provided with this paper. This includes uncropped blots, all quantitative data and the

results of differential expression analysis of RNA-seq and proteomics data, which are included as

supplementary datasets. Additional raw data are available on request from the corresponding author

providing ethical approvals permit sharing of data. The mass spectrometry proteomics data have been

deposited to the ProteomeXchange Consortium[80] via the PRIDE partner repository[81] with the dataset

identifier PXD022523. This can be accessed through the Username: reviewer_pxd022523@ebi.ac.uk

and the Password: UaEXYFKF. Raw RNA sequencing data have been deposited to the European

Genome-Phenome Archive (accession pending). Processed scRNAseq data is available at Zenodo

(https://zenodo.org/record/4564332).

Code availability

Analysis scripts and codes are available at github.com/haniffalab/covid_nasal_epithelium.

Acknowledgements

We acknowledge Public Health England for providing the SARS-CoV-2 isolate and Professor R Randall

(St Andrew's University) for providing Sendai and parainfluenza 3 viruses and the vero E6 cell line. We

thank M Glanville and Dr S Dainty (Newcastle University Infectious Disease Facility) and Dr A Laude

(Bioimaging facility) for assistance. Tissue for this study was provided by the Newcastle Biobank which

is supported by the Newcastle upon Tyne NHS Foundation Trust and Newcastle University. The work

was part-funded by the Barbour Foundation (CJAD, SH, MB, MT, MH, CW), the UK-Coronavirus

Immunology Consortium (CJAD, SH, MH) and the Medical Research Council SHIELD antimicrobial

resistance consortium (JS, AJS). CFH is supported by a Medical Research Council studentship

(MR/NO13840/1) and MB by an MRC Clinician Scientist Fellowship (MR/M008797/1). MT and SH are

funded by Wellcome Investigator Awards (215542/Z/19/Z and 207556/Z/17/Z). CJAD and GR are

Wellcome Clinical Research Career Development Fellows (211153/Z/18/Z and 214539/Z/18/Z). MED

is a Marie Sklodowska Curie Fellow within the European Union's Horizon 2020 research and innovation

programme under the Marie Sklodowska-Curie grant agreement No. 890296. MH is funded by

Wellcome (WT107931/Z/15/Z), The Lister Institute for Preventive Medicine and Newcastle NIHR

Biomedical Research Centre (BRC). JPG, CW and BV were supported by the Medical Research

Foundation (MRF Respiratory Diseases Research Award to JPG; Grant MRF-091-0001-RGGARNE) and

Boehringer Ingelheim. TEM work was supported by a BBSRC Alert17 grant (BB/R013942/1). AJS is a

National Institute for Health Research (NIHR) Senior Investigator. The views expressed in this article

are those of the author(s) and not necessarily those of the NIHR, or the Department of Health and

Social Care. The funders had no role in the study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Author contributions

Conceived the study: MB, CW, CJAD with MH, SH, MT. Experimental design: MH, SH, MB, CW, MT, GR,

CJAD. Nasal model development and patient material: IJH, BV, JS, JPG, SC, JP, AJS, MB, CW. Virology

data generation, analysis and interpretation: CFH, BJT, JSS, FG, AIG, AK, LH, TD, SH, CJAD. Proteomics

data generation, analysis and interpretation: MED, SM, MT. Single-cell sequencing data generation,

analysis and interpretation: RAB, ES, RH, JC, MH, GR. Supervised research: AJS, MH, SH, MB, CW, MT,

GR, CJAD. Drafted the manuscript: CFH and CJAD with ES, BV, MED, TD, MT, GR. Revised the

manuscript: RAB, MED, IJH, JSS, AJS, MH, SH, CW. Approved the manuscript for submission: all

authors.

Declaration of competing interests

SH declares honoraria from CSL Behring and Takeda for teaching and consultancy. The remaining

authors declare no financial or non-financial competing interests.

References

Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278, doi:10.1016/j.cell.2020.02.052 (2020).

Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506, doi:10.1016/S0140-6736(20)30183-5 (2020).

Knight, S. R. *et al.* Risk stratification of patients admitted to hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: development and validation of the 4C Mortality Score. *BMJ* **370**, m3339, doi:10.1136/bmj.m3339 (2020).

Group, R. C. *et al.* Dexamethasone in Hospitalized Patients with Covid-19 - Preliminary Report. *N Engl J Med*, doi:10.1056/NEJMoa2021436 (2020).

Group, R. C. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet* **397**, 1637-1645, doi:10.1016/S0140-6736(21)00676-0 (2021).

Cevik, M., Kuppalli, K., Kindrachuk, J. & Peiris, M. Virology, transmission, and pathogenesis of SARS-CoV-2. *BMJ* **371**, m3862, doi:10.1136/bmj.m3862 (2020).

Chen, M. *et al.* Elevated ACE-2 expression in the olfactory neuroepithelium: implications for anosmia and upper respiratory SARS-CoV-2 entry and replication. *Eur Respir J* **56**, doi:10.1183/13993003.01948-2020 (2020).

Hoagland, D. A. *et al.* Leveraging the antiviral type I interferon system as a first line of defense against SARS-CoV-2 pathogenicity. *Immunity*, doi:10.1016/j.immuni.2021.01.017 (2021).

Sungnak, W. *et al.* SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat Med* **26**, 681-687, doi:10.1038/s41591-020-0868-6 (2020).

Ziegler, C. G. K. *et al.* SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell* **181**, 1016-1035 e1019, doi:10.1016/j.cell.2020.04.035 (2020).

Gamage, A. M. *et al.* Infection of human Nasal Epithelial Cells with SARS-CoV-2 and a 382-nt deletion isolate lacking ORF8 reveals similar viral kinetics and host transcriptional profiles. *PLoS Pathog* **16**, e1009130, doi:10.1371/journal.ppat.1009130 (2020).

Hou, Y. J. *et al.* SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the Respiratory Tract. *Cell* **182**, 429-446 e414, doi:10.1016/j.cell.2020.05.042 (2020).

Pizzorno, A. *et al.* Characterization and Treatment of SARS-CoV-2 in Nasal and Bronchial Human Airway Epithelia. *Cell Rep Med* **1**, 100059, doi:10.1016/j.xcrm.2020.100059 (2020).

Ziegler, C. G. K. *et al.* Impaired local intrinsic immunity to SARS-CoV-2 infection in severe COVID-19. *Cell*, doi:10.1016/j.cell.2021.07.023 (2021).

Onabajo, O. O. *et al.* Interferons and viruses induce a novel truncated ACE2 isoform and not the full-length SARS-CoV-2 receptor. *Nat Genet* **52**, 1283-1293, doi:10.1038/s41588-020-00731-9 (2020).

17 Duncan, C. J. A., Randall, R. E. & Hambleton, S. Genetic Lesions of Type I Interferon
Signalling in Human Antiviral Immunity. *Trends Genet* **37**, 46-58,
doi:10.1016/j.tig.2020.08.017 (2021).
18 Zhang, Q. *et al.* Inborn errors of type I IFN immunity in patients with life-threatening
COVID-19. *Science* **370**, doi:10.1126/science.abd4570 (2020).
19 Pairo-Castineira, E. *et al.* Genetic mechanisms of critical illness in Covid-19. *Nature*,
doi:10.1038/s41586-020-03065-y (2020).
20 Bastard, P. *et al.* Autoantibodies against type I IFNs in patients with life-threatening
COVID-19. *Science* **370**, doi:10.1126/science.abd4585 (2020).
21 Lokugamage, K. G. *et al.* Type I Interferon Susceptibility Distinguishes SARS-CoV-2
from SARS-CoV. *J Virol* **94**, doi:10.1128/JVI.01410-20 (2020).
22 Felgenhauer, U. *et al.* Inhibition of SARS-CoV-2 by type I and type III interferons. *J Biol
Chem* **295**, 13958-13964, doi:10.1074/jbc.AC120.013788 (2020).
23 Lee, J. S. & Shin, E. C. The type I interferon response in COVID-19: implications for
treatment. *Nat Rev Immunol* **20**, 585-586, doi:10.1038/s41577-020-00429-3 (2020).
24 Zhu, N. *et al.* Morphogenesis and cytopathic effect of SARS-CoV-2 infection in human
airway epithelial cells. *Nat Commun* **11**, 3910, doi:10.1038/s41467-020-17796-z
(2020).
25 Fiege, J. K. *et al.* Single cell resolution of SARS-CoV-2 tropism, antiviral responses, and
susceptibility to therapies in primary human airway epithelium. *PLoS Pathog* **17**,
e1009292, doi:10.1371/journal.ppat.1009292 (2021).
26 Vanderheiden, A. *et al.* Type I and Type III Interferons Restrict SARS-CoV-2 Infection of
Human Airway Epithelial Cultures. *J Virol* **94**, doi:10.1128/JVI.00985-20 (2020).
27 Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives Development
of COVID-19. *Cell* **181**, 1036-1045 e1039, doi:10.1016/j.cell.2020.04.026 (2020).
28 Huang, J. *et al.* SARS-CoV-2 Infection of Pluripotent Stem Cell-derived Human Lung
Alveolar Type 2 Cells Elicits a Rapid Epithelial-Intrinsic Inflammatory Response.
bioRxiv, doi:10.1101/2020.06.30.175695 (2020).
29 Mavin, E. *et al.* Real-time measurement of cellular bioenergetics in fully differentiated
human nasal epithelial cells grown at air-liquid-interface. *Am J Physiol Lung Cell Mol
Physiol* **318**, L1158-L1164, doi:10.1152/ajplung.00414.2019 (2020).
30 Vieira Braga, F. A. *et al.* A cellular census of human lungs identifies novel cell states in
health and in asthma. *Nat Med* **25**, 1153-1163, doi:10.1038/s41591-019-0468-5
(2019).
31 Ruiz Garcia, S. *et al.* Novel dynamics of human mucociliary differentiation revealed by
single-cell RNA sequencing of nasal epithelial cultures. *Development* **146**,
doi:10.1242/dev.177428 (2019).
32 Xiu, S. *et al.* Inhibitors of SARS-CoV-2 Entry: Current and Future Opportunities. *J Med
Chem* **63**, 12256-12274, doi:10.1021/acs.jmedchem.0c00502 (2020).
33 Prelli Bozzo, C. *et al.* IFITM proteins promote SARS-CoV-2 infection and are targets for
virus inhibition in vitro. *Nat Commun* **12**, 4584, doi:10.1038/s41467-021-24817-y
(2021).
34 Li, S. *et al.* Metabolic reprogramming and epigenetic changes of vital organs in SARS-
CoV-2-induced systemic toxicity. *JCI Insight* **6**, doi:10.1172/jci.insight.145027 (2021).
35 Moolamalla, S. T. R., Balasubramanian, R., Chauhan, R., Priyakumar, U. D. & Vinod, P.
K. Host metabolic reprogramming in response to SARS-CoV-2 infection: A systems

biology approach. *Microb Pathog* **158**, 105114, doi:10.1016/j.micpath.2021.105114
(2021).

36 Ravindra, N. G. *et al.* Single-cell longitudinal analysis of SARS-CoV-2 infection in human
airway epithelium identifies target cells, alterations in gene expression, and cell state
changes. *PLoS Biol* **19**, e3001143, doi:10.1371/journal.pbio.3001143 (2021).

37 Lei, X. *et al.* Activation and evasion of type I interferon responses by SARS-CoV-2. *Nat*
Commun **11**, 3810, doi:10.1038/s41467-020-17665-9 (2020).

38 Cardoso, R. S. *et al.* Host Retromer Protein Sorting Nexin 2 Interacts with Human
Respiratory Syncytial Virus Structural Proteins and is Required for Efficient Viral
Production. *mBio* **11**, doi:10.1128/mBio.01869-20 (2020).

39 Bergant Marusic, M., Ozbun, M. A., Campos, S. K., Myers, M. P. & Banks, L. Human
papillomavirus L2 facilitates viral escape from late endosomes via sorting nexin 17.
Traffic **13**, 455-467, doi:10.1111/j.1600-0854.2011.01320.x (2012).

40 Duncan, C. J. A. *et al.* Severe type I interferonopathy and unrestrained interferon
signaling due to a homozygous germline mutation in STAT2. *Sci Immunol* **4**,
doi:10.1126/sciimmunol.aav7501 (2019).

41 Monk, P. D. *et al.* Safety and efficacy of inhaled nebulised interferon beta-1a (SNG001)
for treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-
controlled, phase 2 trial. *Lancet Respir Med*, doi:10.1016/S2213-2600(20)30511-7
(2020).

42 Antonelli, G. *et al.* Type I interferons can be detected in respiratory swabs from SARS-
Cov-2 infected patients. *J Clin Virol* **128**, 104450, doi:10.1016/j.jcv.2020.104450
(2020).

43 Cheemarla, N. R. *et al.* Dynamic innate immune response determines susceptibility to
SARS-CoV-2 infection and early replication kinetics. *J Exp Med* **218**,
doi:10.1084/jem.20210583 (2021).

44 Mick, E. *et al.* Upper airway gene expression reveals suppressed immune responses to
SARS-CoV-2 compared with other respiratory viruses. *Nat Commun* **11**, 5854,
doi:10.1038/s41467-020-19587-y (2020).

45 Rebendenne, A. *et al.* SARS-CoV-2 triggers an MDA-5-dependent interferon response
which is unable to control replication in lung epithelial cells. *J Virol*,
doi:10.1128/JVI.02415-20 (2021).

46 Sampaio, N. G. *et al.* The RNA sensor MDA5 detects SARS-CoV-2 infection. *Sci Rep* **11**,
13638, doi:10.1038/s41598-021-92940-3 (2021).

47 Yin, X. *et al.* MDA5 Governs the Innate Immune Response to SARS-CoV-2 in Lung
Epithelial Cells. *Cell Rep* **34**, 108628, doi:10.1016/j.celrep.2020.108628 (2021).

48 Kumar, A. *et al.* SARS-CoV-2 Nonstructural Protein 1 Inhibits the Interferon Response
by Causing Depletion of Key Host Signaling Factors. *J Virol* **95**, e0026621,
doi:10.1128/JVI.00266-21 (2021).

49 Wolff, G., Melia, C. E., Snijder, E. J. & Barcena, M. Double-Membrane Vesicles as
Platforms for Viral Replication. *Trends Microbiol* **28**, 1022-1033,
doi:10.1016/j.tim.2020.05.009 (2020).

50 Finkel, Y. *et al.* SARS-CoV-2 uses a multipronged strategy to impede host protein
synthesis. *Nature* **594**, 240-245, doi:10.1038/s41586-021-03610-3 (2021).

51 Schubert, K. *et al.* SARS-CoV-2 Nsp1 binds the ribosomal mRNA channel to inhibit
translation. *Nat Struct Mol Biol* **27**, 959-966, doi:10.1038/s41594-020-0511-8 (2020).

52   Thoms, M. *et al.* Structural basis for translational shutdown and immune evasion by
the   Nsp1   protein   of   SARS-CoV-2.   *Science*   **369**,   1249-1255,
doi:10.1126/science.abc8665 (2020).
53   Hsu, J. C., Laurent-Rolle, M., Pawlak, J. B., Wilen, C. B. & Cresswell, P. Translational
shutdown and evasion of the innate immune response by SARS-CoV-2 NSP14 protein.
Proc Natl Acad Sci U S A **118**, doi:10.1073/pnas.2101161118 (2021).
54   Robinot, R. *et al.* SARS-CoV-2 infection induces the dedifferentiation of multiciliated
cells and impairs mucociliary clearance. *Nat Commun* **12**, 4354, doi:10.1038/s41467-
021-24521-x (2021).
55   Galani, I. E. *et al.* Untuned antiviral immunity in COVID-19 revealed by temporal type
I/III   interferon   patterns   and   flu   comparison.   *Nat   Immunol*   **22**,   32-40,
doi:10.1038/s41590-020-00840-x (2021).
56   Hadjadj, J. *et al.* Impaired type I interferon activity and inflammatory responses in
severe COVID-19 patients. *Science* **369**, 718-724, doi:10.1126/science.abc6027 (2020).
57   Stephenson, E. *et al.* Single-cell multi-omics analysis of the immune response in
COVID-19. *Nat Med* **27**, 904-916, doi:10.1038/s41591-021-01329-2 (2021).
58   Lopez, J. *et al.* Early nasal type I IFN immunity against SARS-CoV-2 is compromised in
patients   with   autoantibodies   against   type   I   IFNs.   *J   Exp   Med*   **218**,
doi:10.1084/jem.20211211 (2021).
59   Molony, R. D. *et al.* Aging impairs both primary and secondary RIG-I signaling for
interferon   induction   in   human   monocytes.   *Sci   Signal*   **10**,
doi:10.1126/scisignal.aan2392 (2017).
60   Loske, J. *et al.* Pre-activated antiviral innate immunity in the upper airways controls
early SARS-CoV-2 infection in children. *Nat Biotechnol*, doi:10.1038/s41587-021-
01037-9 (2021).
61   Purkayastha, A. *et al.* Direct Exposure to SARS-CoV-2 and Cigarette Smoke Increases
Infection Severity and Alters the Stem Cell-Derived Airway Repair Response. *Cell Stem
Cell* **27**, 869-875 e864, doi:10.1016/j.stem.2020.11.010 (2020).
62   Hartmann, E. *et al.* Analysis of plasmacytoid and myeloid dendritic cells in nasal
epithelium. *Clin Vaccine Immunol* **13**, 1278-1286, doi:10.1128/CVI.00172-06 (2006).
63   Onodi, F. *et al.* SARS-CoV-2 induces human plasmacytoid pre-dendritic cell
diversification via UNC93B and IRAK4. *bioRxiv*, doi:10.1101/2020.07.10.197343
(2021).
64   Cervantes-Barragan, L. *et al.* Plasmacytoid dendritic cells produce type I interferon and
reduce viral replication in airway epithelial cells after SARS-CoV-2 infection. *bioRxiv*,
doi:10.1101/2021.05.12.443948 (2021).
65   Pierce, C. A. *et al.* Natural mucosal barriers and COVID-19 in children. *JCI Insight* **6**,
doi:10.1172/jci.insight.148694 (2021).
66   Ziegler, C. G. K. *et al.* Impaired local intrinsic immunity to SARS-CoV-2 infection in
severe COVID-19. *bioRxiv*, doi:10.1101/2021.02.20.431155 (2021).
67   Rock, J. R. *et al.* Basal cells as stem cells of the mouse trachea and human airway
epithelium. *Proc Natl Acad Sci U S A* **106**, 12771-12775, doi:10.1073/pnas.0906850106
(2009).
68   Hewitt, R. J. & Lloyd, C. M. Regulation of immune responses by the airway epithelial
cell landscape. *Nat Rev Immunol* **21**, 347-362, doi:10.1038/s41577-020-00477-9
(2021).

69 Fujino, N. *et al.* Sensing of apoptotic cells through Axl causes lung basal cell
proliferation in inflammatory diseases. *J Exp Med* **216**, 2184-2201,
doi:10.1084/jem.20171978 (2019).
70 Ordovas-Montanes, J. *et al.* Allergic inflammatory memory in human respiratory
epithelial progenitor cells. *Nature* **560**, 649-654, doi:10.1038/s41586-018-0449-8
(2018).
71 Wu, X. *et al.* Intrinsic Immunity Shapes Viral Resistance of Stem Cells. *Cell* **172**, 423-
438 e425, doi:10.1016/j.cell.2017.11.018 (2018).
72 Meng, Z. *et al.* An experimental trial of recombinant human interferon alpha nasal
drops to prevent COVID-19 in medical staff in an epidemic area. *MedRxiv*,
doi:doi.org/10.1101/2020.04.11.20061473 (2020).
73 Boudewijns, R. *et al.* STAT2 signaling restricts viral dissemination but drives severe
pneumonia in SARS-CoV-2 infected hamsters. *Nat Commun* **11**, 5838,
doi:10.1038/s41467-020-19684-y (2020).
74 Major, J. *et al.* Type I and III interferons disrupt lung epithelial repair during recovery
from viral infection. *Science* **369**, 712-717, doi:10.1126/science.abc2061 (2020).
75 Broggi, A. *et al.* Type III interferons disrupt the lung epithelial barrier upon viral
recognition. *Science* **369**, 706-712, doi:10.1126/science.abc3545 (2020).
76 Fleming, S. J., Marioni, J. C. & Babadi, M. CellBender remove-background: a deep
generative model for unsupervised removal of background noise from scRNA-seq
datasets. *bioRxiv*, doi:10.1101/791699 (2019).
77 Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv*, doi:10.1101/060012
(2021).
78 Holland, C. H. *et al.* Robustness and applicability of transcription factor and pathway
analysis tools on single-cell RNA-seq data. *Genome Biol* **21**, 36, doi:10.1186/s13059-
020-1949-z (2020).
79 Crowe, A. R. & Yue, W. Semi-quantitative Determination of Protein Expression using
Immunohistochemistry Staining and Analysis: An Integrated Protocol. *Bio Protoc* **9**,
doi:10.21769/BioProtoc.3465 (2019).
80 Deutsch, E. W. *et al.* The ProteomeXchange consortium in 2017: supporting the
cultural change in proteomics public data deposition. *Nucleic Acids Res* **45**, D1100-
D1106, doi:10.1093/nar/gkw936 (2017).
81 Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019:
improving support for quantification data. *Nucleic Acids Res* **47**, D442-D450,
doi:10.1093/nar/gky1106 (2019).
1007