Delayed induction of type I and III interferons and nasal epithelial cell permissiveness to

SARS-CoV-2

Nasal interferon responses to SARS-CoV-2

Authors

Catherine F Hatton[1a], Rachel A Botting[2a], Maria Emilia Dueñas[2a], Iram J Haq[1,3a], Bernard Verdon[2a],

Benjamin J Thompson[1], Jarmila Stremenova Spegarova[1], Florian Gothe[1,4], Emily Stephenson[2], Aaron I

Gardner[1], Sandra Murphy[2], Jonathan Scott[1], James P Garnett[1], Sean Carrie[5], Rafiqul Hussain[6], Jonathan

Coxhead[6], Tracey Davey[7], A John Simpson[1], Muzlifah Haniffa[2,8,9,10], Sophie Hambleton[1,11], Malcolm

Brodlie[1,3b], Chris Ward[1b], Matthias Trost[2b], Gary Reynolds[2b], Christopher J A Duncan[1,12 b]*

[a] equal contribution

[b] equal contribution

* corresponding author: christopher.duncan@ncl.ac.uk

Affiliations

1. Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University,

Newcastle upon Tyne, UK.

2. Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK.

3. Paediatric Respiratory Medicine, Great North Children's Hospital, Newcastle upon Tyne Hospitals

NHS Foundation Trust, Newcastle upon Tyne, UK.

4. Department of Pediatrics, Dr. von Hauner Children's Hospital, University Hospital, Ludwig-

Maximilians-Universität Munich, Munich, Germany.

5. Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK.

6. Genomics Core Facility, Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK.

7. Electron Microscopy Research Services, Newcastle University, Newcastle upon Tyne, UK.

8. NIHR Newcastle Biomedical Research Centre, Newcastle Hospitals NHS Foundation Trust, Newcastle

upon Tyne, UK

9. Department of Dermatology, Newcastle Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

10. Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, UK

11. Great North Children's Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, UK.

12. Department of Infection and Tropical Medicine, Newcastle upon Tyne Hospitals NHS Foundation

Trust, UK.

Abstract

The nasal epithelium is a plausible entry point for SARS-CoV-2, a site of pathogenesis and transmission,

and may initiate the host response to SARS-CoV-2. Antiviral interferon responses are critical to

outcome of SARS-CoV-2. Yet little is known about the interaction between SARS-CoV-2 and innate

immunity in this tissue. Here we applied single-cell RNA sequencing and proteomics to a primary cell

model of human primary nasal epithelium differentiated at air-liquid interface. SARS-CoV-2

demonstrated widespread tropism for nasal epithelial cell types. The host response was dominated

by type I and III IFNs and interferon-stimulated gene products. Nevertheless, this response was

notably delayed in onset compared to viral gene expression, and thus failed to impact substantially on

SARS-CoV-2 replication. However, when provided prior to infection, recombinant IFNβ or IFNλ1

induced an efficient antiviral state that potently restricted SARS-CoV-2 viral replication, preserving

epithelial barrier integrity. These data suggest nasal delivery of recombinant IFNs to be a potential

chemoprophylactic strategy.

INTRODUCTION

SARS-CoV-2 is an emergent betacoronavirus responsible for coronavirus disease-19 (COVID-19)[1]. Since its identification in late 2019, global pandemic transmission of SARS-CoV-2 has resulted in over 105 million infections and approximately 2.9 million deaths. SARS-CoV-2 infects target cells via the entry receptor ACE2[2] leading to a spectrum of clinical outcomes, ranging from asymptomatic infection to death[3]. Although multiple host factors (e.g. age, sex, obesity) contribute to adverse clinical outcome[4], the immune response also plays a decisive role, evidenced by the therapeutic benefit of immunomodulatory agents including corticosteroids[5]. Yet much remains to be understood about the immunopathogenesis of COVID-19. Identification of the cells hosting viral entry and characterisation of their response to infection is essential to understanding pathogenesis and improving therapy.

The nasal epithelium is believed to be a key entry point of SARS-CoV-2. Nasal epithelial tropism and efficient viral shedding from the nasopharynx apparently contributes to the high transmissibility of SARS-CoV-2[6], as well as to pathologic features such as anosmia[7]. As an early viral target cell, nasal epithelial cells may also set the tone for the systemic immune response, potentially influencing disease outcome[8]. These factors emphasise the need to study host-virus interaction in human nasal cells. *Ex vivo* single-cell transcriptomic studies indicate that ciliated and/or goblet cells in the nasal mucosa express *ACE2* and *TMPRSS2,* implicating them as probable SARS-CoV-2 target cells[9,10]. This has been confirmed by *in vitro* and *in vivo* studies demonstrating SARS-CoV-2 infection of human nasal epithelial cells[11-13]. Single-cell studies also revealed that nasal cells exhibit basal expression of an antiviral expression programme, characterised by induction of several interferon-stimulated genes (ISGs), suggesting that they may be primed to respond to viral infection[9]. Interestingly, *ACE2* is also regulated by interferons (IFNs) [10,14], implying a complex relationship between IFN signalling and tropism. Type I and type III IFN (IFN-I/III) systems play a critical role in human antiviral innate immunity[15] and have been recently implicated in defence against SARS-CoV-2: susceptibility to severe or life-threatening COVID-19 is associated with deleterious variants in *IFNAR* genes[16,17] and IFN-I blocking autoantibodies[18]. *In vitro*, SARS-CoV-2 appears sensitive to the antiviral properties of IFN-I, at least in

cell lines[19,20], and this activity extends to *in vivo* model systems[8]. These findings motivate studies to

improve understanding of the interaction between SARS-CoV-2 and the IFN-I system in primary human

target cells, providing impetus to clinical trials of recombinant IFNs in treatment or prophylaxis of

COVID-19[21].

Organotypic cultures of primary human nasal epithelium differentiated at air-liquid interface (ALI) are

a translationally-relevant primary cell model for studies of SARS-CoV-2 host-virus interaction[11], with

considerable potential to accelerate our understanding of pathogenesis. A small number of studies

using this model demonstrate that SARS-CoV-2 replicates efficiently in human nasal cells[11-13], yet

important questions concerning cellular tropism and their innate immune response remain

unresolved. Hou and colleagues report that only ciliated cells were permissive to SARS-CoV-2, despite

expression of ACE2 and TMPRSS2 by all cell types[12]. They hypothesised that post-entry factors, such

as innate immunity, might govern tropism. By contrast, Pizzorno and colleagues reported infection in

all major cell types (ciliated, secretory and basal cells)[13], consistent with prior indications from single

cell RNA sequencing (scRNA-seq) data and studies in lower airway models[22,23]. While an IFN response

to SARS-CoV-2 can be detected in nasal cells[11,13], in apparent contrast to bronchial or alveolar

epithelial cells[24-26], the kinetics of induction and the antiviral function of IFNs in nasal epithelium has

not been systematically characterised.

Here we employed a comprehensive range of techniques, including scRNA-seq and proteomics, in

primary human nasal ALI cultures to define: (i) cellular tropism; (ii) the innate immune response to

SARS-CoV-2; and (iii) the antiviral activity of IFN-I/III. We observed broad cellular tropism of SARS-CoV-

2 for nasal epithelial cells, although ciliated cells were the most permissive. Nasal cells mounted a

delayed IFN response that failed to exert control over viral replication. However, SARS-CoV-2

remained highly sensitive to IFN-restriction if exogenous IFN-I/III was applied prior to infection. These

data enrich our understanding of the interaction of SARS-CoV-2 and the human IFN system at the

earliest point of infection, with immediate therapeutic implications.

RESULTS

SARS-CoV-2 robustly infects primary differentiated nasal epithelial cultures

Primary nasal epithelial cultures were established from cryopreserved stocks from six adult donors,

obtained prior to the SARS-CoV-2 pandemic. Cells were expanded, differentiated and then matured

at ALI for 28 days, according to an established protocol[27]. We first sought to address their suitability

as a model of SARS-CoV-2 infection. Visualisation of scRNA-seq data from two representative donors

discerned five major populations of cells (Fig. 1A). These were ciliated, secretory and basal cells,

alongside two rarer populations of FOXN4+ cells and ionocytes. Cells expressed characteristic markers

(Fig. 1B) and corresponded broadly to *ex vivo* single-cell data from nasal biopsies[9,10]. Immunostaining

verified the presence of major cell types in these cultures - including acetylated alpha-tubulin-positive

(AAT) ciliated cells, mucin 5B-positive (MUC5B) secretory cells, and tumour protein 63-positive (TP63)

basal cells (Fig. 1C) - with ciliated cells being the most frequent cell population. Consistent with

published scRNA-seq data, mRNA for key SARS-CoV-2 entry receptors, *ACE2* and *TMPRSS2*, was

expressed, albeit at relatively low levels, alongside other genes implicated in possible alternate modes

of SARS-CoV-2 entry such as *FURIN* and *CTSL* (Fig. S1)[28]. Robust expression of ACE2 and TMPRSS2 at

the protein level was confirmed by immunoblotting of whole-cell lysates prepared from mature ALI

cultures (Fig. 1D). To establish their permissiveness to infection, nasal ALI cultures were inoculated at

the apical surface with a clinical SARS-CoV-2 isolate (England/2/20) at an approximate multiplicity of

infection (MOI) 0.1 - consistent with other studies (0.1-0.5)[11-13] - and monitored for infection over the

next 72 h. Expression of SARS-CoV-2 *nucleocapsid (N)* gene and spike (S) protein increased significantly

over time, indicative of viral replication (Fig. 1E-F). This was accompanied by the release of infectious

viral particles, as determined by plaque assay of apical washes on Vero E6 cells, confirming productive

infection (Fig. 1G). SARS-CoV-2 replication was accompanied by a progressive decline in epithelial

barrier integrity starting from 48 hours post-infection (hpi), reflecting virus-induced epithelial

dysfunction (Fig. 1H). These data established the suitability of the human nasal ALI model for

modelling SARS-CoV-2 infection.

Figure 1. **Robust SARS-CoV-2 infection in a primary differentiated nasal epithelial ALI culture model.**

(A) UMAP visualisation of scRNA-seq data in nasal ALI cultures revealed five major cell types. (B)

Expression of key markers distinguishing cell types. (C) Immunofluorescence analysis of secretory

(MUC5B), ciliated (AAT) and basal (TP63) cells in nasal ALI cultures (representative images from n=6

donors, scale bar = 20 μm), with frequency of cell type summarised in the bar plot. (D) Immunoblot

demonstrating ACE2 and TMPRSS2 expression by each donor, representative of n=3 experiments. (E)

RT-PCR analysis of SARS-CoV-2 nucleocapsid (*N*) gene expression in whole-cell lysate (WCL) (repeat

experiments in n=6 donors). (F) Immunoblot of SARS-CoV-2 spike (S) protein expression in WCL

(representative of repeat experiments in n=4 donors). (G) Production of infectious viral particles

determined by plaque assay of apical washings (repeat experiments in n=6 donors). Dotted line

represents lower limit of detection. (G) Transepithelial resistance (TEER) measurements upon

infection (expressed as % of mock-infected wells, mean of repeat experiments in n=6 donors). Mean

± SEM. * P < 0.05, **** P < 0.001 ANOVA with Dunnett's post-test correction, comparing to 0h (E) or

24h (G-H). MUC5B, mucin 5B; AAT, acetylated alpha tubulin; TP63, tumour protein 63.

Evidence of broad cellular tropism of SARS-CoV-2

To revisit the question of whether individual cell types are more or less permissive to SARS-CoV-2[12,13],

we first examined viral gene expression by scRNA-seq analysis at 24 hpi, selected to represent an early

stage in the progress of infection. Of the viral transcripts, the *N* gene was most abundantly expressed

(Fig. 2A). While all cell types expressed viral transcripts, there were notable differences both in the

proportion of cells infected, and the relative abundance of different viral transcripts within these cells.

A greater proportion of ciliated and secretory cells expressed viral transcripts compared to basal cells

(Fig. 2B). However basal cells are located away from the apical surface; physical inaccessibility to

apically-applied virus at this timepoint might at least partially account for this observation. To

investigate tropism further, we undertook immunofluorescence analysis of viral spike (S) protein

expression at 48 hpi. This analysis revealed broadly similar proportions of ciliated, secretory and basal

cells expressing S protein. However, the mean pixel intensity of S protein was significantly greater in

ciliated cells (Fig. 2C-D). To corroborate these findings, we undertook analysis of intracellular virion-

like structures (VLSs) at 48 hpi by transmission electron microscopy (TEM, Fig 2E), focusing on ciliated

and secretory cells. Intracellular VLSs were observed in both ciliated and secretory cells,

predominantly towards the apical surface (Fig. 2E). Consistent with immunofluorescence analysis of S

protein intensity, there was a significant increase in the number of VLSs per cell in ciliated compared

with secretory cells (Fig. 2E). Collectively, these data suggested that the virus is capable of entering,

and replicating in, all major nasal cell types.

Figure 2. Broad tropism of SARS-CoV-2 for nasal cells. (A) Dot plot of scRNA-seq data showing

magnitude (colour) and proportion (size) of cell types expressing viral transcripts. (B) Relative

proportion of infected cell types based on expression of viral spike (*S)* gene. (C) Quantification of cell-

type specific expression of viral S protein and S intensity at 48 hpi (repeat experiments in n=5 donors).

(D) Representative images of viral S protein expression in ciliated (AAT), secretory (MUC5B) and basal

cells (TP63) shown in (C). Scale bars = 10 μm. (E) Transmission electron micrograph (TEM) analysis of

SARS-CoV-2 infected ciliated and secretory cells at 48 hpi, with quantification of virion-like structures

in the bar plot. Inserts (a) and (b) display VLSs in ciliated and secretory cells respectively. Scale bars =

1 μm. Mean ± SEM ** P < 0.01, *** P < 0.001, **** P < 0.0001, ANOVA with Dunnett's post-test

correction (C) or t-test (E). E = envelope; M = matrix; N = nucleocapsid; S = spike.

Characterisation of nasal cell responses to SARS-CoV-2

Published *ex vivo* single-cell transcriptomic analyses report that nasal cells exhibit the basal expression

of an innate antiviral programme, in the absence of viral infection, characterised by several ISGs[9]. This

suggests that nasal epithelial cells may adopt a constitutive antiviral state. It was also hypothesised

that cellular innate immunity might influence nasal cell tropism of SARS-CoV-2[12].

In epithelial cells, ISGs are predominantly regulated by paracrine signalling mediated by IFN-I/IIIs. IFN-

I/IIIs are produced by virally infected cells, downstream of pattern-recognition receptor (PRR)

signalling, alerting their neighbours to viral infection. Responding cells adopt an antiviral state as a

consequence of the expression of hundreds of antiviral ISGs[29]. Receptors for both IFN-I (IFNAR) and

IFN-III (IFNLR) are expressed by epithelial cells. Ligand binding to these receptors activates a shared

JAK-STAT signalling pathway, culminating in ISG expression[15,29]. However, ISGs can also be regulated

by IFN-independent mechanisms. These include the direct action of constitutively expressed

transcription factors such as IRF3, activated downstream of PRRs[29], or by constitutive ISG expression

- a strategy employed by stem cells for antiviral defence[30].

We examined scRNA-seq data to characterise the innate antiviral response of nasal cells, both at

baseline and in response to SARS-CoV-2 infection at 24 hpi. For this analysis we distinguished cells in

three experimental conditions: unexposed (mock-infected); SARS-CoV-2-exposed but uninfected

(these cells would theoretically be exposed to IFNs and other paracrine signals, but not infected); and

SARS-CoV-2-infected (as defined by expression of *S* transcripts). At baseline, expression of a composite

ISG signature (Fig. 3A) and individual ISGs (Fig. 3B) was observed in unexposed cell types, consistent

with *ex vivo* scRNA-seq findings[9]. Considering that viral infection occurred in all cell types, these data

imply that relative differences in the basal ISG expression programmes did not substantially impact on

SARS-CoV-2 permissiveness, at least under the infection conditions studied here. We next considered

the involvement of paracrine IFNs. In SARS-CoV-2 infected cells, but not their uninfected neighbours,

we observed an enrichment of the ISG signature (Fig. 3A-B). This finding is more suggestive of direct

induction of ISGs in infected cells, rather than paracrine IFN-I/III signalling, at least at this timepoint.

Consistent with this, we observed expression of IFN-I (*IFNB, IFNK, IFNA5)* and IFN-III (*IFNL1-3)* in only

a small minority (0.1%) of infected ciliated cells, and very low levels in other nasal cell types (Fig. 3C).

Considering that over 50% of ciliated cells expressed viral transcripts at this timepoint, these data

suggest that SARS-CoV-2 efficiently avoided IFN induction in the vast majority of infected epithelial

cells, consistent with data in non-nasal airway epithelial cells[25,26]. A range of other chemokines and

cytokines were induced by infected epithelial cells, indicating the preserved capacity to detect viral

infection (Fig. S2).

Figure 3. Characterisation of innate IFN responses in nasal cells. (A) Average expression of interferon-stimulated gene (ISG) composite score signature genes in SARS-CoV-2 unexposed (light blue), SARS-CoV-2 exposed but uninfected (blue), and SARS-CoV-2 infected (purple) cell types at 24 hpi. Boxes represent the median value +/- 1.58*IQR/√(n) and whiskers represent 1.5*IQR. Outliers outside this range were trimmed. Differences between infected and other cell types were significant by FDR-adjusted ANOVA at P < 0.001. (B) Matrix plot demonstrating expression (colour) of individual ISGs

making up composite score in (A). (C) Dot plot of individual differentially-expressed IFNs by cell type

(normalised expression compared to uninfected cells, with colour representing mean expression and

size of the dot representing the proportion of cells expressing each gene).

Kinetics of innate IFN-I/IFN-IIIs response to SARS-CoV-2

To investigate the kinetics of the IFN-I/III response, expression of IFN-I (*IFNA1* and *IFNB*) and IFN-III

(*IFNL1*) was examined by RT-PCR at 6, 24, 48 and 72 hpi (Fig. 4A). Induction of *IFNL1* and *IFNB* was low

at 24 hpi, consistent with scRNA-seq findings, but increased significantly by 48 and 72 hpi. *IFNA1* was

not induced, as observed in our scRNA-seq data. Compared to the timing of initiation of viral gene

expression - which was detectable at 6 hpi and approached its maximum level by 24 hpi (Fig. 1E) - the

induction of IFNs appeared delayed, as suggested by previous studies[11,13]. Infection was accompanied

by progressive upregulation of proinflammatory cytokines such as *IL1B, IL6* and *TNF*, consistent with

initiation of a local inflammatory response (Fig. 4B). To confirm that the attenuated production of

IFNL1 and *IFNB* at early times was not dependent on MOI or due to an intrinsic defect of these cells,

infections were repeated at high MOI (~ 2). Infections at high MOI have been previously reported to

enhance the relatively inefficient IFN-I induction in response to SARS-CoV-2[25]. A preparation of Sendai

virus (SeV) - containing a high proportion of immunostimulatory defective viral genomes (DVGs) - was

used as a positive control. At 6 hpi, a time point at which *IFNB* and *IFNL1* were robustly induced by

SeV DVGs, there was no response to SARS-CoV-2 infection at high MOI (Fig. 4C), despite SARS-CoV-2

gene expression (Fig. 4C). Thus SARS-CoV-2 appears to efficiently evade PRR detection in nasal

epithelial cells. To look for evidence of a paracrine response to IFN-I/III, we undertook analysis of

expression of the ISGs *RSAD2* and *USP18* by RT-PCR and immunoblotting (Fig. 4D-E). This analysis

revealed an increase in ISG expression at later times (48-72 hpi), corresponding to the expression of

IFNB/*IFNL1*. This association is suggestive (although not causal evidence) of paracrine IFN-I/III

signalling. Collectively, these results indicate that nasal epithelial cells are capable of expressing IFN-

I/IIIs during SARS-CoV-2 infection, but that the response is delayed relative to viral replication.

Figure 4. Delayed induction of IFN-I/III signalling in SARS-CoV-2-infected nasal ALI cultures. (A) RT-

PCR analysis of expression of (A) *IFNB*, *IFNL1* and *IFNA1* and (B) *IL6*, *TNFA* and *IL1B* expression in whole-

cell lysates (repeat experiments in n=4 donors). (C) RT-PCR analysis of *IFNB*, *IFNL1* and SARS-CoV-2 *N*

gene expression in nasal ALI cultures exposed to SARS-CoV-2 (MOI = 2) or a DVG-rich stock of Sendai

virus (SeV) for 6 h (repeat experiments in n=2 donors). (D) RT-PCR and (E) immunoblot analysis of

USP18 and *RSAD2* expression in whole-cell lysates (repeat experiments in n=4 donors). ND, Not

detected. Mean ± SEM * P < 0.05, ** P < 0.01, *** P < 0.001 **** P < 0.0001, ANOVA with Dunnett's

post-test correction compared to 0h (A [*IFNB*]) or 24h (A [*IFNL1*], B, D).

IFN-signalling dominates the nasal host response to SARS-CoV-2 at the protein level

To validate and extend these findings, we undertook an unbiased assessment of the host response to

SARS-CoV-2 infection by proteomics analysis. Whole-cell lysates were prepared from SARS-CoV-2 and

mock-infected nasal ALI cultures from six donors at 72 hpi. Lysates were analysed by quantitative mass

spectrometry (Supplementary Dataset S1, quality control data in Fig. S3). Overall, this analysis

detected the differential expression (DE) of 180 proteins including viral proteins such as S, M, N,

ORF1AB, ORF3A and ORF8 (Fig. 5A). The most highly increased host protein was Sorting Nexin 33

(SNX33), an endosomal protein that has not yet been implicated in the life cycle of SARS-CoV-2.

Infected and uninfected cells clustered together by principal component analysis (Fig. 5B). Inspection

of the DE proteins confirmed a robust host innate immune response, dominated by ISG products (Fig.

5A). Functional annotation revealed an enrichment of antiviral response and especially IFN-I signalling

pathways (Fig. 5C, Table S1). These data are consistent with our earlier findings and contrary to prior

reports in cell lines or human bronchial/tracheal epithelial cultures, where a robust endogenous IFN-

I/III response to SARS-CoV-2 was not detected[24-26]. Key ISG proteins identified included IFITM1-3 and

the OAS cluster (OAS1-3), the latter associated with genetic susceptibility to severe COVID-19[17] (Fig.

5C). Significantly downregulated pathways were also identified, including TRIF-dependent toll-like

receptor signalling, as well as RNA polymerase II transcription and endosomal transport (Table S2).

This implied viral subversion of critical host functions, including host gene transcription, protein

 trafficking and viral sensing. Proteins involved in the maintenance of epithelial tight junctions were

 also downregulated, consistent with the loss of barrier integrity observed in earlier experiments (Fig.

 1H).

Figure 5. An ISG response dominates the proteome of SARS-CoV-2 infected nasal ALI cultures.

Differential proteomic profiling of SARS-CoV-2-infected nasal ALI cultures. Mass spectrometry-based proteomics was carried out on whole-cell lysates prepared at 72 hpi (n=6 donors per condition). (A) Volcano plot illustrating 180 differentially expressed proteins with increased (orange points) and decreased (purple points) expression in infected as compared to mock samples. Dotted red lines indicate those proteins with a fold change of < 1.5 and adjusted p values <0.05. (B) Principal component analysis of the whole proteome data set. (C) Functional annotation network of differentially expressed proteins.

Antiviral activity of IFN-I/III towards SARS-CoV-2 infection

Given the prominence of the IFN-I/III response in the proteome of SARS-CoV-2-infected cells, a key question was whether the IFN-I/III response had any impact on SARS-CoV-2 replication. To address this question, nasal ALI cultures were treated with the JAK inhibitor ruxolitinib (RUX). RUX antagonises signalling downstream of IFNAR and IFNLR, owing to the involvement of JAK1 in both signalling pathways. We reasoned that blocking paracrine IFN-I/III signalling would reveal its impact, if any, on SARS-CoV-2 replication. Cells were treated with 10 μM RUX (a dose optimised in prior experiments[31]) or vehicle control (DMSO) in the basal medium for 24 hours prior to infection. Nasal cultures were infected at the apical surface (MOI ~ 0.01) and inhibitors were refreshed every 24h. At 48 hpi, lysates were prepared and analysed by immunoblot. RUX treatment abolished expression of RSAD2 and USP18 in response to SARS-CoV-2 infection (Fig. 6A), indicating that induction of ISG protein products at this timepoint was largely dependent on paracrine IFN-I/III signalling, as previously suggested (Fig. 4D). However, viral S protein expression was not enhanced by RUX treatment (Fig. 6A), nor was release of infectious virus, as measured by plaque assay (Fig. 6B). These data suggested that the endogenous IFN-I/III response failed to impact SARS-CoV-2 replication up to 48 hpi, at least at the MOI tested. These data provided further evidence that SARS-CoV-2 triggers an endogenous paracrine IFN-I/III

response in nasal cells, but showed that this response was insufficient to contain SARS-CoV-2

replication.

An important follow-up question was whether nasal cells could mount an antiviral state to SARS-CoV-

2, providing IFN-I/III was delivered in a timely fashion. To address this, nasal ALI cultures were pre-

treated with exogenous IFNβ or IFNλ1 for 16h to induce an antiviral state, subsequently infected with

SARS-CoV-2 at MOI 0.01, and examined at 48 hpi. Analysis of infection by immunoblotting of whole-

cell lysates for spike (S) protein expression or plaque assay of apical washes demonstrated a significant

reduction in infection with either IFNβ or IFNλ1 pre-treatment (Fig. 6C-D). This was accompanied by

robust induction of antiviral ISG products (Fig. 6C), and preservation of barrier integrity (Fig. S4). It is

worth noting that the ISG expression induced in response to recombinant IFN-I/III was substantially

greater than that induced by SARS-CoV-2 infection (i.e. endogenous IFN-I/III production, Fig. 6C). Thus

exogenous IFN-I/III was capable of inducing in the nasal epithelium an antiviral state that potently

inhibited SARS-CoV-2 infection, providing it was delivered (a) prior to infection, and (b) at sufficient

concentration. This IFN-sensitivity of SARS-CoV-2 contrasts with the relative resistance of SARS-CoV[19].

These data suggest that mucosal delivery of IFNβ is a potential therapeutic strategy for SARS-CoV-2.

In clinical practice, IFNs are unlikely to be used prior to infection, unless this is part of a prophylactic

regimen. To determine the effectiveness of exogenously applied IFN-I once SARS-CoV-2 infection is

underway, cells were treated with IFNβ at 6 or 24 hpi. In this experiment, IFNβ treatment at 6 hpi

continued to impact on SARS-CoV-2 infection, whereas addition after 24 hpi had no effect (Fig. 6E-F).

Interestingly, ISG induction was still observed in response to IFNβ treatment at 24 hpi, albeit at

reduced magnitude (Fig. 6E). These data suggest that SARS-CoV-2 impairs, but does not abolish, JAK-

STAT signalling in infected cells, implying that recombinant IFNs may have a therapeutic role in

established SARS-CoV-2 infection, as recently shown in animal models[8] and in early phase clinical

trials[32].

Figure 6. Impact IFN-I/III signalling on SARS-CoV-2 infection. (A) Immunoblot of nasal ALI cultures

treated with ruxolitinib (RUX, 10 µM) or DMSO vehicle for 24 h prior to infection. Whole-cell lysates

were prepared at 48 hpi. Results representative of experiments in n=4 donors (* = nonspecific band).

(B) Plaque assay showing no increase in infectious particle release at 24 hpi and 48 hpi (same

experimental conditions as A; n=6 donors). (C) Immunoblot of whole-cell lysates prepared from nasal

ALI cultures at 48 hpi (representative of n=4 donors). Cultures were pre-treated for 16 h prior to

infection with IFNβ (1000 IU/mL) or IFNλ1 (100 ng/mL). (D) Plaque assay showing significant reduction

in infectious particle release at 48 hpi if pre-treated with IFNβ (1000 IU/mL) or IFNλ1 (100 ng/mL)

(same experimental conditions as C; n=5 donors). (E) Immunoblot of whole-cell lysates prepared at 48

hpi. Nasal ALI cultures were either pre-treated with IFNβ (1000 IU/mL) for 16h prior to infection with

SARS-CoV-2 or IFNβ (1000 IU/mL) treatment was applied at 6 or 24 hpi. Results representative of

experiments in n=2 donors. (F) Plaque assay on supernatants collected from E (48 hpi). Mean $\pm$ SEM.

**** P < 0.001 n.s. = non-significant. ANOVA with Dunnett's post-test correction compared to vehicle

(B) or untreated control (F).

DISCUSSION

We report the most comprehensive characterisation of the human nasal epithelial response to SARS-

CoV-2 to date, revealing a response dominated by IFN-I/IIIs and their downstream ISG products. While

this is apparently insufficient to contain SARS-CoV-2, recombinant IFN-I/III treatment potently blocked

SARS-CoV-2 replication, suggesting that mucosal delivery of IFNs could be a promising strategy for

post-exposure prophylaxis.

The nasal mucosa is likely to be a main point of entry of SARS-CoV-2. Prior single-cell transcriptomic

studies implied an abundance of target cells in the nasal mucosa and further suggested that they may

be poised to mount an antiviral response[9]. Yet few studies to date have characterised SARS-CoV-2

replication in primary human differentiated nasal cells[11-13], and ours is the first to analyse the host-

virus interaction comprehensively, at single-cell resolution. Our findings indicate that the host

response to SARS-CoV-2 in nasal epithelium is dominated by IFN-I/III, albeit this response is kinetically

delayed. These results are consistent with evidence of IFN-I/III induction in nasal swabs from patients

with COVID-19[33]. However, we found that blockade of the endogenous IFN response did not impact

on SARS-CoV-2 infection. At first glance, this result may appear paradoxical. However, the probable

explanation is the delay of induction of IFN-I/III relative to virus replication. IFN-I/III expression

became detectable 36-48 h after the onset of SARS-CoV-2 replication - at a point at which infectious

virion production was reaching its maximum, and over half of cells in the culture were already infected

- suggesting the production of IFNs was too late to have a meaningful impact on the overall spread of

infection. Consistent with this, there was evidence of downregulation of PRR signalling in the

proteome of infected cells. We conclude that: (i) SARS-CoV-2 appears capable of evading PRR

signalling in nasal epithelial cells in the early stages of viral replication; and (ii) the endogenous IFN-

I/III response of nasal epithelial cells comes effectively "too little, too late" to inhibit SARS-CoV-2 in

the nasal epithelium. The expression of various IFN evasion proteins[34] and the sequestration of viral

replication machinery within cytosolic vesicles[35] underlie the capacity of SARS-CoV-2 to evade early

detection by PRRs, as indicated by our proteomics data. However, an important question is what

molecular patterns are responsible for IFN-I/III induction at later times. Future studies should address

the relative contribution of host damage-associated molecular patterns versus viral pathogen-

associated molecular patterns (e.g. defective viral genomes) accumulating during replication.

Our finding that exogenous IFN-I or IFN-III can potently induce an antiviral state in nasal cells is

consistent with its apparent protective effects in patients[16-18] and in early phase clinical trials[32]. The

main limitation of our data in this nasal epithelial culture system is that it did not account for

professional immune cells present in the nasal mucosa, for example plasmacytoid dendritic cells[36],

which are capable of more rapidly mounting an IFN-I/III response to SARS-CoV-2[37], potentially tipping

the scales in favour of the host. We studied cells derived from adult donors, however it is possible that

nasal cells from paediatric donors may behave differently in terms of their permissiveness to SARS-

CoV-2 and/or the efficiency of their innate immune response. Furthermore, SARS-CoV-2 variants with

mutations in the spike gene have emerged worldwide whilst we were undertaking the experiments

described here; these variants may impact viral replication and/or host immunity, and should be

included in future studies.

Nevertheless, our data, employing a variety of complementary methods, indicate that SARS-CoV-2 has

a relatively broad tropism for nasal epithelial cells, confirming suggestions from prior scRNA-seq

studies[7,9] and other *in vitro* studies of primary nasal[13] and tracheobronchial cells[22]. These findings

contrast with the results of Hou and colleagues, who reported exclusive tropism of SARS-CoV-2 for

ciliated cells in the airway[12]. It is not immediately clear how to reconcile these findings, given that

secretory cells express relevant entry receptors[12]. Hou and colleagues used a fluorescent reporter

virus, the tropism of which might have been slightly narrower than clinical isolates. It is also worth

noting our findings show that while all cell types contained SARS-CoV-2 protein, the intensity of

immunodetection was greater in ciliated cells, which also contained more virion-like structures per

cell. This implies that although all cell types are permissive, there may also be quantitative differences

in the efficiency of viral replication in different cell types. Hou and colleagues previously hypothesised

that post-entry factors such as antiviral immunity might dictate permissiveness. We found no evidence

to support this hypothesis, since while all nasal epithelial cells demonstrated a basal ISG signature -

consistent with *ex vivo* nasal biopsy data[9] - this did not vary substantially between cell types, and was

apparently insufficient to mediate resistance to SARS-CoV-2.

The observation that IFN-I treatment prevents SARS-CoV-2 infection indicates that chemoprophylaxis

with IFN-I may have therapeutic value. This approach has already been tested in a small clinical trial

in China (although the absence of a control group makes it impossible to judge the efficacy of this

approach[38]). Immunisation is the most tractable approach for large-scale primary prevention of

COVID-19. However, owing to incomplete vaccine coverage and/or effectiveness against mildly

symptomatic or asymptomatic infection, and the emergence of variants that may compromise vaccine

efficacy, there will likely continue to be a need for targeted chemoprophylactic therapies to prevent

transmission in specific circumstances. These include post-exposure prophylaxis of contacts - to avoid

the need for self-isolation - as well as pre-exposure prophylaxis for certain high-risk encounters (e.g.

in healthcare settings or prior to long-distance travel). Our data suggest that application of IFNβ to

the nasal mucosa might have an important role to play in this setting and argue for urgent clinical

assessment of this approach. In terms of the therapeutic efficacy of IFNβ in patients with COVID-19,

our findings suggest that early administration will likely be a key factor determining its efficacy.

METHODS

Adult nasal airway epithelial cell culture at air-liquid interface (ALI)

Adult primary human nasal airway epithelial cells were derived from excess clinical material obtained

during routine nasal surgical procedures, processed, differentiated and validated as previously

described[27] with the following modifications: cells were cryopreserved following expansion, then re-

animated and differentiated at ALI in PneumaCult-ALI-S (Stemcell Technologies). The sex and age of

donors are included in Table S3. Ethical approval for sample collection was provided (Research Ethics

Committee Reference 17/NE/0361).

Viruses, cytokines and inhibitors

A clinical isolate of SARS-CoV-2 (England/2/2020) was obtained from Public Health England (PHE). The initial stock was propagated once in Vero E6 cells. The same viral stock was used for all experiments. Sendai virus (Cantell Strain) was obtained from Richard Randall (St Andrew's University). For nasal ALI infections, apical poles were gently washed once with warm Dulbecco's modified Eagle's medium (DMEM; Gibco, USA) and then infected with 60 µL dilution of virus in DMEM, at a MOI of between 2 and 0.01 plaque-forming units per cell for 2 hours, when the virus-containing medium was removed. DMEM was used as inoculum for mock infection. Apical washes (in warm phosphate-buffered saline) were collected at different time-points and stored at -80°C for plaque assays. Plaque assays were undertaken in Vero E6 cells using a 1.2% (w/v) microcrystalline cellulose overlay (Sigma-Aldrich). Cytokines/inhibitors were used at the following concentrations: human recombinant IFNβ1 (1000 ng/mL; Avonex, Biogen Inc, USA); IFNλ1 (100 ng/mL; R&D Systems, USA); and Ruxolitinib (10 µM; Calbiochem, USA) alongside the appropriate dilution of DMSO vehicle. Treatment was applied through basolateral poles.

Single cell RNA sequencing (scRNA-seq)

For droplet-encapsulation cells were processed as described in the supplementary information (SI). Principal component (PC) analysis was undertaken to find the first 20 PCs, which were batch-adjusted using Harmony and used to generate the nearest-neighbour graph. Dimensionality reduction and embedding was performed using Uniform Manifold Approximation and Projection (UMAP), with the neighbourhood graph clustered using the Leiden algorithm. The Wilcoxon Rank Sum test was used to identify differentially expressed genes between clusters, and these were annotated based on expression of markers used in the literature to define populations. A published gene set was used to generate a list of type I ISGs (www.gsea-msigdb.org/gsea/msigdb/cards/GO_RESPONSE_TO_TYPE_I_INTERFERON). This was used to quantify the interferon response within identified cell populations using the AddModuleScore tool in Seurat.

Infected cells were assigned as those cells with detectable SARS-CoV-2 spike RNA ('covid—S'

transcript). Further details are included in the SI.

Proteome sample preparation

The protein concentration was determined by EZQ® protein quantification assay. Protein digestion was

performed using the S-Trap™ sample preparation method and TMT-16 plex labelling was carried out

as per the manufacturer's instructions. Samples were cleaned using MacroSpin columns, and dried

down prior to offline high performance liquid chromatography fractionation. Peptides were

fractionated on a Basic Reverse Phase column on a Dionex Ultimate 3000 off-line LC system. A total of

18 fractions were collected, and each fraction was acidified and dried. Peptides were dissolved in 5%

formic acid, and each sample was independently analysed on an Orbitrap Fusion Lumos Tribrid mass

spectrometer, connected to an UltiMate 3000 RSLCnano System. All spectra were analysed using

MaxQuant 1.6.10.43 and searched against SwissProt *Homo sapiens* and Trembl SARS-CoV-2 FASTA

files. Reporter ion MS3 was used for quantification and the additional parameter of quantitation labels

with 16 plex TMT on N-terminus or lysine was included. A protein and peptide false discovery rate

(FDR) of less than 1% was employed in MaxQuant. Moderated *t*-tests, with patient accounted for in

the linear model, was performed using Limma, where proteins with an adjusted $P < 0.05$ were

considered as statistically significant. All analysis was performed using R. Raw data are present in

Supplementary Dataset 1. A comprehensive description of the methods can be found in the SI.

Quantitative RT-PCR, immunoblotting and immunofluorescence

RT-PCR and immunoblot were done as previously described[31], with further detail in the SI.

Membranes were fixed *in situ* with 4% formaldehyde overnight at 4°C, before sectioning.

Immunofluorescent staining was performed in accordance with published methods[39], with

modifications as summarised in the SI. Images were captured using a Nikon A1 confocal microscope

(Nikon, Japan) and analysed via Fiji (Version 2.0) using the Cell Counter plugin. Primers and probes

(Table S4) and antibodies (Table S5) are described in the SI.

Transmission electron microscopy (TEM)

ALI cultures were fixed with 2% glutaraldehyde (Sigma-Aldrich, MO, USA) with 0.1 M sodium

cacodylate (pH 7.4) buffer overnight at 4°C. TEM resin processing was performed as described

previously[40]. Additional details, including embedding, sectioning and analysis, are included in the SI.

Statistical analysis

Statistical analysis was performed and figures assembled using GraphPad Prism V8 (GraphPad

Software, USA). Data are presented as mean ± SEM of individual donor values (derived typically from

2-3 independent repeat experiments per donor). The donor was used as the unit of experiment for

statistical analysis purposes. Continuous data were normalised or log-transformed prior to analysis

using parametric significance tests. Differences between two groups were compared using an

unpaired, two-tailed Student's *t*-test, whereas differences between more than two groups used

ANOVA, with Dunnett's post-test correction for multiple comparisons. A two-tailed alpha of < 0.05

was the threshold for statistical significance. Statistical analysis of proteomics and transcriptomics

data sets is described in the relevant sections above.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium[41]

via the PRIDE partner repository[42] with the dataset identifier PXD021026. Reviewers can access it

through the Username: reviewer_pxd022523@ebi.ac.uk and the Password: UaEXYFKF. RNA

sequencing data have been deposited to the European Genome-phenome archive (accession

pending). Additional raw data are available on request from the corresponding author.

Acknowledgements

We acknowledge Public Health England for providing the SARS-CoV-2 isolate and Professor R Randall (St Andrew's University) for providing Sendai virus and the Vero E6 cell line. We thank A Khan, M Glanville and S Dainty (Newcastle University Infectious Disease Facility) and A Laude (Bioimaging facility) for assistance. The work was part-funded by the Barbour Foundation (CJAD, SH, MB, MT, MH, CW), the UK-Coronavirus Immunology Consortium (CJAD, SH, MH) and the Medical Research Council SHIELD antimicrobial resistance consortium (JS, AJS). CFH is supported by a Medical Research Council studentship (MR/NO13840/1) and MB by an MRC Clinician Scientist Fellowship (MR/M008797/1). MT and SH are funded by a Wellcome Trust Investigator Awards (215542/Z/19/Z and 207556/Z/17/Z). CJAD and GR are Wellcome Trust Clinical Research Career Development Fellows (211153/Z/18/Z and 214539/Z/18/Z). MED is a Marie Sklodowska Curie Fellow within the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No. 890296. MH is funded by Wellcome (WT107931/Z/15/Z), The Lister Institute for Preventive Medicine and Newcastle NIHR Biomedical Research Centre (BRC). JPG, CW and BV were supported by the Medical Research Foundation (MRF Respiratory Diseases Research Award to JPG; Grant MRF-091-0001-RGGARNE) and Boehringer Ingelheim. TEM work was supported by a BBSRC Alert17 grant (BB/R013942/1). Professor Simpson is a National Institute for Health Research (NIHR) Senior Investigator. The views expressed in this article are those of the author(s) and not necessarily those of the NIHR, or the Department of Health and Social Care.

Author contributions

Conceived the study: MB, CW, CJAD with MH, SH, MT. Experimental design: MH, SH, MB, CW, MT, GR, CJAD. Nasal model development and patient material: IJH, BV, JS, JPG, SC, AJS, MB, CW. Virology data generation, analysis and interpretation: CFH, BJT, JSS, FG, AIG, TD, SH, CJAD. Proteomics data generation, analysis and interpretation: MED, SM, MT. Single-cell sequencing data generation, analysis and interpretation: RAB, ES, RH, JC, MH, GR. Supervised research: AJS, MH, SH, MB, CW, MT,

GR, CJAD. Drafted the manuscript: CFH and CJAD with ES, BV, MED, TD, MT, GR. Revised the

manuscript: RAB, MED, IJH, JSS, AJS, MH, SH, CW. Approved the manuscript for submission: all

authors.

References

1   Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*
382, 727-733, doi:10.1056/NEJMoa2001017 (2020).
2   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by
a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278, doi:10.1016/j.cell.2020.02.052
(2020).
3   Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan,
China. *Lancet* **395**, 497-506, doi:10.1016/S0140-6736(20)30183-5 (2020).
4   Knight, S. R. *et al.* Risk stratification of patients admitted to hospital with covid-19 using the
ISARIC WHO Clinical Characterisation Protocol: development and validation of the 4C
Mortality Score. *BMJ* **370**, m3339, doi:10.1136/bmj.m3339 (2020).
5   Group, R. C. *et al.* Dexamethasone in Hospitalized Patients with Covid-19 - Preliminary Report.
N Engl J Med, doi:10.1056/NEJMoa2021436 (2020).
6   Cevik, M., Kuppalli, K., Kindrachuk, J. & Peiris, M. Virology, transmission, and pathogenesis of
SARS-CoV-2. *BMJ* **371**, m3862, doi:10.1136/bmj.m3862 (2020).
7   Chen, M. *et al.* Elevated ACE-2 expression in the olfactory neuroepithelium: implications for
anosmia and upper respiratory SARS-CoV-2 entry and replication. *Eur Respir J* **56**,
doi:10.1183/13993003.01948-2020 (2020).
8   Hoagland, D. A. *et al.* Leveraging the antiviral type I interferon system as a first line of defense
against SARS-CoV-2 pathogenicity. *Immunity*, doi:10.1016/j.immuni.2021.01.017 (2021).
9   Sungnak, W. *et al.* SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells
together with innate immune genes. *Nat Med* **26**, 681-687, doi:10.1038/s41591-020-0868-6
(2020).
10   Ziegler, C. G. K. *et al.* SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human
Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell* **181**, 1016-
1035 e1019, doi:10.1016/j.cell.2020.04.035 (2020).
11   Gamage, A. M. *et al.* Infection of human Nasal Epithelial Cells with SARS-CoV-2 and a 382-nt
deletion isolate lacking ORF8 reveals similar viral kinetics and host transcriptional profiles.
PLoS Pathog **16**, e1009130, doi:10.1371/journal.ppat.1009130 (2020).
12   Hou, Y. J. *et al.* SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the
Respiratory Tract. *Cell* **182**, 429-446 e414, doi:10.1016/j.cell.2020.05.042 (2020).
13   Pizzorno, A. *et al.* Characterization and Treatment of SARS-CoV-2 in Nasal and Bronchial
Human Airway Epithelia. *Cell Rep Med* **1**, 100059, doi:10.1016/j.xcrm.2020.100059 (2020).
14   Onabajo, O. O. *et al.* Interferons and viruses induce a novel truncated ACE2 isoform and not
the full-length SARS-CoV-2 receptor. *Nat Genet* **52**, 1283-1293, doi:10.1038/s41588-020-
00731-9 (2020).
15   Duncan, C. J. A., Randall, R. E. & Hambleton, S. Genetic Lesions of Type I Interferon Signalling
in Human Antiviral Immunity. *Trends Genet* **37**, 46-58, doi:10.1016/j.tig.2020.08.017 (2021).
16   Zhang, Q. *et al.* Inborn errors of type I IFN immunity in patients with life-threatening COVID-
19. *Science* **370**, doi:10.1126/science.abd4570 (2020).
17   Pairo-Castineira, E. *et al.* Genetic mechanisms of critical illness in Covid-19. *Nature*,
doi:10.1038/s41586-020-03065-y (2020).
18   Bastard, P. *et al.* Autoantibodies against type I IFNs in patients with life-threatening COVID-
19. *Science* **370**, doi:10.1126/science.abd4585 (2020).
19   Lokugamage, K. G. *et al.* Type I Interferon Susceptibility Distinguishes SARS-CoV-2 from SARS-
CoV. *J Virol* **94**, doi:10.1128/JVI.01410-20 (2020).
20   Felgenhauer, U. *et al.* Inhibition of SARS-CoV-2 by type I and type III interferons. *J Biol Chem*
295, 13958-13964, doi:10.1074/jbc.AC120.013788 (2020).
21   Lee, J. S. & Shin, E. C. The type I interferon response in COVID-19: implications for treatment.
Nat Rev Immunol **20**, 585-586, doi:10.1038/s41577-020-00429-3 (2020).

22   Zhu, N. *et al.* Morphogenesis and cytopathic effect of SARS-CoV-2 infection in human airway
epithelial cells. *Nat Commun* **11**, 3910, doi:10.1038/s41467-020-17796-z (2020).
23   Fiege, J. K. *et al.* Single cell resolution of SARS-CoV-2 tropism, antiviral responses, and
susceptibility to therapies in primary human airway epithelium. *PLoS Pathog* **17**, e1009292,
doi:10.1371/journal.ppat.1009292 (2021).
24   Vanderheiden, A. *et al.* Type I and Type III Interferons Restrict SARS-CoV-2 Infection of Human
Airway Epithelial Cultures. *J Virol* **94**, doi:10.1128/JVI.00985-20 (2020).
25   Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives Development of
COVID-19. *Cell* **181**, 1036-1045 e1039, doi:10.1016/j.cell.2020.04.026 (2020).
26   Huang, J. *et al.* SARS-CoV-2 Infection of Pluripotent Stem Cell-derived Human Lung Alveolar
Type 2 Cells Elicits a Rapid Epithelial-Intrinsic Inflammatory Response. *bioRxiv*,
doi:10.1101/2020.06.30.175695 (2020).
27   Mavin, E. *et al.* Real-time measurement of cellular bioenergetics in fully differentiated human
nasal epithelial cells grown at air-liquid-interface. *Am J Physiol Lung Cell Mol Physiol* **318**,
L1158-L1164, doi:10.1152/ajplung.00414.2019 (2020).
28   Xiu, S. *et al.* Inhibitors of SARS-CoV-2 Entry: Current and Future Opportunities. *J Med Chem*
63, 12256-12274, doi:10.1021/acs.jmedchem.0c00502 (2020).
29   Randall, R. E. & Goodbourn, S. Interferons and viruses: an interplay between induction,
signalling, antiviral responses and virus countermeasures. *J Gen Virol* **89**, 1-47,
doi:10.1099/vir.0.83391-0 (2008).
30   Wu, X. *et al.* Intrinsic Immunity Shapes Viral Resistance of Stem Cells. *Cell* **172**, 423-438 e425,
doi:10.1016/j.cell.2017.11.018 (2018).
31   Duncan, C. J. A. *et al.* Severe type I interferonopathy and unrestrained interferon signaling due
to a homozygous germline mutation in STAT2. *Sci Immunol* **4**,
doi:10.1126/sciimmunol.aav7501 (2019).
32   Monk, P. D. *et al.* Safety and efficacy of inhaled nebulised interferon beta-1a (SNG001) for
treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-controlled, phase 2
trial. *Lancet Respir Med*, doi:10.1016/S2213-2600(20)30511-7 (2020).
33   Antonelli, G. *et al.* Type I interferons can be detected in respiratory swabs from SARS-Cov-2
infected patients. *J Clin Virol* **128**, 104450, doi:10.1016/j.jcv.2020.104450 (2020).
34   Lei, X. *et al.* Activation and evasion of type I interferon responses by SARS-CoV-2. *Nat Commun*
11, 3810, doi:10.1038/s41467-020-17665-9 (2020).
35   Wolff, G., Melia, C. E., Snijder, E. J. & Barcena, M. Double-Membrane Vesicles as Platforms for
Viral Replication. *Trends Microbiol* **28**, 1022-1033, doi:10.1016/j.tim.2020.05.009 (2020).
36   Hartmann, E. *et al.* Analysis of plasmacytoid and myeloid dendritic cells in nasal epithelium.
Clin Vaccine Immunol **13**, 1278-1286, doi:10.1128/CVI.00172-06 (2006).
37   Onodi, F. *et al.* SARS-CoV-2 induces human plasmacytoid pre-dendritic cell diversification via
UNC93B and IRAK4. *bioRxiv*, doi:10.1101/2020.07.10.197343 (2021).
38   Meng, Z. *et al.* An experimental trial of recombinant human interferon alpha nasal drops to
prevent COVID-19 in medical staff in an epidemic area. *MedRxiv*,
doi:doi.org/10.1101/2020.04.11.20061473 (2020).
39   Crowe, A. R. & Yue, W. Semi-quantitative Determination of Protein Expression using
Immunohistochemistry Staining and Analysis: An Integrated Protocol. *Bio Protoc* **9**,
doi:10.21769/BioProtoc.3465 (2019).
40   Faitg, J., Davey, T., Turnbull, D. M., White, K. & Vincent, A. E. Mitochondrial morphology and
function: two for the price of one! *J Microsc* **278**, 89-106, doi:10.1111/jmi.12891 (2020).
41   Deutsch, E. W. *et al.* The ProteomeXchange consortium in 2017: supporting the cultural
change in proteomics public data deposition. *Nucleic Acids Res* **45**, D1100-D1106,
doi:10.1093/nar/gkw936 (2017).

42   Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019: improving
support for quantification data. *Nucleic Acids Res* **47**, D442-D450, doi:10.1093/nar/gky1106
(2019).
632