Chromatin accessibility regulates intron retention in a cell type-specific manner

Veronika Petrova[1,2], Renhua Song[3,4], DEEP Consortium, Karl J.V. Nordström[5], Jörn Walter[5], Justin J.-L. Wong[3,4], Nicola J. Armstrong[6], John E.J. Rasko[2,4,7]*†, Ulf Schmitz[1,2,4]*†

[1] Computational BioMedicine Laboratory Centenary Institute, The University of Sydney, Camperdown 2050, Australia

[2] Gene and Stem Cell Therapy Program Centenary Institute, The University of Sydney, Camperdown 2050, Australia

[3] Epigenetics and RNA Biology Program Centenary Institute, The University of Sydney, Camperdown 2050, Australia

[4] Faculty of Medicine and Health, The University of Sydney, Camperdown 2050, Australia

[5] Laboratory of EpiGenetics, Saarland University, Campus A2 4, 66123 Saarbrücken, Germany

[6] Mathematics and Statistics, Curtin University, Bentley, WA 6102, Australia

[7] Cell and Molecular Therapies, Royal Prince Alfred Hospital, Camperdown 2050, Australia

*To whom correspondence should be addressed:

Dr. Ulf Schmitz; email: u.schmitz@centenary.org.au or Prof. John Rasko, email: j.rasko@centenary.org.au

†These authors should be regarded as joint last authors.

Abstract

Background: The phenomenon of widespread and dynamic intron retention (IR) programs in cells of vertebrate species has recently gained increasing attention. It has been shown that IR is involved in a multitude of cell-physiological processes, while aberrant IR profiles have been associated with numerous human diseases including several cancers.

Gap: Despite consistent reports about intrinsic sequence features that predispose introns to become retained, conflicting findings about cell type or condition-specific IR regulation by *trans*-regulatory and epigenetic mechanisms demand an unbiased and systematic analysis of IR in a controlled experimental setting.

Methods: We integrated matched mRNA sequencing (RNA-seq), whole genome bisulfite sequencing (WGBS), nucleosome occupancy methylome sequencing (NOMe-Seq), and chromatin immunoprecipitation sequencing (ChIP-seq) data from primary human myeloid and lymphoid cells. Using these multi-omics data and machine learning we trained two complementary models to determine the role of epigenetic factors in the regulation of IR in cells of the innate immune system.

Results: Our results show that intrinsic characteristics are key for introns to evade splicing and that some epigenetic marks are associated with IR. However, cell type-specific IR profiles are largely marked by changes in chromatin accessibility, whereby predisposed introns in permissive chromatin regions are more likely to be retained.

Conclusion: This study demonstrates the important role of chromatin architecture in IR regulation. Our results have profound implications for the analysis of other forms of alternative splicing as well. Since an increasing number of studies describe pathogenic alterations in splicing regulation and novel therapeutic approaches that target aberrant splicing, our findings will inform the development of novel epigenetic therapies.

Keywords: nucleosome occupancy, intron retention, epigenetics, alternative splicing, histone marks, CpG methylation, chromatin accessibility

Introduction

The role of introns in mammalian genomes remains largely unexplained. Given the time and energy required for the transcription and subsequent excision of introns from pre-mRNA, it was important to recognise in recent years that introns can be selectively retained in mature mRNA transcripts and thereby contribute significantly to transcriptomic complexity [1,2]. Intron retention (IR) is a form of alternative splicing that was assumed to occur due to the failure of the spliceosome to excise an intron from a pre-mRNA transcript. However, growing evidence suggests that IR is highly regulated by multiple complementary factors [3].

IR is widespread across all human tissues and affects more than 80% of protein-coding genes [4]. For example, dynamic IR profiles have been identified in key genes involved in hematopoietic cell differentiation and activation [1,5-8]. Fates of intron-retaining transcripts can be diverse and include (i) nonsense-mediated decay triggered by intronic premature termination codons, (ii) detention in the nucleus or nuclear degradation, and (iii) translation into alternative protein isoforms or creation of neoepitopes [3,9,10]. A better understanding of how IR is regulated is crucial to determine factors leading to aberrant IR, which has been associated with multiple diseases including cancer [11-13]

Despite numerous studies that describe the role of retained introns in key biological functions in animals and in human diseases [3,9,11], a comprehensive understanding of their regulation is still lacking. Retained introns have conserved intrinsic characteristics such as a higher GC content, shorter lengths, and weaker splice sites in comparison to their non-retained counterparts [2,3,14]. These features predispose introns to retention but cannot explain the dynamic IR profiles observed in numerous biological processes.

The regulation of alternative splicing has been the focus of many studies. Evidence suggests that alternative splicing is regulated at least at two levels: (i) locally, where *trans*-acting splicing regulators interact with *cis*-acting regulatory elements, and (ii) globally, through the structure of chromatin, which is largely governed by epigenetic factors, including nucleosome assembly, histone modifications and CpG methylation [15].

Previous reports have shown that, apart from intrinsic sequence-based features, intron expression can be regulated through (i) *cis*-regulatory elements, such as sequence motifs attracting *trans*-acting splicing-regulatory RNA binding proteins [4], (ii) core components of the splicing machinery [1], and (iii) change in the RNA Pol II elongation rate [16]. Moreover, an increasing number of studies have found links between epigenetic profiles and IR; reporting that IR is associated with reduced CpG methylation [8,17-19] and various histone modifications [20,21]. However, these reports have typically established the association of IR with only one epigenetic factor at a time. In general, the question of whether there are dominant epigenetic factors that underpin IR regulation remain unanswered.

In the quest to find a splicing regulatory 'code' several studies have used machine learning methods to train models that predict exon usage with increasing precision [22,23]. Moreover, some models were developed to predict cryptic splicing events caused by genetic variations and to link these to human diseases [24-26]. However, the computational prediction of IR events has not been attempted to date and the role of epigenetic marks has rarely been considered in computational models of splicing regulation [3,27].

In this study, to sought to systematically elucidate the role of epigenetic marks in the regulation of IR. We analysed genome-wide profiles of 6 histone modifications, CpG methylation and nucleosome occupancy at single-base resolution in primary lymphoid and myeloid cells. Using machine learning, we developed two models that predict IR in primary human immune cells. More specifically, we trained a logistic regression with elastic net (EN) classifier and a conditional Random Forest (RF) classifier with matched transcriptomics and epigenomics data from monocytes, macrophages, naïve T-cells, T-central memory, and T-effector memory cells (Figure 1).

92 Our results show that intrinsic characteristics are key for introns to evade splicing and that epigenetic
93 marks may modulate IR levels in a cell type-specific manner, where the dominant factor for dynamic
94 IR regulation is chromatin organisation.

Results

Intrinsic features of retained introns are consistent across cell types

To investigate how IR is regulated in primary immune cells (CD4+ T-cells, monocytes, and
macrophages), we integrated transcriptomics (mRNA-Seq) data with epigenomics data including
genome-wide CpG methylation (WGBS), histone modifications (ChIP-Seq), and nucleosome
occupancy (NOMe-Seq) (Table S1). The cells were isolated from peripheral blood of 2 healthy donors,
except for the monocyte-derived macrophages. Using the IR identification software IRFinder (4), we
quantified IR events of expressed genes (FPKM>1) in five cell types across myeloid and lymphoid
cells, representing two modes of differentiation: monocyte-to-macrophage differentiation and naïve T-
cell differentiation into central memory (CM) and effector memory (EM) T-cells. Introns that were
present in at least 10% of a gene's mature mRNA transcripts ($IR_{ratio} \geq 0.1$) with an overall intron depth
≥ 10 were considered retained. Non-retained introns were defined as those with an $IR_{ratio} \leq 0.01$ and
intron depth < 10.

We identified a total of 26,147 retained introns in 12,379 genes, some of which were retained in both
myeloid and lymphoid cells while others were cell type-specific (Figure S1A). Consistent with previous
reports, retained introns in our dataset are shorter in length, exhibit a higher GC content and weaker
splice site strengths compared to non-retained introns (Figure S1B-E). Our analysis revealed diverse
splicing patterns in myeloid and lymphoid cells. While 40% of the retained introns in myeloid cells
were significantly differentially retained ($\Delta IR \geq 0.1$; $p < 0.05$ Audic-Claverie test) between monocytes
and macrophages (571/1425), T cells displayed greater stability in regard to IR with only 8% of introns
classified as differentially retained (146/1812 in naïve T vs CM, and 80/969 in CM vs EM). In contrast
to the monocyte-to-macrophage differentiation, where we observed a reduction in IR events (Figure
2A), the overall number of retained introns remained consistent in all CD4+ T cells. These patterns
coincide with fewer changes in gene expression during T cell differentiation in contrast to major gene
expression changes in monocyte-to-macrophage differentiation (Figure S1F).

Most retained introns in our analysis overlapped with histone marks (HM) or with a nucleosome free
region (NFR, predicted from NOMe-seq data) around their 5′ and 3′ splice sites (+/100 bp) as well as
the middle of an intron (Figure S2A). Interestingly, many non-retained introns (~50%) lacked such
epigenetic marks in lymphoid cells (as opposed to only 20-30% of retained introns). H3K36me3 was
the most frequently observed histone modification followed by NFR peaks. In retained introns, between
30% and 60% of H3K36me3 signals were classified as strong (see Methods), whilst in non-retained
introns the proportion of overlap with the regions of strong signal ranged between 2% and 18%. Again,
the patterns of signal strength varied between the cell types (Table S3).

CpG methylation profiles (extracted from WGBS data) for retained and non-retained introns displayed
a characteristic bimodal distribution with two distinct peaks at 0% and 100%. Differential methylation
was predominantly found at the splice sites when we compared regions of genomic DNA associated
with IR and no IR. At the 5′ splice sites, we observed higher methylation levels in retained compared
to non-retained introns in all five cell types. However, this trend was reversed in the lymphoid cells at
the 3′ splice sites and in the middle of introns (Figure S2B).

M.CviPI enzyme, used in NOMe-seq experiment, methylates cytosine dyads in GC sequence and GCH
methylation levels (where H is any nucleobase except guanine) provide information about chromatin
accessibility. Unlike endogenous CpG methylation, GC dinucleotides are rarely fully methylated,
therefore the mid-range levels (anywhere between 20 to 50%) are usually sufficient to indicate open
chromatin regions. In our data, chromatin accessibility (i.e. GCH methylation) increased from
monocytes to macrophages with slightly higher levels in retained introns, while lymphoid cells had

increased chromatin accessibility (GCH methylation levels 15-35%) but with lower levels in retained
introns compared to non-retained introns (Figure S2C).

To determine important factors of IR regulation, we compiled sequence-based and epigenetic features:
(i) sequence-based features: intron length, GC content, splice site strength, CpG density (also referred
to as intrinsic features), (ii) transcriptomics features: percent spliced-in (PSI) values of the flanking
exons, and (iii) epigenomics features extracted from the WGBS, ChIP-Seq (H3K9me3, H3K27me3,
H3K27ac, H3K36me3, H3K4me1, H3K4me3), and NOMe-Seq data (Table S2). We then used these
features (n=46) to train EN models for each cell type and predict whether introns are either retained or
non-retained. The performance of our models was assessed based on the area under the receiver
operating characteristic curve (AUC) values, which ranged between 0.87 and 0.95 (Figure 2B) and
values for the area under the prediction-recall curve (accuracy) ranging between 0.85 and 0.95 (Figure
2C). The consistently high values suggest that the model choice was appropriate for the task.

Next, in order to evaluate whether the learned relationship between the model features and IR was
generalizable across cell types we trained our model with data from one cell type and tested it with data
from another cell type. For all training/test data pairs, the AUC and accuracy metrics were comparable
to those models that were trained and tested on the same cell type (Table S4).

The EN model assumes a monotonic linear relationship between the class variable and the model
features. To determine whether this assumption is adequate for IR classification, we also trained
conditional random forest (cRF) models, which do not make any prior assumption about the relationship
between the outcome of interest and the model features. Comparing the results from both types of
models, we found that cRF performed slightly better than EN with AUC values ranging between 0.91
and 0.98 (Figures 2D, S3A) and PR values between 0.87 and 0.95 (Figure S3B).

To assess which features contribute most to the model performance (and thus, the relevance of a feature
to IR), we used variable-importance measures (VIM). For EN, these are the regression coefficients
ordered from lowest to highest, where parameters with larger values have a greater effect. For cRF,
variable importance was calculated as the mean decrease in accuracy after permutation of each model
feature (Figure 2E). Given the known properties of retained introns it was no surprise that intrinsic
features, such as length, GC content and CpG density were ranked as the top predictors with a high
level of agreement across all cell types analysed. Again, we observed consistency between the EN and
cRF models, except for minor variations in the order that important features were ranked in.

Epigenetic features were also ranked among the top 5 predictors across all models and cell types,
however their nature and relative importance varied between cell types (Figure 2E). Overall EN models
ranked epigenetic features as moderately to very important (VIM between 0.4 and 0.8), which is
comparable to the intrinsic features (ranging between 0.3 and 1). In contrast, cRF identified epigenetic
features as somewhat important with VIM mostly below 0.50 (Figures S3C, S3D). Nevertheless,
intrinsic features were consistently identified as most relevant for correctly classifying IR, suggesting
that these features predispose introns to being retained irrespective of cell or tissue type.

177
Chromatin accessibility is predicted to be the strongest regulator of IR

In the previous section we classified IR on a cell type-specific basis and determined the intrinsic features
as having the strongest association with IR outcomes. However, we often find that an intron is retained
in one cell type but not in another. In those cases, factors beyond intrinsic features are the likely drivers
of this transition.

To find these IR determinants, we modified our initial modelling approach by focusing only on the
dynamic introns - those that changed their retention status between cell types (Figure 3A). In total,
1,540 introns matched this criterion with various IR patterns (Figure 3B). We used these introns to train
EN and cRF models with both epigenetic and intrinsic features. The cRF model was performed superior
to the EN model achieving AUCs of 0.85 and 0.76, respectively (Figure 3C). cRF also achieved a higher
area under the precision-recall curve value (0.83) than EN (0.73) (Figure 3D). The poorer performance
of EN might be a reflection of the model's inability to fully utilise complex structures within the omics

data, thus supporting the notion that a relationship between chromatin modifiers and IR is indeed non-
linear, as previously suggested (28).

Evaluation of feature rankings revealed that, despite varying model performances, both EN and cRF
models identified features related to chromatin accessibility as most important for correct IR
classification (Figure 3E). These features include GCH methylation and GCH (i.e. nucleosome)
occupancy and the presence of nucleosome free regions (NFRs). GCH methylation at the 5′ and 3′ splice
sites were determined as most important features discriminating retained form non-retained introns in
both models. The cRF classifier also identified CpG methylation as somewhat important for IR
classification, which has a known relationship with chromatin accessibility (29-31). Interestingly, the
cRF model also identified GC content as a moderately important contributor to IR outcomes, whilst the
EN model included histone marks (H3K27ac and H3K36me3) in their top 10 predictors (Figure S4A).

201
Epigenetic IR regulation is independent of gene expression regulation

It is reasonable to assume that changes in the epigenomic landscape might not directly affect IR but
rather gene expression (32). To confirm that the features identified as relevant to IR are independent
from gene expression regulation, we split dynamically retained introns into three groups: (i) host gene
expression is reduced along with the change in IR status, (ii) host gene expression remained stable ($\log_2$
FC FPKM $\leq$ 2), and (iii) host gene expression increased (Figure 4A). For most of the dynamic introns
the host gene expression remained unchanged (N = 1,220), whilst down- and upregulated host genes
were associated with 73 and 247 alternately retained introns, respectively. We repeated the
classification analysis on the group of introns where the IR changes were not accompanied by host gene
expression changes. Since the relationship between IR and epigenetic model features is not linear, as
was established in the previous section, we only used the cRF algorithm.

The model fitted on this data subset achieved an AUC of 0.83 (Figure 4B) and an area under the
precision-recall curve value of 0.78 (Figure S4B). The features that were selected as important were
GCH methylation at the 5′ and 3′ splice sites and GC content in the same order as in the model trained
on all dynamically retained introns (Figure 4C). This observation held true for both highly and lowly
expressed host genes (Figures S4C). We therefore concluded that the observed epigenetic changes
associated with IR modulation are independent from gene expression regulation. In Figure 4D, we show
two exemplary introns where greater chromatin accessibility was associated with an increase in IR:
Phosphatidylinositol Glycan Anchor Biosynthesis Class T (PIGT) helps building the
glycosylphosphatidylinositol-anchor which is found on the surface of various blood cells (Figure 4D,
left). PIGT is known to express many isoforms through alternative splicing including IR. The nucleotide
binding protein SEPTIN8 is a regulator of cytoskeletal organization, which has multiple alternatively
spliced transcript variants as well (Figure 4D, right).

225
Dynamic changes in chromatin structure are responsible for cell type-specific IR

As chromatin accessibility was identified as the strongest predictive factor for differential IR, we closely
examined its relationship with retained and non-retained introns. We identified 5 distinct GCH
methylation profiles in the +/- 200 bp region around the 5′ splice site of retained introns (Figure 5A,
left). Similar clustering profiles were identified in the region around 3′ splice sites and the middle of
introns (Figure S5). To understand changes in chromatin status in the context of differential IR, we
plotted the GCH methylation values of the same introns when they were not retained (Figure 5A). The
associated heatmap shows that GCH methylation is widely depleted in non-retained introns, with no
distinct clustering. In retained intron, however, we observed a clear increase in GCH methylation
immediately upstream or downstream from the 5′ splice site (Figure 5B, clusters 1, 3 and 4). We also
identified a group of retained introns with relatively low levels of GCH methylation (cluster 2) and
another with particularly strong GCH methylation (cluster 5).

Upon visualising the intronic regions that changed their IR status between cell types, we observed
greater chromatin accessibility levels in retained introns (Figure 6A). Moreover, for the majority of
introns, we found that IR gain was accompanied with a reduction in H3K36me3 signal (Figure 6A).

Based on the observed patterns, we hypothesise that there is an association between chromatin dynamics
and IR: chromatin is more likely to be in a permissive state (high GCH methylation) in the vicinity of
retained introns and more compact (low GCH methylation) around constitutively spliced introns.
Indeed, we observed that chromatin becomes more accessible as introns become retained (65% of
observations). In other cases, the IR status changes without any change to the chromatin state (35% of
observations).

Based on the observations concerning chromatin accessibility, we sought to assess the relationship
between IR and epigenetic factors in the context of changing chromatin states, i.e. differential GCH
methylation (Figure 6B), and stable chromatin status, i.e. non-differential GCH methylation (Figure
6C). In our analysis, we separated first introns from other introns to detach epigenetic signals associated
with gene promoters.

The patterns of CpG methylation, H3K27ac, H3K4me3 and H3K4me1 levels in retained and non-
retained introns were similar in both chromatin modes (dynamic and stable). First non-retained introns
displayed enrichment for histone marks and reduced CpG methylation levels, while first retained introns
had negligible levels of histone marks and were marked by the absence of CpG methylation (Figure 6B
and 6C, top rows). In contrast, the above-mentioned histone marks were silenced in the internal introns
irrespective of the IR status, while the H3K36me3 signal increased. Interestingly, H3K36me3 levels
were reduced in retained introns associated with dynamic chromatin (Figure 6B, 2nd row, far right),
while they remained similar in retained- and non-retained introns associated with stable chromatin
(Figure 6C, 2nd row, far right).

A most interesting result of this analysis was that there are no differences in epigenetic marks between
internal retained and non-retained introns when a stable chromatin state is maintained (Figure 6C,
bottom row). This suggests that there must be unknown factors that are independent of chromatin
accessibility responsible for modulating IR. Thus, further investigations are required to identify
additional factors that impact on IR in haematopoietic cells.

Discussion

In this study, we have employed a machine-learning approach to determine regulators of IR in primary
hematopoietic cells. Previous studies have described features that are associated with retained introns,
including a higher intronic GC content, shorter intron lengths, weaker 5′ and 3′ splice site strengths, and
some epigenetic marks (2,14,17). However, these studies have focused on single or paired omics layers
only and often used individual cell lines for their analyses. In contrast, we have for the first time
integrated matched transcriptomic, nucleosome occupancy, CpG methylation and 6 histone
modification profiles from 5 primary human cell types representing 2 independent systems of
haematopoietic cell differentiation.

We approached this task as a supervised learning problem using EN and conditional RF algorithms.
Unlike deep learning methods, that are very capable of identifying complex relationship patterns but do
not provide tools to determine how exactly an outcome was determined (33), multivariate modelling
allows the identification of features that contribute most to the outcome of interest. Such modelling
strategy is "data-independent" and can be applied to other forms of alternative splicing as well. For
example, RF has been used to study the importance of chromatin modifications in the interaction
between topologically associated domains (34) and EN was used to model prognostic alternative
splicing signatures in breast cancer (35).

A functional link between gene expression regulation and chromatin organisation has been previously
described. Nucleosome free regions at a transcription start site are strongly associated with transcription
initiation (36). Nucleosomes were also reported to be preferentially positioned in exons to facilitate

their identification among flanking introns by the splicing machinery (37,38). However, it is important to note that these findings were made using the micrococcal nuclease digestion with deep sequencing (MNase-seq) protocol, which is more susceptible to GC content bias. Kelly et al. (39) showed that nucleosome enrichment in exons vs. introns was not observed in NOMe-seq data, which they attributed to the technical differences between the two experimental approaches. NOMe-seq data includes the percentage of methylated reads at a given position as opposed to the count of mapped reads in MNase-seq data. Similarly, our NOMe-seq based analysis of chromatin accessibility, quantified by GCH methylation, did not reveal a specific preference for nucleosomes to be positioned in exons rather than introns. However, nucleosomes were preferentially positioned immediately upstream or downstream of retained introns in contrast to non-retained introns.

Nucleosome positioning has been linked to alternative splicing before, where RNA Pol II elongation rates increase upon histone depletion and pre-mRNA splicing is delayed (40). Previous studies identified nucleosomes as physical barriers to efficient transcription elongation *in vitro*, however *in vivo* they are efficiently removed from transcribed chromatin (41). Enrichment for GCH methylated loci might be an indication of euchromatin formation, which is usually decorated with certain histone modifications and characterised by nucleosome depletion and reduced CpG methylation levels (42). High GCH methylation levels in retained introns, that we observed in our analysis, are indicative of NFRs, regions of possible nucleosome eviction that are characterised by a high density of methylated GCH sites and unmethylated CpG sites (43). Interestingly, You et al. showed that a loss of nucleosome depleted regions accompanied by nucleosome occupancy precedes changes in endogenous CpG methylation in OCT4 and NANOG genes in embryonic carcinoma cell line NCCIT (44). Formation of an NFR upstream from the 5′ exon/intron boundary led to DNA hypomethylation and the depletion of H3K36me3 in SETD2 deficient tumours (45). It is therefore reasonable to conclude that alteration of the epigenetic landscape attributed to IR initially starts with changes in nucleosome architecture and subsequent transcriptome rewiring.

Strong histone modification levels in the proximity of transcription start sites (like we observed for H3K4me3 and H3K27ac) indicate a well-positioned nucleosome (46), while reduced histone modification levels, particularly reduced H3K4me3, are associated with transcription factor (TF) binding (47). TF binding sites can undergo nucleosome remodelling (48) in the form of nucleosome shifts or nucleosome eviction and the formation of an NFR with associated changes to RNA polymerase II elongation rates. We propose that IR in first introns might be a biproduct of functional histone modifications and nucleosome remodelling for the purpose of TF recruitment in the regions proximal to transcription start sites. In regions further downstream of transcription start sites, nucleosome positioning becomes less stable (36), which may explain less pronounced differences in H3K36me3 signal over internal introns.

In summary, our analysis provides new insights into the role of epigenetic modifications in the regulation of IR and highlights the unanticipated strong contribution of nucleosome occupancy in IR modulation. Moreover, the models generated in this study can be adapted to study epigenetic gene expression and alternative splicing regulation in other cell systems or species.

Methods

To investigate how IR is regulated in primary immune cells, we integrated epigenomics and transcriptomics data from the German Epigenome Program (DEEP). Primary monocytes, monocyte-derived macrophages, and primary T-cells (naïve, central memory, effector memory) were retrieved from 2 healthy donors. Cell isolation, differentiation, DNA/RNA extraction and library preparation for mRNA-Seq, WGBS, NOMe- and ChIP-Seq experiments are described in detail in these articles (49,50).

mRNA-Seq data processing and identification of IR events

RNA-Seq reads (FASTQ format) of each technical replicate were tested for quality using FastQC v.0.11.5 (github.com/s-andrews/FastQC). Further processing, including adaptor trimming, was

335 performed within the IRFinder algorithm for IR quantification (4). Sequencing reads were mapped to
336 the human reference genome (GRCh38) using STAR v2.7 with default parameters (51). IR-ratios, a
337 quantitative measure of IR levels, were determined as:

$$338 \quad IRratio = \frac{Intronic\ Abundance}{Intronic\ Abundance\ +\ Exonic\ Abundance},$$

where the Intronic Abundance is defined as the trimmed mean of the reads that map to an intron, after
having excluded features that overlap the intron, with the highest and lowest 30% of values being
excluded. Exonic Abundance is defined as the number of reads that map across an exon-exon junction.
Library size normalisation (between-sample normalisation) was not required as the ratio between
intronic and exonic abundance is determined from within the same transcriptome (4).

Introns that were present in at least 10% of a gene's mature mRNA transcripts ($IR_{ratio} \geq 0.1$) with an
overall intron depth ≥ 10 were considered retained. Non-retained introns were defined as those with an
$IR_{ratio} \leq 0.01$ and intron depth < 10.

We used Cufflinks v2.1.1 (52) to estimate gene abundance in fragments per kilobase per million
(FPKM). Only introns from host genes with FPKM ≥ 1 were selected for the downstream analyses.

WGBS data processing

Raw WGBS FASTQ files were assessed for quality using FastQC v.0.11.5 (github.com/s-
andrews/FastQC). Standard Illumina adaptors used for the library preparation were trimmed using
cutadapt v.1.10 with a quality cutoff of 20 base pairs (bp) and minimum read length of 30 bp. Trimmed
reads were mapped to the GRCh38 reference genome, duplicate reads removed, and methylation calling
performed using Bismark v.0.19.0 (53). Only CpG sites with a coverage of more than 5 reads were
retained for further analysis.

ChIP-Seq data processing

ChIP-Seq data for six histone modifications (H2K27ac, H3K27me3, H3K36me3, H3K4me1,
H3K4me3, H3K9me3) were aligned to the human GRCh38 reference genome using STAR v2.7 (51).
Duplicate reads were removed using Picard v.2.18.4 (broadinstitute.github.io/picard/) and further
processed using MACS2 v.2.2.6 (54) to identify histone modification peaks, with default parameters
and q-value cut-off of 0.01. All histone modifications were processed in the "narrow peak" mode in
order to extract peak summit coordinates. For visualisation in IGV, we generated coverage tracks using
`bamCoverage` from deepTools2 with the following parameters `--binSize 1 --`
`normalizeUsing BPM --effectiveGenomeSize 2913022398 --extendReads 200`.
For HM line plots, we substracted ChIP-Seq Input from a respective HM ChIP-seq read counts and
normalised based on Bins Per Million (BPM) mapped reads using bamCompare and parameters `--`
`binSize 1 --scaleFactorsMethod readCount --effectiveGenomeSize`
`2913022398 --operation subtract --normalizeUsing BPM`.

NOMe-Seq data processing

Raw FASTQ files were assessed for quality using FastQC v.0.11.5 (github.com/s-andrews/FastQC).
Reads were mapped to the GRCh38 reference genome, duplicate reads removed, and methylation
calling performed using Bismark v.0.19.0 (53). GCH methylation information was extracted with the
`coverage2cytosine` utility with `--nome` parameter.

NFRs were predicted using gNOMeHMM tool (github.com/karl616/gNOMePeaks/) with default
parameters, which include 4,000 bp up- and downstream from each peak for background signal
calculation and the maximum distance between GpC sites of 150 bp (50). We used the same algorithms
to predict nucleosome positioning by substituting GCH methylation, as required input, with GCH
occupancy ($1 - GCH\ methylation$) and reducing the background region to 1,000 bp up- and
downstream from each peak and the distance between GCH sites to 20bp.

Feature selection

Model features were associated with three genomic regions around retained and non-retained introns: (i) +/- 100 bp from the 5´splice site, (ii) +/- 100 bp from the 3´splice site, and (iii) +/- 100bp from the middle of an intron, each region being 200 bp long. GC content was extracted using bedtools () `nuc` command. For splice site strength calculations, we used MaxEnt scan. CpG density values was obtained using epiR. The percent spliced in (PSI) index of flanking exons was calculated as described in (55). Exons with PSI ≥ 0.9 were considered as included.

To generate epigenetic features, we overlapped three regions of interest with the pre-processed epigenetic data. NFR regions were defined as regions greater than 40bp in length with p-value ≤ 0.05 (Fisher test comparing CpG methylation in the NFR to the surrounding background). Presence or absence of an NFR was dichotomised as "yes" – 1 and "no" – 0. Information about nucleosome location was included into the model in the similar manner (nucleosomes were defined as regions greater than 140bp in length with p-value ≤ 0.05).

The relationship between histone modification and IR was included into the model through the presence or absence of an overlap with a histone signal region. It was categorised as 0 – no overlap, 1 – overlap with a region of HM signal, 2 – overlap with a region of strong signal (strong signal = mean (HM pile-up) + sd (HM pile-up)). The full list of features is presented in Table S1.

Elastic Net and Conditional Random Forest Modelling

To identify features important for IR, we constructed binary classification models using EN and cRF algorithms. We approached our problem in a naïve manner, i.e. we did not impose any prior assumptions about the factors that might potentially play a role and therefore an equal penalty factor was applied to all features. To avoid overfitting, we ranked the features' importance using both EN and cRF techniques (56).

EN classification was performed in the *caret* R package (57) using *glmnet* method (58) for a binary outcome. The group imbalance, due to the different number of retained and non-retained introns identified suitable for modelling, was handled by down-sampling, using *downSample* command. Parameter λ was optimised by 10-fold cross validation procedure. Features were ranked based on the absolute values of the model coefficients.

In cRF, unlike standard RF where the first split variable is randomly selected, an association test between the outcome and the model predictors is performed first. The ranked p-values are then used to identify the covariate with the strongest association to the outcome, which is later used for the first binary split at cutpoint c for a continuous covariate or at category C for a categorical covariate. cRF classification was also performed in *caret* using *cforest* method as implemented in the *party* R package (59). The cRF model provides an unbiased measure of variable importance, which we used to rank the most important features for IR prediction.

Statistical Analysis

Statistical analyses were performed in R v.4.0, unless stated otherwise. P-values ≤ 0.05 were considered significant, no multiple testing was conducted.

Data Availability

Sequencing data are deposited at the European Genome-Phenome Archive under the accession number EGAS00001001608. Access is subject to an application process as per the EGA requirements.

Acknowledgements

We thank Benedikt Brors and Roland Eils from DKFZ Heidelberg and Alf Hamann from DRFZ Berlin, Wie Chen, Nikolaus Rajewsky and Sascha Sauer from MDC Berlin, Ho-Ryun Chung and Martin

Vingron from MPI-MG Berlin, Thomas Jenuwein, Thomas Manke and Andrew Pospisilik from MPI-IE Freiburg, Philip Rosenstiel and Stefan Schreiber from CAU Kiel, Jan G. Hengstler from IfADo Dortmund, Thomas Lengauer from MPI-INF Saarbrücken, Bernhard Horsthemke from Universität Duisburg-Essen, Alexandra Kiemer from Universität des Saarlandes Saarbrücken, Thomas Pap from WWU Münster and Gerd Schmitz from Universität Regensburg who were involved in the work with biological samples, sequencing and generation of WGBS, NOMe-Seq, ChIP-Seq and RNA-Seq data for the DEEP Consortium.

The authors acknowledge the technical assistance provided by the Sydney Informatics Hub, a Core Research Facility of the University of Sydney.

Author Contributions: J.E.J.R., J.J.-L.W. and U.S. designed the study and supervised the project, V.P. and R.S. performed bioinformatic analyses, V.P. performed statistical analysis and data modelling, N.J.A. advised on statistical methodology, DEEP Consortium provided sequencing data, J.W. designed and coordinated sequencing experiments, K.J.V.N. data management, V.P. and U.S. wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding

This work was supported by the National Health and Medical Research Council (Investigator Grant #1177305 to J.E.J.R., Project #1128175 and #1129901 to J.E.J.R. and J.J.-L.W., #1126306 to J.J.-L.W.; the NSW Genomics Collaborative Grant (J.E.J.R. and J.J.-L.W.); Cure the Future (J.E.J.R.), and an anonymous foundation (J.E.J.R.). U.S. and J.J.-L.W. hold Fellowships from the Cancer Institute of New South Wales. U.S. also received support from the Australian Academy of Science in form of an Australia-India Early and Mid-Career Fellowship. This research was funded by the Cancer Council NSW Project Grants (RG11-12, RG14-09, RG20-12) to J.E.J.R. and U.S. K.V.J.N. and J.W. were supported by the German Epigenome Program (DEEP) funded by the Ministry of Education and Research in Germany (BMBF 01KU1216).

Conflict of Interest: J.E.J.R. has received honoraria or speakers' fees (GSK, Miltenyi, Takeda, Gilead, Pfizer, Spark, Novartis, Celgene, bluebird bio); Director of Pathology (Genea); equity ownership (Genea, Rarecyte); consultant (Rarecyte, Imago); chair, Gene Technology Technical Advisory, OGTR, Australian Government. K.J.V.N. is currently employed by AstraZeneca. The remaining authors declare no competing financial interests.

References

1. Wong, J.J., Ritchie, W., Ebner, O.A., Selbach, M., Wong, J.W., Huang, Y., Gao, D., Pinello, N., Gonzalez, M., Baidya, K. *et al.* (2013) Orchestrated intron retention regulates normal granulocyte differentiation. *Cell*, **154**, 583-595.
2. Schmitz, U., Pinello, N., Jia, F., Alasmari, S., Ritchie, W., Keightley, M.C., Shini, S., Lieschke, G.J., Wong, J.J. and Rasko, J.E.J. (2017) Intron retention enhances gene regulatory complexity in vertebrates. *Genome Biol*, **18**, 216.
3. Monteuuis, G., Wong, J.J.L., Bailey, C.G., Schmitz, U. and Rasko, J.E.J. (2019) The changing paradigm of intron retention: regulation, ramifications and recipes. *Nucleic Acids Res*, **47**, 11497-11513.
4. Middleton, R., Gao, D., Thomas, A., Singh, B., Au, A., Wong, J.J., Bomane, A., Cosson, B., Eyras, E., Rasko, J.E. *et al.* (2017) IRFinder: assessing the impact of intron retention on mammalian gene expression. *Genome Biol*, **18**, 51.
5. Edwards, C.R., Ritchie, W., Wong, J.J., Schmitz, U., Middleton, R., An, X., Mohandas, N., Rasko, J.E. and Blobel, G.A. (2016) A dynamic intron retention program in the mammalian megakaryocyte and erythrocyte lineages. *Blood*, **127**, e24-e34.

6.   Ni, T., Yang, W., Han, M., Zhang, Y., Shen, T., Nie, H., Zhou, Z., Dai, Y., Yang, Y., Liu, P. *et*
al. (2016) Global intron retention mediated gene regulation during CD4+ T cell activation.
Nucleic Acids Res, **44**, 6817-6829.
7.   Ullrich, S. and Guigo, R. (2020) Dynamic changes in intron retention are tightly associated
with regulation of splicing factors and proliferative activity during B-cell development. *Nucleic*
Acids Res, **48**, 1327-1340.
8.   Green, I.D., Pinello, N., Song, R., Lee, Q., Halstead, J.M., Kwok, C.T., Wong, A.C.H., Nair,
S.S., Clark, S.J., Roediger, B. *et al.* (2020) Macrophage development and activation involve
coordinated intron retention in key inflammatory regulators. *Nucleic Acids Res*, **48**, 6513-6529.
9.   Wong, J.J., Au, A.Y., Ritchie, W. and Rasko, J.E. (2016) Intron retention in mRNA: No longer
nonsense: Known and putative roles of intron retention in normal and disease biology.
Bioessays, **38**, 41-49.
10.   Smart, A.C., Margolis, C.A., Pimentel, H., He, M.X., Miao, D., Adeegbe, D., Fugmann, T.,
Wong, K.K. and Van Allen, E.M. (2018) Intron retention is a source of neoepitopes in cancer.
Nat Biotechnol, **36**, 1056-1058.
11.   Monteuuis, G., Schmitz, U., Petrova, V., Kearney, P.S. and Rasko, J.E.J. (2020) Holding on to
junk bonds: intron retention in cancer and therapy. *Cancer Res*.
12.   Hershberger, C.E., Moyer, D.C., Adema, V., Kerr, C.M., Walter, W., Hutter, S., Meggendorfer,
M., Baer, C., Kern, W., Nadarajah, N. *et al.* (2020) Complex landscape of alternative splicing
in myeloid neoplasms. *Leukemia*.
13.   Dvinge, H., Guenthoer, J., Porter, P.L. and Bradley, R.K. (2019) RNA components of the
spliceosome regulate tissue- and cancer-specific alternative splicing. *Genome Res*, **29**, 1591-
1604.
14.   Braunschweig, U., Barbosa-Morais, N.L., Pan, Q., Nachman, E.N., Alipanahi, B.,
Gonatopoulos-Pournatzis, T., Frey, B., Irimia, M. and Blencowe, B.J. (2014) Widespread intron
retention in mammals functionally tunes transcriptomes. *Genome Res*, **24**, 1774-1786.
15.   Zhou, H.L., Luo, G., Wise, J.A. and Lou, H. (2014) Regulation of alternative splicing by local
histone modifications: potential roles for RNA-guided mechanisms. *Nucleic Acids Res*, **42**, 701-
713.
16.   Fong, N., Kim, H., Zhou, Y., Ji, X., Qiu, J., Saldi, T., Diener, K., Jones, K., Fu, X.D. and
Bentley, D.L. (2014) Pre-mRNA splicing is facilitated by an optimal RNA polymerase II
elongation rate. *Genes Dev*, **28**, 2663-2676.
17.   Wong, J.J., Gao, D., Nguyen, T.V., Kwok, C.T., van Geldermalsen, M., Middleton, R., Pinello,
N., Thoeng, A., Nagarajah, R., Holst, J. *et al.* (2017) Intron retention is regulated by altered
MeCP2-mediated splicing factor recruitment. *Nat Commun*, **8**, 15134.
18.   Kim, D., Shivakumar, M., Han, S., Sinclair, M.S., Lee, Y.J., Zheng, Y., Olopade, O.I., Kim, D.
and Lee, Y. (2018) Population-dependent Intron Retention and DNA Methylation in Breast
Cancer. *Mol Cancer Res*, **16**, 461-469.
19.   Gao, D., Pinello, N., Nguyen, T.V., Thoeng, A., Nagarajah, R., Holst, J., Rasko, J.E. and Wong,
J.J. (2019) DNA methylation/hydroxymethylation regulate gene expression and alternative
splicing during terminal granulopoiesis. *Epigenomics*, **11**, 95-109.
20.   Guo, R., Zheng, L., Park, J.W., Lv, R., Chen, H., Jiao, F., Xu, W., Mu, S., Wen, H., Qiu, J. *et*
al. (2014) BS69/ZMYND11 reads and connects histone H3.3 lysine 36 trimethylation-
decorated chromatin to regulated pre-mRNA processing. *Mol Cell*, **56**, 298-310.
21.   Wei, G., Liu, K., Shen, T., Shi, J., Liu, B., Han, M., Peng, M., Fu, H., Song, Y., Zhu, J. *et al.*
(2018) Position-specific intron retention is mediated by the histone methyltransferase SDG725.
BMC Biol, **16**, 44.
22.   Barash, Y., Calarco, J.A., Gao, W., Pan, Q., Wang, X., Shai, O., Blencowe, B.J. and Frey, B.J.
(2010) Deciphering the splicing code. *Nature*, **465**, 53-59.
23.   Leung, M.K., Xiong, H.Y., Lee, L.J. and Frey, B.J. (2014) Deep learning of the tissue-regulated
splicing code. *Bioinformatics*, **30**, i121-129.
24.   Xiong, H.Y., Alipanahi, B., Lee, L.J., Bretschneider, H., Merico, D., Yuen, R.K., Hua, Y.,
Gueroussov, S., Najafabadi, H.S., Hughes, T.R. *et al.* (2015) RNA splicing. The human splicing
code reveals new insights into the genetic determinants of disease. *Science*, **347**, 1254806.

25. Jaganathan, K., Kyriazopoulou Panagiotopoulou, S., McRae, J.F., Darbandi, S.F., Knowles, D.,
Li, Y.I., Kosmicki, J.A., Arbelaez, J., Cui, W., Schwartz, G.B. *et al.* (2019) Predicting Splicing
from Primary Sequence with Deep Learning. *Cell*, **176**, 535-548 e524.
26. Baeza-Centurion, P., Minana, B., Schmiedel, J.M., Valcarcel, J. and Lehner, B. (2019)
Combinatorial Genetics Reveals a Scaling Law for the Effects of Mutations on Splicing. *Cell*,
176, 549-563 e523.
27. Pacini, C. and Koziol, M.J. (2018) Bioinformatics challenges and perspectives when studying
the effect of epigenetic modifications on alternative splicing. *Philos Trans R Soc Lond B Biol*
Sci, **373**.
28. Singer, M., Kosti, I., Pachter, L. and Mandel-Gutfreund, Y. (2015) A diverse epigenetic
landscape at human exons with implication for expression. *Nucleic Acids Research*, **43**, 3498-
3508.
29. Lay, F.D., Liu, Y., Kelly, T.K., Witt, H., Farnham, P.J., Jones, P.A. and Berman, B.P. (2015)
The role of DNA methylation in directing the functional organization of the cancer epigenome.
Genome Res, **25**, 467-477.
30. Farlik, M., Halbritter, F., Muller, F., Choudry, F.A., Ebert, P., Klughammer, J., Farrow, S.,
Santoro, A., Ciaurro, V., Mathur, A. *et al.* (2016) DNA Methylation Dynamics of Human
Hematopoietic Stem Cell Differentiation. *Cell Stem Cell*, **19**, 808-822.
31. Taberlay, P.C., Statham, A.L., Kelly, T.K., Clark, S.J. and Jones, P.A. (2014) Reconfiguration
of nucleosome-depleted regions at distal regulatory elements accompanies DNA methylation
of enhancers and insulators in cancer. *Genome Research*, **24**, 1421-1432.
32. Jaenisch, R. and Bird, A. (2003) Epigenetic regulation of gene expression: how the genome
integrates intrinsic and environmental signals. *Nat Genet*, **33 Suppl**, 245-254.
33. Rauschert, S., Raubenheimer, K., Melton, P.E. and Huang, R.C. (2020) Machine learning and
clinical epigenetics: a review of challenges for diagnosis and classification. *Clinical*
Epigenetics, **12**, 51.
34. Dixon, J.R., Jung, I., Selvaraj, S., Shen, Y., Antosiewicz-Bourget, J.E., Lee, A.Y., Ye, Z., Kim,
A., Rajagopal, N., Xie, W. *et al.* (2015) Chromatin architecture reorganization during stem cell
differentiation. *Nature*, **518**, 331-336.
35. Wang, L., Wang, Y., Su, B., Yu, P., He, J., Meng, L., Xiao, Q., Sun, J., Zhou, K., Xue, Y. *et al.*
(2020) Transcriptome-wide analysis and modelling of prognostic alternative splicing signatures
in invasive breast cancer: a prospective clinical study. *Scientific Reports*, **10**, 16504.
36. Radman-Livaja, M. and Rando, O.J. (2010) Nucleosome positioning: How is it established, and
why does it matter? *Developmental Biology*, **339**, 258-266.
37. Schwartz, S., Meshorer, E. and Ast, G. (2009) Chromatin organization marks exon-intron
structure. *Nature Structural & Molecular Biology*, **16**, 990-995.
38. Tilgner, H., Nikolaou, C., Althammer, S., Sammeth, M., Beato, M., Valcárcel, J. and Guigó, R.
(2009) Nucleosome positioning as a determinant of exon recognition. *Nature Structural &*
Molecular Biology, **16**, 996-1001.
39. Kelly, T.K., Liu, Y., Lay, F.D., Liang, G., Berman, B.P. and Jones, P.A. (2012) Genome-wide
mapping of nucleosome positioning and DNA methylation within individual DNA molecules.
Genome research, **22**, 2497-2506.
40. Jimeno-González, S., Payán-Bravo, L., Muñoz-Cabello, A.M., Guijo, M., Gutierrez, G., Prado,
F. and Reyes, J.C. (2015) Defective histone supply causes changes in RNA polymerase II
elongation rate and cotranscriptional pre-mRNA splicing. *Proceedings of the National*
Academy of Sciences, **112**, 14840-14845.
41. Saldi, T., Cortazar, M.A., Sheridan, R.M. and Bentley, D.L. (2016) Coupling of RNA
Polymerase II Transcription Elongation with Pre-mRNA Splicing. *J Mol Biol*, **428**, 2623-2635.
42. Portela, A. and Esteller, M. (2010) Epigenetic modifications and human disease. *Nature*
Biotechnology, **28**, 1057-1068.
43. Nordström, K.J.V., Schmidt, F., Gasparoni, N., Salhab, A., Gasparoni, G., Kattler, K., Müller,
F., Ebert, P., Costa, I.G., consortium, D. *et al.* (2019) Unique and assay specific features of
NOMe-, ATAC- and DNase I-seq data. *Nucleic Acids Research*, **47**, 10580-10596.
44. You, J.S., Kelly, T.K., De Carvalho, D.D., Taberlay, P.C., Liang, G. and Jones, P.A. (2011)
OCT4 establishes and maintains nucleosome-depleted regions that provide additional layers of

epigenetic regulation of its target genes. *Proceedings of the National Academy of Sciences*, **108**,
14497-14502.

45. Simon, J.M., Hacker, K.E., Singh, D., Brannon, A.R., Parker, J.S., Weiser, M., Ho, T.H., Kuan,
P.-F., Jonasch, E., Furey, T.S. *et al.* (2014) Variation in chromatin accessibility in human
kidney cancer links H3K36 methyltransferase loss with widespread RNA processing defects.
Genome research, **24**, 241-250.

46. Andersson, R., Enroth, S., Rada-Iglesias, A., Wadelius, C. and Komorowski, J. (2009)
Nucleosomes are well positioned in exons and carry characteristic histone modifications.
Genome research, **19**, 1732-1741.

47. Wu, J.N., Pinello, L., Yissachar, E., Wischhusen, J.W., Yuan, G.-C. and Roberts, C.W.M.
(2015) Functionally distinct patterns of nucleosome remodeling at enhancers in glucocorticoid-
treated acute lymphoblastic leukemia. *Epigenetics & Chromatin*, **8**, 53.

48. Ballaré, C., Castellano, G., Gaveglia, L., Althammer, S., González-Vallinas, J., Eyras, E.,
Le Dily, F., Zaurin, R., Soronellas, D., Vicent, Guillermo P. *et al.* (2013) Nucleosome-Driven
Transcription Factor Binding and Gene Regulation. *Molecular Cell*, **49**, 67-79.

49. Durek, P., Nordström, K., Gasparoni, G., Salhab, A., Kressler, C., de Almeida, M., Bassler, K.,
Ulas, T., Schmidt, F., Xiong, J. *et al.* (2016) Epigenomic Profiling of Human CD4(+) T Cells
Supports a Linear Differentiation Model and Highlights Molecular Regulators of Memory
Development. *Immunity*, **45**, 1148-1161.

50. Wallner, S., Schroder, C., Leitao, E., Berulava, T., Haak, C., Beisser, D., Rahmann, S., Richter,
A.S., Manke, T., Bonisch, U. *et al.* (2016) Epigenetic dynamics of monocyte-to-macrophage
differentiation. *Epigenetics Chromatin*, **9**, 33.

51. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M. and Gingeras, T.R. (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**,
15-21.

52. Trapnell, C., Williams, B.A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M.J., Salzberg,
S.L., Wold, B.J. and Pachter, L. (2010) Transcript assembly and quantification by RNA-Seq
reveals unannotated transcripts and isoform switching during cell differentiation. *Nat
Biotechnol*, **28**, 511-515.

53. Krueger, F. and Andrews, S.R. (2011) Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics*, **27**, 1571-1572.

54. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,
Myers, R.M., Brown, M., Li, W. *et al.* (2008) Model-based analysis of ChIP-Seq (MACS).
Genome Biol, **9**, R137.

55. Schafer, S., Miao, K., Benson, C.C., Heinig, M., Cook, S.A. and Hubner, N. (2015) Alternative
Splicing Signatures in RNA-seq Data: Percent Spliced in (PSI). *Curr Protoc Hum Genet*, **87**,
11161-111614.

56. Ding, M.Q., Chen, L., Cooper, G.F., Young, J.D. and Lu, X. (2018) Precision Oncology beyond
Targeted Therapy: Combining Omics Data with Machine Learning Matches the Majority of
Cancer Cells to Effective Therapeutics. *Mol Cancer Res*, **16**, 269-278.

57. Kuhn, M. (2008) Building Predictive Models in R Using the caret Package. *Journal of
Statistical Software*, **28**, 1-26.

58. Friedman J, H.T., Tibshirani R. (2010) Regularization Paths for Generalized Linear Models via
Coordinate Descent. *Journal of Statistical Software*, **33**, 1-22.

59. Strobl C, B.A., Kneib T, Augustin T, Zeileis A. (2008) Conditional Variable Importance for
Random Forests. *BMC Bioinformatics*, **9**.

Figure legends

Figure 1 Experimental design and workflow to determine regulators of IR. Raw high-throughput data were processed for each biological replicate and amalgamated by cell type from the indicated number of samples (n). The output was used for feature extraction: IR events were treated as a binary outcome and we trained an Elastic Net (EN) regression model and a conditional Random Forest model with a total of 46 sequence-based and epigenetic features. Using feature ranking, we identified the factors that were most strongly associated with IR outcomes and compared the performances of both modelling strategies. These steps were repeated for each cell type.

Figure 2 IR prediction and model feature association analyses. (A) Scatter plot of differential IR events (Sig blue – significant; Not Sig yellow – not significant) between monocytes (Mo) vs macrophages (Ma) (left), Naïve (TN) vs Central Memory (CM) T cells (middle), and Central Memory vs Effector Memory (EM) T cells (right). **(B)** Receiver operating characteristic (ROC) curves and **(C)** precision recall (PR) curves comparing the performance of the EN classifier in five cell types. **(D)** Comparison of AUC values between EN and cRF algorithms, error bars show 95% confidence interval. **(E)** Variable importance scores for the top 10 features identified by EN and conditional RF algorithms. The scores were scaled to values that add up to 1.0 and the size of a bar corresponds to the effect size.

Figure 3. Analysis of dynamics intron retention. (A) Modified modelling strategy from Figure 1. Only introns that were found to be in retained and non-retained states in different cell types were included in the analysis. **(B)** Alluvial plot illustrating the dynamics of IR states among the five cell types. **(C)** ROC and **(D)** PR curves comparing the performance of cRF (brown) and EN (black). **(E)** Variable importance scores for the top 5 features identified by EN and conditional RF algorithms, scaled between 0 and 1.

Figure 4 Analysis of introns from genes with non-differential expression levels. (A) Scatter plot of host gene expression for introns that change their IR status. **(B)** ROC curve indicating the performance of a conditional RF model fitted on the data from non-differentially expressed genes (GE, gene expression). **(C)** Ranking of the features based on the scaled variable importance scores. **(D)** Integrative Genomics Viewer (IGV) plots revealing higher density and hypermethylation levels of GCH sites in the splice site regions of differentially retained introns in both highly- and lowly- expressed gene examples (NFR – Nucleosome Free Region, GCH Methylation – methylation levels of GC dinucleotides followed by any nucleobase except guanine).

Figure 5 GCH methylation clustering in differentially retained introns. **(A)** Clustering of GCH methylation in the +/- 200 bp region around the 5′ splice site (ss). Each line corresponds to one intron that is either in a retained (left) or non-retained state (right). **(B)** Line plots showing average GCH methylation values (i.e. chromatin accessibility) in retained vs non-retained introns across 5 clusters.

Figure 6 Interplay between chromatin accessibility, CpG methylation and histone modifications. **(A)** IGV plots of mRNA-seq, H3K36me3 ChIP-seq, NOMe-seq, and WGBS-seq data indicating different levels of GCH methylation between retained and non-retained introns and higher prevalence of NFRs in the regions proximal to IR. **(B)** Line graphs show the average levels of GCH methylation, CpG methylation, and the difference between ChIP-seq H3K4me3, H3K27ac, H3K4me1, and H3K36me3 signals and ChIP-Seq Input, normalised to the Bins Per Million (BPM), in retained (red) and non-retained (blue) introns associated with chromatin status. The first row shows epigenetic signals at the 5′ splice site of first introns (close to the promoter region) and the second row represents all other introns. **(C)** The same analysis performed in (B) is repeated for introns where the chromatin status remains the same, i.e. non-differential GCH methylation.

Main Figures

Figure 1 Experimental design and workflow to determine regulators of IR. Raw high-throughput data were processed for each biological replicate and amalgamated by cell type from the indicated number of samples (n). The output was used for feature extraction: IR events were treated as a binary outcome and we trained an Elastic Net (EN) regression model and a conditional Random Forest model with a total of 46 sequence-based and epigenetic features. Using feature ranking, we identified the factors that were most strongly associated with IR outcomes and compared the performances of both modelling strategies. These steps were repeated for each cell type.

Figure 2 IR prediction and model feature association analyses. (A) Scatter plot of differential IR events (Sig blue – significant; Not Sig yellow – not significant) between monocytes (Mo) vs macrophages (Ma) (left), Naïve (TN) vs Central Memory (CM) T cells (middle), and Central Memory vs Effector Memory (EM) T cells (right). **(B)** Receiver operating characteristic (ROC) curves and **(C)** precision recall (PR) curves comparing the performance of the EN classifier in five cell types. **(D)** Comparison of AUC values between EN and cRF algorithms, error bars show 95% confidence interval. **(E)** Variable importance scores for the top 10 features identified by EN and conditional RF algorithms. The scores were scaled to values that add up to 1.0 and the size of a bar corresponds to the effect size.

Figure 3 Analysis of dynamics intron retention. (A) Modified modelling strategy from Figure 1. Only introns that were found to be in retained and non-retained states in different cell types were included in the analysis. **(B)** Alluvial plot illustrating the dynamics of IR states among the five cell types. **(C)** ROC and **(D)** PR curves comparing the performance of cRF (brown) and EN (black). **(E)** Variable importance scores for the top 5 features identified by EN and conditional RF algorithms, scaled between 0 and 1.

Figure 4 Analysis of introns from genes with non-differential expression levels. (A) Scatter plot of host gene expression for introns that change their IR status. **(B)** ROC curve indicating the performance of a conditional RF model fitted on the data from non-differentially expressed genes (GE, gene expression). **(C)** Ranking of the features based on the scaled variable importance scores. **(D)** Integrative Genomics Viewer (IGV) plots revealing higher density and hypermethylation levels of GCH sites in the splice site regions of differentially retained introns in both highly- and lowly- expressed gene examples (NFR – Nucleosome Free Region, GCH Methylation – methylation levels of GC dinucleotides followed by any nucleobase except guanine).

Figure 5 GCH methylation clustering in differentially retained introns. (A) Clustering of GCH methylation in the +/- 200 bp region around the 5′ splice site (ss). Each line corresponds to one intron that is either in a retained (left) or non-retained state (right). **(B)** Line plots showing average GCH methylation values (i.e. chromatin accessibility) in retained vs non-retained introns across 5 clusters.

Figure 6 Interplay between chromatin accessibility, CpG methylation and histone modifications. (A) IGV plots of mRNA-seq, H3K36me3 ChIP-seq, NOMe-seq, and WGBS-seq data indicating different levels of GCH methylation between retained and non-retained introns and higher prevalence of NFRs in the regions proximal to IR. **(B)** Line graphs show the average levels of GCH methylation, CpG methylation, and the difference between ChIP-seq H3K4me3, H3K27ac, H3K4me1, and H3K36me3 signals and ChIP-Seq Input, normalised to the Bins Per Million (BPM), in retained (red) and non-retained (blue) introns associated with chromatin status. The first row shows epigenetic signals at the 5′ splice site of first introns (close to the promoter region) and the second row represents all other introns. **(C)** The same analysis performed in (B) is repeated for introns where the chromatin status remains the same, i.e. non-differential GCH methylation.