Process-explicit models reveal pathway to extinction for woolly mammoth using pattern-oriented validation

Damien A. Fordham[1,2]*, Stuart C. Brown[1], H. Reşit Akçakaya[3], Barry W. Brook[4], Sean

Haythorne[1], Andrea Manica[5], Kevin T. Shoemaker[6], Jeremy J. Austin[1], Benjamin Blonder[7],

Julia Pilowsky[1,2], Carsten Rahbek[2,8,9,10], David Nogues-Bravo[2].

[1]The Environment Institute and School of Biological Sciences, University of Adelaide,
Adelaide, South Australia, Australia.

[2]Center for Macroecology, Evolution, and Climate, GLOBE Institute, University of
Copenhagen, Copenhagen, Denmark

[3]Department of Ecology and Evolution, Stony Brook University, Stony Brook, New York,
United States of America

[4]School of Natural Sciences and ARC Centre of Excellence for Australian Biodiversity and
Heritage, University of Tasmania, Hobart, Tasmania, Australia

[5]Department of Zoology, University of Cambridge, Cambridge, England

[6]Department of Natural Resources and Environmental Science, University of Nevada, Reno,
Nevada, United States of America

[7]Department of Environmental Science, Policy, and Management, University of California,
Berkeley, California, United States of America

[8]Department of Life Sciences, Imperial College London, Ascot, England

[9]Danish Institute for Advanced Study, University of Southern Denmark, Odense, Denmark.

[10]Institute of Ecology, Peking University, Beijing, China

*Corresponding author: Damien Fordham, University of Adelaide, Adelaide, South

Australia, Australia, 5005 **(email:** <u>damien.fordham@adelaide.edu.au; ph:</u> +61 8 8313 6711;

<u>fax:</u> +61 8 83134347).

Running Title: Decline and extinction of woolly mammoth

Article Type: Letter

Statement of authorship: D.A.F, D.N.B. and C.R. conceived the idea for the paper. D.A.F. and S.C.B. did the analysis with support from H.R.A, B.W.B, S.H., A.M., K.T.S, J.J.A., J.P. and B.B. All authors contributed to writing the manuscript

Word Count: 155 (abstract); 4,543 (main text)

References, Figures and Tables: 60 references; 5 figures; 0 tables

Abstract

Pathways to extinction start long before the death of the last individual. However, causes of early-stage population declines and the susceptibility of small residual populations to extirpation are typically studied in isolation. Using validated process-explicit models, we disentangle the ecological mechanisms and threats that were integral in the initial decline and later extinction of the woolly mammoth. We show that reconciling ancient DNA data on woolly mammoth population decline with fossil evidence of location and timing of extinction requires process-explicit models with specific demographic and niche constraints, and a constrained synergy of climatic change and human impacts. Validated models needed humans to hasten climate-driven population declines by many millennia, and to allow woolly mammoths to persist in mainland Arctic refugia until the mid-Holocene. Our results show that the role of humans in the extinction dynamics of woolly mammoth began well before the Holocene, exerting lasting effects on the spatial pattern and timing of its range-wide extinction.

Introduction

Despite knowledge from the early 19[th] century (Cuvier 1807) that species go extinct, ecological mechanisms that underpin extinctions remain poorly resolved (Beissinger 2000; Channell & Lomolino 2000; Traill *et al.* 2007). This is because pathways to extinction can begin long before the extinction event, resulting from driver-state relationships that are difficult to detect and disentangle (Soulè 1983; Caughley 1994). In contrast, the demographic, ecological and genetic processes that make small populations susceptible to eventual extinction are better established (Lande 1993; Frankham 2005). Here we develop a process-explicit modelling framework that integrates the declining and small population paradigms, central to ecology and conservation biology (Caughley 1994), using pattern-oriented validation (Grimm *et al.* 2005). We apply it to the range collapse and extinction of the woolly mammoth (*Mammuthus primigenius*) to develop a more holistic understanding of spatiotemporal extinction dynamics. The woolly mammoth was one of many large mammals that went extinct during the Pleistocene-Holocene transition (Barnosky *et al.* 2004; Cooper *et al.* 2015).

Global warming following the Last Glacial Maximum, which extended from the late Pleistocene to the early Holocene, resulted in regional temperature increases of 4 to > 10 °C (Clark *et al.* 2012). During this deglaciation phase, many megafaunal species (terrestrial taxa > 45 kg) became extinct (Barnosky *et al.* 2004) and many others suffered regional extirpations (Cooper *et al.* 2015). This biotic simplification radically changed the structure and function of ecosystems (Gill *et al.* 2009; Doughty *et al.* 2016). At the same time, Palaeolithic human populations were spreading and becoming more ubiquitous, facilitated by increases in primary productivity associated with climatic change (Eriksson *et al.* 2012; Timmermann & Friedrich 2016). What has remained fiercely contested is the relative role of human hunting and climate change, or a synergy of these impacts, on the fate of the

megafauna (Barnosky *et al.* 2004; Stuart *et al.* 2004; Stuart 2005; Nogués-Bravo *et al.* 2008;

Lorenzen *et al.* 2011; MacDonald *et al.* 2012; Cooper *et al.* 2015). Major barriers to a

resolution have included a sparse and uncertain fossil record (Haile *et al.* 2009), a lack of

high-resolution spatiotemporal projections of climatic change and human abundances

(Fordham *et al.* 2018), reliance on correlative rather than process-based approaches to infer

drivers of extinction from ecological and molecular data (Fordham *et al.* 2020), and a focus

on extinctions of populations once at critically small thresholds (Palkopoulou *et al.* 2015;

Graham *et al.* 2016; Rogers & Slatkin 2017), rather than the causes of smallness itself

(Caughley 1994).

Recent developments in macroecological modelling are enabling the drivers and

processes of megafauna extinctions to be unravelled from the point of initial population

decline to the final extinction event, using a wide body of evidence from paleo-archives

(Fordham *et al.* 2020). These novel approaches, which simulate the dynamics of an

ecological system as explicit driver-state relationships (Rangel *et al.* 2018), offer fresh

opportunities to disentangle the mechanisms responsible for ecological responses to climate-

and human-driven changes in species distributions and abundances across space and time

(Fordham *et al.* 2016a). Unlike correlative approaches, fossil and molecular signatures of past

demographic change are used as independent, objective targets for directly evaluating a

model's structural adequacy and parameterization (Grimm *et al.* 2005). This new pattern-

oriented approach to refining and validating process-explicit models of species' range

dynamics allows relevant global change drivers and ecological processes to be simulated and

tested, revealing the most likely chains of causality that lead to extinction (Fordham *et al.*

2021).

The iconic woolly mammoth was present on earth for more than half a million years

(Bevan *et al.* 2017) before going extinct in the mid-Holocene (Stuart *et al.* 2004). During this

time, woolly mammoths co-existed with Neanderthals (*Homo neanderthalensis*) and modern

humans (*H. sapiens*) for many millennia, and were exploited for meat, skins, bones and ivory

(Stuart *et al.* 2004; Stuart 2005; Nogués-Bravo *et al.* 2008). As the earth warmed rapidly

during the last deglaciation (approximately 19 to 11 ka (thousand years ago) BP), boreal

forests spread throughout Eurasia, replacing tundra grassland and forbs (Binney *et al.* 2017),

the preferred habitat for woolly mammoths.

Previous process-explicit examinations of the extinction dynamics of woolly

mammoth focused largely on threats to persistence for populations at already critical

thresholds (Palkopoulou *et al.* 2015; Graham *et al.* 2016; Rogers & Slatkin 2017), concluding

that genomic meltdown through inbreeding caused its extinction (Palkopoulou *et al.* 2015;

Rogers & Slatkin 2017). Where long-lasting roles of climate and humans have been

considered, their function in setting the location and timing of woolly mammoth extinction

(and extinctions of other large mammals) during the Pleistocene-Holocene transition has been

inferred from snapshots (points in time ≥ 12,000 years apart) of projected historical range

movement (Nogués-Bravo *et al.* 2008; Lorenzen *et al.* 2011) and analysis of time-binned

fossil and archaeological data (Stuart *et al.* 2004; MacDonald *et al.* 2012), using correlative

(not process-explicit) modelling methods. Consequently, the spatiotemporal dynamics of

extinction forces, and resultant long-term patterns in population and range collapse, remain

unclear.

To unravel the determinants of early-stage population declines, and subsequent range

collapse and extinction of woolly mammoth in Eurasia, we built process-explicit spatially

dynamic macroecological models that simulate how ecological requirements and

demographic processes interact with climate change and human pressures to affect the

geographical range, population dynamics and the range-wide extirpation pattern of woolly

mammoths from 21 ka BP (kilo annum Before Present). We explored more than 90,000

scenarios, deriving best estimates of model parameters with pattern-oriented validation

methods. As validation targets, we used spatiotemporal inferences of local extinctions and

demographic change from hundreds of radiocarbon dated fossils and ancient DNA sequences.

These validation targets identified models with the combinations of ecological processes

(niche and demographic constraints for movement and extinction) and rates of global change

(climate-change, human-impact and their interaction) that best reconcile with ancient DNA

data on the timing and rate of population decline, along with fossil evidence of the extirpation

pattern, and location and timing of eventual species extinction. In doing so, our new

macroecological modelling approach was able to directly disentangle in space and time the

processes and threats crucial to the initial population decline and later extinction of the

woolly mammoth, revealing that its pathway to extinction started long before the final

extinction event.

Materials and Methods

The process-explicit macroecological model of climate-human-woolly mammoth interactions

we built was designed to reconstruct the pattern of range collapse, population decline and

extinction of woolly mammoth in Eurasia using pattern-oriented modelling methods (Grimm

et al. 2005) and spatiotemporal evidence from hundreds of radiocarbon dated fossils and

ancient DNA sequences (Fig 1 in S1 Text). Driver-state relationships simulated the effects of

climate change and hunting by humans on key ecological processes of extinction: niche

lability, dispersal, population growth and Allee effect. Models were coded in Program R

(available here: https://doi.org/10.5281/zenodo.5567859) and are described in detail in the

Methods in S1 Text.

Woolly mammoth niche

Radiocarbon dated and georeferenced fossils for the woolly mammoth (*M. primigenius*)

during the Late Pleistocene and Holocene were sourced from publicly accessible databases

and published literature (Fordham & Brown 2020). Their age reliability was assessed

(Barnosky & Lindsey 2010) and all reliable ages were calibrated using OxCal (Bronk

Ramsey 2009) and the IntCal13 calibration curve (Reimer *et al.* 2013). See Methods in S1

Text.

The TraCE-21 simulation of the transient climate of the last 21,000 years was used to

generate monthly mean climatic parameters from 21 ka BP to 0 BP (Fordham *et al.* 2017).

HadCM3 paleoclimate simulations from 60 to 21 ka BP (Singarayer & Valdes 2010) were

harmonized with TraCE-21 simulations and resampled to a $1 \times 1°$ resolution (Methods in S1

Text). We intersected fossil locations and time periods (calibrated age $\pm$ 1 SD) with

paleoclimate simulations of climatic parameters that affect the population dynamics of large

herbivores in polar regions (Sæther 1997) and characterized an n-dimensional hypervolume

of climatic suitability through time (Fig 2 in S1 Text), generating a biologically relevant

representation of the climate history over which woolly mammoths were present at fossil site

(Nogués-Bravo 2009). The resulting hypervolume of climate suitability, which approximates

the fundamental niche of the woolly mammoth (Nogués-Bravo 2009), was exhaustively

subsampled (Methods in S1 Text). This allowed the realized niche of the woolly mammoth

(Fordham *et al.* 2016a) to be identified using process-explicit macroecological modelling (see

below). Subsampling was done with Outlier Mean Index (OMI) analysis, using plausible

bounds of climatic specialization and niche breadth (Dolédec *et al.* 2000).

Spatial projections of climate suitability for woolly mammoths in Eurasia were

generated from 21 ka to 0 BP at 25-year time steps for each realized niche (n = 1,862

subsamples of the full hypervolume of climate suitability), using a standard maximum

entropy method (Phillips *et al.* 2006), accounting for latitudinal variation in grid-cell size and

temporal variation in the proportion of a cell that are land or sea ice (Methods in S1 Text).

Humans

The peopling of Eurasia by Palaeolithic humans, and their relative abundance, was modelled

using a Climate Informed Population Genetics Model (CISGEM) that accurately reconstructs

arrival times of anatomically modern humans and current-day distributions of global and

regional genetic diversity (Eriksson *et al.* 2012; Raghavan *et al.* 2015). CISGEM simulates

local effective population sizes (N_e) as a function of genetic history, local demography,

paleoclimate and vegetation. Like other numerical models of early human migration

(Timmermann & Friedrich 2016), arrival, occupancy and density (here N_e) are forced by

spatiotemporal estimates of climate and sea level changes over the past 125 k years. To

account for parameter uncertainty in spatiotemporal projections of N_e we ran thousands of

different models, each with a unique combination of parameter settings (based on established

upper and lower confidence bounds; Eriksson *et al.* 2012), selected using Latin hypercube

sampling (McKay *et al.* 1979). These N_e values were scaled between 0 and 1 and used in the

process-explicit macroecological model as potential spatiotemporal measures of relative

abundance (n = 10,000 potential scenarios of migration and population growth). CISGEM

and its application is described in detail in the Methods in S1 Text.

Climate-human-woolly mammoth interactions

Extinction and colonization dynamics were simulated as landscape-level population

processes, operating at 25-year generational time steps from 21 ka BP. Models centred on

'best estimates' for demographic processes (population growth rate and its variance,

dispersal, Allee effect), environmental attributes (niche breadth and climatic specialisation)

and threats (human abundance and rates of exploitation) were varied across wide but

plausible ranges using Latin hypercube sampling of uniform probability distributions, to

provide a robust coverage of multi-dimensional parameter space (Fordham *et al.* 2016b). This

procedure produced > 90,000 conceivable models (parameterizations) with different

combinations of rates of climate change, hunting by humans and demographic processes.

Each model was run for a single replicate (Prowse *et al.* 2016). The structure and parameters

of the process-explicit model are described in detail in the Methods in S1 Text.

Pattern-oriented modelling (Grimm *et al.* 2005) was used to evaluate different model

versions and parameterization, by cross-matching simulations with inferences from paleo-

archives (Fordham *et al.* 2020), using Approximate Bayesian Computation (ABC; Csilléry *et*

al. 2010). Specifically, simulations of range and extinction dynamics were validated against a

four-parameter multivariate target, consisting of trend in total population size (based on N_e),

inferred from ancient DNA (Lorenzen *et al.* 2011); time and location of range-wide

extinction, inferred from the fossil record (corrected for the Signor-Lipps effect; Saltré *et al.*

2015); and occupancy at fossil sites. The top 1% of feasible parameterizations of climate-

human-woolly mammoth interactions were retained (n = 900) and used to generate ensemble

averaged estimates of spatial abundance, timing of extirpation (extinction at the grid cell),

total population size, probability of occupancy and hunting rates. Estimates were weighted by

the Euclidean distance of the model from the idealized targets.

Our process-explicit modelling approach also permits possible alternatives to past

events to be simulated and the biological consequences assessed (Fordham *et al.* 2020). For

example, we held human-hunting parameters constant at zero harvesting in the best 1% of

validated models and then analysed the effect of this constraint on dynamical processes and

emergent patterns, and compared these to model simulations of climate and human

interactions with the woolly mammoth.

Statistical analysis

To identify and evaluate the processes and drivers that caused the initial population collapse

of woolly mammoth, and later the susceptibility of small residual populations to eventual

extinction (declining and small population paradigms; Caughley 1994), we discretised results

from the best 900 models into three distinct climatic periods (T1 = 21-15 ka BP, T2 = 15-11

ka BP and T3 = 11-5 ka BP) (Clark *et al.* 2012) and three sub-regions (Europe, Asia,

Beringia). For each period and region (including all of Eurasia) we computed the magnitude

and pace of climatic change, human population growth and expansion (Fordham & Brown

2020). We calculated Expected Minimum Abundance (EMA) at the termination of each

period (T1-T3) for each region. EMA quantifies risks of both overall population decline

(quasi-extinction) and total extinction (McCarthy & Thompson 2001). Statistical learning

models (Wright & Ziegler 2017) were used to identify spatiotemporal determinants of

extinction risk (Pearson *et al.* 2014). Spatiotemporal autocorrelation between climatic and

human drivers of extinction was calculated using the Lee's L Statistic (Lee 2001). Phase

synchrony and peak coincidence was calculated using the synchrony package for R (Gouhier

& Guichard 2014).

Results

Successfully simulating vital aspects of the range collapse and extinction of woolly

mammoth, inferred from paleo-archives, required highly constrained environmental attributes

(climatic requirements) and demographic mechanisms in our process-explicit model (Fig 1).

This indicates that ecological niche requirements and individual fitness at small population

size are likely to have been crucial drivers of the extinction dynamics of woolly mammoth in

continental Eurasia. Our prior-to-posterior checks show that non-marginal niches (niches

with low specialisation), low maximum abundances and a small Allee effect is needed to

simulate both the timing and rate of population decline (based on ancient DNA) and the

location and timing of range-wide extinction (based on the fossil record) (Table 1 in S1 Text).

By comparison, posterior distributions for exploitation parameters (including harvest rate)

more closely matched prior distributions (Fig 1).

The best models (top 1% of all models, which most closely matched the validation

targets) reconstructed a north-eastward range contraction for woolly mammoth from 19 ka

BP, with extirpation in most of Europe by ~ 14 ka BP (Fig 2), with the exception of refugial

areas in what is now Britain, northern France and Belgium (as well as pockets in the

Netherlands and Denmark), where patches of steppe-tundra ecosystems are likely to have

persisted in favourable climatic areas until the termination of the Pleistocene (Binney *et al.*

2017). These best models simulated an accelerated rate of range collapse in Asia following

rapid warming at ~ 15 ka BP (S1 Movie), with populations persisting within the Siberian

refugia of Taimyr, Beringia and the Yamal Peninsula, in accordance with fossil remains

(Stuart *et al.* 2004; Stuart 2005). Furthermore, these models accurately projected a steep

decline in total population size during the Late Pleistocene, as inferred here from ancient

DNA estimates of effective population size (Fig 2). Our simulations, however, did not project

a continent-wide extirpation during the early Holocene.

To converge on the validation targets, simulations needed woolly mammoth to persist

in Beringia, along the Kara Sea (including the Taimyr Peninsula) and, possibly in northern

Fennoscandia (following the retreat of glacial ice sheets), until at least the mid-Holocene (Fig

2). Model agreement for persistence in these mainland Arctic refugia until the mid-Holocene

was generally high in the best 1% of models (S1 Movie), pinpointing locations of Holocene-

age refugia. These refugial locations are likely, given the incompleteness of the fossil record

(Fig 3 in S1 Text) and low numbers of mammoths projected during the Holocene (Fig 2). In

contrast, refugia projected in the high elevation plateaux of southern Asia, and in Svalbard,

during the Holocene had low model agreement: on average < 9% probability of occurrence

(S1 Movie).

Spatiotemporal determinants of extinction

Across Eurasia, climatic change during the last deglaciation and the Holocene affected the

geography of human and woolly-mammoth abundance differently based on the best models

of climate-human-woolly mammoth interactions (Fig 3). Accelerated warming following

Heinrich Event 1 (~ 17.5 ka BP) resulted in climatically preferred conditions (temperature

and precipitation) for mammoths and humans becoming decoupled (mean Lee's L statistic of

spatial autocorrelation = – 0.07 between 17.5 and 15 ka BP), due likely to humans responding

to environmental change by colonizing and remaining resident in new niches (Giampoudakis

et al. 2017), and the woolly mammoth retreating to the coldest areas of its climatic niche,

where conditions for people were most harsh.

Using statistical learning analysis to identify spatiotemporal determinants of

extinction risk from these best models, we show that during the last deglaciation, climatic

shifts alone explained 52 and 57 % of the variance in expected minimum abundance (EMA)

of woolly mammoth in Eurasia at 15 ka BP and 11 ka BP, respectively (Fig 3). Larger

magnitudes of change in the climatic conditions suitable for woolly mammoth persistence,

along with a faster pace of loss of these conditions, resulted in increased extinction risk

during the periods 21-15 ka BP (T1) and 15-11 ka BP (T2) (Fig 4 in S1 Text). The magnitude

and pace of human population growth became as important as climatic change during the

Holocene in influencing the EMA of woolly mammoth in Eurasia at 5 ka BP (Fig 3). A lower

explained variance in the Holocene compared to earlier time periods occurs because pattern-

oriented validation procedures retain models that simulate mechanisms of Late Pleistocene

range collapse and population decline (those occurring in T1 and T2) having a long-lasting

effect on the timing and location of extinction in the Holocene (T3; 11-5 ka BP).

Demographic responses of woolly mammoths to impacts of humans and climate were

spatiotemporally heterogenous, with these differences being essential for setting the time and

place for extinction. Populations of woolly mammoths in distinct regions of Eurasia

experienced very different magnitudes of climatic and human impacts over time (Fig 5 in S1

Text), suggesting that the dynamics of extinction determinants were labile. Accurately

reconstructing inferences of range collapse and population change from fossils and ancient

DNA required that humans impacted woolly mammoth prior to the Holocene in Europe.

The magnitude and pace of climatic and human impacts during the last deglaciation in

Europe explained relatively similar amounts of variance in EMA of woolly mammoth (Fig 4).

During this period, humans and mammoths responded differently to climatic change in

Europe (mean Lee's L = 0.21 for the last deglaciation, indicating low autocorrelation),

confirming direct and important roles of hunting and climatic change on mammoth EMA.

Together these results indicate that humans had a sustained effect on the population dynamics

and extirpation of woolly mammoths in Europe. For the few populations in northern

Fennoscandia that were simulated to have colonized the region following the retreat of glacial

ice sheets (S2 Movie), gradual climatic change during the Holocene was the greatest threat to

simulated persistence (Fig 4) not hunting by humans, which remained relatively low (S3

Movie), owing to small population densities of humans in northern Fennoscandia (S4 Movie).

In Asia and Beringia, the role of humans on the range collapse and extirpation of

woolly mammoth more closely mirrored the pattern of continent-wide extinction observed for

Eurasia, with humans having a proportionately larger threatening influence on EMA by the

Holocene (Fig 5 in S1 Text). In Beringia, climatic changes during the last deglaciation and

Holocene affected humans and mammoths more similarly than for other regions of Eurasia

(Fig 5 in S1 Text; mean Lee's L = 0.44 for the last deglaciation; and 0.35 for the Holocene),

indicating greater synchrony in shifts in range and abundance.

Climate-human interactions

Counterfactual models with no mammoth exploitation by humans provide further evidence

that humans acted to hasten the timing of range collapse and extinction of woolly mammoths.

We show that in the absence of humans and their interactions with climatic change, woolly

mammoths would have been more abundant across time (including during the Late

Pleistocene), and populations would have persisted for much longer, perhaps even avoiding

extinction within climatic refugia (Fig 5). Mid-to-late Holocene population sizes of woolly

mammoths were much larger in simulations in the absence of human harvesting (Table 2 in

S1 Text), causing a 24% increase in persistence beyond 3.8 ka BP; the estimated time of

extinction (95% confidence interval = 4089 to 3450 BP) based on the fossil record (Methods

in S1 Text). Although some model simulations with humans on the landscape also did not

result in range-wide extinction by 3.8 ka BP (Fig 6 in S1 Text), confidence intervals for EMA

intersect zero (Table 2 in S1 Text).

While simulations of mammoth abundance, with and without humans, were

asynchronous (except for Beringia prior to 15 ka BP), the extreme peaks and troughs in

abundance generally occurred at similar times in the simulations (Table 3 in S1 Text). This

was not the case for Europe during the early part of the last deglaciation (21 – 15 ka BP) and

Asia during the latter part of the last deglaciation (15 – 11 ka BP), confirming that the

strength of the regulatory role of humans on the extinction dynamics of woolly mammoth

was regionally and temporally variable.

Discussion

Macroecological models of mechanistic interactions of climate and humans with the woolly

mammoth show that its pathway to extinction was long and lasting, starting many millennia

before the final extinction event with successive regional extirpations rather than a rapid

range-wide collapse. We show that reconciling spatiotemporal evidence of the decline and

extinction of woolly mammoths from fossils and ancient DNA requires quite tight

demographic and niche constraints, and spatiotemporally variant rates of climatic change and

human impacts (and their synergy) since the Last Glacial Maximum. Our simulations reveal

that human population growth and northward migrations during the Late Pleistocene led to

the premature extirpation of populations of woolly mammoth in areas of Eurasia that were

climatically suitable into the Holocene, hastening climate-driven declines by up to 4,000

years in some regions. They also show that simulating important aspects of the extinction

dynamics of woolly mammoths as inferred from paleo-archives requires persistence in

mainland Arctic refugia until the mid-Holocene. Thus process-explicit models that

continuously simulate the causes of early-stage population declines, and later the

susceptibility of small residual populations to eventual extinction, provide new opportunities

to unravel ecological mechanisms of extinction that occurred in the ancient past.

The role of humans as a causative agent in the extinction of woolly mammoths is

likely to have been both direct and indirect. In addition to exploitation-driven changes in

demographic processes, the climate-change-facilitated co-occupation of steppe and forb

habitats by humans and woolly mammoths could have affected metapopulation structures and

processes by interrupting sub-population connectivity, affecting movement between resource-

rich zones (Cooper *et al.* 2015). Posterior distributions for exploitation parameters (including

harvest rate) for Eurasia-wide simulations more closely matched prior distributions when

compared to some ecological niche and demographic parameters, indicating reduced

parameter importance. The reduced importance of the role of human hunting in the extinction

dynamics of woolly mammoth at a continental scale likely reflects variable rates of hunting

of woolly mammoths by humans in space and time and a likelihood that, to some extent,

humans impacted mammoth metapopulation processes independent of hunting (e.g., via

habit change, with hunting being most damaging at specific 'pinch-points' such as

movement corridors).

Correctly simulating inferences of extinction dynamics from fossils and ancient DNA,

required the survival of woolly mammoth in mainland Arctic refugia until the mid-Holocene,

some 5,000 years longer than previously thought based on fossil evidence alone (Stuart *et al.*

2004; Nikolskiy *et al.* 2011; Dehasque *et al.* 2021). Extirpation and extinction events for

megafauna are commonly revised as younger fossils and environmental DNA are discovered,

often causing persistence to be extended by several millennia (Haile *et al.* 2009; Wang *et al.*

submitted). This is because actual extinction dates are often underestimated using date of last

fossil appearance, which usually records the last time a species was abundant not last

occurrence (Mann *et al.* 2019). This raises the real possibility that populations of woolly

mammoth persisted in mainland arctic refugia until the mid-Holocene, as predicted by our

model. Especially given that population abundances of mammoths during the Holocene

would have been low (Fig 2), indicator species of mammoth persisted in Siberia during the

mid-Holocene (Boeskorov 2020), the refugial areas that we pinpoint remain poorly sampled

(Fig 2 in S1 Text), and environmental DNA evidence of woolly mammoths in Beringia and

the Taimyr Peninsula to the mid-Holocene (Wang *et al.* submitted). The recent discovery of

persistence of woolly mammoths in Siberia to 3.9 ka using environmental metagenomics

(Wang *et al.* submitted) provides an important independent validation of our process-based

model (Grimm *et al.* 2005), indicating a strong ability of the simulations to detect hidden

refugia and unveil spatiotemporal pathways to extinction.

Although equifinality was avoided in our analyses, using multiple validation targets,

our finding that mammoths likely persisted in mainland refugia in Eurasia until the mid-

Holocene was dependent on currently available inferences of extinction from the fossil

record, estimates of population decline from ancient DNA, and projections of spatially and

temporally variant rates of climatic change and human impacts. While the fossil record for

woolly mammoth, during the Pleistocene-Holocene transition, is more complete then for

many other megafauna species, there are still vast areas of Eurasia which remain poorly

represented by fossil samples (Fig 3 in S1 Text). Likewise, projections of population growth

and migrations of people during the Late Pleistocene and Holocene do not yet account for

topographical processes and cultural changes (Eriksson *et al.* 2012; Timmermann & Friedrich

2016), which could potentially influence our results. Therefore, uncertainty in our estimates

of the extinction dynamics of woolly mammoths in space and time is likely to be reduced

through new fossil discoveries, and higher spatial resolution projections of the peopling of

Eurasia.

The prior and posterior parameter distributions of some demographic and exploitation

parameters in the process-explicit models were similar, indicating that a wide range of values

for these parameters will reconstruct spatiotemporal evidence of the decline and extinction of

woolly mammoths. For these non-identifiable parameters, it is important to consider whether

the empirical targets used for pattern-oriented modelling best fit the study animal and system

(Gelman *et al.* 2013), which they do for the woolly mammoth. Nevertheless, it does mean

that a variety of different parameter combinations can give a close fit to inferences of

extinction dynamics from the fossil record and ancient DNA, potentially resulting in different

ecological interpretations.

Here we opened a window into late Quaternary biodiversity dynamics using process-

based macroecological models, in order to synthesize disparate evidence from paleo-archives

(*sensu* Fordham et al. 2020), establishing that ecological pathways to extinction can start

many millennia before populations are at critically small thresholds. This reinforces the need

for long-term perspectives when testing theories and making generalisations regarding the

spatial dynamics of range collapses and extinctions of species. Our results emphasise that

extinctions can only be explained by combining the declining and small population

paradigms (Caughley 1994). They also highlight that range shifts during the late Quaternary

offer distinct opportunities to test the circumstances under which geographic ranges collapse

first along the periphery versus those which start within the range interior (Channell &

Lomolino 2000), particularly if they are reconstructed using process-explicit models and

pattern-oriented approaches (Fordham *et al.* 2021).

Our analyses strengthen and better resolves the case for human impacts as a crucial

and chronic driver of early-stage population declines of megafauna, revealing an essential

role of humans in population declines of mammoths in Eurasia during the Late Pleistocene; a

period when climatic conditions warmed rapidly. In doing so, it refutes a prevalent theory

that the role of humans in the extinction dynamics of woolly mammoths was limited to a mid-

Holocene coup de grâce (Nogués-Bravo *et al.* 2008), and highlights the importance of

disentangling demographic responses to varying biotic and abiotic stressors for

metapopulations at regional scales, particularly when assessing species survival under climate

change.

Acknowledgments

F. Saltré, K. Giampoudakis, A. Flórez-Rodríguez assisted with collating and analysing the

fossil record and ancient DNA.

References

Barnosky, A.D., Koch, P.L., Feranec, R.S., Wing, S.L. & Shabel, A.B. (2004). Assessing the Causes of Late Pleistocene Extinctions on the Continents. *Science*, 306, 70.

Barnosky, A.D. & Lindsey, E.L. (2010). Timing of Quaternary megafaunal extinction in South America in relation to human arrival and climate change. *Quaternary International*, 217, 10-29.

Beissinger, S.R. (2000). Ecological mechanisms of extinction. *Proc Natl Acad Sci U S A*, 97, 11688.

Bevan, A., Colledge, S., Fuller, D., Fyfe, R., Shennan, S. & Stevens, C. (2017). Holocene
fluctuations in human population demonstrate repeated links to food production and
climate. *Proc Natl Acad Sci U S A*, 114, E10524.

Binney, H., Edwards, M., Macias-Fauria, M., Lozhkin, A., Anderson, P., Kaplan, J.O. *et al.*
(2017). Vegetation of Eurasia from the last glacial maximum to present: Key
biogeographic patterns. *Quaternary Science Reviews*, 157, 80-97.

Boeskorov, G.G. (2020). Survival of indicator species of the mammoth fauna large mammals
in the Holocene of Yakutia (East Siberia, Russia). *IOP Conference Series: Earth and
Environmental Science*, 438, 012004.

Bronk Ramsey, C. (2009). Bayesian Analysis of Radiocarbon Dates. *Radiocarbon*, 51, 337-
360.

Caughley, G. (1994). Directions in Conservation Biology. *J Anim Ecol*, 63, 215-244.

Channell, R. & Lomolino, M.V. (2000). Dynamic biogeography and conservation of
endangered species. *Nature*, 403, 84-86.

Clark, P.U., Shakun, J.D., Baker, P.A., Bartlein, P.J., Brewer, S., Brook, E. *et al.* (2012).
Global climate evolution during the last deglaciation. *Proc Natl Acad Sci U S A*, 109,
E1134-1142.

Cooper, A., Turney, C., Hughen, K.A., Brook, B.W., McDonald, H.G. & Bradshaw, C.J.
(2015). Abrupt warming events drove Late Pleistocene Holarctic megafaunal
turnover. *Science*, 349, 602-606.

Csilléry, K., Blum, M.G.B., Gaggiotti, O.E. & François, O. (2010). Approximate Bayesian
Computation (ABC) in practice. *Trends Ecol Evol*, 25, 410-418.

Cuvier, M. (1807). XXXVII. Additional memoir upon living and fossil elephants. *The
Philosophical Magazine*, 29, 244-254.

Dehasque, M., Pečnerová, P., Muller, H., Tikhonov, A., Nikolskiy, P., Tsigankova, V.I. *et al.*
(2021). Combining Bayesian age models and genetics to investigate population
dynamics and extinction of the last mammoths in northern Siberia. *Quaternary
Science Reviews*, 259, 106913.

Dolédec, S., Chessel, D. & Gimaret-Carpentier, C. (2000). Niche separation in community
analysis: A new method. *Ecology*, 81, 2914-2927.

Doughty, C.E., Roman, J., Faurby, S., Wolf, A., Haque, A., Bakker, E.S. *et al.* (2016). Global
nutrient transport in a world of giants. *Proc Natl Acad Sci U S A*, 113, 868-873.

Eriksson, A., Betti, L., Friend, A.D., Lycett, S.J., Singarayer, J.S., von Cramon-Taubadel, N.
et al. (2012). Late Pleistocene climate change and the global expansion of
anatomically modern humans. *Proc Natl Acad Sci U S A*, 109, 16089-16094.

Fordham, D. & Brown, S.C. (2020). Appendix 1 - 3 for Process-explicit models reveal
pathway to extinction for woolly mammoth using pattern-oriented validation.
Available at: http://doi.org/10.25909/5f22592242ca2.

Fordham, D.A., Akcakaya, H.R., Alroy, J., Saltre, F., Wigley, T.M.L. & Brook, B.W.
(2016a). Predicting and mitigating future biodiversity loss using long-term ecological
proxies. *Nat Clim Change*, 6, 909-916.

Fordham, D.A., Haythorne, S. & Brook, B.W. (2016b). Sensitivity Analysis of Range
Dynamics Models (SARDM): Quantifying the influence of parameter uncertainty on
forecasts of extinction risk from global change. *Environmental Modelling & Software*,
83, 193-197.

Fordham, D.A., Haythorne, S., Brown, S.C., Buettel, J.C. & Brook, B.W. (2021). poems: R
package for simulating species' range dynamics using pattern-oriented validation.
Methods Ecol Evol.

Fordham, D.A., Jackson, S.T., Brown, S.C., Huntley, B., Brook, B.W., Dahl-Jensen, D. *et al.*
(2020). Using paleo-archives to safeguard biodiversity under climate change. *Science*,
369, eabc5654.

Fordham, D.A., Saltré, F., Brown, S.C., Mellin, C. & Wigley, T.M.L. (2018). Why decadal to
century timescale palaeoclimate data are needed to explain present-day patterns of
biological diversity and change. *Glob. Change Biol.*, 24, 1371-1381.

Fordham, D.A., Saltre, F., Haythorne, S., Wigley, T.M.L., Otto-Bliesner, B.L., Chan, K.C. *et*
al. (2017). PaleoView: a tool for generating continuous climate projections spanning
the last 21 000 years at regional and global scales. *Ecography*, 40, 1348-1358.

Frankham, R. (2005). Genetics and extinction. *Biol. Conserv.*, 126, 131-140.

Gelman, A., Carlin, J.B., Stern, H.S., Dunson, D.B., Vehtari, A. & Rubin, D.B. (2013).
Bayesian Data Analysis. Third edn. Chapman and Hall/CRC.

Giampoudakis, K., Marske, K.A., Borregaard, M.K., Ugan, A., Singarayer, J.S., Valdes, P.J.
et al. (2017). Niche dynamics of Palaeolithic modern humans during the settlement of
the Palaearctic. *Glob. Ecol. Biogeogr.*, 26, 359-370.

Gill, J.L., Williams, J.W., Jackson, S.T., Lininger, K.B. & Robinson, G.S. (2009). Pleistocene
megafaunal collapse, novel plant communities, and enhanced fire regimes in North
America. *Science*, 326, 1100-1103.

Gouhier, T.C. & Guichard, F. (2014). Synchrony: quantifying variability in space and time.
Methods Ecol Evol, 5, 524-533.

Graham, R.W., Belmecheri, S., Choy, K., Culleton, B.J., Davies, L.J., Froese, D. *et al.*
(2016). Timing and causes of mid-Holocene mammoth extinction on St. Paul Island,
Alaska. *Proc Natl Acad Sci U S A*, 113, 9310-9314.

Grimm, V., Revilla, E., Berger, U., Jeltsch, F., Mooij, W.M., Railsback, S.F. *et al.* (2005).
Pattern-Oriented Modeling of Agent-Based Complex Systems: Lessons from
Ecology. *Science*, 310, 987.

Haile, J., Froese, D.G., Macphee, R.D., Roberts, R.G., Arnold, L.J., Reyes, A.V. *et al.* (2009).
Ancient DNA reveals late survival of mammoth and horse in interior Alaska. *Proc*
Natl Acad Sci U S A, 106, 22352-22357.

Lande, R. (1993). Risks of Population Extinction from Demographic and Environmental
Stochasticity and Random Catastrophes. *The American Naturalist*, 142, 911-927.

Lee, S.-I. (2001). Developing a bivariate spatial association measure: An integration of
Pearson's *r* and Moran's *I. Journal of Geographical Systems*, 3, 369-385.

Lorenzen, E.D., Nogues-Bravo, D., Orlando, L., Weinstock, J., Binladen, J., Marske, K.A. *et*
al. (2011). Species-specific responses of Late Quaternary megafauna to climate and
humans. *Nature*, 479, 359-364.

MacDonald, G.M., Beilman, D.W., Kuzmin, Y.V., Orlova, L.A., Kremenetski, K.V., Shapiro,
B. *et al.* (2012). Pattern of extinction of the woolly mammoth in Beringia. *Nat*
Commun, 3, 893.

Mann, D.H., Groves, P., Gaglioti, B.V. & Shapiro, B.A. (2019). Climate-driven ecological
stability as a globally shared cause of Late Quaternary megafaunal extinctions: the
Plaids and Stripes Hypothesis. *Biological Reviews*, 94, 328-352.

McCarthy, M.A. & Thompson, C. (2001). Expected minimum population size as a measure
of threat. *Anim. Conserv.*, 4, 351-355.

McKay, M.D., Beckman, R.J. & Conover, W.J. (1979). A Comparison of Three Methods for
Selecting Values of Input Variables in the Analysis of Output from a Computer Code.
Technometrics, 21, 239-245.

Nikolskiy, P.A., Sulerzhitsky, L.D. & Pitulko, V.V. (2011). Last straw versus Blitzkrieg
overkill: Climate-driven changes in the Arctic Siberian mammoth population and the
Late Pleistocene extinction problem. *Quaternary Science Reviews*, 30, 2309-2328.

Nogués-Bravo, D. (2009). Predicting the past distribution of species climatic niches. *Glob.*
Ecol. Biogeogr., 18, 521-531.

Nogués-Bravo, D., Rodríguez, J., Hortal, J., Batra, P. & Araújo, M.B. (2008). Climate
Change, Humans, and the Extinction of the Woolly Mammoth. *PLoS Biol*, 6, e79.

Palkopoulou, E., Mallick, S., Skoglund, P., Enk, J., Rohland, N., Li, H. *et al.* (2015).
Complete genomes reveal signatures of demographic and genetic declines in the
woolly mammoth. *Curr Biol*, 25, 1395-1400.

Pearson, R.G., Stanton, J.C., Shoemaker, K.T., Aiello-Lammens, M.E., Ersts, P.J., Horning,
N. *et al.* (2014). Life history and spatial traits predict extinction risk due to climate
change. *Nat Clim Change*, 4, 217-221.

Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006). Maximum entropy modeling of
species geographic distributions. *Ecol. Model.*, 190, 231-259.

Prowse, T.A.A., Bradshaw, C.J.A., Delean, S., Cassey, P., Lacy, R.C., Wells, K. *et al.* (2016).
An efficient protocol for the global sensitivity analysis of stochastic ecological
models. *Ecosphere*, 7, e01238.

Raghavan, M., Steinrücken, M., Harris, K., Schiffels, S., Rasmussen, S., DeGiorgio, M. *et al.*
(2015). Genomic evidence for the Pleistocene and recent population history of Native
Americans. *Science*, 349, aab3884.

Rangel, T.F., Edwards, N.R., Holden, P.B., Diniz-Filho, J.A.F., Gosling, W.D., Coelho,
M.T.P. *et al.* (2018). Modeling the ecology and evolution of biodiversity:
Biogeographical cradles, museums, and graves. *Science*, 361, eaar5452.

Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B. *et al.* (2013).
IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP.
Radiocarbon, 55, 1869-1887.

Rogers, R.L. & Slatkin, M. (2017). Excess of genomic defects in a woolly mammoth on
Wrangel island. *PLoS Genetics*, 13, e1006601.

Sæther, B.-E. (1997). Environmental stochasticity and population dynamics of large
herbivores: a search for mechanisms. *Trends Ecol Evol*, 12, 143-149.

Saltré, F., Brook, B.W., Rodríguez-Rey, M., Cooper, A., Johnson, C.N., Turney, C.S.M. *et al.*
(2015). Uncertainties in dating constrain model choice for inferring extinction time
from fossil records. *Quatenary Science Reviews*, 112, 128-137.

Singarayer, J.S. & Valdes, P.J. (2010). High-latitude climate sensitivity to ice-sheet forcing
over the last 120 kyr. *Quaternary Science Reviews*, 29, 43-55.

Soulè, M.E. (1983). What do we really know about extinction? In: *Genetics and
Conservation: a reference for managing wild animals and plant populations* (ed. C.
M. Schonewald-Cox et al). Benjamin/Cummings Publishing.

Stuart, A.J. (2005). The extinction of woolly mammoth (*Mammuthus primigenius*) and
straight-tusked elephant (*Palaeoloxodon antiquus*) in Europe. *Quaternary
International*, 126-128, 171-177.

Stuart, A.J., Kosintsev, P.A., Higham, T.F.G. & Lister, A.M. (2004). Pleistocene to Holocene
extinction dynamics in giant deer and woolly mammoth. *Nature*, 431, 684-689.

Timmermann, A. & Friedrich, T. (2016). Late Pleistocene climate drivers of early human
migration. *Nature*, 538, 92-95.

Traill, L.W., Bradshaw, C.J.A. & Brook, B.W. (2007). Minimum viable population size: A
meta-analysis of 30 years of published estimates. *Biol. Conserv.*, 139, 159-166.

Wang, Y., Pedersen, M.W., Alsos, I.G., De Sanctis, B., Racimo, F., Coissac, E. *et al.* 2021.
Late Quaternary dynamics of Arctic biota revealed by ancient environmental
metagenomics. Nature, https://doi.org/10.1038/s41586-021-04016-x

Wright, M.N. & Ziegler, A. (2017). ranger: A Fast Implementation of Random Forests for
High Dimensional Data in C++ and R. *Journal of Statistical Software; Vol 1, Issue 1*
(2017).

Figures

610
611
Fig. 1. Posterior distribution of parameter values. Distributions for prior values (grey),
and posterior values for ecological niche requirements (green), exploitation parameters (blue)
and demographic parameters (orange). Values on the y-axis have been scaled so that the
maximum height of each density plot is 1. Values on the x-axis are centred and scaled. Box
plots show the 25th, 50th and 75th percentiles. Whiskers show maximum values. Ecological
niche requirements are distance between average climatic conditions of the occupied niche
and the average climatic condition in the study region (Marginality); and breadth of climatic
conditions the species can occupy (Breadth). Harvest parameters are percentage of the
population that is harvested (Harvest); extent to which harvest follows a Type II to Type III
functional response (Func. resp.); and maximum human abundance in a grid cell (Hum.
dens.). Demographic processes are Allee effect, maximum abundance (Density); maximum
dispersal distance (Long disp.); mean dispersal rate (Mean disp.); maximum population
growth rate (Pop growth); and variation in population growth rate (Var growth). See Methods
in S1 Text and Appendix 3 in Fordham and Brown (2020) for more details.

Fig 2. Population decline and extinction of the woolly mammoth in Eurasia. (A) Simulated timing of extirpation (model weighted) for woolly mammoth in Eurasia (ka BP). **(B)** Ancient DNA estimates of effective population size (N_e) (blue line = mean, blue banding = 95% CI) and simulated total population size (black line = weighted mean, grey lines = range) rescaled between 0 and 1. Green line shows change in mean annual temperature for Eurasia. Vertical dashed lines show Heinrich 1 and 14.7 k climatic events. Red band shows estimated time of mammoth extinction.

Fig 3. Effects of humans and climate on the decline of woolly mammoth across Eurasia.
Drivers of expected minimum for the periods 21-15 ka BP (**A**), < 15-11 ka BP (**B**), < 11-5 ka
BP (**C**) for the 1% best validated models. Pie charts show variance explained (%) by climate
(blue), humans (orange) and area of occupancy at the start of each period (grey). Histograms
show contribution to explained variance for magnitude (M) or pace (P) of variables detailed
in Appendix 1 of Fordham and Brown (Fordham & Brown 2020). White areas of pie chart
represent unexplained variance. (**D**) shows the relationship (Lee's L statistic) between human
abundance and climate suitability for mammoths across time for Eurasia (black line =
weighted mean, grey lines = individual model runs). Positive values indicate positive
correlation, whilst negative values indicate a decoupling of the variables (negative
correlation). Fig 7 in S1 Text shows variable importance for area of occupancy.

648
649 **Fig 4. Extinction risk in Europe for woolly mammoth due to humans and climate. (A**)
650 Variable importance (histograms) and variance explained (pie charts) for climate (top row)
651 and human parameters (middle row). Fig 2 describes the legend. (**B**) Lee's L statistic of
652 autocorrelation between human abundance and climate suitability for mammoths.
653

654
Fig 5. Footprint of humans on the extinction dynamics of woolly mammoths. Relative
change in expected minimum abundance of woolly mammoth in response to climate change
and exploitation from 21 ka BP (Humans) and a counterfactual no-exploitation scenario (No
humans) for Europe (**A**), Asia (**C**) and Beringia (**E**). Maps show the difference in timing of
extirpation for woolly mammoths in the absence of hunting by humans (i.e. No humans –
Humans) for Europe (**B**), Asia (**D**) and Beringia (**F**). Positive values indicate a later
extirpation date in the absence of humans. Differences of ± 500 years are not shown. Pink
lines mark regional boundaries.
663

Supporting Information

S1 Text. Supporting methods, figures, and tables

S1 Movie. Probability of occupancy for woolly mammoth in Eurasia from 21,000 BP

S2 Movie. Density estimates of woolly mammoth in Eurasia from 21,000 BP

S3 Movie. Harvest rates of woolly mammoth in Eurasia from 21,000 BP

S4 Movie. Effective population size of humans in Eurasia from 21,000 BP