Title: Caffeinated soda intake in children is associated with neurocognitive vulnerabilities of

substance misuse and is predictive of a higher risk of alcohol sipping after 12 months

Authors:

Mina Kwon[1], Hyeonjin Kim[1], Jaeyeong Yang[1], Yoseph Lee[1], Jihyun K. Hur[1,2], Tae-Ho Lee[3], and

Woo-Young Ahn[1,4]

[1]Department of Psychology, Seoul National University, Seoul, Korea 08826

[2]Department of Psychology, Yale University, New Haven, CT 06511

[3]Department of Psychology, Virginia Tech, Blacksburg, VA 24060

[4]Department of Brain and Cognitive Sciences, Seoul National University, Seoul, Korea 08826

Corresponding author:

Woo-Young Ahn, Ph.D.

Department of Psychology

Seoul National University

Seoul, Korea 08826

Tel: +82-2-880-2538, Fax: +82-2-877-6428. E-mail: wahn55@snu.ac.kr

Running head: Soda intake in children and alcohol sipping

Word count:

Number of Figures & Tables: 4 / 1

Number of words in the introduction: 662

Number of words in the method: 530

Number of words in the results: 779

Number of words in the discussion: 1,431

Declaration of interests: None to declare

Funding:

This work was supported by the Basic Science Research Program through the National Research

Foundation (NRF) of Korea funded by the Ministry of Science, ICT, & Future Planning (NRF-

2018R1C1B3007313 and NRF-2018R1A4A1025891); the Institute for Information &

Communications Technology Planning & Evaluation (IITP) grant funded by the Korea government

(MSIT) (No. 2019-0-01367, BabyMind); the National Research Foundation of Korea (NRF) grant

funded by the Korea government (MSIT) (No. 2021M3E5D2A0102249311); and the Creative-

Pioneering Researchers Program through Seoul National University to W.-Y.A.

ABSTRACT

Background and Aims Caffeinated soda contains two addictive substances, sugar and caffeine, and is the most preferred means of caffeine consumption among children. However, it remains unexplored if habitual caffeinated soda intake in childhood is associated with a higher risk of alcohol misuse in the future. Here, we investigated the neurocognitive correlates of caffeinated soda consumption and examined whether caffeinated soda intake is associated with a higher risk of alcohol initiation in children. **Methods** Using the Adolescent Brain Cognitive Development Study, we first investigated the relationships between frequent caffeinated soda intake and well-known risk factors of substance misuse: impaired working memory, high impulsivity, and blunted reward processing. We next examined whether caffeinated soda intake predicts more alcohol sipping after 12 months, and whether caffeinated soda intake mediates the link between neurocognitive risk factors and future alcohol sipping. **Results** Frequent consumption of caffeinated soda was associated with neurocognitive risk factors for substance misuse. Daily soda intake was associated with hyperactivity in the striatal and parietal regions and hypoactivity in the inferior frontal gyrus (IFG) during the emotional N-Back Task. In addition, the striatum, IFG, and anterior cingulate cortex (ACC) showed aberrant activity during the stop signal task, and the medial prefrontal cortex and the ACC exhibited increased and decreased activity, respectively, during the monetary incentive delay task. Furthermore, caffeinated soda intake predicted greater alcohol sipping after 12 months, and mediated the relationship between impulsivity and future alcohol sipping. **Conclusions** This study addresses the question that many parents are asking: "Is frequent caffeinated soda intake in childhood associated with a higher risk of substance misuse in the future?" We showed that caffeinated soda consumption in childhood was associated with impaired neurocognitive functioning and may predict future alcohol intake, which has significant public health implications.

INTRODUCTION

Soft drinks are commonly consumed even by children, and a vast majority of sodas contain caffeine [1–3]. Caffeinated soda refers to soft drinks containing caffeine as well as sweeteners and artificial flavorings. Thus, caffeinated soda combines the addictive substances of sugar and caffeine, both of which affect neurocognitive function and cause physical side effects, although the concept of sugar addiction remains controversial [4–6]. Not surprisingly, the consequences of excessive consumption of both sugar and caffeine have been well documented [7], including a strong association between caffeinated beverage consumption in adolescents and future substance use [8–13]. In prospective studies tracking the effects of substance use, the percentage of regular energy drink users who became alcohol or marijuana users after 1–2 years was approximately five times higher than that of non-energy drink users [9,12,13]. Others have shown that coffee or energy drink consumption in children significantly predicts future substance use, such as tobacco and alcohol use [8,10,12].

Previous studies examining the association between caffeine consumption and later substance use have focused more on adolescents who drink multiple caffeinated beverages on a daily basis, partially because the consumption level of coffee and energy drinks is extremely low in children (0.4% for coffee and < 0.1% for energy drinks in 9–10-year-old children) [14]. However, caffeinated soda is the most preferred and accessible mode of caffeine intake in children [1,4,15]. Moreover, it is well known that adolescence is the most common period for initiating substance use, and an earlier onset of substance use predicts greater addiction severity [16–19]. Therefore, there is a critical need to investigate the association between caffeinated soda intake during preadolescent childhood and the risk of substance use disorders (SUDs) in the future, including vulnerabilities, such as associated neurocognitive deficits or early exposure to substances of abuse. Environmental factors, particularly the role of family, should be examined together, considering the previous studies that suggested a critical role of parental interventions in youth soft drink consumption and substance use [20–26]. Few studies have directly examined the effects of frequent caffeinated soda consumption in adolescents and children, and they either

targeted only behavioral aspects using surveys or self-reports without obtaining neural data, or

targeted only adolescents and not children [9,27–29].

Here, we addressed the unanswered question of whether frequent consumption of

caffeinated soda in children indicates a higher risk of later substance misuse, using the

Adolescent Brain Cognitive Development (ABCD) Study. First, we examined the relationship

between caffeinated soda intake and three well-known neurocognitive risk factors for SUDs in

children: impaired working memory (WM), high impulsivity, and altered reward processing [30].

These are the three main risk factors targeted in the ABCD study due to their significant

implications for addiction (see [30] for more detail). We used psychological, behavioral, and

neurological indices related to each risk factor, from the baseline data of the ABCD study (**Figure**

1). We applied a machine learning approach to the indices so that we could identify multivariate

risk factors for SUDs associated with caffeinated soda intake [31]. We next examined whether

caffeinated soda intake could predict future alcohol sipping, as (1) alcohol sipping has been

reported to predict future alcohol abuse [32–34], and (2) alcohol sipping is the most common

gateway behavior toward other substances of abuse [19,35]. Lastly, we examined mediation

models explaining how the interaction between daily caffeinated soda intake and the SUD risk

factors leads to future alcohol sipping behavior. Based on the previous literature suggesting that

caffeinated beverage consumption predicts future substance use [8–10,12,13], we hypothesized

that caffeinated soda intake would mediate the relationship between neurocognitive risk factors

and future alcohol sipping (**Figure 1A**). We also evaluated family risk factors as mediators of this

relationship, based on the literature that family risk factors, including family substance use or

family support, are risk factors for substance use in adolescents [24–26]. Taken together, we

aimed to elucidate the potential underlying risks of frequent caffeinated soda intake during

childhood.

METHODS

Participants

We included 4,529 participants without any missing data pertaining to soda intake, risk factors for SUDs, alcohol sipping, and all the confounding variables. The participants were divided into four groups based on their caffeinated soda consumption: the non-drinking group (N=1,945), monthly drinking group (N=1,941), weekly drinking group (N=484), and daily drinking group (N=159) [36]. See **Supplementary Materials** for more details on the measures and participants.

Analysis

Caffeinated soda intake and risk factors for SUDs

To identify a multivariate profile of risk factors for caffeinated soda consumption, we performed a least absolute shrinkage and selection operator (LASSO) regression analysis [37], a machine learning algorithm, to distinguish the daily drinking group (N=159) from the non-drinking group (N=1,945). We focused on these two extreme groups to ensure adequate separation between habitual drinkers (daily drinking group) and non-drinkers (non-drinking group). We used all of the risk factors and confounding variables as input (i.e., predictors) in the LASSO-based prediction model to classify each individual into the daily drinking group (coded as "1") or non-drinking group (coded as "0). More specifically, candidate predictors included every measure for the three cognitive factors as well as other control variables. See **Supplementary Materials** for more details on the predictors and LASSO regression.

Caffeinated soda intake and future alcohol sipping

To rigorously examine the association between caffeinated soda intake and future alcohol sipping, we used univariate (hierarchical regression) and multivariate (machine learning) approaches. Hierarchical logistic regression analysis was used to test whether caffeinated soda intake at baseline can predict alcohol sipping after 12 months. To ensure that the main predictor

accounted for more variance than the confounding variables, we set a three-level hierarchical

model (**Table 1**). See **Supplementary Materials** for more details on the hierarchical regression.

We also performed multivariate LASSO regression analysis to confirm the association

between caffeinated soda intake and alcohol sipping. We used all possible variables, including

the three neurocognitive risk factors (WM, impulsivity, and reward processing) and the

confounding variables as candidate predictors collected at baseline. Using these inputs (i.e.,

predictors) and the LASSO-based model, we aimed to classify individuals who reported alcohol

sipping after 12 months (coded as "1") and those who reported no alcohol sipping (coded as "0).

All of the procedures, except for the predictors and dependent variables, are identical to those in

the former analysis (see **METHODS - *Caffeinated soda intake and risk factors for SUDs*).**

Caffeinated soda intake, alcohol sipping, and risk factors for SUDs

We performed the mediation analysis to provide a more integrated insight into how the

interactions between caffeinated soda intake and the SUD risk factors lead to future alcohol

sipping in children. We fit three mediation models for each SUD risk factor, with a mediation

design involving the following four elements: an independent variable (one neurocognitive risk

factor for one model - WM, impulsivity, and reward processing), two mediators (caffeinated soda

intake and family risk factors; covariance of these two mediators was included in the model), and

a dependent variable (alcohol sipping after 12 months). This design was chosen to identify a

neurocognitive risk factor that is mediated by caffeinated soda intake in predicting alcohol sipping.

See **Supplementary Materials** for more details on the mediation analysis.

RESULTS

Caffeinated soda intake and risk factors for SUDs

Our first question was whether caffeinated soda intake is associated with well-known

neurocognitive risk factors for SUDs (**Figure 1A**). **Figure 2A** shows the multivariate profiles from

the LASSO regression distinguishing the daily-drinking group from the non-drinking group. Sex

(being male) and low parental education were the two strongest predictors of daily consumption,

along with a family history of drug use, high Behavioral Activation System (BAS) scores [38], lack

of sleep, and low family income. **Figure 2B** shows a receiver-operation characteristic (ROC)

curve and its mean area under the curve (AUC) for classification of the daily drinking and non-

drinking groups. The mean AUC values were 0.82 and 0.73 for the training and test sets,

respectively.

Among the risk factors for SUDs, high impulsivity measured using the BAS was the most

predictive of daily caffeinated soda intake (**Figure 2C**). A higher Urgency–Premeditation–

Perseverance–Sensation Seeking–Positive Urgency (UPPS-P) impulsive behavior score [39] and

a lower Behavioral Inhibition System (BIS) score [38] were related to daily intake. Additionally,

hyperactivation in the nucleus accumbens (NAc) and pars triangularis in the inferior frontal gyrus

(IFG) and hypoactivation in the caudal anterior cingulate cortex (ACC) and caudate nucleus

during the stop signal task (SST) predicted the classification of the daily drinking group. Among

the WM measures, poor performance on the list sorting test, card sort test, and 0-back

performance in the emotional N-Back (EN-back) task predicted daily intake of caffeinated soda. In

addition, hypoactivation of the pars orbitalis of the IFG and greater activation of the NAc, caudate

nucleus, inferior parietal lobule (IPL), and thalamus proper during the EN-back task (2-back vs. 0-

back) also predicted daily caffeinated soda intake (**Figure 2C**). The daily drinking group showed a

smaller difference in accuracy between the reward and neutral trials on the monetary incentive

delay (MID) task compared to the non-drinking group, suggesting reduced motivation to achieve

the monetary reward. Additionally, hyperactivity in the medial orbitofrontal cortex (OFC) and

hypoactivity in the rostral ACC during the MID task (reward vs. neutral anticipation) predicted the

classification into the daily drinking group.

These results support that low WM, high impulsivity, and blunted reward anticipation are

significantly associated with daily caffeinated soda consumption. Along with some other

demographic factors (male sex, family history of drug use, low socioeconomic status (SES), low

parental monitoring, high externalizing behaviors, less sleep, high body mass index (BMI), older

age, and low physical activity), the three neurocognitive risk factors for SUDs distinguished the

daily drinking group from the non-drinking group.

Caffeinated soda intake and future alcohol sipping

We first conducted a three-level hierarchical logistic regression analysis to address our

second main question of whether caffeinated soda intake predicts future alcohol sipping, (**Table**

1). Specifically, we examined whether caffeinated soda intake accounted for additional variance

than did other confounding variables. The adjusted R-square value of Model 3 revealed a

significant difference compared with Model 1 (F(4, 8,854)=177.58, p<2.20e-16) and Model 2

(F(3,8,854)=6.61, p<0.001), supporting the hypothesis that caffeinated soda drinking behavior in

childhood can predict future alcohol drinking (**Figure 3A**). Higher caffeinated soda consumption

significantly predicted alcohol sipping after 12 months even after controlling for confounding

variables and baseline alcohol sipping (**Table 1**).

Next, we performed LASSO regression predicting alcohol sipping after 12 months, using

all of the neurocognitive risk factors and confounding variables collected at baseline. As shown in

Figure 3B, caffeinated soda intake (mean estimate of coefficients=0.138, 95% confidence

interval (CI)=[0.123, 0.155]) was identified as a predictor that distinguished alcohol sipping after

12 months, even after including alcohol sipping at baseline as predictors. The mean AUC values

of the predictive model were 0.81 and 0.76 for the training and test sets, respectively (**Figure 3C**).

Caffeinated soda intake, risk factors for SUDs, and future alcohol sipping

We examined the integrated relationships among the risk factors for SUDs, caffeinated

soda intake, and alcohol sipping after 12 months. We tested three mediation models using each

neurocognitive risk factor (WM, impulsivity, and reward processing) as the independent variable.

The models that used WM or reward processing as an independent variable showed inconclusive

total effects (WM model: B=−0.007, 95% CI=[−0.013. −0.001], *p*=0.232; reward processing

model: B=0.004, 95% CI=[−0.002. 0.010], *p*=0.457). However, as shown in **Figure 4**, the model

using impulsivity as an independent variable revealed significant total effects (*B*=0.023, 95%

CI=[0.018, 0.028], *p*<0.001). The model detected significant indirect effects of impulsivity on

alcohol sipping after 12 months, with caffeinated soda intake and family risk factors mediating the

relationships (caffeinated soda intake: B=0.002, 95% CI=[0.001, 0.003], p=0.013; family risk

factor: $B=-0.002$, 95% CI=[-0.003, -0.001], $p=0.006$). These results suggest caffeinated soda

intake mediates impulsivity as the SUD risk factors toward future alcohol sipping.

DISCUSSION

In this study, we investigated whether caffeinated soda intake in childhood is associated

with a higher risk of alcohol use in the future. Using the large dataset from the ABCD study

($N=4,529$), we identified the relationships between caffeinated soda intake and the neurocognitive

risk factors for SUDs, evaluated the predictive ability of caffeinated soda on alcohol sipping after

12 months, and tested the hypothesis that caffeinated soda intake mediates neurocognitive risk

factors and alcohol sipping after 12 months. Our findings suggest that frequent consumption of

caffeinated soda in children is closely related to the neurocognitive risk factors for SUDs and can

predict future alcohol sipping, particularly by mediating impulsivity.

The machine learning approach revealed a strong association between caffeinated soda

intake and the neurocognitive risk factors for SUDs. High impulsivity scores, low WM

performance, and blunted responses to monetary reward distinguished daily caffeinated soda

drinkers from non-drinkers (**Figure 2**). In particular, we showed higher self-reported impulsivity in

the daily drinking group based on the UPPS-P and BIS/BAS scores along with altered activation

in regions implicated in impulsivity, i.e., hypoactivity in the ACC and caudate nucleus and

hyperactivity in the IFG and NAc during the SST. Consistent with our findings, reduced activities

in the ACC and caudate nucleus are commonly reported in children with attention deficit

hyperactivity disorder (ADHD) during response inhibition tasks [40–44], and activation of the

ventral striatum is greater in individuals with SUDs, implying dysregulation of impulsivity [45–47].

We detected hyperactivity in the IFG among daily soda drinkers compared with non-drinkers.

Previous studies have reported reduced activity in the IFG as neural markers of substance abuse

and ADHD [48–50]. However, increased activity in the IFG has also been reported in individuals

with tobacco use disorder [51], and increased IFG activity suggests that more preparatory effort is

required for successful inhibition [52].

We also showed behavioral WM impairments in the daily drinking group on the list sorting test, card sort test, and the 0-back accuracy in the EN-back task, accompanied by hypoactivation in the IFG and hyperactivation in the NAc, caudate nucleus, IPL, and thalamus during the EN-back task. These results are consistent with the developmental mismatch hypothesis, which states that subcortical structures of affect and reward processing develop earlier than prefrontal structures involved in the top-down regulation of behavior in children and adolescents [53,54]. Based on the previous studies that cognitive control during the EN-back task is related to WM capacity in the prefrontal cortex and simultaneous emotional regulation in the limbic system [30,55], the low WM performance in the daily drinking group may be related to increased activation in the limbic system and decreased activation in the IFG activation during the WM task.

Blunted reward anticipation was associated with caffeinated soda intake. Along with a smaller motivation to achieve the reward, activation in the ACC was lower and activation in the medial OFC was higher during the MID task in the daily drinking group than in the non-drinking group. Hyperactivation in the medial prefrontal cortex (mPFC) regions during reward anticipation is supported by previous literature suggesting increased prefrontal region activity, although these findings do not always replicate [56–60].

We showed that frequent consumption of caffeinated soda predicted alcohol sipping after 12 months using hierarchical regression (**Table 1**) and LASSO regression (**Figure 3**). Among the significant predictors in the LASSO regression, caffeinated soda intake was the second strongest predictor of future alcohol sipping, following the baseline alcohol sipping (**Figure 3B**). Other survived predictors include predictors for daily caffeinated soda intake (**Figure 2A**) while others showed the opposite direction in predicting daily caffeinated soda intake and alcohol sipping after 12 months. For example, higher SES, greater physical activity, no family history of drug use, younger age, and reduced activity in the ACC during the EN-back task and in the medial OFC during the MID task were associated with the participants who experienced alcohol sipping. Interestingly, predictors of caffeinated soda intake were similar to the risk factors for SUDs [18,61], suggesting that caffeinated soda intake during childhood and SUDs share similar neurocognitive vulnerabilities. Note that the majority of the cohort was substance-naïve [62], and

1   we evaluated alcohol sipping measures instead of actual administration of the substance. Thus, it

2   would be informative to track the predictive ability of caffeinated soda and alcohol use and

3   investigate changes in the direction of the predictors as the children get older.

4   The mediation analysis revealed that caffeinated soda consumption is a mediator

between impulsivity and future alcohol sipping (**Figure 4**). This finding suggests that caffeinated

soda itself may not directly cause children to initiate substance use, but the drinking behavior of

the beverages could indicate high impulsivity, which may be linked to a high risk of initiating

substance use in the near future. Because caffeinated soda contains two addictive substances,

sugar and caffeine, it is somewhat natural for children to prefer and seek the beverages. While

this taste preference could lead to a seeking behavior and habitual consumption, previous studies

have shown impulsivity as one of the most common traits of SUDs [63–65]. Therefore, the

behavior of frequently consuming caffeinated soda could indicate a high risk of initiating

substance use in the future, although further research is needed to disentangle this complex

relationship. Along with caffeinated soda intake, family risk factors also mediated the relationship

between impulsivity and future alcohol sipping; higher impulsivity was associated with having

more family risk factors, which led to less alcohol sipping after 12 months. Notably, some

previous studies suggested that family risk factors, such as low SES, increase the risk of future

alcohol use [66–69], while others suggested the opposite [70,71].

To our knowledge, this is the first study to investigate the direct link between caffeinated

soda intake in childhood and the risk of substance use. The results are consistent with studies on

caffeine consumption during adolescence and its association with future substance misuse

[10,13], supporting a higher risk of caffeinated soda consumption in childhood, particularly

regarding vulnerability to future substance misuse. Such information is invaluable, as caffeinated

soda is incomparably the most common medium for caffeine consumption in childhood, and the

risk of substance misuse should be detected before adolescence, the most common period of

substance use onset.

Our results have important implications for public health recommendations, as our study

provides novel insight into the neurocognitive correlates of caffeinated soda consumption in

children, which has rarely been evaluated. At the same time, a few limitations of our research and

future directions for further investigations should be discussed. First, we did not perform

functional connectivity analyses or multivoxel pattern analyses, which may have provided

additional insight into the effects of caffeinated soda intake. Second, as the 9–10-year-old

children in this study had not yet started other substances except for alcohol, such as tobacco or

marijuana, future work using the longitudinal 10-year follow-up data of the ABCD study should

examine whether frequent consumption of caffeinated soda is associated with alcohol or other

substance misuse. Third, we acknowledge that multiple variables other than caffeinated soda

intake and family risk factors may mediate the relationship between neurocognitive risk factors

and future alcohol use; thus, extensive investigation of how caffeinated soda intake interacts with

other SUD risk/protective factors is needed in the future. Lastly, the ABCD dataset included only

a small set of measures of food/drink consumption. Future studies would benefit by including

more diverse measures of foods or drinks containing caffeine or sugar; for example, by

determining if similar results are found with non-caffeinated soda or other sugary beverages.

In conclusion, our results revealed the potential risks of caffeinated soda consumption in

children by investigating the associations between caffeinated soda consumption and risk factors

for SUDs, examining the ability of caffeinated soda consumption to predict future alcohol sipping,

and revealing that caffeinated soda consumption mediates impulsivity and future alcohol sipping,

using the large ABCD dataset. While previous research on the side effects of caffeinated soda

consumption has been limited to negative physical consequences, the present results strongly

suggest that caffeinated soda drinking in children is also associated with changes in

neurocognitive function and can predict alcohol sipping after 12 months. Our study further

suggests a strong need to develop evidence-based recommendations for caffeinated soda [2], as

there is no consensus on a safe dose of caffeinated soda in children, and some children are at

higher risk of adverse events from frequent caffeinated soda intake. Further clarification on the

causal relationships and neuro-developmental evidence are needed to determine whether

caffeinated soda is a warning sign for future substance misuse and whether it induces

neurocognitive impairments in children.

ACKNOWLEDGEMENTS

We thank the families and children who participated, and continue to participate, in the ABCD

study, as well as staff at the study sites who are involved in data collection and curating the data

release. Data used in the preparation of this article were obtained from the Adolescent Brain

Cognitive Development (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive

(NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-

10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the

National Institutes of Health and additional federal partners under award numbers U01DA041048,

U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987,

U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988,

U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148,

U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is

available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a

complete listing of the study investigators can be found at

https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and

implemented the study and/or provided data but did not necessarily participate in analysis or

writing of this report. This manuscript reflects the views of the authors and may not reflect the

opinions or views of the NIH or ABCD consortium investigators. The ABCD data repository grows

and changes over time. The ABCD data used in this report came from [NIMH Data Archive Digital

Object Identifier (10.15154/1503209)]. DOIs can be found at http://dx.doi.org/10.15154/1503209.

REFERENCES

1. Temple JL. Caffeine use in children: What we know, what we have left to learn, and why we should worry. Neurosci Biobehav Rev. 2009;33(6):793–806.

2. Temple JL. Trends, Safety, and Recommendations For Caffeine Use in Children and Adolescents. J Am Acad Child Adolesc Psychiatry. 2018;58(1):36–45.

3. Ooi JY, Wolfenden L, Sutherland R, Nathan N, Oldmeadow C, Mclaughlin M, et al. A Systematic Review of the Recent Consumption Levels of Sugar-Sweetened Beverages in Children and Adolescents From the World Health Organization Regions With High Dietary–Related Burden of Disease. Asia Pac J Public Heal. 2022;34(1):11–24.

4. Sylvetsky AC, Parnarouskis L, Merkel PE, Gearhardt AN. Children's Sugar-Sweetened Beverage Consumption: Striking Parallels With Substance Use Disorder Symptoms. Frontiers Pediatrics. 2020;8:594513.

5. Nehlig A. Are we dependent upon coffee and caffeine? A review on human and animal data. Neurosci Biobehav Rev. 1999;23(4):563–76.

6. Avena NM, Rada P, Hoebel BG. Evidence for sugar addiction: Behavioral and neurochemical effects of intermittent, excessive sugar intake. Neurosci Biobehav Rev. 2008;32(1):20–39.

7. Porciúncula LO, Sallaberry C, Mioranzza S, Botton PHS, Rosemberg DB. The Janus face of caffeine. Neurochem Int. 2013;63(6):594–609.

8. Arria AM, Caldeira KM, Kasperski SJ, Vincent KB, Griffiths RR, O'Grady KE. Energy Drink Consumption and Increased Risk for Alcohol Dependence. Alcohol Clin Exp Res. 2011;35(2):365–75.

9. Miyake ER, Marmorstein NR. Energy drink consumption and later alcohol use among early adolescents. Addict Behav. 2015;43:60–5.

10. Barrense-Dias Y, Berchtold A, Akre C, Surís J. Consuming energy drinks at the age of 14 predicted legal and illegal substance use at 16. Acta Paediatr. 2016;105(11):1361–8.

11. Kristjansson AL, Kogan SM, Mann MJ, Smith ML, Juliano LM, Lilly CL, et al. Does early exposure to caffeine promote smoking and alcohol use behavior? A prospective analysis of middle school students. Addiction. 2018;113(9):1706–13.

12. Marmorstein NR. Investigating associations between caffeinated beverage consumption and later alcohol consumption among early adolescents. Addict Behav. 2018;90(Psychopharmacology 203 2009):362–8.

13. Leal WE, Jackson DB. The role of energy drink consumption in the intention to initiate marijuana use among adolescents. Addict Behav. 2019;93(Postgraduate Medicine 127 3 2015):240–5.

14. Lisdahl KM, Tapert S, Sher KJ, Gonzalez R, Nixon SJ, Ewing SWF, et al. Substance use patterns in 9-10 year olds: Baseline findings from the adolescent brain cognitive development (ABCD) study. Drug Alcohol Depen. 2021;108946.

15. Vercammen KA, Koma JW, Bleich SN. Trends in Energy Drink Consumption Among U.S. Adolescents and Adults, 2003–2016. Am J Prev Med. 2019;56(6):827–33.

16. Chambers RA, Taylor JR, Potenza MN. Developmental Neurocircuitry of Motivation in Adolescence: A Critical Period of Addiction Vulnerability. Am J Psychiat. 2003;160(6):1041–52.

17. Wagner FA, Anthony JC. From First Drug Use to Drug Dependence: Developmental Periods of Risk for Dependence upon Marijuana, Cocaine, and Alcohol. Neuropsychopharmacol. 2002;26(4):479–88.

18. Jordan CJ, Andersen SL. Sensitive periods of substance abuse: Early risk for the transition to dependence. Dev Cogn Neuros-neth. 2017;25:29–44.

19. Barry AE, King J, Sears C, Harville C, Bondoc I, Joseph K. Prioritizing Alcohol Prevention: Establishing Alcohol as the Gateway Drug and Linking Age of First Drink With Illicit Drug Use. J School Health. 2016;86(1):31–8.

20. Schneider S, Schilling L, Osenbrügge N. Determinants of soft drink consumption among children and adolescents in developed countries - a systematic review. Cent Eur J Publ Heal. 2021;29(4):290–300.

21. Lopez NV, Ayala GX, Corder K, Eisenberg CM, Zive MM, Wood C, et al. Parent Support and Parent-Mediated Behaviors Are Associated with Children's Sugary Beverage Consumption. J Acad Nutr Diet. 2012;112(4):541–7.

22. Zahid A, Davey C, Reicks M. Beverage Intake among Children: Associations with Parent and Home-Related Factors. Int J Environ Res Pu. 2017;14(8):929.

23. Gilligan C, Kypri K. Parent attitudes, family dynamics and adolescent drinking: qualitative study of the Australian parenting guidelines for adolescent alcohol use. Bmc Public Health. 2012;12(1):491–491.

24. Underwood JM, Brener N, Thornton J, Harris WA, Bryan LN, Shanklin S, et al. Youth Risk Behavior Surveillance — United States, 2019. Centers for Disease Control and Prevention. 2020 Aug 21;

25. Muchiri BW, Santos MML dos. Family management risk and protective factors for adolescent substance use in South Africa. Subst Abus Treat Prev Policy. 2018;13(1):24.

26. Wills TA, Yaeger AM. Family Factors and Adolescent Substance Use. Curr Dir Psychol Sci. 2003;12(6):222–6.

27. Solnick SJ, Hemenway D. Soft drinks, aggression and suicidal behaviour in US high school students. Int J Inj Control Sa. 2013;21(3):266–73.

28. James JE, Kristjánsson ÁL, Sigfúsdóttir ID. Adolescent substance use, sleep, and academic achievement: Evidence of harm due to caffeine. J Adolescence. 2011;34(4):665–73.

29. Suglia SF, Solnick S, Hemenway D. Soft Drinks Consumption Is Associated with Behavior Problems in 5-Year-Olds. J Pediatrics. 2013;163(5):1323–8.

30. Casey BJ, Cannonier T, Conley MI, Cohen AO, Barch DM, Heitzeg MM, et al. The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. Dev Cogn Neuros-neth. 2018;32(Biol. Psychiatry 69 2011):43–54.

31. Volkow ND, Koob G, Baler R. Biomarkers in Substance Use Disorders. Acs Chem Neurosci. 2015;6(4):522–5.

32. Watts AL, Wood PK, Jackson KM, Lisdahl KM, Heitzeg MM, Gonzalez R, et al. Incipient alcohol use in childhood: Early alcohol sipping and its relations with psychopathology and personality. Dev Psychopathol. 2021;33(4):1338–50.

33. Jackson KM, Barnett NP, Colby SM, Rogers ML. The Prospective Association Between Sipping Alcohol by the Sixth Grade and Later Substance Use. J Stud Alcohol Drugs. 2015;76(2):212–21.

34. Donovan JE, Molina BSG. Childhood Risk Factors for Early-Onset Drinking*. J Stud Alcohol Drugs. 2011;72(5):741–51.

35. Kirby T, Barry AE. Alcohol as a gateway drug: a study of US 12th graders. J Sch Heal. 2012;82(8):371–9.

36. Mullee A, Romaguera D, Pearson-Stuttard J, Viallon V, Stepien M, Freisling H, et al. Association Between Soft Drink Consumption and Mortality in 10 European Countries. Jama Intern Med. 2019;179(11).

37. Tibshirani R. Regression Shrinkage and Selection Via the Lasso. J Royal Statistical Soc Ser B Methodol. 1996;58(1):267–88.

38. Carver CS, White TL. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. J Pers Soc Psychol. 1994;67(2):319.

39. Barch DM, Belden AC, Tillman R, Whalen D, Luby JL. Early Childhood Adverse Experiences, Inferior Frontal Gyrus Connectivity, and the Trajectory of Externalizing Psychopathology. J Am Acad Child Adolesc Psychiatry. 2018;57(3):183–90.

40. Lei D, Du M, Wu M, Chen T, Huang X, Du X, et al. Functional MRI Reveals Different Response Inhibition Between Adults and Children With ADHD. Neuropsychology. 2015;29(6):874–81.

41. Cortese S, Kelly C, Chabernaud C, Proal E, Martino AD, Milham MP, et al. Toward Systems Neuroscience of ADHD: A Meta-Analysis of 55 fMRI Studies. Am J Psychiat. 2012;169(10):1038–55.

42. Hart H, Radua J, Nakao T, Mataix-Cols D, Rubia K. Meta-analysis of Functional Magnetic Resonance Imaging Studies of Inhibition and Attention in Attention-deficit/Hyperactivity Disorder: Exploring Task-Specific, Stimulant Medication, and Age Effects. Jama Psychiat. 2013;70(2):185–98.

43. Zarka D, Cebolla AM, Cevallos C, Palmero-Soler E, Dan B, Cheron G. Caudate and cerebellar involvement in altered P2 and P3 components of GO/NoGO evoked potentials in children with attention-deficit/hyperactivity disorder. Eur J Neurosci. 2021;53(10):3447–62.

44. Mahmood OM, Goldenberg D, Thayer R, Migliorini R, Simmons AN, Tapert SF. Adolescents' fMRI activation to a response inhibition task predicts future substance use. Addict Behav. 2013;38(1):1435–41.

45. Bjork JM, Smith AR, Hommer DW. Striatal sensitivity to reward deliveries and omissions in substance dependent patients. Neuroimage. 2008;42(4):1609–21.

46. Volkow ND, Fowler JS, Wang GJ, Swanson JM, Telang F. Dopamine in Drug Abuse and Addiction: Results of Imaging Studies and Treatment Implications. Arch Neurol-chicago. 2007;64(11):1575–9.

47. Heitzeg MM, Nigg JT, Yau WYW, Zucker RA, Zubieta JK. Striatal Dysfunction Marks Preexisting Risk and Medial Prefrontal Dysfunction Is Related to Problem Drinking in Children of Alcoholics. Biol Psychiat. 2010;68(3):287–95.

48. Schmidt A, Walter M, Gerber H, Schmid O, Smieskova R, Bendfeldt K, et al. Inferior Frontal Cortex Modulation with an Acute Dose of Heroin During Cognitive Control. Neuropsychopharmacol. 2013;38(11):2231–9.

49. Fu L ping, Bi G hua, Zou Z tong, Wang Y, Ye E mao, Ma L, et al. Impaired response inhibition function in abstinent heroin dependents: An fMRI study. Neurosci Lett. 2008;438(3):322–6.

50. Morein-Zamir S, Dodds C, Hartevelt TJ, Schwarzkopf W, Sahakian B, Müller U, et al. Hypoactivation in right inferior frontal cortex is specifically associated with motor response inhibition in adult ADHD. Hum Brain Mapp. 2014;35(10):5141–52.

51. Kalhan S, Chen LPE, Garrido MI, Hester R. People with tobacco use disorder exhibit more prefrontal activity during preparatory control but reduced anterior cingulate activity during reactive control. Addict Biol. 2022;27(2):e13159.

52. DeVito EE, Meda SA, Jiantonio R, Potenza MN, Krystal JH, Pearlson GD. Neural Correlates of Impulsivity in Healthy Males and Females with Family Histories of Alcoholism. Neuropsychopharmacol. 2013;38(10):1854–63.

53. Somerville LH, Hare T, Casey BJ. Frontostriatal maturation predicts cognitive control failure to appetitive cues in adolescents. J Cognitive Neurosci. 2010;23(9):2123–34.

54. Mills KL, Goddings AL, Clasen LS, Giedd JN, Blakemore SJ. The Developmental Mismatch in Structural Brain Maturation during Adolescence. Dev Neurosci-basel. 2014;36(3–4):147–60.

55. Gee DG, Humphreys KL, Flannery J, Goff B, Telzer EH, Shapiro M, et al. A developmental shift from positive to negative connectivity in human amygdala-prefrontal circuitry. J Neurosci Official J Soc Neurosci. 2013;33(10):4584–93.

56. Beck A, Schlagenhauf F, Wüstenberg T, Hein J, Kienast T, Kahnt T, et al. Ventral Striatal Activation During Reward Anticipation Correlates with Impulsivity in Alcoholics. Biol Psychiat. 2009;66(8):734–42.

57. Balodis IM, Kober H, Worhunsky PD, Stevens MC, Pearlson GD, Potenza MN. Diminished
Frontostriatal Activity During Processing of Monetary Rewards and Losses in Pathological
Gambling. Biol Psychiat. 2012;71(8):749–57.

58. Swartz JR, Weissman DG, Ferrer E, Beard SJ, Fassbender C, Robins RW, et al. Reward-
Related Brain Activity Prospectively Predicts Increases in Alcohol Use in Adolescents. J Am Acad
Child Adolesc Psychiatry. 2019;59(3):391–400.

59. Elsayed NM, Kim MJ, Fields KM, Olvera RL, Hariri AR, Williamson DE. Trajectories of Alcohol
Initiation and Use During Adolescence: The Role of Stress and Amygdala Reactivity. J Am Acad
Child Adolesc Psychiatry. 2018;57(Environ Psychol Nonverbal Behav 1 1 1976):550–60.

60. Luijten M, Schellekens AF, Kühn S, Machielse MWJ, Sescousse G. Disruption of Reward
Processing in Addiction : An Image-Based Meta-analysis of Functional Magnetic Resonance
Imaging Studies. Jama Psychiat. 2017;74(4):387.

61. Kilpatrick DG, Acierno R, Saunders B, Resnick HS, Best CL, Schnurr PP. Risk Factors for
Adolescent Substance Abuse and Dependence: Data From a National Sample. J Consult Clin
Psych. 2000;68(1):19–30.

62. Lisdahl KM, Sher KJ, Conway KP, Gonzalez R, Ewing SWF, Nixon SJ, et al. Adolescent brain
cognitive development (ABCD) study: Overview of substance use assessment methods. Dev
Cogn Neuros-neth. 2018;32:80–96.

63. Cyders MA, Flory K, Rainer S, Smith GT. The role of personality dispositions to risky behavior
in predicting first-year college drinking. Addiction. 2009;104(2):193–202.

64. Whiteside SP, Lynam DR. Understanding the Role of Impulsivity and Externalizing
Psychopathology in Alcohol Abuse: Application of the UPPS Impulsive Behavior Scale. Exp Clin
Psychopharm. 2003;11(3):210–7.

65. Magid V, Colder CR. The UPPS Impulsive Behavior Scale: Factor structure and associations
with college drinking. Pers Indiv Differ. 2007;43(7):1927–37.

66. Calling S, Ohlsson H, Sundquist J, Sundquist K, Kendler KS. Socioeconomic status and
alcohol use disorders across the lifespan: A co-relative control study. Plos One.
2019;14(10):e0224127.

67. Carroll HA, Heleniak C, Witkiewitz K, Lewis M, Eakins D, Staples J, et al. Effects of parental
monitoring on alcohol use in the US and Sweden: A brief report. Addict Behav. 2016;63:89–92.

68. Collins SE. Associations Between Socioeconomic Factors and Alcohol Outcomes. Alcohol
Res Curr Rev. 2016;38(1):83–94.

69. Goodwin DW. Alcoholism and Heredity: A Review and Hypothesis. Arch Gen Psychiat.
1979;36(1):57–61.

70. Liu Y, Lintonen T, Tynjälä J, Villberg J, Välimaa R, Ojala K, et al. Socioeconomic differences in
the use of alcohol and drunkenness in adolescents: Trends in the Health Behaviour in School-
aged Children study in Finland 1990–2014. Scand J Public Healt. 2016;46(1):102–11.

71. Kendler KS, Gardner CO, Hickman M, Heron J, Macleod J, Lewis G, et al. Socioeconomic
Status and Alcohol-Related Behaviors in Mid- to Late Adolescence in the Avon Longitudinal
Study of Parents and Children. J Stud Alcohol Drugs. 2014;75(4):541–5.

72. Destrieux C, Fischl B, Dale A, Halgren E. Automatic parcellation of human cortical gyri and
sulci using standard anatomical nomenclature. Neuroimage. 2010;53(1):1–15.

Figure 1. An overview of the study design

A. Diagram of the research aims. First, the association between the risk factors for substance use

disorders (SUDs) and caffeinated soda intake was examined. Then, we assessed whether

caffeinated soda intake predicts alcohol sipping after 12 months, and whether caffeinated soda

intake works as a mediator between the risk factors and future alcohol sipping. **B.** The variables

used to measure the three main neurocognitive risk factors: working memory, impulsivity, and

reward processing. Working memory was measured from the performance of list sorting test, card

sort test, and 2-back in the emotional N-Back Task (EN-back) performed inside a functional

magnetic resonance imaging (fMRI) scanner. The fMRI data from the EN-back task were

analyzed based on the contrast of 2-back versus 0-back, using regions in the frontoparietal and

fronto-thalamic network as the regions of interest (ROIs). Impulsivity was measured by Urgency-

Premeditation-Perseverance-Sensation Seeking-Positive Urgency (UPPS-P), and Behavioral

Inhibition System/ Behavioral Activation System (BIS/BAS), and the stop signal reaction time

(SSRT) during the stop signal task (SST) performed inside the fMRI scanner. The fMRI data from

the SST were analyzed based on the contrast of correct stop versus correct go, using regions in

the lateral prefrontal cortex, anterior cingulate cortex (ACC), and striatum as the ROIs. Reward

processing was measured by cash choice task, and also by comparing the success rate of

reward versus neutral conditions during the monetary incentive delay (MID) task performed inside

the fMRI scanner. The fMRI data from the MID task were analyzed based on the contrast of

reward anticipation versus neutral anticipation, using regions in the ventral striatum and medial

frontal cortex as the ROIs. The ROIs were selected based on the Destrieux atlas [72].

Figure 2. Results of LASSO regression predicting daily caffeinated soda intake (daily drinking

group vs. non-drinking group)

A. Estimates of coefficients of the survived variables (x axis: coefficient estimates; y axis: predictor).

The site variables were excluded for clarify (see **Figure S1** for the site variables). **B.** Distribution of

the area under the curve (AUC) values (left) and a representative receiver-operation characteristic

(ROC) curve (right) for the training and test datasets. **C.** Regions of interest (ROIs) identified as

having significant estimates of coefficients during the emotional N-Back Task (EN-back), stop signal

task (SST), and monetary incentive delay (MID) task.

Abbreviations. LASSO, least absolute shrinkage and selection operator; BMI, body mass index;

BIS, Behavioral Inhibition System; BAS, Behavioral Activation System; UPPS-P, Urgency-

Premeditation-Perseverance-Sensation Seeking-Positive Urgency; SST, Stop Signal Task; SSRT,

Stop Signal Reaction Time; MID, Monetary Incentive Delay Task; NAc, Nucleus Accumbens; IPL,

Inferior Parietal Lobule; IFG, Inferior Frontal Gyrus; ACC, Anterior Cingulate Cortex; OFC,

Orbitofrontal Cortex.

Figure 3. Results of LASSO regression predicting alcohol sipping after 12 months

A. Rate of alcohol sipping after 12 months in each group. **B.** Estimates of coefficients of the survived

variables (x axis: coefficient estimates; y axis: predictor). Site variables were excluded for a clarity

(see **Figure S2** for the site variables). **C.** Distribution of the area under the curve (AUC) values (left)

and a representative receiver-operation characteristic (ROC) curve (right) for the training and test

datasets.

*: $p<0.05$, **: $p<0.01$; ***: $p<0.001$

Figure 4. Caffeinated soda consumption at baseline as a mediator of the relationship between impulsivity and alcohol sipping after 12 months

Along with family risk factors, caffeinated soda intake was a significant mediator between

impulsivity and future alcohol sipping. Significant paths are flagged with asterisks (*: *p*<0.05, **:

p<0.01; ***: *p*<0.001). Indirect effects of caffeinated soda intake and family risk factors on the

relationship between impulsivity and alcohol sipping after 12 months were significant (caffeinated

soda intake: B=0.002, 95% CI=[0.001, 0.003], p=0.013; family risk factors: B=−0.002, 95%

CI=[−0.003, −0.001], p=0.006).

Table 1. Results of hierarchical regression analysis predicting alcohol sipping after 12 months

Variables	Model 1	Model 2	Model 3
Sex (Male)	-0.020 (-0.031 - -0.010)***	-0.015 (-0.025 - -0.005)**	-0.013 (-0.023 - -0.003)**
Race (Caucasian)	0.032 (0.011 - 0.053)**	0.022 (0.002 - 0.042)*	0.022 (0.002 - 0.042)*
Race (African American)	0.001 (-0.024 - 0.026)	0.003 (-0.021 - 0.027)	0.003 (-0.021 - 0.027)
Family income	0.007 (0.000 - 0.015)*	0.005 (-0.002 - 0.012)	0.005 (-0.002 - 0.012)
Parental education	0.009 (0.002 - 0.015)*	0.007 (0.001 - 0.014)*	0.009 (0.002 - 0.015)*
Parental monitoring	-0.003 (-0.009 - 0.002)	-0.002 (-0.007 - 0.003)	-0.001 (-0.006 - 0.004)
Family history of drug	-0.003 (-0.015 - 0.009)	-0.005 (-0.017 - 0.007)	-0.005 (-0.017 - 0.006)
Family history of alcohol	-0.008 (-0.019 - 0.003)	-0.010 (-0.020 - 0.001)	-0.010 (-0.021 - 0.001)
BMI	0.007 (0.002 - 0.012)**	0.006 (0.001 - 0.011)*	0.006 (0.001 - 0.011)*
Physical activity	0.007 (0.001 - 0.012)*	0.005 (0.000 - 0.010)	0.005 (0.000 - 0.011)
Lack of sleep	0.000 (-0.005 - 0.006)	0.002 (-0.004 - 0.007)	0.001 (-0.004 - 0.007)
Externalizing behaviors	0.009 (0.003 - 0.015)**	0.007 (0.001 - 0.013)*	0.006 (0.000 - 0.013)*
Internalizing behaviors	-0.003 (-0.009 - 0.004)	-0.003 (-0.009 - 0.003)	-0.003 (-0.009 - 0.003)
Alcohol sipping (Baseline)		0.174 (0.161 - 0.187)***	0.172 (0.159 - 0.185)***
Monthly caffeinated soda intake			0.021 (0.010 - 0.032)***
Weekly caffeinated soda intake			0.024 (0.008 - 0.041)**
Daily caffeinated soda intake			0.036 (0.010 - 0.062)**

Note. The values are standardized regression coefficients (95% confidence intervals) from each step in the hierarchical analyses predicting alcohol sipping after 12 months. Significant variables predicting alcohol sipping are flagged with asterisks (*: $p<0.05$, **: $p<0.01$; ***: $p<0.001$). Higher consumption of caffeinated soda intake (Model 3) predicted a higher tendency for alcohol sipping after 12 months even after controlling for the confounding variables (13 variables from the top and 21 data collection sites in Model 1) and alcohol sipping collected at baseline (Model 2).

Abbreviations. BMI, Body Mass Index.