Title: Daily caffeinated soda intake in children is associated with neurocognitive vulnerabilities of

substance misuse and predicts higher risk of alcohol sipping in 12 months

Abbreviated title: Daily soda intake in children and alcohol sipping

Authors:

Mina Kwon[1], Hyeonjin Kim[1], Jaeyeong Yang[1], Yoseph Lee[1], Jihyun K. Hur[2], Tae-Ho Lee[3], and

Woo-Young Ahn[1,4]

[1]Department of Psychology, Seoul National University, Seoul, Korea 08826

[2]Department of Psychology, Yale University, New Haven, CT 06511

[3]Department of Psychology, Virginia Tech, Blacksburg, VA 24060

[4]Department of Brain and Cognitive Sciences, Seoul National University, Seoul, Korea 08826

Corresponding author:

Woo-Young Ahn, Ph.D.

Department of Psychology

Seoul National University

Seoul, Korea 08826

Tel: +82-2-880-2538, Fax: +82-2-877-6428. E-mail: wahn55@snu.ac.kr

Number of pages: 43

Number of Figures & Tables: 4 / 3

Number of words in the abstract: 247

Number of words in the introduction: 624

Number of words in the discussion: 1,098

Conflict of Interest: The authors declare no competing financial interest.

Abstract

Caffeinated soda contains two addictive substances, sugar and caffeine, and is the most preferred means of caffeine consumption among children. However, it remains unexplored if habitual caffeinated soda intake in childhood is associated with a higher risk of alcohol misuse in the future. Here, we investigated the neurocognitive correlates of daily consumption of caffeinated soda and examined whether daily caffeinated soda intake is associated with a higher risk of alcohol initiation in children. By applying machine learning and hierarchical regression to a large dataset from the Adolescent Brain Cognitive Development (ABCD) Study, we investigated the relationship between daily caffeinated soda intake and the well-known risk factors of substance misuse, including impaired working memory, high impulsivity, and altered reward processing. We next examined whether daily caffeinated soda intake could predict greater alcohol sipping in 12 months. The results showed that daily consumption of caffeinated soda was associated with impaired working memory, higher impulsivity, and altered reward processing, with increased amygdala activation during the emotional working memory task and hypoactivity in the nucleus accumbens and posterior cingulate cortex during reward processing tasks. Daily drinkers reported higher rates of alcohol sipping compared to non-drinkers after 12 months, and daily intake of caffeinated soda mediated the relationship between the neurocognitive risk factors and future alcohol sipping. These results suggest that daily caffeinated soda intake in childhood (1) is associated with impaired neurocognitive functioning, and (2) may work as a predictor for alcohol initiation, which has significant implications for public health recommendations.

Significance Statement

Is daily caffeinated soda intake associated with higher risks of substance misuse? If so, which specific risk factors of substance misuse are involved? Moreover, can daily caffeinated soda intake predict future alcohol initiation? These are the questions that many parents are asking but surprisingly there is no clear guideline, and its potential vulnerabilities of substance misuse remain largely unknown. Here, by applying machine learning and hierarchical regression

1   approaches, we demonstrate that daily caffeinated soda intake is associated with neurocognitive

2   risk factors of substance misuse including impaired working memory, higher impulsivity, and

3   altered reward processing. We also present that daily caffeinated soda consumption could predict

4   higher alcohol sipping rates in 12 months. These results have significant implications for public

5   health recommendations.

Introduction

Caffeinated soda combines two addictive substances, sugar and caffeine, both of which

affect neurocognitive functions and cause physical side effects (Nehlig, 1999; Avena et al., 2008;

Sylvetsky et al., 2020). Not surprisingly, the consequences of excessive consumption of both

sugar and caffeine have been well documented (Porciúncula et al., 2013), including a strong

association between caffeinated beverages consumption in adolescents and future substance

use (Arria *et al.*, 2011; Miyake and Marmorstein, 2015; Barrense-Dias *et al.*, 2016; Kristjansson *et*

al., 2018; Marmorstein, 2018; Leal and Jackson, 2019). In prospective studies tracking the effects

of substance use, the percentage of regular energy drink users who became alcohol or marijuana

users after 1-2 years was approximately five times higher than that of non-users (Miyake and

Marmorstein, 2015; Marmorstein, 2018; Leal and Jackson, 2019). Others have also shown that

coffee or energy drink consumption in developing children could significantly predict future

substance use such as tobacco and alcohol (Arria et al., 2011; Barrense-Dias et al., 2016;

Marmorstein, 2018).

Existing literature examining the associations between caffeine consumption and later

substance use focused more on adolescents who drink multiple caffeinated beverages on a daily

basis, partially due to the fact that consumption level of coffee and energy drink is extremely low

in children, 0.4% for coffee and less than 0.1% for energy drinks in 9-10 years old (Lisdahl et al.,

2021). However, caffeinated soda is the most preferred and accessible mode of caffeine intake in

children (Temple, 2009; Vercammen et al., 2019; Sylvetsky et al., 2020). Moreover, it is well

known that adolescence is the most common period of substance use initiation, and an earlier

onset of substance use predicts greater addiction severity (Wagner and Anthony, 2002;

Chambers et al., 2003; Barry et al., 2016; Jordan and Andersen, 2017). Therefore, there is a

critical need to investigate the link between caffeinated soda intake during preadolescence

childhood and the risk of substance use disorders (SUDs) in the future, including vulnerabilities

such as associated neurocognitive deficits or early exposure to substances of abuse. Even

though few studies have directly examined the effects of frequent caffeinated soda intake in

adolescents and children, they either targeted only behavioral aspects using surveys or self-

reports without obtaining neural data, or did not target children but only adolescents (James et al.,

2011; Solnick and Hemenway, 2013; Suglia et al., 2013; Miyake and Marmorstein, 2015).

Here, we addressed the unanswered question of whether daily intake of caffeinated soda

in children could indicate a higher risk of later substance misuse, using the Adolescent Brain

Cognitive Development (ABCD) Study. First, we examined the relationship between daily

caffeinated soda intake and three well-known neurocognitive risk factors of SUDs in children:

impaired working memory (WM), high impulsivity, and altered reward processing (Casey et al.,

2018). From the baseline data of the ABCD study, we used psychological, behavioral, and

neurological indices related to each risk factor (**Figure 1A**), and employed both univariate

(hierarchical regression) and multivariate (machine learning) approaches to rigorously investigate

the associations between daily caffeinated soda intake and the risk factors of SUDs. **Figure 1B**

shows the two different analytical methods that have different predictive directionalities. We next

examined whether daily caffeinated soda intake could predict future alcohol sipping, in that (1)

alcohol sipping has been reported to predict future alcohol abuse (Donovan and Molina, 2011;

Jackson et al., 2015; Watts et al., 2021), and (2) alcohol sipping is the most common gateway

behavior toward other substances of abuse (Kirby and Barry, 2012; Barry et al., 2016). Lastly, we

identified the best model explaining how the interaction of daily caffeinated soda intake and the

risk factors of SUDs leads to later alcohol sipping behavior. Together, we tried to provide insight

into possible underlying risks of daily caffeinated soda intake in childhood.

Method

Participants

The ABCD study has collected data from 11,875 children aged 9- and 10-years-old. The

participants were recruited through school systems from 21 different sites in the US. More

information about the recruitment and study design is available in Garavan et al. (2018). Further

details of the demographic, physical, and mental health assessments are described in Barch et

al. (2018).

To include as many samples as possible, we used different sample sizes for each

analysis. For the first analysis examining the relationship between caffeinated soda intake and

the risk factors of SUDs, 3,966 participants at the baseline data collection (Data Release 2.0)

without any missing data in the following curated datasets were included: task-based functional

magnetic resonance imaging (fMRI) data for three tasks (Emotional N-Back (EN-Back) Task

(Cohen et al., 2015), Stop Signal Task (SST) (Logan et al., 1984), and Monetary Incentive Delay

(MID) Task (Knutson et al., 2000)), behavioral data obtained from cognitive tasks (the List Sorting

Test, Card Sort Test, and Cash Choice Task) and from psychological surveys, and demographic

data. The task-based fMRI data were collected from three different MRI scanners (Siemens,

General Electric, and Philips); data from the Philips scanner were incorrectly processed in the

curated dataset and were therefore excluded from the analysis (N=527). For the second analysis

testing the link between caffeinated soda intake and future alcohol sipping, we included 8,953

participants without any missing data in soda intake measure and control variables (i.e.,

psychological surveys and demographic information) in the baseline data and in alcohol sipping

measure in the year 1 data (Data Release 3.0). For the third analysis testing the link between

daily caffeinated soda intake, alcohol sipping, and the risk factors of SUDs, 3,827 participants

were analyzed based on the same criteria as the first analysis, in addition to having a complete

set of data in the year 1 data.

Measures

Caffeinated soda intake

In every analysis sample, the participants were divided into four groups based on their

caffeinated soda consumption over the last 6 months, which was assessed using the participants'

self-report in response to the question, "Typically, how many drinks of the following beverages

have you had per week in the past 6 months? – soda with caffeine (Mountain Dew, Jolt, Coke,

Pepsi, Dr. Pepper, Barq's root beer)". The non-drinking group reported 0 cans per week, the

monthly-drinking group reported <1 can per week, the weekly-drinking group reported 1–7 cans

per week, and the daily-drinking group reported ≥7 cans per week. These criteria for group

allocation were defined by referring to the population-based cohort study of Mullee et al. (2019).

For the main analyses, we focused on the daily-drinking group and non-drinking group to ensure

adequate separation between habitual drinkers (daily-drinking group) and non-drinkers (non-

drinking group).

Risk factors of SUDs

As shown in **Figure 1**, we focused on the three well-known psychological risk factors for

SUDs (WM, impulsivity, and reward processing), measured by self-report surveys, behavioral

tasks, and fMRI, as described below.

To obtain behavioral measures of WM, we used the List Sorting Working Memory Test

(List Sorting Test) and Dimensional Change Card Sort Test (Card Sort Test) from the NIH

Toolbox (Tulsky et al., 2013), and the EN-Back task. The List Sorting Test requires participants to

sort and sequence visual and auditory stimuli. For the Card Sort Test, participants are asked to

sort cards according to either color (when the target card has a contour) or shape (when the card

without a contour is shown). The EN-Back task (Cohen et al., 2015), which measures WM and

emotion regulation, was conducted inside the MRI scanner. The task consists of two different

blocks: the 0-back block requires participants to respond when the current stimulus is the same

as a target, and the 2-back block requires participants to respond when the current stimulus is the

same as the one shown in two trials before. Due to the difference in WM load between the two

blocks, accuracy during the 2-back block is commonly used as a behavioral measure of WM, and

the 2-back block vs. 0-back block contrast is commonly used to examine the neural correlates of

WM (Casey et al., 2018). Unlike the traditional N-Back task, in which the stimuli are numbers, the

EN-Back task displays emotional (happy, fearful, and neutral) face stimuli and place (non-

emotional) stimuli. Therefore, the task also allows us to test the effect of emotional processing in

WM. The regions of interest (ROIs) of the EN-Back task include the amygdala, a region that is

well-known for emotional processing and has also been linked with working memory function

(Banks et al., 2007), as well as the hippocampus and dorsolateral prefrontal cortex (DLPFC),

which are critical for WM (Barbey et al., 2013). See Casey et al. (2018) for more details about the

task.

Impulsivity, particularly trait impulsivity, was measured using the short form Urgency-

Premeditation-Perseverance-Sensation Seeking-Positive Urgency (UPPS-P) impulsive behavior

scale for children (the 20-item short version for youths, Barch et al., 2018) and parental report of

ABCD Youth Behavioral Inhibition System/ Behavioral Activation System (BIS/BAS). The BAS is

related to goal-directed efforts, such as motor activation in response to impending rewards, while

the BIS is engaged when inhibition toward a goal occurs, such as the avoidance of punishment

(Carver and White, 1994). The existing literature suggests that Behavioral Inhibition is associated

with depression and anxiety, while Behavioral Activation is associated with impulsive behaviors,

compulsive behaviors, substance misuse, and aggression (Newman et al., 2005).

To identify the neural correlates of impulsivity, participants completed the SST (Logan et

al., 1984) inside the scanner. The SST measures impulsivity related to impulse control or

response inhibition when performing an action. The SST requires participants to withhold motor

responses to a "Go" stimulus when a "Stop" signal is unpredictably presented after a short delay.

The delay between the "Go" stimulus presentation and the "Stop" signal presentation is termed

the stop signal delay (SSD). The SSD was initially set to 50ms. Following a successful inhibition

or an unsuccessful inhibition, the SSD was increased or decreased by 50ms to make the task

more difficult or easier, respectively. Each run contained 180 trials, and each trial began with the

presentation of the "Go" cue. Participants were instructed to respond to the cue as quickly and

accurately as possible. Thirty trials from a single run (16.67%) presented a "Stop" signal, which

was an upward arrow displayed for 300 ms. Stop signal reaction time (SSRT) was used as a

behavioral measure of impulsivity, calculated by subtracting the mean SSD from the mean "Go"

response time (Logan et al., 1984; Schachar and Logan, 1990); a shorter SSRT means better

response inhibition. For the fMRI analysis, a "correct stop vs. correct go" contrast was used, and

the caudal anterior cingulate cortex (ACC) and the inferior frontal gyrus (IFG) were selected as

ROIs; both areas have been associated with impulsivity and impulse control (Stadler et al., 2007;

Jacobson et al., 2011). More information about the SST is available in Casey et al. (2018).

Reward processing is closely linked to impulsivity, in that impulsive people often show

reward-seeking behavior (Zuckerman, 2001). Previous literature suggests that people with SUDs

show hyperresponsiveness toward drug-related cues (Robinson and Berridge, 2008; Bechara et

al., 2019) while showing blunted responsiveness toward generalized reward (e.g. monetary

reward) (Balodis and Potenza, 2015; Luijten et al., 2017). We used the Cash Choice Task

(Luciana et al., 2018), which measures delayed gratification by assessing impulsivity in receiving

a monetary reward when asked the following single-item question: "Let's pretend a kind person

wanted to give you some money. Would you rather have $75 in three days or $115 in 3 months?"

The former choice reflects poor delayed gratification, with a higher delay discounting rate, while

the latter choice implies a smaller delay discounting rate. Responses were scored as 1 ($75 in

three days) or 2 ($115 in 3 months).

The MID task (Knutson et al., 2000) is widely used to measure the neural correlates of

reward processing, such as the anticipation of monetary rewards and losses. Each trial of the

MID task begins with an incentive cue of five possible trial types (Win $20, Win $5, Lose $0.2,

Lose $5, and $0 – no money at stake), followed by a delay (1500 ms~4000 ms) and a

presentation of a target during which the participant chooses to either win money or avoid losing

money, and ends with feedback on whether the participant won or lost money. Each participant

performed 40 reward anticipation, 40 loss anticipation, and 20 no money anticipation trials,

resulting in a total of 100 trials. The differences in accuracy between reward and neutral trials,

and loss and neutral trials indicate motivation to win reward or avoid loss. The contrast of

"Reward vs. Neutral" at the cue onset, which reflects reward anticipation, was used for fMRI

analysis, with the ROIs as the nucleus accumbens (NAc) and the posterior cingulate cortex

(PCC), which are known to play a key role in reward processing, particularly in reward

anticipation (Day and Carelli, 2007). See Casey et al. (2018) for more details about the MID task.

Alcohol sipping

Alcohol sipping was measured with a self-report from youth using iSay Sip Inventory,

which is being collected once a year and asked only to children who had heard of alcohol (see

Lisdahl *et al.* (2021) for more details). From the year 1 data (Data Release 3.0), we used a binary

response to a single question asking if they have sipped alcohol outside of a religious ceremony.

We considered those who have not heard of it as having yet to sip alcohol (i.e., no alcohol

sipping).

Confounding variables

We included the following variables as control variables in data analysis: Family history of

substance use, parental monitoring, sleep deprivation, and externalizing/internalizing behaviors.

Family history of substance use was measured using a parental yes/no response to the following

question: "Has any blood relative of your child ever had any problems due to drugs, such as:

Marital separation or divorce; Fired from work; Arrests or DUI (Driving Under Influence); Adverse

health effects; In a drug treatment program; Suspended or expelled from school two or more

times; Isolated oneself from the family, caused arguments, or often intoxicated." To measure the

family history of alcohol use, the same question was asked, except that "drug" was replaced with

"alcohol".

Parental monitoring was measured using the ABCD Parental Monitoring Survey, in which

children answered five questions ("How often do your parents/guardians know where you are?",

"How often do your parents know who you are with, or when you are not at school and away from

home?", "If you are at home when your parents or guardians are not, how often do you know how

to get in touch with them?", "How often do you talk to your parent or guardian about your plans for

the coming day, such as your plans about what will happen at school or what you are going to do

with friends?", "In an average week, how many times do you eat dinner with your

parents/guardians?"). Responses ranged from 1 (Never) to 5 (Always or Almost Always), and the

mean of all five responses was used for the analysis. The SES measures the total combined

family income and parents' highest education level. Parents' education was calculated as the

mean of both parents, or of the responding parent only in the case of single parents.

The sleep deprivation was measured using the ABCD Parent Sleep Disturbance Scale for

Children, for which parents responded to the question, "How many hours of sleep does your child

get on most nights?" (1=9–11 hours, 2=8–9 hours, 3=7–8 hours, 4=5–7 hours, 5=less than 5

hours). Physical activity was measured using the ABCD Parent Sports and Activities Involvement

Questionnaire, which asks whether the child has ever continuously participated in specific sports

and activities, such as football, field hockey, climbing, or basketball, for four months or more. The

number of "yes" responses to 29 questions was summed for the analysis. BMI was calculated

according to the children's body weight and height.

Externalizing and internalizing behaviors were measured using the parental report of the

Child Behavior Checklist (Achenbach, 2009), which comprises 113 items about child behavior

over the past six months. Internalizing and externalizing scores were measured based on the

following syndrome scores: anxious/depressed, withdrawn/depressed, somatic complains, social

problems, though problems, rule-breaking behavior, and aggressive behavior (Thompson et al.,

2018).

MRI Protocol

Imaging-derived phenotype scores provided by the ABCD study were used. Data

acquisition protocol was matched for the three 3T scanners used (Siemens Prisma, General

Electric 750, and Philips). We analyzed beta weights within each ROI for the 2-back vs. 0-back

contrast in the EN-Back task, the correct stop vs. correct go contrast in the SST, and the reward

vs. neutral anticipation contrast in the MID task. More details about fMRI imaging protocols such

as scanning parameters in the ABCD study are reported by Casey et al. (2018).

Analysis

Daily caffeinated soda intake and risk factors of SUDs

To identify a multivariate profile of risk factors that can differentiate the daily-drinking

group (*N*=1,640) from the non-drinking group (*N*=141), we performed the least absolute shrinkage

and selection operator (LASSO) regression analysis (Tibshirani, 1996), a machine learning

algorithm, with all possible variables from the baseline data. LASSO regression offers automatic

variable selection through a penalized least squares method, which increases the interpretability

of the findings especially when the multiple variables share a common characteristic to some

extent. For example, each task measure (e.g., the EN-Back task, List Sorting Test, and Card Sort

Test) within cognitive risk factors are often moderately correlated with one another because they

share common cognitive processing (e.g., WM). In this case, LASSO regression shrinks

coefficients of variables with less explanatory power to zero and selects only variables with the

highest power for further analyses (Tibshirani, 1996, 2011).

In the main LASSO-based prediction model, we used all the risk factors and the

confounding variables as an input (i.e., predictor) to classify each individual into either daily

drinking group (codes as "1") or non-drinking group (coded as "0") (**Figure 1B**), given that the

primary interest was to identify the potential risk of daily soda-drinking behavior. More specifically,

forty candidate predictors included every measure for the three cognitive factors as well as other

control variables (y-axis of **Figure 2A**). We used an R package called "easyml" (v0.1.1), a toolkit

to easily build and evaluate machine learning models using the glmnet package (Friedman et al.,

2010) for LASSO regression and several other machine learning algorithms

(https://github.com/CCS-Lab/easyml; Ahn et al., 2017). The mixing parameter alpha was set to 1

to conduct the LASSO (1 for LASSO, 0 for ridge, and a value between 0 and 1 for elastic net). We

trained the model using k-fold cross-validation (k=10) within the training set, and this generated

an estimated coefficient 100 times for a particular train-test split, whereby we calculated the mean

and standard deviation of the estimated coefficients. By replicating predictions many times for a

particular train-test split, averaging the predictions across 10 iterations and using 1000 divisions

of train-test splits, we could avoid intrinsic random errors (see Ahn et al., 2014, 2017 & Ahn and

Vassileva, 2016 for more details). Given that previous results have revealed that the elastic net

often performs better for highly correlated predictors than for LASSO regression (Zou and Hastie,

2005), we tested the elastic net by setting alpha to a value between 0 and 1. However, there were

no significant differences between the LASSO and elastic net results, so we only report the

LASSO results.

In addition to the multivariate approach, we also performed the univariate level of

regression analysis to confirm the link between each risk factor and daily caffeinated soda intake.

To this end, we conducted a hierarchical regression analysis using the "lm" function in R and

tested whether daily caffeinated soda intake was a significant predictor of each of the three

cognitive factors (**Figure 1B**). Each measure was set as the dependent variable, including the

brain activation of each ROI (**Figure 3**). The main predictor of interest was the caffeinated soda

intake group, whether the individual belongs to the daily-drinking or non-drinking groups. To

ensure that the caffeinated soda intake accounted for more variance than did the confounding

variables, we set a three-level hierarchical model. At the first level, predictors only included the

confounding variables, including sex, age, race, family income, parental education, physical

activity, parental monitoring, family history of drug and alcohol abuse, and externalizing and

internalizing behaviors. At the second level, we included sleep deprivation as an additional

predictor, which is strongly linked to caffeine intake (Roehrs and Roth, 2008). Finally, at the third

level, the caffeinated soda intake group (daily-drinking vs non-drinking group) was added as a

predictor to examine the effect of daily consumption of caffeinated soda. To conclude that

caffeinated soda intake is responsible for group differences in each cognitive factor, two

conditions should be satisfied. First, we conducted an analysis of variance (ANOVA) between the

third-level analysis and the other two lower-level analyses; the third level should show a higher

significance than the previous levels. Also, the categorical caffeinated soda intake regressor

should be statistically significant ($p<0.05$) in the third-level analysis. We tested for these

conditions to determine whether caffeinated soda intake was a significant predictor of the

dependent variable in question.

Daily caffeinated soda intake and future alcohol sipping

Hierarchical logistic regression analysis was used to test whether caffeinated soda intake

at baseline could predict alcohol sipping after 12 months. All the procedures, including the first,

second, and third models, and the criteria of significance are identical to the first analysis (see

Method – Daily caffeinated soda intake and risk factors of SUDs). The only difference was that

the dependent variable was alcohol sipping after 12 months. Among 8,952 participants, 356 were

daily-drinking group, and 3,574 were the non-drinking group. Therefore, 3,930 samples were

used in hierarchical regression analysis.

Daily caffeinated soda intake, alcohol sipping, and risk factors of SUDs

To provide a more integrated insight into how the interactions between daily caffeinated

soda intake and the risk factors of SUDs lead children to later alcohol sipping, we performed the

mediation analysis. Among the risk factors, we selected significant predictors of daily caffeinated

soda intake from the LASSO regression (**Figure 2A**), which results in eighteen variables including

four WM measures (e.g. List Sorting Test, [EN-back] 0-back, [EN-back] L amygdala, [EN-back] R

amygdala), three impulsivity measures (e.g. BAS, UPPS-P, [SST] SSRT), four reward processing

measures (e.g. [MID] earned money, [MID] L NAc, [MID] reward vs. neutral, [MID] L PCC), three

family characteristics (e.g. parental monitoring, family income, parental education), three

demographic measures (e.g. BMI, sex, race), and sleep deprivation. We then performed a

principal component analysis (PCA) and extracted the first principal component (PC1) for the risk

factors that could be grouped as the following six clusters: (1) neurocognitive risk factors

consisting of eleven measures related to WM, impulsivity, and reward processing; (2)

environmental/demographic risk factors with seven measures other than the neurocognitive risk

factors; (3) WM-related risk factors; (4) impulsivity-related risk factors; (5) reward-processing-

related risk factors; (6) family-related risk factors. This leads to twenty-four indices of risk factors

(eighteen variables survived from LASSO regression and six PC1 scores), which could be

included in the models, along with a categorical index of caffeinated soda intake and the single

measure of alcohol sipping collected after 12 months.

With selected variables, we tried to find the best fitting model using the "lavaan" package

(v0.6-8) in R with four elements mediation design: independent variable (X), two mediators (M1,

M2), and dependent variable (Y) (**Figure 4B**). This design was chosen to examine how the

interactions between the three types of risk factors (e.g., caffeinated soda intake, neurocognitive

risk factors, and environmental/demographic risk factors) lead to future alcohol sipping. The

dependent variable was fixed to alcohol sipping, while all possible combinations of the other three

elements were tested (e.g. daily caffeinated soda intake as X and the other two as mediators).

Unlike daily caffeinated soda intake which consists of a single measure, neurocognitive risk

factors and environmental/demographic risk factors consist of multiple indices; therefore, one

index was selected for one model. This results in a total of 405 models for 3 combinations: $_{15}C_1$

(neurocognitive risk factors) * $_9C_1$ (environmental/demographic risk factors) = 135 models for one

combination. See **Table 4-1** in the Supplemental Material to check the models.

For all possible models, we computed the Akaike Information Criterion (AIC) and the

Bayesian Information Criterion (BIC) values, common measures of model comparison; lower

AIC/BIC values suggest a model having a better fit to the data than other models with higher

values (Gana and Broc, 2018). We then determined the one best fitting model with the lowest

AIC/BIC values that best explains the relationship between caffeinated soda intake, alcohol

sipping in 12 months, and neurocognitive and environmental/demographic risk factors. See the

Supplemental Material for examples of the tested models; among 405 models, we selected (1)

models with significant total effect (123 out of 405 models), and (2) models with at least one

significant indirect effect (63 out of 123 models; see **Figure 4-1** and **Table 4-1**).

Results

Sample Characteristics

To characterize group differences as a function of caffeinated soda intake level, we

performed a one-way ANOVA with group factor (four levels of soda intake: non-drinking, monthly-

drinking, weekly-drinking, daily-drinking groups) on all the risk factors and confounding variables.

As a result, ANOVA revealed a significant main effect of group on soda intake per week

($F_{(3, 3962)}$=1161, $p<0.001$) and on all demographic factors, including family income ($F_{(3, }$

$_{3962)}$=46.06, $p<0.001$), parental education ($F_{(3, 3962)}$=51.50, $p<0.001$), parental monitoring

($F_{(3, 3962)}$=7.88, $p<0.001$), BMI ($F_{(3, 3962)}$=19.37, $p<0.001$), physical activity ($F_{(3, }$

$_{3962)}$=11.47, $p<0.001$), sleep deprivation ($F_{(3, 3962)}$=33.19, $p<0.001$), externalizing behaviors

($F_{(3, 3962)}$=7.74, $p<0.001$), and internalizing behaviors ($F_{(3, 3962)}$=3.31, $p=0.019$).

For the neurocognitive risk factors, all the behavioral measures of WM showed a main

effect of group (List Sorting Test: F(3, 3962)=14.16, p<0.001; Card Sort Test: F(3, 3962)=5.68,

p<0.001; [EN-Back] 2-back: F(3, 3962)=3.74, p=0.011; [EN-Back] 0-back: F(3, 3962)=5.60,

p<0.001). Except for the left amygdala activation (F(3, 3962)=4.69, p<0.001), there was no main

group effect on fMRI activation during the EN-Back task (R hippocampus: F(3, 3962)=1.46,

p=0.224; L hippocampus: F(3, 3962)=1.08, p=0.356; R amygdala: F(3, 3962)=2.56, p=0.054; R

DLPFC: F(3, 3962)=0.16, p=0.923; L DLPFC: F(3, 3962)=0.14, p=0.936). For impulsivity, there

was a significant main effect on all behavioral measures except for the SSRT and BAS (SSRT:

F(3, 3962)=1.70, p=0.165; BIS: F(3, 3962)=0.29, p=0.836; UPPS-P: F(3, 3962)=12.62, p<0.001;

BAS: F(3, 3962)=16.67, p<0.001), but not on any neural data during the SST (R caudal ACC: F(3,

3962)=0.58, p=0.631; L caudal ACC: F(3, 3962)=0.81, p=0.489; R IFG: F(3, 3962)=0.53,

p=0.661; L IFG: F(3, 3962)=2.12, p=0.095). Lastly, for the reward processing measures, there

was no group difference in behavioral data ([MID] difference in hit rates between reward and

neutral trials: F(3, 3962)=2.34, p=0.072; [MID] difference in hit rates between loss and neutral

trials: F(3, 3962)=1.99, p=0.113), but main effect of group was found for all neural data except for

the activity in the right NAc (L PCC: F(3, 3962)=6.77, p<0.001; R PCC: F(3, 3962)=5.79, p<0.001;

L NAc: F(3, 3962)=3.21, p=0.022; R NAc: F(3, 3962)=0.72, p=0.539).

Table 1 shows Tukey's posthoc comparison results of demographic factors among the

four groups (non-drinking, monthly-drinking, weekly-drinking, daily-drinking). Several

demographic factors were significantly different among the four groups. Higher caffeinated soda

intake was associated with a lower SES (parental income & parental education), lower parental

monitoring, higher BMI, lower amount of physical activity, and less sleep. Low WM (lower

performance in the List Sorting Test, Card Sort Test, and EN-Back task), high impulsivity (higher

score on the UPPS-P and BAS), and indifference toward monetary rewards (smaller difference in

performance between the reward vs. neutral trials and loss vs. neutral trials) were found in the

daily-drinking group compared to the non-drinking group. Among the ROIs, the amygdala showed

hyperactivation in the daily-drinking group (compared to the non-drinking group) during the EN-

Back task.

Daily caffeinated soda intake and risk factors of SUDs

ANOVA results suggest that daily consumption of caffeinated soda might be a risk factor and a predictor of future substance misuse. However, such group differences could be driven by confounding variables, not by daily consumption of caffeinated soda. Thus, we performed LASSO regression to identify significant predictors of daily caffeinated soda intake. LASSO offers automatic variable selection, which increases the interpretability of the findings especially when some of the variables are significantly correlated with each other. **Figure 2A** shows the multivariate profiles from LASSO regression classifying the daily-drinking group from the non-drinking group. Low parental education was the strongest predictor of daily consumption, along with greater sleep deprivation and low family income. Being male, African American, and/or with a high BMI also strongly predicted allocation to the daily-drinking group.

Among the risk factors of SUDs, high impulsivity measured with the BAS was most predictive of daily caffeinated soda intake. A higher SSRT, indicative of delayed response inhibition, and a higher UPPS-P score were related to the daily intake. Among the WM measures, poor performance in the List Sorting Test and 0-back performance during the EN-Back task predicted daily intake of caffeinated soda. In addition, bilateral amygdala activation during the EN-Back task (2-back vs. 0-back) also predicted daily caffeinated soda intake (**Figure 2C**). For reward processing, the daily-drinking group showed a smaller accuracy difference between reward trials and neutral trials in the MID task, implying a reduced motivation to achieve the monetary reward. Also, hypoactivity in the left PCC and left NAc during the MID task (reward vs. neutral anticipation) predicted the membership of the daily-drinking group (**Figure 2D**). Only the left PCC was negatively correlated with the accuracy difference between reward and neutral trials (left PCC: $r(3964)=-0.03$, $p<0.05$, 95% CI=[-0.06, 0]; left Nac: $r(3964)=0.008$, $p=0.6$, 95% CI=[-0.02, 0.039]), but the correlation was not significant when analyzed separately by group (daily-drinking group: $r(139)=-0.19$, $p=0.1$, 95% CI=[-0.3, 0.03]; non-drinking group: $r(1638)=-0.02$, $p=0.52$, 95% CI=[-0.06, 0.03]). The left Nac was negatively correlated with the UPPS-P score only in the daily-drinking group ($r(139)=-0.2$, $p<0.05$, 95% CI=[-0.357, -0.04]), but not in the non-drinking group ($r(1638)=0.04$, $p=0.07$, 95% CI=[-0.004, 0.093]. **Figure 2B** shows the receiver-

operation characteristic (ROC) curve and its mean area under the curve (AUC) for the

classification of daily-drinking and non-drinking groups. The mean AUC was 0.77 and 0.71 for the

training and test sets.

Next, we performed hierarchical regression analysis to test whether daily caffeinated

soda intake was a significant predictor of each measure of WM, impulsivity, and reward

processing, including the brain activation of each ROI (**Figure 3)** while controlling for confounding

variables (see **Methods**). Specifically, we examined whether caffeinated soda intake accounted

for additional variance than did the other confounding variables. **Table 2** shows the results of the

hierarchical regression analysis using the binary caffeinated soda intake (whether the subjects

are daily drinkers or non-drinkers). Daily consumption of caffeinated soda was negatively

associated with the List Sorting Test score and positively associated with the trait impulsivity

scores of the UPPS-P, the BAS score, and SSRT. The left amygdala showed hyperactivation

during the EN-Back task (2-back vs. 0-back) in the daily-drinking group; although the effect size is

small due to the heterogeneous nature of the ABCD dataset (Karcher et al., 2019; Paulus et al.,

2019; Rosenberg et al., 2020; Cheng et al., 2021), activation of the left amygdala was negatively

correlated with WM capacity, measured by 2-back performance (r(3964)=-0.03, p=0.027, 95%

confidence interval (CI)=[-0.06, -0.0004]) and by the List Sorting Test score (r(3964)=-0.04,

p=0.003, 95% CI=[-0.07, -0.016]).

These results suggest that low WM and high impulsivity are significantly associated with

daily caffeinated soda consumption in both analytic approaches (**Figure 1B**). Furthermore, we

found from the LASSO regression that the indifference toward reward during the MID task, and

some other demographic factors (low SES, less sleep, being male, being African American, and

high BMI) could also distinguish the daily-drinking group from the non-drinking group.

Daily caffeinated soda intake and future alcohol sipping

We tested whether daily caffeinated soda intake could predict alcohol sipping after 12

months by conducting hierarchical logistic regression analysis. Even after controlling the

confounding variables (see **Methods**), daily caffeinated soda consumption could significantly

predict alcohol sipping after 12 months (**Table 3**). Along with lower parental education (*B*=- *i.*041,

p<0.05), lower family income (*B*=0.011, *p*<0.01), and higher BMI (*B*=0.04, *p*<0.05), daily

consumption of caffeinated soda at baseline (*B*=0.06, *p*<0.001) could predict alcohol sipping after

12 months. Also, the adjusted R-squared of the Model 2 and Model 3 showed a significant

difference (model 2 vs. model 3: $F(1, 3916)=12.04$, *p*<0.001), supporting the hypothesis that

future alcohol initiation could be predicted by caffeinated soda drinking behavior in childhood

(**Figure 4A**). These results remained significant even after controlling the alcohol sipping at

baseline (daily caffeinated soda intake: *B*=0.043, *p*<0.01; model 2 vs. model 3: $F(1, 3915)=7.14$,

p<0.01).

Daily caffeinated soda intake, alcohol sipping, and risk factors of SUDs

12   Lastly, we examined all possible mediation models explaining the relationships between

caffeinated soda intake, alcohol sipping in 12 months, and the risk factors of SUDs. The finally

selected best model based on the AIC and BIC values among 405 possible models (see

Methods and **Figure 4-1** and **Table 4-2** in the Supplemental Material) showed significant indirect

effects of neurocognitive risk factors on alcohol sipping after 12 months through family risk factors

and daily caffeinated soda intake mediating the relationships (indirect effect of family risk factor:

B=0.019, SE=0.002, *p*<0.05; indirect effect of caffeinated soda intake: *B*=-0.008, SE=0.036,

p<0.05; direct effect: *B*=-0.059, SE=0.025, *p*<0.01). These results also support our hypothesis

that daily caffeinated soda intake could predict future alcohol sipping, and suggest possible

mechanisms of alcohol use initiation and the risk factors from SUDs.

Discussion

24   The present results demonstrated that daily caffeinated soda consumption in childhood is

25   associated with well-known neurocognitive risk factors of SUDs, and could predict future alcohol

26   sipping. The results of this study are consistent with previous studies of caffeine consumption

27   during adolescence and its association with future substance misuse (Barrense-Dias et al., 2016;

28   Leal and Jackson, 2019), supporting a higher risk of daily caffeinated soda consumption in

childhood, particularly regarding vulnerabilities to future substance misuse. However, this is the first study investigating the direct link between caffeinated soda intake in childhood and the risks of substance use. Such information is invaluable since caffeinated soda is incomparably the most common medium of caffeine consumption in childhood and the risk of substance misuse should be detected before adolescence, the most common period of the onset of substance use.

The first evidence suggesting the risks of daily consumption of caffeinated soda was its strong association with the neurocognitive risk factors of SUDs. For the robustness of the results, we utilized multiple analytic approaches with two analytical methods (LASSO regression and hierarchical regression; **Figure 1B**). Low WM performance (List Sorting Test score) and high impulsivity scores (high UPPS-P and BAS) were consistently associated with greater caffeinated soda consumption across all different approaches. In particular, we showed hyperactivity of the amygdala during the EN-Back task in children with daily caffeinated soda consumption, and found greater behavioral WM impairments during the task in the daily-drinking group than in the non-drinking group. These results are consistent with the developmental mismatch hypothesis, which denote that subcortical structures of affect and reward processing develop earlier than prefrontal structures involved in the top-down regulation of behavior in children and adolescents (Somerville et al., 2010; Mills et al., 2014). Based on the previous literature that cognitive control during the EN-Back task is related to WM capacity in the prefrontal cortex and simultaneous emotional regulation in the limbic system (Gee et al., 2013; Casey et al., 2018), it is possible that the hyperactivation of the amygdala and the negative association between the left amygdala and WM capacity observed in daily caffeinated soda drinkers was due to the unsuccessful suppression of emotional processes during WM processing. We also found altered reward processing in the daily-drinking group; that is, lower activation in the left NAc was associated with higher trait impulsivity in the daily-drinking group. This finding supports previous literature suggesting that blunted reward sensitivity in the ventral striatum might motivate impulsive or risky behaviors, such as the use of substances or pathological gambling, to receive sufficient satisfaction (Beck et al., 2009; Balodis et al., 2012).

The second evidence was acquired from examining whether daily intake of caffeinated

soda could predict alcohol sipping in 12 months. In the hierarchical regression controlling other

confounding variables, daily consumption of caffeinated soda could significantly predict future

alcohol sipping. This relationship could be explained by our third analysis, where we explored all

possible mediation models with daily caffeinated soda consumption, future alcohol sipping, and

the risk factors of SUDs that came out to be significant predictors of caffeinated soda intake in

LASSO regression. The best model (**Figure 4B**) suggests that daily intake of caffeinated soda

could work as a mediator between the neurocognitive risk factors of SUDs and future alcohol

sipping. These findings suggest that caffeinated soda itself would not directly cause children to

initiate substance use, but the daily drinking behavior of the beverages could indicate a high risk

of substance initiation in the near future. Because caffeinated soda combines two addictive

substances, sugar and caffeine, it is somewhat natural for children to prefer and seek the

beverages. While the preference of the taste could lead to a seeking behavior and habitual

consumption, previous literature has shown impulsivity as one of the most common traits of SUDs

(Whiteside and Lynam, 2003; Magid and Colder, 2007; Cyders et al., 2009). Therefore, the

behavior of daily consuming caffeinated soda daily could indicate a high risk of substance

initiation in the future, though further research should clearly disentangle the complex

relationships.

Along with daily caffeinated soda intake, family risk factors (PC1 of parental monitoring,

family income, and parental education) also mediated the relationship between neurocognitive

risk factors and future alcohol sipping (**Figure 4B**). This finding is in line with previous literature,

suggesting that parental monitoring and high socioeconomic status could have protective effects

on future alcohol use or alcohol-related problems (Kendler et al., 2014; Carroll et al., 2016;

Collins, 2016; Calling et al., 2019). We acknowledge that multiple mediators could exist between

neurocognitive risk factors and future alcohol use other than caffeinated soda intake, so future

research should extensively investigate how caffeinated soda intake interacts with other

risk/protective factors of SUDs.

1   Our results have important implications for public health recommendations since our

2 study provides novel insights into the neurocognitive correlates of daily caffeinated soda

3 consumption in children, which has been rarely tested yet. At the same time, we should mention a

4 few limitations in our research and future directions for further investigations. First, because we

5 used an ROI-based approach to analyze the neuroimaging data, we could not perform functional

6 connectivity analyses or multivoxel pattern analysis, which might provide additional valuable

7 insights into the effects of daily caffeinated soda intake. Second, since the samples of 9-10-year-

8 old children have not yet started other substances except for alcohol, such as tobacco or

9 marijuana, future work using the longitudinal 10-year follow-up data of the ABCD study should

10 examine whether daily caffeinated soda consumption is associated with alcohol or other

11 substance misuses as well.

12   In conclusion, our results revealed the potential risks of caffeinated soda consumption in

13 children by (1) investigating the associations between daily caffeinated soda consumption and

14 SUD risk factors, and (2) examining the predictive abilities of daily caffeinated soda consumption

15 for future alcohol sipping, in a large dataset of the ABCD study. While previous research

16 regarding the side effects of caffeinated soda has been mostly limited to negative physical

17 consequences, the present results strongly suggest that daily caffeinated soda drinking in

18 children is also associated with alterations in neurocognitive functions, and can predict alcohol

19 sipping in 12 months. Our study further suggests that there is a strong need to develop evidence-

20 based recommendations on caffeinated soda (Temple, 2018), as there is no consensus on a safe

21 dose of caffeinated soda for children and a possibility that certain children are at higher risk for

22 adverse events from habitual caffeinated soda intake. Further clarification on causal relationships

23 and neuro-developmental evidence is needed to determine whether caffeinated soda is indeed a

24 warning sign for future substance misuse and whether it induces neurocognitive impairments in

25 children.

Acknowledgments

We thank the families and children who participated, and continue to participate, in the ABCD study, as well as staff at the study sites who are involved in data collection and curating the data release. Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators. The ABCD data repository grows and changes over time. The ABCD data used in this report came from [NIMH Data Archive Digital Object Identifier (10.15154/1503209)]. DOIs can be found at http://dx.doi.org/10.15154/1503209.

Funding

This work was supported by the Basic Science Research Program through the National Research Foundation (NRF) of Korea funded by the Ministry of Science, ICT, & Future Planning (NRF-2018R1C1B3007313 and NRF-2018R1A4A1025891); the Institute for Information & Communications Technology Planning & Evaluation (IITP) grant funded by the Korea government (MSIT) (No. 2019-0-01367, BabyMind); the National Research Foundation of Korea (NRF) grant

1   funded by the Korea government (MSIT) (No. 2021M3E5D2A0102249311); and the Creative-

2   Pioneering Researchers Program through Seoul National University to W.-Y.A.

References

1 Achenbach TM (2009) Achenbach system of empirically based assessment (aseba): Development, findings, theory, and applications. University of Vermont, Research Center of Children, Youth & Families.

Ahn W-Y, Hendricks P, Haines N (2017) Easyml: Easily Build and Evaluate Machine Learning Models.

Ahn W-Y, Kishida KT, Gu X, Lohrenz T, Harvey A, Alford JR, Smith KB, Yaffe G, Hibbing JR, Dayan P, Montague PR (2014) Nonpolitical Images Evoke Neural Predictors of Political Ideology. Curr Biol 24:2693–2699.

Ahn W-Y, Vassileva J (2016) Machine-learning identifies substance-specific behavioral markers for opiate and stimulant dependence. Drug Alcohol Depen 161:247–257.

Arria AM, Caldeira KM, Kasperski SJ, Vincent KB, Griffiths RR, O'Grady KE (2011) Energy Drink Consumption and Increased Risk for Alcohol Dependence. Alcohol Clin Exp Res 35:365–375.

Avena NM, Rada P, Hoebel BG (2008) Evidence for sugar addiction: Behavioral and neurochemical effects of intermittent, excessive sugar intake. Neurosci Biobehav Rev 32:20–39.

Balodis IM, Kober H, Worhunsky PD, Stevens MC, Pearlson GD, Potenza MN (2012) Diminished Frontostriatal Activity During Processing of Monetary Rewards and Losses in Pathological Gambling. Biol Psychiat 71:749–757.

Balodis IM, Potenza MN (2015) Anticipatory Reward Processing in Addicted Populations: A Focus on the Monetary Incentive Delay Task. Biol Psychiat 77:434–444.

Banks SJ, Eddy KT, Angstadt M, Nathan PJ, Phan KL (2007) Amygdala–frontal connectivity

during emotion regulation. Soc Cogn Affect Neur 2:303–312.

Barbey AK, Koenigs M, Grafman J (2013) Dorsolateral prefrontal contributions to human working

memory. Cortex 49:1195–1205.

Barrense-Dias Y, Berchtold A, Akre C, Surís J (2016) Consuming energy drinks at the age of 14

predicted legal and illegal substance use at 16. Acta Paediatr 105:1361–1368.

Barry AE, King J, Sears C, Harville C, Bondoc I, Joseph K (2016) Prioritizing Alcohol Prevention:

Establishing Alcohol as the Gateway Drug and Linking Age of First Drink With Illicit Drug Use.

J School Health 86:31–38.

Bechara A, Berridge KC, Bickel WK, Morón JA, Williams SB, Stein JS (2019) A Neurobehavioral

Approach to Addiction: Implications for the Opioid Epidemic and the Psychology of Addiction.

Psychol Sci Publ Int 20:96–127.

Beck A, Schlagenhauf F, Wüstenberg T, Hein J, Kienast T, Kahnt T, Schmack K, Hägele C,

Knutson B, Heinz A, Wrase J (2009) Ventral Striatal Activation During Reward Anticipation

Correlates with Impulsivity in Alcoholics. Biol Psychiat 66:734–742.

Calling S, Ohlsson H, Sundquist J, Sundquist K, Kendler KS (2019) Socioeconomic status and

alcohol use disorders across the lifespan: A co-relative control study. Plos One 14:e0224127.

Carroll HA, Heleniak C, Witkiewitz K, Lewis M, Eakins D, Staples J, Andersson C, Berglund M,

Larimer ME (2016) Effects of parental monitoring on alcohol use in the US and Sweden: A

brief report. Addict Behav 63:89–92.

Carver CS, White TL (1994) Behavioral inhibition, behavioral activation, and affective responses

to impending reward and punishment: The BIS/BAS Scales. J Pers Soc Psychol 67:319.

Casey BJ et al. (2018) The Adolescent Brain Cognitive Development (ABCD) study: Imaging

acquisition across 21 sites. Dev Cogn Neuros-neth 32:43–54.

Chambers RA, Taylor JR, Potenza MN (2003) Developmental Neurocircuitry of Motivation in

Adolescence: A Critical Period of Addiction Vulnerability. Am J Psychiat 160:1041–1052.

Cheng W, Rolls E, Gong W, Du J, Zhang J, Zhang X-Y, Li F, Feng J (2021) Sleep duration, brain

structure, and psychiatric and cognitive problems in children. Mol Psychiatr 26:3992–4003.

Cohen AO, Breiner K, Steinberg L, Bonnie RJ, Scott ES, Taylor-Thompson K, Rudolph MD,

Chein J, Richeson JA, Heller AS, Silverman MR, Dellarco DV, Fair DA, Galván A, Casey BJ

(2015) When Is an Adolescent an Adult? Assessing Cognitive Control in Emotional and

Nonemotional Contexts. Psychol Sci 27:549–562.

Collins SE (2016) Associations Between Socioeconomic Factors and Alcohol Outcomes. Alcohol

Res Curr Rev 38:83–94.

Cyders MA, Flory K, Rainer S, Smith GT (2009) The role of personality dispositions to risky

behavior in predicting first-year college drinking. Addiction 104:193–202.

Day JJ, Carelli RM (2007) The Nucleus Accumbens and Pavlovian Reward Learning. Neurosci

13:148–159.

Donovan JE, Molina BSG (2011) Childhood Risk Factors for Early-Onset Drinking*. J Stud

Alcohol Drugs 72:741–751.

Friedman J, Hastie T, Tibshirani R (2010) Regularization Paths for Generalized Linear Models via

Coordinate Descent. J Stat Softw 33:1–22.

Gana K, Broc G (2018) Structural Equation Modeling with lavaan.

Garavan H, Bartsch H, Conway K, Decastro A, Goldstein R, Heeringa S, Jernigan T, Potter A, Thompson W, Zahs D (2018) Recruiting the ABCD Sample: Design Considerations and Procedures. Dev Cogn Neuros-neth 32:16–22.

Gee DG, Humphreys KL, Flannery J, Goff B, Telzer EH, Shapiro M, Hare TA, Bookheimer SY, Tottenham N (2013) A developmental shift from positive to negative connectivity in human amygdala-prefrontal circuitry. J Neurosci Official J Soc Neurosci 33:4584–4593.

Jackson KM, Barnett NP, Colby SM, Rogers ML (2015) The Prospective Association Between Sipping Alcohol by the Sixth Grade and Later Substance Use. J Stud Alcohol Drugs 76:212–221.

Jacobson L, Javitt DC, Lavidor M (2011) Activation of Inhibition: Diminishing Impulsive Behavior by Direct Current Stimulation over the Inferior Frontal Gyrus. J Cognitive Neurosci 23:3380–3387.

James JE, Kristjánsson ÁL, Sigfúsdóttir ID (2011) Adolescent substance use, sleep, and academic achievement: Evidence of harm due to caffeine. J Adolescence 34:665–673.

Jordan CJ, Andersen SL (2017) Sensitive periods of substance abuse: Early risk for the transition to dependence. Dev Cogn Neuros-neth 25:29–44.

Karcher NR, O'Brien KJ, Kandala S, Barch DM (2019) Resting-State Functional Connectivity and Psychotic-like Experiences in Childhood: Results From the Adolescent Brain Cognitive Development Study. Biol Psychiat 86:7–15.

Kendler KS, Gardner CO, Hickman M, Heron J, Macleod J, Lewis G, Dick DM (2014) Socioeconomic Status and Alcohol-Related Behaviors in Mid- to Late Adolescence in the Avon Longitudinal Study of Parents and Children. J Stud Alcohol Drugs 75:541–545.

Kirby T, Barry AE (2012) Alcohol as a gateway drug: a study of US 12th graders. J Sch Heal

82:371–379.

Knutson B, Westdorp A, Kaiser E, Hommer D (2000) FMRI Visualization of Brain Activity during a

Monetary Incentive Delay Task. Neuroimage 12:20–27.

Leal WE, Jackson DB (2019) The role of energy drink consumption in the intention to initiate

marijuana use among adolescents. Addict Behav 93:240–245.

Lisdahl KM et al. (2021) Substance use patterns in 9-10 year olds: Baseline findings from the

adolescent brain cognitive development (ABCD) study. Drug Alcohol Depen:108946.

Logan GD, Cowan WB, Davis KA (1984) On the ability to inhibit simple and choice reaction time

responses: A model and a method. J Exp Psychology Hum Percept Perform 10:276–291.

Luciana M, Bjork JM, Nagel B, Barch DM, Gonzalez R, Nixon SJ, Banich MT (2018) Adolescent

Neurocognitive Development and Impacts of Substance Use: Overview of the Adolescent

Brain Cognitive Development (ABCD) Baseline Neurocognition Battery. Dev Cogn Neuros-

neth 32:67–79.

Luijten M, Schellekens AF, Kühn S, Machielse MWJ, Sescousse G (2017) Disruption of Reward

Processing in Addiction : An Image-Based Meta-analysis of Functional Magnetic Resonance

Imaging Studies. Jama Psychiat 74:387.

Magid V, Colder CR (2007) The UPPS Impulsive Behavior Scale: Factor structure and

associations with college drinking. Pers Indiv Differ 43:1927–1937.

Marmorstein NR (2018) Investigating associations between caffeinated beverage consumption

and later alcohol consumption among early adolescents. Addict Behav 90:362–368.

Mills KL, Goddings A-L, Clasen LS, Giedd JN, Blakemore S-J (2014) The Developmental

Mismatch in Structural Brain Maturation during Adolescence. Dev Neurosci-basel 36:147–160.

Miyake ER, Marmorstein NR (2015) Energy drink consumption and later alcohol use among early adolescents. Addict Behav 43:60–65.

Mullee A et al. (2019) Association Between Soft Drink Consumption and Mortality in 10 European Countries. Jama Intern Med 179.

Nehlig A (1999) Are we dependent upon coffee and caffeine? A review on human and animal data. Neurosci Biobehav Rev 23:563–576.

Newman JP, MacCoon DG, Vaughn LJ, Sadeh N (2005) Validating a Distinction Between Primary and Secondary Psychopathy With Measures of Gray's BIS and BAS Constructs. J Abnorm Psychol 114:319–323.

Paulus MP, Squeglia LM, Bagot K, Jacobus J, Kuplicki R, Breslin FJ, Bodurka J, Morris AS, Thompson WK, Bartsch H, Tapert SF (2019) Screen media activity and brain structure in youth: Evidence for diverse structural correlation networks from the ABCD study. Neuroimage 185:140–153.

Porciúncula LO, Sallaberry C, Mioranzza S, Botton PHS, Rosemberg DB (2013) The Janus face of caffeine. Neurochem Int 63:594–609.

Robinson TE, Berridge KC (2008) The incentive sensitization theory of addiction: some current issues. Philosophical Transactions Royal Soc B Biological Sci 363:3137–3146.

Roehrs T, Roth T (2008) Caffeine: Sleep and daytime sleepiness. Sleep Med Rev 12:153–162.

Rosenberg MD, Martinez SA, Rapuano KM, Conley MI, Cohen AO, Cornejo MD, Hagler DJ, Meredith WJ, Anderson KM, Wager TD, Feczko E, Earl E, Fair DA, Barch DM, Watts R, Casey BJ (2020) Behavioral and Neural Signatures of Working Memory in Childhood. J Neurosci 40:5090–5104.

Schachar R, Logan GD (1990) Impulsivity and inhibitory control in normal development and

childhood psychopathology. Dev Psychol 26:710–720.

Solnick SJ, Hemenway D (2013) Soft drinks, aggression and suicidal behaviour in US high school

students. Int J Inj Control Sa 21:266–273.

Somerville LH, Hare T, Casey BJ (2010) Frontostriatal maturation predicts cognitive control

failure to appetitive cues in adolescents. J Cognitive Neurosci 23:2123–2134.

Stadler C, Sterzer P, Schmeck K, Krebs A, Kleinschmidt A, Poustka F (2007) Reduced anterior

cingulate activation in aggressive children and adolescents during affective stimulation:

Association with temperament traits. J Psychiatr Res 41:410–417.

Suglia SF, Solnick S, Hemenway D (2013) Soft Drinks Consumption Is Associated with Behavior

Problems in 5-Year-Olds. J Pediatrics 163:1323–1328.

Sylvetsky AC, Parnarouskis L, Merkel PE, Gearhardt AN (2020) Children's Sugar-Sweetened

Beverage Consumption: Striking Parallels With Substance Use Disorder Symptoms. Frontiers

Pediatrics 8:594513.

Temple JL (2009) Caffeine use in children: What we know, what we have left to learn, and why

we should worry. Neurosci Biobehav Rev 33:793–806.

Temple JL (2018) Trends, Safety, and Recommendations For Caffeine Use in Children and

Adolescents. J Am Acad Child Adolesc Psychiatry 58:36–45.

Thompson WK, Barch D, Bjork J, Gonzalez R, Nagel B, Nixon SJ, Luciana M (2018) The

Structure of Cognition in 9 and 10 year-old Children and Associations with Problem Behaviors:

Findings from the ABCD Study's Baseline Neurocognitive Battery. Dev Cogn Neuros-neth

36:100606.

Tibshirani R (1996) Regression Shrinkage and Selection Via the Lasso. J Royal Statistical Soc

Ser B Methodol 58:267–288.

Tibshirani R (2011) Regression shrinkage and selection via the lasso: a retrospective. J Royal

Statistical Soc Ser B Statistical Methodol 73:273–282.

Tulsky DS, Carlozzi NE, Chevalier N, Espy KA, Beaumont JL, Mungas D (2013) V. NIH Toolbox

Cognition Battery (CB): measuring working memory. Monogr Soc Res Child 78:70–87.

Vercammen KA, Koma JW, Bleich SN (2019) Trends in Energy Drink Consumption Among U.S.

Adolescents and Adults, 2003–2016. Am J Prev Med 56:827–833.

Wagner FA, Anthony JC (2002) From First Drug Use to Drug Dependence: Developmental

Periods of Risk for Dependence upon Marijuana, Cocaine, and Alcohol.

Neuropsychopharmacol 26:479–488.

Watts AL, Wood PK, Jackson KM, Lisdahl KM, Heitzeg MM, Gonzalez R, Tapert SF, Barch DM,

Sher KJ (2021) Incipient alcohol use in childhood: Early alcohol sipping and its relations with

psychopathology and personality. Dev Psychopathol 33:1338–1350.

Whiteside SP, Lynam DR (2003) Understanding the Role of Impulsivity and Externalizing

Psychopathology in Alcohol Abuse: Application of the UPPS Impulsive Behavior Scale. Exp

Clin Psychopharm 11:210–217.

Zou H, Hastie T (2005) Regularization and variable selection via the elastic net. J Royal

Statistical Soc Ser B Statistical Methodol 67:301–320.

Zuckerman M (2001) International Encyclopedia of the Social & Behavioral Sciences. Dev

Psychology Personality Dev Motivational Emot Psychology Overv Classif Methods Personality

Psychology Self Personality Risk Article Titles S:13905–13908.

Figure 1. An overview of the study design

A. Variables used to measure three main cognitive functions: working memory, impulsivity, and

reward processing. Working Memory was measured from the performance of 2-back in the

Emotional N-Back Task (EN-Back) performed inside the functional magnetic resonance imaging

(fMRI) scanner, List Sorting Test, and Card Sort Test. The EN-Back task's fMRI data was

analyzed based on the contrast of 2-back vs. 0-back, with hippocampus, amygdala, and

dorsolateral prefrontal cortex as the regions of interest (ROIs). Impulsivity was measured by stop

signal reaction time (SSRT) during the Stop Signal Task (SST) performed inside the fMRI

scanner, Urgency-Premeditation-Perseverance-Sensation Seeking-Positive Urgency (UPPS-P),

and Behavioral Inhibition System/ Behavioral Activation System (BIS/BAS). The SST's fMRI data

was analyzed based on the contrast of correct stop vs. correct go, with posterior cingulate cortex

and inferior frontal gyrus as the ROIs. Reward processing was measured by comparing the

success rate of reward vs. neutral and loss vs. neutral during the Monetary Incentive Delay (MID)

Task performed inside the fMRI scanner and cash choice task. The MID task's fMRI data was

analyzed based on the contrast of reward anticipation vs. neutral anticipation, anterior cingulate

cortex and nucleus accumbens as the ROIs. **B.** Two different analytic methods were performed.

Hierarchical regression examines whether caffeinated soda intake is a significant predictor of

each neurocognitive variables, while a machine learning approach examines which variables are

significant predictors of caffeinated soda intake.

Figure 2. Results of LASSO regression with categorical caffeinated soda intake (non-drinking group vs. daily-drinking group).

A. The estimates of coefficients (x axis: coefficient estimates; y axis: predictor). **B.** Representative receiver-operation characteristic (ROC) curve (left) and a distribution of area under the curve (AUC) scores (right) in training and test datasets. **C.** Regions of interest (ROIs) identified as having significant estimates of coefficients during the Emotional N-Back Task (EN-Back). **D.** ROIs identified as having significant estimates of coefficients during the Monetary Incentive Delay (MID) Task. The bar graph at the top of **C** and **D** displays activation pattern of entire four groups (error bars represent ±1 standard error of the mean). The color bar indicates the estimates of coefficients of LASSO regression.

Abbreviations. LASSO, least absolute shrinkage and selection operator; AA, African American; BMI, body mass index; BIS, Behavioral Inhibition System; BAS, Behavioral Activation System; UPPS-P, Urgency-Premeditation-Perseverance-Sensation Seeking-Positive Urgency; SST, Stop Signal Task; SSRT, Stop Signal Reaction Time; DLPFC, Dorsolateral Prefrontal Cortex; NAc, Nucleus Accumbens; PCC, Posterior Cingulate Cortex; IFG, Inferior Frontal Gyrus; ACC, Anterior Cingulate Cortex.

Figure 3. Outline of hierarchical regression analysis

Independent variables were added hierarchically: control variables in the first level, sleep

deprivation added in the second level, and caffeinated soda intake in the third level. Dependent

variables were put into hierarchical model one by one. Therefore, a total of 24 models were

tested, including each behavior measure (BIS and BAS each), and left and right regions of

interest of fMRI data.

Abbreviations. SES, Socioeconomic status; BMI, body mass index; EN-Back, Emotional N-Back

task; SST, Stop Signal Task; SSRT, Stop Signal Reaction Time; UPPS-P, Urgency-

Premeditation-Perseverance-Sensation Seeking-Positive Urgency; BIS/BAS, Behavioral Inhibition

System/ Behavioral Activation System; MID, Monetary Incentive Delay Task; DLPFC,

Dorsolateral Prefrontal Cortex; ACC, Anterior Cingulate Cortex; IFG, Inferior Frontal Gyrus; PCC,

Posterior Cingulate Cortex; NAc, Nucleus Accumbens.

Figure 4. Association between caffeinated soda consumption at baseline and alcohol sipping after

12 months

A. Caffeinated soda intake at baseline predicting alcohol sipping after 12 months. **B.** The best

model explaining the relationships between daily caffeinated soda intake and alcohol sipping after

12 months. See **Figure 4-1** and **Table 4-1** in the Supplemental Material for more examples of the

tested models.

*: $p<0.05$, **: $p<0.01$; ***: $p<0.001$

 Table 1. Baseline characteristics of participants

variable	Non-drinking (0 can)	Monthly-drinking (< 1 can /week)	Weekly-drinking (< 7 can / week)	Daily-drinking (≥ 7 can / week)
Sample size	1640	1721	464	141
Soda intake / week	0	0.46 (0.01)***	2.86 (0.06)***	14.92 (1.32)***
demographic variables				
Women, No. (%)	917 (0.56)	820 (0.48)	173 (0.37)	66 (0.47)
Age	9.5 (0.01)	9.55 (0.01)**	9.64 (0.02)***	9.49 (0.04)
Family income	8.06 (0.04)	7.8 (0.05)***	7.07 (0.11)***	6.65 (0.21)***
Parental education	17.4 (0.05)	16.9 (0.06)***	16.19 (0.11)***	15.83 (0.22)***
Parental monitoring	4.47 (0.01)	4.44 (0.01)	4.37 (0.02)***	4.35 (0.05)*
Family history (drug)	0.29 (0.01)	0.3 (0.01)	0.36 (0.02)*	0.34 (0.04)
Family history (alcohol)	0.43 (0.01)	0.48 (0.01)**	0.46 (0.02)	0.42 (0.04)
BMI	17.9 (0.09)	18.5 (0.09)***	19.11 (0.2)***	19.6 (0.45)***
Physical activity	4.20 (0.06)	3.94 (0.06)*	3.52 (0.11)***	3.44 (0.20)**
Sleep deprivation	1.50 (0.02)	1.61 (0.02)***	1.81 (0.04)***	1.93 (0.08)***
Externalizing behaviors	44.01 (0.23)	44.12 (0.23)	46.14 (0.48)***	45.93 (0.91)
Internalizing behaviors	44.61 (0.26)	44.10 (0.25)*	45.84 (0.51)	46.62 (0.93)
Working memory				
Behavioral measures				
List Sorting Test	104.61 (0.34)	103.54 (0.33)	101.59 (0.63)***	97.91 (1.34)***
Card Sort Test	99.88 (0.38)	99.39 (0.37)	95.64 (1.23)*	95.57 (1.15)**
[EN-Back] (2-back)	0.81 (0.002)	0.80 (0.002)	0.80 (0.004)	0.79 (0.007)*
[EN-Back] (0-back)	0.88 (0.002)	0.88 (0.002)	0.87 (0.004)	0.85 (0.008)***
fMRI data				
[EN-Back] Right hippocampus	-0.07 (0.005)	-0.07 (0.005)	-0.06 (0.01)	-0.047 (0.02)
[EN-Back] Left hippocampus	-0.079 (0.005)	-0.082 (0.005)	-0.07 (0.01)	-0.057 (0.018)
[EN-Back] Right DLPFC	0.062 (0.006)	0.059 (0.006)	0.064 (0.012)	0.051 (0.021)
[EN-Back] Left DLPFC	0.042 (0.005)	0.04 (0.006)	0.044 (0.012)	0.031 (0.02)
[EN-Back] Right amygdala	-0.071 (0.007)	-0.075 (0.007)	0.043 (0.014)	-0.026 (0.029)
[EN-Back] Left amygdala	-0.066 (0.006)	-0.075 (0.006)	-0.044 (0.012)	-0.003 (0.024)*

Impulsivity

   Behavioral measures

UPPS-P	39.70 (0.18)	40.82 (0.18)***	41.56 (0.37)***	41.94 (0.63)**
BIS	9.43 (0.09)	9.53 (0.09)	9.52 (0.17)	9.57 (0.32)
BAS	19.58 (0.16)	20.54 (0.16)***	21.35 (0.31)***	22.53 (0.58)***
[SST] SSRT	299.51 (1.62)	302.29 (1.55)	297.27 (2.87)	308.76 (5.59)

  fMRI data

[SST] Right caudal ACC	0.068 (0.003)	0.073 (0.003)	0.072 (0.007)	0.062 (0.012)
[SST] Left caudal ACC	0.062 (0.003)	0.068 (0.003)	0.07 (0.007)	0.061(0.01)
[SST] Right IFG	0.103 (0.004)	0.109 (0.004)	0.101 (0.007)	0.104 (0.014)
[SST] Left IFG	0.057 (0.003)	0.069 (0.003)*	0.058 (0.006)	0.059 (0.013)

Reward processing

  Behavioral measures

Cash Choice Task	1.61 (0.01)	1.59 (0.01)	1.61 (0.02)	1.60 (0.04)
[MID] earned money	21.23 (0.30)	22.14 (0.29)*	21.983(0.58)	22.34 (1.11)
[MID] reward vs. neutral	0.12 (0.003)	0.11 (0.003)	0.106 (0.006)	0.09 (0.01)*
[MID] loss vs. neutral	0.095 (0.003)	0.093 (0.003)	0.083 (0.006)	0.071(0.012)*

  fMRI data

[MID] Right PCC	0.024 (0.004)	-8.7e-05 (0.004)***	0.006 (0.008)*	0.004 (0.015)
[MID] Left PCC	0.014 (0.004)	-0.011 (0.004)***	-0.005 (0.008)*	-0.013 (0.014)
[MID] Right nucleus accumbens	0.060 (0.005)	0.05 (0.005)	0.057 (0.01)	0.063 (0.02)
[MID] Left nucleus accumbens	0.082 (0.005)	0.063 (0.005)	0.050 (0.02)	0.044 (0.017)

1 *Note.* Data are Mean (Standard Error); *: $p<0.05$, **: $p<0.01$; ***: $p<0.001$ (each soda drinking

2 groups (monthly-drinking, weekly-drinking, daily-drinking) compared with the non-drinking group)

3 *Abbreviations.* BMI, Body Mass Index; EN-Back, Emotional N-Back Task; DLPFC, Dorsal Lateral

4 Prefrontal Cortex; SST, Stop Signal Task; ACC, Anterior Cingulate Cortex; IFG, Inferior Frontal

5 Gyrus; MID, Monetary Incentive Delay Task; [MID] reward vs. neutral, difference in percentage of

1   correct trials between reward trial and neutral trial; [MID] loss vs. neutral, difference in percentage

2   of correct trials between loss trial and neutral trial; PCC, Posterior Cingulate Cortex.

Table 2. Results of hierarchical regression analysis predicting each neurocognitive risk factor of SUDs

variable	Working Memory		Impulsivity		
	List sorting Test	[EN-Back] L amygdala	SSRT	UPPS-P	BAS
sex	-.051*	-.008	.021	-.125***	-.112***
race (African American)	-.133*	-.018	-.112	-.023	.052
race (others)	.042	-.023	.03	.031	-.038
parental education	.161***	-.081**	-.014	.015	-.028
family income	.024	-.021	.03	-.013	-.055
parental monitoring	.057*	-.042	-.022	-.185***	.055
body mass index (BMI)	-.012	.007	.009	-.029	.01
family history (drug)	.001	-.019	-.046	.022	.04
family history (alcohol)	.009	-.01	.05	-.013	-.04
externalizing behaviors	-.032	.012	.025	.212***	.118***
internalizing behaviors	-.028	-.024	-.077**	-.112***	-.084**
sleep deprivation	-.042	-.023	-.009	.033	.086***
daily caffeinated soda intake	-.221**	.146*	.146*	.160*	.232**

Note. The values are standardized regression coefficients from the third step in the hierarchical

analyses among the 24 dependent variables, the five variables were significantly associated with

daily consumption of caffeinated soda, even after controlling the controlled variables (eleven

variables from the top) and the sleep deprivation. Higher score in the List sorting Test score was

associated with lower intake of caffeinated soda, while other four variables (left amygdala

activation during the Emotional N-Back Task (EN-Back), stop signal reaction time (SSRT),

Urgency-Premeditation-Perseverance-Sensation Seeking-Positive Urgency (UPPS-P), and

Behavioral Activation System (BAS) were associated with higher intake of caffeinated soda.

*: $p<0.05$, **: $p<0.01$; ***: $p<0.001$

Table 3. Results of hierarchical regression analysis predicting alcohol sipping after 12 months

Variables	Model 1	Model 2	Model 3
sex	-.03*	.032*	-.029
race (African American)	-.036	-.038	-.045
race (others)	-.147	-.149	-.137
parental education	.034	.035	-.041*
family income	.079*	.08*	.011*
parental monitoring	.002*	.002	.028
BMI	.005	.005*	.004*
family history (drug)	-.03	-.029	-.03
family history (alcohol)	.341	.34	.346
externalizing behaviors	.025	.024	.02
internalizing behaviors	-.001	-.001	-.001
amount of sleep		.004	.003
daily caffeinated soda intake			.1***

1 *Note.* The values are standardized regression coefficients from each step in the hierarchical

2 analyses predicting alcohol sipping after 12 months. Daily consumption of caffeinated soda intake

3 could predict higher alcohol sipping tendency after 12 months even after controlling the controlled

4 variables (eleven variables in Model 1) and the sleep deprivation.

5 *: $p<0.05$, **: $p<0.01$; ***: $p<0.001$