Title: Caffeinated soda intake in children is associated with neurobehavioral risk factors

for substance misuse

Authors:

Mina Kwon[1], Hyeonjin Kim[1], Jaeyeong Yang[1], Yoseph Lee[1], Jihyun K. Hur[1,2], Tae-Ho

Lee[3], James M. Bjork[4], and Woo-Young Ahn*[1,5]

[1]Department of Psychology, Seoul National University, Seoul, Korea 08826

[2]Department of Psychology, Yale University, New Haven, CT 06511

[3]Department of Psychology, Virginia Tech, Blacksburg, VA 24060

[4]Department of Psychiatry, Virginia Commonwealth University, Richmond, VA 23298

[5]Department of Brain and Cognitive Sciences, Seoul National University, Seoul, Korea

08826

Corresponding author:

Woo-Young Ahn, Ph.D.

Department of Psychology

Seoul National University

Seoul, Korea 08826

Tel: +82-2-880-2538, Fax: +82-2-877-6428. E-mail: wahn55@snu.ac.kr

Abstract

Background and Objectives: Use of psychotropic substances in childhood has been associated with both impulsivity and other manifestations of poor executive function as well as escalation over time to use of progressively stronger substances. However, how this relationship may start in earlier childhood has not been well explored. Here, we investigated the neurobehavioral correlates of daily caffeinated soda consumption in preadolescent children and examined whether caffeinated soda intake is associated with a higher risk of subsequent alcohol initiation. **Methods:** Using Adolescent Brain Cognitive Development study data, we first investigated cross-sectional relationships between frequent caffeinated soda intake and well-known risk factors of substance misuse: impaired working memory, high impulsivity, and aberrant reward processing. We then examined whether caffeinated soda intake at baseline predicts more alcohol sipping at 12 months follow-up using a machine learning algorithm. **Results:** Daily consumption of caffeinated soda was cross-sectionally associated with neurobehavioral risk factors for substance misuse such as higher impulsivity scores and lower working memory performance. Furthermore, caffeinated soda intake predicted greater alcohol sipping after 12 months even after controlling for rates of baseline alcohol sipping; children who drink caffeinated soda daily are twice as likely to start sipping alcohol after one year compared to those who do not drink caffeinated soda at all. **Conclusions:** These findings suggest that previous linkages between caffeine and substance use in adolescence also extend to younger initiation, and may stem from core neurocognitive features thought conducive to substance initiation.

Keywords: caffeinated soda, alcohol sipping, risk factors of substance use, impulsivity, working memory, ABCD study

Soft drinks are commonly consumed even by children, and a vast majority of

sodas contain caffeine (Temple, 2009, 2018). Moreover, caffeinated sodas typically

contain sweeteners such as high-fructose corn syrup, which can affect neurocognitive

function and cause physical side effects, for an instance, by affecting microbiome

(Ettinger, 2022). Not surprisingly, the consequences of excessive consumption of both

sugar and caffeine have been well documented (Ooi et al., 2022; Porciúncula et al., 2013;

Temple et al., 2017), including a strong association between caffeinated beverage

consumption in adolescence and future substance use (Arria et al., 2011; Barrense-Dias et

al., 2016; Kristjansson et al., 2018; Leal & Jackson, 2019; Marmorstein, 2018; Miyake &

Marmorstein, 2015). In prospective studies tracking the effects of substance use, the

percentage of regular energy drink users who became alcohol or marijuana users after 1–

2 years was approximately five times higher than that of non-energy drink users (Leal &

Jackson, 2019; Marmorstein, 2018; Miyake & Marmorstein, 2015). Others have shown

that coffee or energy drink consumption in adolescents or young adults significantly

predicts future substance use, such as tobacco and alcohol use (Arria et al., 2011;

Barrense-Dias et al., 2016; Kristjansson et al., 2018, 2022; Marmorstein, 2018).

In light of the potential for a problematic progression from drinking caffeinated

beverages to using harder substances, there is a critical need to investigate aspects of this

progression as early as possible. As adolescence is the most common period for initiating

substance use and an earlier onset of substance use predicts greater addiction severity

(Chambers et al., 2003; Jordan & Andersen, 2017), examining these relationships in

preadolescence is crucial in that caffeinated soda intake in childhood could provide useful

predictive information on future substance use. In preadolescent children, caffeinated

soda is the most preferred and accessible mode of caffeine intake (Temple, 2009, 2018).

Only a handful of studies, however, have directly examined the effects of frequent

caffeinated beverage consumption in preadolescent children, due to their lower rates of

consumption (0.4% for coffee and < 0.1% for energy drinks in 9–10-year-old children)

(Lisdahl et al., 2021). Most of the previous studies examining the association between

caffeine consumption and later substance use have focused primarily on adolescents who

drink multiple caffeinated beverages on a daily basis (Temple, 2009, 2018). Moreover,

while a few existing studies have examined behavioral risk factors associated with

caffeinated soda intake, they have not assessed neural underpinnings of such risk factors

(James et al., 2011; Miyake & Marmorstein, 2015; Solnick & Hemenway, 2013; Suglia et

al., 2013).

Here, we addressed the unanswered question of whether frequent consumption of

caffeinated soda in preadolescent children indicates a higher risk of future alcohol

experimentation, using data from the Adolescent Brain Cognitive Development (ABCD)

Study (Bjork et al., 2017). In addition, we wished to explore the potential

neurobehavioral mechanisms of such relationships. In light of previous linkages between

activity in neurocircuits germane to motivation and inhibition (as detected from

functional magnetic resonance imaging (fMRI)) and substance use in adolescents (Lees et

al., 2021), we examined the relationship between daily caffeinated soda intake and both

behavioral and neuroimaging markers of neurobehavioral risk factors for substance use

disorders (SUDs) in children. These include impaired working memory (WM), high

impulsivity, and altered mesolimbic reward processing (**Fig. 1A**). These are the three

primary neurocircuit functions targeted in the ABCD study due to their significant

implications for addiction (see Casey et al. (2018) for more detail). We applied a machine

learning approach (a least absolute shrinkage and selection operator (LASSO) regression;

Tibshirani, 1996) to the measures to identify multivariate risk factors for SUDs associated

with daily caffeinated soda intake and select features highly associated with caffeinated

soda intake (Volkow et al., 2015). Using the same approach, we then examined whether

caffeinated soda intake could predict future alcohol sipping, as (1) alcohol sipping has

been reported to predict future alcohol abuse (Jackson et al., 2015; Watts et al., 2021),

and (2) alcohol sipping is the most common gateway behavior toward other substances of

abuse (Barry et al., 2016). Based on the previous literature suggesting that caffeinated

beverage consumption predicts future substance use (Arria et al., 2011; Barrense-Dias et

al., 2016; Leal & Jackson, 2019; Marmorstein, 2018; Miyake & Marmorstein, 2015), we

hypothesized that daily caffeinated soda intake in the baseline ABCD assessment

(children age 9-10) would predict future alcohol sipping at the year 1 follow up (**Fig. 1B**).

Taken together, we aimed to elucidate the potential neurobehavioral underpinnings of

frequent caffeinated soda intake during childhood.

Methods

Participants

The ABCD study collected data from 11,878 children aged 9–10 years. The

participants were recruited via school systems from 21 different sites in the United States.

The Institutional Review Board (IRB) at the University of California, San Diego,

approved all the research protocol of the ABCD study at most sites per reliance

agreement, with local IRB approval at the remaining sites (Auchter et al., 2018). All

participants provided written assent, and their parents or guardians provided written

consent (Auchter et al., 2018). More information about the recruitment and study design

is available in Garavan et al. (2018). Further details of the demographic, physical, and

mental health assessments are described in Barch et al., (2018).

Out of 11,878 children from ABCD release 2.0, we excluded those with any

missing data regarding either caffeinated soda intake, neurobehavioral risk factors for

SUDs, future alcohol sipping, and confounding variables (see **Measures**). We also

excluded outliers based on the measures of caffeinated soda intake with the cutoff

larger/smaller than the mean ±5 standard deviations. Of these, 147 reported drinking

more than 7 cans of caffeinated soda per week (daily-drinkers) and 1,945 reported not

drinking a single can (non-drinkers) in the past 6 months. Therefore, we included 2,092

participants for the main analyses (see **Fig. S1** for a flowchart of the selection process).

To examine the potential for selection bias, we compared the baseline characteristics

among children who were included and excluded in the analyses (**Table S1**).

Measures

Caffeinated soda intake

Caffeinated soda intake was assessed from self-report of the participants in

response to the question, "How many drinks of the following beverages have you had per

week in the past 6 months? – soda with caffeine (Mountain Dew, Jolt, Coke, Pepsi, Dr.

Pepper, Barq's root beer)". The participants who reported consuming more than 7 cans

per week (i.e., corresponds to more than 1 can per day in average) were allocated to

daily-drinking group (*N*=147), and those who reported consuming 0 can per week were

allocated to the non-drinking group (*N*=1,945). See **Table S2** for a comparison of the

(average) daily-drinkers and non-drinkers on the following behavioral measures, and see

Table S3 for a comparison of the neural measures.

Neurobehavioral risk factors for SUDs

As shown in **Fig. 1A**, we focused on three well-known neurobehavioral risk

factors for SUDs: poor WM, increased impulsivity, and aberrant reward processing.

These constructs were collectively measured by self-report surveys, behavioral tasks, and

fMRI, respectively, as described below. Spatial regions of interest (ROIs) for each task

were restricted *a priori* to the canonical activations initially reported using the ABCD

samples (Casey et al., 2018). The curated data used were based on the Destrieux atlas

(Destrieux et al., 2010).

To obtain behavioral measures of WM, we used the List Sorting Working

Memory Test (list sorting test; Tulsky et al., 2013) and Dimensional Change Card sort

Test (card sort test; Tulsky et al., 2013), as well as response accuracy during the 2-back

condition of the emotional N-back (EN-back) task (Cohen et al., 2015) (see

Supplementary materials). The EN-back task was conducted inside the MRI scanner.

The contrast of "2-back versus 0-back" was used for the fMRI analysis, and the ROIs

were selected based on activation maxima of the contrast in the initial subset of

participants (Casey et al., 2018): the rostral middle frontal gyrus (MFG), caudal MFG,

inferior frontal gyrus (IFG; pars triangularis and pars orbitalis), lateral orbitofrontal

cortex (OFC), superior parietal lobule and inferior parietal lobule (IPL) in the

frontoparietal network, and the caudate nucleus, putamen, nucleus accumbens (NAc),

rostral anterior cingulate cortex (ACC), caudal ACC, thalamus proper, ventral

diencephalon, amygdala, and hippocampus in the fronto-thalamic network.

Due to the multi-faceted construct of impulsivity, wherein laboratory and self-report assessments of impulsivity are thought to capture different components (Sharma et al., 2014), we analyzed each of self-report trait-like impulsivity as well as rapid-response and decision-based impulsivity. Trait impulsivity was measured using the short form Urgency–Premeditation–Perseverance–Sensation Seeking–Positive Urgency (UPPS-P) impulsive behavior scale for children (the 20-item short version for youths; Barch et al., 2018) and a parental report of the ABCD Youth Behavioral Inhibition System/ Behavioral Activation System (BIS/BAS; Carver & White, 1994). The BAS is related to goal-directed efforts, such as motor activation in response to an impending reward, while the BIS is engaged when inhibition toward a goal occurs, such as the avoidance of punishment (Carver & White, 1994). Excessive behavioral inhibition is associated with depression and anxiety, while excessive behavioral activation is associated with impulsive behaviors, compulsive behaviors, substance misuse, and aggression (Newman et al., 2005). To identify the neurocircuit correlates of impulsivity, participants completed the stop signal task (SST; Logan et al., 1984) during fMRI. The SST measures impulsivity related to impulse control or response inhibition when performing an action. The contrast of "correct stop versus correct go" was used for the fMRI analysis. ROIs were the lateral prefrontal cortex (rostral MFG, pars orbitalis in IFG, and lateral OFC), rostral and caudal ACC, and striatum (caudate, putamen, and NAc). These areas have

been associated with impulsivity and impulse control (Aron et al., 2014; Casey et al., 2018). See **Supplementary materials** for more details of the SST.

Reward processing is closely linked to impulsivity, as impulsive people often show immediate reward-seeking behavior (Zuckerman, 2001). For our offline metric of reward sensitivity, we used the Cash Choice Task (Luciana et al., 2018), which assesses willingness to delay gratification (see **Supplementary materials**). Previous studies have suggested that people with SUDs exhibit hyper-responsiveness of mesolimbic motivational neurocircuitry toward drug-related cues (Bechara et al., 2019) while showing blunted responsiveness toward cues for a non-drug reward (e.g., monetary reward) (Luijten et al., 2017). The monetary incentive delay (MID) task (Knutson et al., 2000) is widely used to measure the neural correlates of the anticipation of monetary rewards and losses. The contrast of "reward versus neutral" at the cue onset, which reflects reward anticipation, was used for the fMRI analysis, with the ROIs in the ventral striatum (i.e., NAc), rostral ACC, and medial OFC in the medial prefrontal cortex, which play a key role in reward processing, particularly in reward anticipation (Bartra et al., 2013). See **Supplementary materials** for more details about the MID task.

Future alcohol sipping

Alcohol sipping was measured by self-report from the participants using the iSay Sip Inventory, which was performed once a year and asked only in children who had heard of alcohol (see Lisdahl et al. (2018) for more details). From the year 1 data (data release 3.0), we used a binary response to a single question asking if the participants had

sipped alcohol outside of a religious ceremony. We considered those who had not heard

of alcohol as having yet to sip alcohol (i.e., no alcohol sipping).

Confounding variables

In light of the potential for socioeconomic and structural factors to influence

health behaviors, we included the following variables as confounding variables in the

data analysis: socioeconomic status (SES), family history of substance use, parental

monitoring, sleep deprivation, externalizing/internalizing symptomatology, data

collection sites, and type of MRI scanner. See **Supplementary materials** for more detail.

Analysis

Caffeinated soda intake and risk factors for SUDs

To identify a multivariate profile of risk factors for caffeinated soda

consumption, we performed a binomial LASSO regression analysis (Tibshirani, 1996), a

machine learning algorithm, to distinguish the daily soda-drinking group (N=147) from

the non-drinking group (N=1,945). We used all of the risk factors and confounding

variables as input (i.e., predictors) in the LASSO-based prediction model to classify each

individual into the daily-drinking group (coded as "1") or non-drinking group (coded as

"0"). More specifically, candidate predictors included every measure for the three

cognitive factors as well as other control variables. As LASSO regression offers feature

selection (Tibshirani, 1996), we aimed to identify features that could differentiate the

daily-drinking group from the non-drinking group. See **Supplementary materials** for

more details on LASSO regression.

Caffeinated soda intake and future alcohol sipping

We performed a binomial LASSO regression analysis to test the association between daily caffeinated soda intake and alcohol sipping. We used all possible variables, including the three primary neurobehavioral risk factors (WM, impulsivity, and reward processing) along with the confounding variables as candidate predictors collected at baseline. Using these inputs (i.e., predictors), the LASSO-based model was used to classify individuals who reported alcohol sipping after 12 months (coded as "1"; *N*=90) and those who reported no alcohol sipping (coded as "0"; *N*=2,002). We aimed to select predictors that could distinguish individuals who reported alcohol sipping from those who did not. All of the procedures, except for the predictors and dependent variables, are identical to those in the former analysis (see "**Methods - *Caffeinated soda intake and risk factors for SUDs"***). Furthermore, we performed a chi-square test to compare the ratio of alcohol sipping after 12 months between the two groups. Lastly, we estimated the risk ratio of the daily-drinking group by dividing the cumulative incidence of alcohol sipping in the daily-drinking group by the cumulative incidence in the non-drinking group.

To rigorously examine the association between caffeinated soda intake and future alcohol sipping, we conducted three additional analyses. First, we used continuous measure for caffeinated soda intake instead of using the categorical measure of daily- and non-drinkers. Using data from 4,517 participants (see **Fig. S1**), we carried out the same binomial LASSO regression analysis mentioned earlier (**Fig. S3**). Second, we used only the behavioral measures of risk factors in predicting future alcohol sipping, to include as many participants as possible. Using data from 8,939 participants (see **Fig. S1**), we again

performed the binomial LASSO regression analysis (**Fig. S4**). Lastly, using the sample of

8,939 participants, we ran a hierarchical logistic regression analysis to confirm whether

continuous measures of caffeinated soda intake at baseline can predict alcohol sipping

after 12 months (**Table. S4**). For more details, please refer to the **Supplementary**

materials.

Results

Caffeinated soda intake and neurobehavioral risk factors for SUDs

Our first question was whether daily caffeinated soda intake is cross-sectionally

associated with well-known neurobehavioral risk factors for SUDs. **Fig. 2A** shows the

multivariate profiles from binomial LASSO regression analysis distinguishing the daily-

drinking group from the non-drinking group. Family history of drug use and low parental

education were the two strongest predictors of daily consumption, along with sex (being

male), lack of sleep, low family income, racial identities (as African American), high

body mass index (BMI), high externalizing behaviors, and low parental monitoring. **Fig.**

2B shows a receiver-operation characteristic (ROC) curve and its mean area under the

curve (AUC) for classification of the daily-drinking and non-drinking groups. The mean

AUC values were 0.80 and 0.72 for the training and test sets, respectively.

Among our *a priori* candidate neurobehavioral risk factors for SUDs, high

impulsivity measured from the BAS was most strongly associated with daily caffeinated

soda intake (**Fig. 2C**). A higher UPPS-P score was also related to daily intake.

Additionally, hypoactivation in the caudal ACC during the SST predicted the

classification of the daily-drinking group. Among the WM measures, poor performance

on the list sorting test and 0-back performance in the EN-back Task predicted daily intake

of caffeinated soda. In addition, hypoactivation of the pars orbitalis of the IFG and

greater activation of the IPL by working memory demands during the EN-back task (2-

back vs. 0-back) also predicted daily caffeinated soda intake (**Fig. 2C**). However,

variables related to reward processing were not found to be predictors of daily-drinking

group.

These results suggest that high impulsivity and low WM are significantly

associated with daily caffeinated soda consumption. Along with some other demographic

factors (family history of drug use, male sex, low SES, low parental monitoring, high

externalizing behaviors, less sleep, and high BMI), the two neurobehavioral risk factors

for SUDs distinguished the daily-drinking group from the non-drinking group.

Caffeinated soda intake and future alcohol sipping

To address our second question of whether daily caffeinated soda intake predicts

future alcohol sipping, we conducted binomial LASSO regression predicting alcohol

sipping after 12 months, using all of the neurobehavioral risk factors and confounding

variables collected at baseline. As shown in **Fig. 3A**, daily caffeinated soda intake was

identified as one of the predictors that distinguished alcohol sipping after 12 months

(mean estimate of coefficients=0.122, 95% confidence interval (CI)=[0.010, 0.235]),

even after including alcohol sipping at baseline as a predictor. The mean AUC values of

the predictive model were 0.90 and 0.72 for the training and test sets, respectively (**Fig.**

3B). Moreover, the ratio of alcohol sipping after 12 months was twice as high in the

daily-drinking group compared to the non-drinking group (daily-drinking group: 0.082%

(12 out of 147 children); non-drinking group: 0.040% (78 out of 1,945 children); $X^2 =$

5.802, $p = 0.016$) (**Fig. 3C**). This finding suggests that the daily-drinking group had 2.04

times the risk of sipping alcohol compared to the non-drinking group. Our results were

further supported by the binomial LASSO regression analyses with larger samples (**Fig.**

S4 & Fig. S5) and hierarchical logistic regression analysis (**Table S4**) which can be

found in the **Supplementary materials**.

Discussion

In this study, we investigated whether caffeinated soda intake in preadolescent

childhood is associated with a higher risk of alcohol experimentation in the future. Using

the large dataset from the ABCD study, we first clarified cross-sectional relationships

between daily caffeinated soda intake and well-known neurobehavioral risk factors for

SUDs at study baseline; then evaluated whether caffeinated soda intake at baseline was

predictive of alcohol sipping after 12 months. Our findings suggest that frequent

consumption of caffeinated soda in children is closely related to previously-established

neurobehavioral risk factors for SUDs and can predict future alcohol sipping.

Our machine learning approach partially supported our hypothesized association

between caffeinated soda intake and the neurobehavioral risk factors for SUDs. Notably,

high impulsivity scores and low WM performance each distinguished daily caffeinated

soda drinkers from non-drinkers even after considering the effects of other confounding

factors (**Fig. 2**). However, there was no significant association observed for mesolimbic

recruitment by prospective rewards.

In relation to impulsivity, we found higher self-reported impulsivity in the daily-drinking group based on the UPPS-P and BAS scores along with altered activation in the ACC, a brain region implicated in cognitive control and impulsivity (Kerns et al., 2004; Shenhav et al., 2016). Reduced activities in the ACC during response inhibition or behavior-monitoring are commonly reported in children with attention deficit hyperactivity disorder (ADHD) (Cortese et al., 2012; Hart et al., 2013; McTeague et al., 2017) and individuals with SUDs (Luijten et al., 2014; Nestor et al., 2011; Yücel et al., 2007). Thus, hypoactivation of the ACC during response inhibition in daily-drinking group further seems to strengthen the association between daily soda consumption and elevated levels of impulsivity.

We also found WM impairments in the daily-drinking group on the list sorting test and the 0-back accuracy in the EN-back task, accompanied by hypoactivation in the IFG and hyperactivation in the IPL by working memory demands during the EN-back task. Prior studies have shown that greater activation in the prefrontal cortex is related to greater WM capacity (Casey et al., 2018; Owen et al., 2005), and increased activation in the IPL is associated with higher WM load (Baldo & Dronkers, 2006; Veltman et al., 2003). Taken together, these findings strongly suggest an association between daily-soda consumption and WM deficits.

In contrast, we did not find a significant association between neurocognitive measures of reward processing and daily caffeinated soda intake. Aberrant reward processing is a commonly observed neurocognitive feature in addiction (Zeng et al., 2023), with individuals either displaying increased or decreased sensitivity to reward (Berridge & Robinson, 2016; Blum et al., 2000; Demidenko et al., 2020; Robinson &

Berridge, 1993). Altered reward processing has been also reported in children with a high risk of alcohol use, such as those with a family history of alcohol use problems (Bjork et al., 2008; Martz et al., 2022). One possibility for the lack of association is that the effect of reward processing variables on SUD may depend on population or context (Bjork, 2020), or may have been masked by the strong effect of family history of drug use, since our study controlled for the familial risk of alcohol and drug use. Additionally, it is worth considering that monetary rewards might have differential effects compared to drug rewards (Nestor & Ersche, 2023), which could explain the absence of significant associations. Consistent with our findings, a recent study comparing reward processing of alcohol dependent patients, first-degree relatives, and healthy controls could not find any significant group differences both in monetary reward and loss anticipation when controlling for age (Musial et al., 2023). Thus, further research is required to reconcile the mix findings related to reward processing in youth with a risk of addiction, perhaps by studying more narrowly defined subgroups and by examining both drug and non-drug rewards.

After examining the link between the neurobehavioral risk factors and caffeinated soda intake, we showed that frequent consumption of caffeinated soda predicted alcohol sipping after 12 months using LASSO regression (**Fig. 3**). Even after controlling for other well-established variables for alcohol sipping, including baseline alcohol sipping, caffeinated soda intake remained predictive of future alcohol sipping (**Fig. 3A**). Other survived predictors of future sipping include predictors for daily caffeinated soda intake itself (**Fig. 2A**), such as higher behavioral impulsivity score (i.e., UPPS-P), BMI, and hypoactivation of the IFG during the EN-back task. Conversely,

reduced activity in the NAc and greater activity in the amygdala during the EN-back task, and reduced activity of the medial OFC during the MID task were associated with the participants who experienced alcohol sipping, but did not survive as predictors for daily caffeinated soda intake. Higher family income was associated with the participants who experienced alcohol sipping, while it was associated with the participants who do not drink caffeinated soda. Interestingly, some predictors of caffeinated soda intake were similar to the risk factors for SUDs (Jordan & Andersen, 2017), suggesting that caffeinated soda intake during childhood and SUDs share similar neurobehavioral vulnerabilities. Note that the majority of the cohort was substance-naïve (Lisdahl et al., 2018), and we evaluated alcohol sipping measures instead of substantial self-administration of alcohol. Thus, it would be informative to track the predictive ability of caffeinated soda and alcohol use and investigate changes in the direction of the predictors as the children get older.

Our findings suggest that caffeinated soda in children is predictive of substance use in the near future. The longitudinal associations between the use of more benign psychotropic substances early in life and the use of "harder" substances later in adolescence or emerging adulthood have been characterized for decades (Kandel, 1975), and have been attributed to two competing (but not mutually-exclusive) theories. The "gateway hypothesis" (Kandel & Yamaguchi, 2002) generally implies that exposure to the earlier-used substance itself, such as nicotine or cannabis, induces a toxicological effect on brain which renders the individual more sensitive to reinforcing effects of harder drugs. Evidence for this theory is supported by controlled animal model intervention studies (e.g., (Collins & Izenwasser, 2004)). Animal studies on the effects of

caffeine intake on increasing later alcohol consumption (Hou et al., 2016; Kunin et al., 2000; SanMiguel et al., 2019) supports that the gateway hypothesis could also be applied to the association between caffeine and alcohol. Thus, one possible explanation of higher alcohol sipping rate of the daily-drinking group in our study is that the substances contained in caffeinated soda (caffeine and sugar) may have induced neurophysiological effects and reinforced regular soda drinkers to try alcohol after 12 months.

Conversely, the "common liability" hypothesis posits that the progression from softer to harder substances results primarily from the intersection of a genetically-regulated under-controlled temperament with progressively expanding *access* to harder substances with advancement into young adulthood (Vanyukov et al., 2012). Cross-sectional observations supported the common liability hypothesis, as children and adolescents with disruptive behavior disorders such as conduct disorder, which is strongly linked to SUDs, showed deficits in executive function (e.g., impaired impulse control) (Matthys et al., 2013). Longitudinal studies also support the common liability hypothesis (Debenham et al., 2021), wherein impaired impulse control and high sensation-seeking in young adolescents are predictive of increased drinking over time (Castellanos-Ryan et al., 2011). Similarly, caffeinated soda intake itself may not directly cause children to initiate substance use, but the drinking behavior of the beverages could indicate high impulsivity, which may be linked to a high risk of initiating substance use in the near future, consistent with the common liability hypothesis (Vanyukov et al., 2012). Because caffeinated soda contains two addictive substances, sugar and caffeine, it is somewhat natural for children to prefer and seek the beverages (Temple, 2009). While this taste preference could lead to a seeking behavior and habitual consumption, previous

studies have shown impulsivity as one of the most common traits of SUDs (Cyders et al.,
2009; Magid & Colder, 2007). Therefore, the behavior of frequently consuming
caffeinated soda could indicate a high risk of initiating substance use in the future, due to
the common risk factors (e.g., high impulsivity) between the two behaviors, although
further research is needed to disentangle this complex relationship.

To our knowledge, this is the first study to investigate the direct link between
caffeinated soda intake in earlier childhood and the risk of substance use. The results are
consistent with studies on caffeine consumption during adolescence and its association
with future substance misuse (Barrense-Dias et al., 2016; Leal & Jackson, 2019),
supporting a higher risk of caffeinated soda consumption in childhood, particularly
regarding vulnerability to future substance misuse. Such information is invaluable, as
caffeinated soda is incomparably the most common medium for caffeine consumption in
childhood, and the risk of substance misuse should be detected before adolescence, the
most common period of substance use onset.

Our results have important implications for public health recommendations, as
our study provides novel insight into the neurobehavioral correlates of caffeinated soda
consumption in children, which has rarely been evaluated. At the same time, a few
limitations of our research and future directions for further investigations should be
discussed. First, we did not perform functional connectivity analyses or multivoxel
pattern analyses, which may have provided additional insight into the effects of
caffeinated soda intake. Second, as the 9–10-year-old children in this study had not yet
started other substances except for alcohol, such as tobacco or marijuana, future work
using the longitudinal 10-year follow-up data of the ABCD study should examine

whether frequent consumption of caffeinated soda is associated with alcohol or other

substance misuse. Third, we acknowledge that multiple variables other than caffeinated

soda intake may mediate the relationship between neurobehavioral risk factors and future

alcohol use; thus, extensive investigation of how caffeinated soda intake interacts with

other SUD risk/protective factors is needed in the future. In addition, the effects of acute

caffeine intake could have influenced the task performance of the soda drinking groups

(Graczyk et al., 2018). Thus, future studies investigating the longitudinal effects of

caffeinated soda intake should control for the acute caffeine consumption of children.

Lastly, the ABCD dataset included only a small set of measures of food/drink

consumption. We relied on a self-report measure consisting of a single item to assess

caffeinated soda intake. For a more comprehensive understanding of the risks associated

with caffeinated soda intake, it is critical for future research to integrate more objective

and detailed assessments of soda consumption. Moreover, it is essential to differentiate

the effects of caffeine and sweeteners by including more diverse measures of caffeine or

sugar consumption.

In conclusion, our results revealed the potential risks of caffeinated soda

consumption in children by investigating the associations between caffeinated soda

consumption and risk factors for SUDs and examining the ability of caffeinated soda

consumption to predict future alcohol sipping, using the large ABCD dataset. While

previous research on the side effects of caffeinated soda consumption has been limited to

negative physical consequences, the present results strongly suggest that caffeinated soda

drinking in children is also associated with altered neurobehavioral function and can

predict alcohol sipping after 12 months. Our study further suggests a strong need to

develop evidence-based recommendations for caffeinated soda consumption in minors

(Temple, 2018), as there is no consensus on a safe dose of caffeinated soda in children,

and some children are at higher risk of adverse events from frequent caffeinated soda

intake. Further clarification on the causal relationships and neuro-developmental

evidence are needed to determine whether caffeinated soda is a warning sign for future

substance misuse and whether it induces neurobehavioral impairments in children.

Acknowledgements

We thank the families and children who participated, and continue to participate, in the

ABCD study, as well as staff at the study sites who are involved in data collection and

curating the data release. Data used in the preparation of this article were obtained from

the Adolescent Brain Cognitive Development (ABCD) Study (https://abcdstudy.org),

held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed

to recruit more than 10,000 children age 9-10 and follow them over 10 years into early

adulthood. The ABCD Study® is supported by the National Institutes of Health and

additional federal partners under award numbers U01DA041048, U01DA050989,

U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987,

U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134,

U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120,

U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123,

U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-

partners.html. A listing of participating sites and a complete listing of the study

investigators can be found at https://abcdstudy.org/consortium_members/. ABCD

consortium investigators designed and implemented the study and/or provided data but

did not necessarily participate in analysis or writing of this report. This manuscript

reflects the views of the authors and may not reflect the opinions or views of the NIH or

ABCD consortium investigators. The ABCD data repository grows and changes over

time. The ABCD data used in this report came from [NIMH Data Archive Digital Object

Identifier (10.15154/1503209)]. DOIs can be found at

http://dx.doi.org/10.15154/1503209.

Declaration of interest

The authors declare no competing interests.

References

Aron, A. R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior frontal cortex: one decade on. *Trends in Cognitive Sciences*, *18*(4), 177–185. https://doi.org/10.1016/j.tics.2013.12.003

Arria, A. M., Caldeira, K. M., Kasperski, S. J., Vincent, K. B., Griffiths, R. R., & O'Grady, K. E. (2011). Energy Drink Consumption and Increased Risk for Alcohol Dependence. *Alcoholism: Clinical and Experimental Research*, *35*(2), 365–375. https://doi.org/10.1111/j.1530-0277.2010.01352.x

Auchter, A. M., Mejia, M. H., Heyser, C. J., Shilling, P. D., Jernigan, T. L., Brown, S. A., Tapert, S. F., & Dowling, G. J. (2018). A Description of the ABCD Organizational Structure and Communication Framework. *Developmental Cognitive Neuroscience*, *32*, 8–15. https://doi.org/10.1016/j.dcn.2018.04.003

Baldo, J. V., & Dronkers, N. F. (2006). The Role of Inferior Parietal and Inferior Frontal Cortex in Working Memory. *Neuropsychology*, *20*(5), 529–538. https://doi.org/10.1037/0894-4105.20.5.529

Barch, D. M., Albaugh, M. D., Avenevoli, S., Chang, L., Clark, D. B., Glantz, M. D., Hudziak, J. J., Jernigan, T. L., Tapert, S. F., Yurgelun-Todd, D., Alia-Klein, N., Potter, A. S., Paulus, M. P., Prouty, D., Zucker, R. A., & Sher, K. J. (2018). Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Developmental Cognitive Neuroscience*, *32*(Biol. Psychol. 124 2017), 55–66. https://doi.org/10.1016/j.dcn.2017.10.010

Barrense-Dias, Y., Berchtold, A., Akre, C., & Surís, J. (2016). Consuming energy drinks at the age of 14 predicted legal and illegal substance use at 16. *Acta Paediatrica*, *105*(11), 1361–1368. https://doi.org/10.1111/apa.13543

Barry, A. E., King, J., Sears, C., Harville, C., Bondoc, I., & Joseph, K. (2016). Prioritizing Alcohol Prevention: Establishing Alcohol as the Gateway Drug and Linking Age of First Drink With Illicit Drug Use. *Journal of School Health*, *86*(1), 31–38. https://doi.org/10.1111/josh.12351

Bartra, O., McGuire, J. T., & Kable, J. W. (2013). The valuation system: A coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *NeuroImage*, *76*, 412–427. https://doi.org/10.1016/j.neuroimage.2013.02.063

Bechara, A., Berridge, K. C., Bickel, W. K., Morón, J. A., Williams, S. B., & Stein, J. S. (2019). A Neurobehavioral Approach to Addiction: Implications for the Opioid Epidemic and the Psychology of Addiction. *Psychological Science in the Public Interest*, *20*(2), 96–127. https://doi.org/10.1177/1529100619860513

Berridge, K. C., & Robinson, T. E. (2016). Liking, Wanting, and the Incentive-Sensitization Theory of Addiction. *American Psychologist*, *71*(8), 670–679. https://doi.org/10.1037/amp0000059

Bjork, J. M. (2020). The Ups and Downs of Relating Nondrug Reward Activation to Substance Use Risk in Adolescents. *Current Addiction Reports*, *7*(3), 421–429. https://doi.org/10.1007/s40429-020-00327-7

Bjork, J. M., Knutson, B., & Hommer, D. W. (2008). Incentive-elicited striatal activation in adolescent children of alcoholics. *Addiction*, *103*(8), 1308–1319. https://doi.org/10.1111/j.1360-0443.2008.02250.x

Bjork, J. M., Straub, L. K., Provost, R. G., & Neale, M. C. (2017). The ABCD Study of Neurodevelopment: Identifying Neurocircuit Targets for Prevention and Treatment of Adolescent Substance Abuse. *Current Treatment Options in Psychiatry*, *4*(2), 196–209. https://doi.org/10.1007/s40501-017-0108-y

Blum, K., Braverman, E. R., Holder, J. M., Lubar, J. F., Monastra, V. J., Miller, D., Lubar, J. O., Chen, T. J. H., & Comings, D. E. (2000). The Reward Deficiency Syndrome: A Biogenetic Model for the Diagnosis and Treatment of Impulsive, Addictive and Compulsive Behaviors. *Journal of Psychoactive Drugs*, *32*(sup1), 1–112. https://doi.org/10.1080/02791072.2000.10736099

Carver, C. S., & White, T. L. (1994). Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. *Journal of Personality and Social Psychology*, *67*(2), 319. https://doi.org/10.1037/0022-3514.67.2.319

Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M., Soules, M. E., Teslovich, T., Dellarco, D. V., Garavan, H., Orr, C. A., Wager, T. D., Banich, M. T., Speer, N. K., Sutherland, M. T., Riedel, M. C., Dick, A. S., Bjork, J. M., Thomas, K. M., … Workgroup, the A. I. A. (2018). The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Developmental Cognitive Neuroscience*, *32*(Biol. Psychiatry 69 2011), 43–54. https://doi.org/10.1016/j.dcn.2018.03.001

Castellanos-Ryan, N., Rubia, K., & Conrod, P. J. (2011). Response Inhibition and Reward Response Bias Mediate the Predictive Relationships Between Impulsivity and Sensation Seeking and Common and Unique Variance in Conduct Disorder and Substance Misuse. *Alcoholism: Clinical and Experimental Research*, *35*(1), 140–155. https://doi.org/10.1111/j.1530-0277.2010.01331.x

Chambers, R. A., Taylor, J. R., & Potenza, M. N. (2003). Developmental Neurocircuitry
of Motivation in Adolescence: A Critical Period of Addiction Vulnerability. *American*
Journal of Psychiatry, *160*(6), 1041–1052. https://doi.org/10.1176/appi.ajp.160.6.1041

Collins, S. L., & Izenwasser, S. (2004). Chronic nicotine differentially alters cocaine-
induced locomotor activity in adolescent vs. adult male and female rats.
Neuropharmacology, *46*(3), 349–362.
https://doi.org/10.1016/j.neuropharm.2003.09.024

Cortese, S., Kelly, C., Chabernaud, C., Proal, E., Martino, A. D., Milham, M. P., &
Castellanos, F. X. (2012). Toward Systems Neuroscience of ADHD: A Meta-Analysis
of 55 fMRI Studies. *American Journal of Psychiatry*, *169*(10), 1038–1055.
https://doi.org/10.1176/appi.ajp.2012.11101521

Cyders, M. A., Flory, K., Rainer, S., & Smith, G. T. (2009). The role of personality
dispositions to risky behavior in predicting first-year college drinking. *Addiction*,
104(2), 193–202. https://doi.org/10.1111/j.1360-0443.2008.02434.x

Debenham, J., Birrell, L., Champion, K., Lees, B., Yücel, M., & Newton, N. (2021).
Neuropsychological and neurophysiological predictors and consequences of cannabis
and illicit substance use during neurodevelopment: a systematic review of longitudinal
studies. *The Lancet Child & Adolescent Health*, *5*(8), 589–604.
https://doi.org/10.1016/s2352-4642(21)00051-1

Demidenko, M. I., Huntley, E. D., Jahn, A., Thomason, M. E., Monk, C. S., & Keating,
D. P. (2020). Cortical and subcortical response to the anticipation of reward in high
and average/low risk-taking adolescents. *Developmental Cognitive Neuroscience*, *44*,
100798. https://doi.org/10.1016/j.dcn.2020.100798

Destrieux, C., Fischl, B., Dale, A., & Halgren, E. (2010). Automatic parcellation of
human cortical gyri and sulci using standard anatomical nomenclature. *NeuroImage*,
53(1), 1–15. https://doi.org/10.1016/j.neuroimage.2010.06.010

Ettinger, S. (2022). Diet, Gut Microbiome, and Cognitive Decline. *Current Nutrition*
Reports, 1–10. https://doi.org/10.1007/s13668-022-00435-y

Garavan, H., Bartsch, H., Conway, K., Decastro, A., Goldstein, R., Heeringa, S.,
Jernigan, T., Potter, A., Thompson, W., & Zahs, D. (2018). Recruiting the ABCD
Sample: Design Considerations and Procedures. *Developmental Cognitive*
Neuroscience, *32*(Journal of Studies on Alcohol and Drugs 76 2015), 16–22.
https://doi.org/10.1016/j.dcn.2018.04.004

Graczyk, A. M., Ziegler, A. M., Bendlin, A., Sion, T., Vattana, K., & Temple, J. L.
(2018). Effects of Caffeine Administration on Reaction Time, Attention, and
Inhibitory Control in Children and Adolescents. *Journal of Cognitive Enhancement*,
2(3), 276–286. https://doi.org/10.1007/s41465-018-0074-3

Hart, H., Radua, J., Nakao, T., Mataix-Cols, D., & Rubia, K. (2013). Meta-analysis of Functional Magnetic Resonance Imaging Studies of Inhibition and Attention in Attention-deficit/Hyperactivity Disorder: Exploring Task-Specific, Stimulant Medication, and Age Effects. *JAMA Psychiatry*, *70*(2), 185–198. https://doi.org/10.1001/jamapsychiatry.2013.277

Hou, F., Lesmes, L. A., Kim, W., Gu, H., Pitt, M. A., Myung, J. I., & Lu, Z.-L. (2016). Evaluating the performance of the quick CSF method in detecting contrast sensitivity function changes. *Journal of Vision*, *16*(6), 18. https://doi.org/10.1167/16.6.18

Jackson, K. M., Barnett, N. P., Colby, S. M., & Rogers, M. L. (2015). The Prospective Association Between Sipping Alcohol by the Sixth Grade and Later Substance Use. *Journal of Studies on Alcohol and Drugs*, *76*(2), 212–221. https://doi.org/10.15288/jsad.2015.76.212

James, J. E., Kristjánsson, Á. L., & Sigfúsdóttir, I. D. (2011). Adolescent substance use, sleep, and academic achievement: Evidence of harm due to caffeine. *Journal of Adolescence*, *34*(4), 665–673. https://doi.org/10.1016/j.adolescence.2010.09.006

Jordan, C. J., & Andersen, S. L. (2017). Sensitive periods of substance abuse: Early risk for the transition to dependence. *Developmental Cognitive Neuroscience*, *25*, 29–44. https://doi.org/10.1016/j.dcn.2016.10.004

Kandel, D. (1975). Stages in adolescent involvement in drug use. *Science*, *190*(4217), 912–914. https://doi.org/10.1126/science.1188374

Kandel, D. B., & Yamaguchi, K. (2002). *Stages and Pathways of Drug Involvement*. 65–89. https://doi.org/10.1017/cbo9780511499777.006

Kerns, J. G., Cohen, J. D., III, A. W. M., Cho, R. Y., Stenger, V. A., & Carter, C. S. (2004). Anterior Cingulate Conflict Monitoring and Adjustments in Control. *Science*, *303*(5660), 1023–1026. https://doi.org/10.1126/science.1089910

Knutson, B., Westdorp, A., Kaiser, E., & Hommer, D. (2000). FMRI Visualization of Brain Activity during a Monetary Incentive Delay Task. *NeuroImage*, *12*(1), 20–27. https://doi.org/10.1006/nimg.2000.0593

Kristjansson, A. L., Kogan, S. M., Mann, M. J., Smith, M. L., Juliano, L. M., Lilly, C. L., & James, J. E. (2018). Does early exposure to caffeine promote smoking and alcohol use behavior? A prospective analysis of middle school students. *Addiction*, *113*(9), 1706–1713. https://doi.org/10.1111/add.14261

Kristjansson, A. L., Mann, M. J., Smith, M. L., Kogan, S. M., Lilly, C. L., & James, J. E. (2022). Caffeine consumption and onset of alcohol use among early adolescents. *Preventive Medicine*, *163*, 107208. https://doi.org/10.1016/j.ypmed.2022.107208

Kunin, D., Gaskin, S., Rogan, F., Smith, B. R., & Amit, Z. (2000). Caffeine promotes
ethanol drinking in rats Examination using a limited-access free choice paradigm.
Alcohol, *21*(3), 271–277. https://doi.org/10.1016/s0741-8329(00)00101-4

Leal, W. E., & Jackson, D. B. (2019). The role of energy drink consumption in the
intention to initiate marijuana use among adolescents. *Addictive Behaviors*,
93(Postgraduate Medicine 127 3 2015), 240–245.
https://doi.org/10.1016/j.addbeh.2019.02.008

Lees, B., Garcia, A. M., Debenham, J., Kirkland, A. E., Bryant, B. E., Mewton, L., &
Squeglia, L. M. (2021). Promising vulnerability markers of substance use and misuse:
A review of human neurobehavioral studies. *Neuropharmacology*, *187*, 108500.
https://doi.org/10.1016/j.neuropharm.2021.108500

Lisdahl, K. M., Sher, K. J., Conway, K. P., Gonzalez, R., Ewing, S. W. F., Nixon, S. J.,
Tapert, S., Bartsch, H., Goldstein, R. Z., & Heitzeg, M. (2018). Adolescent brain
cognitive development (ABCD) study: Overview of substance use assessment
methods. *Developmental Cognitive Neuroscience*, *32*, 80–96.
https://doi.org/10.1016/j.dcn.2018.02.007

Lisdahl, K. M., Tapert, S., Sher, K. J., Gonzalez, R., Nixon, S. J., Ewing, S. W. F.,
Conway, K. P., Wallace, A., Sullivan, R., Hatcher, K., Kaiver, C., Thompson, W.,
Reuter, C., Bartsch, H., Wade, N. E., Jacobus, J., Albaugh, M. D., Allgaier, N.,
Anokhin, A. P., … Consortium, A. (2021). Substance use patterns in 9-10 year olds:
Baseline findings from the adolescent brain cognitive development (ABCD) study.
Drug and Alcohol Dependence, 108946.
https://doi.org/10.1016/j.drugalcdep.2021.108946

Logan, G. D., Cowan, W. B., & Davis, K. A. (1984). On the ability to inhibit simple and
choice reaction time responses: A model and a method. *Journal of Experimental
Psychology: Human Perception and Performance*, *10*(2), 276–291.
https://doi.org/10.1037/0096-1523.10.2.276

Luciana, M., Bjork, J. M., Nagel, B., Barch, D. M., Gonzalez, R., Nixon, S. J., & Banich,
M. T. (2018). Adolescent Neurocognitive Development and Impacts of Substance
Use: Overview of the Adolescent Brain Cognitive Development (ABCD) Baseline
Neurocognition Battery. *Developmental Cognitive Neuroscience*, *32*, 67–79.
https://doi.org/10.1016/j.dcn.2018.02.006

Luijten, M., Machielsen, M. W. J., Veltman, D. J., Hester, R., Haan, L. de, & Franken, I.
H. A. (2014). Systematic review of ERP and fMRI studies investigating inhibitory
control and error processing in people with substance dependence and behavioural
addictions. *Journal of Psychiatry and Neuroscience*, *39*(3), 149–169.
https://doi.org/10.1503/jpn.130052

Luijten, M., Schellekens, A. F., Kühn, S., Machielse, M. W. J., & Sescousse, G. (2017). Disruption of Reward Processing in Addiction : An Image-Based Meta-analysis of Functional Magnetic Resonance Imaging Studies. *JAMA Psychiatry*, *74*(4), 387. https://doi.org/10.1001/jamapsychiatry.2016.3084

Magid, V., & Colder, C. R. (2007). The UPPS Impulsive Behavior Scale: Factor structure and associations with college drinking. *Personality and Individual Differences*, *43*(7), 1927–1937. https://doi.org/10.1016/j.paid.2007.06.013

Marmorstein, N. R. (2018). Investigating associations between caffeinated beverage consumption and later alcohol consumption among early adolescents. *Addictive Behaviors*, *90*(Psychopharmacology 203 2009), 362–368. https://doi.org/10.1016/j.addbeh.2018.11.033

Martz, M. E., Hardee, J. E., Cope, L. M., McCurry, K. L., Soules, M., Zucker, R. A., & Heitzeg, M. M. (2022). Nucleus Accumbens Response to Reward among Children with a Family History of Alcohol Use Problems: Convergent Findings from the ABCD Study® and Michigan Longitudinal Study. *Brain Sciences*, *12*(7), 913. https://doi.org/10.3390/brainsci12070913

Matthys, W., Vanderschuren, L. J. M. J., & Schutter, D. J. L. G. (2013). The neurobiology of oppositional defiant disorder and conduct disorder: Altered functioning in three mental domains. *Development and Psychopathology*, *25*(1), 193–207. https://doi.org/10.1017/s0954579412000272

McTeague, L. M., Huemer, J., Carreon, D. M., Jiang, Y., Eickhoff, S. B., & Etkin, A. (2017). Identification of Common Neural Circuit Disruptions in Cognitive Control Across Psychiatric Disorders. *American Journal of Psychiatry*, *174*(7), 676–685. https://doi.org/10.1176/appi.ajp.2017.16040400

Miyake, E. R., & Marmorstein, N. R. (2015). Energy drink consumption and later alcohol use among early adolescents. *Addictive Behaviors*, *43*, 60–65. https://doi.org/10.1016/j.addbeh.2014.12.009

Musial, M. P. M., Beck, A., Rosenthal, A., Charlet, K., Bach, P., Kiefer, F., Vollstädt-Klein, S., Walter, H., Heinz, A., & Rothkirch, M. (2023). Reward Processing in Alcohol-Dependent Patients and First-Degree Relatives: Functional Brain Activity During Anticipation of Monetary Gains and Losses. *Biological Psychiatry*, *93*(6), 546–557. https://doi.org/10.1016/j.biopsych.2022.05.024

Nestor, L. J., & Ersche, K. D. (2023). Abnormal Brain Networks Related to Drug and Nondrug Reward Anticipation and Outcome Processing in Stimulant Use Disorder: A Functional Connectomics Approach. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *8*(5), 560–571. https://doi.org/10.1016/j.bpsc.2022.08.014

Nestor, L. J., Ghahremani, D. G., Monterosso, J., & London, E. D. (2011). Prefrontal
hypoactivation during cognitive control in early abstinent methamphetamine-
dependent subjects. *Psychiatry Research: Neuroimaging*, *194*(3), 287–295.
https://doi.org/10.1016/j.pscychresns.2011.04.010

Newman, J. P., MacCoon, D. G., Vaughn, L. J., & Sadeh, N. (2005). Validating a
Distinction Between Primary and Secondary Psychopathy With Measures of Gray's
BIS and BAS Constructs. *Journal of Abnormal Psychology*, *114*(2), 319–323.
https://doi.org/10.1037/0021-843x.114.2.319

Ooi, J. Y., Wolfenden, L., Sutherland, R., Nathan, N., Oldmeadow, C., Mclaughlin, M.,
Barnes, C., Hall, A., Vanderlee, L., & Yoong, S. L. (2022). A Systematic Review of
the Recent Consumption Levels of Sugar-Sweetened Beverages in Children and
Adolescents From the World Health Organization Regions With High Dietary–
Related Burden of Disease. *Asia Pacific Journal of Public Health*, *34*(1), 11–24.
https://doi.org/10.1177/10105395211014642

Owen, A. M., McMillan, K. M., Laird, A. R., & Bullmore, E. (2005). N-back working
memory paradigm: A meta-analysis of normative functional neuroimaging studies.
Human Brain Mapping, *25*(1), 46–59. https://doi.org/10.1002/hbm.20131

Porciúncula, L. O., Sallaberry, C., Mioranzza, S., Botton, P. H. S., & Rosemberg, D. B.
(2013). The Janus face of caffeine. *Neurochemistry International*, *63*(6), 594–609.
https://doi.org/10.1016/j.neuint.2013.09.009

Robinson, T. E., & Berridge, K. C. (1993). The neural basis of drug craving: An
incentive-sensitization theory of addiction. *Brain Research Reviews*, *18*(3), 247–291.
https://doi.org/10.1016/0165-0173(93)90013-p

SanMiguel, N., López-Cruz, L., Müller, C. E., Salamone, J. D., & Correa, M. (2019).
Caffeine modulates voluntary alcohol intake in mice depending on the access
conditions: Involvement of adenosine receptors and the role of individual differences.
Pharmacology Biochemistry and Behavior, *186*, 172789.
https://doi.org/10.1016/j.pbb.2019.172789

Sharma, L., Markon, K. E., & Clark, L. A. (2014). Toward a Theory of Distinct Types of
"Impulsive" Behaviors: A Meta-Analysis of Self-Report and Behavioral Measures.
Psychological Bulletin, *140*(2), 374–408. https://doi.org/10.1037/a0034418

Shenhav, A., Cohen, J. D., & Botvinick, M. M. (2016). Dorsal anterior cingulate cortex
and the value of control. *Nature Neuroscience*, *19*(10), 1286–1291.
https://doi.org/10.1038/nn.4384

Solnick, S. J., & Hemenway, D. (2013). Soft drinks, aggression and suicidal behaviour in
US high school students. *International Journal of Injury Control and Safety*
Promotion, *21*(3), 266–273. https://doi.org/10.1080/17457300.2013.815631

Suglia, S. F., Solnick, S., & Hemenway, D. (2013). Soft Drinks Consumption Is Associated with Behavior Problems in 5-Year-Olds. *The Journal of Pediatrics*, *163*(5), 1323–1328. https://doi.org/10.1016/j.jpeds.2013.06.023

Temple, J. L. (2009). Caffeine use in children: What we know, what we have left to learn, and why we should worry. *Neuroscience & Biobehavioral Reviews*, *33*(6), 793–806. https://doi.org/10.1016/j.neubiorev.2009.01.001

Temple, J. L. (2018). Trends, Safety, and Recommendations For Caffeine Use in Children and Adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*, *58*(1), 36–45. https://doi.org/10.1016/j.jaac.2018.06.030

Temple, J. L., Bernard, C., Lipshultz, S. E., Czachor, J. D., Westphal, J. A., & Mestre, M. A. (2017). The Safety of Ingested Caffeine: A Comprehensive Review. *Frontiers in Psychiatry*, *8*, 80. https://doi.org/10.3389/fpsyt.2017.00080

Tibshirani, R. (1996). Regression Shrinkage and Selection Via the Lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, *58*(1), 267–288. https://doi.org/10.1111/j.2517-6161.1996.tb02080.x

Tulsky, D. S., Carlozzi, N. E., Chevalier, N., Espy, K. A., Beaumont, J. L., & Mungas, D. (2013). V. NIH Toolbox Cognition Battery (CB): measuring working memory. *Monographs of the Society for Research in Child Development*, *78*(4), 70–87. https://doi.org/10.1111/mono.12035

Vanyukov, M. M., Tarter, R. E., Kirillova, G. P., Kirisci, L., Reynolds, M. D., Kreek, M. J., Conway, K. P., Maher, B. S., Iacono, W. G., Bierut, L., Neale, M. C., Clark, D. B., & Ridenour, T. A. (2012). Common liability to addiction and "gateway hypothesis": Theoretical, empirical and evolutionary perspective. *Drug and Alcohol Dependence*, *123*, S3–S17. https://doi.org/10.1016/j.drugalcdep.2011.12.018

Veltman, D. J., Rombouts, S. A. R. B., & Dolan, R. J. (2003). Maintenance versus manipulation in verbal working memory revisited: an fMRI study. *NeuroImage*, *18*(2), 247–256. https://doi.org/10.1016/s1053-8119(02)00049-6

Volkow, N. D., Koob, G., & Baler, R. (2015). Biomarkers in Substance Use Disorders. *ACS Chemical Neuroscience*, *6*(4), 522–525. https://doi.org/10.1021/acschemneuro.5b00067

Watts, A. L., Wood, P. K., Jackson, K. M., Lisdahl, K. M., Heitzeg, M. M., Gonzalez, R., Tapert, S. F., Barch, D. M., & Sher, K. J. (2021). Incipient alcohol use in childhood: Early alcohol sipping and its relations with psychopathology and personality. *Development and Psychopathology*, *33*(4), 1338–1350. https://doi.org/10.1017/s0954579420000541

Yücel, M., Lubman, D. I., Harrison, B. J., Fornito, A., Allen, N. B., Wellard, R. M., Roffel, K., Clarke, K., Wood, S. J., Forman, S. D., & Pantelis, C. (2007). A combined spectroscopic and functional MRI investigation of the dorsal anterior cingulate region in opiate addiction. *Molecular Psychiatry*, *12*(7), 691–702. https://doi.org/10.1038/sj.mp.4001955

Zeng, J., You, L., Yang, F., Luo, Y., Yu, S., Yan, J., Liu, M., & Yang, X. (2023). A meta-analysis of the neural substrates of monetary reward anticipation and outcome in alcohol use disorder. *Human Brain Mapping*, *44*(7), 2841–2861. https://doi.org/10.1002/hbm.26249

Zuckerman, M. (2001). International Encyclopedia of the Social & Behavioral Sciences. *Developmental Psychology: Personality Development: Motivational/Emotional Psychology: Overview, Classification, and Methods: Personality Psychology and Self: Personality: Risk: Article Titles: S*, 13905–13908. https://doi.org/10.1016/b0-08-043076-7/01772-1

Figure 1. An overview of the analytical approach

A. The assessments used to capture the three main categories of neurobehavioral risk
factors: working memory, impulsivity, and reward processing. Working memory was
measured from the performance of list sorting test, card sort test, and 2-back in the
emotional N-Back Task (EN-back) performed inside a functional magnetic resonance
imaging (fMRI) scanner. The fMRI data from the EN-back task were analyzed based on
the contrast of 2-back versus 0-back, using regions in the frontoparietal and fronto-
thalamic network as the regions of interest (ROIs). Impulsivity was measured by
Urgency-Premeditation-Perseverance-Sensation Seeking-Positive Urgency (UPPS-P),
and Behavioral Inhibition System/ Behavioral Activation System (BIS/BAS), and the
stop signal reaction time (SSRT) during the stop signal task (SST) performed inside the
fMRI scanner. The fMRI data from the SST were analyzed based on the contrast of
correct stop versus correct go in lateral prefrontal cortex, anterior cingulate cortex (ACC),
and striatum ROIs. Reward processing was measured behaviorally by the cash choice
task, and also by comparing the success rate of reward versus neutral conditions during
the monetary incentive delay (MID) task performed inside the fMRI scanner. The fMRI
data from the MID task were analyzed based on the contrast of reward anticipation versus
neutral anticipation, using regions in the ventral striatum and medial frontal cortex as the
ROIs. The ROIs were selected based on the Destrieux atlas (Destrieux et al., 2010). See
"Methods" for more details on the variables. **B.** Diagram of the research aims. First, the
associations between the risk factors for substance use disorders (SUDs) and caffeinated
soda intake were examined cross-sectionally using the baseline ABCD data. Then, we
assessed whether caffeinated soda intake would predict alcohol sipping after 12 months
while controlling other risk factors of SUDs.

Figure 2. Results of binomial LASSO regression predicting daily caffeinated soda intake (daily-drinking group vs. non-drinking group)

A. Estimates of coefficients of the survived variables (x axis: standardized coefficient estimates; y axis: predictor). The site variables were excluded for clarification (see **Supplementary Fig. S2** for results including all variables). Each dot indicates mean of the coefficient, positive coefficient in red and negative coefficient in green. Each error bar indicates 95% confidence interval. **B.** Distribution of the area under the curve (AUC) values (left) and a representative receiver-operation characteristic (ROC) curve (right) for the training and test datasets. **C.** Regions of interest (ROIs) identified as having significant estimates of coefficients during the stop signal task (SST) and emotional N-Back Task (EN-back).

1   *Abbreviations*. LASSO, least absolute shrinkage and selection operator; BMI, body mass
2   index; BAS, Behavioral Activation System; UPPS-P, Urgency-Premeditation-
3   Perseverance-Sensation Seeking-Positive Urgency; IPL, Inferior Parietal Lobule; IFG,
4   Inferior Frontal Gyrus; ACC, Anterior Cingulate Cortex.

Figure 3. Results of binomial LASSO regression predicting alcohol sipping after 12 months

A. Estimates of coefficients of the survived variables (x axis: coefficient estimates; y axis: predictor). The site variables were excluded for clarification (see **Supplementary Fig. S3** for results including all variables). Each dot indicates mean of the coefficient, positive coefficient in red and negative coefficient in green. Each error bar indicates 95% confidence interval. **B.** Distribution of the area under the curve (AUC) values (left) and a representative receiver-operation characteristic (ROC) curve (right) for the training and test datasets. **C.** Rate of alcohol sipping at 12-month follow-up in each group. The asterisk indicates significance from the chi-square test ($p < 0.05$). *Abbreviations*. LASSO, least absolute shrinkage and selection operator; BMI, body mass index; UPPS-P, Urgency-Premeditation-Perseverance-Sensation Seeking-Positive Urgency; EN-Back, emotional N-Back Task; MID, Monetary Incentive Delay Task; NAc, Nucleus Accumbens; OFC, Orbitofrontal cortex.