Plant responses to multiple antagonists are mediated by order of attack and phytohormone crosstalk

Saumik Basu[1*], Robert E. Clark[1], Sayanta Bera[2], Clare L. Casteel[2], and David W. Crowder[1]

[1] Department of Entomology, Washington State University, Pullman, WA, USA, 99164

[2] School of Integrative Plant Science, Plant Pathology and Plant-Microbe Biology Section, Cornell University, Ithaca, NY, USA 14850

Abstract

Plants are often attacked by multiple antagonists, and traits of the attacking organisms, and their order of arrival onto hosts, may affect plant defenses. However, few studies have assessed how multiple antagonists, and varying attack order, affect plant defense or nutrition. To address this, we assessed defensive and nutritional responses of *Pisum sativum* plants after attack by a vector herbivore (*Acrythosiphon pisum*), a non-vector herbivore (*Sitona lineatus*), and a pathogen (*Pea enation mosaic virus,* PEMV). We show PEMV-infectious *A. pisum* induced several pathogen-specific plant defense signals, but these defenses were inhibited when *S. lineatus* was present in peas infected with PEMV. In contrast, feeding by *S. lineatus* induced anti-herbivore defense signals, but these defenses were enhanced by PEMV. *Sitona lineatus* also increased abundance of plant amino acids, but only when they attacked after PEMV-infectious *A. pisum.* Our results suggest that diverse communities of biotic antagonists alter defense and nutritional traits of plants through complex pathways that depend on the identity of attackers and their order of arrival onto hosts. Moreover, we show interactions among a group of biotic stressors can vary along a spectrum from antagonism to enhancement/synergism based on the identity and order of attackers, and these interactions are mediated by a multitude of phytohormone pathways.

Keywords: disease ecology, plant-insect-pathogen interactions, plant defensive chemistry, plant nutrients, species interactions

*Correspondence:

  Saumik Basu

  saumik.basu@wsu.edu

1 INTRODUCTION

Plant hosts have defenses to counter attacks from antagonists such as herbivores and pathogens (Pandey, Ramegowda, & Senthil-Kumar, 2015; Miller, Costa Alves, & Van Sluys, 2017). For example, the jasmonic acid pathway often regulates plant defenses against herbivores, while the salicylic acid pathway often regulates defenses against pathogens (Koornneef & Pieterse, 2008; Thaler, Humphrey, & Whiteman, 2012). While many studies have assessed plant responses to particular antagonists, plants are often challenged by many stressors concurrently, and plant defenses can depend on the order in which antagonists arrive on plants (Thaler et al., 2012; Nejat & Mantri, 2017). For example, herbivores often limit plant defenses against pathogens when they arrive first on plants, but herbivores often have few impacts on plant defenses against pathogens when they arrive after pathogens on plants (Okada, Abe, & Arimura, 2015; Lin et al., 2019). In other contexts, certain organisms 'prime' pathways, promoting defense against subsequent organisms activating the same pathway, such that attack order may not matter (Mauch-Mani, Baccelli, Luna, & Flors, 2017; Ramírez-Carrasco, Martínez-Aguilar, & Alvarez-Venegas, 2017).

Biotic stressors may also alter the nutritional quality of plants by regulating amino acid metabolism (Casteel et al., 2014; Zhou, Lou, Tzin, & Jander, 2015), which can alter the feeding behavior and nutrient uptake by subsequent herbivores (Behmer, 2009; Zhu, Poelman, & Dicke, 2014). For example, the composition of free amino acids constitutively changes in leaves of soybean plants in response to soybean aphids (Chiozza, O'Neal, & MacIntosh, 2010). *Tomato yellow leaf curl virus* also alters the nutritional quality of tomato plants by affecting free amino acid levels in phloem, which alters the amino acid composition of whitefly (*Bemisia tabaci*) honeydew (Guo et al., 2019). However, few studies have explored how the diversity and identity of attacking organisms, and variation in the order of attack, affect nutritional traits of plants.

While there has been considerable research on the jasmonic and salicylic acid pathways, to understand complexities of plant defense it is necessary to assess how biotic antagonists mediate other signaling pathways (e.g., Lacerda, Vasconcelos, Pelegrini, & Grossi de Sa, 2014; Suzuki, 2016). Moreover, it is key to assess how changes in plant defense correlate with plant nutrients. For example, plants in low-nitrogen soil often adopt carbon-based defenses, while plants grown with fertilizer often accumulate more nitrogenous toxins (Cipollini, Walters, & Voelckel, 2017). Nitrogen in plants may also affect both pathogens and herbivores through synthesis of defensive metabolite, nitric oxide, and by nitrogen mobilization (War et al., 2012; Mur, Simpson, Kumari, Gupta, & Gupta, 2017). However, few studies have correlated effects of multiple biotic stressors on both plant chemical signaling and nutritional properties (Petek et al., 2014; Su et al., 2016).

We addressed these knowledge gaps by assessing the response of *Pisum sativum* plants to attack from a piercing-sucking vector herbivore, the pea aphid (*Acrythosiphon pisum*), a chewing non-vector herbivore, the pea leaf weevil (*Sitona lineatus*), and an aphid-borne pathogen, *Pea-enation mosaic virus* (PEMV). These organisms co-occur in ecosystems of eastern Washington and northern Idaho, USA, and interactions between them can affect plant traits and signaling pathways affecting insects and pathogens (Chisholm, Eigenbrode, Clark, Basu, & Crowder, 2019; Bera, Blundell, Liang, Crowder, & Casteel, 2020). However, the order in which herbivores and pathogens arrive on hosts, which varies across sites (Chisholm et al., 2019), may impact plant traits and defenses. To address this, we varied the diversity, identity, and order of attack among this community of biotic antagonists and assessed resulting changes in gene expression and phytohormones related to plant defense and nutrition. Our study revealed how plant responses to diverse stressors can mediate complex species interactions within a pathosystem.

2 MATERIALS AND METHODS

2.1 Study system

The Palouse region of eastern Washington and northern Idaho, USA, is home to many legumes

including *P. sativum* (Black et al., 1998). In *P. sativum* fields, *S. lineatus*, a chewing herbivore,

co-occurs with *A. pisum*, a phloem-feeding herbivore that can transmit pathogens such as PEMV

(Chisholm et al., 2019). PEMV is one of several viruses that infects *P. sativum*, and this

pathogen is obligately transmitted by aphids in a persistent manner (Chisholm et al., 2019).

Sitona lineatus adults overwinter outside of *P. sativum* fields and migrate into fields in late

spring before *A. pisum* arrive (Cárcamo et al., 2018). After *S. lineatus* eggs hatch, larvae burrow

into the soil to feed and pupate before emerging as adults in the summer (Cárcamo et al., 2018);

these second-generation adults often occur on plants under attack from *A. pisum* and PEMV

(Chisholm et al., 2019). Thus, *S. lineatus* attacks plants in the field both before and after *A.*

pisum and PEMV. However, it is unknown if responses of *P. sativum* differ based on the number

of stressors, and their order of attack. Moreover, molecular mechanisms that mediate interactions

among these stressors are largely unknown (Chisholm et al., 2019; Bera et al., 2020).

To address these questions, we conducted greenhouse assays to assess interactions between

S. lineatus, *A. pisum,* and PEMV on *P. sativum* plants, and molecular mechanisms affecting these

interactions. First-generation adult *S. lineatus* for experiments were collected from commercial

P. sativum fields, or wild patches of *Vicia villosa*, immediately prior to experiments. Colonies of

infectious *A. pisum* with PEMV, and uninfectious *A. pisum*, were started from Palouse field-

collected individuals (Chisholm et al., 2019) and were maintained on *P. sativum* plants in a

greenhouse (21-24°C during day cycle, 16-18°C during dark cycle, 16:8 h light:dark).

2.2 Experimental design

We conducted a 3×2 greenhouse (21-24°C day cycle, 16-18°C dark cycle, 16:8 h light:dark)

experiment that varied *S. lineatus*, *A. pisum*, and PEMV (Fig. 1). There were three *S. lineatus*

treatments: (i) control: no adults prior to *A. pisum* treatments (none), (ii) two adults that fed for

48 h prior to *A. pisum* treatments (first), and (iii) two adults that fed for 48 h after *A. pisum*

treatments (second). The two *A. pisum* treatments were: (i) sham: 10 5-d old uninfectious adults

that fed for 48 h and (ii) PEMV: 10 5-d old PEMV-infectious adults that fed for 48 h. For

treatments with *S. lineatus* first, they were removed by hand prior to *A. pisum* treatments; for

treatments with *A. pisum* first, they were removed by aspirator prior to *S. lineatus* treatments.

Treatments were conducted on individual *P. sativum* plants in mesh 'bug dorms' ($0.6 \times 0.6 \times 0.6$

m), with six replicates randomly assigned to each treatment in a factorial design (3 *S. lineatus*

treatments $\times$ 2 *A. pisum* treatments). After insects were removed, plants were allowed to develop

for 7 d before we harvested tissue to assess viral titer, gene expression, and nutrients. Tissue

samples from the whole aboveground portion plants were collected and flash frozen in liquid

nitrogen and stored in a -80°C freezer until processing. Viral titer samples confirmed that 100%

of plants in the PEMV treatments became infected over the course of the experiment.

2.3 Analysis of plant defense and biosynthetic genes

Plant tissue was processed using liquid nitrogen in sterilized mortars and pestles. Powdered

tissue samples (50 to 100 mg) were used for RNA extraction with Promega SV total RNA

isolation kits (Promega, Madison, WI). The quantity and quality of RNA was estimated on a

NanoDrop1000 and agarose gel electrophoresis, respectively and 1 µg of total RNA from each

sample was used for cDNA synthesis (Bio-Rad iScript cDNA Synthesis kits). Gene specific

primers (Table S3) for qRT-PCR were designed using the IDT Primer Quest Tool. Each qRT-

PCR reaction (10 µl) was set up containing 3 µl of ddH2O, 5 µl of iTaq Univer SYBR Green

Supermix (Bio-Rad), 1 µl of specific primer mix (forward and reverse [concentration 10µM]),

and 1 µl of diluted (1: 25) cDNA template. Reactions were set up in triplicates for each sample

and ran on a CFX96 qRT-PCR machine (Biorad). The qRT-PCR program included an initial

denaturation for 3 min at 95°C, followed by 40 denaturation cycles for 15 s at 95°C, annealing

for 30 s at 60°C, and extension for 30 s at 72°C. For melting curve analysis, a dissociation step

cycle (55°C for 10 s and then 0.5°C for 10 s until 95°C) was added. The comparative $2^{-\Delta\Delta Ct}$

method (Livak & Schmittgen, 2001; Kozera & Rapacz, 2013) was used to calculate the relative

expression level of each gene, with β-tubulin as an endogenous control.

We assessed expression of seven genes associated with defense in peas. Gene sequences

were obtained using accession numbers (available genes) or using Pea Marker Database

(Kulaeva et al., 2017) and blast searching through the reference genome (Kreplak et al., 2019).

Four genes were associated with plant hormone biosynthesis: (i) *Isochorismate synthase1* (*ICS1*)

(salicylic acid), (ii) *Lipoxygenase 2* (*LOX2*) (jasmonic acid), (iii) *Aldehyde oxidase 3* (*AO3*)

(abscisic acid), and (iv) *Gibberellin 2-oxidase (GA2ox)* (gibberellic acid). *ICS1* converts

chorismate to isochorismate, a precursor of salicylic acid biosynthesis (Seguel et al., 2018),

while *LOX2* is a precursor to jasmonic acid biosynthesis (Wasternack & Hause, 2013). *AO3*

catalyzes abscisic acid biosynthesis by oxidizing abscisic aldehyde, and *GA2OX* catalyzes

bioactive giberrelic acids or their immediate precursors to inactive forms (Zdunek-Zastocka &

Sobczak, 2013; Serova, Tsyganova, Tikhonovich, & Tsyganov, 2019; He et al., 2019). All of

these gene transcripts can affect plant defense and plant-microbe interactions (Lee et al., 2012;

Yergaliyev et al., 2016).

The three additional genes examined were associated with defense response transcripts that occur downstream from hormone induction. One of these genes, *Pathogenesis-related protein 1* (*PR1*) affects systemic acquired resistance-mediated defense signaling and occurs downstream in the salicylic acid pathway (Fondevilla, Küster, Krajinski, Cubero, & Rubiales, 2011; Miranda et al., 2017). The second defense response transcript was an antimicrobial defensin peptide called *Disease resistance response* gene *(DRR230),* which has been reported to provide resistance in peas against various pathogens (Lacerda et al., 2014; Selim, Sanssené, Rossard, & Courtois, 2017). The third defense response transcript assessed was *Lectin (PsLectin)*. Plant lectins are a group of carbohydrate binding proteins, and *Lectin* genes can be induced by salicylic acid, jasmonic acid, and herbivores to stimulate phytoalexin and pistatin production in peas (Fondevilla et al., 2011; Armijo et al., 2013; Macedo, Oliveira, & Oliveira, 2015).

2.4 Measurement of plant phytohormones

Plant tissue samples were assessed for three phytohormones: jasmonic acid, salicylic acid, and abscisic acid following procedures of Patton, Bak, Sayre, Heck, & Casteel (2019). Briefly, tissue samples were first flash frozen in liquid nitrogen before being lyophilized and weighed. Hormones were extracted in iso-propanol:H_2O:HCL_{1MOL} (2:1:0.005) with 100 µl of internal standard solution (1000 pg of each). Samples were evaporated to dryness, resuspended in 100 µl of MeOH, filtered, and 10 µl of each sample was injected into an Agilent Technologies 6420 triple quad liquid chromatography-tandem mass spectrometry instrument (Agilent, Santa Clara, CA). A Zorbax Extend-C18 column 3.0×150mm (Agilent, Santa Clara, CA) was used with 0.1% formic acid in water (A) and 0.1% (v/v) formic acid in acetonitrile (B) at a flow rate of 600 mL min^{-1}. The gradient used was 0–1 min, 20% B; 1–10 min, linear gradient to 100% B; 10-

13 min, 100% A. Retention times were: jasmonic acid (D5) standard (5.740 min), jasmonic acid

(5.744 min), salicylic acid D4 standard (4.677 min), salicylic acid (4.720 min).

2.5 Analysis of plant nutritional components

For amino acid analysis, leaf tissue was lyophilized, weighed, and extracted with 20mM of HCL

(Patton et al., 2019). Derivation was done using AccQTag reagents following the manufacturer's

instructions (Waters, Shinagawa-ku, Tokyo), and derivatised samples (10 μl) were then injected.

Ground tissue was extracted with 100 μl of 20 mM HCl, centrifuged, and the supernatant was

saved. Amino acids were derivatized using AccQ-Fluor reagent kits (Waters, Milford, MA), with

L-Norleucine as an internal standard. 10 μl from each sample were injected with an Agilent 1260

Infinity pump with a Nova-Pak C18 column and fluorescence detector, and Agilent Chemstation

software for data recording. Amino acid derivatives were detected with an excitation wavelength

of 250 nm and an emission wavelength of 395 nm. Peak areas were compared to a standard curve

made from a serial dilution of amino acid standards (Sigma-Aldrich, St. Louis, MO). injected

into a Agilent 1260 Infinity HPLC (Agilent, Santa Clara, CA) with a Nova-Pak C18 column

(Casteel et al., 2014). Solvent A, AccQ-Tag Eluent A, was premixed from Waters; Solvent B

was acetonitrile:water (60:40). The gradient used was 0–0.01 min, 100% A; 0.01–0.5 min, linear

gradient to 3% B; 0.5–12 min, linear gradient to 5% B; 12–15 min, linear gradient to 8% B; 15–

45 min, 35% B; 45–49 min, linear gradient to 35% B; 50–60 min, 100% B. The flow rate was

1.0 ml min^{-1}. Amino acid derivatives were measured with an Agilent fluorescence detector with

an excitation wavelength of 250 nm and an emission wavelength of 395 nm. For concentration

calculations, standard curves were generated for each amino acid using dilutions of the standard.

2.6 Data analysis

To evaluate effects of our treatments on host-plant defenses and host-plant quality, we ran a

series of multivariate models using R ver. 3.5.2 (R Working Group, 2018). First, gene expression

was evaluated with *ICS1*, *LOX2*, *GA2ox*, *AO3*, *PR1*, *DRR230*, and *PsLectin* as the responses,

with MANOVA to assess treatment effects on relative gene expression ($2^{-\Delta\Delta Ct}$) based on cycle

threshold values for each observed gene transcript. Estimated marginal mean of Ct values, and

standard error of the mean, were generated using the emmeans package in R (Lenth, 2016). The

methodology for $2^{-\Delta\Delta Ct}$ followed modified recommendations from Rao, Huang, Zhou, & Lin

(2013) and Kozera & Rapacz (2013), using housekeeping gene *β-tubulin* to normalize expression

and a sham aphid (non-infective Pea aphid and no weevil addition) treatment as a control.

Hormone levels in plants were evaluated using MANOVA, with salicylic acid, jasmonic

acid, and abscisic acid as responses (3 variables). Total amino acid content was evaluated using a

generalized linear model (GLM) with total concentration among all amino acids as the response.

All models assessed treatment effects, using *S. lineatus* addition, *A. pisum* infection status, and

their interaction as predictors. Finally, changes in the amino acid profile was evaluated using

non-metric multidimensional scaling (NMDS) with the vegan package (Oksanen et al., 2019)

following Ceulemans, Hulsmans, Ende, & Honnay (2017).

3 RESULTS

3.1 Effects of multiple antagonists and attack order on plant gene transcripts

Transcription of plant genes associated with hormone biosynthesis across four pathways:

(i) salicylic acid (*ICS1*), (ii) jasmonic acid (*LOX2*), (iii) abscisic acid (*AO3*), and (iv) gibberellic

acid (*GA2ox*), were induced by PEMV when *S. lineatus* was not present (Fig. 2A-D, Table S1,

Pillai = 0.942, *P* = 0.002). However, there was no induction of any of these biosynthesis genes

in response to PEMV when *S. lineatus* was present, indicating that *S. lineatus* inhibited plant

defense against PEMV (Table S1, A x W interaction, *Pillai* = 1.509, *P* = 0.021, Fig. 2). In both

the presence and absence of PEMV, *S. lineatus* induced transcription of *LOX2*, but *S. lineatus*

did not directly modify the expression of ICS1, AO3 or GA2ox (Fig. 2B-D, Table S1). When

PEMV-infectious *A. pisum* attacked following *S. lineatus*, there was greater induction of LOX2

compared to when *S. lineatus* attacked alone (Table S1, A x W interaction, *Pillai* = 1.509, *P* =

0.021, Fig. 2). In contrast to the antagonism exerted by *S. lineatus* on the response to PEMV, this

represents enhancement of plant defense when PEMV infection followed attack by *S. lineatus*.

All three defense response transcripts (*PR1, DDR230, PsLectin*) were induced by PEMV

when *S. lineatus* was not present (Fig. 3); similarly, each transcript was induced by *S. lineatus*

when PEMV was not present (Fig. 3, Table S1; A ⬚ W interaction, *F* = 2.64, *P* = 0.111). When

S. lineatus attacked second, the expression level of *PR1 and Lectin* did not change compared to

when weevils were absent. The effects of PEMV on the transcripts was modified by the presence

of *S. lineatus* and attack order. While *DDR230* was induced by PEMV (Table S1, *F* = 47.181, *P*

< 0.001), this effect diminished when *S. lineatus* was present after PEMV (Fig. 3B). Similarly,

the effects of PEMV on *PR1* were inhibited when *S. lineatus* attacked second (Fig. 3), whereas

that the induction of lectin by PEMV was not altered by *S. lineatus* in either order (Fig. 3).

3.2 Effects of multiple antagonists and attack order on plant phytohormones

We observed variation in phytohormones in response to *A. pisum* (Table S2, *Pillai* = 0.95, *P* <

0.001) and *S. lineatus* (Table S2, *Pillai* = 1.195, P < 0.001). PEMV-infectious *A. pisum* strongly

induced salicylic acid (Table S2, *F* = 254.2, *P* < 0.001), but this was inhibited when *S. lineatus*

attacked after PEMV (Fig. 4A, Tukey HSD). PEMV did not affect jasmonic acid (Table S2, $F =$

0.97, P = 0.34), but the order of *S. lineatus* did (Table S2, $F = 5.30$, $P = 0.018$). Both *S. lineatus*

(Table S2, $F = 4.10$, P = 0.037) and infectious *A. pisum* induced abscisic acid (Table S2, $F =$

9.96, P = 0.006) and this effect was contingent on the attack order (Table S2, A ⬚ W, $F = 4.32$, P

= 0.032, Fig. 4, Tukey HSD). Jasmonic acid levels were suppressed by *S. lineatus* when

attacking prior to non-infectious sham *A. pisum*, but not on plants already attacked by PEMV

(Fig. 4B, Tukey HSD).

3.3 Effects of multiple antagonists and attack order on plant nutrients

Feeding by *S. lineatus* increased the total amino acid levels (GLM, $\chi 2 = 9.19$, $P = 0.01$, Fig 5),

but PEMV-infectious *A. pisum* did not (GLM, $\chi^2 = 0.044$, $P = 0.83$), and this effect was not

modified depending on attack order (GLM, A ⬚ W interaction, $\chi^2 = 0.24$, $P = 0.63$). Non-metric

multidimensional scaling (NMDS) analysis of amino acid composition also showed that changes

to amino acid availability was most different among treatments for alanine, arginine, lysine, and

glycine (Ordination plot, Fig S1).

5 DISCUSSION

Assessing interactions between biotic stressors and plants in food webs is critical to understand

the dynamics of these interactions. Plant responses to stressors are often specific to the attacker,

and both phytochemical responses and plant nutritional status can affect susceptibility to specific

stressors (van Geem, Gols, Raaijmakers, & Harvey, 2016*; Shikano, 2017). We show plants

responded in complex ways to unique biotic stressors, including a piercing-sucking herbivore (*A.*

pisum), a chewing herbivore (*S. lineatus*), and a virus (PEMV). Our study is among the first to

assess how the order of attack, and diversity of stressors, mediate the defensive responses of

plants and plant nutritional status (see also Vos, Moritz, Pieterse, & Van Wees, 2015). Our

results show that plants traits varied in response to the type of attacker, the number of stressors,

and their order of arrival. Moreover, we show that assessment of multiple gene transcripts,

phytohormones, and plant nutrients provides a more comprehensive perspective on mechanisms

driving plant-insect-pathogen interactions than any isolated response.

We found PEMV caused broad defensive responses in *P. sativum* by inducing specific gene

transcripts and phytohormones (Figs. 2, 3 & 4). Biotropic pathogens such as PEMV are known to

activate salicylic acid signaling (Singh, Swain, Singh, & Nandi, 2018; Chisholm, Sertsuvalkul,

Casteel, & Crowder, 2018). This was reflected by increased expression of the *ICS1* biosynthesis

gene, increased salicylic acid hormone levels, and increased expression of the downstream

defensive transcript *PR1* when PEMV was present. However, effects of PEMV were not limited

to salicylic acid, as PEMV induced gene transcripts often associated with biosynthesis of abscisic

acid *(AO3)*, and giberrellic acid (*GA2ox*), while also affecting defense response genes that occur

downstream from induction of these hormones (*PR1, DRR230*, *PsLectin*). However, PEMV

alone can not induce gene transcript associated with biosynthesis of JA (*LOX2*), but PEMV

infection after weevil herbivory induced *LOX2* accumulation.   Similar results in *P. sativum* have

been observed in response to fungal infection by *Mycosphaerella pinodes* and *Phoma koolunga*,

where infections induce defense related genes across multiple signaling pathways (Fondevilla et

al., 2011; Tran, You, & Barbetti, 2018). However, increased expression of *LOX2* and *AO3* gene

transcripts (Fig. 2) were not reflected by increased levels of jasmonic acid or abscisic acid (Fig.

4). This suggests that measuring phytohormones, or gene transcripts, in isolation may fail to

reveal more complex pathways by which plants respond to stress (Kazan & Lyons, 2014).

While PEMV had broad effects on plants, *S. lineatus* attenuated these responses. When *S.*

lineatus was present, before or after PEMV, the expression of three out of four biosynthesis gene

transcripts (with the exception of *LOX2*) in response to infectious aphids were comparatively

weaker (Fig. 2). Increased expression of *LOX2* may be due to *S. lineatus* inducing expression of

defensive transcripts associated with JA-mediated chewing herbivore attack and the effect was

further enhanced by PEMV infection after *S. lineatus* feeding (Fig. 2). Effects of PEMV on plant

defense genes (*PR1*, *DDR230*, *PsLectin*) were also affected by *S. lineatus*, but varied with attack

order. Overall, in this study the order of attack seemed to have stronger effects on downstream

plant defense gene transcripts than on hormone biosynthesis gene transcripts.

While *S. lineatus* increased expression of two (*LOX2* and *GA2ox*) of the four biosynthesis

gene transcripts studied when PEMV was not present, expression of *LOX2* was enhanced when

PEMV was also present (Fig. 2). In contrast, PEMV caused decreased expression of two genes

(*PR1*, *DRR230*) that were induced by *S. lineatus* was present alone (Fig. 2). For plants attacked

first by either PEMV or *S. lineatus*, we observed the strongest evidence for mutual antagonism at

the gene transcript level rather than for phytohormones (Figs. 2-4). Our study shows interactions

among a set of stressors can vary based on attack order. Here, we found that *S. lineatus* feeding

following PEMV infection inhibited plant defense (mutual antagonism), while *LOX2* expression

was enhanced when PEMV infection followed *S. lineatus* feeding (synergistic effect). While the

first effect is in line with studies showing "mutual antagonism" between chewing herbivores and

biotropic pathogens (Thaler, Agrawal, & Halitschke, 2010; Vos et al., 2015), our study suggests

the order of attack can lead to variation along a spectrum from antagonism to enhancement.

Our results provide evidence that the order of arrival of biotic stressors on plants can play a

crucial role in determining plants' response to these attackers. While mutual antagonism between

S. lineatus and PEMV was common, for some genes these effects only occurred when *S. lineatus*

attacked first, and for others when *S. lineatus* attacked second (Figs. 2-4). Mutual antagonism has

most often been studied as effects of a prior attacker affecting a subsequent attacker, such as

when a herbivore alters gene activation or phytohormones in ways that attenuate performance of

subsequent attackers (Kessler & Halitschke, 2007; Erb, Robert, Hibbard, & Turlings, 2011;

Stam, Mantelin, McLellan, & Thilliez, 2014; Huang et al., 2017). However, our results suggest

that a second attacker may also mitigate defensive responses against the first attacker in ways

that might affect plant defense and propagation of pathogens. For example, we show that plants

infected by PEMV had decreased defenses when subsequently attacked by *S. lineatus* (Fig. 3),

which should promote PEMV replication. Moreover, our results suggest that, PEMV infection

induces pathogen defense and *S. lineatus* inhibits that if they appear on plants after the infection

has been established. This may be more strongly expressed as variation in gene transcripts rather

than hormone levels, a result that has similarly been seen in *Arabidopsis* in response to pathogen

infection (Anderson et al., 2004).

Mutual antagonism in plant signaling pathways has most commonly been examined in

regard to tradeoffs between jasmonic acid and salicylic acid. Our results show these tradeoffs

extend to other signaling pathways. For example, jasmonic acid exhibits antagonism with

abscisic acid in *Arabidopsis* following attack from *Fusarium oxysporum* (Anderson et al., 2004).

Mutual antagonism between jasmonic acid and gibberellic acid, and jasmonic acid and abscisic

acid, have also been reported (Yang, Yang, & He, 2013; Okada et al*.,* 2015; Liu & Hou, 2018).

For example, jasmonic acid facilitates defense over growth by repressing degradation of DELLA

protein in rice and *Arabidopsis*, but elevated DELLA proteins interfere with the gibberellic acid

pathway by binding to growth promoting transcription factors associated with gibberellic acid

signaling (Yang et al., 2012, 2013; Okada et al., 2015). Antagonistic relationships between

giberellic acid and abscisic acid have also been reported in both mono and dicot plants and

regulated by various transcription factor regulators in response to diverse environmental cues

(Liu & Hou, 2018). However, antagonisms between salicylic acid and abscisic acid may actually

lead to synergism between jasmonic acid and abscisic acid, where elevated abscisic acid levels

following infection with *Pseudomonas syringae* induce jasmonic acid in *Arabidopsis*, which in

turn limits the levels of salicylic acid (Fan, Hill, Crooks, Doerner, & Lamb, 2009). Overall, these

results suggest that a broad examination of genes and hormones are needed to elucidate pathways

underlying plant-insect-pathogen interactions in *P. sativum* and other plants.

Our results suggest mutual antagonism may also occur among defense gene transcripts that

are associated with a single signaling pathway. For example, the induction of *PR1,* a salicylic

acid-responsive gene, was mitigated by *S. lineatus* attack after PEMV infection, as may be

expected with antagonism between jasmonic acid and salicylic acid. However, the expression of

ICS1, another gene associated with the biosynthesis of salicylic acid, was not responsive to *S.*

lineatus. This has been seen in other studies where *ICS1* was not induced by caterpillar feeding

although other genes associated with salicylic acid were (Onkokesung, Reichelt, van Doorn,

Schuurink, & Dicke 2016). These results suggest that a plant's response to multiple stressors is

unlikely to result from simple crosstalk but rather from interactions among multiple signaling

pathways that may exhibit complex responses.

In addition to affecting plant gene expression and phytohormones, plant pathogens such as

viruses can also alter nutritional quality of their host plants in ways that affect vectors (Mauck,

Bosque☐Pérez, Eigenbrode, Moraes, & Mescher, 2012; Wang, Senthil-Kumar, Ryu, Kang, &

Mysore, 2012; Patton et al., 2019). Similarly, non-vector herbivores may strongly affect the

quantity and quality of plant nutrients (Ángeles-López, Rivera-Bustamante, & Heil, 2016). For

example, *pepper golden mosaic virus* (PGMV) infection in *Capsicum annuum* increased levels

of the amino acids proline, tyrosine, valine but decreased levels of histidine and alanine. In the

same system, the greenhouse whitefly, *Trialeurodes vaporarioum*, reversed the levels of these

amino acids (Ángeles-López et al., 2016). Arrival of *S. lineatus* before PEMV infection

suppressed the amount of total amino acids in peas, while enhanced amino acid level was

detected if *S. lineatus* damaged peas after PEMV infected was established. This suggests the

intriguing possibility that antagonism between a pathogen and non-vector herbivore can occur at

the level of amino acid production in plants.

Overall, our study provides example of complex interactions between a vector-borne plant

pathogen and a non-vector herbivore that varies from antagonism to enhancement and manifest

as changes in plant gene transcripts, phytohormones levels, and plant nutrients. However, we

show that assessing the order of attack is necessary to best understand the complexity and

mechanisms of plant-insect-pathogen interactions. Moreover, our study suggests complete

pathways must be characterized as differences are evident even when a few transcripts and

metabolites are analyzed., often measured with associated gene transcripts (Bedini, Mercy,

Schneider, Franken, & Lucic-Mercy, 2018; Ángeles-López el al., 2016; Shi et al., 2019), may

fail to capture mechanisms by which plants interact with multiple stressors. Our results

demonstrate both the order of arrival, and the diversity of interactions, determine plant responses

to stress through the combined action of defense gene activation, phytohormone accumulation,

and modification of plant nutrients. Characterizing the pathways by which plants respond to

single and multiple stressors, with varying attack order, can shed light on the mechanisms that

shape food web interactions among plants, herbivores, and pathogens.

ACKNOWLEDGEMENTS

We thank the many undergraduates who helped in various experiments, data collection, and O.

Hill for assistance with data analysis. This research was supported by USDA☐NIFA Grants

2016☐67011☐24693, 2017-67013-26537, and Hatch project 1014754.

REFERENCES

Anderson, J. P., Badruzsaufari, E., Schenk, P. M., Manners, J. M., Desmond, O. J., Ehlert, C., …

Kazan, K. (2004). Antagonistic interaction between abscisic acid and jasmonate-ethylene

signaling pathways modulates defense gene expression and disease resistance in

Arabidopsis. *The Plant Cell*, *16*(12), 3460–3479. doi: 10.1105/tpc.104.025833

Ángeles-López, Y. I., Rivera-Bustamante, R. F., & Heil, M. (2016). Colonization by phloem-

feeding herbivore overrides effects of plant virus on amino acid composition in phloem of

chili plants. *Journal of Chemical Ecology*, *42*(10), 985–988. doi: 10.1007/s10886-016-

0747-2

Armijo, G., Salinas, P., Monteoliva, M. I., Seguel, A., García, C., Villarroel-Candia, E., …

Holuigue, L. (2013). A salicylic acid-induced lectin-like protein plays a positive role in the

effector-triggered immunity response of *Arabidopsis thaliana* to *Pseudomonas syringae*

Avr-Rpm1. *Molecular Plant-Microbe Interactions: MPMI*, *26*(12), 1395–1406. doi:

10.1094/MPMI-02-13-0044-R

Bedini, A., Mercy, L., Schneider, C., Franken, P., & Lucic-Mercy, E. (2018). Unraveling the

initial plant hormone signaling, metabolic mechanisms and plant defense triggering the

endomycorrhizal symbiosis behavior. *Frontiers in Plant Science*, *9*. doi:

10.3389/fpls.2018.01800

Behmer, S. T. (2009). Insect herbivore nutrient regulation. *Annual Review of Entomology*, *54*(1), 165–187. doi: 10.1146/annurev.ento.54.110807.090537

Bera, S., Blundell, R., Liang, D., Crowder, D. W., & Casteel, C. L. (2020). The oxylipin signaling pathway is required for increased aphid attraction and retention on virus-infected plants. *Journal of Chemical Ecology*. doi: 10.1007/s10886-020-01157-7

Black, A. E., Strand, E., Wright, R. G., Scott, J. M., Morgan, P., & Watson, C. (1998). Land use history at multiple scales: implications for conservation planning. *Landscape and Urban Planning*, *43*(1), 49–63. doi: 10.1016/S0169-2046(98)00096-6

Cárcamo, H. A., Vankosky, M. A., Olfert, O. O., Wijerathna, A., Evenden, M. L., & Meers, S.B. (2018). Progress toward integrated pest management of pea leaf weevil: a review. *Annals of the Entomological Society of America*, *111(4)*, 144–153. (doi:10.1093/aesa/say007)

Casteel, C. L., Yang, C., Nanduri, A. C., De Jong, H. N., Whitham, S. A., & Jander G. (2014). The NIa-Pro protein of *Turnip mosaic virus* improves growth and reproduction of the aphid vector, *Myzus persicae* (green peach aphid). *Plant Journal*, 77, 653–663. doi: 10.1111/tpj.12417

Ceulemans, T., E. Hulsmans, W. V. Ende, & O. Honnay. (2017). Nutrient enrichment is associated with altered nectar and pollen chemical composition in *Succisa pratensis* (Moench) and increased larval mortality of its pollinator *Bombus terrestris* L. *PLOS ONE* *12*(4), e0175160. doi: 10.1371/journal.pone.0175160

Chiozza, M. V., O'Neal, M. E., & MacIntosh, G. C. (2010). Constitutive and induced differential accumulation of amino acid in leaves of susceptible and resistant soybean plants in response to the soybean aphid (Hemiptera: Aphididae). *Environmental Entomology*, *39*(3), 856–864. doi: 10.1603/EN09338

Chisholm, P. J., Eigenbrode, S. D., Clark, R. E., Basu, S., & Crowder, D. W. (2019). Plant-

mediated interactions between a vector and a non-vector herbivore promote the spread of a

plant virus. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1911), 20191383.

doi: 10.1098/rspb.2019.1383

Chisholm, P. J., Sertsuvalkul, N., Casteel, C. L., & Crowder, D. W. (2018). Reciprocal plant-

mediated interactions between a virus and a non-vector herbivore. *Ecology*, *99*(10), 2139–

2144. doi: 10.1002/ecy.2449

Cipollini, D., Walters, D., & Voelckel, C. (2017). Costs of resistance in plants: from theory to

evidence. In *Annual Plant Reviews online* (pp. 263–307). doi:

10.1002/9781119312994.apr0512

Erb, M., Robert, C. A. M., Hibbard, B. E., & Turlings, T. C. J. (2011). Sequence of arrival

determines plant-mediated interactions between herbivores. *Journal of Ecology*, *99*(1), 7–

15. doi: 10.1111/j.1365-2745.2010.01757.x

Fan, J., Hill, L., Crooks, C., Doerner, P., & Lamb, C. (2009). Abscisic acid has a key role in

modulating diverse plant-pathogen interactions. *Plant Physiology*, *150*(4), 1750–1761. doi:

10.1104/pp.109.137943

Fondevilla, S., Küster, H., Krajinski, F., Cubero, J. I., & Rubiales, D. (2011). Identification of

genes differentially expressed in a resistant reaction to *Mycosphaerella pinodes* in pea

using microarray technology. *BMC Genomics*, *12*, 28. doi: 10.1186/1471-2164-12-28

Guo, L., Su, Q., Yin, J., Yang, Z., Xie, W., Wang, S., … Zhang, Y. (2019). Amino acid

utilization may explain why *Bemisia tabaci* Q and B differ in their performance on plants

infected by the *Tomato yellow leaf curl virus*. *Frontiers in Physiology*, *10*. doi:

10.3389/fphys.2019.00489

He, H., Liang, G., Lu, S., Wang, P., Liu, T., Ma, Z., Zuo, C., Sun, X., Chen, B., & Mao, J.

(2019). Genome-wide identification and expression analysis of GA2ox, GA3ox, and

GA20ox are related to gibberellin oxidase genes in grape (*VitisVinifera* L.). *Genes, 10*(9),

680. https://doi.org/10.3390/genes10090680

Huang, W., Robert, C. A. M., Hervé, M. R., Hu, L., Bont, Z., & Erb, M. (2017). A mechanism

for sequence specificity in plant-mediated interactions between herbivores. *New*

Phytologist, *214*(1), 169–179. doi: 10.1111/nph.14328

Kazan, K., & Lyons, R. (2014). Intervention of phytohormone pathways by pathogen effectors.

The Plant Cell, *26*(6), 2285–2309. doi: 10.1105/tpc.114.125419

Kessler, A., & Halitschke, R. (2007). Specificity and complexity: the impact of herbivore-

induced plant responses on arthropod community structure. *Current Opinion in Plant*

Biology, *10*(4), 409–414. doi: 10.1016/j.pbi.2007.06.001

Koornneef, A., & Pieterse, C. M. J. (2008). Cross talk in defense signaling. *Plant Physiology*,

146(3), 839–844. doi: 10.1104/pp.107.112029

Kozera, B., & Rapacz, M. (2013). Reference genes in real-time PCR. *Journal of Applied*

Genetics, *54*(4), 391–406. doi: 10.1007/s13353-013-0173-x

Kreplak, J., Madoui, M.-A., Cápal, P., Novák, P., Labadie, K., Aubert, G., … Burstin, J. (2019).

A reference genome for pea provides insight into legume genome evolution. *Nature*

Genetics, *51*(9), 1411–1422. doi: 10.1038/s41588-019-0480-1

Kulaeva, O. A., Zhernakov, A. I., Afonin, A. M., Boikov, S. S., Sulima, A. S., Tikhonovich, I.

A., & Zhukov, V. A. (2017). Pea Marker Database (PMD) – A new online database

combining known pea (*Pisum sativum* L.) gene-based markers. *PLOS ONE*, *12*(10),

e0186713. doi: 10.1371/journal.pone.0186713

Lacerda, A. F., Vasconcelos, E. A. R., Pelegrini, P. B., & Grossi de Sa, M. F. (2014). Antifungal

defensins and their role in plant defense. *Frontiers in Microbiology*, *5*, 116. doi:

10.3389/fmicb.2014.00116

Lee, Y., Kim, Y.-C., Kim, S. Y., Lee, I.-J., Choi, D., Paek, K.-H., … Park, J. M. (2012). A novel

gibberellin 2-oxidase gene *CaGA2ox1* in pepper is specifically induced by incompatible

plant pathogens. *Plant Biotechnology Reports*, *6*(4), 381–390. doi: 10.1007/s11816-012-

0235-2

Lenth, R.V. 2016. Least-Squares Means: The R package lsmeans. Journal of Statistical Software,

69:1-33.

Lin, D., Xu, Y., Wu, H., Liu, X., Zhang, L., Wang, J., & Rao, Q. (2019). Plant defense responses

induced by two herbivores and consequences for whitefly *Bemisia tabaci*. *Frontiers in*

Physiology, *10*. doi: 10.3389/fphys.2019.00346

Liu, X., & Hou, X. (2018). Antagonistic regulation of ABA and GA in metabolism and signaling

pathways. *Frontiers in Plant Science*, *9*. doi: 10.3389/fpls.2018.00251

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-

time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif.)*,

25(4), 402–408. doi: 10.1006/meth.2001.1262

Macedo, M. L. R., Oliveira, C. F. R., & Oliveira, C. T. (2015). Insecticidal activity of plant

lectins and potential application in crop protection. *Molecules*, *20*(2), 2014–2033. doi:

10.3390/molecules20022014

Mauch-Mani, B., Baccelli, I., Luna, E., & Flors, V. (2017). Defense priming: an adaptive part of

induced resistance. *Annual Review of Plant Biology*, *68*(1), 485–512. doi: 10.1146/annurev-

arplant-042916-041132

Mauck, K., Bosque☐Pérez, N. A., Eigenbrode, S. D., Moraes, C. M. D., & Mescher, M. C.

(2012). Transmission mechanisms shape pathogen effects on host–vector interactions:

evidence from plant viruses. *Functional Ecology*, *26*(5), 1162–1175. doi: 10.1111/j.1365-

2435.2012.02026.x

Miller, R. N. G., Costa Alves, G. S., & Van Sluys, M.-A. (2017). Plant immunity: unravelling

the complexity of plant responses to biotic stresses. *Annals of Botany*, *119*(5), 681–687.

doi: 10.1093/aob/mcw284

Miranda, V. de J., Porto, W. F., Fernandes, G. da R., Pogue, R., Nolasco, D. O., Araujo, A. C.

G., … Franco, O. L. (2017). Comparative transcriptomic analysis indicates genes

associated with local and systemic resistance to *Colletotrichum graminicola* in maize.

Scientific Reports, *7*(1), 2483. doi: 10.1038/s41598-017-02298-8

Mur, L. A. J., Simpson, C., Kumari, A., Gupta, A. K., & Gupta, K. J. (2017). Moving nitrogen to

the centre of plant defence against pathogens. *Annals of Botany*, *119*(5), 703–709. doi:

10.1093/aob/mcw179

Nejat, N., & Mantri, N. (2017). Plant immune system: crosstalk between responses to biotic and

abiotic stresses the missing link in understanding plant defence. *Current Issues in*

Molecular Biology, *23*, 1–16. doi: 10.21775/cimb.023.001

Okada, K., Abe, H., & Arimura, G. (2015). Jasmonates induce both defense responses and

communication in monocotyledonous and dicotyledonous plants. *Plant & Cell Physiology*,

56(1), 16–27. doi: 10.1093/pcp/pcu158

Oksanen, F.J. Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan

McGlinn, Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry H.

Stevens, Eduard Szoecs and Helene Wagner (2019). vegan: Community ecology package.

R package version 2.5-5. https://CRAN.R-project.org/package=vegan

Onkokesung, N., Reichelt, M., van Doorn, A., Schuurink, R. C., & Dicke, M. (2016).

Differential Costs of Two Distinct Resistance Mechanisms Induced by Different Herbivore

Species in Arabidopsis. Plant physiology, 170(2), 891–906. doi:10.1104/pp.15.01780

Pandey, P., Ramegowda, V., & Senthil-Kumar, M. (2015). Shared and unique responses of

plants to multiple individual stresses and stress combinations: physiological and molecular

mechanisms. *Frontiers in Plant Science*, *6*. doi: 10.3389/fpls.2015.00723

Patton, M. F., Bak, A., Sayre, J. M., Heck, M. L., & Casteel, C. L. (2020). A polerovirus, *Potato*

leafroll virus, alters plant–vector interactions using three viral proteins. *Plant, Cell &*

Environment, *43*(2), 387–399. doi: 10.1111/pce.13684

Petek, M., Rotter, A., Kogovšek, P., Baebler, S., Mithöfer, A., & Gruden, K. (2014). *Potato virus*

Y infection hinders potato defence response and renders plants more vulnerable to Colorado

potato beetle attack. *Molecular ecology*, 23(21), 5378–5391. doi: org/10.1111/mec.12932

Ramírez-Carrasco, G., Martínez-Aguilar, K., & Alvarez-Venegas, R. (2017). Transgenerational

defense priming for crop protection against plant pathogens: A hypothesis. *Frontiers in*

Plant Science, *8*. doi: 10.3389/fpls.2017.00696

Rao, X., X. Huang, Z. Zhou, and X. Lin. 2013. An improvement of the $2^{\wedge}$(–delta delta CT)

method for quantitative real-time polymerase chain reaction data analysis. *Biostatistics,*

bioinformatics and biomathematics, *3*(3), 71–85.

Seguel, A., Jelenska, J., Herrera-Vásquez, A., Marr, S. K., Joyce, M. B., Gagesch, K. R., …

Holuigue, L. (2018). PROHIBITIN3 forms complexes with ISOCHORISMATE

SYNTHASE1 to regulate stress-induced salicylic acid biosynthesis in *Arabidopsis*. *Plant Physiology*, *176*(3), 2515–2531. doi: 10.1104/pp.17.00941

Selim, S., Sanssené, J., Rossard, S., & Courtois, J. (2017). Systemic induction of the defensin and phytoalexin pisatin pathways in pea (*Pisum sativum*) against *Aphanomyces euteiches* by acetylated and nonacetylated oligogalacturonides. *Molecules*, *22*(6), 1017. doi: 10.3390/molecules22061017

Serova, T. A., Tsyganova, A. V., Tikhonovich, I. A., & Tsyganov, V. E. (2019). Gibberellins inhibit nodule senescence and stimulate nodule meristem bifurcation in pea (*Pisum sativum* L*.*). *Frontiers in Plant Science*, *10*. doi: 10.3389/fpls.2019.00285

Shi, J.-H., Sun, Z., Hu, X.-J., Jin, H., Foba, C. N., Liu, H., … Wang, M.-Q. (2019). Rice defense responses are induced upon leaf rolling by an insect herbivore. *BMC Plant Biology*, *19*(1), 514. doi: 10.1186/s12870-019-2116-0

Shikano, I. (2017). Evolutionary ecology of multitrophic interactions between plants, insect herbivores and entomopathogens. *Journal of Chemical Ecology*, *43*(6), 586–598. doi: 10.1007/s10886-017-0850-z

Singh, N., Swain, S., Singh, A., & Nandi, A. K. (2018). AtOZF1 positively regulates defense against bacterial pathogens and npr1-independent salicylic acid signaling. *Molecular Plant-Microbe Interactions*, *31*(3), 323–333. doi: 10.1094/MPMI-08-17-0208-R

Stam, R., Mantelin, S., McLellan, H., & Thilliez, G. (2014). The role of effectors in nonhost resistance to filamentous plant pathogens. *Frontiers in Plant Science*, *5*. doi: 10.3389/fpls.2014.00582

Su, Q., Mescher, M.C., Wang, S., Chen, G., Xie, W., Wu, Q., Wang, W., & Zhang, Y. *Tomato yellow leaf curl virus* differentially influences plant defence responses to a vector and a

non-vector herbivore. (2016). *Plant, Cell & Environment*, 39(3), 597-607. doi: 10.1111/pce.12650

Suzuki, N. (2016). Hormone signaling pathways under stress combinations. *Plant Signaling & Behavior*, *11*(11), e1247139. doi: 10.1080/15592324.2016.1247139

Thaler, J. S., Agrawal, A. A., & Halitschke, R. (2010). Salicylate-mediated interactions between pathogens and herbivores. *Ecology*, *91*(4), 1075–1082. doi: 10.1890/08-2347.1

Thaler, J. S., Humphrey, P. T., & Whiteman, N. K. (2012). Evolution of jasmonate and salicylate signal crosstalk. *Trends in Plant Science*, *17*(5), 260–270. doi: 10.1016/j.tplants.2012.02.010

Tran, H. S., You, M. P., & Barbetti, M. J. (2018). Expression of defence-related genes in stems and leaves of resistant and susceptible field pea (*Pisum sativum*) during infection by *Phoma koolunga*. *Plant Pathology*, *67*(1), 156–166. doi: 10.1111/ppa.12709

van Geem, M., Gols, R., Raaijmakers, C. E., & Harvey, J. A. (2016). Effects of population-related variation in plant primary and secondary metabolites on aboveground and belowground multitrophic interactions. *Chemoecology*, *26*(6), 219–233. doi: 10.1007/s00049-016-0222-0

Vos, I. A., Moritz, L., Pieterse, C. M. J., & Van Wees, S. C. M. (2015). Impact of hormonal crosstalk on plant resistance and fitness under multi-attacker conditions. *Frontiers in Plant Science*, *6*. doi: 10.3389/fpls.2015.00639

Wang, K., Senthil-Kumar, M., Ryu, C.-M., Kang, L., & Mysore, K. S. (2012). Phytosterols play a key role in plant innate immunity against bacterial pathogens by regulating nutrient efflux into the apoplast. *Plant Physiology*, *158*(4), 1789–1802. doi: 10.1104/pp.111.189217

War, Abdul Rashid, Paulraj, M. G., Ahmad, T., Buhroo, A. A., Hussain, B., Ignacimuthu, S., &

Sharma, H. C. (2012). Mechanisms of plant defense against insect herbivores. *Plant*

Signaling & Behavior, *7*(10), 1306–1320. doi: 10.4161/psb.21663

Wasternack, C., & Hause, B. (2013). Jasmonates: biosynthesis, perception, signal transduction

and action in plant stress response, growth and development. An update to the 2007 review

in Annals of Botany. *Annals of Botany*, *111*(6), 1021–1058. doi: 10.1093/aob/mct067

Yang, D.-L., Yang, Y., & He, Z. (2013). Roles of plant hormones and their interplay in rice

immunity. *Molecular Plant*, *6*(3), 675–685. doi: 10.1093/mp/sst056

Yang, D.-L., Yao, J., Mei, C.-S., Tong, X.-H., Zeng, L.-J., Li, Q., … He, S. Y. (2012). Plant

hormone jasmonate prioritizes defense over growth by interfering with gibberellin

signaling cascade. *Proceedings of the National Academy of Sciences of the United States of*

America, *109*(19), E1192-1200. doi: 10.1073/pnas.1201616109

Yergaliyev, T. M., Nurbekova, Z., Mukiyanova, G., Akbassova, A., Sutula, M., Zhangazin, S.,

… Omarov, R. T. (2016). The involvement of ROS producing aldehyde oxidase in plant

response to Tombusvirus infection. *Plant Physiology and Biochemistry: PPB*, *109*, 36–44.

doi: 10.1016/j.plaphy.2016.09.001

Zdunek-Zastocka, E., & Sobczak, M. (2013). Expression of *Pisum sativum PsAO3* gene, which

encodes an aldehyde oxidase utilizing abscisic aldehyde, is induced under progressively but

not rapidly imposed drought stress. *Plant Physiology and Biochemistry: PPB*, *71*, 57–66.

doi: 10.1016/j.plaphy.2013.06.027

Zhou, S., Lou, Y.-R., Tzin, V., & Jander, G. (2015). Alteration of plant primary metabolism in

response to insect herbivory. *Plant Physiology*, *169*(3), 1488–1498. doi:

10.1104/pp.15.01405

Zhu, F., Poelman, E. H., & Dicke, M. (2014). Insect herbivore-associated organisms affect plant

responses to herbivory. *New Phytologist*, *204*(2), 315–321. doi: [10.1111/nph.12886](10.1111/nph.12886)

DATA ACCESSIBILITY

610   Data are publicly available from Figshare: https://doi.org/10.6084/m9.figshare.14227097.v1

AUTHOR CONTRIBUTIONS

613   S.B. and D.W.C. conceived the ideas and designed the methodology; S.B., R.E.C., S.B. and

614   C.L.C. collected the data; R.E.C., S.B., C.L.C and D.W.C. analyzed the data; all authors

615   contributed critically to the drafts and gave final approval for publication.

Figure legends

Figure 1. Schematic representation of 2×3 factorial design. Green-colored aphids indicate sham (non-infective) *A. pisum*, while blue-colored aphids indicate PEMV-infective *A. pisum*. Slashes indicate order of *S. lineatus* treatments (*S. lineatus* first, *A. pisum* first, or no *S. lineatus*).

Figure 2. Relative transcript accumulation of plant hormone biosynthesis genes associated with four hormonal signaling pathways: (A) *ICS1* (salicylic acid), (B) *LOX2* (jasmonic acid), (C) *AOX3* (abscisic acid), and (D) *GA2ox* (gibberellic acid) following attack with various combinations of *S. lineatus*, *A. pisum*, and PEMV. Within each panel, bars separated by a different letter were significant different based on MANOVA (Tukey HSD, $\alpha = 0.05$). Bar height and error bars indicate marginal mean and standard error of the regression coefficient for each respective treatment.

Figure 3. Relative transcript accumulation of plant defense response transcripts: (A) *PR1,* (B) *DDR230,* and (C) *PsLectin* following attack with various combinations of *S. lineatus*, *A. pisum*, and PEMV. Within each panel, bars separated by a different letter were significant different based on MANOVA (Tukey HSD, $\alpha = 0.05$). Bar height and error bars indicate marginal mean and standard error of the regression coefficient for each respective treatment.

Figure 4. (A) Salicylic acid, (B) jasmonic acid, and (C) abscisic acid phytohormone levels in *P. sativum* plants following attack with various combinations of *S. lineatus*, *A. pisum*, and PEMV. Within each panel, bars not connected by the same letter are significantly different (Tukey HSD, $\alpha = 0.05$). Bar height and error bars indicate marginal mean and standard error of the regression coefficient for each respective treatment.

Figure 5. Nutritional analysis (total amino acid) in *P. sativum* following attack with various combinations of *S. lineatus*, *A. pisum*, and PEMV. *S. lineatus* increase total amino acid concentration in plants (GLM, $\chi2 = 9.194$, $P = 0.01$). There was no "sham-none" treatment

642   combination so that could not be estimated. Within each panel, bars not connected by the same

643   letter are significantly different (Tukey HSD, $\alpha = 0.05$). Bar height and error bars indicate

644   marginal mean and standard error of the regression coefficient for each respective treatment.

646 **Fig. 1**

647
648

649   **Fig. 2**

652   **Fig. 3**

  Fig. 4

657   **Fig. 5**