Peak shifts and extinction under sex-specific selection

Stephen P. De Lisle

Evolutionary Ecology Unit, Department of Biology

Lund University

Sölvegatan 37 223 62

Lund, Sweden

Email: stephen.de_lisle@biol.lu.se

Keywords: adaptive landscape, sexual dimorphism, stabilizing selection, microevolution,

macroevolution

Abstract

A well-known property of sexual selection combined with a cross sex genetic correlation

(r_{mf}), is that it can facilitate a peak shift on the adaptive landscape. How do these diversifying

effects of sexual selection + r_{mf} balance with the constraints imposed by such sexual

antagonism, to affect macroevolution of sexual dimorphism? Here, I extend existing

quantitative genetic models of evolution on complex adaptive landscapes. Beyond recovering

classical predictions for the conditions promoting a peak shift, I show that when r_{mf} is

moderate to strong, relatively weak sexual selection is required to induce a peak shift in

males only. Increasing the strength of sexual selection leads to a sexually-concordant peak

shift, suggesting that macroevolutionary rates of sexual dimorphism may be largely

decoupled from the strength of within-population sexual selection. Accounting explicitly for

demography further reveals that sex-specific peak shifts may be more likely to be successful

than concordant shifts in the face of extinction, especially when natural selection is strong.

An overarching conclusion is that macroevolutionary patterns of sexual dimorphism are

unlikely to be readily explained by within-population estimates of selection or constraint

alone.

Introduction

A longstanding dilemma in evolutionary biology lies in understanding how populations can evolve from one phenotypic optimum to another. When a population is under net-stabilizing selection and in the vicinity of the optimum trait value (a 'peak'), selection will pull the population towards the nearby optimum (1), leaving alternative optima seemingly inaccessible (2). For a peak shift to occur, some force must allow a population mean phenotype to transcend the pull of the nearby optimum, and cross a fitness valley to climb a peak beyond (3-6).

Candidate phenomena that may facilitate crossing a natural-selection valley include genetic drift, a change in the environment, or sexual selection. Wright famously proposed a key role for drift in valley crossing (3, 5, 6), although drift alone will only facilitate such a crossing with exceptional rarity, requiring very weak selection (a shallow valley) and small population size (7-9). A change in the environment seems a likely explanation, although also an incomplete one as rates of phenotypic macroevolution do not seem to be coupled to environmental upheaval in any obvious way (10, 11). Finally, sexual selection can readily pull a population off a natural-selection fitness optimum (12), resulting in a peak shift even across quite deep valleys (13). If only one sex is under significant sexual selection, whether or not a peak shift occurs in the other sex depends on the magnitude of the cross sex genetic correlation (r_{mf}) for the trait (13). If r_{mf} is high enough, sexual selection in one sex will pull both sexes off of their optimum, leading to a peak shift in both sexes. More generally, directional selection on any trait can induce a peak shift in other, genetically correlated traits that themselves reside on an optimum (9). In this way, sexual selection coupled with cross-sex genetic correlations has been proposed as a likely mechanism facilitating peak shifts and thus promoting the origin of diversity (14).

Two open questions remain in light of sexual selection's likely role in driving peak

shifts. First, how do sexual selection-induced peak shifts manifest evolution of sexual

dimorphism, when adaptive landscapes are complex; That is, when sexually dimorphic and

sexually-monomorphic trait optima exist, what are the conditions that promote or constrain

likelihood of a peak shift to each optimum? Second, how do we reconcile the diversifying

effects of sexual selection and r_{mf} with the constraining effects that r_{mf} is expected to have on

male and female adaptation? Put another way, although we know sexual selection + r_{mf} can

lead to a peak shift, we also know that this condition of sexual conflict constrains adaptation.

In this note I extend Lande's (12) model of sexual and natural selection towards a

single optimum to the case of multiple optima. This model is similar to one analyzed by

Lande and Kirkpatrick (13), but is agnostic to female preference evolution, a major focus of

their model. My analysis reveals two underappreciated features of peak shift models. First,

large regions of parameter space exist in which relatively weak sexual selection is required to

induce a sex-specific peak shift; strong sexual selection is expected to lead to evolution along

a line of sexual monomorphism at the macroevolutionary scale. Second, the latter type of

peak shift is severely limited by extinction. Although these effects are consistent with results

of previous work (9, 13), to my knowledge they have rarely been appreciated, particularly in

the context of macroevolution of sexual dimorphism.

The Model – Two-sex 'Twin Peaks'

Lande (12) considered a model of evolution of sexual dimorphism by natural and

sexual selection, in which natural selection favors a single optimum value for the male trait z

and female trait y. His model of natural selection corresponds to a single adaptive peak on

the two-sex adaptive landscape (where $\overline{W} = \overline{W}_m \overline{W}_f$, Figure S1A). Directional sexual

selection acts independently of natural selection to redistribute fitness across individuals

without changing mean absolute fitness. This scenario represents a constant mating bias for

one sex, arising from, e.g., female mating preference or male-male competition.

We can expand natural selection fitness function of the Lande model to consider a

scenario where more than one optimum for a trait exists, using a mixture of Gaussian

functions (15) termed the 'Twin Peak' model by Price et al. (9). For males, this leads to the

following function relating population mean fitness $\overline{W}_m$ to population mean phenotype $\bar{z}$:

$$\overline{W}_m \propto \exp\left\{-\left(\bar{z} - \theta_{z,1}\right)^2\left(\omega_{z,1} + P_z\right)^{-1}\right\} \qquad 1$$

$$+ \exp\left\{-\left(\bar{z} - \theta_{z,2}\right)^2\left(\omega_{z,2} + P_z\right)^{-1}\right\}$$

where $\theta_{z,1}$ and $\theta_{z,2}$ are two phenotypic optima, P_z is the phenotypic variance, and $\omega_{z,1}$ $\omega_{z,2}$

are the strengths of stabilizing selection (assumed equal throughout). Such an adaptive

landscape corresponds to an individual fitness surface (16) with two separate optima. This

model is bimodal, with optima in the vicinity of θ, for $2\theta^2 > \omega + P$. The adaptive

landscape for a scenario where only one sex has multiple optima is illustrated in Figure S1B.

If instead we assume females also have multiple optima, using an expression analogous to

equation 1 but relating $\overline{W}_f$ to the female mean trait value $\bar{y}$, we obtain the 'Two-sex Twin

Peak' model, illustrated in Figure S1C and Figure 1.

Given $\beta_m = \nabla_m ln\ \overline{W}_m + c_m$ (12), where ∇_m is the gradient $\partial/\partial\bar{z}$ and c_m is sexual

selection arising from some mating bias independent of the strength of natural selection, we

can define β_m under the twin peak model in equation 1 as:

$$\beta_m = \overline{W}_m^{-1}\left[2\frac{(\theta_{z,1}-\bar{z})}{(\omega_{z,1}+P_z)}\exp\left\{-\frac{(\theta_{z,1}-\bar{z})^2}{(\omega_{z,1}+P_z)}\right\} + 2\frac{(\theta_{z,2}-\bar{z})}{(\omega_{z,2}+P_z)}\exp\left\{-\frac{(\theta_{z,2}-\bar{z})^2}{(\omega_{z,2}+P_z)}\right\}\right] + c_m \qquad 2$$

With an analogous expression for β_f. For simplicity and consistency with past work, I focus

on the scenario of sexual selection in males only ($c_f = 0$, $c_m \geq 0$). Evolution of the male

and female mean phenotypes, $\bar{z}$ and $\bar{y}$, depends not only on selection within each sex but also

on correlated response to selection in the other sex mediated by the cross sex genetic

correlation r_{mf}:

$$\Delta\bar{z} = \frac{1}{2}\beta_m G_m + \frac{1}{2}\beta_f B$$

$$\Delta\bar{y} = \frac{1}{2}\beta_f G_f + \frac{1}{2}\beta_m B$$

128   where G_m and G_f are the male and female genetic variances and $r_{mf} = B/\sqrt{(G_m G_f)}$.

130   *Population growth*

131   Population growth can be considered a function of mean fitness, such that

132   maladaptation carries demographic cost. Following Lande (12), we can describe change in

133   census population size N as

$$N_{t+1} = k\,\bar{W}_m \bar{W}_f N_t$$

where k is constant or function defining per capita birth rates. For simplicity, I assume k as a

constant corresponding to density-independent population growth. In equation 4, adaptation

in both sexes contributes to population growth, corresponding to a biological scenario where,

for example, parental care is shared across the sexes. Alternatively, in many species male

adaptation may contribute little to population growth rates, and we can instead define change

in census size as

$$N_{t+1} = k\,\bar{W}_f N_t$$

140   which assumes that there are always at least enough males to fertilize females in the

141   population. Equations 4 and 5 represent two ends of a continuum of in which male

142   adaptation may contribute to population growth rates.

Drift

145 The role of random genetic drift has been considered extensively in peak shift models, and I

146 address it only briefly here for completeness. The rate of evolution by genetic drift will be

147 proportional to $\mathbf{G}/N_e$ (17), and in a two-sex population,

$$N_e = \frac{4N_m N_f}{N_m + N_f}$$

149 (18). Drift occurs after natural selection such that each sex is represented proportional to

150 their own mean fitness,

$$N_m \propto \frac{N}{2}\overline{W}_m, \quad N_f \propto \frac{N}{2}\overline{W}_f$$

where N represents the number of individual (e.g., zygotes) before selection, with N/2 of

each sex under a Fisherian primary sex ratio (1).

I used numerical simulations to explore how sexual selection and r_{mf} influence peak

shifts and extinction. In order to understand how sexual dimorphism evolves from an

ancestral condition of sexual monomorphism, each simulation was started with the

population mean at a sexually-monomorphic optimum of $\bar{z} = \bar{y} = \theta_{z,2} = \theta_{y,2} = -2$, with

$\theta_{z,1} = \theta_{y,1} = 2$. In this scenario there are three unoccupied optima, two of which are

accessible deterministically under positive sexual selection and r_{mf}; one new optimum is

along a line of male-female isometry, and represents a shift from one sexually monomorphic

peak to another (henceforth, 'sexually-monomorphic', or concordant, peak shift). A second

accessible peak involves a male-only shift, and so represents a 'sexually dimorphic' peak

shift. All ω were assumed equal, and phenotypic variance was set at unity and G = P/2,

assumed constant. Starting N was set to 10,000; population growth was assumed as in

equation 3 or 4, although population size was capped at 10,000 with an arbitrary extinction

threshold of N = 20. The growth constant k was assumed 1.05. Thus growth was exponential

and density independent up to the upper bound. Assuming unbounded growth, changing

population size, k, or genetic parameters did not change qualitative conclusions. Complete R

script is provided as supplemental material.

Results

When there is only a single optimum, sexual and natural selection jointly determine

the equilibrium trait value for males, which are displaced from their peak proportional to

sexual selection c_m and the strength of stabilizing selection ω (Figure S1D; 12). Female

equilibrium trait values are unaffected by the strength of sexual selection in males, but their

path towards their optimum is affected by r_{mf} (Figure S1D). When multiple optima exist, the

combination of sexual selection and r_{mf} determines if a peak shift occurs and which

alternative optima is reached. Two scenarios are shown in Figure 1, illustrating that

increasing the strength of sexual selection while holding r_{mf} constant, and vice versa, have

similar effects. Unlike the single optimum case, when multiple optima exist, female

equilibrium trait values depend on sexual selection in males and r_{mf}; for a peak shift to occur

in females, their product must be high enough for female mean phenotype to be displaced

beyond the critical value required for a peak shift. Note that a 'monomorphic peak' shift

(Figure 1A) still entails some sexual dimorphism in the traits will be observed at equilibrium

(because males will be displaced optimum), although the equilibrium magnitude of sexual

dimorphism will be far higher when a peak shift is restricted to males only (a 'dimorphic

shift'; Figure 1). Valley crossing carries substantial demographic costs, illustrated in Figure

S2.

190   Whether a peak shift occurs for males depends only on the strength of sexual

selection c_m. However, whether the shift is dimorphic (males only) or monomorphic depends

also on r_{mf}, such that for wide ranges of moderate to strong genetic correlations, increasing

the strength of sexual selection while holding r_{mf} constant yields first a transition from no

peak shift to a dimorphic shift, and then to a monomorphic shift, and then in some cases

extinction (Figure 2, left panels). Increasing the strength of stabilizing selection (smaller ω)

exacerbates these effects. These effects hold regardless of the demographic model assumed,

although extinction is far more likely to be observed when population growth rates are a

function of both male and female adaptation (Figure 2, right panels). These conclusions

remain qualitatively unchanged when accounting for genetic drift, although the outcomes

become probabilistic due to the stochastic nature of drift in the fitness valley (Figure S3).

Discussion

Although the broad factors that may possibly affect peak shifts are well understood, how

these factors jointly determine the probability of actual peak shifts remains a puzzle (6). Sex-

specific selection, in particular sexual selection, provides one example of the uncertainties

around peak shifts. Sexual selection and genetic correlations may be key in facilitating peak

shifts (12, 13), yet this diversifying effect (14) is seemingly at odds with the constraining role

that such a condition must also manifest (12, 19). The point of this paper is to explore how

sexual selection and r_{mf} interact to influence the likelihood of a peak shift occurring, as well

as which alternative optima are most likely to be reached. This modest revisit to classical

peak shift models yields two underappreciated phenomenon that have important implications

for the evolution of sexual dimorphism.

First, although increasing the strength of sexual selection increases the likelihood of a

peak shift, whether that peak shift results in substantial sexual dimorphism depends

somewhat counter intuitively on the magnitude of sexual selection.   For wide ranges of r_{mf},

weaker sexual selection produces a peak shift in males only; increasing sexual selection

further increases the correlated response in females to the point where females also shift

peaks (see also figure 4A in Price et al. 1993). The implication is that when sexually

monomorphic optima exist, strong sexual selection is expected to lead to peak shifts along a

line of male-female isometry when r_{mf} is non-zero. Only when sexual selection is weak

enough in magnitude that its interaction with r_{mf} fails to push females beyond the critical

displacement required for a peak shift, yet strong enough to directly push males beyond their

own critical displacement, will dimorphic optima be reached. This suggests that

macroevolutionary patterns of sexual dimorphism may depend far more on nuances of the

two-sex adaptive landscape than on the magnitude of within-population sexual selection.

Second, the demographic cost of valley crossing affects which peaks can be reached.

When population growth rates depend on adaptation in both sexes, or when natural selection

is strong, sex-specific peak shifts are more likely to occur successfully in the face of

extinction than are concordant ('monomorphic') shifts. This demographic effect is the result

of higher demographic costs for crossing a valley in both sexes. When population growth

rates depend only on female adaptation, valley crossing in males-only carries no demographic

cost, and so may be more likely to be observed. These effects are amplified under strong

selection (deep valleys), and indicate that extinction-generated survivorship bias (20) may

make sex-specific (dimorphic) peak shifts more likely to be observed than monomorphic

shifts.

These two features of sex-specific peak shifts may explain several puzzling phenomena

observed in macroevolutionary studies of sexual dimorphism. First, a large number of

studies have investigated the link between proxies for the strength of within-population

sexual selection and among-lineage patterns of sexual dimorphism. Often, these studies find

only weak relationships between these sexual-selection proxies and the magnitude of sexual

dimorphism (21-24). This finding of little correlation between the magnitude (yet some

correlation between the sign; 25) of sexual dimorphism and sexual selection proxies is

consistent with the results presented here. Concomitantly, any association between r_{mf} and

the strength of sexual selection within populations (as some data implies could be the case;

26, 27) could have important macroevolutionary consequences.

Further, when male adaptation contributes little to population growth rate, this may

lead to a male bias in macroevolutionary rate if male valleys are easier to cross without

extinction. In many clades, males are observed to exhibit higher rates of body size evolution

than females (28-31). Although the opposite (higher female rates) is observed in some groups

(32), it does appear to be rarer.

Evolution on real adaptive landscapes, with mixtures of peaks of varying height,

varying distance from each other, will surely be more complex than the model presented

here. None the less, whether a peak shift to a dimorphic or monomorphic peak occurs will

depend in part on an interaction between sexual selection and genetic correlations that may

decouple the strength of sexual selection from the magnitude of resulting sexual dimorphism

at equilibrium, and crossing two valleys at once may be demographically costly.

Acknowledgements I thank David Punzalan, Locke Rowe, Erik Svensson, Beatriz Willink,

Masahito Tsuboi, and Benjamin Jarrett for discussion and/or comments on the manuscript.

Funding was provided by grants from the Swedish Research Council (VR registration

number 2019- 03706) and Kungl. Fysiografiska Sällskapet i Lund to S.P.D.

Figure 1. Peak shifts induced by sexual selection in the Two-sex Twin Peaks model.

Panel **A** shows two deterministic trajectories, starting from the lower left optimum,

corresponding to two different values of the cross-sex genetic correlation r_{mf}. Although

sexual selection is strong enough to induce a peak shift, which new optimum reached

depends upon the value of r_{mf}. Panel **B** illustrates 20 replicates of stochastic evolution (i.e.,

with drift) under the same parameter values as A and assuming female demographic

dominance ($N_{t+1} = k\,\overline{W}_f\,N_t$). Round circles illustrate the mean values after 10000

generations of evolution; extinction events are denoted with a crossed circle. Panel **C**

contrasts trajectories under two strengths of sexual selection c_m, with otherwise identical

parameter values. Panel **D** illustrates stochastic evolution under the same parameter values as

C. White dashed line illustrates the line of sexual monomorphism, for reference. In A and B,

$c_m = 0.2$; B and C $r_{mf} = 0.8$; other parameter values as described in text.

Figure 2. Deterministic outcomes as a function of sexual selection and r_{mf}. Panels show

deterministic (i.e. no drift) outcomes of 1000 generations of evolution, calculated

numerically, corresponding to three different strengths of stabilizing selection. Left panels

assume female demographic dominance, right panels assume male and female fitness both

283   determine population growth rate. 'Dimorphic shift' refers to a male-only peak shift,

284   'Monomorphic shift' refers to a peak shift in both sexes (illustrated in Fig 2A). No peak shift

285   means the lineage remains in the vicinity of the ancestral optimum. Small gaps between

286   regions represent plotting limitations of overlaid contour plots.

Literature Cited

1.   Fisher RA. The genetical theory of natural selection. Oxford: The Clarendon press;

1930. xiv, 272 p p.

2.   Frank SA. Wright's Adaptive Landscape Versus Fisher's Fundamental Theorem. In:

Svensson EI, Calsbeek R, editors. The Adaptive Landscape in Evolutionary Biology: Oxford

University Press; 2013.

3.   Wright S. The roles of mutation, inbreeding, crossbreeding and selection in evolution.

Proceedings of the Sixth Annual Conference of Genetics. 1932;1:356-66.

4.   Coyne JA, Orr HA. Speciation. Sunderland, MA: Sinaur; 2004.

5.   Coyne JA, Barton NH, Turelli M. Perspective: a critique of Sewall Wright's shifting

balance theory of evolution. Evolution. 1997;51:643-71.

6.   Goodnight CJ. Wright's Shifting Balance Theory and Factors Affecting the

Probability of Peak Shifts. In: Svensson EI, Calsbeek R, editors. The Adaptive Landscape in

Evolutionary Biology: Oxford University Press; 2013.

7.   Lande R. Expected time for random drift of a population between stable phenotypic

states. Proceedings of the National Academy of Sciences. 1985;82:7641-5.

8.   Barton NH, Charlesworth B. Genetic revolutions, founder effects and speciation.

Annual Review of Ecology and Systematics. 1984;15:133-64.

9.   Price T, Turelli M, Slatkin M. Peak shifts produced by correlated response to

selection. Evolution. 1993;47(1):280-90.

10.   Uyeda JC, Hansen TF, Arnold SJ, Pienaar J. The million year wait for

macroevolutionary bursts. Proceedings of the National Academy of Sciences of the United

States of America. 2011;108:15908-13.

11.   Gingerich PD. Rates of evolution on the time scale of the evolutionary process.

Genetica. 2001;112:127-44.

12.   Lande R. Sexual dimorphism, sexual selection, and adaptation in polygenic

characters. Evolution. 1980;34(2):292-305.

13.   Lande R, Kirkpatrick M. Ecological speciation by sexual selection. Journal of

Theoretical Biology. 1988;133:85-98.

14.   Bonduriansky R. Sexual selection and sexual conflict as engines of ecological

diversification. American Naturalist. 2011;178:729-45.

15.   Felsenstein J. Excursions along the interface between disruptive and stabalizing

seleciton. Genetics. 1979;93:773-95.

16.   Arnold SJ, Pfrender ME, Jones AG. The adaptive landscape as a conceptual bridge

between micro- and macroevolution. Genetica. 2001;112/113:9-32.

17.   Lande R. Natural selection and random genetic drift in phenotypic evolution.

Evolution. 1976;30:314-34.

18.   Wright S. Evolution in Mendelian populations. Genetics. 1931;16:97-159.

19.   Matthews G, Hangartner S, Chapple DG, Connallon T. Quantifying maladaptation

during the evolution of sexual dimorphism. Proceedings of the Royal Society of London

Series B. 2019;286:20191372.

20.   De Lisle SP, Punzalan D, Rollinson N, Rowe L. Extinction and the temporal

distribution of macroevolutionary bursts. Journal of Evolutionary Biology. 2020;in press.

21.   Cox RM, Skelly SL, John-Alder HB. A comparative test of adaptive hypotheses for

sexual size dimorphism in lizards. Evolution. 2003;57:1653-69.

22.   Shine R. Ecological causes for the evolution of sexual dimorphism: a review of the

evidence. Quarterly Review of Biology. 1989;64(4):419-61.

23.   Cox RM, Calsbeek R. Sexually antagonistic selection, sexual dimorphism, and the

resolution of intralocus sexual conflict. American Naturalist. 2009;173(2):176-87.

24.   Nali RC, Zamudio KR, Haddad CFB, Prado CPA. Size-dependent selective

mechanisms on males and females and the evolution of sexual size dimorphism in frogs. The

American Naturalist. 2014;184:727-40.

25.   Janicke T, Häderer IK, Lajeunesse MJ, Anthes N. Darwinian sex roles confirmed

across the animal kingdom. Science Advances. 2016;2:e1500983.

26.   Poissant J, Wilson AJ, Coltman DW. Sex-specific genetic variance and the evolution

of sexual dimorphism: a systematic review of cross-sex genetic correlations. Evolution.

2009;64(1):97-107.

27.   Bonduriansky R, Rowe L. Sexual selection, genetic architecture, and the condition

dependence of body size and shape in the sexually dimorphic fiy *Prochyliz xanthostoma*

(Diptera: Piophilidae). Evolution. 2005(59):138-51.

28.   Rensch B. Evolution Above the Species Level. New York: Columbia University

Press; 1960. 490 p.

29.   Fairbairn DJ. Allometry for sexual size dimorphism: Pattern and process in the

coevolution of body size in males and females. Annual Review of Ecology and Systematics.

1997;28:659-78.

30.   Stephens PR, Wiens JJ. Evolution of sexual size dimorphisms in emydid turtles:

ecological dimorphism, Rensch's rule, and sympatric divergence. Evolution. 2009;63:910-25.

31.   Colwell RK. Rensch's rule crosses the line: Convergent allometry of sexual size

dimorphism in hummingbirds and flower mites. American Naturalist. 2000;156(5):495-510.

32.   De Lisle SP, Rowe L. Correlated evolution of allometry and sexual dimorphism

across higher taxa. American Naturalist. 2013;182(5):630-9.

384
385

Figure S1. Models of stabilizing selection in two sexes. Panels illustrate the adaptive

landscape, the function $\overline{W}_m\overline{W}_f$. In Lande's (1980) original model of evolution of sexual

dimorphism by natural and sexual selection, a single multivariate optimum, corresponding to

one optimum for the male trait and one for the female trait, is assumed (**A**). The 'Twin Peaks'

model extends this to consider two optima in one trait (sex), with only a single optimum for

the other trait (sex); (**B**). Extending the Twin Peak model to both sexes results in two optima

for each trait, or four multivariate optima on the two-sex adaptive landscape (**C**). As

parameterized in this manuscript, the Two-sex Twin Peak model contains two sexually

monomorphic optima, and two sexually dimorphic optima. Of interest is when and how peak

shifts from an ancestral monomorphic optimum occur. In this model sexual selection leads to

a displacement of males from their optimum (**C**), the magnitude of which is proportional to

the strength of stabilizing natural selection. Two trajectories are shown starting from the

optimum; one where $r_{mf} = 0$ (blue) and one where $r_{mf} = 0.9$ (blue).

414

Figure S2 Demographic cost of valley crossing. Figures show population size from

trajectories plotted in Figure 1A and 1C. Panel A contrasts two values of the cross-sex

genetic correlation, r_{mf} = 0.8 (red line) or r_{mf} = 0.9 (blue line). Panel B contrasts two values

of the strength of sexual selection, cm = 0.2 (red line) or cm = 0.3 (blue line). Demographic

costs of a monomorphic peak shift, corresponding to the blue line in both panels, are higher

than for a dimorphic (male-only peak shift). Other parameter values are as described in

Figure 1.

Figure S3 Stochastic outcomes as a function of sexual selection and r_{mf}. Panels show

stochastic (i.e., evolution under both selection and drift) outcomes of 1000 generations of

evolution, calculated numerically, corresponding to the scenario of strong natural selection

and female demographic dominance illustrated in Figure 2C. Thus, qualitative conclusions

on peak shifts remain similar to the deterministic case, and this holds for other parameter

values shown in Figure 2.