Peak shifts and extinction under sex-specific selection

Stephen P. De Lisle

Evolutionary Ecology Unit, Department of Biology

Lund University

Sölvegatan 37 223 62

Lund, Sweden

Email: stephen.de_lisle@biol.lu.se

Keywords: adaptive landscape, sexual dimorphism, stabilizing selection, microevolution,

macroevolution

Abstract

A well-known property of sexual selection combined with a cross sex genetic correlation

(r_{mf}), is that it can facilitate a peak shift on the adaptive landscape. How do these diversifying

effects of sexual selection + r_{mf} balance with the constraints imposed by such sexual

antagonism, to affect macroevolution of sexual dimorphism? Here, I extend existing

quantitative genetic models of evolution on complex adaptive landscapes. Beyond recovering

classical predictions for the conditions promoting a peak shift, I show that when r_{mf} is

moderate to strong, relatively weak sexual selection is required to induce a peak shift in

males only. Increasing the strength of sexual leads to a sexually-concordant peak shift,

suggesting that macroevolutionary rates of sexual dimorphism may be largely decoupled

from the strength of within-population sexual selection. Accounting explicitly for

demography further reveals that sex-specific peak shifts may be more likely to be successful

than concordant shifts in the face of extinction, especially when natural selection is strong.

An overarching conclusion is that macroevolutionary patterns of sexual dimorphism are

unlikely to be readily explained by within-population estimates of selection or constraint

alone.

Introduction

A longstanding dilemma in evolutionary biology lies in understanding how populations can evolve from one phenotypic optimum to another. When a population is under net-stabilizing selection and in the vicinity of the optimum trait value (a 'peak'), selection will pull the population towards the nearby optimum (Fisher 1930), leaving alternative optima seemingly inaccessible (Frank 2013). For a peak shift to occur, some force must allow a population mean phenotype to transcend the pull of the nearby optimum, and cross a fitness valley to climb a peak beyond (Wright 1932, Coyne et al. 1997, Coyne and Orr 2004, Goodnight 2013). The problem is fundamental, because such peak shifts are required for ecological speciation and must also underlie much of what is considered macroevolutionary rate.

Candidate phenomenon that may facilitate a peak shift include genetic drift, a change in the environment (so that the population mean phenotype is no longer on a fitness optimum), or sexual selection. Wright famously proposed a key role for drift in valley crossing (Wright 1932, Coyne et al. 1997, Goodnight 2013), although drift alone will only facilitate such a crossing with exceptional rarity, requiring very weak selection (a shallow valley) and small population size (Barton and Charlesworth 1984, Lande 1985, Price et al. 1993). A change in the environment seems a likely explanation, although also an incomplete one as rates of phenotypic macroevolution do not seem to be coupled to environmental upheaval in any obvious way (Gingerich 2001, Uyeda et al. 2011). Finally, sexual selection can readily pull a population off of a natural-selection fitness optimum (Lande 1980), resulting in a peak shift even across quite deep valleys (Lande and Kirkpatrick 1988). If only one sex is under significant sexual selection, whether or not a peak shift occurs in the other sex depends on the magnitude of the cross sex genetic correlation (r_{mf}) for the trait (Lande and Kirkpatrick 1988). If r_{mf} is high enough, sexual selection in one sex will pull both sexes off of their optimum, leading to a peak shift in both sexes. More generally, directional

selection on any trait can induce a peak shift in other, genetically correlated traits that

themselves reside on an optimum (Price et al. 1993). In this way, sexual selection coupled

with cross-sex genetic correlations has been proposed as a likely mechanism facilitating peak

shifts and thus promoting the origin of diversity (Bonduriansky 2011).

Two open questions remain in light of sexual selection's likely role in driving peak

shifts. First, how do sexual selection-induced peak shifts manifest evolution of sexual

dimorphism, when adaptive landscapes are complex; That is, when sexually dimorphic and

sexually-monomorphic trait optima exist, what are the conditions that promote or constrain

likelihood of a peak shift to each optimum? Second, how do we reconcile the diversifying

effects of sexual selection in combination with r_{mf}, with the constraining effects that r_{mf} is

expected to have on male and female adaptation? Put another way, although we know sexual

selection + r_{mf}, can lead to a peak shift, we also know that this condition of sexual conflict

can reduce population mean fitness, which reduces likelihood of successful valley crossing.

In this note I extend Lande's (1980) model of sexual and natural selection towards a

single optimum to the case of multiple optima, thereby creating a two-sex analog of Price et

al.'s (1993) 'Twin Peak' model, that incorporates sexual selection. This model is similar to

one analyzed by Lande and Kirkpatrick (1988), but is agnostic of female preference

evolution, a major focus of their model. I take a simulation approach that allows explicit

demographics, including drift and extinction, to be readily incorporated. My analysis reveals

two underappreciated features of peak shift models that have important consequences for the

evolution of sexual dimorphism. First, large regions of parameter space exist in which

relatively weak sexual selection is required to induce a sex-specific peak shift; strong sexual

selection is expected to lead to evolution along a line of sexual monomorphism at the

macroevolutionary scale. Second, the latter type of peak shift is severely limited by

extinction. Although these effects are consistent with results of previous work (Lande and

Kirkpatrick 1988, Price et al. 1993), to my knowledge they have rarely been appreciated,

particularly in the context of macroevolution of sexual dimorphism.

The Model – Two-sex 'Twin Peaks'

Lande (1980) considered a model of evolution of sexual dimorphism by natural and

sexual selection, in which natural selection favors a single optimum value for the male trait z

and female trait y. His model of natural selection corresponds to a single adaptive peak on

the two-sex adaptive landscape (where $\overline{W} = \overline{W}_m \overline{W}_f$, Figure 1A). Directional sexual selection

acts independently of natural selection to redistribute fitness across individuals without

changing mean absolute fitness. This scenario represents a constant mating bias for one sex,

arising from, e.g., female mating preference or male-male competition.

We can expand the Lande model, specifically the natural selection fitness function, to

consider a scenario where more than one optimum for a trait exists, using a mixture of

Gaussian functions (Felsenstein 1979) termed the 'Twin Peak' model by Price et al. (1993).

For males, this leads to the following function relating population mean fitness $\overline{W}_m$ to

population mean phenotype $\bar{z}$:

$$
\begin{aligned}
\overline{W}_m \propto\ & \exp\left\{-\left(\bar{z} - \theta_{z,1}\right)^T \left(\omega_{z,1} + \mathrm{P}_z\right)^{-1} \left(\bar{z} - \theta_{z,1}\right)\right\} \\
& + \exp\left\{-\left(\bar{z} - \theta_{z,2}\right)^T \left(\omega_{z,2} + \mathrm{P}_z\right)^{-1} \left(\bar{z} - \theta_{z,2}\right)\right\}
\end{aligned}
\qquad 1
$$

where $\theta_{z,1}$ and $\theta_{z,2}$ are two phenotypic optima, P_z is the phenotypic variance, and $\omega_{z,1}\ \omega_{z,2}$

are the strengths of stabilizing selection (assumed equal throughout). Such an adaptive

landscape corresponds to an individual fitness surface (Arnold et al. 2001) with two separate

optima. This model is bimodal, with optima in the vicinity of θ, for $2\theta^2 > \omega + \mathrm{P}$. The

adaptive landscape for a scenario where only one sex has multiple optima is illustrated in

Figure 1B, a fitness function analogous to that of Price et al (1993) but for separate sexes. If

instead we assume females also have multiple optima, using an expression analogous to

equation 1 but relating $\bar{W}_f$ to the female mean trait value $\bar{y}$, we obtain the 'Two-sex Twin

Peak' model, illustrated in Figure 1C.

127   Given $\beta_m = \nabla_m ln\ \bar{W}_m + c_m$ (Lande 1980), where ∇_m is the gradient operator $\partial/\partial\bar{z}$

128   and c_m is sexual selection arising from some mating bias independent of the strength of

129   natural selection, we can define β_m under the twin peak model in equation 1 as:

$$\beta_m = \ \bar{W}_m^{-1}\left[2\frac{\left(\theta_{z,1} - \bar{z}\right)}{\left(\omega_{z,1} + P_z\right)}\exp\left\{-\frac{\left(\theta_{z,1} - \bar{z}\right)^2}{\left(\omega_{z,1} + P_z\right)}\right\}\right.$$
$$\left. + \ 2\frac{\left(\theta_{z,2} - \bar{z}\right)}{\left(\omega_{z,2} + P_z\right)}\exp\left\{-\frac{\left(\theta_{z,2} - \bar{z}\right)^2}{\left(\omega_{z,2} + P_z\right)}\right\}\right] + \ c_m$$

130   With an analogous expression for β_f. For simplicity and consistency with past work, I focus

131   on the scenario of sexual selection in males only ($c_f = 0,\ c_m \geq 0$). Evolution of the male

132   and female mean phenotypes, $\bar{z}$ and $\bar{y}$, depends not only on selection within each sex but also

133   on correlated response to selection in the other sex mediated by the cross sex genetic

134   correlation r_{mf}:

$$\Delta\bar{z} = \ \frac{1}{2}\beta_m G_m + \ \frac{1}{2}\beta_f B$$

$$\Delta\bar{y} = \ \frac{1}{2}\beta_f G_f + \ \frac{1}{2}\beta_m B$$

135   where G_m and G_f are the male and female genetic variances and $r_{mf} = B/\sqrt{(G_m G_f)}$.

137   *Population growth*

138   Population growth can be considered a function of mean fitness, such that

139   maladaptation carries demographic cost. Following Lande (1980), we can describe change in

140   census population size N as

$$N_{t+1} = k\,\overline{W}_m\overline{W}_f N_t \qquad\qquad 4$$

where k is constant or function defining per capita birth rates. For simplicity, I assume k as a

constant corresponding to density-independent population growth. In equation 4, adaptation

in both sexes contributes to population growth, corresponding to a biological scenario where,

for example, parental care is shared across the sexes. Alternatively, in many species male

adaptation may contribute little to population growth rates, and we can instead define change

in census size as

$$N_{t+1} = k\,\overline{W}_f N_t \qquad\qquad 5$$

which assumes that there are always at least enough males to fertilize females in the

population. Equations 4 and 5 represent two ends of a continuum of in which male

adaptation may contribute to population growth rates.

Drift

The role of random genetic drift has been considered extensively in peak shift models, and I

address it only briefly here for completeness. The rate of evolution by genetic drift will be

proportional to $\mathbf{G}/N_e$ (Lande 1976), where $\mathbf{G}$ is the genetic variance covariance matrix

containing sex-specific genetic variances on the diagonal and the cross sex genetic

covariance ($r_{mf}\sqrt{G_m G_f}$.) on the off diagonal. In a two-sex population, the effective

population size is:

$$N_e = \frac{4N_m N_f}{N_m + N_f} \qquad\qquad 6$$

159   (Wright 1931). Drift occurs after natural selection such that each sex is represented

160   proportional to their own mean fitness,

$$N_m \propto \frac{N}{2} \bar{W}_m, \quad N_f \propto \frac{N}{2} \bar{W}_f \qquad\qquad 7$$

where N represents the number of individual (e.g., zygotes) before selection, with N/2 of

each sex under a Fisherian primary sex ratio (Fisher 1930). Thus, male maladaptation may

contribute to genetic drift even if census population size is purely a function of female mean

fitness.

I used numerical simulations to explore how sexual selection and r_{mf} influence peak

shifts and extinction. In order to understand how sexual dimorphism evolves from an

ancestral condition of sexual monomorphism, each simulation was started with the

population mean at a sexually-monomorphic optimum of $\bar{z} = \bar{y} = \theta_{z,2} = \theta_{y,2} = -2$, with

$\theta_{z,1} = \theta_{y,1} = 2$. In this scenario there are three unoccupied optima, two of which are

accessible deterministically under positive sexual selection and r_{mf}; one new optimum is

along a line of male-female isometry, and represents a shift from one sexually monomorphic

peak to another (henceforth, 'sexually-monomorphic', or concordant, peak shift). A second

accessible peak involves a male-only shift, and so represents a 'sexually dimorphic' peak

shift. Although I have assumed the adaptive landscape is centered along the line of sexual

monomorphism and an ancestral state on a sexually-monomorphic peak, concordant and

dimorphic peak shifts have the same effect on macroevolutionary change in sexual

dimorphism even if the ancestral peak were itself dimorphic. Thus the concepts of a

'monomorphic' and 'dimorphic' peak shift hold regardless of the ancestral state. All ω were

assumed equal, and phenotypic variance was set at unity and G = P/2, corresponding to a

heritability of 0.5 assumed constant. For demographically explicit simulations, starting N

was set to 10,000; population growth was assumed as in equation 3 or 4, although population

size was capped at 10,000 with an arbitrary extinction threshold of N = 20. The growth

constant k was assumed 1.05. Thus growth was exponential and density independent up to

the upper bound. Assuming unbounded growth, changing population size, k, or genetic

parameters did not change qualitative conclusions. Complete R script is provided as

supplemental material.

Results

When there is only a single optimum, sexual and natural selection jointly determine

the equilibrium trait value for males, which are displaced from their peak proportional to

sexual selection c_m and the strength of stabilizing selection ω (Figure 1D; Lande 1980).

Female equilibrium trait values are unaffected by the strength of sexual selection in males,

but their path towards their optimum is affected by r_{mf} (Figure 1D). When multiple optima

exist, the combination of sexual selection and r_{mf} determines if a peak shift occurs and which

alternative optima is reached. Two scenarios are shown in Figure 2, illustrating that

increasing the strength of sexual selection while holding r_{mf} constant, and vice versa, have

similar effects. Unlike the single optimum case, when multiple optima exist, female

equilibrium trait values depend on sexual selection in males and r_{mf}; for a peak shift to occur

in females, their product must be high enough for female mean phenotype to be displaced

beyond the critical value required for a peak shift. Although a peak shift to a 'monomorphic

peak' (Figure 2A) requires that at equilibrium sexual dimorphism in the traits will be

observed (because males will be displaced from the new monomorphic optimum proportional

to c_m), the equilibrium magnitude of sexual dimorphism will be far higher when a peak shift

is restricted to males only (a 'dimorphic shift'; Figure 2). Valley crossing carries substantial

demographic costs, illustrated in Figure S1.

Numerical outcomes after 1000 generations of evolution are shown in Figure 3.

Whether a peak shift occurs for males depends only on the strength of sexual selection c_m.

However, whether the shift is dimorphic (males only) or monomorphic depends also on

r_{mf}, such that for wide ranges of moderate to strong genetic correlations, increasing the

strength of sexual selection while holding r_{mf} constant yields first a transition from no peak

shift to a dimorphic shift, and then to a monomorphic shift, and then in some cases extinction

(Figure 3, left panels). Increasing the strength of stabilizing selection (smaller ω)

exacerbates these effects. These effects hold regardless of the demographic model assumed,

although extinction, instead of a successful peak shift, is far more likely to be observed when

population growth rates are a function of both male and female adaptation (Figure 3, right

panels). These conclusions remain qualitatively unchanged when accounting for genetic

drift, although the outcomes become probabilistic due to the stochastic nature of drift in the

fitness valley (Figure S2).

Discussion

Although the broad factors that may possibly affect peak shifts are well understood, how

these factors jointly determine the probability of actual peak shifts remains a puzzle

(Goodnight 2013). Sex-specific selection, in particular sexual selection, provides one

example of the uncertainties around peak shifts. Sexual selection may be key in facilitating

peak shifts because it can readily pull at least one sex away from a natural selection optimum

(Lande 1980, Lande and Kirkpatrick 1988). Yet, the diversifying role of sexual selection

combined with genetic correlations (Bonduriansky 2011) is seemingly at odds with the

constraining role that such a condition must also manifest; cross-sex genetic correlations slow

evolutionary rate in response to sex-specific selection (Lande 1980, Matthews et al. 2019),

and any lineage that wallows long in a fitness valley may face extinction, even if on its way

to climbing a new peak (De Lisle et al. 2020). The point of this paper is to explore how

sexual selection and r_{mf} interact to influence the likelihood of a peak shift occurring, as well

as which alternative optima are most likely to be reached. This modest revisit to classical

peak shift models yields two underappreciated phenomenon that have important implications

for the evolution of sexual dimorphism.

First, although increasing the strength of sexual selection in males increases the

likelihood of a peak shift, whether that peak shift occurs only in males (leading to the

evolution of sexual dimorphism) or also in females (leading to concordant evolution; sexual

monomorphism) depends somewhat counterintuitively on the magnitude of sexual selection.

For wide ranges of r_{mf}, weaker sexual selection produces a peak shift in males only;

increasing sexual selection further increases the correlated response in females to the point

where females also shift peaks (see also figure 4A in Price et al. 1993). The implication is

that when sexually monomorphic optima exist, strong sexual selection is expected to lead to

peak shifts along a line of male-female isometry when r_{mf} is non-zero. Only when sexual

selection is weak enough in magnitude that its interaction with r_{mf} fails to push females

beyond the critical displacement required for a peak shift, yet strong enough to directly push

males beyond their own critical displacement, will dimorphic optima be reached. This

suggests that macroevolutionary patterns of sexual dimorphism may depend far more on

nuances of the two-sex adaptive landscape – specifically the location of male and female

optima, or ecological niches – than on the magnitude of within-population sexual selection.

The above result suggests sex-specific peak shifts may be somewhat rare, at least

when selection and/or genetic correlations are high in magnitude. However, incorporating

demography changes this picture. When population growth rates depend on adaptation in

both sexes, or when natural selection is strong, sex-specific peak shifts are more likely to

occur successfully in the face of extinction than are concordant ('monomorphic') shifts. This

demographic effect is in part the result of higher demographic costs for crossing a valley in

both sexes, compared to only one. When population growth rates depend only on female

adaptation, valley crossing in males only carries no demographic cost, and so may be more

likely to be observed. These effects are amplified under strong selection (deep valleys), and

indicate that extinction-generated survivorship bias (De Lisle et al. 2020) may make sex-

specific (dimorphic) peak shifts more likely to be observed than monomorphic shifts.

These two features of sex-specific peak shifts may explain several puzzling phenomena

observed in macroevolutionary studies of sexual dimorphism. First, a large number of

studies have investigated the link between proxies for the strength of within-population

sexual selection and among-lineage patterns of sexual dimorphism. Often, these studies find

only weak relationships between these sexual-selection proxies and the magnitude of sexual

dimorphism (Shine 1989, Cox et al. 2003, Cox and Calsbeek 2009, Nali et al. 2014). This

finding of little correlation between the magnitude (yet some correlation between the sign;

Janicke et al. 2016) of sexual dimorphism and sexual selection proxies is consistent with the

results presented here; when the two-sex adaptive landscape is complex, macroevolutionary

patterns of sexual dimorphism may be largely decoupled from the strength of sexual

selection, as strong sexual selection may often simply lead to concordant peak shifts when

such peaks exist. Concomitantly, any association between r_{mf} and the strength of sexual

selection within populations (as some data implies could be the case; Bonduriansky and

Rowe 2005, Poissant et al. 2009) could have important macroevolutionary consequences.

Similar weak and varying relationships between sexual selection proxies and lineage

diversification and extinction rates may also be consistent with the conclusions of the peak

shift model presented here (Morrow and Pitcher 2003, Morrow et al. 2003, Morrow and

Fricke 2004). Peak shifts are required for ecological speciation, and sexual selection can

readily induce such peak shifts. Yet, ultimately, the occurrence and final outcome of a peak

shift depends not just on sexual selection, but rather its interaction with genetic correlations

and ecological opportunity.

Accounting explicitly for demographic effects of maladaptation reveals that sex-

specific peak shifts may be more likely to be successful in the face of extinction, compared to

concordant peak shifts. This effect would be predicted to lead to a sex-bias in

macroevolutionary rate. When male adaptation contributes little to population growth rate,

this may lead to a male bias in macroevolutionary rate if male valleys are easier to cross

without extinction. Such effects would be expected to be especially strong for traits that are

expected to be tightly coupled with population growth. For example, in many clades, males

are observed to exhibit higher rates of body size evolution than females (Rensch 1960,

Fairbairn 1997, Colwell 2000, Stephens and Wiens 2009). Although the opposite (higher

female rates) is observed in some groups (De Lisle and Rowe 2013), it does appear to be

rarer. Although past explanations for male-bias in macroevolutionary rates of body size

evolution have often focused on strong sexual selection, it may be the case that these biased

rates are in part an outcome of survivorship bias.

My analysis has focused on a simplistic case of multiple optima on a two-sex adaptive

landscape, where the height of each optimum is the same. Evolution on real adaptive

landscapes, with mixtures of peaks of varying height, varying distance from each other, will

surely be more complex. None the less, the general conclusions presented here are expected

to hold; whether a peak shift to a dimorphic or monomorphic peak occurs depends in part on

an interaction between sexual selection and genetic correlations that may decouple the

strength of sexual selection from the magnitude of resulting sexual dimorphism at

equilibrium, and crossing two valleys at once may be demographically costly.

Acknowledgements I thank David Punzalan, Locke Rowe, Erik Svensson, Beatriz Willink,

Masahito Tsuboi, and Benjamin Jarrett for discussion and/or comments on the manuscript.

Funding was provided by grants from the Swedish Research Council and Royal

Physiographical Society of Lund to S.P.D.

Figure 1. Models of stabilizing selection in two sexes. Panels illustrate the adaptive

landscape, the function $\overline{W}_m\overline{W}_f$. In Lande's (1980) original model of evolution of sexual

dimorphism by natural and sexual selection, a single multivariate optimum, corresponding to

one optimum for the male trait and one for the female trait, is assumed (**A**). The 'Twin Peaks'

model extends this to consider two optima in one trait (sex), with only a single optimum for

the other trait (sex); (**B**). Extending the Twin Peak model to both sexes results in two optima

for each trait, or four multivariate optima on the two-sex adaptive landscape (**C**). As

parameterized in this manuscript, the Two-sex Twin Peak model contains two sexually

monomorphic optima, and two sexually dimorphic optima. Of interest is when and how peak

shifts from an ancestral monomorphic optimum occur. In this model sexual selection leads to

336   a displacement of males from their optimum (**C**), the magnitude of which is proportional to

337   the strength of stabilizing natural selection. Two trajectories are shown starting from the

338   optimum; one where $r_{mf} = 0$ (blue) and one where $r_{mf} = 0.9$ (blue).

Figure 2. Peak shifts induced by sexual selection in the Two-sex Twin Peaks model.

Panel **A** shows two deterministic trajectories, starting from the lower left optimum,

corresponding to two different values of the cross-sex genetic correlation r_{mf}. Although

sexual selection is strong enough to induce a peak shift, which new optimum reached

depends upon the value of r_{mf}. Panel **B** illustrates 20 replicates of stochastic evolution (i.e.,

with drift) under the same parameter values as A and assuming female demographic

dominance ($N_{t+1} = k \, \overline{W}_f N_t$). Round circles illustrate the mean values after 10000

generations of evolution; extinction events are denoted with a crossed circle. Panel **C**

contrasts trajectories under two strengths of sexual selection c_m, with otherwise identical

parameter values. Panel **D** illustrates stochastic evolution under the same parameter values as

C. White dashed line illustrates the line of sexual monomorphism, for reference. In A and B,

$c_m = 0.2$; B and C $r_{mf} = 0.8$; other parameter values as described in text.

Figure 3. Deterministic outcomes as a function of sexual selection and r_{mf}. Panels show

deterministic (i.e. no drift) outcomes of 1000 generations of evolution, calculated

numerically, corresponding to three different strengths of stabilizing selection. Left panels

assume female demographic dominance, right panels assume male and female fitness both

373   determine population growth rate. 'Dimorphic shift' refers to a male-only peak shift,

374   'Monomorphic shift' refers to a peak shift in both sexes (illustrated in Fig 2A). No peak shift

375   means the lineage remains in the vicinity of the ancestral optimum. Small gaps between

376   regions represent plotting limitations of overlaid contour plots.

398   **Literature Cited**

399   Arnold, S. J., M. E. Pfrender, and A. G. Jones. 2001. The adaptive landscape as a conceptual

400   bridge between micro- and macroevolution. Genetica **112/113**:9-32.

401   Barton, N. H., and B. Charlesworth. 1984. Genetic revolutions, founder effects and

402   speciation. Annual Review of Ecology and Systematics **15**:133-164.

403   Bonduriansky, R. 2011. Sexual selection and sexual conflict as engines of ecological

404   diversification. American Naturalist **178**:729-745.

405   Bonduriansky, R., and L. Rowe. 2005. Sexual selection, genetic architecture, and the

406   condition dependence of body size and shape in the sexually dimorphic fiy *Prochyliz*

407   *xanthostoma* (Diptera: Piophilidae). Evolution:138-151.

408   Colwell, R. K. 2000. Rensch's rule crosses the line: Convergent allometry of sexual size

409   dimorphism in hummingbirds and flower mites. American Naturalist **156**:495-510.

410   Cox, R. M., and R. Calsbeek. 2009. Sexually antagonistic selection, sexual dimorphism, and

411   the resolution of intralocus sexual conflict. American Naturalist **173**:176-187.

412   Cox, R. M., S. L. Skelly, and H. B. John-Alder. 2003. A comparative test of adaptive

413   hypotheses for sexual size dimorphism in lizards. Evolution **57**:1653-1669.

414   Coyne, J. A., N. H. Barton, and M. Turelli. 1997. Perspective: a critique of Sewall Wright's

415   shifting balance theory of evolution. Evolution **51**:643-671.

416   Coyne, J. A., and H. A. Orr. 2004. Speciation. Sinaur, Sunderland, MA.

417   De Lisle, S. P., D. Punzalan, N. Rollinson, and L. Rowe. 2020. Extinction and the temporal

418   distribution of macroevolutionary bursts. Journal of Evolutionary Biology **in press**.

419   De Lisle, S. P., and L. Rowe. 2013. Correlated evolution of allometry and sexual dimorphism

420   across higher taxa. American Naturalist **182**:630-639.

Fairbairn, D. J. 1997. Allometry for sexual size dimorphism: Pattern and process in the

coevolution of body size in males and females. Annual Review of Ecology and

Systematics **28**:659-678.

Felsenstein, J. 1979. Excursions along the interface between disruptive and stabalizing

seleciton. Genetics **93**:773-795.

Fisher, R. A. 1930. The genetical theory of natural selection. The Clarendon press, Oxford.

Frank, S. A. 2013. Wright's Adaptive Landscape Versus Fisher's Fundamental Theorem.*in* E.

I. Svensson and R. Calsbeek, editors. The Adaptive Landscape in Evolutionary

Biology. Oxford University Press.

Gingerich, P. D. 2001. Rates of evolution on the time scale of the evolutionary process.

Genetica **112**:127-144.

Goodnight, C. J. 2013. Wright's Shifting Balance Theory and Factors Affecting the

Probability of Peak Shifts.*in* E. I. Svensson and R. Calsbeek, editors. The Adaptive

Landscape in Evolutionary Biology. Oxford University Press.

Janicke, T., I. K. Häderer, M. J. Lajeunesse, and N. Anthes. 2016. Darwinian sex roles

confirmed across the animal kingdom. Science Advances **2**:e1500983.

Lande, R. 1976. Natural selection and random genetic drift in phenotypic evolution.

Evolution **30**:314-334.

Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.

Evolution **34**:292-305.

Lande, R. 1985. Expected time for random drift of a population between stable phenotypic

states. Proceedings of the National Academy of Sciences **82**:7641-7645.

Lande, R., and M. Kirkpatrick. 1988. Ecological speciation by sexual selection. Journal of

Theoretical Biology **133**:85-98.

Matthews, G., S. Hangartner, D. G. Chapple, and T. Connallon. 2019. Quantifying

maladaptation during the evolution of sexual dimorphism. Proceedings of the Royal

Society of London Series B. **286**:20191372.

Morrow, E. H., and C. Fricke. 2004. Sexual selection and the risk of extinction in mammals.

Proceedings of the Royal Society B: Biological Sciences **271**:2395-2401.

Morrow, E. H., and T. E. Pitcher. 2003. Sexual selection and the risk of extinction in birds.

Proceedings of the Royal Society of London Series B-Biological Sciences **270**:1793-

1799.

Morrow, E. H., T. E. Pitcher, and G. Arnqvist. 2003. No evidence that sexual selection is an

'engine of speciation' in birds. Ecology Letters **6**:228-234.

Nali, R. C., K. R. Zamudio, C. F. B. Haddad, and C. P. A. Prado. 2014. Size-dependent

selective mechanisms on males and females and the evolution of sexual size

dimorphism in frogs. The American Naturalist **184**:727-740.

Poissant, J., A. J. Wilson, and D. W. Coltman. 2009. Sex-specific genetic variance and the

evolution of sexual dimorphism: a systematic review of cross-sex genetic correlations.

Evolution **64**:97-107.

Price, T., M. Turelli, and M. Slatkin. 1993. Peak shifts produced by correlated response to

selection. Evolution **47**:280-290.

Rensch, B. 1960. Evolution Above the Species Level. Columbia University Press, New York.

Shine, R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the

evidence. Quarterly Review of Biology **64**:419-461.

Stephens, P. R., and J. J. Wiens. 2009. Evolution of sexual size dimorphisms in emydid

turtles: ecological dimorphism, Rensch's rule, and sympatric divergence. Evolution

63:910-925.

Uyeda, J. C., T. F. Hansen, S. J. Arnold, and J. Pienaar. 2011. The million year wait for

macroevolutionary bursts. Proceedings of the National Academy of Sciences of the

United States of America **108**:15908-15913.

Wright, S. 1931. Evolution in Mendelian populations. Genetics **16**:97-159.

Wright, S. 1932. The roles of mutation, inbreeding, crossbreeding and selection in evolution.

Proceedings of the Sixth Annual Conference of Genetics **1**:356-366.

Supplemental Figures

Figure S1 Demographic cost of valley crossing. Figures show population size from

trajectories plotted in Figure 2A and 2C. Panel A contrasts two values of the cross-sex

genetic correlation, r_{mf} = 0.8 (red line) or r_{mf} = 0.9 (blue line). Panel B contrasts two values

of the strength of sexual selection, cm = 0.2 (red line) or cm = 0.3 (blue line). Demographic

costs of a monomorphic peak shift, corresponding to the blue line in both panels, are higher

than for a dimorphic (male-only peak shift). Other parameter values are as described in

Figure 2.

Figure S2 Stochastic outcomes as a function of sexual selection and r_{mf}. Panels show

stochastic (i.e., evolution under both selection and drift) outcomes of 1000 generations of

evolution, calculated numerically, corresponding to the scenario of strong natural selection

and female demographic dominance illustrated in Figure 3C. Thus, qualitative conclusions

on peak shifts remain similar to the deterministic case, and this holds for other parameter

values shown in Figure 3.