Recovering sedimentary ancient DNA of harmful dinoflagellates off Eastern Tasmania, Australia, over the last 9 000 years

Authors:

Linda Armbrecht[1,2*], Bradley Paine[1], Christopher J.S. Bolch[1], Alan Cooper[3], Andrew McMinn[1], Craig Woodward[4] and Gustaaf Hallegraeff[1]

Affiliations:

[1] Institute for Marine and Antarctic Studies, University of Tasmania, Battery Point TAS 7004, Australia

[2] Australian Centre for Ancient DNA, School of Biological Sciences, Faculty of Sciences, The University of Adelaide, Adelaide SA 5005, Australia

[3] Gulbali Institute, Agriculture, Water and Environment, Charles Sturt University, Albury NSW 2640, Australia

[4] Australian Nuclear Science and Technology Organisation, Locked Bag 2001, Kirrawee DC NSW

*Correspondence: linda.armbrecht@utas.edu.au

Highlights

- Metagenomic *sed*aDNA and hybridisation capture enabled analyses of harmful dinoflagellates off Tasmania
- Sequence validation was used to confirm the presence of *Alexandrium* spp., *Gymnodinium* spp. and *Noctiluca scintillans*
- *Alexandrium* and *Gymnodinium* have been present in Tasmanian waters during the past ~9 000 years
- *Noctiluca scintillans sed*aDNA derived relative abundance correlates with its recorded increase since 2010

Abstract

Harmful algal blooms (HABs) have significantly impacted the seafood industry along the Tasmanian east coast over the past four decades. To investigate the history of regional HABs, we applied sedimentary ancient DNA analyses (*sed*aDNA) to coastal sediments up to ~9 000 years old collected inshore and offshore Maria Island, Tasmania. We used metagenomic

shotgun sequencing combined with a hybridisation capture array ('HABbaits1') to target harmful dinoflagellates of the genera *Alexandrium, Gymnodinium,* and *Noctiluca scintillans.* Bioinformatic analyses were used to verify *sed*aDNA sequences and their presence in older layers, especially for microreticulate cyst forming species including *Gymnodinium catenatum* due to its important role in shellfish toxicity. Our results show that the *Alexandrium* genus (up to 854 and 20 reads per sample inshore and offshore, respectively, based on capture-data) has been present off eastern Tasmania within the last ~8 307 years. For *G. catenatum* we detected a total of only 9 unambiguously verified reads sporadically between ~7 638 years ago and the present in the offshore core. We recovered verified *sed*aDNA of the fragile, non-fossilising species *N. scintillans*, along with evidence of increased relative abundance from 2010, consistent with plankton surveys. This study identifies challenges regarding *sed*aDNA sequence validation of some species (in particular, for *G. catenatum*), and provides guidance for the development of tools to monitor past and present HAB species and events, and to improve future HAB event predictions.

Key words:
Alexandrium; Gymnodinium catenatum; Noctiluca scintillans; biotoxins; ballast water; seafloor

Abbreviations:
harmful algal blooms, HAB; heuristic operations for pathogen screening, HOPS, hydrofluoric acid, HF; internally transcribed spacer, ITS; Paralytic Shellfish Poisoning, PSP; Paralytic Shellfish Toxin, PST; sedimentary ancient DNA, *sed*aDNA; small subunit ribosomal rRNA, SSU; large subunit ribosomal rRNA (LSU)

1   Introduction

An increasingly important issue is the extent to which harmful algal blooms (HABs) are expanding in distribution, magnitude, and frequency due to climate change, eutrophication, and anthropogenic transport. The negative impacts of HABs on tourism, aquaculture, fisheries, and human health, mean that the appearance of novel HAB phenomena regularly raise the question of whether the species responsible is a recent introduction (e.g., via ballast water; McMinn et al., 1997, Bolch and de Salas 2007), or a previously cryptic endemic species newly stimulated by changing environmental conditions (Hallegraeff et al., 2021) or extreme climate events (Trainer et al., 2019). Few studies have examined the "cryptic species" hypothesis by investigating long-term dynamics over thousands of years (e.g., Thorsen et al., 1995, *Gymnodinium nolleri* misidentified as *G. catenatum* in Scandinavia; Klouch et al., 2016, *Alexandrium minutum* in the Bay of Brest).

The ocean environment off eastern Tasmania is a well-documented climate change hotspot
characterised by a strengthening East Australian Current and rapidly increasing ocean
temperatures (2.3 °C increase since the 1940s, Ridgway and Hill, 2009). The consequences of
this oceanographic change are being detected in coastal marine communities, including
changes in plankton and HAB species composition (Thompson et al., 2009, Condie et al.,
2019). Particular examples are the HAB dinoflagellates *Gymnodinium catenatum*, *Noctiluca*
scintillans, and *Alexandrium* species, which we focus on here.

G. catenatum produces Paralytic Shellfish Toxin (PST) and is the only toxic member of a
phylogenetically distinct lineage of *Gymnodinium* species (*G. catenatum, G, inusitatum, G.*
microreticulatum, G. nolleri, and *G. trapeziforme*) that produce fossilisable resting cysts with
distinctive surface reticulation – referred to hereafter as 'microreticulate species'. Thought to
have been introduced to Tasmania in the 1970s by shipping ballast water (McMinn et al. 1997),
G. catenatum first bloomed in the mid-1980s, caused PST contamination up to 250-fold above
acceptable limits, and extensive shellfish farm closures in the Derwent-Huon estuaries from
1986 to 1993 (Hallegraeff et al., 1995). Supporting evidence for a recent introduction includes
detection of resting cysts in ships ballast tanks (Hallegraeff and Bolch, 1992), a lack of cysts
in dated marine sediment cores prior to the 1970's (McMinn et al., 1997), and molecular
evidence from rRNA gene sequencing that links Australasian populations to source populations
from the Seto inland sea in southern Japan (Bolch and de Salas, 2007).

Noctiluca scintillans was first documented in Australia from Sydney Harbour (Bennett, 1860),
but since the 1990s has increasingly caused highly visible red tides and bioluminescence,
resulting in frequent temporary closures of popular Sydney tourist beaches (Hallegraeff et al.,
2020). Amongst the suggested causes of *N. scintillans* blooms are eutrophication and coastal
upwelling generating more diatom prey (Dela Cruz et al., 2002, 2003). *Noctiluca* was first
observed in Tasmanian waters in 1994, and is presumed to have dispersed southward with the
East Australian Current, presenting a new threat to the salmonid fish farm industry from 2002
(Hallegraeff et al., 2019). In 2010, *N. scintillans* was detected in the Southern Ocean for the
first time, 240 km south of Tasmania, raising concerns of grazing impacts on iconic krill-based
food webs (McLeod et al., 2012).

Since 2012, further PST-caused seafood harvest closures of Tasmanian bivalves, abalone, and
rock lobsters, along with public health warnings have been generated by winter-spring blooms
of the cool-temperate dinoflagellate *Alexandrium catenella*. These events have resulted in PST
contamination up to 150 mg saxitoxin (STX) equiv./kg (Condie et al., 2019). The
morphologically similar but genetically distinct species *A. australiense* and *A. pacificum*
(Bolch and DeSalas, 2007) were previously known from Tasmanian waters, but *A. catenella*
(= *A. tamarense* Group 1; =*A. fundyense*; John et al., 2014), had not been previously detected
in Australasian waters. Microsatellite DNA analyses of regional populations show that
Tasmanian *A. catenella* are divergent from other populations (John et al. 2018), indicating it is
either endemic or dispersed naturally to the area millenia ago. Thus, *A. catenella* may be a

previously cryptic population recently stimulated by changing environmental conditions, such
as increased winter water column stratification (Trainer et al., 2019). At least five cases of non-
fatal human paralytic shellfish poisonings (four from *A. catenella* and one from *G. catenatum*)
have been formally reported from Tasmania in the past 10 - 20 years (Turnbull et al., 2013;
Edwards et al., 2018), but it remains unknown if paralytic shellfish poisonings occurred in this
region prior to the arrival of Europeans.

Marine sediment archives containing highly resistant sub-fossil resting cysts are an important
resource to examine long-term dynamics of harmful dinoflagellate species and address key
questions about their introduction, disappearance, and reappearance in a region. Analysis of
sedimentary ancient DNA (*sed*aDNA) from similar samples offers distinct advantages for the
analysis of non-cyst forming or less-well preserved cysts that are currently not accounted for
in microfossil-based ecosystem reconstructions (Armbrecht, 2020; Bolch, 2022). For example,
Noctiluca does not produce cysts but has been detected using *sed*aDNA previously (Shaw et
al., 2019). However, *sed*aDNA analysis is not straightforward as the DNA is highly fragmented
(typically <100 bp), occurs at low concentrations, and any one target species represents only a
small fraction of the total DNA present. As a result, it is important to enrich target species
*sed*aDNA via methods such as hybridisation capture (Armbrecht, et al., 2021), and to use
bioinformatic assessment of DNA damage patterns typical of degraded ancient DNA to
authenticate the results (Hübler et al., 2019). Accordingly, an RNA probe array ('HABbaits1')
was designed previously to bind and retain taxonomic marker genes (18S rRNA, 28S rRNA,
Internal transcribed spacer (ITS), ribulose-bisphosphate carboxylase (rbcL), cytochrome c
oxidase subunit 1 (COI)) of harmful dinoflagellates (and other plankton groups), including
above described *Alexandrium* spp.*, G. catenatum*, and *N. scintillans* (for full details of the
HABbaits1 design see Armbrecht et al., 2021).

In this study, we combine metagenomic shotgun sequencing and hybridisation capture using
the HABbaits1 array to investigate the presence and palaeo-historical patterns of *Alexandrium,*
Gymnodinium catenatum, and *Noctiluca scintillans* from Tasmanian marine sediments over the
past ~9 000 years. We then use this information to support ecological interpretation of regional
community change in Tasmanian waters.

2   Materials and Methods

2.1   Sediment core collection and preparation
An approximately 3 m long sediment core (gravity core, designated 'GC2S1') was collected in
May 2018 during *RV Investigator* voyage IN2018_T02 in 104 m water depth close to the
continental shelf edge, east of Maria Island, Tasmania (Site 1; 148.240 °E; 42.845 °S) (Fig. 1).
Since gravity cores can disturb surface sediments, and the latter can be lost due to the horizontal

tipping after collection, a shorter (12 cm) parallel core was obtained at the same site using a
KC Denmark Multi-Corer (designated 'MCS1-T6', where 'T6' refers to 'Tube 6' of the multi-
corer). A 35 cm long multi-core was also obtained from within Mercury Passage, west of Maria
Island (Site 3; 42.550 °S, 148.014 °E) in a water depth of 68 m (designated 'MCS3-T2'). All
cores were immediately capped, sealed, labelled and transported in their original PVC coring
tubes to the Australian Nuclear Science and Technology Organisation (ANSTO), Lucas
Heights, Australia, where they were kept at 4°C. The cores GC2S1, MCS1-T6, and MCS3-T2
were opened, split, scanned (using a multi-function core scanning instrument (ITRAX) with
X-ray fluorescence (XRF), radiographic X-ray, optical imaging and magnetic susceptibility
measurements), and subsampled for *sed*aDNA analyses in October 2018 (i.e., as fast as possible
considering the logistics involved). To minimise contamination during core splitting and
sampling, working benches and cutting knives were cleaned with 3% bleach and 70% ethanol,
gloves changed immediately when contaminated with sediment, and appropriate PPE was worn
at all times (gloves, facemask, hairnet, disposable coverall). Sampling of GC2S1 was
conducted by first removing the outer ~1 cm of the working core-half and then taking
subsamples by pressing sterile 15 mL centrifuge tubes ~3 cm into the sediment at 5 cm intervals
(working from bottom to the top of the core at each step). Sampling of MCS1-T6 was
conducted as for GC2S1 except at finer intervals of 2 cm. Mercury Passage multicore (MCS3-
T2) sampling was conducted at 2 cm depth intervals in the top 8 cm and at 5 cm intervals
below. All *sed*aDNA samples were immediately stored at -20 °C, respectively. Hereafter, we
refer to sediment depths as 0, 2 cm, *etc.*, however it should be noted that due to the diameter of
the sample collection centrifuge tubes (~1.5 cm diameter), this refers to the minimum depth of
the sample interval which was actually 0 - 1.5 cm, 2 – 3.5 cm, etc., respectively.
171

172
173 **Figure 1: Sediment coring sites near Maria Island, Tasmania, Australia.** Overview of
174 coring locations offshore (Gravity Core Site 1, GC2S1, and Multi-Core Site 1 Tube 6, MCS1-
175 T6) and inshore Maria Island in the Mercury Passage (Multi-Core Site 3 Tube 2, MCS3-T2).
176 GC2S1 and MCS3-T6 were collected adjacent to each other and are depicted as one dot. Map
177 created in ODV (Schlitzer, R., Ocean Data View, https://odv.awi.de, 2018).
178

2.2   Sediment dating
ITRAX scanning, XRF, radiographic X-ray, optical imaging and magnetic susceptibility
measurements confirmed excellent undisturbed preservation of the cores. An age profile was
generated for MCS3-T2 and MCS1-T6 based on ^{210}Pb measurements (8 and 6 dates,
respectively), while GC2S1 was based on both ^{210}Pb (7 dates) and ^{14}C (3 dates). A Bayesian
age-depth model was constructed for each site based on these ^{210}Pb and ^{14}C measurements
using rbacon (Blaauw et al., 2019) on the R platform (R Core Team, 2013) with the SHCal20
curve for radiocarbon age calibration (Hogg et al., 2020). Details on the construction of the
age-depth model are provided within the Supplementary Material (Supplementary Material
Note 1, Supplementary Material Fig. 1, Supplementary Material Table 1).
189

2.3   Sedimentary ancient (sedaDNA) extractions
Extractions of *sed*aDNA took place within ultraclean ancient (GC2S1) and forensic (MCS1-
T6, MCS3-T2) facilities (ACAD, The University of Adelaide) following recommended ancient

DNA decontamination standards (Willerslev and Cooper, 2005). The extraction technique
followed the 'combined' protocol developed for marine eukaryote *sed*aDNA (Armbrecht et al.,
2020). This method combines a gentle EDTA incubation step to isolate fragile eukaryote DNA
(Slon et al., 2017) with bead-beating to extract intracellular DNA from robust spores and cysts
(Shaw et al., 2019). Smaller DNA fragments ($\geq$27 base pairs, bp) characteristic of ancient DNA
are targeted using in-solution silica binding (Brotherton et al., 2013), along with magnetic
beads to size-select DNA fragments under 500 bp (Armbrecht et al., 2020). DNA extracts were
prepared from 42 sediment samples (GC2S1, MCS1-T6, MCS3-T2) and 7 extraction blank
controls (see also Armbrecht et al., 2021).
202

2.4 Shotgun sequencing library preparations
Metagenomic shotgun libraries were prepared from 42 sediment samples (GC2S1, MCS1-T6,
MCS3-T2) and 7 extraction blank controls using established techniques (Armbrecht et al.,
2020, 2021). Sequencing was undertaken using an Illumina NextSeq sequencing platform (2 x
75 bp cycle) at the Australian Cancer Research Foundation Cancer Genomics Facility & Centre
for Cancer Biology (Adelaide, Australia) and the Garvan Institute of Medical Research, KCCG
Sequencing Laboratory Kinghorn Centre for Clinical Genomics (Darlinghurst, Australia).
210

2.5 Hybridization capture and sequencing library preparations
In order to maximise the yield of *sed*aDNA from our target dinoflagellate species a
hybridization-capture technique was developed and applied. An RNA array or 'bait set' termed
'HABbaits1' was developed in collaboration with Arbor Biosciences, USA, to target the
harmful dinoflagellates *Alexandrium* groups I – IV, *Gymnodinium catenatum* and *Noctiluca*
scintillans. Details of the design, protocol optimisation, and application of HABbaits1 to
marine *sed*aDNA have been published previously (Armbrecht et al., 2021). A final pool of a
selection of 30 multiplexed sequencing libraries (GC2S1 and MCS3-T2) prepared from
HABbaits1 was sequenced using Illumina HiSeq XTen (2 x 150 bp cycle) at the KCCG,
Darlinghurst, Australia.
221

2.6 Bioinformatics
Bioinformatic processing of sequence data followed published protocols (Armbrecht et al.
2020), using software and analytical parameters from Armbrecht et al. (2021). After filtering
to remove low-complexity and duplicate reads, each dataset was processed without the
standardisation step (*i.e.,* without rarefying) to retain the maximum number of reads, which is
crucial for ancient DNA damage analysis (see below). However, to assess our data semi-
quantitatively, we also standardised the filtered shotgun data by subsampling (*i.e.,* rarefying)
to the lowest number of reads detected in a sample, *i.e.,* 2.2 million (Mio) (using seqtk version
1.2), and continued to process this data in parallel to the non-rarefied data. Subsampling

reduced the number of reads significantly therefore this data is provided in the Supplementary
Material only.

After quality control (FastQC v.0.11.4, MultiQC v1.8), the NCBI Nucleotide database
(ftp://ftp.ncbi.nlm.nih.gov/blast/db/FASTA/nt.gz, downloaded November 2019) was used as
the reference database to build a MALT index (Step 3) and sequences aligned using MALT
(version 0.4.0; semiglobal alignment) (Herbig et al., 2016). All resulting .blastn files were
converted to .rma6 files using the Blast2RMA tool in MEGAN (version 6_18_9, Huson et al.,
2016). Subtractive filtering (*i.e.,* subtracting reads for species identified in EBCs from samples)
was conducted separately for the shotgun and HABbaits1 data (see Armbrecht et al., 2021
Supplementary Material, for a comprehensive list of eukaryote contaminants), however, no
Dinophyceae taxa were detected in EBCs. For both datasets, read counts of all Dinophyceae
nodes (from MEGAN6 v. 18.10) were exported for downstream analyses.

To test *sed*aDNA damage, the 'MALTExtract' and 'Postprocessing' tools of the HOPS v0.33-
2 pipeline (Hübler et al., 2020) were run for our three target dinoflagellates using the same
configurations as used in Armbrecht et al. (2021), eg. taxalist *'b'*, including *Alexandrium* spp.,
Gymnodinium spp., and *N. scintillans* on shotgun and HABbaits1 *sed*aDNA data. The
MALTExtract output, *i.e.,* reads categorised as ancient (showing damage) or default (passing
stringent filtering criteria but not showing damage) for the three target taxa was exported and
the proportion of *sed*aDNA damage per taxon was determined. Also, *sed*aDNA damage
profiles were generated for these three taxa using MALTExtract Interactive Plotting
Application (MEx-IPA, https://github.com/jfy133/MEx-IPA), however, due to low read
numbers these graphics were not informative and excluded from downstream analyses.

To test the reliability of short sequence *sed*aDNA assignments to the three targets groups, we
created three dummy sample datasets, each containing sequences of one of the three
dinoflagellate groups downloaded from NCBI (see Supplementary Material Note 3). Each
sequence was included with its complete length (Supplementary Material Table 2) and split
into 56 bp fragments (corresponding to the average sequence length of our filtered shotgun
data, see Results). If the last fragment was <25 bp it was excluded from the dummy sample
(mimicking a minimum cut-off of 25 bp during sample data processing). Dummy samples were
converted from .fasta to .fastq files (see Supplementary Material Note 3) and processed via the
same analytical pipeline used for sample sequencing.

The initial assessment of s*ed*aDNA assigned to *Gymnodinium catenatum* indicated a significant
proportion of sequences mapped to regions of the LSU-rRNA gene where there is limited
database coverage of gymnodinoid dinoflagellates (see Supplementary Material Note 4,
Supplementary material Figs. 4,5). Due to the potential for reference bias, we undertook
additional sequencing of the D3 to D10 region (~1 850 bp) of the LSU-rRNA of related
microreticulate species available to us in culture, *G. microreticulatum* (strain CAWD191) and

G. nolleri (strain K-0626) (Supplementary Material Note 4) and carried out more detailed
validation of these sequences Firstly, gymnodinoid-assigned reads were mapped to *G.*
catenatum reference sequence (DQ785882) including comparative alignment with related
Gymnodinium species (default alignment and assembly; Geneious Prime 2021). Direct
match/mismatches to species in the alignment were used to determine the nearest-neighbour
taxon and base-pair mismatches from the reference sequence for each read (Supplementary
Material Fig. 5). Secondly, all *Gymnodinium* reads assigned to *G. catenatum* after the initial
NCBI run were then re-run against an in-house *Gymnodinium*-focused reference sequence
database with addition of newly generated *G. microreticulatum* and *G. nolleri* LSU-rRNA D3-
D10 sequences (see Supplementary Material Note 5).
282
283

3 Results

3.1 Age model
Our age model revealed that the inshore core MCS3-T2 dates to 77 years before 1950 at 34.5
centimetres below seafloor (cmbsf), *i.e.,* the year 1873 (mean value, ~144 years before sample
collection in 2017). The offshore gravity core GC2S1 dates to 8 878 years before 1950 at 268
cmbsf (~8 945 years before sample collection in 2017). Comparing the ^{210}Pb activity profiles
from both GC2S1 and the adjacent multicore MCS1-T6, we estimated that ~3.5 cm were
missing from the top of GC2S1, representing the last ca. 30 years (Supplementary Material Fig.
1).
293

3.2 Validation of Alexandrium, Gymnodinium., and Noctiluca scintillans short read species
assignments
Dummy dataset analyses revealed short reads of *Alexandrium* spp. could be reliably assigned
to genus level, thus we interpret this target taxon exclusively at genus level (see Supplementary
Material Note 3, Supplementary Material Table 2, Supplementary Material Fig. 3). For *N.*
scintillans, the majority of dummy sequences were correctly assigned to this species, and
correctly back-mapped to the NCBI sequence used to create our dummy sample (GQ380592.1).
Some reads were assigned to a different *N. scintillans* 18S rRNA reference sequence available
on NCBI (AF022200.1), but species-level assignment remained correct therefore we consider
identification of *N. scintillans* from short *sed*aDNA fragments to be robust and reliable. For
Gymnodinium catenatum, most dummy reads were correctly assigned but few reads were
erroneously assigned to the apicomplexan genus *Eimeria*, the Streptophytina, the
Membracoidea, other dinoflagellates such as *Durinskia baltica, Symbiodinium* sp. AW-2009,
or the related microreticulate species *G. microreticulatum*, suggesting *Gymnodinium* species
assignment of short reads (56 bp) of 18S rRNA should be interpreted with caution. In most
cases assignment uncertainty could be resolved by further inspection and verification.

Similarly, rare mis-assignment of short 18S rRNA reads from *Alexandrium* were made to
Apicomplexa and Apocrita, and for *N. scintillans* to *Symbiodinium* and Apocrita.
312
Reference alignment mapping of HABbaits1 reads assigned to *Gymnodinium* from core GC2S1
(352 reads, see Results), showed the majority (99%) mapped to the rRNA genes
(Supplementary Material Note 4, Supplementary Material Fig. 5). Of these, 65% were within
3 bp of exact match for the *G. catenatum* reference sequence (Supplementary Material Table
3) and 97% could be unambiguously assigned to the microreticulate-group of *Gymnodinium*
species that includes *G. catenatum*. More than half (57%) of the reads were closest to the *G.*
catenatum reference sequence, 36% unambiguously identified as *G. catenatum*, and 12.5% as
G. microreticulatum (Supplementary Material Fig. 5). Twenty-one percent mapped to rRNA
regions where *G. catenatum* could not be resolved from its closest relative *G. nolleri*; 30% to
regions with insufficient sequence variation to distinguish microreticulate-group species
(Supplementary Material Figure 5).
324
325   Re-assignment of the 352 gymnodinoid reads against our supplemented in-house database
326   supported these observations and resulted in more conservative species-level assignment. Of
327   216 reads assigned, 76% could only be assigned at genus level, 28% to *G. nolleri*, 6% to *G.*
328   *microreticulatum*, and 4% to *G. catenatum* (Supplementary Material Note 5, Supplementary
329   Material Table 4).
330

3.3   Representation of Dinophyceae in shotgun data
From the 42 shotgun samples we retrieved a total of 824 503 filtered sequences across the three
domains (Bacteria, Archaea and Eukaryotes), 149 892 of which were assigned to Eukaryota
(18%). A total of 529 reads were assigned to Dinophyceae (*i.e.,* Dinophyceae represented
0.06% of the domains, and 0.35% of all eukaryotes). Harmful dinoflagellate taxa sequences
were detected in low abundance (19 *Alexandrium* spp., 13 *Gymnodinium* spp., and no *Noctiluca*
sequences; Fig. 2). *Alexandrium* spp. were primarily detected in the upper 15 cmbsf at MCS3-
T2 (last ~39 years), and few reads were assigned to this genus between 55 and 189 cmbsf at
GC2S1 (~2 600 and 7 200 years ago, respectively). Based on our *Alexandrium* dummy run
(Section 3.2) we used a conservative approach, at genus level (Fig. 2). Three sequences were
assigned to microreticulate *Gymnodinium* group in GC2S1 at 95 cmbsf (~4,400 years ago), and
219 cmbsf (~7,800 years ago) (Fig. 2C). While microreticulate gymnodinoids were not
detected in the younger sediments of GC2S1, they were present in the surface sediments of the
shorter MCS1-T6 core (Fig. 2B), demonstrating their presence in more recently deposited
offshore sediments.
346
347

Figure 2: Relative Dinophyceae abundance (shotgun). Dinophyceae detected in non-rarefied shotgun data at Maria Island coring sites A) MCS3-T2, B) MCS1-T6, and C) GC2S1. All non-target taxa are grouped as Dinophyceae (grey), while *Alexandrium* and Gymnodiniales and its subgroups are shown separately (in colour). Identity of *G. catenatum* should be interpreted as representing all microreticulate *Gymnodinium* species, and the genus *Alexandrium* (see Section 3.2). *N. scintillans* was not detected in the shotgun data. Total read count (all sites): 529.

3.4   Representation of Dinophyceae using HABbaits1
Hybridisation capture of our 30 selected *sed*aDNA extracts with the HABbaits1 array resulted
in 27 successfully enriched samples from GCS1 and MCS3-T2 with a total of 872 774 reads
across the Bacteria, Archaea and Eukaryote domains, of which 613 853 were assigned to
Eukaryota (70%). A total of 32 075 of the eukaryote reads were assigned to Dinophyceae (*i.e.,*
3.68% of all domains, and 5.23% of all eukaryotes), representing a 61- and 15-fold increase in
Dinophyceae reads relative to shotgun data, demonstrating the efficacy of target enrichment
achieved with HABbaits1.
365
366   The genus *Alexandrium* were most abundant at MCS3-T2, with the maximum number of reads
367   (854 sequences) found inshore at MCS3-T2 12 - 13.5 cmbsf (~190 years ago) (Fig. 3A). At
368   GC2S1, *Alexandrium* spp. were detected sporadically at different depths at low abundance (<20
369   reads per sample) (Fig. 3B).
370
Several reads were assigned to the microreticulate *G. catenatum*-group at both MCS3-T2 and
GC2S1 (352 reads in total). A relatively high abundance (85 reads) of this group was identified
inshore in MCS3-T2 2 - 3.5 cm (~2 years ago, i.e., 2015), while at GC2S1, the microreticulate
G. catenatum-group appeared sporadically offshore in low abundance (≤22 reads) in the upper
section of GC2S1 (above 75 – 76.5 cmbsf; ~3,500 years ago), while being slightly more
abundant (≤123 reads) in the lower core section (below 169 – 170.5 cmbsf, ~6,500 years ago;
Fig. 3B). Re-assignment of the 352 reads initially identified as *G. catenatum* to the new
Gymnodinium database provided 216 reads assigned to *Gymnodinium* spp. (all others remained
unassigned). Most reads were assigned on genus level (165 reads), and 28, 14 and 9 reads were
assigned to *G. nolleri, G. microreticulatum* and *G. catenatum,* respectively (Fig. 4). *G. nolleri*
was identified sporadically through GCS1, while *G. microreticulatum* and *G. catenatum* were
found primarily in surface sediments at MCS3 and/or GC2S1 as well as sporadically in deeper
samples at GC2S1 (e.g., 209 cmbsf, ~7,638 years ago) (Fig. 4).
384
385   Hybridisation capture also allowed the identification of *N. scintillans*, primarily in the upper
386   section of MCS3-T2 (above 15 cmbsf), reaching maximum relative abundance (based on 59
387   sequences) at 6 - 7.5 and 12 - 13.5 cmbsf (~7 and 15 years ago, *i.e.,* around 2010 and 2002)
388   (Fig. 3A). *N. scintillans* was also detected in the two top samples at GC2S1 (≤8 reads each,
389   Fig. 3B).
390
391   All three target dinoflagellate taxa were detected in low relative abundance in the deepest
392   sample (Fig. 3B), but we cannot rule out the possibility of seawater contamination of the bottom
393   core sample during core retrieval. Additionally, dummy sample validation indicates the
394   possibility that shorter *Gymnodinium* and *Noctiluca* sequences may be misassigned to
395   *Symbiodinium* (see Section 2.3), increasing relative abundance of these taxa by ~1% (Fig. 3).
396
397

Figure 3: Relative Dinophyceae abundance (HABbaits1). Dinophyceae detected in non-rarefied HABbaits1 data at Maria Island coring sites A) MCS3-T2 and B) GC2S1, with target taxa highlighted. Highlighted are *Alexandrium* spp. (pink), the microreticulate gymnodiniods (yellow) and broader genus *Gymnodinium* spp. (orange), and *N. scintillans* (red) within the broader group Noctilucales/*Noctiluca* (total of 1 and 2 reads only, respectively). All other Dinophyceae are summarised (dark grey), with *Symbiodiniaceae* shown separately (light grey) due to potential misassignments with *Gymnodinium* or *Noctiluca* (section 3.2). Total number of reads (both sites): 32 075.

412
Figure 4: Microreticulate *Gymnodinium* species. Re-assignment of 352 *Gymnodinium* spp.
at Site MCS3 (A) and GC2S1 (B) using an in-house *Gymnodinium* spp. database supplemented
with additional microreticulate rRNA gene sequences. A total of 216 reads were assigned to
genus and/or species level. The deepest sample at GC2S1 (267 cmbsf) should be interpreted
with caution due to potential seawater contamination during core retrieval.
418
419

3.5 sedaDNA damage analysis and authentication

Application of the HOPS DNA damage approach to the HABbaits1 data at site MCS3-T2
identified 30, 4, and 1 reads with ancient characteristics (*i.e.,* showed clear signs of DNA
damage) for *Alexandrium, G. catenatum*, and *N. scintillans*, respectively. At site GC2S1, and
22, 95, and 11 'ancient' reads were identified for *Alexandrium, G. catenatum*, and *N.
scintillans*, respectively (Table 1). A relatively high number of reads passed the default filtering
criteria in HOPS (Table 1) - 2 and 15% of the *Alexandrium* reads were classified as ancient in
MCS3-T2 and GC2S1, respectively (Fig. 5). For *G. catenatum*-group, 4 and 27% of reads were
classified as ancient in MCS3-T2 and GC2S1, respectively, and for *N. scintillans*, 0.5 and 18%
were classified as ancient (Table 1, Fig. 5). As such, reads assigned to the three target
dinoflagellates showed much higher *sed*aDNA damage at the offshore site GC2S1 relative to
the inshore site MCS3-T2.
432
433

Figure 5: Ancient and default reads identified for three target HAB taxa in MCS3-T2 and GC2S1. Shown are the proportion of *sed*aDNA reads classified by HOPS analysis as ancient and default (% DNA damage) per sample based on the HABbaits1 capture sequences. The number of reads underlying these proportions are indicated within the bars.

Table 1: *sed*aDNA damage of reads assigned to *Alexandrium* genus, *G. catenatum*-group and *N. scintillans*. The total number and proportion of reads classified into ancient and default via HOPS *sed*aDNA damage analysis per site (based on HABbaits1). The proportion of ancient reads is a measure of '% *sed*aDNA damage' for each of the three species (in italics).

Total	MCS3		GC2S1	
Taxa	**Ancient**	**Default**	**Ancient**	**Default**
Alexandrium spp.	30	1381	22	129
microreticulate G. catenatum-group	4	97	95	252
N. scintillans	1	213	11	49
Proportion (%)	**MCS3**		**GC2S1**	
Taxa	**Ancient**	**Default**	**Ancient**	**Default**
Alexandrium spp.	*2*	98	*15*	85
microreticulate G. catenatum-group	*4*	96	*27*	73
N. scintillans	*0.5*	99.5	*18*	82

4   Discussion

Our analysis of dinoflagellate *sed*aDNA shows that ancient DNA from the genera *Alexandrium* and *Gymnodinium* is preserved in marine sediment for thousands of years. In contrast, *N. scintillans* DNA was detected primarily in recent sediments. The up to 60-fold increase in dinoflagellate sequences achieved using hybridisation capture enrichment with the HABbaits1 array demonstrates the value of this approach for HAB focused *sed*aDNA studies. Additional optimisation, such as adjusting the hybridisation protocol (hybridisation temperature and time, Horn et al., 2012; MyBaits, 2018), and increased sequencing depth, will make it possible to maximise the ancient genetic signal of individual marine species over millennial timescales.

4.1 Considerations for interpretation of Alexandrium, Gymnodinium and Noctiluca
sedaDNA detection

Due to the short length of ancient DNA reads (mean ~56 bp; shotgun data post-filtering), we
expected a degree of uncertainty in their assignment to reference sequences, at a species level.
Our dummy data processing runs containing fragments of taxonomic marker genes of the three
target dinoflagellates *Alexandrium* spp., *Gymnodinium* spp., and *N. scintillans* provided a
useful means to assess the level of species assignment uncertainty on a taxon-by-taxon basis.
For *Alexandrium,* the dummy sample analyses indicated our short sequences could not be
confidently assigned to species, resulting in our use of a conservative genus-level
interpretation. In Tasmanian waters, this genus includes *A. affine, A. australiense, A. catenella,*
A. margalefi, A. ostenfeldii, A. pacificum and *A. pseudogonyaulax* (Hallegraeff et al. 1991,
Bolch & de Salas 2007, Ruvindy et al. 2018). Resolving this complex at species or even
genotype-level could provide valuable information on the history and distribution of toxic
members of this genus. However, given the low confidence in assigning the *Alexandrium*
dummy sequences below genus level, this is beyond the scope of this study and would require
optimisation of RNA bait-design, hybridisation capture (*e.g.*, increased input material), and/or
gene assembly to generate longer DNA fragments, in order to achieve more reliable taxonomic
assignments.

It was possible to make more confident species level assignments for *N. scintillans* due to the
uniquely monospecific nature of this genus (Gómez et al., 2010). The few mis-assignments
noted were 18S rRNA sequences of our target dinoflagellates (Supplementary Material Note
3, Supplementary Material Fig. 3), reinforcing our conclusions that this marker gene is too
conserved to confidently assign very short dinoflagellate reads. While the HABbaits1 design
included multiple marker genes for harmful dinoflagellates occurring off Tasmania, future
RNA bait designs focused on the LSU-rRNA and rRNA-ITS are likely to be more effective
and lead to higher confidence in species-level assignment of *sed*aDNA from this region (and
other locations; Bolch, 2022).

The target gene regions included in the HABbaits1 array were sufficiently variable to
distinguish gymnodinoid dinoflagellates, including the microreticulate group of species that
includes *G. catenatum*. While exact sequence matches to *G. catenatum* reference sequence
DQ785882 were rare, >60% were within 3 bp of this reference, and in regions with sufficient
variation to distinguish between microreticulate-group species, including the most closely
related *G. nolleri* and *G. microreticulatum* (Supplementary Material Fig. 5, Supplementary
Material Table 3). The reference sequence DQ785882 was derived from a Korean *G. catenatum*
isolate, therefore, the observed differences may partly be attributed to global variations in
rRNA sequence of *G. catenatum* (Bolch and De Salas, 2007; Band-Schmidt et al. 2008). An
alternative is that processed (de-duplicated) *sed*aDNA data effectively represents a single
fragment from the original genome of the target species. Dinoflagellate genomes possess many
thousands to millions of rRNA gene loci (Ruvindy et al., 2023) considerable intra-genomic

sequence heterogeneity (Gribble and Anderson, 2006; Miranda et al., 2012), multiple sequence
classes and the presence of pseudogenes (Scholin and Anderson, 1994). Numerous low
frequency sequence variants have also been detected in cultures established from single cells
(Miranda et al. 2012). As a result, the single-based variation we noted among *sed*aDNA reads
may result from the *sed*aDNA pipeline faithfully retaining different single nucleotide mutations
at different rRNA loci from the original genomes.

Both our dummy data and manual fragment mapping indicated an assignment to the
microreticulate group of gymnodinoid species at high confidence, but the resolution of the five
species is partly dependent on both fragment length and gene region/location of each *sed*aDNA
read. The net effect is that only a proportion (up to 33%) of microreticulate species reads could
be unambiguously assigned to an individual species, but that those that could be assigned to
species-level using the *Gymnodinium*-only database are robust. Based on our validation studies
(Supplementary Material Notes 4,5), HABbaits1 can be feasibly applied to gymnodinoids more
generally, and other dinoflagellate genera with similar levels of rRNA-ITS and LSU-rRNA
sequence variation between species. The mapping to reference sequences also highlighted the
potential for assignment artefacts arising from limited database coverage. Robust species
assignments of microreticulate gymnodinoids, including toxic *G. catenatum*, was only possible
after addition of the new D3-D10 LSU-rRNA reference sequences. The improved target gene
coverage in the database allowed detection of the non-toxic related species *G.
microreticulatum*, which is uncommon but widely distributed in both Australian and
Tasmanian waters (Bolch and Hallegraeff, 1990; Bolch and Reynolds, 2002), and sequences
assigned to *G. nolleri* which had not been previously identified from the Australasian region
by microscopy. While *G. nolleri* appears confined primarily to coastal European waters, it is
conceivable this species is distributed very widely but is rare/cryptic in Tasmanian waters. It
may also have been potentially more abundant in the past, as shown in the Baltic Sea where
increased abundance of *G. nolleri* resting cysts coincided with the Medieval Warm Period,
affecting the north Atlantic region from circa 800-1250 AD (Thorsen et. al., 1995).

The *sed*aDNA approach used here allowed the authenticity of the ancient sequences recovered
for the three target taxa to be assessed. While % *sed*aDNA damage was low ($\leq$4%) at the
inshore site MCS3-T2, this is consistent with the earlier studies showing that eukaryote
*sed*aDNA damage is very low in the upper ~35 cm of sediment at this site (Armbrecht et al.,
2021). At GC2S1 (longer offshore core), the increased *sed*aDNA damage observed (up to 27%
for *G. catenatum*) indicate that the recovered sequences from *Alexandrium*, the microreticulate
gymnodinoids and *N. scintillans* are consistent with authentic *sed*aDNA, except for the samples
from the bottom of the core, which were potentially contaminated by seawater during core
retrieval.

4.2   Alexandrium
The distribution of *Alexandrium sed*aDNA reads throughout the offshore core, including the
deepest samples, indicates that *Alexandrium* has been present in the area during the past ~9 000
years. The HABbaits1 data also show that *Alexandrium* has recently increased in relative
abundance in inshore waters (from ~15 years ago; circa 2003), but has remained in low relative
abundance offshore (<3%).
542
Compared to microreticulate gymnodinoid species that are typically rare in coastal sediments
(<2% of cysts; Bolch and Hallegraeff, 1990), *Alexandrium* species can be prolific producers of
resting cysts (approximately 40% of vegetative cells; Anderson et al., 2014). Live resting cysts
are resistant to chemical and biological attack, protecting DNA from degradation during
transport to the bottom sediment. Some cysts can remain dormant but viable in anoxic
sediments for >100 years (Feifel et al., 2012), indicating functional and undegraded DNA
survival for centuries. The patterns of abundance observed in *sed*aDNA mirror trends observed
in microscopy-based counts of *Alexandrium* resting cysts from the same cores (Paine et al., in
review). The highest *Alexandrium* cyst concentrations were observed in the top 35 cmbsf at
MCS3-T2 (inshore), while low concentrations of *Alexandrium*-like cysts were observed
throughout the GC2S1 core (offshore). Very few *Alexandrium* sequences were detected
offshore, but the trend towards increased *sed*aDNA damage with depth (especially at GC2S1)
supports the authenticity of the ancient reads.
556
Unfortunately, our HABbaits1 enriched *sed*aDNA was unable to discriminate between
Alexandrium species (see section 4.1). However, the relative abundance patterns support the
view that toxic *Alexandrium* species (*A. catenella* and *A. pacificum*) are endemic in the
Tasmanian region, but are cryptic, low abundance species not previously distinguished from
morphologically similar and generally non-toxic *A. australiense*. Their recent increased
relative abundance in inshore cores also support the view that emergence of blooms from 2012
onwards have been stimulated by climate-induced changes in environmental conditions
(Condie et al., 2019).
565

566   *4.3   Gymnodinium catenatum and microreticulate species*
567
The HABbaits1 array detected microreticulate gymnodinoid sequences, including *G.
catenatum*, in both cores (MCS3-T2; GC2S1). Inshore, presence was detected from the most
recent sediments (MCS1 2 cmbsf, ~2 years ago, *i.e.,* 2015), in contrast to the offshore core
where they were detected from 5 and 20 cmbsf (~100 and ~700 years ago, respectively), and
sporadically throughout GC2S1 up to 209 cmbsf (~7 638 years ago). In 1987 (30 years before
the sediment core collection), *G. catenatum* was reported as representing 1% of total
dinoflagellate cysts in surface sediments of Spring Bay, close to the inshore core site MCS3
(Bolch and Hallegraeff, 1990). The closest sample we analysed to this was MCS3-T2 15 cmbsf
(~30 years ago), however, no *G. catenatum sed*aDNA was detected, raising the possibility that

any *G. catenatum sed*aDNA preserved was too low to be detected. In future, more thorough
sequencing efforts could resolve this issue.
579
After refining species assignments using the new *Gymnodinium*-only reference database and
detailed read verification, we detected *G. catenatum sed*aDNA sporadically through core
GC2S1 (5, 20, 50, 209 cmbsf, corresponding to ~100, ~700, ~2 300, and ~7 638 years ago;
deepest sample excluded). This extends the likely presence of this species in the region well
beyond the proposed 1970s introduction based on analysis of resting cyst abundance in dated
cores from the neighbouring Huon River (McMinn et al., 1997). Instead the *sed*aDNA approach
indicates *G. catenatum* is more likely to be endemic but previously cryptic. While rDNA-ITS
sequence polymorphisms indicate a link between Australasian populations and the Seto Inland
Sea in Japan (Bolch and DeSalas, 2007), our findings favour proposed alternative hypotheses
that recent dispersal between these areas could equally have been from Australasia to Japan
(Bolch and DeSalas, 2007), or, more likely, the rDNA polymorphisms are part of a wider global
biogeographical pattern of natural dispersal over many thousands of years (John et al. 2018).
While the number of verified *G. catenatum* reads was very low (9 reads), their unambiguous
identification highlights the known problem of overlooking low-cyst producing cryptic species
via microscopy. In a companion palynological survey of the same sediment samples only 3
microreticulate cysts were found out of a total of 4 273 dinocysts counted, and was confined to
0 – 30 cmbsf of MCS3-T2 (inshore; Paine et al, in review).
597

598 *4.4   Noctiluca scintillans*
599
The HABbaits1 approach has revealed the presence of *N. scintillans* inshore of Maria Island
during the last ~30 years, as well as traces of this dinoflagellate in recent sediments offshore
(0 – 1.5 and 5 – 6.5 cmbsf; <100 years old). Inshore, the highest relative abundances of *N.*
scintillans were detected in recent sediments (from the year 2010 onwards; 6 – 7.5 and 12 –
13.5 cmbsf), indicating blooms occurred within the last 21 years. This is consistent with
observational records that *N. scintillans* was first detected in Tasmania in 1994 and from 2010
onwards blooms have increased in both frequency and intensity (Hallegraeff et al., 2019).
607

To our knowledge, this is the first authenticated *N. scintillans sed*aDNA from a coastal
ecosystem, demonstrating the sensitivity of the HABbaits1 approach for detecting plankton
community change of fragile non-fossilising species from the marine sediment record. The
presence of *N. scintillans sed*aDNA in recent sediments at the offshore site confirmed that its
DNA has been preserved at the seafloor, after sinking through a 104 m water column. The
absence of *N. scintillans sed*aDNA from older sediments (>100 years) at the offshore site
(excluding the presumably contaminated deepest sample of GC2S1), indicates that DNA of
this soft-bodied, non-cyst-forming dinoflagellate likely does not preserve well and may pose
limitations to its analysis using *sed*aDNA records (see also Supplementary Material Note 6,
Supplementary Material Fig. 6). Future investigations could target sediment cores from

Sydney, where this species has been present closer to shore for longer, and the rate of damage
of *N. scintillans sed*aDNA might be estimated over longer time scales. The application of
similar methods to the putative range expansion of green *Noctiluca* (with green algal
symbionts) into the Arabian Sea (Gómez et al., 2014) would also be of considerable interest.
622
623

Acknowledgements
We are grateful to Brian Brunelle from Arbor Biosciences, USA, for his expert assistance with
designing the HABbaits1 capture array. We thank Oscar Estrada-Santamaria, Steve Richards,
Holly Heiniger, Nicole Moore and Steve Johnson for their help and advice during DNA
extractions, library preparation, and hybridization capture. We are grateful to Raphael
Eisenhofer, Vilma Pérez, Yassine Souilmi, Yichen Liu, and Ron Hübler for their help with
bioinformatic analyses. We thank the Marine National Facility, the crew of RV *Investigator*
voyage IN2018_T02 and the scientific voyage team for their support during field work [2018
MNF Grant H0025318]. We thank Mr Shikder Saiful Islam (IMAS, University of Tasmania)
for generating additional rRNA gene sequencing from microreticulate species of
Gymnodinium. We thank Henk Heijnis (ANSTO) for logistical support and his contributions
to secure funding for this project. This study was funded through the Australian Research
Council [ARC Discovery Project DP170102261]. A.C. was funded by ARC Laureate
Fellowship FL140100260. LA was supported by an Australian Research Council (ARC)
Discovery Early Career Researcher (DECRA) Fellowship (DE210100929).

Author contributions
L.A. designed and carried out *sed*aDNA laboratory work, bioinformatic analyses and
coordinated the manuscript writing. G.H., C.B., and C.W. collected and provided the core
samples. B.P. carried out dinocysts analyses. L.A., G.H. and B.P. wrote the first draft of the
manuscript. C.B. led the supplementary *Gymnodinium* rRNA gene sequencing, carried out
reference sequence alignments, *sed*aDNA fragment mapping, species assignment validation
and additional figures, and revised subsequent manuscript drafts. C.W. provided sediment core
dating. A.C. contributed guidance on hybridization capture design and ancient DNA analyses.
A.C., G.H., A.M. and C.B. developed the conceptual approach and secured funding for the
project. All co-authors contributed to writing, data interpretation, comment and revision on
manuscript drafts, and final editing before submission for publication.
651

652   **Competing Interests:**
653   The authors declare that there are no competing financial interests in relation to the work
654   described.

References:

Anderson, D.M., Jacobson, D.M., Bravo, I., Wrenn, J.H. 1988. The unique, microreticulate cyst of the naked dinoflagellate *Gymnodinium catenatum* Graham. Journal of Phycology 24, 255-262.

Anderson, D.M., Keafer, B.A., Kleindist, J.L., McGillicuddy, D.J.,Martin, J.L., Norton, K, Pilskaln, C.H., Smith, J.L., Sherwood, C.R., Butman, B. 2014. *Alexandrium fundyense* cysts in the Gulf of Maine: long-term time series of abundance and distribution, and linkages to past and future blooms. Deep Sea Research Part 2 Topical Studies in Oceanography 103, 6–26.

Armbrecht, L. 2020. The potential of sedimentary ancient DNA to reconstruct past ocean ecosystems. Oceanography, https://doi.org/10.5670/oceanog.2020.211.

Armbrecht, L., Herrando-Pérez, S., Eisenhofer, R., Hallegraeff, G.M., Bolch, C.J.S., Cooper, A. 2020. An optimized method for the extraction of ancient eukaryote DNA from marine sediments. Molecular Ecology Resources 20, 906-919.

Armbrecht, L. Hallegraeff, G., Bolch, C.J.S., Woodward, C., Cooper, A. 2021. Hybridisation capture allows DNA damage analysis of ancient marine eukaryotes, Scientific Reports 11, 3220.

Bennett, G. 1860 Gatherings of a naturalist in Australasia, Currawong Press, Australia, (1982) ISBN.

Blaauw, M., Christen J.A., Lopez, M.A., Vazquez J.E., Gonzalez O.M., Belding T., Theiler J., Gough, B., Karney, C. 2019. Age-Depth Modelling using Bayesian Statistics v. 2.3.6, https://CRAN.R-project.org/package=rbacon.

Blackburn, S.I., Bolch, C.J., Haskard, K.A., Hallegraeff, G.M. 2001. Reproductive compatibility among four global populations of the toxic dinoflagellate *Gymnodinium catenatum* (Dinophyceae). Phycologia 40, 78-87.

Bolch C.J.S. 2022. Impact of molecular approaches on dinoflagellate taxonomy and systematics. *In: L. Clementson, R. Eriksen & A. Willis (Eds) Advances in Phytoplankton Ecology: Applications of Emerging Technologies*, pp. 81-117. Elsevier, Amsterdam (ISBN: 9780128228616; eBook ISBN: 9780128230299).

Bolch, C.J., Hallegraeff, G.M., 1990. Dinoflagellate cysts in recent marine sediments from Tasmania, Australia. Botanica Marina 33, 173-192.

Bolch, C.J., Negri, A.P., Hallegraeff, G.M. 1999. *Gymnodinium microreticulatum* sp.nov.

698   (Dinophyceae): a naked, microreticulate cyst producing dinoflagellate, distinct from
699   *Gymnodinium catenatum* Graham and *Gymnodinium nolleri* Ellegaard et Moestrup.
700   Phycologia 38, 301-313.
701
702   Bolch, C.J., Reynolds, M.J. 2002. Species resolution and global distribution of microreticulate
703   dinoflagellate cysts. Journal of Plankton Research 24, 565-578.
704
705   Bolch, C.J., de Salas, M.F. 2007. A review of the molecular evidence for ballast water
706   introduction of the toxic dinoflagellates *Gymnodinium catenatum* and the *Alexandrium*
707   *"tamarensis* complex" to Australasia. Harmful Algae 6, 465-485.
708
709   Condie, S.A., Oliver, E.C.J., Hallegraeff, G.M. 2019. Environmental drivers of unprecedented
710   *Alexandrium catenella* dinoflagellate blooms off eastern Tasmania, 2012–2018. Harmful
711   Algae 87, 101628.
712
713   Dela-Cruz, J., Ajani, P., Lee, R., Pritchard, T. and Suthers, I., 2002. Temporal abundance
714   patterns of the red tide dinoflagellate *Noctiluca scintillans* along the southeast coast of
715   Australia. Marine Ecology Progress Series 236, 75-88.
716
717   Dela-Cruz, J., Middleton, J.H. and Suthers, I.M., 2003. Population growth and transport of the
718   red tide dinoflagellate, *Noctiluca scintillans*, in the coastal waters off Sydney Australia, using
719   cell diameter as a tracer. Limnology and Oceanography 48, 656-674.
720
721   Edwards, L., Wilson, K., Veitch, M.G.K., 2018. An outbreak of Paralytic Shellfish Poisoning
722   in Tasmania. Communicable Diseases Intelligence 42 (PII: S2209-6051(18) 00004-0)
723
724   Feifel, K.M., Moore, S.M. and Horner, T.A., 2012. An *Alexandrium* spp. cyst record from
725   Sequim Bay, Washington State, USA, and its relation to past climate variability. Journal of
726   Phycology 48, 550-558
727
728   Gomes, H., Goes, J.I., Matodkar, S.G.P., Buskey, E.J., Basu, S., Parab, S., Thoppil, P. 2014.
729   Massive outbreaks of *Noctiluca scintillans* blooms in the Arabian Sea due to spread of hypoxia.
730   Nature Communications 5, 4862.
731
732   Gribble, K.E., Anderson, D.M. 2006. Molecular phylogeny of the heterotrophic
733   dinoflagellates, *Protoperidinium*, *Diplopsalis* and *Preperidinium* (Dinophyceae), inferred from
734   large subunit rDNA. Journal of Phycology 42, 1081-1095.
735
736   Gómez, F., Moreira, D., López-Garcia, P. 2010. Molecular phylogeny of noctilucoid
737   dinoflagellates (Noctilucales, Dinophyceae). Protist 161, 466-478.
738

Hallegraeff, G.M., Albinsson, M.E., Dowdney, J., Holmes, A.K., Mansour, M.P., Seger, A. 2019. Prey preference, environmental tolerances and ichthyotoxicity by the red-tide dinoflagellate *Noctiluca scintillans* cultured from Tasmanian waters. Journal of Plankton Research 41, 407-418.

Hallegraeff, G.M., Anderson, D.M., Belin, C., Bottein, M-Y., Bresnan, E., Chinain, M., Enevoldsen, H., Iwataki, M., McKenzie, C.H., Sunesen, I., Pitcher, G.C., Provoost, P., Richardson, A.J., Schweibold, L., Tester, P.A., Trainer, V.L., Yñiguez, A.T., Zingone, A. 2021. Perceived global increase in algal blooms is attributable to intensified monitoring and emerging bloom impacts. Communications Earth & Environment 2, 117. https://doi.org/10.1038/s43247-021-00178-8

Hallegraeff, G.M, Bolch, C.J. 1992. Transport of diatom and dinoflagellate resting spores in ships' ballast water: implications for plankton biogeography and aquaculture. Journal of Plankton Research 14, 1067-1084.

Hallegraeff, G.M., Bolch, C.J., Blackburn, S.I., Oshima, Y. 1991. Species of the toxigenic dinoflagellate genus *Alexandrium* in Southeastern Australian waters. Botanica Marina 34, 575-587.

Hallegraeff, G.M., McCausland, M.A. and Brown, R.K. 1995. Early warning of toxic dinoflagellate blooms of *Gymnodinium catenatum* in southern Tasmanian waters. Journal of Plankton Research 17, 1163-1176.

Hallegraeff, G.M., Schweibold, L., Jaffrezic, E., Rhodes, L., MacKenzie, L., Hay, B., Farrell, H., 2020. Overview of Australian and New Zealand harmful algal species occurrences and their societal impacts in the period 1985 to 2018, including a compilation of historic records. Harmful Algae, 101848.

Head, M.J., Lewis, J., De Vernal, A. 2006. The cyst of the calcareous dinoflagellate *Scrippsiella trifida*: Resolving the fossil record of its organic wall with that of *Alexandrium tamarense*. Journal of Paleontology 80, 1-18.

Herbig, A., Maixner, F., Bos, K.I., Zink, A., Krause, J. and Huson, D.H., 2016. MALT: Fast alignment and analysis of metagenomic DNA sequence data applied to the Tyrolean Iceman. BioRxiv, 050559.

Hogg, A.G., Heaton, T.J., Hua, Q., Palmer, J.G., Turney, C.S., Southon, J., Bayliss, A., Blackwell, P.G., Boswijk, G., Ramsey, C.B., Pearson, C. 2020. SHCal20 Southern Hemisphere Calibration, 0–55,000 years cal BP. Radiocarbon 62, 759-778.

Horn, S. 2012. Target enrichment via DNA hybridization capture. In B. Shapiro and M.
Hofreiter (Eds.), Ancient DNA, methods and protocols (pp. 177–188). New York, Dordrecht,
Heidelberg, London: Springer.

Hübler, R., Key, F.M., Warinner, C., Bos, K.I., Krause, J. and Herbig, A., 2019. HOPS:
Automated detection and authentication of pathogen DNA in archaeological remains. Genome
Biology 20, 1-13.

Huson, D. H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., Tappu, R. 2016.
MEGAN Community Edition - Interactive exploration and analysis of large-scale microbiome
sequencing data. PLOS Computational Biology 12,
e1004957.https://doi.org/10.1371/journal.pcbi.1004957.

John, U., Alperman, T., Nagai, S., Litaker, R.W., Murray, S., Anderson, D.M., Bolch, C., 2018.
Detailed insights into *Alexandrium catenella* (Dinophyceae) (Group 1 genotype) population
structure and evolution. Abstracts 18th International Conference on Harmful Algae, Abstract
Book. p. 324.

John, U., Litaker, R.W., Montresor, M., Murray, S., Brosnahan, M.L., Anderson, D.M. 2014.
Formal revision of the *Alexandrium tamarense* species complex (Dinophyceae) taxonomy: The
introduction of five species with emphasis on molecular-based (rDNA) classification. Protist
165, 779-804.

Jones, R., 1978. Why did the Tasmanians stop eating fish. In: Gould RA (ed), Explorations in
Ethno-Archaeology, pp.11-47. University of New Mexico Press.

Klouch, K.Z., Schmidt, S., Andrieux-Loyer, F., Le Gac, M, Hervio-Heath, D., Qui-Minet, Z.N.,
Quere J, Bigeard, E., Guillou, L., Siano, R. 2016. Historical records from dated sediment cores
reveal the multidecadal dynamic of the toxic dinoflagellate *Alexandrium minutum* in the Bay
of Brest (France). FEMS Microbiology Ecology 92, DOI 10.1093/femsec/fiw101.

McLeod, D. J., Hallegraeff, G. M., Hosie, G.W., Richardson, A. J. 2012. Climate driven range
expansion of the red-tide dinoflagellate *Noctiluca scintillans* into the Southern Ocean. Journal
of Plankton Research 34, 332–337.

McMinn, A.1992. Recent and late Quaternary dinoflagellate cyst distribution on the continental
shelf and slope of southeastern Australia. Palynology 16, 13-24.

McMinn, A., Hallegraeff, G.M., Thompson, P., Jenkinson, A.V., Heijnis, H. 1997. Cyst and
radionucleotide evidence for the recent introduction of the toxic dinoflagellate *Gymnodinium*
catenatum into Tasmanian waters. Marine Ecology Progress Series 161,165-172.

Miranda L.N., Zhuang, Y., Zhang, H., Lin, S. 2012. Phylogenetic analysis guided by
intragenomic SSU rDNA polymorphism refines classification of "*Alexandrium tamarense*"
species complex. Harmful Algae 16, 35-48.

MyBaits   Manual   v.4.01—Hybridization   Capture   for   Targeted   NGS,   2018.
https://arborbiosci.com/wp-content/uploads/2019/08/myBaits-Manual-v4.pdf.

Paine, B., Armbrecht, L., Bolch, C., McMinn, A., Hallegraeff, G.M., 2023. Dinoflagellate cyst
distribution over the past 9 kyrs BP on the east coast Tasmania, southeast Australia.
Palynology, in review.

R Core Team. 2013. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.

Ribeiro, S., Amorim, A., Andersen, T.J., Abrantes, F. and Ellegaard, M. 2012. Reconstructing
the history of an invasion: the toxic phytoplankton species *Gymnodinium catenatum* in the
Northeast Atlantic. Biological Invasions 14, 969-985.

Ridgway, K. and Hill, K. 2009. Ridgway, K. and Hill, K. (2009) The East Australian Current.
In A Marine Climate Change Impacts and Adaptation Report Card for Australia 2009 (Eds.
E.S. Poloczanska, A.J. Hobday and A.J. Richardson), NCCARF Publication 05/09, ISBN 978-
1-921609-03-9.

Ruvindy, R., Bolch, C.J., MacKenzie, L., Smith, K., Murray, S.A. 2018. qPCR assays for the
detection and quantification of multiple paralytic shellfish toxin-producing species. Frontiers
in Microbiology, DOI 10.3389/fmicb.2018.03153.

Ruvindy, R., Barua, A., Bolch, C.J.S, Sarowar, C., Savela, H., Murray, S.A. 2023. Genomic
copy number variability at the genus, species and population levels impacts in situ ecological
analyses of dinoflagellates and harmful algal blooms. ISME Communications **3**, 70.

Scholin, C.A., Anderson, D.M. 1994. Identification of group and strain-specific genetic
markers for globally distributed *Alexandrium* (Dinophyceae). I. RFLP analysis of SSU rRNA
genes. Journal of Phycology 30, 744–54.

Shaw, L.A., Weyrich, L.S., Hallegraeff, G.M., Cooper, A. 2019. Retrospective eDNA
assessment of potentially harmful algae in historical ship ballast tank and marine port
sediments. Molecular Ecology 28, 2476-2485.

Slon, V., Hopfe, C., Weiß, C. L., Mafessoni, F., De la Rasilla, M., Lalueza-Fox, C., … Stewart,
J. R. (2017). Neandertal and Denisovan DNA from Pleistocene sediments. Science 356, 605–
608.

Stockmarr, J. 1971. Tablets with spores used in absolute pollen analysis. Pollen et Spores 13,
615-621.

Taylor, R., 2007. The polemics of eating fish in Tasmania: the historical evidence revisited.
Aboriginal History 31, 1-26.

Thompson, P.A., Baird, M.E., Ingleton, T., Doblin, M.A. 2009. Long-term changes in
temperate Australian coastal waters: implications for phytoplankton. Marine Ecology Progress
Series 394, 75–89.

Thorsen, T., Dale, B., Nordberg, K. 1995. Blooms of the toxic dinoflagellate *Gymnodinium*
catenatum as evidence of climate fluctuations in the late Holocene of southwest Scandinavia.
The Holocene 5, 435-446.

Trainer, V.L., Moore, S.K., Hallegraeff, G., Kudela, R.M., Clement, A., Mardones, J.I.,
Cochrane, W.P. 2019. Pelagic harmful algal blooms and climate change: Lessons from nature's
experiments with extremes. Harmful Algae 91, 101591.

Turnbull, A, Harrison, R., McKeown, S. 2013. Paralytic shellfish poisoning in South Eastern
Tasmania. Communicable Diseases Intelligence 37: E52-4.

Willerslev, E., Cooper, A. 2005. Ancient DNA. Proceedings of the Royal Society of London
B: Biological Sciences 272, 3–16.

Yamaguchi, M., Itakura, S., Imai, I., Ishida, Y. 1995. A rapid and precise technique for
enumeration of resting cysts of *Alexandrium* spp. (Dinophyceae) in natural sediments.
Phycologia 34, 207-214.