Title

Retrograde Control of Cytosolic Translation Targets Synthesis of Plastid Proteins and
Nuclear Responses for High-Light Acclimation.

Marten Moore[1,2], Aaron B. Smith[2], Melanie Wegener[1], Corinna Wesemann[1], Sonja Schmidtpott[1],
Muhammad Ansar Farooq[1,3], Diep Ray Ganguly[2,4], Thorsten Seidel[1], Barry J. Pogson[2§] and Karl-
Josef Dietz[1§].

[1]Biochemistry and Physiology of Plants, Bielefeld University, 33501 Bielefeld, Germany.

[2]ARC Centre of Excellence in Plant Energy Biology, Research School of Biology, The Australian
National University, Canberra, ACT 2601, Australia.

[3]Institute of Environmental Sciences and Engineering (IESE), School of Civil and Environmental
Engineering (SCEE), National University of Science and Technology (NUST), Sector H-12,
Islamabad 44000, Pakistan

[4]CSIRO Synthetic Biology Future Science Platform, Canberra, ACT 2601, Australia

Keywords: High light, chloroplast, retrograde signalling, stress acclimation, transcriptome,
translational control,

§ Joint Senior Authorship, Addresses for Correspondence

Karl-Josef Dietz
Biochemistry and Physiology of Plants – W5-134
Bielefeld University
33501 Bielefeld, Germany
Tel. +49 521 1065589
Fax. +49 521 1066039
E-mail: karl-josef.dietz@uni-bielefeld.de
24
Barry James Pogson
Research School of Biology
Australian National University
Canberra ACT 2601 Australia
Tel. +61 261255629
E-mail barry.pogson@anu.edu.au

List of Abbreviation: ABA: abscisic acid; *A. thaliana*: *Arabidopsis thaliana*; AKIN10: protein
kinase being part of the sucrose non-fermenting kinase complex; Asc: ascorbate; CaM:
calmodulin; CML: calmodulin-like; DEG: differentially expressed gene; DHA:
dehydroascorbate; DREB2A: Dehydration-responsive element binding protein 2; FC: fold
change; GAPC: glyceraldehyde-3-phosphate dehydrogenase 2; GCN2: general control non-
repressible 2; GO: gene ontology; GSH: glutathione; GSSG: diglutathione disulfide; HL: high-
light (800 μmol photons m^{-2} s^{-1}); H_2O_2: hydrogen peroxide; HSE: heat shock element; HSF: heat
shock factor; HSP: heat shock protein; IRES: independent ribosomal entry sites; JA: jasmonic
acid; LL: low light (8 μmol photons m^{-2} s^{-1}); $\log_2$ FC: logarithmic fold change to base 2; MAPK:
mitogen-activated protein kinase; mbSUS: mating-based Split-ubiquitin-System; MBF1c:
Multiprotein Bridging Factor 1C; OPDA: 12-oxo-phytodienoic acid; ORF: open reading frame;
PEG: polyethylene glycol; PET: photosynthetic electron transport; PTM: post-translational
modification; RBP: RNA binding protein; ROS: reactive oxygen species; RuBisCO: ribulose
bisphosphate carboxylase oxygenase; SAP: stress associated protein; SnRK1: SNF1□related
kinase 1; TE: translation efficiency; TF: transcription factor; TOR: target of rapamycin; TPT2:
triose phosphate/phosphate translocator 2; uORF: upstream open reading frame; UTR:
untranslated region; VTC2: GDP-L-galactose phosphorylase;

Abstract

Canonical retrograde signalling comprises information transmission from organelles to the nucleus and in particular controls gene expression for organellar proteins. The need to re-assess this paradigm was suggested by discrepancies between de novo protein synthesis and transcript abundance in response to excess light. Here we uncover major components of a translation-dependent retrograde signalling pathway that first impacts translation and then transcription. The response realization depends on the kinases Mitogen-activated protein kinase 6 (MPK6) and Sucrose non-fermenting 1-related kinase (SnRK1) subunit, AKIN10. Global ribosome foot-printing revealed differential ribosome association of 951 transcripts within 10 min after transfer from low to high light. Despite predominant translational repression, 15 % of transcripts were increased in translation and enriched for chloroplast-localized photosynthetic proteins. About one third of these transcripts, including Stress associated proteins (SAP) 2 and 3, share regulatory motifs in their 5`-UTR that act as binding sites for glyceraldehyde-3-phosphate dehydrogenase (GAPC) and light responsive RNA binding proteins (RBPs). SAP2 and 3 are both translationally regulated and interact with the calcium sensor Calmodulin-like 49 (CML49), which promotes relocation to the nucleus inducing a translation-dependent nuclear stress response. Thus, translation-dependent retrograde signalling bifurcates to directly regulate a translational circuit of chloroplast proteins and simultaneously initiate a nuclear circuit synchronizing retrograde and anterograde response pathways, serving as a rapid mechanism for functional acclimation of the chloroplast.

Introduction

Light impacts plant morphogenesis, development, and metabolism. Gains in biomass and fitness strongly correlate with light-dependent photosynthesis and starch accumulating during the light phase being essential for metabolism and growth during the dark phase. However, excess or high light (HL) leads to an over-reduction of the photosynthetic electron transport (PET) chain and enhances production of reactive oxygen species (ROS) [1]. The natural light environment fluctuates continually on time scales of seconds, hours, days, and seasons. To cope with rapid light fluctuations, plants control their capacity to dissipate excess energy and minimize photo-oxidative damage [2]. ROS produced in excess light are quickly detoxified, for example H_2O_2 by the water-water cycle in an ascorbate-dependent or ascorbate-independent process utilizing ascorbate peroxidases and peroxiredoxins, respectively [3, 4]. On longer time scales, changes in light quantity alter photosystem stoichiometry and increase the photosystem repair capacity [5].

Proteins of the photosynthetic machinery are encoded in the plastid and the nuclear genomes while all antioxidant enzymes are exclusively encoded in the nucleus, as is the majority of plastidial proteins. Transcription, stability, and translation of chloroplast-encoded transcripts strongly depends on nuclear-encoded sigma factors and RNA binding proteins (RBPs) [6]. The reorganization of photosystems and readjustment of the antioxidant systems, therefore, rely on a coordinated synthesis of nucleus- and chloroplast-encoded proteins [7]. Coordination is realized by transmitting information on the metabolic state of the chloroplast as retrograde signals to the nucleus. Alterations in light quality and quantity result in fluctuations of metabolites and signal export from the chloroplast with impacts on transcript abundance within minutes [8-10].

As the majority of chloroplast proteins are nuclear-encoded, several signaling pathways exist for developmental and operational retrograde control of nuclear gene expression. They include signals from the plastoquinone and carotenoid pools, tetrapyrrole biosynthesis, dihydroxyacetone phosphate (DHAP), ROS, ascorbate/dehydroascorbate (Asc/DHA) and glutathione/glutathione disulphide (GSH/GSSG), hormones like 12-oxo-phytodienoic acid (OPDA) that also functions as precursor for jasmonic acid (JA), and xanthoxin as precursor of abscisic acid (ABA), the SAL1/PAP pathway and sugars [3, 9, 11-18]. Rapid light induced transcription requires export of triose phosphates, such as DHAP, via the Triose phosphate/phosphate translocator (TPT2) and Mitogen activated protein kinase 6 (MPK6), which is coupled to the upregulation of a subset of

Apetala 2/Ethylene response factor (AP2/ERF) transcription factors [9]. It is worth noting that

TPT2 co-controls the transmission of the DHAP retrograde signal [19].

To date, retrograde signaling pathways are considered with respect to nuclear gene regulation.

However, direct retrograde control of cytosolic translation remains mostly unexplored, especially

during rapid responses to stimuli. A few studies focused on the effects of translational control

under the effects of the controversially discussed gun mutants [20], or on targeted transcripts

without whole genome analysis [21, 22], which allow a limited scope of regulative mechanism of

cytosolic translation as a consequence of retrograde signalling.

In addition to regulation by phosphorylation/dephosphorylation, redox regulation of the

translational apparatus has been suggested as one such translational control mechanism [23].

Recently, light-induced ROS production has been linked to rapid activation of eIF2α kinase

General control non derepressable-2 kinase (GCN2), thus inhibiting translation initiation [24].

This process also affected the translation of transcripts encoding mitochondrial ATP synthesis

and chloroplast thylakoid proteins.

Post-transcriptional regulation of translation and mRNA stability is known from studies on

circadian rhythmicity and involves light entrainment [25]. Transcripts encoding enzymes of

central metabolism associate with the ribosomes in in a light- and circadian-dependent manner.

The impact of circadian rhythmicity on the steady-state pool of polysome-recruited transcripts

was lost if the shift to darkness or re-illumination occurred out of phase [26]. The energy sensor

kinase AKIN10 entrains the circadian clock depending on light input [27]. AKIN10 is part of the

Sucrose non-fermenting 1-related kinase 1 (SnRK1) complex and acts as a central regulator of

transcription and translation networks via the Target of rapamycin (TOR) complex [28-30].

In addition to temporal retrograde signaling cascades affecting translational regulation,

consideration to spatial effects should be given. Photosynthetic cells in flowering plants usually

have multiple chloroplasts but a single nucleus. The distance between individual chloroplasts and

the nucleus often spans several tens of micrometers, potentiating heterogeneity of nuclear gene

regulation between chloroplasts of varying vicinity to the nucleus. However, each chloroplast has

a surrounding cytosol with translation-competent ribosomes. Retrograde control of translation

may be a potent mechanism for immediate spatial and temporal adjustment of the chloroplast

proteome. Therefore, there is a fundamental need to address the significance and mechanism of translational regulation of light acclimation by retrograde signalling.

Here we applied ribosome foot-printing to measure global protein synthesis in arabidopsis leaves following a 10 min exposure to HL. Results revealed a rapid light-induced reorganization of the translatome, which was perturbed in *mpk6* and *akin10* mutants. A cluster of transcripts were preferentially polysome-associated in HL and encoded primarily for cytosolic proteins involved in translational and chloroplast-targeted proteins involved in assembly, function, and maintenance of the chloroplast. A second cluster revealed simultaneous increases in transcript abundance and polysome association, containing stress-associated transcripts such as *HSF2A*, *DREB2A*, and *HSPs*. Transcripts of these clusters contained conserved sequence motifs in their 5'-UTRs, which was found to increase translation *in vivo*. Focusing on one motif, we identified light-dependent recognition by GAPC and scrutinized the downstream effects of the translational reorganization following two selected transcripts *SAP2* and *SAP3*, whose proteins interact with the calcium sensor CML49 co-regulating further stress responses. The study evidences the effects and identifies regulators of rapid retrograde regulation of cytosolic translation.

Results

Rapid retrograde signaling for translational reprogramming

To investigate the impacts of severe light stress on translation, we combined ribosome profiling with an established light shift protocol to compare differing fold-increases in light intensity on cumulative translation (Figure 1) (*9, 31, 32*). Plants were transferred from normal growth light (NL, 80 µmol photons m^{-2} s^{-1}) or low light (LL, 8 µmol photons m^{-2} s^{-1} for 10 d) to HL (800 µmol photons m^{-2} s^{-1}, NL→HL or LL→HL) representing a 10-fold or 100-fold radiation increase, respectively. Transcripts associated with multiple ribosomes (polysomes) were isolated as described [31] under presence of translational inhibitors cyclohexamide, chloroamphenicol and lincomycin for combinatorial arrest of cytosolic and plastidial ribosomes. Extracts were loaded and run through sucrose gradients, and polysomal profiles recorded during fractionation (Figure 1a). This revealed rapid reorganization of ribosome-associated RNA within 10 min of a 100-fold light increase. Comparison of LL control with 10 and 60 min LL→HL treatments revealed cumulative dynamic changes in ribosome-associated polysome profiles (Figure 1a). The 80S monosome peak (one ribosome per mRNA) was greatly reduced after 10 min of LL→ HL, but

less so by 60 min LL→HL. The polysomal area was unchanged between LL control and 10 min

LL→HL, but increased after 60 min LL→HL. Contrarily to the 100-fold light shift, no time- and

dose-dependent changes in the polysomal profiles could be observed in 10-fold NL→HL and

NL→HL→NL shifted plants.

Quantification of ribosome-associated mRNA isolated from ribosomal fractions in Figure 1a

revealed a progressive repression of polysome association during LL→HL up to 30 min (Figure

1b). To confirm reversibility of the repression, we also quantified ribosome-associated mRNA

during recovery from 10 and 50 minutes light stress (i.e. LL→HL→LL). We observe equally

rapid re-initiation of translation represented by increased ribosome-associated mRNAs within 10

minutes of transferal back to LL following either 10 or 50 min LL→HL (LL→HL→LL). This

indicated that the dynamic regulation is rapidly initiated within 10 min as first timepoint of

significance, which is expressed in a 0.79-fold decrease in 80S (monosomes) peak area and 0.85-

fold decrease in 60S peak area (Figure 1c), resulting in a cumulative down regulation of

ribosome- associated mRNA. Ribosome integrity was explored by staining fractions of

monosomes, low molecular polysomal mass (2 ribosomes) and high polysomal mass ($\geq$3

ribosomes) and imaging via transmission electron microscopy (TEM). Low and high molecular

mass polysomes could be imaged, confirming successful separation of intact monosomes and

polysomes.

Regulatory pathways connect retrograde signals and cytosolic translation

To investigate the upstream regulation leading to the dynamic shift in cytosolic translation, we

queried two potential interactors of translational regulation that operate in the same rapid time

frame. Previously, we demonstrated phosphorylation of MPK6 in response to retrograde

signaling within less than 10 min following the same LL→HL shift treatment [9]. Intriguingly,

MPK6 activity is potentially linked to phosphorylation state of the SnRK1 subunit AKIN10 [29,

32], which acts as negative feedback loop to TOR signaling and the initiation of translation [33,

34]. Therefore, we hypothesized a link between retrograde signaling and translation, through

light-responsive activation of MPK6 and AKIN10 [29]. To test this, we compared changes of

ribosome profiles in *mpk6* and *akin10* and Col-0 wild-type (WT) background after 10 min

LL→HL shift (Figure 2). Under LL control conditions *mpk6* revealed a mildly reduced 60S and

80S peak area in comparison to WT control (0.91-fold for each) and a 33% reduction in

polysome-bound mRNA (Figure 2a). Contrarily, *akin10* exhibited a strong reduction of the 60S

and 80S peak area in comparison to WT (0.67 and 0.52-fold, respectively) with no significant

difference in polysomal area. After 10 min LL→HL, *mpk6* exhibited a closer to WT ribosomal

profile, but with decreases in 60S and 80S area of 45% and 49%, respectively (compared to 15%

and 21% in WT) (Figure 2b). In addition, a strong decline of 43% in polysomal area was

recorded. The profiles of *akin10* did not differ between LL and LL→HL, but regardless of

experimental conditions exhibited a lower 60S and 80S peak area in comparison to WT with an

unaltered level of polysomal area.

In conclusion, the loss of MPK6 and SnRK1 subunit AKIN10 showed alterations to different

aspects of polysomal profiles that were genotype- and light-dependent, cumulatively altering the

translational profile. While *akin10* lost the WT reaction to LL→HL treatment, *mpk6* reflected the

WT reaction in regards of 60S and 80S area, with an additional reduction of polysomal area after

LL→HL treatment.

Regulation of translation outweighs transcriptional changes in early light acclimation

We quantified the cumulative changes in translation in response to LL→HL regulation by RNA

sequencing on polysome-associated transcripts. Ribosome footprints (RPF) were generated from

three independent experiments of LL and LL→HL-treated plants. RNase If treatment digested the

unprotected sequence stretches [35]. Both the RPF (translatome) and total purified RNA

(transcriptome) were quantified by ribosomal RNA-depleted RNA sequencing [36]. The

generation of footprints in the presence of CHX, CHL and LIN resulted in oligonucleotide read

lengths with two local optima of 24-26 (footprint size from 70S ribosomes in plastids and

mitochondria) and 28-34 nt lengths (cytosolic footprint size from 80S ribosomes), documenting

sufficient capture of arrested plastidial and cytosolic footprints (Suppl. Fig 1a). The read frame

periodicity of the footprints (Suppl. Fig 1b), as well as the read frame at start and end sites that

corresponded to independent studies using a combination of CHX and CHL for translation

inhibition [37, 38].

Performing differential gene expression analysis on ribosome-footprints confirmed the

observation of global repression of translation with 793 transcripts (76 % of total DEGs) that

were significantly down regulated. A set of 92 transcripts (9 %) remained unchanged and

abundance of 158 (15 %) increased (Suppl. Fig 2a). Conversely, total RNA remained mostly

unchanged with only 121 and 83 transcripts being transcriptionally up- and down-regulated,

respectively. Based on our analysis 4.8-fold more transcripts were regulated at the translational

than transcriptional level after 10 min LL→HL.

Evidence for a translational retrograde circuit enriched in photosynthesis proteins

In order to reveal distinct regulative patterns not following the cumulative translational response,

we further investigated the 24 % of transcripts showing significant ribosome association after

LL→HL. The transcripts showing significantly differential expression in the ribosome-bound or

free mRNA were examined for cluster formation by plotting their relative change in ribosome-

association versus the change in total amount (FC ribosomal mRNA LL→HL/LL / FC total

mRNA LL→HL/LL). This evaluation allowed for tentative distinction of four subsets of

transcripts with variant regulatory responses (Figure 3a). Transcripts in Cluster 1 revealed

increased ribosome association without changes in total mRNA abundance (119 DEGs). Clusters

2 and 3 comprise transcripts that upon transfer to HL simultaneously changed in the ribosome-

associated and total mRNA, 62 up and 123 down regulated DEGs, respectively (Figure 3a;

clusters 2 and 3). Association with ribosomes of Cluster 4 transcripts decreased in HL, but were

not significantly altered in total RNA fold change (FC) (466 DEGs).

Cluster 1 transcripts that were solely increased in ribosome association were five-fold enriched in

transcripts coding for chloroplast localized proteins according to SUBA4 [39]. GO analysis of

cluster 1 revealed a 22.9-fold enrichment for the GO term 'photosynthesis, light reaction' in

comparison to the whole genome (FDR p<4.32E- 07). It encompassed transcripts coding in

particular for photosystem light harvesting proteins, including minor (AT3G47470) and major

light harvesting proteins (AT4G10340), subunits of photosystem I (AT2G46820), one of the two

nuclear encoded cytochrome b6f subunits (AT2G26500) and NADPH oxidoreductase subunits

participating in electron transport (AT1G06690) as well as small subunits of ribulose

bisphosphate carboxylase oxygenase (RuBisCO, AT5G38410) (see Suppl. Dataset 1, cluster 1).

In addition to photosynthetic proteins, this set included transcripts for proteins participating in

assembly, damage, and repair. For example chloroplast localized HSC70-1 (AT5G02500), which

is involved in protein import; chloroplast vesiculation factors (AT4G32150) participating in

partitioning damaged proteins for proteolysis; and PYG7/YCF37 (AT2G23670) which is required

for PSI assembly/repair. The transcripts coding for cytosolic proteins were 9.36-fold enriched in

GO terms for translation (FDR p<4.19E- 07), e.g., proteins of ribosomal large subunit

(AT3G08520).

Transcripts of cluster 2 with increased ribosome-associated and total mRNA (Figure 3a), were
enriched in 'abiotic stress'-related GO terms, namely response to heat (71.25-fold enriched in
comparison to genome, FDR p< 4.53E- 41); response to hydrogen peroxide (94.93-fold, FDR
p < 1.32E-16); cellular response to unfolded protein (76.03-fold, FDR p<1.52E- 06); and 'de
novo' protein folding (49.13, FDR p< 2.28E- 04). This cluster included small (HSP20-likes) and
large heat shock proteins (HSP70, HSP101), their organizing factors (HOP) as well as key
regulatory transcription factors HSFA2 (AT2G26150), HSF2BA (AT5G62020), HSFB2B
(AT4G11660) and DREB2A (AT5G05410) involved in high light acclimation (see Suppl.
Dataset 1, Cluster 2).

Cluster 3 consisted of 123 transcripts with decreased share in the ribosomal and total fractions
and was enriched in GO terms for 'negative regulation of transcription' (8.91-fold enriched in
comparison to the whole genome, FDR p<5.73E- 04); 'response to salicylic acid' (9.89-fold
enriched, FDR p<1.75E- 02) and 'response to oxygen levels' (10.32-fold enriched, FDR
p<5.34E-05). These transcripts included transcription factors WRKY70 (AT3G56400), ERF107
(AT5G61590), as well as growth regulators OFP15 and OFP16 (AT2G36050, AT2G32100,
respectively) (see Suppl. Dataset 1, Cluster 2).

Transcripts in Cluster 4 had decreased ribosome association but lacked significant transcriptional
changes (Figure 3a). This largest subset among all clusters with 465 transcripts was enriched in
multiple GO terms, e.g., 'response to auxin' (6.39-fold, FDR p<4.00E-03), 'tetrapyrrole
metabolic process (8.98-fold, FDR p<2.32E-04) and 'response to far red light' (4.07-fold, FDR
p<8.24E-03), but a predominant response as observed in cluster 1 and 2 was undetectable.

Plastid translation regulated in response to rapid light shift

While cytosolic translation preferentially recruited photosynthetic and stress-related transcripts in
HL, analysis of plastome-encoded genes revealed only the chloroplast-encoded PSBA coding for
the PSII subunit D1 as translationally upregulated. Mitochondrial transcripts were missing from
the translationally upregulated group. Conversely, 36 plastome- and 16 mitochondriome-encoded
DEGs, representing 42.9 % and 34 % of the detected organellar transcripts, displayed a lower
ribosome association in cluster 4. The translationally down regulated plastome-encoded
transcripts were 50-fold enriched in GO terms for translational processes (30 % in comparison to
0.6 % of genome representation) and 122-fold in ribosome assembly (11 % compared to 0.09 %),

but also 100-fold in photosynthetic light reactions (11 % to 0.11 %) represented by YCF3 and
YCF4 (ATCG00360, ATCG00520), PSII CP47 reaction center protein (ATCG00680), and NDH-
H (ATCG01110). No enrichment in particular GO terms could be detected in the mitochondrial
DEGs, but 6 of the 16 DEGs encoded for NADH dehydrogenases.

Cluster 1 and 2 transcripts eludes global translational repression and MPK6 and AKIN10
regulation

We explored the relationship of transcript abundance (sum of CPM across all samples) and
ribosome association (FC ribosome-associated mRNA) of the transcripts for each cluster to query
how the mRNA abundance, or availability impacts ribosome association (Figure 3b). Contrarily
to the classical translation efficiency (TE) term, this allowed to de-convolute the effects of
ribosome loading and unloading from differential transcription (Suppl. Fig. 2c,d). Transcripts of
cluster 1 and 4 displayed an inverse correlation between FC of ribosome association and total
transcript amount; less abundant transcripts displayed a higher FC in ribosome association
(Figure 3b). On the other hand, this relationship was absent in clusters 2 and 3 as FC in ribosome
association were independent of total transcript abundance, indicating that ribosome association
was dependent on specific transcripts and not purely their abundance.

Given the altered cumulative translational response observed in the ribosome profiles during
LL→HL in *mpk6* and *akin10*, we assessed whether they specifically contribute to cluster 1 and 2
responses. The role of MPK6 and AKIN10 on ribosome association of representative transcripts
of cluster 1 and 2 after 10 min LL→HL was quantified via qPCR of total and ribosome-
associated transcripts in *mpk6* and *akin10* (Suppl. Fig. 3). Interestingly, no uniform pattern of
deregulation could be observed for this subset of transcripts in the mutants (Suppl. Fig. 3a-d).
Querying genotype- and treatment-specific effects via principal component analysis indicated
that the treatment and not the genotype accounted for ribosomal association of cluster 1 and 2
representatives (Suppl. Fig. 3e-f). Thus, cluster 3 and 4 represent the cumulative ribosomal
regulation as seen in Figure 1c, while cluster 1 and 2 are not represented in the cumulative
changes and are unaffected by AKIN10 and MPK6.

Analyzing cluster 1 and 2 transcripts for differential ribosomal association at upstream open
reading frames (uORF) revealed only 5 transcripts with shifts to uORFs (Suppl. Table 3), thus no
common pattern of changed reading frames of cluster 1 and 2 could be observed. Interestingly,

comparing the total RNA-seq read depth with the RPF read depth for enriched maxima of cluster
1 & 2 transcripts revealed enriched ribosomal occupation at certain positions, indicative for
ribosome stalling sites (Suppl. Fig. 4).

The increased ribosome association of transcripts of cluster 1 and cluster 2 is in juxtaposition to
the global down-regulation of translation after 10 min of LL→HL. This together with the strong
overrepresentation of distinct GO terms in both clusters and the decoupling of transcript
abundance and FC in ribosome association, led to the hypothesis that transcript-specific
recruitment is taking place. We hypothesized that inherent properties of the mRNA must be co-
regulating preferential translation. Therefore, we considered the 5'- and 3'-UTRs of the cluster 1
and 2 transcripts, separately, and combined as translationally upregulated transcripts, for
detecting conserved mRNA structures and sequences.

Translationally regulated transcripts are enriched in RNA motifs in their UTRs

Reorganization of translation requires unloading and reloading of ribosomes with different
transcripts. This process is instigated by translation initiation factors acting in concert with
mRNA regulatory sequences and/or structures. The 5'-UTR and 3'-UTR possess features that
influence the initiation rate, like independent ribosomal entry sites (IRES) [40, 41]. Therefore, the
5'- and 3'-UTRs of the cluster 1 and 2 were screened for conserved motifs enriched in
comparison to all TAIR annotated 5'- or 3'-UTRs using the MEME-suite [42]. The search
identified 52 motifs in the 5'-UTRS of transcripts in clusters 1 and 2, of which 21 were in
common. Three motifs were selected based on their over-representation in cluster 1 and 2 (M1:
TCTCCGGAGAA, M2: TTAGGGTT and M3: TCGCCG, Figure 4).

Motif M2 was previously identified as stress-responsive 5'-UTR element that confers preferential
translation in vitro [43]. M1 and M2 were present 40- and 34-times in the 5'- and 3'-UTRs of
cluster 1 and 2 transcripts, accounting for 31% and 26% of transcripts, respectively (Suppl. Fig.
5b). M2 is present in transcripts that are exclusively translationally regulated and not
transcriptionally (cluster 1), whereas M1 and M3 are also present in transcriptionally regulated
transcripts (cluster 2). Transcripts carrying M1 and M2 had a median FC of relative ribosome
association of 2.09 and 1.97, respectively, compared to the median of 2.96 for all cluster 1
transcripts. Transcripts carrying M3 had a higher FC of 3.57. Analysis of motif position in the

respective 5'-UTRs of motif-carrying transcripts revealed a preferred position at the 3'-end in longer 5'-UTRs, while short 5'-UTRs showed no positional preference (Suppl. Fig. 5a).

Calculating the secondary structure via RNA Shapes Studio [44] revealed that the motifs do not overlap with any conserved secondary structure (Dot-Bracket annotation, Suppl. Fig. 5a, Suppl. Fig. 6). Heat-responsive transcripts like *HSFB2B* and *HSP20*-like proteins, as well as other stress-dependent transcripts, were overrepresented among the motif-carrying transcripts. Interestingly the 5'-UTRs of the *Stress Associated Protein 3* (*SAP3*) carries all three motifs and that of *SAP2* carries M2 (Suppl. Fig. 5a,b). SAPs belong to the family of zinc finger proteins that contain A20/AN1 zinc-binding domains. They are structurally and functionally conserved among plants [45]. *SAP* transcript amounts increase in response to multiple stresses, and overexpression of specific SAPs in planta enhances biotic or abiotic stress tolerance or photosynthetic capacity [46]. To assess importance of the motifs on an evolutionary scale, we queried their presence in the rice genome. *OsSAP8* and *OsSAP9* were identified as putative homologues of *AtSAP2* and *AtSAP3*, in which we detected minor variants of all three motifs in the corresponding 5'-UTRs (Suppl. Fig. 8a). Additionally, we observed the preferential association of transcripts for *OsSAP8* and *OsSAP9* with polysomes upon L→H transfer in rice (Suppl. Fig. 8b).

Enriched 5' UTR motifs enhance HL-dependent protein synthesis and are targeted by RNA binding proteins

To confirm a regulatory role of the three motifs towards translational regulation, we performed a protoplast-based luciferase assay. Cassettes carrying the M1, M2, or M3 motifs in triplicate repeats were analyzed for translationally regulatory elements and secondary structure and inserted into the *AtUBI10* 5'-UTR (Suppl. Fig. 6) upstream of *luciferin 4-monooxygenase* (*Photinus pyralis*). The verified constructs were transiently co-transfected with pGL4:: *luciferin 2-monooxygenase* (*Renilla reniformis*), as an internal reference, into *A. thaliana* mesophyll protoplasts (Figure 4a). The protoplasts were treated analogously to the whole plant experiments, comparing protoplasts exposed to L→H-transfer for 10 min to those at LL. The normalized luciferase activity from lysed protoplasts transformed with the ubiquitin 5'-UTR without motifs increased 2.5-fold after L→H-treatment (Figure 4a). All motifs conveyed a light induced higher signal than the ubiquitin control with M1-construct causing the highest increase of 5-fold, 1.75-fold for M3 and insignificantly increased for M2 (1.52-fold).

We performed RNA-EMSAs to query for potential RBP-binding properties of the motifs.
Biotinylated M3-RNA was incubated with leaf protein extracts that were either prepared from
plants under LL or exposed to 10 min L→H. Addition of protein extracts from LL exposed leaves
resulted in a pronounced shift of signals in comparison to the biotinylated M3-RNA control
(Suppl. Fig. 7a). Addition of unlabeled competitor M3-RNA strongly suppressed the shift.
Supplementation with extract from 10 min LL→HL-treated plants resulted in a signal
intensification that again was out-competed by competitor M3-RNA. This result suggests RBP
binding capability of M3 with differences in protein signature in extracts from L-light plants and
10 min LL→HL-plants. To identify the potential RBPs interacting with M1 and M2, an
explorative affinity pulldown using total protein extracts from plants under LL and LL→HL and
subsequent SDS-PAGE was performed. The respective enriched bands (Suppl. Fig 7b) were then
analyzed via MALDI-TOF. The identified peptides were filtered for peptides matching the
respective SDS-PAGE band size and a Mascot-Score of > 60 resulting in GAPC2 and ATP
synthase subunit alpha as matches for M1 and M2, respectively.

To confirm this interaction, we performed an RNA-EMSA incubating recombinant GAPC1 with
free consensus M1- and M2-RNA (Figure 4b). As demonstrated by a concentration-dependent
band shift, GAPC1 could be confirmed as an RBP with binding affinity to consensus M1- and
M2-RNA Additionally, GAPC1 did not bind to the motif free 5'-UTR of a transcript that was
transcriptionally upregulated during LL→HL, demonstrating a motif-specific mRNA interaction
(Figure 4b). To confirm RNA-RBP interaction with endogenous M1, recombinant GAPC1,
GAPC2, or GAPDH isolated from rabbit muscle was incubated with native M1 motif of the *SAP3*
5'-UTR with concentration dependent shifts, which could be suppressed by adding unlabeled
competitor RNA (Suppl. Fig. 7c). To further test *ex vivo* interaction in a light dependent context,
native M1 motif of the *SAP3* 5'-UTR was used in an affinity pull down with total protein extract
form LL and LL→HL-treated plants, with subsequent immunodetection of GAPC1 and GAPC2
(Figure 4c). Interestingly, a strong GAPC signal could be detected in the affinity pull down from
LL total protein extract. In a converse manner, the signal from HL protein extract was weak, with
no detectable differences in total GAPC1/C2 content between LL and HL total protein extracts,
strongly suggesting a light-dependent interaction of M1 and GAPC1/C2 (Figure 4c).

Synchronization of translational signaling and nuclear transcription: SAPs interact with
CML49 altering nuclear transcription

GO analysis of cluster 2 transcripts revealed an enrichment for stress-responsive transcription factors and co-factors like *MBF1c*, *HSF2A*, and *DREB2A*. These are known to transcriptionally regulate major stress response genes like *HSPs* and *APX2* [47, 48]. Therefore, we hypothesized that rapid translational regulation of cluster 2 transcripts could integrate *de novo* protein synthesis and transcriptional regulation during HL. Next to the afore mentioned well researched transcription factor involved in major stress responses, other targets of translational control were *SAP2* and *SAP3*. With SAP2 fused to the GAL4-DNA-binding domain as bait, six clones were recovered from prey consisting of *A. thaliana* cDNA library fused to the GAL4 activating domain. Five of these clones coded for Calmodulin-like 49 (CML49). The interaction between CML49, and SAP2 and SAP3 was confirmed using the mating-based-Split-Ubiquitin (mbSUS) yeast-two-hybrid system (Figure 5a). Additionally, we performed fluorescence resonance energy transfer (FRET) confirming interactions between SAP2, SAP3, and CML49 with a high FRET efficiency (> 5-fold of negative control) proving their interaction *in vivo* (Figure 5b,c). Interaction of the SAPs with CML49 resulted in trans-localization to the nucleus (Figure 5b). This interaction was also confirmed for OsSAP8/OsSAP9 and OsCML49a/OsCML49b in rice (Suppl. Fig. 9a,b), thus this interaction appears to be a mechanism conserved in flowering plants.

The preferential association of *SAP2* and *SAP3* with polysomes after LL→HL-transfer and the interaction of SAP2 and SAP3 with CML49 led us to hypothesize that CML49 and SAP2/SAP3 function as regulatory module in this specific environmental context. This is underlined by the stress-dependent expression pattern for *SAP2* and *SAP3* and the constitutively expressed CML49 within arabidopsis and rice (Suppl. Fig. 10a-f). In addition, the similar response of *SAP2* and *SAP3* in the LL→HL-response might indicate some functional redundancy. With these two data-guided hypotheses in mind, we hypothesized that the CML49 knockout should have stronger effects on downstream events during H-light acclimation than deletion of SAP3. To investigate the importance of SAP3 and CML49 during HL acclimation, we determined the transcriptional responses of *cml49* and, *sap3* to 60 min LL→HL (Suppl. Data Set 1). A set of 32 transcripts was more than 3-fold increased or decreased in *sap3* relative to WT, while the corresponding set comprised 214 transcripts in *cml49*.

The data set was examined for transcripts that exhibited stronger downregulation in *cml49* than in *sap3* upon LL→HL-transfer, since *SAP2* and *SAP3* exhibit similar LL→HL responses which could be indicative for functional redundancy. This filter revealed 49 transcripts with the

expected response pattern (Figure 6a). The predominant GO terms belonged to the categories
'response to temperature' and 'response to light stimulus', each of which represented 22 % of the
genes, followed by 'response to hydrogen peroxide' with 11 % and 'circadian rhythmicity' with 6
% (Suppl. Fig. 11). Heat shock proteins and one heat shock factor (*HSFA2*) belonged to the group
of heat responsive elements, *sigma factor 5* and *Pseudo response regulator 9* (*PRR9*) to the light
responsive elements, *Purple acid phosphstase 17* (*PAP17*) and *GDP-L-galactose phosphorylase*
(VTC2) to the group of 'response to hydrogen peroxide'. *Circadian clock associated D 1* (*CCA1*)
and *PRR9* represent two genes associated with the circadian clock. The occurrence of transporters
for phosphate, sugar and sulphate (*PS3*, *SWEET 13*, *GPT2*, *APR3*) in this list indicates a function
of CML49 and SAP3 in regulating metabolite and nutrient distribution. Interestingly, the
Translation elongation factor 1B (*eEF1B*) was also found in this list possibly pointing to
SAP3/CML49-dependent amplification effects in the HL-dependent control of translation.

The next attempt to validate the *SAP3* and *CML49* transcriptional regulation and the kinetics of
HL induction (Figure 6b-f) included more detailed analysis of five targets that were selected from
the list of co-regulated transcripts. *VTC2* showed a transient four-fold upregulation at 10 min
LL→HL in WT, which dissipated by 30 min. Conversely, *VTC2* was steadily downregulated in
sap3 and *cml49* to four-fold below the L-light level (Figure 6 b-f). *HSP20*-like rapidly increased
in WT upon LL→HL-transfer reaching its maximum of 64-fold induction at 30 min LL→HL.
The mutants showed a slower and lower upregulation than WT with significantly reduced
responses at 10 and 30 min LL→HL in *sap3*, and at 60 min in both mutants (Figure 6b-f). The
same trend appeared in WT for *PS3*, *SIG5* and *eEF1B* with a four-fold upregulation after 10 until
60 min of LL→HL shift. The mutants showed a strong and consistent deregulation compared to
WT (Figure 6b-f).

Common cis-elements in promotor region of co-regulated genes

The 49 co-regulated target genes were screened for common cis-elements in the promotor region
in order to find commonalities, such as involved transcription factors required for co-regulation.
A total of 9,459 common cis-elements were identified in the 1000-bp upstream of transcriptional
start site of the 49 co-regulated genes (Suppl. Table 3). Most abundant common cis-elements
were the WBOX, targeting sequences for the WRKY-TFs (2112), followed by sulfur response
(SURE, 1536) and MYB and MYB-related cis-element sequences (1368). Stress related cis-
elements like the sequences overrepresented in light-induced promotors (SORLIPs) and early

responsive to dehydration (ABRE) were also highly abundant with 1278 and 1148, respectively.

bZIP cis-elements (C-box, 232), phosphate starvation (PHR1, 224) and salicylic acid inducible

(LS7, 173) were less abundant.

Is there a translational retrograde – anterograde circuit?

After 10 min LL→HL all plastidial translation was repressed with the sole exception of *PSBA*.

This correlated with increased ROS within chloroplasts and subsequent inhibition of plastidial

translation [9, 49]. De-repression of plastidial transcription and translation during stress is

mitigated via sigma factors like SIG5 which contributes towards transcription of the psbD operon

(PSII D2) [50]. Translational regulation of stress responsive transcription factors like HSF2A,

DREB2A, and the SAP2/3 interaction with CML49 leads to translation-dependent nucleus

synchronization after 10 min LL→HL. Plastid-targeted proteins like SIG5 and GPT2 and VTC2

were transcriptionally upregulated in CML49- and SAP3-dependent manner but not preferentially

translated after 10 min LL→HL (Figure 6b-f, Suppl. Dataset 1). It is tempting to hypothesize that

the SAP-CML49 co-regulated expression of SIG5 contributes to de-repression of chloroplast

genes, especially since PSII D2 transcription is 1.4-fold increased after 60 min of LL→HL.

Discussion

Regulation of actively translated transcripts described herein adds an entirely new mechanism to

retrograde communication in the cell. We identified components of the novel pathway of

translation-dependent retrograde signaling (TraDeRS) that bifurcates into direct translation of

chloroplast-targeted proteins independently of nuclear transcription (Figure 7 I). In addition,

retrograde control of cytosolic translation affects synthesis of a suite of transcription factors that,

in turn, promote nuclear transcription (Figure 7 II). Both mechanisms culminate to an anterograde

response that enables acclimation. Collectively, the suite of chloroplast-targeted and chloroplast-

encoded proteins controlled by TraDeRS includes those involved in protein import and folding,

chloroplast transcription, photosynthesis and photoprotection.

Evidence for translation-dependent retrograde signalling (TraDeRS)

HL stress perturbs and cumulatively represses global translation, while simultaneously increasing

translation of stress-specific proteins within minutes of LL→HL-transfer (Figure 1a-b, Figure

3a). Our work demonstrates that this process is rapidly and reversibly controlled in a light dose-

dependent manner (Figure 1a-b). Most other immediate short-term responses to HL are

chloroplast-located and post-translational modulated like non-photochemical quenching [51],

malate valve stimulation [52], or activation of antioxidant metabolism [53]. In contrast to the

chloroplast-localized processes, TraDeRS takes place in the cytosol and potentially integrates

multiple cues for changes in *de novo* protein synthesis. Previously, we demonstrated that rapid

transcriptional responses in the nucleus affect a network of transcription factors within 20 s after

the LL→HL-transfer [9, 54]. Only later peaks the H_2O_2 burst at 30 min after the LL→HL. The

decreased polysome association of transcripts displayed a reciprocal behavior with the H_2O_2 burst

after the LL→HL shift. The maximal H_2O_2 accumulation after 30 min is followed by a return to

basal levels after 60 min [9]. Parallel to the decreasing H_2O_2-content, general ribosome

association of mRNA recovered after one hour of HL (Figure 1b and as reported [43]).

Retrograde-dependent translational control involves MPK6 and AKIN10

Ribosome unloading and preferential loading of transcripts are decisive steps in translational

control in the cycle of translation initiation, elongation, and termination [55]. The question is how

does HL perceived in the chloroplast modulate translation? Protein phosphorylation, thiol redox-

related changes and acetylation are reported post-translational modifications (PTM) that regulate

translation initiation, elongation and termination factors [56]. Formation of the translational pre-

initiation complex (PIC) is inhibited by the GCN2 [57], and its activation has been shown in to be

linked to MAPK signaling pathways [58]. Consequently, our results showed that mutants lacking

MPK6 exhibit an altered ribosomal profile. Although the global association of transcripts to

ribosomes were altered in *mpk6* (Figure 2) hinting at altered translation initiation, none of the

marker transcripts from the increased translational cluster were disturbed in their preferential

ribosome association upon LL→HL-transfer (Suppl. Fig. 3a-d). This is indicative of a stress-

dependent separation of global inhibition of translation initiation via MPK6 and the increased

ribosome association of specific transcripts through a distinct mechanism.

The second rate-limiting mechanism of translation initiation is via eIF4F complex formation and

subsequent mRNA binding regulated via 4EBPs. In mammals 4EBPs are activated by the

mammalian Target of rapamycin (mTOR) kinase [59] mediating phosphorylation of the

eukaryotic Initiation factor 4E (eIF4E), disabling interaction with initiation factor eIF4G, and

thus limiting eIF4F formation [60]. Homologous 4eBPs being target of TOR signaling pathways

in plants could not be identified yet. However, phosphorylation of the Translation initiation

factors eIF4E and eIFiso4E via SnRK1, an interactor of TOR, inhibited translation in a plant

system [61]. Herein we show that the lack of AKIN10, a subunit of SnRK1, altered the ribosomal

profile, under control and stress, and interfered with the increased ribosome association of the

marker transcript *SAP2* (Suppl Fig. 3a). A future question is what activates this pathway? ROS or

redox-mediated regulation are candidates, as are GCN2 and TOR-signaling via AKIN10 and

MPK6, and DHAP. We previously reported that TPT2-mediated export of DHAP from the

chloroplast was required for rapid phosphorylation of MPK6 after 10 min LL→HL-transfer. In

summary, the results show that both MPK6 and AKIN10 are required for stress-dependent

translational control in a retrograde signaling context, most likely via a phosphorylation-

dependent pathway. Importantly, the increased ribosome association of specific transcripts

appears to occur independently of this pathway.

Transcripts affected by bifurcating translation-dependent signaling pathways

The set of preferentially loaded transcripts could be grouped into purely translationally regulated

and both, translation- and transcription-regulated transcripts, with each group representing

distinct pathways of the bifurcated acclimation responses. Transcripts related to photosynthesis,

heat stress and H_2O_2 dominated the set of preferentially polysome-loaded and motif-carrying

transcripts (Suppl. Dataset 1, cluster2). Photosynthesis and chloroplast repair-related transcripts

were highly enriched among the purely translationally regulated transcripts (Suppl. Dataset 1,

Cluster1). The 119 transcripts in the translation-only regulatory cluster were 5-fold enriched in

chloroplast-localized proteins according to SUBA4 [39]. This set includes transcripts encoding

proteins critical for assembly, function, and maintenance of the chloroplast. Minor and major

Light harvesting proteins, Cytochrome b_6f (*AT2G26500*) and NADP oxidoreductase

(*AT1G06690*) as well as one small subunit of RuBisCO (AT5G38420). Additionally, PSI

assembly factor PYG7/YCF37 (*AT2G23670*) and import chaperones like HSC70-1 (*AT5G02500*)

and chloroplast vesiculation factors (*AT2G25625*). Our results strongly suggest the existence of

transcription-independent regulation of the photosystem repair mechanism that is under rapid

translation-dependent control, forming a direct retrograde circuit (Figure 7, I. Translational

Circuit).

Complementary to photosystem repair, translational control of nuclear transcription factors

(HSFs, DREBs) synchronizes the gene expression with the cellular demand for subsequent stress

acclimation. The identified HSF2A, DREB2A and MBF1c play major roles in transcriptional

control of HL responses [48, 62]. Interestingly, transcripts for these translation factors are already

present under LL but not preferentially translated (Suppl. DataSet 1). Preferential translation after

stress perception might serve as first activator of these transcription factors that are known to be

up-regulated by oxidative stresses, such as HL.

Regulatory elements in transcript UTRs controlling light-dependent ribosome association

Transcript-specific translation impacts *de novo* synthesis of distinct protein classes and must be

inherently based on regulatory RNA sequence or structural properties. To identify conserved

sequences that might serve as such regulatory property, we conducted *de novo* motif detection in

UTRs of translationally regulated transcripts (cluster 1 and 2) revealing 42 motifs. We could

demonstrate *in vivo* impacts of three identified motifs during HL-stress on translation and provide

evidence suggesting that they serve as RBP binding sites for GAPC1/GAPC2 (Figure 4, Suppl.

Fig. 7a-c). Indeed, RNA-EMSA shifts with leaf extracts from LL and LL→HL leaves indicate

that M1 acts as an mRNP binding site (Figure 4c). Conditional RBP-binding and stimulation of

translation could be shown for motifs M1>>M3>M2 in the arabidopsis protoplast assay (Figure

4a). The motif M2 (TAGGGTTT) has previously been identified in the 5'-UTRs of a set of

preferentially polysome-associated transcripts in young seedlings treated with light for 30 min.

Translation of transcripts encoding reporters carrying M2 *in vitro* occurred at higher rates than

constructs lacking the sequence [43]. The motif was also overrepresented in preferentially

polysome-associated transcripts under hypoxic stress [63], suggesting a role for motif M2 under

various stresses. However, prior to this study its mode of action had not been reported.

The motif-binding RBPs were investigated by RNA-based affinity chromatography and mass-

spectrometry. This identified ATP synthase subunit alpha and GAPC2 as candidate RBPs for M1

and M2, respectively. The other identified peptides were either not matching in predicted protein

size, or had a too low Mascot-Score and were excluded as background or unspecific binding.

GAPDH functions as non-canonical RBP in animal systems [64, 65] and this role was only

recently discussed in plants as well [66]. We validated the GAPDH / motif interaction using the

native M1 in the *SAP3* 5'-UTR via *ex-vivo* RNA-affinity chromatography, and subsequent

immunodetection, and *in vitro* by RNA-EMSA. The interaction was strong with LL protein

extracts and almost lost in extracts from LL→HL shifted leaves, pointing towards a light-

dependent dissociation of GAPC2 (Figure 4c). Therefore, this dissociation upon LL→HL shift

appears to release repression of ribosome association, for transcripts harboring the motifs. As

protein amounts of GAPDH (C1/C2) were unaltered by LL→HL treated plants, conditional RNA
binding could be mediated by PTM or the availability of co-factors or substrates. This may link
translational control by RBPs to the cellular metabolic state, thereby enabling the cell to rapidly
adjust protein synthesis according to metabolic requirements.

SAP2 and _SAP3_ contain motifs 1 and 2 and their proteins interact with CML49

Motifs M1 and M2 are both present in the 5'-UTRs of _SAP2_ and _SAP3_ from arabidopsis and also
in their homologues in the monocot rice. The transcripts in rice showed the same polysomal
association during L→H as in arabidopsis. Interestingly, re-analyzing a dataset of arabidopsis
seedlings re-illuminated after darkness for 10 min [26] for translationally regulated transcripts in
the same manner as performed for this study, revealed _SAP2_ as one of only five transcripts
overlapping with the purely translationally regulated subgroup (cluster1, Suppl. Dataset 1).

The SAPs belong to a zinc-finger multigene family with a highly conserved structure, leading to
similarity among several plant species [45, 46]. Despite little molecular knowledge, two
regulatory roles have been attributed to SAPs, namely ubiquitin ligase activity and redox sensor
activity regulating gene expression under abiotic stress [46, 67-69]. Previous work demonstrated
that SAP12 undergoes reversible redox-dependent conformational changes induced during
oxidative stress [67]. Given the high level of sequence conservation in the SAP-protein family we
analyzed SAP2 and SAP3 for disulfide bonds with the DiANNA [70] tool suit. The analysis
revealed a predicted disulfide bond in the AN1 domains of SAP2 and SAP3 at Cys117 and
Cys135, respectively. SAP2 had one additional disulfide bond predicted between Cys8 and
Cys22. Thus, SAP2 and SAP3 might also function as redox sensors under HL. Furthermore,
SAP5 interacts with MBF1c to regulate acclimatory responses, during heat stress via interaction
with HSF and DREB transcription factors, ultimately influencing transcription [71]. Given the
structural similarity between SAP2, SAP3 and SAP5, the interaction of SAPs with MBF1c might
co-regulate transcript specificity while interaction between HSFs, DREBs and MBF1c
synchronize transcriptional demand, but this remains subject to further study. Nonetheless, as a
consequence of retrograde signaling the abundance of SAP2 and SAP3 is increased and, in
parallel, promotes their re-location to the nucleus to promote transcription.

SAP2 and SAP3 interact with the Calmodulin-like-protein-49 (CML49). Several CMLs display
an electrophoretic mobility shift in the presence of Ca^{2+}, indicating that they act as Ca^{2+} sensors

[72, 73]. Most CMLs lack known functional domains that impede the prediction of biochemical

functions. Nonetheless, CMLs are important in signaling processes and are activated under stress

conditions like heat, cold, drought, and heavy metal accumulation [74, 75]. The Ca^{2+}-dependent

binding of CaM7 to a Z-/G-box is involved in regulating expression of light-responsive genes

[76]. H_2O_2 triggers an increase in cytosolic Ca^{2+} by activating Ca^{2+} channels [77]. Ca^{2+}-liganded

CaM activates MPK8 which suppresses ROS accumulation via transcriptional control of *RBOHD*

[78]. *CML49* occurs in almost all tissues and was not transcriptionally or translationally induced

after 1 h of LL→HL indicating that *CML49* functions as constitutive signal transmitter.

Interestingly, the rice homologues of *SAP2* and *SAP3* (*OsSAP8* and *OsSAP9*) also interact with

CML49 homologues indicating a conserved regulatory mechanism potentially integrating ROS

cues with Ca^{2+} signaling (Suppl. Fig. 8). But this discussed function needs further experimental

confirmation.

SAP3 and CML49 interaction co-regulates light-induced transcription

Both *sap3* and *cml49* knockouts display smaller sets of light-induced transcripts compared to

WT. We propose that CML49 acts as general interactor with both SAPs, while each SAP acts as a

stress-specific partner. First, CML49 is constitutively expressed while *SAP2* and *SAP3* are

preferentially translated under HL. Newly translated SAPs can then interact with available

CML49 resulting in relocation of the complex to the nucleus. Second, of the 215 transcripts

downregulated in *cml49* plants after 60 min of LL→HL transfer a stress subset of 32 are

downregulated in *sap3*. Thus, the stress-responsive protein, SAP3, is providing specificity to

what would be otherwise a broader CML49 response (Figure 6a). Cumulatively, the preferential

translation of the SAPs and their interaction with CML49, but also other TFs like DREB2A and

HSF2A, form a translation-reliant signal pathway affecting nuclear transcription.

This translation-reliant nuclear circuit regulates two major clusters of genes. As noted above, the

majority of the transcripts regulated by the proposed CML49-SAP3 module code for stress

responsive transcripts. A subset which includes cytosolic *HSP70*, *HOP3*, *ROF2*, *HSFA2*,

HSP90.1, *HSP20-like* that increase after 1 h of LL→HL shift (Figure 6 a). Interestingly, *HSP90.1*

and *ROF1* interaction is needed for HSFA2-dependent heat memory and also regulates *short*

HSPs (*sHSPs*) [79]. HSFs have also been shown to translocate to the nucleus in a redox-regulated

manner [80]. Taken together, we propose that this mechanism forms a feed-forward loop that can

amplify the transcriptional responses over time, the translation-reliant retrograde stress signaling
pathway (Figure 7, II. Translation-Reliant Nuclear Circuit).

A second subset of nuclear-encoded transcripts co-regulated by CML49 and SAP3 takes part in
an anterograde signaling pathway that collectively could contribute to chloroplast acclimation.
Among the transcripts were *VTC2*, encoding GDP-L-galactose phosphorylase, which catalyzes
the rate limiting enzymatic step in ascorbate synthesis [81]. With respect to anterograde
signaling, plastidic *Sigma factor 5* (*SIG5*) is utilized by chloroplast-encoded RNA-polymerase to
tran

scribe the chloroplast-encoded *psbD* gene, encoding the D2 protein of the reaction center of
photosystem II [6], that is induced by HL [82]. *SIG5* and *D2* transcripts were increased in
response to the LL→HL, and *SIG5* was reduced in the *cml49* and *sap3* knockouts, providing
evidence for an anterograde circuit potentially affecting chloroplast transcription at later time-
points in response to translational-dependent retrograde signaling.

Conclusion

This work reveals a translation-dependent retrograde signaling pathway and identifies major
participants across key stages that regulate rapid acclimatory responses to LL→HL transfer. The
translatome reorganization initiated by LL→HL involves phosphorylation cascades involving
MPK6 and AKIN10, which inhibit cumulatively global translation while simultaneously
preferential ribosome association occurs for specific functionally distinct transcripts by a
different mechanism. This may serve as a very efficient local control mechanism (i.e.
circumventing the nucleus) and is possibly faster than transcriptional regulation for synthesizing
proteins required for chloroplast acclimation. The preferentially loaded transcripts directly code
for repair and photosynthesis-related proteins forming a direct translational retrograde circuit
(Figure 7). Preferential ribosome association is facilitated by conserved motifs in their 5'-UTR
acting as RBP binding sites. The motifs, together with GAPC2, function in the direct circuit in a
light-dependent manner. A translation-reliant nuclear circuit regulates, next to well-studied stress
related transcription factors, the translation of SAP2 and SAP3 in response to LL→HL together
with their interaction with the Ca^{2+} sensor CML49 to promote accumulation of stress-responsive
transcripts and anterograde signaling via SIG5, which functions in plastid transcription. The
elucidated multiple components of a bifurcating translation-dependent retrograde signaling

pathway allows for rapid tuning of protein synthesis to account for the demand of the
photosynthesizing cell.

Materials and Methods

Plant growth and treatments. *A. thaliana* wild-type (ecotype Columbia) and mutants (*cml49-1*
(SALK_090873), *cml49-2* (SALK_035905), *sap3-2 (SALK_022340), sap3-3 (SALK_136520C),*
sap2-1 (SALK_008677), *sap2-2* (SAIL_1182_C11), *mpk6-3* [83], and *akin 10* (GABI_579E09)
were grown under short day conditions (10 h light with 80-100 µmol photons m^{-2} s^{-1} (NL), and
21°C/ 14 h dark, and 18°C, 50 % relative humidity) for three weeks prior to transfer to 8 µmol
photons m^{-2} s^{-1} (LL) for 10 days. Plants were transferred to 800 µmol photons m^{-2} s^{-1} (HL) for
time periods as indicated, while control plants remained in LL. Whole rosettes were harvested
under control and HL conditions in the midday point of the light period to exclude circadian
effects and immediately frozen in liquid nitrogen and stored at -80°C.

Polysome profiling and sucrose cushion for polysome isolation. Total ribosomes and
ribosome-bound mRNA were extracted for polysome profiling and quality check via TEM after
separation by sucrose gradient centrifugation as described [31]. Polysomal profiles were blank
gradient normalized and the peaks corresponding to 60S, 80S and polysomes integrated before
normalization to WT values. Significant differences were calculated via two sided Students's t-
test p≤0.05. Sucrose cushion centrifugation was performed prior to foot print analysis. 3 g of
pulverized plant material was used with 6 mL polysomal extraction buffer (PEB; 0.2 M Tris, pH
9.0, 0.2 M KCl, 25 mM EGTA, 35 mM MgCl, 1 % Brij, 1 % Triton X-100, 1 % Tween 20, 1 %
Igepal CA630, 1% sodium deoxycholate (DOC), 1 % polyethylene-10-tridecylether (PTE), 5 mM
DTT, 1 mM PMSF, 100 µg/mL cycloheximide, 100 µg/mL chloramphenicol, 100 µg/mL
lincomycin). The suspension was passed twice through Miracloth^{TM} and subsequently cleared by
centrifugation. Extracts were loaded on a sucrose cushion (0.2 M Tris, pH 9.0, 0.2 M KCl,
0.025 M EGTA, 0.035 M MgCl, 1.75 M sucrose, 5 mM DTT, 50 µg/mL cycloheximide,
50 µg/mL chloramphenicol, 50 µg/mL lincomycin) and centrifuged for 18 h at 100,000 xg in a
Beckmann SW71Ti rotor. The sediment was suspended in 100 µL RNase digestion buffer
(20 mM Tris-HCl, pH 8.0, 140 mM KCl, 35 mM $MgCl_2$, 50 µg/µL cycloheximide, 50 µg/µL
chloramphenicol, 50 µg/µL lincomycin) and used immediately for RNase If digestion.

Transmission Electron Microscopy (TEM) Imaging. TEM images were generated at the Centre for Advanced Microscopy at ANU using the JEM F200 (JEOL) by using 2 µL of undiluted sucrose gradient from fractions representing monosomal or polysomal fractions and negative staining as described in Booth et. al. [84].

RNase If digestion. RNase If digest was performed with 2,000 relative absorbance units OD_{260} of the in RNase digestion buffer suspended polysomes by adding 50 U of RNase If (New England Biolabs) in 250 µL total volume by incubation at room temperature with constant rotation for 1 h. In parallel same amounts of polysomes were treated with 5 µL SUPERase RNase inhibitor (Thermo) and kept on ice as control sample. The digestion was stopped by adding 5 µL SUPERase RNase inhibitor and the RNA precipitated by adding 1 volume of isopropanol, 300 mM NaOAc pH 5.2 and 20 µg glycogen at -80 over-night. The supernatant was discarded after centrifugation at 16,000 xg at 4 °C for 30 min and the sediment was washed twice with 75 % ethanol and resuspended in RNA loading buffer for size exclusion electrophoresis.

Library sequencing and analysis of mRNAs. Total RNA was isolated using Tri Reagent (Sigma-Aldrich) from extracts suspended in polysomal extraction buffer (see sucrose cushion for polysome isolation, above). Briefly, 500 µL of extract was combined with 1 mL of TriReagent, followed twice by extraction of the organic phase with 200 µL chloroform. RNA was precipitated with equal volume 100% isopropanol and incubated overnight at -20°C. The RNA was recovered by centrifugation and washed with 75% ethanol before resuspension in water. RNA quality was assessed using a LabChip GXII (Perkin-Elmer). Total RNA was DNase-treated using TURBO DNase (Invitrogen) following the manufacturer's protocol. Preparation of total RNAseq libraries was carried out using Illumina TruSeq Stranded Total RNA with Ribo-Zero plant, scaled at half reaction volumes.

Libraries of ribosome-protected fragments (RPFs) were prepared from three biological replicates per time-point and sample following RNase If digestion (above) according to Juntawong et al. (2014) with minor modifications. RNA linker, rRNA subtraction, and PCR oligonucleotides were ordered from Integrated DNA Technologies (IDT) and are listed in Suppl. Table 2. Oligonucleotides for rRNA subtraction were adapted from those listed in [36] and [85]. Additionally, rRNA1c, rRNA1 and rRNA2 depletion primers were custom designed targeting the highest abundant rRNA detected during method establishment sequencing.

Total RNA and RPF sequencing libraries were quantified using the Qubit fluorometer (Invitrogen) and size distribution was assessed using the LabChip GXII. Samples were pooled and sequenced (75 bp single end) on the NextSeq500 at the Biomolecular Resource Facility (BRF) at the Australian National University.

Quality control of sequencing data was performed with FastQC (v.0.11.5). Total RNA data had adapters removed using scythe (v.0.991) with a prior of 0.01, and quality trimmed with sickle (v.1.33) with a quality threshold of 20 and a minimum trimmed length of 20. Mapping to TAIR10 was performed with subjunc (v.1.4.5), reporting only reads with a single unambiguous mapping location [86]. RPF data was trimmed for adapter contamination and read quality using Cutadapt (v.1.9.1) with a minimum trimmed length of 20 bp and error rate of 0.1 [87]. Trimmed reads were mapped to TAIR10 using Bowtie2 (v2.2.9) requiring end-to-end alignment and a substring length of 10 [88]. Total and RPF-mapped reads were summarised using featureCounts [89]. Counts data was converted to reads per kilobase million (RPKM), and counts per million reads mapped (CPM) in R using edgeR [90, 91]. Only transcripts with counts of RPKM above 7 for all three replicates were considered for subsequent analysis.

DEG definition and transcript clustering. Cut-offs for assignment as Differentially Expressed Genes (DEGs) were chosen in a data-driven process, according to upper and lower quantiles from a random distribution analysis of all regulated polysome-associated and total transcripts (total mRNA: 1.5 and 0.75 and polysomal: 1.1 and 0.75; 0.05 error margin). Clusters were derived for transcripts significantly changed up and down in both polysomal and total RNA (cluster 1 and 4). Cluster 2 and 3 were selected by filtering for only significantly up- and down-regulated in the polysomal set and not significantly changed in total RNA. Other transcripts were not assigned to clusters, such as 33 DEGs that were changed in total mRNA but not in the ribosomal fraction and termed as buffered transcripts.

Whole genome ATH1 microarray. Total RNA was isolated from 1 h LL→HL-treated *sap3*, *cml49* and WT plants using RNeasy mini Kit (Qiagen). Controls were kept in LL. The hybridization of the Affymetrix ATH1 genome Array was commissioned to KFB (Regensburg, Germany). Each condition was hybridized three times from independent experiments for *sap3* and WT and once for *cml49*. Microarray data were analysed with ROBIN (MPI Golm, Potsdam, Germany) and normalized with RMA algorithm [92]. Transcripts were considered as differentially regulated if the FC was >2 and the adjusted *p*-value <0.05 [93, 94].

Transcript profiling. RNA isolation and cDNA synthesis were performed as in [95].
Quantitative real-time PCR (qPCR) of total and polysomal transcripts was conducted as
described in [96] with the difference of using KAPA SYBR® FAST qPCR Kit (Kapa Biosystems,
South Africa). *ACTIN2* (*AT3G18780*) and *TUBULIN5* (*AT1G20010*) were used as reference
genes.

RNA electrophoretic mobility-shift assay (RNA-EMSA). 5'-biotinylated RNA and unmodified
RNA of Motif 3 (5'-AAAUCGCCGGAG-3' 5'-Bio-AAAUCGCCGGAG-3') were synthesized
by Eurofins (Hamburg, Germany) and used for LightShift™ Chemiluminescent RNA EMSA Kit
(Thermo Scientific). Total protein was extracted from LL and 10 min LL→HL-treated plants
using extraction buffer (25 mM Tris, pH 7.8, 75 mM NaCl, 15 mM EGTA, 15 mM
glycerophosphate, 15 mM 4-nitrophenyl pyrophosphate, 10 mM $MgCl_2$, 1 mM DTT, 1 mM NaF,
0.5 mM NA_3VO_4, 0.5 mM PMSF, 10 µg/mL leupeptin, 10 µg/mL aprotinin, 0.1 % Tween-20).
Prior to separation by non-denaturing polyacrylamide gel electrophoresis, RNAs were incubated
with extracts at room temperature for 30 min. Biotinylated RNA was detected by
chemiluminescence with streptavidin-HRP. For targeted EMSA AtGAPC1 and ArGAPC2 were
expressed in *E.coli* BL21 (DE3) pLysS and purified as described in [97] with slight
modifications. Buffers were not supplemented with NAD^+ and purified protein was stored in 10
mM NaPi, pH 7.4. AtGAPC1, AtGAPC2 and GAPDH from rabbit muscle (SIGMA Aldrich)
were used for mobility shift assays with 5'-biotinylated M1-RNA (5'-BioTEG-
AUAUAUUCUCCGGAGAA-3'),   5'   biotinylated   M2-RNA   (5'-BioTEG-
AUAUAUGAGAUUAGGGUUU-3'),   5'-biotinylated   SAP3   M1   (5'-Bio-
ACCAGAUUUUUUCGCCGGAGUUUGUUUUGAUCCA-3'),   and   random   5'UTR
(*AT5G16470*)   with   mutated   M2-sequence   (5'-Bio-
UUAGCGUUUUAAU*GUUGAGUGAGUA*AGUUAAGGGGUU-3').

Reporter gene construct and transient protoplast assay. For the reporter gene construct assay
solely *UBI* 10 5'-UTR as control (CCATGGCTTGATCACGGTAGAGAGAATTG–
AGAGAAAGTTTTTAAGATTTTGAGAAATTGAAATTCTCG) or *UBI* 10 5'-UTR with
motifs (M1:   CCATGGCTTGATCACGGTAGAGAGAATTGAGAGAAAGTTTTTTTC–
TCCGGAGATTCTCCGGAGATTCTCCGGAGAAAGATTTTGAGAAATTGAAATTCTCGA
G;   M2:   CCATGGCTTGATCACGGTAGAGAGAATTGAGAGAAAGTTTTTAAACCC–
TAATCTCAAACCCTAATCTCAAACCCTAATCTCAAGATTTTGAGAAATTGAAATTCTC

GAG; M3: CCATGGCTTGATCACGGTAGAGAGAATTGAGAGAAAGTTTTTCTCCGG–

CGACTCCGGCGACTCCGGCGAAAGATTTTGAGAAATTGAAATTCTCGAGA) were

cloned into MS129 luciferase reporter [98], and 5 µg plasmid DNA was co-transfected with same

amount of low-rate constitutively expressed pRL-TK *Renilla* luciferase vector (Promega) into

mesophyll protoplasts isolated from *A. thaliana* leaves by the downsized polyethylene glycol-

mediated transfection method (Seidel et al., 2004). After 12 h of acclimation protoplast were

transferred to LL and half of the transfected protoplasts were transferred to HL for 10 min and

subsequently flash frozen in $N_{2\ liq}$ for cell extracts. Luciferase activity of cell extracts was

measured with Sirius L Tube Luminometer (Berthold) and the luciferase assay system (Promega).

RBP pulldown and identification. Affinity pulldown with motif-carrying biotinylated RNA for

detection and identification of potential protein-RNA interactors was conducted as described in

[99] with minor differences. In short: 500 µg (or 250 µg for pull down followed by

immunodetection of cytosolic GAPDH) of total protein from LL and LL→HL-treated plants were

incubated for 30 min at RT with 1 µg biotinylated M1- or M2-RNA (3 µg for 5'-Bio-*SAP3*-M1),

1 µL aprotinin (25 µg/µL), 1 µL leupeptin (25 µg/µL), 2 µL RNasin (Promega), 250 µL TENT

buffer (0.02 M Tris-HCl, pH 8.0), 0.002 M EDTA (pH 8.0), 0.5 M NaCl, 1 % (v/v) Triton X-100)

and adjusted to 500 µL with TENT buffer. 50 µL Dynabeads (Thermo Fisher) were added and

incubated under slow shaking for 30 min at RT. After precipitation and three washing steps with

TENT buffer, Dynabeads were dissolved in 1x protein loading buffer and incubated for 10 min at

95 °C, precipitated and the supernatant loaded on SDS-PAGE for separation and band

identification via mass-spectrometry or immunodetection of GAPC1/C2 after blotting the gel-

separated proteins on nitrocellulose membrane by semi dry electro blotting technique.

Promotor and uORF analysis. The +50 to -1000-bp region upstream of the ATG start codon

and +500 downstream of the transcription stop site of the CML49 and SAP3 co-regulated genes

were analysed for common *cis*-elements using PlantPAN2.0 [100]. For uORF analysis the web

tool uPEPperoni was used [101] and the eFP Browser 2.0 for general expression pattern [102].

mbSUS-analysis. Vector construction of the mating-based Split-Ubiquitin-System analysis was

performed with the Gateway® cloning system (InvitrogenTM). The coding sequences of SAP2,

SAP3 and CML49 were fused to attB recombination sites and cloned into the pDONRTM221-

vector by a BP-reaction. Subsequently, each entry clone was used in a LR-reaction with the

destination vectors pX-NubWTgate (Nub) and pMetYCgate (Cub). Controls were the

transformed diploid yeast with NubI- (pNubWtX-gate, positive control) and the NubG-vector

(pNX35-Dest, negative control) as prey. The resulting vectors were transformed via lithium

acetate transformation into *Saccharomyces cerevisiae* strains THY.AP5 and THY.AP4,

respectively [103]. Yeast cells were grown in 20 mL selective Nub/Cub medium at 30°C/140 rpm

for 5-7 days. To start the mating step, THY.AP5 and THY.AP4 were transferred to YPD plates

and grown for 1 d at 30°C. For the methionine assays, mated yeast was cultivated in selective

liquid diploid medium for 3 d (30°C, 140 rpm). OD600 of selected yeast cells were measured and

diluted to OD600 of 0.1 and 1.0 respectively in sterile water. Then 6 µL drops of each dilution

were transferred to agar plates containing no (Met 0 µM) or 0.075 % (v/v) methionine (Met

50 µM). Subsequently, the yeast was grown for 72 h at 30°C and results documented

photographically [103, 104].

Subcellular localization and FRET measurements. The coding sequences of the genes of

interest were fused in frame to yellow or cyan fluorescent protein as indicated and fluorescence

resonance energy transfer (FRET)-experiments were performed as described by [105] with

CLSM 780 (Zeiss). More than 20 protoplasts were measured twice for each set of PEG-based

transformation and the experiment was repeated independently

Acknowledgements

The authors acknowledge support by the Deutsche Forschungsgemeinschaft (Di 346/19; SP 1935) and for funding by the Australian Government through the Australian Research Council (DP220103640). Barry Pogson is the recipient of an Australian Research Council Australian Laureate Fellowship (FL190100056) funded by the Australian Government. Special thanks to Renate Scheibe for providing the pET16b construct for recombinant expression of GAPC1/GAPC2. We acknowledge the ACRF Biomolecular Resource Facility, a service node of Bioplatforms Australia, and the Centre for Biodiversity Analysis Ecogenomics and Bioinformatics Lab for provision of resources and expertise to perform Illumina sequencing. We acknowledge the facilities and the scientific and technical assistance of Microscopy Australia at the Centre for Advanced Microscopy, Australian National University, a facility that is funded by the University and the Federal Government.

The authors declare that they have no competing interests.

Data availability

All relevant data is accessible in this manuscript or in the supplemental material. A preprint version is accessible on bioRxiv under: doi: https://doi.org/10.1101/2021.02.18.431817.

Authors contributions

MM designed the research and performed the polysomal gradients, TEM microscopy, transcript profiling, microarrays, RPF-seq and total RNAseq and bioinformatic data analysis, transient protoplast reporter gene assays, RNA-EMSA, protein affinity pull downs and wrote the paper. AS did the RPF-library preparation and sequencing and performed bioinformatic analyses. CW prepared polysomal gradients and transcript profiling. SS performed Y2H experiments and cellular localization and FRET. MW performed RBP affinity pull downs and gel shift assays on SAP3 motifs. AF measured FRET and transcript profiles of OsSAP and OsCML49. DRG discussed and analyzed transcriptomics data. TS supervised and performed FRET measurements. BP supervised and discussed the results and wrote the paper. KJD designed the study, supervised and discussed the results, coordinated the group and wrote the paper.

Figure legends

Figure 1: Time- and dose-dependent changes in ribosome associated transcripts during high light treatment. (a) Growth light (NL) or low light (LL)-acclimated plants were transferred to high light (HL) for a 10-fold light shift (NL→HL) or a 100-fold light shift (LL→HL) for indicated time. Depicted are representative baseline-normalised polysomal profiles in sucrose gradients of n=3 biological replicates. (b) Association of RNA with polysomes is dynamically and reversibly regulated during a 100-fold light shift. Plants were kept in LL (0') or transferred to HL (LL→HL) for indicated time periods and then transferred back to LL for the indicated time for recovery. Ribosome association was determined by comparing ribosome-associated mRNA quantity from polysomal fractions of sucrose gradients to total mRNA on a fresh weight (FW) basis (mean ± SE from n=3 independent experiments; letters indicate significance of difference as determined with ANOVA post-hoc Tukey p<0.5). (c) Analysis of polysome status from LL and 10 min LL→HL treated plants via polysomal profiling and TEM imaging of ribosomes (arrows) of the corresponding fractions revealed presence of monosomes (80S), low molecular polysomes (2x) and high molecular polysomes (3-5x) (green bar represents 100 nm).

Figure 2: Mutants involved in retrograde signalling exhibit genotype-dependent LL→HL translational response. (a). Differential ribosome association of transcripts in retrograde mutants as detected in polysomal profiles of LL grown and 10 min LL→HL treated plants (b). 10 min LL→HL shift repressed ribosome association of transcripts. Depicted are representative polysomal profiles of WT, *mpk6* and *akin10* KO background after sucrose gradient centrifugation of n=3 biological replicates (60S, 80S and polysomal fractions indicated). (c) Quantification of genotype dependent LL→HL response shows repression of ribosome association in WT, a stronger decrease in *mpk6* and a loss of regulation on akin*10* background. Polysome profiles were background normalized and the peaks corresponding to 60S, 80S and polysomal area integrated and normalized to WT control (* represents significant differences to WT, two sided Student's t-test p<0.05).

Figure 3: Translatome after 10 min LL→HL reveals distinct regulatory pattern. Significantly regulated transcripts are depicted as dots with the size indicating transcript abundance. There are 4 distinct clusters of transcripts: translationally upregulated (Cluster 1), transcriptionally and translationally upregulated (Cluster 2), transcriptionally and translationally downregulated (Cluster 3), and translationally downregulated (Cluster 4). Each cluster is significantly enriched in transcripts coding for different GO annotations, exemplar genes are listed. Interestingly, Cluster 2 codes only for stress genes (most prominently HSF2A, DREB2A) while Cluster 1 encompasses almost exclusively photosynthetic genes and translation processes (e.g. LHCBG1.1 and R40S_S16-1)**.** (b) The fold change response of each cluster depends on transcript abundance. Comparison of

transcript ribosomal mRNA fold change with total measured transcripts revealed an inverse correlation for Cluster 1 and 4, although they are not transcriptionally changed. Surprisingly, this inverse correlation was not clearly seen for the transcriptionally and translationally changed transcripts in Cluster 2 and 3, indicating a less pronounced impact of transcript abundance (i.e. availability) on ribosome association.

Figure 4: Identification and functional validation of three motifs in the 5'UTR of preferentially ribosome-associated transcripts. These motifs are hypothesized to confer preferential translation after 10 min LL→HL. (a) In 338 5'-UTRs of the preferentially ribosome-associated transcripts 42 enriched motifs were identified using the MEME algorithm against a randomized 5'-UTR set. Motifs 1 and 3 were present in 32% of those transcripts, and motif 2 in 13% with an additional 8% occurrence in their respective 3' UTR. The 5'-UTR of *UBI10* was cloned upstream of the luciferase (LUC) reporter under control of the ubiquitin promotor. Triplet motifs were introduced into the 5'-UTR of ubiquitin mRNA. *A. thaliana* mesophyll protoplasts were co-transfected with one of each construct and with the pGL4::RLUC reference construct. Depicted are schematics of constructs used in the reporter assays and relative LUC signal in lysed mesophyll protoplasts treated with HL for 10' normalized to reference construct. The construct containing M1 showed a strong increase in LUC signal after LL→HL. Data are means ± SD from two independent experiments with 10 replicates each, * and + indicate significant difference to ubiquitin control, two sided Student's t-test: p<0.05 and 0.1 , respectively. (b) In vitro GAPC1-binding activity to 5'-UTR motifs and random sequences. M1 and M2 were incubated with indicated µg of heterologously expressed AtGAPC1 and subsequently analyzed for RNA electrophoretic mobility-shifts. AtGAPC1 and RNA interaction can be detected by separating at higher apparent size (bound). RNA with M1 and M2 were bound by AtGAPC1 in a dose-dependent manner. This could not be observed for the random RNA oligomer (Random RNA). (c) *Ex vivo* light-dependent interaction of GAPC1/2 with *SAP3* M1. Total protein extract from LL and LL→HL-treated plants was incubated with biotinylated M1 motif RNA and bound to streptavidin magnetic beads. The input GAPC1/2 protein was detected via SDS PAGE and Western Blot using a GAPC1/2 antibody as well as Ponceau staining of RuBisCO. The magnetic beads were washed from unbound protein and RNA interaction (Wash steps). The bound protein was eluted and a strong GAPC1/2 signal only appeared in the LL protein sample.

Figure 5: Interaction of SAP2 or SAP3 with CML49 in the split-ubiquitin yeast-two-hybrid system and by Förster resonance energy transfer (FRET) of SAP2:CFP or SAP3:CFP with CML49:YFP in *A. thaliana* protoplasts. (a) Quantification of interaction strength between SAP2 or SAP3 and CML49 using a methionine-dependent split-ubiquitin-assay. Depicted are colonies after 72 h at 30 °C in different ODs (OD = 0.1 and 1.0). Growth was monitored either on Met 0 µM

(no methionine, left colonies) or Met 50 µM (50 µM methionine, right colonies) medium plates. Diploid yeast was transformed with NubI- (pNubWtX-gate) as positive control and as negative control with the NubG-vector (pNX35-Dest) as prey (depicted are representative images of n=3 independent experiments). (b) *A. thaliana* mesophyll protoplasts were transfected with SAP2:CFP, SAP3:CFP or CML49:YFP vector DNA. Fluorescence images taken by CLSM for subcellular localization. Following transfection with SAP2:CFP or SAP3:CFP and CML49:YFP constructs, FRET efficiencies between CFP and YFP were quantified. FRET was also determined with CFP:YFP as positive control and 35S::CFP with 35S::YFP as negative control. For each set of transformation more than 20 protoplast were measured twice in n=3 independent transfection experiments.

Figure 6. Deregulated HL-response in *sap3* and *cml49* compared to WT. (a) Microarrays were hybridized with cDNA from WT, *cml49* and *sap3* after 60 min of LL→HL transfer. Assuming a concerted function of SAP2 and SAP3 together with CML49 in the HL response, the data were filtered for co-regulated targets that were stronger deregulated in *cml49* than *sap3* background. The expression level corresponds to the colour intensity with yellow = -2 and blue = 4 ($\log_2$ fold). (b-f) Validation of transcript responses to HL of putative CML49 and SAP3 targets. VTC2, HSP20-like, PS3, SIG5 and eEF1B were selected based on the transcriptome response, and their transcript levels quantified by qRT-PCR following the LL→HL-transfer. Data are means ± SD of n=4 independent experiments, asterisks indicate significant difference, two sided Student's t-test p<0.05.

Figure 7: Schematics of translation-dependent retrograde signalling (TraDeRS). Rapid regulation of polysome loading and unloading after 10 min of LL→HL shift was partially disturbed in *mpk6* and *akin10* mutants indicating retrograde regulation of translation. Regulation involves transcription-independent changes of translational efficiency of majorly chloroplast-targeted proteins involved in photosynthesis as well as chloroplast-associated assembly, repair and import of proteins, thus forming a direct feedback circuit regulated via translation (I. Direct Translational Circuit). The preferential association of transcripts from transcription factors as well as *SAP2* and *SAP3* are part of the translation-reliant synchronization of the nucleus with the changed chloroplast demand (II. Translation-Reliant Nuclear Circuit). SAP2 and SAP3 interact with CML49 and this regulatory module affects transcript abundance of, e.g. *VTC2* and *SIG5*, participating in anterograde signalling, and other major stress response genes like *ANAC2*, *HSP101* and *HOP3*.

Supplementary materials

Suppl. Fig. 1: Size distribution of RPF-Seq and frame analysis of start/end sites. (a) The size distribution of RPF-sequencing reads from LL and LL→HL-treated plants exhibited two distinct optima at 23-25 and 28-32 nucleotides matching the reported sizes for stalled plastidial/mitochondrial versus cytosolic footprints, respectively. (b,c) Periodicity of translation frames agrees with reported literature using combined application of cycloheximide and lincomycin in plant systems. Depicted are the mean normalized counts in RPKM for each coding frame (0, 1 or 2) at the relative nucleotide distance to the 3'- and 5'- end and start site, respectively. No difference in coverage could be detected between LL and LL→HL conditions.

Suppl. Fig. 2: Translationally regulated transcripts after 10 min LL→HL. (a) The volcano-plot of ribosome associated transcripts revealed 951 significantly (adj. p-value <0.05) changed transcripts after 10 min LL→HL transfer. 76 % (793) of the significantly changed ribosome-associated transcripts were down-regulated with 15% (158) being upregulated and 8% (92) unchanged in their fold change response. (b) Transcriptional changes were less prevalent with only 204 total transcripts being significantly changed with 38% (83) down-regulated and 55% (121) upregulated total mRNA FC response. This corresponds to a 4.6-fold higher translational than transcriptional control of gene expression after 10 min LL→HL transfer. (c,d) Despite being primarily categorized by ribosome association and not by altered transcription, cluster 2 is not distinguished when assessed using the by common translation efficiency (TE) calculations (TE = FC ribosome-associated mRNA / FC total mRNA). In fact, following TE calculation for this dataset (TE; as Ribo10/Total10 /Ribo 0/Total 0) confounds transcriptional and translational regulation and might incorporate different biological processes. For all TE-upregulated transcripts were the ribosome association ($\log_2$ Ribosomal mRNA FC (LL→HL/LL)) and transcriptional changes ($\log_2$ Total mRNA FC (LL→HL/LL)) plotted. The majority of TE upregulated transcripts are transcriptionally downregulated (95%; 456) with only 85% (404) of TE-upregulated transcripts actually having a positive fold change in ribosome association after 10 min LL→HL transfer. More importantly, 15 % (73) of the TE-upregulated transcripts are less associated with ribosomes as is indicated by the downregulation of fold change in ribosome association, thus the group of TE upregulated transcripts consists out of positive association and negative association with ribosomes during stress.

Suppl. Fig. 3: Cluster 1&2 representatives avoid global translational down-regulation in *akin10* and *mpk6* background. (a-d) Four transcripts were selected as representatives from Cluster 1&2 and subjected to qPCR analysis in wildtype (WT), *mpk6* and *akin10* mutants. Data are means ± SE of n=5 independent experiments, * indicate significance difference relative to LL control, greek and

capital letters indicate significant differences between genotypes for LL and LL→HL samples, respectively. Two factor ANOVA with Tukey post hoc test: p <0.05, respectively. A significant increase of total transcript in LL was observed in *mpk6* and *akin10* in comparison to WT for *SAP2* as representative for Cluster 1 (a) and *SAP3* and *HSP70* as representatives for Cluster 2 (b,c), but not for *DREB2A* (d). Total transcript abundance of *SAP2* and *HSP70* was significantly higher in *mpk6* and *akin10* after LL→HL-transfer than in WT, but not for *SAP3* and *DREB2A*. In *mpk6* under LL conditions, ribosome-associated transcript abundances were not significantly different for *DREB2A* and *SAP2* but significantly higher in *mpk6* than in WT for *SAP3* and *HSP70*. Ribosome-associated transcripts after LL→HL transfer were significantly lower in *mpk6* than WT for *SAP2* and significantly higher for HSP70, with no significant difference for *DREB2A* and *SAP3*. In *akin10*, no significant difference to WT after LL→HL for *SAP2* and *SAP3*, but a significant increased relative abundance for HSP70 and *DREB2A* ribosome-associated transcripts could be detected. The 10 min LL→HL response was similar for all four ribosome-associated transcripts in mutants and WT with slightly changed amplitude of relative increase for *HSP70* and *SAP2*. (e,f) Differences between transcriptional and translational effects of mutants on LL and LL→HL treated plants prompted us to conduct a principal component analysis using the qPCR data from (a-d) to query genotype-specific and treatment-specific effects. (e) The three different genotypes overlap in their median variances (large symbols) for the whole experiment, indicating that the variance in transcript response is independent on the genotype. (f) Including the treatment in the PCA analysis revealed no overlap between the median principal component between LL and LL→HL transcripts, indicating that the treatment accounted for the variance in the transcriptional response

Suppl. Fig. 4: Translated regions of representative transcripts from cluster 1 & 2. Genome Browser (IGV) views of RPF-seq (green) and total RNA-seq (blue) abundance (read depth) for loci of representative transcripts from cluster 1 & 2. Tracks are colored according to sample/time-point (each track represents overlay of n=3 biological replicates). Black arrow indicate potential ribosomal stalling sites. Gene model is symbolized as blue box with introns represented as dashed line, CDS as bold, and 5'- and 3'UTR as thin line with arrows indicating read direction.

Suppl. Fig. 5: Distribution and secondary structure of motif-carrying 5'UTRs. (a) Motif 1, 2 and 3 occurrences in their respective transcripts as revealed by MEME-suit. The secondary structure is given in Dot-Bracket notation and was calculated using the software *RNA Shapes Studio [44]*, with Motifs drawn at corresponding site(s) for motif 1 (green), 2 (yellow) and 3 (red), respectively. (b) Distribution of motif-carrying transcripts in translatomes are indicated as dots for transcripts carrying motif 1 (green), 2 (yellow) or 3 (red).

Suppl. Fig. 6: Reporter-construct analysis for translational regulatory elements and secondary structure. (a) Motif 1, 2 and 3 occurrences in their respective transcripts as revealed by MEME-suit. The secondary structure is given in Dot-Bracket notation and was calculated using the software RNA Shapes Studio *[44]*, with motifs drawn at corresponding site(s) for motif 1 (green), 2 (yellow) and 3 (red), respectively. (b) Distribution of motif-carrying transcripts in translatomes are indicated as dots for transcripts carrying motif 1 (green), 2 (yellow) or 3 (red).

Suppl. Fig. 7: Motifs are potential RBP binding sites. (a) 5'-biotinylated M3-RNA with and without unlabelled competitor M3-RNA was incubated with total protein extracts of LL or 10 min LL→HL shifted plants, respectively. Depicted is a representative blot. (b) RBP 1-D acrylamide gel. Biotinylated M1 and M2 RNAs were coupled to strep-tagged beads and incubated with protein extracts from LL treated plants. After washing, samples were eluted and loaded on one-dimensional acrylamide gels. The enriched bands (arrows) were cut out and analysed via MALDI. (c) RNA-EMSA was used to validate dose dependent interaction (ratio of bound to unbound) of *SAP3*-M1-RNA with arabidopsis cytosolic AtGapC1, AtGapC2 or GAPDH isolated from rabbit muscle GAPDH isoform C1 *in vitro*.

Suppl. Fig. 8: Motifs 1-3 and SAP2, SAP3 translational regulation are conserved between *A. thaliana* and Rice. OsSAP8 and OsSAP9 were identified as closest homologues of SAP2 and SAP3 from *A. thaliana*. (a) M1, M2 and M3 could be identified in the 5'-and 3'-UTR of the rice homologues with only slight variation. (b,c) qPCR-based quantification of *OsSAP8* and *OsSAP9* transcripts of total and polysomal RNA fractions after 10 min of L→H-transfer confirms conserved translational regulation of SAP2 and SAP3 between *A. thaliana* and Rice. Data represent means ± SD of n=2 independent experiments, asterisks indicate significance relative to control, two sided Student's t-test: $p < 0.05$.

Suppl. Fig. 9: Subcellular localization of rice OsSAP8, OsSAP9, OsCML49a and OsSAP49b in transfected arabidopsis protoplasts and their molecular interaction shown by FRET. (a) *A. thaliana* mesophyll protoplasts were transfected with OsSA8:YFP, OsSAP3:YFP, OsCML49b:CFP or OsCML49a:CFP vector DNA. Fluorescence images taken by CLSM for subcellular localization. (b) Following transfection with constructs, FRET efficiencies between CFP and YFP were quantified indicating conserved interaction between arabidopsis and rice SAP and CML49.

Suppl. Fig. 10: Gene expression map of *AtSAP2*, *AtSAP3*, *AtCML49* and *OsSAP8, OsSAP9* and *OsCML49a*. (a) Transcript abundance and expression pattern of *AtSAP2*, (b) *AtSAP3* and (f) *AtCML49* in tissues retrieved from the eFP Browser. (d) Transcript abundance and expression

pattern of *SAP8*, (e) *SAP9* and (f) *CML49a* in different rice tissues retrieved from the Rice eFP Browser. *CML49b* was missing in group 1 data source of Rice eFP Browser.

Suppl. Fig. 11: GO-Analysis of *CML49* & *SAP3* Co-Expressed Genes during HL Acclimation. (a) List of co-regulated targets of SAP3 and CML49 that are stronger deregulated in *cml49* than in *sap3* background were subjected to GO-term analysis via Cytoscape BiNGO. Dark red nodes represent categories that are significantly overrepresented in context of genome abundance. The size of node represents gene abundance in the co-regulated gene set to the corresponding A strong overrepresentation of response to high light intensity, response to heat and response to hydrogen peroxide was detected.

Suppl. Table 1. Oligonucleotides.

Target gene	ID number	Forward Sequence (5'–3')	Reverse Sequence (5' – 3')
HSP20-like	AT1G53540	GTTCGAAGGTTTCTTGACGC	TGTCTCCCTCCAATCCACTT
HSP70	AT3G12580	TCGACTGCTCAGACGACTAT	GAACCTAGCACGAGTGATGG
SAP2	AT1G51200	TGAGCCGATGACTGTTTCTG	TTGGTTTCGCAGCTTCTTCT
SAP3	AT2G27580	GGTGTGGTTTCGTTTTCTGC	GCTTTCACAATCGGATTCGC
VTC2	AT4G26850	GACTGCTGTGTTTGCCTTCA	CGGGCTCACTTCACCTAGAG
eEF1B	AT5G12110	ATCTCCACACAGAGCGAGGT	TTGCTAGCATTGGGAAAACC
SIG5	AT5G24120	TCCTCCTCGTGAGCAAGTTT	GGAGCCGAATTATGTTCCAA
PS3	AT3G47420	TTGGTTAACGGTCCTTACGC	CGTTACCGTAGCCAAAGCTC
OsSAP8	Os06G41010	CGATGTGGCAACTTGTACTG	TTAGCCTTGGCAATAGCATC
OsSAP9	Os07G07350	AAACAACTGTGGCTTCTTCG	CGTAAATGCCTTCTTCTCCA
OsActin	Os03G07181	AGGCCGTCCTCTCTCTGTAT	GGGGAGAGCATATCCTTCAT
Osα-tubulin	Os03G51600	TCTGAACGTGGATGTCAATG	GAGATCACTGGAGCGTAGGA
OsSAP8NotI		AAAAGCGGCCGCATGGAGCACAAGGAGACT	
OsSAP8EcoRI		TTTTGAATTCCTAAATTTTGTCAAGTTT	
OsSAP9NotI		AAAAGCGGCCGCATGGCACAAGAAAGCTGG	
OsSAP9EcoRI		TTTTGAATTCTCAGATCTTGTTGATCTT	
OsCML49NotI		AAAAGCGGCCGCATGGCCGGCTACCCGCCA	
OsCML49EcoRI		TTTTGAATTCTTATGCAATGAGAAAAGG	
rRNA depletion	rRNA1b	CAGCTGAGGCATCCTAACAGACCGGTAGACTTG	
rRNA depletion	rRNA21	AGTAAGCCCACCCCAAGATGAGTGCTCTC	
rRNA depletion	rRNA27	ACGACTCTCGGCAACGGATATCTCGGCTCTCGCATCGATGAAGAACG	
rRNA depletion	rRNA29	GGCCGAGGGCACGTCTGCCTGGGTGTCAC	
rRNA depletion	rRNA30	GACTGTGAAACTGCGAATGGCTCATTAA	
rRNA depletion	rRNA44	TAGCGACGTTGCTTTTTGATCCTTCG	
rRNA depletion	rRNA1c	GATAACCGTAGTAATTCTAGAG	
rRNA depletion	rRNA1	CATAAACGATGCCGACCAGGGATCAGCGG	
rRNA depletion	rRNA2	TTTATTAGATAAAAGGTCGACG	

Suppl. Table 2: Overview of *cis*-elements within the promoter regions of *CML49* and *SAP3* co-regulated genes.
The total number and type of *cis*-elements within the co-regulated subgroup was searched -1.000 bp upstream of the ATG region using PlantPAN2.0. The largest group of *cis*-elements belong to the WRKY-group, followed by MYB, MYB-related, SURE (sulfur-responsive element) and ABRE-like (early responsive to dehydration). Only few putative sites belong to the LS7 (salicylic acid-inducible element), C-box (bZIP factors), PHR1 (phosphate starvation response) and some represent cis-elements for light-dependent SORLIPs (Sequences Over represented in Light-Induced Promoters).

AGI	Name	MYB/ MYB-related	WBOX	SURE	PHR1	C-box	ABRElike	LS7	SORLIPs	Other
AT4G13800	unknown	29	37	35	4	2	20	4	17	24
AT5G52570	BETA-OHASE 2	24	36	29	6	2	27	4	30	26
AT5G49480	CP1	24	41	30	1	2	16	1	19	21
AT4G04020	FIB	22	39	18	5	3	19	3	26	16
AT5G58770	CPT4	35	35	22	3		23	3	30	29
AT3G56290	unknown	23	47	31	3	2	25	3	35	34
AT1G67360	REF	41	46	32	5	8	30	6	27	14
AT1G61800	GPT2	28	36	23	7	9	18	3	17	26
AT5G20150	SPX1	22	42	20	13	4	14	1	10	23
AT3G21890	BBX31	22	41	27	2	6	22	3	14	30
AT2G46830	CCA1	31	43	47	1	6	21	1	44	14
AT5G52640	HSP90.1	35	38	43	4	4	23	3	15	14
AT2G36800	DOGT1	32	48	44	5	6	28	3	30	14
AT5G48570	ROF2	24	49	30	2	4	22	1	15	14
AT1G52690	LEA7	27	44	27	5	8	32	4	28	25
AT2G24540	AFR	28	45	41	6	9	28	4	28	31
AT3G09350	Fes1A	34	41	40	3	9	26	9	25	21
AT4G21990	APR3	22	40	26	3	6	21	1	16	20
AT4G12400	HOP3	33	37	25	3	1	24	6	12	21
AT5G50800	ATSWEET13	26	43	32	4	7	24	5	30	19
AT2G37180	RD28	23	43	21	7	4	22	3	20	34
AT4G34550	PCH1-like	26	35	30	4	3	16	3	19	34
AT3G17609	HYH	27	38	27	6	5	24	3	28	30
AT5G64510	TIN1	34	44	34	4	12	27	9	30	30
AT1G01520	ASG4	33	34	45	3	2	29	7	38	26
AT1G32900	GBSS1	31	52	31	5	9	28	2	36	30
AT2G26150	HSFA2	20	60	46	0	4	19	3	30	34
AT5G12110	eEF1B	41	53	34	2	6	17	4	32	31
AT4G26850	VTC2	20	40	31	8	8	23	4	20	36
AT2G33380	RD20	29	42	26	6	7	38	3	22	29
AT3G21150	BBX32	26	41	39	5	6	17	5	30	23
AT3G09450	unknown	38	34	42	6	1	29	3	43	29
AT2G21320	BBX18	26	57	40	7	4	20	1	20	27
AT1G58520	RXW8	20	28	26	2	3	17	4	18	45
AT5G24120	SIG5	14	52	31	1	4	22	1	16	25
AT5G65380	MATE	24	47	31	4	8	32	7	33	29
AT3G53830	RCC1	24	40	35	6	4	31	2	18	27
AT3G17790	PAP17	37	44	34	7	8	28	6	31	29
AT1G54050	HSP20-like	44	40	30	6	5	15	3	38	31
AT1G62180	ATAPR2	29	53	24	5	3	26	3	27	28
AT3G21670	ATNPF6.4	22	44	27	3	1	25	2	21	37
AT3G12580	HSP70	20	42	24	4	1	18	2	11	32
AT2G46790	PRR9	22	43	26	6	3	19	1	44	33
AT1G77450	NAC032	31	45	35	4	6	33	8	35	30
AT2G15020	unknown	20	34	26	5	0	24	3	34	36
AT3G47420	PS3	30	42	10	7	2	21	5	18	27
AT5G63130	Phox/Bem1p	20	49	19	4	5	15	3	31	37
AT1G01720	ANAC2	29	39	50	2	6	22	2	36	23
AT4G26150	CGA1	26	34	27	5	2	15	2	14	36
AT1G74310	HSP101	20	35	33	5	2	13	1	17	34
	Total	1368	2112	1556	224	232	1148	173	1278	1368

Suppl. Table 3. uORF analysis of motif-carrying transcripts. The motif-carrying transcripts were screened for upstream open reading frames (uORF) coding for upstream peptides (uPEPs) using the online tool uPEPperoni.

Target gene	ID number	uPEPs Sequence (5'–3')	Position (start-end)
PHT3	AT2G17270	MRTLPTYTLFAFPFIAFL	(182, 235)
Methyltransferase superfamily protein	AT3G13440	MGLGRNLQTIRSSYKIRVF	(23, 79)
PDX1.2	AT3G16050	MEAHLVSFFCIRRRFAIPP	(14, 70)
RBL14	AT3G17611	MRVKRRSLWRICFRRDQAAAI	(8, 70)
Unknown protein	AT4G27010	MPDFELSASPICFVLLNPKTKTPRREKSSL	(53, 142)
HSFB1	AT4G36990	MEETKRNSDLLRSRVFLSGFYCWDWE	(267, 344)
SIG5	AT5G4088	MMMPSSSCFHFGLRFFISSIRDPPIRIDTQRSRVSTDMDMDKS	(159, 287)

References:

1.   Barber, J. and B. Andersson, *Too much of a good thing: light can be bad for photosynthesis.* Trends in Biochemical Sciences, 1992. **17**(2): p. 61-66.
2.   Niyogi, K.K., *Safety valves for photosynthesis.* Current Opinion in Plant Biology, 2000. **3**(6): p. 455-460.
3.   Baier, M. and K.J. Dietz, *Chloroplasts as source and target of cellular redox regulation: a discussion on chloroplast redox signals in the context of plant physiology.* J Exp Bot, 2005. **56**(416): p. 1449-62.
4.   Asada, K., *THE WATER-WATER CYCLE IN CHLOROPLASTS: Scavenging of Active Oxygens and Dissipation of Excess Photons.* Annu Rev Plant Physiol Plant Mol Biol, 1999. **50**: p. 601-639.
5.   Pfannschmidt, T., et al., *A novel mechanism of nuclear photosynthesis gene regulation by redox signals from the chloroplast during photosystem stoichiometry adjustment.* J Biol Chem, 2001. **276**(39): p. 36125-30.
6.   Nagashima, A., *The Multiple-Stress Responsive Plastid Sigma Factor, SIG5, Directs Activation of the psbD Blue Light-Responsive Promoter (BLRP) in Arabidopsis thaliana.* Plant and Cell Physiology, 2004. **45**(4): p. 357-368.
7.   Pogson, B.J., et al., *Plastid signalling to the nucleus and beyond.* Trends Plant Sci, 2008. **13**(11): p. 602-9.
8.   Häusler, R.E., et al., *How sugars might coordinate chloroplast and nuclear gene expression during acclimation to high light intensities.* Mol Plant, 2014. **7**(7): p. 1121-37.
9.   Vogel, M.O., et al., *Fast retrograde signaling in response to high light involves metabolite export, MITOGEN-ACTIVATED PROTEIN KINASE6, and AP2/ERF transcription factors in Arabidopsis.* Plant Cell, 2014. **26**(3): p. 1151-65.
10.   Crisp, P.A., et al., *Rapid recovery gene downregulation during excess-light stress and recovery in Arabidopsis.* Plant Cell, 2017.
11.   Pfalz, J. and T. Pfannschmidt, *Essential nucleoid proteins in early chloroplast development.* Trends Plant Sci, 2013. **18**(4): p. 186-94.
12.   Pfannschmidt, T., A. Nilsson, and J.F. Allen, *Photosynthetic control of chloroplast gene expression.* Nature, 1999. **397**(6720): p. 625-628.
13.   Beck, C.F., *Signaling pathways from the chloroplast to the nucleus.* Planta, 2005. **222**(5): p. 743-56.
14.   Koussevitzky, S., et al., *Signals from chloroplasts converge to regulate nuclear gene expression.* Science, 2007. **316**(5825): p. 715-9.
15.   Estavillo, G.M., et al., *Evidence for a SAL1-PAP chloroplast retrograde pathway that functions in drought and high light signaling in Arabidopsis.* Plant Cell, 2011. **23**(11): p. 3992-4012.
16.   Xiao, Y., et al., *Retrograde signaling by the plastidial metabolite MEcPP regulates expression of nuclear stress-response genes.* Cell, 2012. **149**(7): p. 1525-35.
17.   Park, S.W., et al., *Cyclophilin 20-3 relays a 12-oxo-phytodienoic acid signal during stress responsive regulation of cellular redox homeostasis.* Proc Natl Acad Sci U S A, 2013. **110**(23): p. 9559-64.
18.   Dietz, K.J., *Redox regulation of transcription factors in plant stress acclimation and development.* Antioxid Redox Signal, 2014. **21**(9): p. 1356-72.

19.   Schmitz, J., et al., *The essential role of sugar metabolism in the acclimation response of Arabidopsis thaliana to high light intensities.* Journal of Experimental Botany, 2014. **65**(6): p. 1619-1636.

20.   Wu, G.Z., et al., *Extensive Posttranscriptional Regulation of Nuclear Gene Expression by Plastid Retrograde Signals.* Plant Physiol, 2019. **180**(4): p. 2034-2048.

21.   Floris, M., et al., *Post-transcriptional control of light-harvesting genes expression under light stress.* Plant Mol Biol, 2013. **82**(1-2): p. 147-54.

22.   Petracek M.E., D.L.F., Huber S.C., Thompson W.F., *Light-Regulated Changes in Abundance and Polyribosome Association of Ferredoxin mRNA Are Dependent on Photosynthesis* Plant Cell, 1997. **9**(12): p. 2291-300.

23.   Moore, M., N. Gossmann, and K.J. Dietz, *Redox Regulation of Cyosolic Translation in Plants.* Trends in Plant Science, 2016. **21**(5): p. 388-397.

24.   Lokdarshi, A., et al., *Light Activates the Translational Regulatory Kinase GCN2 via Reactive Oxygen Species Emanating from the Chloroplast.* Plant Cell, 2020.

25.   Piques, M., et al., *Ribosome and transcript copy numbers, polysome occupancy and enzyme dynamics in Arabidopsis.* Mol Syst Biol, 2009. **5**: p. 314.

26.   Juntawong, P. and J. Bailey-Serres, *Dynamic Light Regulation of Translation Status in Arabidopsis thaliana.* Front Plant Sci, 2012. **3**: p. 66.

27.   Shin, J., et al., *The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner.* Plant Cell Environ, 2017. **40**(7): p. 997-1008.

28.   Robaglia, C., M. Thomas, and C. Meyer, *Sensing nutrient and energy status by SnRK1 and TOR kinases.* Curr Opin Plant Biol, 2012. **15**(3): p. 301-7.

29.   Nietzsche, M., et al., *A protein–protein interaction network linking the energy-sensor kinase SnRK1 to multiple signaling pathways in Arabidopsis thaliana.* Current Plant Biology, 2016. **5**: p. 36-44.

30.   Baena-Gonzalez, E., et al., *A central integrator of transcription networks in plant stress and energy signalling.* Nature, 2007. **448**(7156): p. 938-42.

31.   Mustroph, A., P. Juntawong, and J. Bailey-Serres, *Isolation of plant polysomal mRNA by differential centrifugation and ribosome immunopurification methods.* Methods Mol Biol, 2009. **553**: p. 109-26.

32.   Cho, H.Y., et al., *Quantitative phosphoproteomics of protein kinase SnRK1 regulated protein phosphorylation in Arabidopsis under submergence.* J Exp Bot, 2016. **67**(9): p. 2745-60.

33.   Nukarinen, E., et al., *Quantitative phosphoproteomics reveals the role of the AMPK plant ortholog SnRK1 as a metabolic master regulator under energy deprivation.* Sci Rep, 2016. **6**: p. 31697.

34.   Schepetilnikov, M. and L.A. Ryabova, *Recent Discoveries on the Role of TOR (Target of Rapamycin) Signaling in Translation in Plants.* Plant Physiol, 2018. **176**(2): p. 1095-1105.

35.   Ingolia, N.T., et al., *The ribosome profiling strategy for monitoring translation in vivo by deep sequencing of ribosome-protected mRNA fragments.* Nat Protoc, 2012. **7**(8): p. 1534-50.

36.   Juntawong, P., et al., *Translational dynamics revealed by genome-wide profiling of ribosome footprints in Arabidopsis.* Proc Natl Acad Sci U S A, 2014. **111**(1): p. E203-12.

37.   Bazin, J., et al., *Global analysis of ribosome-associated noncoding RNAs unveils new modes of translational regulation.* Proc Natl Acad Sci U S A, 2017. **114**(46): p. E10018-E10027.

38. Hsu, P.Y., et al., *Super-resolution ribosome profiling reveals unannotated translation events in Arabidopsis.* Proc Natl Acad Sci U S A, 2016. **113**(45): p. E7126-E7135.

39. Hooper, C.M., et al., *SUBA4: the interactive data analysis centre for Arabidopsis subcellular protein locations.* Nucleic Acids Res, 2017. **45**(D1): p. D1064-D1074.

40. Jimenez-Gonzalez, A.S., et al., *Functional and structural analysis of maize hsp101 IRES.* PLoS One, 2014. **9**(9): p. e107459.

41. Yamasaki, S., et al., *Identification of 5'-untranslated regions that function as effective translational enhancers in monocotyledonous plant cells using a novel method of genome-wide analysis.* Plant Biotechnol (Tokyo), 2018. **35**(4): p. 365-373.

42. Bailey, T.L. and C. Elkan, *Fitting a mixture model by expectation maximization to discover motifs in biopolymers.* Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology, 1994: p. 28-36.

43. Liu, M.J., et al., *Widespread translational control contributes to the regulation of Arabidopsis photomorphogenesis.* Mol Syst Biol, 2012. **8**: p. 566.

44. Janssen, S. and R. Giegerich, *The RNA shapes studio.* Bioinformatics, 2015. **31**(3): p. 423-5.

45. Vij, S. and A.K. Tyagi, *Genome-wide analysis of the stress associated protein (SAP) gene family containing A20/AN1 zinc-finger(s) in rice and their phylogenetic relationship with Arabidopsis.* Mol Genet Genomics, 2006. **276**(6): p. 565-75.

46. Wang, F., et al., *Overexpression of OsSAP16 Regulates Photosynthesis and the Expression of a Broad Range of Stress Response Genes in Rice (Oryza sativa L.).* PLoS One, 2016. **11**(6): p. e0157244.

47. Nishizawa, A., et al., *Arabidopsis heat shock transcription factor A2 as a key regulator in response to several types of environmental stress.* Plant J, 2006. **48**(4): p. 535-47.

48. Suzuki, N., et al., *Identification of the MBF1 heat-response regulon of Arabidopsis thaliana.* Plant J, 2011. **66**(5): p. 844-51.

49. Nishiyama, Y., S.I. Allakhverdiev, and N. Murata, *Protein synthesis is the primary target of reactive oxygen species in the photoinhibition of photosystem II.* Physiol Plant, 2011. **142**(1): p. 35-46.

50. Belbin, F.E., et al., *Integration of light and circadian signals that regulate chloroplast transcription by a nuclear-encoded sigma factor.* New Phytol, 2016.

51. Niyogi, K.K., *PHOTOPROTECTION REVISITED: Genetic and Molecular Approaches.* Annu Rev Plant Physiol Plant Mol Biol, 1999. **50**: p. 333-359.

52. Hebbelmann, I., et al., *Multiple strategies to prevent oxidative stress in Arabidopsis plants lacking the malate valve enzyme NADP-malate dehydrogenase.* J Exp Bot, 2012. **63**(3): p. 1445-59.

53. Oelze, M.L., et al., *Efficient acclimation of the chloroplast antioxidant defence of Arabidopsis thaliana leaves in response to a 10- or 100-fold light increment and the possible involvement of retrograde signals.* Journal of Experimental Botany, 2012. **63**(3): p. 1297-1313.

54. Dietz, K.J., M.O. Vogel, and A. Viehhauser, *AP2/EREBP transcription factors are part of gene regulatory networks and integrate metabolic, hormonal and environmental signals in stress acclimation and retrograde signalling.* Protoplasma, 2010. **245**(1-4): p. 3-14.

55. Browning, K.S. and J. Bailey-Serres, *Mechanism of cytoplasmic mRNA translation.* Arabidopsis Book, 2015. **13**: p. e0176.

56. Moore, M., N. Gossmann, and K.J. Dietz, *Redox Regulation of Cytosolic Translation in Plants.* Trends Plant Sci, 2016. **21**(5): p. 388-97.

57. Zhang, Y., et al., *GCN2-dependent phosphorylation of eukaryotic translation initiation factor-2alpha in Arabidopsis.* J Exp Bot, 2008. **59**(11): p. 3131-41.

58. Wek, R.C., H.Y. Jiang, and T.G. Anthony, *Coping with stress: eIF2 kinases and translational control.* Biochemical Society Transactions, 2006. **34**(1): p. 7-11.

59. Clemens, M.J., A. Elia, and S.J. Morley, *Requirement for the eIF4E binding proteins for the synergistic down-regulation of protein synthesis by hypertonic conditions and mTOR inhibition.* PLoS One, 2013. **8**(8): p. e71138.

60. Richter, J.D. and N. Sonenberg, *Regulation of cap-dependent translation by eIF4E inhibitory proteins.* Nature, 2005. **433**(7025): p. 477-80.

61. Bruns, A.N., et al., *Phosphorylation of Arabidopsis eIF4E and eIFiso4E by SnRK1 Inhibits Translation.* FEBS J, 2019.

62. Rossel, J.B., I.W. Wilson, and B.J. Pogson, *Global changes in gene expression in response to high light in Arabidopsis.* Plant Physiol, 2002. **130**(3): p. 1109-20.

63. Branco-Price, C., et al., *Selective mRNA translation coordinates energetic and metabolic adjustments to cellular oxygen deprivation and reoxygenation in Arabidopsis thaliana.* Plant J, 2008. **56**(5): p. 743-55.

64. Chang, C.H., et al., *Posttranscriptional control of T cell effector function by aerobic glycolysis.* Cell, 2013. **153**(6): p. 1239-51.

65. White, M.R. and E.D. Garcin, *The sweet side of RNA regulation: glyceraldehyde-3-phosphate dehydrogenase as a noncanonical RNA-binding protein.* Wiley Interdiscip Rev RNA, 2016. **7**(1): p. 53-70.

66. Marondedze, C., et al., *Changes in the Arabidopsis RNA-binding proteome reveal novel stress response mechanisms.* BMC Plant Biol, 2019. **19**(1): p. 139.

67. Stroher, E., et al., *Redox-Dependent Regulation of the Stress-Induced Zinc-Finger Protein SAP12 in Arabidopsis thaliana.* Molecular Plant, 2009. **2**(2): p. 357-367.

68. Giri, J., et al., *Rice A20/AN1 zinc-finger containing stress-associated proteins (SAP1/11) and a receptor-like cytoplasmic kinase (OsRLCK253) interact via A20 zinc-finger and confer abiotic stress tolerance in transgenic Arabidopsis plants.* New Phytol, 2011. **191**(3): p. 721-32.

69. Kang, M., et al., *Arabidopsis SAP5 functions as a positive regulator of stress responses and exhibits E3 ubiquitin ligase activity.* Plant Mol Biol, 2011. **75**(4-5): p. 451-66.

70. Ferre, F. and P. Clote, *DiANNA: a web server for disulfide connectivity prediction.* Nucleic Acids Res, 2005. **33**(Web Server issue): p. W230-2.

71. Kim, G.D., Y.H. Cho, and S.D. Yoo, *Regulatory functions of evolutionarily conserved AN1/A20-like Zinc finger family proteins in Arabidopsis stress responses under high temperature.* Biochem Biophys Res Commun, 2015. **457**(2): p. 213-20.

72. Vanderbeld, B. and W.A. Snedden, *Developmental and stimulus-induced expression patterns of Arabidopsis calmodulin-like genes CML37, CML38 and CML39.* Plant Mol Biol, 2007. **64**(6): p. 683-97.

73. Dobney, S., et al., *The calmodulin-related calcium sensor CML42 plays a role in trichome branching.* J Biol Chem, 2009. **284**(46): p. 31647-57.

74. Wu, H.C., et al., *Heat shock-induced biphasic Ca(2+) signature and OsCaM1-1 nuclear localization mediate downstream signalling in acquisition of thermotolerance in rice (Oryza sativa L.).* Plant Cell Environ, 2012. **35**(9): p. 1543-57.

75. Wu, H.C. and T.L. Jinn, *Oscillation regulation of Ca2+ /calmodulin and heat-stress related genes in response to heat stress in rice (Oryza sativa L.).* Plant Signal Behav, 2012. **7**(9): p. 1056-7.

76.   Kushwaha, R., A. Singh, and S. Chattopadhyay, *Calmodulin7 plays an important role as transcriptional regulator in Arabidopsis seedling development.* Plant Cell, 2008. **20**(7): p. 1747-59.

77.   Rentel, M.C. and M.R. Knight, *Oxidative stress-induced calcium signaling in Arabidopsis.* Plant Physiol, 2004. **135**(3): p. 1471-9.

78.   Takahashi, F., et al., *Calmodulin-dependent activation of MAP kinase for ROS homeostasis in Arabidopsis.* Mol Cell, 2011. **41**(6): p. 649-60.

79.   Meiri, D. and A. Breiman, *Arabidopsis ROF1 (FKBP62) modulates thermotolerance by interacting with HSP90.1 and affecting the accumulation of HsfA2-regulated sHSPs.* Plant J, 2009. **59**(3): p. 387-99.

80.   Jung, H.S., et al., *Subset of heat-shock transcription factors required for the early response of Arabidopsis to excess light.* Proc Natl Acad Sci U S A, 2013. **110**(35): p. 14474-9.

81.   Linster, C.L. and S.G. Clarke, *L-Ascorbate biosynthesis in higher plants: the role of VTC2.* Trends Plant Sci, 2008. **13**(11): p. 567-73.

82.   Mellenthin, M., et al., *Expression of the Arabidopsis Sigma Factor SIG5 Is Photoreceptor and Photosynthesis Controlled.* Plants (Basel), 2014. **3**(3): p. 359-91.

83.   Bethke, G., et al., *Flg22 regulates the release of an ethylene response factor substrate from MAP kinase 6 in Arabidopsis thaliana via ethylene signaling.* Proceedings of the National Academy of Sciences, 2009. **106**(19): p. 8067-8072.

84.   Booth, D.S., A. Avila-Sakar, and Y. Cheng, *Visualizing proteins and macromolecular complexes by negative stain EM: from grid preparation to image acquisition.* J Vis Exp, 2011(58).

85.   Chotewutmontri, P., et al., *Ribosome Profiling in Maize.* Methods Mol Biol, 2018. **1676**: p. 165-183.

86.   Liao, Y., G.K. Smyth, and W. Shi, *The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote.* Nucleic Acids Res, 2013. **41**(10): p. e108.

87.   Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.* EMBnet.journal, 2011. **17**(1): p. 10.

88.   Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat Methods, 2012. **9**(4): p. 357-9.

89.   Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program for assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-30.

90.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for differential expression analysis of digital gene expression data.* Bioinformatics, 2010. **26**(1): p. 139-40.

91.   McCarthy, D.J., Y. Chen, and G.K. Smyth, *Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation.* Nucleic Acids Res, 2012. **40**(10): p. 4288-97.

92.   Irizarry, R.A., et al., *Summaries of Affymetrix GeneChip probe level data.* Nucleic Acids Research, 2003. **31**(4): p. 15e-15.

93.   Benjamini, Y. and Y. Hochberg, *Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing.* Journal of the Royal Statistical Society, 1995. **57**(1): p. 289-300.

94.   Benjamini, Y. and Y. Hochberg, *On the Adaptive Control of the False Discovery Rate in Multiple Testing With Independent Statistics.* Journal of Educational and Behavioral Statistics, 2000. **25**(1): p. 60-83.

95.   Wormuth, D., et al., *Regulation of gene expression by photosynthetic signals triggered through modified CO2 availability.* BMC Plant Biol, 2006. **6**: p. 15.

96.   Oelze, M.L., et al., *Efficient acclimation of the chloroplast antioxidant defence of Arabidopsis thaliana leaves in response to a 10- or 100-fold light increment and the possible involvement of retrograde signals.* J Exp Bot, 2012. **63**(3): p. 1297-313.

97.   Holtgrefe, S., et al., *Regulation of plant cytosolic glyceraldehyde 3-phosphate dehydrogenase isoforms by thiol modifications.* Physiol Plant, 2008. **133**(2): p. 211-28.

98.   Sprenger-Haussels, M. and B. Weisshaar, *Transactivation properties of parsley proline-rich bZIP transcription factors.* The Plant Journal, 2000. **22**(1): p. 1-8.

99.   Panda, A., J. Martindale, and M. Gorospe, *Affinity pulldown of biotinylated RNA for detection of protein-RNA complexes.* Bio-Protocol, 2016. **6**(24).

100.   Chow, C.N., et al., *PlantPAN 2.0: an update of plant promoter analysis navigator for reconstructing transcriptional regulatory networks in plants.* Nucleic Acids Res, 2016. **44**(D1): p. D1154-60.

101.   Skarshewski, A., et al., *uPEPperoni: an online tool for upstream open reading frame location and analysis of transcript conservation.* BMC Bioinformatics, 2014. **15**: p. 36.

102.   Winter, D., et al., *An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets.* PLoS One, 2007. **2**(8): p. e718.

103.   Lalonde, S., et al., *A membrane protein/signaling protein interaction network for Arabidopsis version AMPv2.* Front Physiol, 2010. **1**: p. 24.

104.   Obrdlik, P., et al., *K+ channel interactions detected by a genetic system optimized for systematic studies of membrane protein interactions.* Proc Natl Acad Sci U S A, 2004. **101**(33): p. 12242-7.

105.   Seidel, T., D. Golldack, and K.J. Dietz, *Mapping of C-termini of V-ATPase subunits by in vivo-FRET measurements.* FEBS Lett, 2005. **579**(20): p. 4374-82.

Figure 1

a   Time- and Dose-Dependent Ribosome Association

100-fold Light Shift   10-fold Light Shift

— LL
– – 10' LL→ HL
······ 60' LL→ HL

— NL
– – 30' NL→ HL
---- 60' NL→ HL

Abs (256 nm)

Sucrose Density Fractionation ->   Sucrose Density Fractionation ->

b   Ribosome Association of RNA During LL→HL→LL Shift

Ribosome associated RNA
[(ribosome mRNA /total mRNA)/FW]

| 0' | 10' | 30' | 10' | 50' |
| Low Light (LL) | Low Light to High Light (LL→ HL) | | Recovery (LL→ HL→ LL) | |

c   Ribosomal Fractions at 10 min LL→ HL

— LL
– – 10' LL→ HL

80S

60S

Polysome

2x   3-5x

Abs (256 nm)

Sucrose Density Fractionation ->

Ratio (↓LL→HL/↓LL)	60S	80S	Polysome
	0.85	0.79	1.01

| TEM Image of Fraction | 80S | Polysome (2x) | Polysome (3-5x) |

Figure 2

a   **Retrograde Signalling Mutants: LL**

LL Control

Abs (256 nm)

60S   80S

Polysome

WT
mpk 6
akin 10

Sucrose Density Fractionation

b   **Retrograde Signalling Mutants: LL→HL Shift**

LL→ HL

Abs (256 nm)

60S   80S

Polysome

WT
mpk 6
akin 10

Sucrose Density Fractionation

c   **Genotype Dependent LL→HL Response**

Polysome Profile Area ($\int$Treatment/$\int$Control$_{WT}$)

Genotype	Treatment	60S	80S	Polysome
WT	LL	1.00	1.00	1.00
	LL→ HL	0.85*	0.79*	1.01
mpk 6	LL	0.91	0.91	0.77*
	LL→ HL	0.55*	0.51*	0.57*
akin 10	LL	0.67*	0.52*	0.98
	LL→ HL	0.71*	0.62*	0.99

* Indicating significant difference to WT LL

Figure 3

a Regulatory Patterns of Translatomic Changes in LL→HL Leaves

Cluster 1

GO: Photosynthesis enriched

YCF37

PSI LHCB1.1

RuBisCO S2B/S3B

PSI sub. P

Cyt b6f

chl. HSP 70-1

SAP2

Cluster 2

GO: Abiotic stress

GRXC11

HSF2A

DREB2A

HSP70

SAP3

Cluster 3

GO: Negative regulation of transcription

OFP15

WRKY70

RAV2

Cluster 4

GO: Response to Auxin

Ribosome Association
(FC Ribosomal mRNA LL→HL / LL)

Transcriptional Changes
(FC total mRNA LL→HL / LL)

b Fold Change Response of Cluster Depends on Transcript Abundance

Ribosome Association
(FC Ribosomal mRNA
LL→ HL / LL [RPKM])

Cluster 1 Cluster 2 Cluster 3 Cluster 4

Total measured Transcript (Abundance [CPM])

Figure 4

a

5'-UTR Motifs and their Effect on Translation in *in vivo* Reporter Assay

b *In vitro* GAPC1-Binding Activity to 5'-UTR Sequences

c *Ex vivo* Light-Dependent Interaction of GAPC1/2 with SAP3 Motif1

SAP2 and SAP3 Interaction and Subcellular Repartitioning with CML49

a Y2H SAP2 and SAP3 Interaction with CML49

b *In-vivo* Co-Localization of SAP2, SAP3 and CML49

c FRET Interaction of SAP2, SAP3 and CML49

Figure 6

a Co-Regulated Transcripts between *sap3* and *cml49*

Gene ID	Target	WT	sap3 KO	cml49 KO	compartment
AT4G13800	unknown				membrane, plasma membrane
AT5G52570	BETA-OHASE 2				chloroplast, chloroplast membrane
AT5G49480	CP1				cytoplasm
AT4G04020	FIB				chloroplast
AT5G58770	CPT4				nucleus
AT3G56290	unknown				mitochondrium
AT1G67360	REF				cytoplasm, vacuola
AT1G61800	GPT2				chloroplast
AT5G20150	SPX1				cytoplasm, nucleus
AT3G21890	BBX31				cytoplasm, nucleus
AT2G46830	CCA1				nucleus
AT5G52640	HSP90.1				cytoplasm
AT2G36800	DOGT1				chloroplast, membrane
AT5G48570	ROF2				cytoplasm, vacuola
AT2G24540	AFR				cytoplasm
AT3G09350	Fes1A				cytoplasm, nucleus
AT4G21990	APR3				chloroplast
AT4G12400	HOP3				cytoplasm, nucleus
AT5G50800	ATSWEET13				membrane, plasma membrane
AT2G37180	RD28				membrane, plasma
AT4G34550	PCH1-like				chromosome, cytoplasm, nucleus
AT3G17609	HYH				nucleus
AT5G64510	TIN1				chloroplast, endoplasmic reticulum
AT1G01520	ASG4				nucleus
AT1G32900	GBSS1				chloroplast
AT2G26150	HSFA2				chloroplast, cytoplasm, nucleus
AT5G12110	eEF1B				cytoplasm
AT4G26850	VTC2				cytoplasm, nucleus
AT2G33380	RD20				chloroplast membrane, endoplasmic reticulum
AT3G21150	BBX32				cytoplasm, nucleus
AT3G09450	unknown				membrane
AT2G21320	BBX18				nucleus
AT1G58520	RXW8				chloroplast
AT5G24120	SIG 5				chloroplast, mitochondrium
AT5G65380	MATE				membrane
AT3G53830	RCC1				cytoplasm, nucleus
AT3G17790	PAP17				cell surface
AT1G54050	HSP20-like				cytoplasm
AT1G62180	ATAPR2				chloroplast
AT3G21670	ATNPF6.4				membrane
AT3G12580	HSP70				membrane
AT2G46790	PRR9				nucleus
AT1G77450	NAC032				nucleus
AT2G15020	unknown				mitochondrium
AT3G47420	PS3				membrane
AT5G63130	Phox/Bem1p				nucleus
AT1G01720	ANAC2				nucleus
AT4G26150	CGA1				nucleus
AT1G74310	HSP101				chloroplast stroma, cytoplasm, cytosol, nucleus

(log₂ FC)   -2 ... 4

Kinetics of Selected Co-Regulated Transcripts between SAP3 and CML49

Figure 7