Titel

Retrograde Control of Cytosolic Translation Targets Synthesis of Plastid Localized Proteins

and Nuclear Responses for Efficient Light Acclimation.

Marten Moore[1,2], Aaron Smith[2], Corinna Wesemann[1], Sonja Schmidtpott[1], Melanie Wegener[1],

Muhammad Ansar Farooq[1], Thorsten Seidel[1], Barry Pogson[2] and Karl-Josef Dietz[1§].

[1]Biochemistry and Physiology of Plants, Bielefeld University, 33501 Bielefeld, Germany.

[2]ARC Centre of Excellence in Plant Energy Biology, The Australian National University,

Canberra, ACT 2601, Australia.

Keywords: High light, chloroplast, retrograde signalling, stress acclimation, transcriptome,

translational control,

§ Address for correspondence

Karl-Josef Dietz§
Biochemistry and Physiology of Plants – W5-134
Bielefeld University
33501 Bielefeld
Germany
Tel. +49 521 1065589
Fax. +49 521 1066039
E-mail: karl-josef.dietz@uni-bielefeld.de

List of Abbreviation: ABA: abscisic acid; *A. thaliana*: *Arabidopsis thaliana*; AKIN10: protein kinase being part of the sucrose non-fermenting kinase complex; Asc: ascorbate; CaM: calmodulin; CML: calmodulin-like; DEG: differentially expressed gene; DHA: dehydroascorbate; GAPC: glyceraldehyde-3-phosphate dehydrogenase 2; GCN2: general control non-repressible 2; GO: gene-ontology; GSH: glutathione; GSSG: glutathione disulfide; H-light: Highlight (800 μmol photons m^{-2} s^{-1}); H_2O_2: hydrogen peroxide; HSE: heat shock element; HSF: heat shock factor; HSP: heat shock protein; IRES: independent ribosomal entry sites; JA: jasmonic acid; L-light: Lowlight (8 μmol photons m^{-2} s^{-1}); $\log_2$: logarithmic fold change to base 2; MAPK: mitogen-activated protein kinase; mbSUS: mating-based Split-ubiquitin-System; *O. sativa*: *Oryza sativa*; OPDA: 12-oxo-phytodienoic acid; ORF: open reading frame; PEG: polyethylene glycol; PET: photosynthetic electron transport; RBP: RNA binding protein; PPLT: preferentially polysome-loaded transcript; PPUT: preferentially polysome-unloaded transcript; ROS: reactive oxygen species; SAP: stress associated protein; SnRK1: SNF1-related kinase 1; TE: translation efficiency; TF: transcription factor; TOR: target of rapamycin; TPT: triosophosphate/phosphate translocator; UTR: untranslated region; VTC2: GDP-L-galactose phosphorylase;

Abstract

Canonical retrograde signaling is the transmission of information from organelles to the nucleus. Discrepancies between protein accumulation and transcript abundance in response to oxidative stress were suggestive of protein translation responding to retrograde signaling. Here we uncover multiple components of a translation-dependent retrograde signaling pathway that impact translation efficiency and gene expression, including the kinases, MPK6 and the SnRK1 subunit, AKIN10. Global ribosome foot-printing demonstrated rapid differential loading of 939 of transcripts from polyribosomes within 10 min after transfer from Low to High-light. Translationally regulated transcripts shared motifs in their 5`-UTR that act as binding sites for RBPs such as GAPC. The Stress Associated Proteins 2 and 3 carry such motifs in their UTRs and interact with the calcium sensor Calmodulin-like 49, relocating to the nucleus to co-regulate a translation-dependent transcriptional response. Translation dependent retrograde signaling bifurcates into a direct translational circuit and a translation-reliant nuclear circuit synchronizing translation, nuclear and anterograde response pathways, which may serve as a just in time-provision of needed proteins to the plastids.

Introduction

Introduction

Light impacts plant morphogenesis, development and metabolism. Gains in biomass and fitness strongly correlates with light-dependent photosynthesis, with starch accumulating during the light phase being essential for metabolism and growth during the dark phase. However, excess light leads to an over-reduction of the photosynthetic electron transport (PET) chain and enhances production of reactive oxygen species (ROS) (1). The natural light environment fluctuates continually on the time scales of seconds, hours and days. To cope with rapid light fluctuations, plants control their capacity to dissipate excess energy and minimize photo-oxidative damage (2). ROS accumulating in excess light are detoxified by the water-water cycle in an ascorbate-dependent or ascorbate-independent process utilizing ascorbate peroxidases and peroxiredoxins, respectively (3, 4). On longer time scales, changes in light quantity alter photosystem stoichiometry and increase the photosystem repair capacity.

Proteins of the photosynthetic machinery are encoded in the plastome and the nuclear genome while all antioxidant enzymes are exclusively encoded in the nucleus, as is the majority of the plastidial proteins. Transcription, stability and translation of chloroplast mRNA strongly depends on nuclear genome-encoded sigma factors and RNA binding proteins (RBPs) (5). The reorganization of photosystems and readjustment of the antioxidant systems, therefore, rely on a coordinated production of nucleus- and chloroplast-encoded proteins (6). Coordination is realized by transmitting information on the metabolic state of the chloroplast as retrograde signals to the nucleus. Alterations in light quality and quantity result in fluctuations of metabolites and signal export from the chloroplast with impact on accumulation and stability of mRNA.

Several signaling pathways are involved in developmental and operational retrograde control of nuclear gene expression. They include signals from the plastoquinone and carotenoid pools, tetrapyrrole biosynthesis, dihydroxyacetone phosphate (DHAP), reactive oxygen species (ROS), ascorbate/dehydroascorbate (Asc/DHA) and glutathione/glutathione disulphide (GSH/GSSG), hormones like 12-oxo-phytodienoic acid (OPDA) that also functions as precursor for jasmonic acid (JA), and xanthoxin as precursor of abscisic acid (ABA), the SAL1/PAP pathway and sugars (3, 6-16). Rapid regulation of transcription in response to excess light involves export of triosephosphates, such as DHAP, via the triosephosphate/phosphate translocator (TPT2) and

Mitogen Activated Protein Kinase 6 (MPK6) and couples to the upregulation of a set of

Apetala2/Ethylene Response Factor (AP2/ERF) transcription factors (16). It is worth noting that

TPT2 co-controls the export of the DHAP retrograde signal (17). Transcriptome analyses (RNA-

seq, qPCR, sRNA-seq, and PARE) during repeated excess-light stress and recovery revealed

global, rapid and specific changes in transcript accumulation and rate of decay in response to

changing light regimes (18).

The retrograde signaling pathways identified to date target transcriptional control, while cytosolic

translation as a retrograde target of photosynthesizing chloroplasts is mostly unexplored,

particularly at rapid signaling time scales. While there are descriptive studies focusing on the

effects of translational control under the effects of the highly discussed gun mutants (19), very little

evidence of the mechanisms of cytosolic control of translation as a consequence of retrograde

signaling is available. This gap in knowledge needs to be closed since translational control has the

potential for more immediate spatial and temporal adjustment of the chloroplast proteome and may

better account for proteins whose abundance is regulated post-transcriptionally. Redox regulation

of the translational apparatus has been suggested as one such translational control mechanism (20).

Recently, a link between the light stress-associated increase in ROS leading and the rapid activation

of GCN2 kinase, thus inhibiting the translation initiation pathway of GCN2-eIF2α, has been

reported (21). This process also affected the translation of transcripts encoding mitochondrial ATP

synthesis and chloroplast thylakoid proteins.

Post-transcriptional regulation of mRNA stability and translation is known from circadian

rhythmicity and involves light entrainment (22). mRNAs encoding enzymes of central metabolism

associate with the ribosomes in dependence on light and input from the circadian clock. The impact

of circadian rhythmicity on the steady-state pool of polysome-recruited transcripts was lost if the

shift to darkness or re-illumination occurred out of phase (23). The energy sensor kinase AKIN10

entrains the circadian clock depending on light input (24). AKIN10 is part of the Sucrose Non-

Fermenting 1-Related Kinase 1 (SnRK1) complex and acts as a central regulator of transcription

and translation networks via the TOR complex (25-27).

We felt that there is a fundamental need to address the significance and mechanism of translational

regulation of light acclimation by retrograde signaling. Retrograde control of vicinal translation is

hypothesized as straightforward mechanism for chloroplasts to influence their protein complement

in a much faster way or even totally independent on nuclear gene expression.

To challenge this hypothesis, global polysome loading was used as readout of current protein

synthesis of Arabidopsis leaves following a 10 min exposure to H-light. Results revealed a rapid

reorganization of the translatome and identified a set of preferentially polysome-associated

mRNAs. Among the stress-associated proteins with Zn-binding domains, SAP2 and SAP3 attracted

our attention. The calmodulin-like protein CML49 was an interacting partner of SAP2 and SAP3.

Transcript profiling of *sap3* and *cml49* knockout lines following H-light exposure revealed

downstream target genes critically involved in light acclimation. Finally, the centrality of the

AKIN10 kinase in coordinating energy availability, metabolic activity and stress placed this kinase

in the focus of this work on retrograde control of translation. This study provides first conclusive

evidence for rapid retrograde effects on cytosolic translation.

Results

Rapid initiation of retrograde translational signaling

The rapid reorganization of the translatome was addressed in Arabidopsis whole rosettes following

a sudden change in photosynthetic irradiance. We used our established shift treatment of low (L-

light) (8 µmol photons m^{-2} s^{-1}) to H-light (800 µmol photons m^{-2} s^{-1}) for 0 or 10 min (L→H-transfer)

or by returning to L-light for 10 and 50 min (L→H→L) prior to harvest consistent with established

analyses (16, 28). To assess the effect of retrograde signaling on translation, mRNAs associated

with multiple ribosomes (polysomes) were isolated to identify actively translated mRNA, via

comparison with the global mRNA population (total RNA). To this end, ribosomes were arrested

on their correspondent mRNA and isolated through sucrose gradient centrifugation, subsequently

RNase I-digested and the resulting ribosome-protected fragments processed for polysome-

protected fragment sequencing (translatome) in parallel to total RNA sequencing (total

transcriptome) as described (29). The traditional ribosome stalling reagents for this protocol are

cycloheximide and chloramphenicol which mainly stall the 80S ribosome. Additionally, the 70S

ribosome inhibitor lincomycin was used during polysome extraction and following protocol steps.

This change in protocol allowed for the first time the simultaneous assessment of cytosolic

translation as well as translation in mitochondria and chloroplasts. The fraction of polysome-loaded

mRNA strongly declined after 10 min of L→H-transfer, but exceeded the control level after

transfer from L➔H➔L for 10 or 50 min (Figure 1 a, Suppl. Fig. 1 f-g.). The results indicate a rapid

unloading and reloading of polysomes in response to changing light conditions.

Translation efficiency (TE), defined as relative ribosome loading, was calculated by dividing the

abundance of a given transcript in the ribosome-associated fraction by its abundance in the total

RNA pool, normalized by library size. The TE is therefore a measure of the proportion of

transcripts for a given gene undergoing active translation at a given time. A set of 939 transcripts

(false discovery rate (FDR) adjusted p-value (padj) < 0.05) was differentially loaded after short-

term exposure to H-light (Figure 1 b, Suppl. Dataset 1). The group of TE-downregulated transcripts

(571) exceeded that of TE-upregulated (368) transcripts by a factor of 1.55 (Figure 1 b), further

supporting the finding of global downregulation of translation after L➔H-transfer. The comparison

of transcripts with altered polysome association relative to total abundance unveiled more

regulation at the translational than transcriptional level. Sets of 289 and 442 transcripts with

increased or decreased abundance responded only translationally, whereas much smaller sets of

128 (up) and 51 (down) transcripts were exclusively transcriptionally regulated (Figure 1 c).

The key question is how excess light triggers a change in translational efficiency and was addressed

by scrutinizing known translation-regulating factors. MAP kinase-dependent phosphorylation of

the SnRK1 subunit AKIN10 leads to regulation of translation via TOR signaling and we have

previously shown that MPK6 phosphorylation depends on functional TPT, which exports DHAP

as retrograde signal (30). The dynamic unloading of ribosomes during L➔H seen in the WT

genotype was absent in *akin10* and was dampened in *mpk6* background as could be visualized via

polysomal profiling and fractioning (Suppl. Fig. 1 f-g.). To assess the shift of polysome-associated

mRNAs not only quantitatively but also qualitatively, we selected 89 transcripts whose

translational efficiency increased. This set contained photosynthesis and stress response genes such

as *SAP2* and *SAP3*. In addition, we selected three genes from the 442 transcripts whose translational

efficiency decreased, namely *ERF105*, *HSP70* and *DREB2A*. All five transcripts showed increased

polysome-loading in response to 10 min H-light but either showed increased or decreased

translational efficiency due to changing total mRNA amount. Compared to WT, TE of 4 of the 5

transcripts was significantly altered in *akin10* background, with *SAP2* and *HSP70* also being

significantly altered in *mpk6* background (Suppl. Fig. 1 a-e. and Suppl. Table 1). Interestingly, the

increase of TE of *SAP2*, *SAP3* and *HSP20* was diminished in *akin10*, while the decrease of TE in

ERF105, *DREB2A* and *HSP70* was less accentuated.

Evidence for a translational retrograde circuit enriched in photosynthesis proteins

Clustering of the differentially accumulating transcripts in the polysomal versus total RNA fractions allowed discrimination of subsets that revealed distinctive regulatory responses (Figure 1 d.). Cluster 1 includes transcripts that exclusively increase their polysome loading, while total mRNA abundance was insignificantly changed (119 DEGs; preferentially polysome-loaded transcripts; PPLT). Transcripts of cluster 2 are unloaded from polysomes (466 DEGs; preferentially polysome-unloaded transcripts PPUT). Both clusters were significantly altered in polysomal fold-change but not significantly altered in total RNA fold-change. Clusters 3 and 4 comprise transcripts that upon transfer to H-light simultaneously changed in the polysomal and total RNA fraction, 62 up and 123 down regulated DEGs (cluster 3 and 4, respectively; Figure 1 d; Poly and Total). Others were not assigned to clusters, e.g., 33 DEGs that were changed in total mRNA but not in the polysomal fraction and termed as buffered transcripts.

The polysome-loaded cluster (PPLT, Cluster 1) was five-fold enriched in transcripts coding for chloroplast-localised proteins according to SUBA4 (31). Among those, GO terms for photosynthesis are highly overrepresented (8.7 % in comparison to 0.5 % of genome representation), in particular for photosystem II light harvesting (1.9 %). Those include minor (AT3G47470) and major light harvesting proteins (AT4G10340), subunits of photosystem I (AT2G46820), one of the two nuclear encoded cytochrome b6f subunits (AT2G26500) and NADPH oxidoreductase subunits participating in electron transport (AT1G06690) as well as *small subunits of RuBisCO* (AT5G38410) (see Suppl. Dataset 1, PPLT). Apart from photosynthetic proteins, this set included proteins participating in assembly, damage and repair, e.g., chloroplast-localised *HSC70-1* (AT5G02500), which is involved in protein import; chloroplast vesiculation factors (AT4G32150) taking part in partitioning damaged proteins for proteolysis; and *PYG7/YCF37* (AT2G23670) which is required for PSI assembly/repair. The transcripts coding for cytosolic proteins were enriched in GO terms for translation (15 % in comparison to 4.9 % of genome representation), e.g., proteins of *ribosomal large subunit* (AT3G08520).

Cluster 2 comprising solely translationally downregulated DEGs (PPUT) (Figure 1 d) was enriched in multiple GO terms, e.g., response to auxin 5.7 %, chlorophyll biosynthetic process 2.0 % and response to far red light 2.6 %, but a prevailing response was undetectable. The transcripts upregulated both in the polysomal and total fraction in cluster 3 (Figure 1 d, poly and total) were enriched in abiotic stress GO terms: heat stress (48.1 % in comparison to 0.5 % of genome

representation); response to hydrogen peroxide (20.3 %); and light stimulus (20.3 %; Suppl. Dataset 1). This cluster included small (*HSP20-likes*) and large heat shock proteins (*HSP70*, *HSP101*), their organizing factors (*HOP*) as well as their key regulatory transcription factors *HSFA2* (AT2G26150), *HSF2BA* (AT5G62020), *HSFB2B* (AT4G11660) and *DREB2A* (AT5G05410).

Plastid translation regulated in response to rapid light shift

While cytosolic translation was increased for photosynthetic transcripts, plastid translation revealed as translationally upregulated transcript only chloroplast-encoded PSBA coding for the PSII subunit D1. No mitochondrial transcripts were detected as translationally upregulated. Interestingly, 36 and 16 DEGs of chloroplast and mitochondria representing 42.9 % and 34 % of detected plastidial transcripts, respectively, were decreased in translation (PPUT cluster). The translational downregulated transcripts were highly enriched in GO terms for translational processes (30 % in comparison to 0.6 % of genome representation) and ribosome assembly (11 % compared to 0.09 %) and photosynthetic light reaction (11 % to 0.11 %) represented by *YCF3* and *4* (ATCG00360, ATCG00520), PSII *CP47* reaction center protein (ATCG00680) and *NDH H* (ATCG01110). No enrichment in particular GO terms could be detected in the mitochondrial DEGs, but 6 of the 16 DEGs encoded for NADH dehydrogenases.

Translationally regulated transcripts are enriched in regulatory short mRNA motifs

Efficient reorganisation of translation requires unloading of ribosomes and reloading with preferably required transcripts. Translation initiation factors instigate translation in concert with mRNA regulatory sequences and/or structures. Often the 5'-UTR and 3'-UTR possess features that influence the initiation rate, like independent ribosomal entry sites (IRES). Therefore, the 5'- and 3'-UTRs of the clusters PPLT, PPUT and poly and total transcripts were screened for conserved motifs enriched against all TAIR annotated 5'-UTRs or 3'-UTRs using the MEME-suite (32). The search identified 40 motifs in the UTRs from which three were selected based on their overrepresentation in the PPLT (Cluster 1) and poly and total cluster (Cluster 3) (M1: TCTCCGGAGAA, M2: TTAGGGTT and M3: TCGCCG). M3 is a subsequence of M1 (Figure 2). M2 has previously been identified as stress-responsive 5'-UTR element that confers preferential translation in vitro (33). M1 and M2 were present 40- and 34-times in the 5'- and 3'-UTRs of the transcripts preferentially associating with ribosomes in H-light, accounting for 31% and 26%,

respectively. M2 is present in transcripts that are exclusively translationally regulated and not transcriptionally, whereas M1 and M3 are also present in transcriptionally regulated transcripts (Suppl. Fig. 2 c). Transcripts carrying M1 and M2 had a median relative translation rate of 2.09 and 1.97, respectively, and relative to the median translation rate of 2.96 of all PPLT transcripts. Transcripts carrying M3 had a higher relative translation rate of 3.57 (Suppl. Fig. 2 d). Heat-responsive transcripts like *HSFB2B* and *HSP20*-like proteins, as well as other stress-dependent transcripts, were overrepresented among the motif-carrying transcripts. Interestingly the UTRs of the *Stress Associated Protein 3* (*SAP3*) carries all three motifs and *SAP2* UTR carries M2 (Suppl. Fig. 2 a, b). SAPs belong to the family of zinc finger proteins that contain A20/AN1 zinc-binding domains. They are structurally and functionally conserved among diverse plant species (34). *SAP* transcript amounts increase in response to multiple stresses, and overexpression of specific SAPs in planta enhances biotic or abiotic stress tolerance or photosynthetic capacity (35, 36). A significant increase in ribosome loading after 10 min of L→H could be verified via qPCR for the transcripts of *HSP70*, *ERF105*, *DREB2A*, *SAP3* and *SAP2* with no significant increase in total mRNA for *SAP2* (Suppl. Fig. 1). SAP2 was highly enriched in the polysomal fraction, hence preferentially recruited to the polysomes. To assess importance of the motifs on an evolutionary scale we queried the rice genome and identified *OsSAP8* and *OsSAP9* as putative rice homologues of *SAP2* and *SAP3* and detected variations of all three motifs in their respective 5'-UTRs (Suppl. Fig. 4). Additionally, we observed the preferential association of transcripts for *OsSAP8* and *OsSAP9* with polysomes upon L→H transfer in rice (Suppl. Fig. 5).

Retrograde translational signaling is mediated by conserved motifs in the 5' UTR that serve as potential target for RNA binding proteins

Position analysis in the respective 5'-UTR of motif-carrying transcripts revealed a preferred position at the 3'-end in longer UTRs, while short UTRs showed no positional preference (Suppl. Fig. 2 b). Calculating the secondary structure via RNA Shapes Studio revealed that the motifs do not overlap with in any conserved secondary structure (Suppl. Fig. 2 b).

The hypothesized role in translational regulation of the three motifs was tested in a cell-based luciferase reporter assay. Cassettes carrying the M1, M2 or M3 motifs in triplicate repeats were inserted into the ubiquitin 5'-UTR in front of firefly luciferase reporter and the verified constructs transiently co-transfected together with pGL4::Renilla Luciferase as reference reporter into A. thaliana mesophyll protoplasts (Figure 2 b). The protoplasts were treated analogously to the whole

plant experiments with L→H-transfer for 10 min. The normalized luciferase activity from lysed
protoplasts transformed with the ubiquitin 5'-UTR without motifs increased 2.5-fold after L→H-
treatment. All motifs conveyed a higher signal than the ubiquitin control with M1-construct causing
the highest increase of 5-fold, 1.75-fold for M3 and insignificant for M2 (1.52-fold) (Figure 2 c).

The potential mRNP-binding properties of the motifs were tested via RNA-EMSA. Biotinylated
M3-RNA was incubated with leaf protein extracts that were either prepared from L-light-
acclimated plants or plants exposed to L→H transfer for 10 min (Suppl. Fig. 3 a). Addition of
protein extracts from L-light exposed leaves resulted in a pronounced shift of signals in comparison
to the biotinylated M3-RNA control. Addition of unlabeled competitor M3-RNA strongly
suppressed the shift. Supplementation with extract from 10 min L→H-treated plants resulted in a
signal intensification that again was out-competed by competitor M3-RNA. This result suggests
mRBP binding capability of M3 with differences in protein signature in polysomal fractions
between L-light plants and 10 min L→H-plants. Proteins interacting with M1 and M2 were
detected via affinity pulldown purification and subsequent SDS-PAGE revealed enrichment of
specific bands of which two were identified as GAPC2 and ATP synthase subunit alpha binding to
M2 and M1, respectively (Suppl. Fig 3 b-d).

Targeted RNA-EMSA with recombinant AtGAPC1 as RBP candidate showed a concentration
dependent band shift of free consensus M1- and M2-RNA (Suppl. Fig 3 e). To scrutinize these first
indications, interaction with the native M1 motif of the SAP3 5'UTR was used for affinity pull
down of potential RBPs with total protein extracts from L-light- or L→H-light-treated plants.
Immunodetection of AtGAPDH isoforms C1 and C2 revealed a strong signal in the elution fraction
derived from L-light leaf extract, whereas the signal in the H-light fraction was comparatively
weak. For reference, the GapC1/C2 content showed no difference in the input extract from L-light-
compared to H-light-plants. The same probe (SAP3 M1) was also used for RNA-EMSA with
recombinant AtGAPC1, C2 or GAPDH isolated from rabbit muscle to validate the interaction
between the motif-carrying RNA and GAPDH in vitro. Like above, adding increasing amounts of
protein led to a band shift of the free RNA. This shift was suppressed by adding unlabeled
competitor RNA.

Synchronization of translational signaling and nuclear transcription: CML49 interacts with SAPs to integrate translational, redox and calcium signaling

Rapid translational regulation of stress-responsive transcription factors and co-factors like MBF1c, HSF2A and DREB2A suggests integration of cytosolic translation in synchronizing plastid demand and nuclear transcription. These factors are known to be involved in transcriptional regulation of major stress response genes like HSPs and APX2 (37, 38). Other targets of translational control were the transcripts of *SAP2* and *SAP3*. Our previous studies revealed redox-dependent conformational changes of SAP12, another member of the SAP gene family (39). Analyzing SAP2 and SAP3 for disulfide bonds by DiANNA (40) revealed a predicted disulfide bond in the AN1 domains of SAP2 and SAP3 at Cys117 and Cys135. SAP2 compared to SAP3 has one additional cysteine at position Cys8 which is predicted to form a disulfide bond with Cys22. Thus, SAP2 and SAP3 might also function as redox sensory system in the translationally initiated retrograde signaling network. The preferential association of SAP2 and SAP3 mRNA with polysomes after L→H-transfer led us to search for interacting partners of these proteins. A yeast-two-hybrid screen with SAP2 fused to the GAL4-DNA-binding domain as bait recovered six clones from the prey consisting of Arabidopsis cDNA fused to the GAL4 activating domain, five of which coded for Calmodulin-like 49 (CML49). The interaction between CML49 and SAP2 and SAP3 was confirmed using the mating-based-Split-Ubiquitin (mbSUS) yeast-two-hybrid system (Figure 3 a) and by fluorescence resonance energy transfer (FRET) (Figure 3 b, c). SAP2 and CML49 and SAP3 and CML49 co-localized in the nucleus (Figure 3 b) and showed a high FRET efficiency (> 5-fold of negative control) proving their interaction in vivo (Figure 3 c). This interaction was also confirmed for the SAP2/3 and CML49 homologues from rice (Suppl. Fig 6), thus seems to be a conserved mechanism across multiple species.

The preferential association of *SAP2* and *SAP3* with polysomes after L→H-transfer and the interaction of SAP2 and SAP3 with CML49 led us to hypothesize that CML49 and SAP2/SAP3 function as regulatory module in this specific environmental context. In addition, the similar response of SAP2 and SAP3 in the L→H-response might indicate some functional redundancy. With these two data-guided hypotheses in mind, we hypothesized that the CML49 knockout should have stronger effects on downstream events during H-light acclimation than deletion of SAP3. To investigate the importance of SAP3 and CML49 in H-light acclimation, we determined the transcriptomes of *cml49*, *sap3* and WT in L-light and after 60 min L→H-transfer and calculated

323 the fold change as ratio of L→H/L (Suppl. Data Set 1). A set of 32 transcripts was more than 3-

324 fold increased or decreased in *sap3* relative to WT, while the corresponding set comprised 214

325 transcripts in *cml49*.

326 The data set was examined for transcripts that exhibited stronger downregulation in *cml49* than in

327 *sap3* upon L→H-transfer. This filter revealed 49 transcripts with the expected response pattern

328 (Figure 4 a). The predominant gene ontology groups belonged to the categories 'response to

329 temperature' and 'response to light stimulus', each of which represented 22 % of the genes,

330 followed by 'response to hydrogen peroxide' with 11 % and 'circadian rhythmicity' with 6 %. Heat

331 shock proteins and one heat shock factor (*HSFA2*) belonged to the group of heat responsive

332 elements, the *sigma factor 5* and *PSEUDO-RESPONSE REGULATOR 9* (*PRR9*) to the light

333 responsive elements *PURPLE ACID PHOSPHATASE 17* (*PAP17*) and the *GDP-L-galactose*

334 *phosphorylase VTC2* to the group of 'response to hydrogen peroxide'. *CIRCADIAN CLOCK*

335 *ASSOCIATED 1* (*CCA1*) and *PRR9* represent two genes associated with the circadian clock. The

336 occurrence of transporters for phosphate, sugar and sulphate (*PS3*, *AtSweet 13*, *GPT2*, *APR3*) in

337 this list indicates a function of CML49 and SAP3 in regulating metabolite and nutrient distribution.

338 Interestingly, the *translation elongation factor 1B* (*eEF1B*) was also found in this list possibly

339 pointing to SAP3/CML49-dependent amplification effects in the H-light dependent control of

340 translation.

341 The next experiments aimed to validate the deregulation of some of the identified transcripts and

342 to resolve the kinetics of their H-light induction (Figure 4 b-f). To this end, *sap3*, *cml49* and WT

343 plants were shifted from L-light to H-light for 10, 30 and 60 min. Plants left in L-light conditions

344 were used as control. Five targets were selected from the list of co-regulated transcripts for more

345 detailed analysis: *VTC2* showed a two-fold (log2) upregulation at t=10 in WT, followed by a

346 decrease to L-light levels at 30 and 60 min of L→H shift. In a converse manner, the response of

347 *sap3* and *cml49* displayed a steady decrease of transcript down to four-fold below the L-light level

348 (Figure 4 b-f). *HSP20*-like rapidly increased in WT upon L→H-transfer reaching its maximum of

349 six-fold (log2) induction at t=30 min and stayed constant until t=60 min. The mutants showed a

350 lower upregulation than WT with significantly reduced responses at t=10 and 30 min in *sap3* and

351 at 60 min in both mutants (Figure 4 b-f). The same trend appeared in WT for *PS3*, *SIG5* and *eEF1B*

352 with a two-fold upregulation after 10 until 60 min of L→H shift. The mutants showed a strong and

353 consistent deregulation compared to WT (Figure 4 b-f).

Common cis-elements in promotor region of co-regulated genes

The 49 co-regulated target genes were screened for common cis-elements in the promotor region in order to find commonalities such as involved transcription factors (TFs) needed for expression of the co-regulated gene set. A total of 9459 common cis-elements were identified in the 1000-bp upstream region of the 49 genes (Suppl. Table 2). Most abundant common cis-elements were the WBOX (2112), targeting sequences for the WRKY-TFs, followed by sulfur response (SURE) with 1536 and MYB and MYB-related with 1368 cis-element sequences. Stress related cis-elements like the sequences overrepresented in light-induced promotors (SORLIPs) and early responsive to dehydration (ABRE) were also highly abundant with 1278 and 1148, respectively. bZIP cis-elements (C-box, 232), phosphate starvation (PHR1, 224) and salicylic acid inducible (LS7, 173) were less abundant.

Is there a translational retrograde – anterograde circuit?

After 10 min L→H all plastidial translation is repressed with the sole exception of *PSBA*. This correlated with increased ROS within chloroplasts and subsequent inhibition of plastidial translation (16, 41). De-repression of plastidial transcription and translation during stress is mitigated via sigma factors like SIG5 which contributes towards transcription of the psbD operon (PSII D2) (42). Translational regulation of stress responsive transcription factors like HSF2A, DREB2A, and the SAP2/3 interaction with CML49 leads to translation-dependent nucleus synchronization after 10 min L→H. Plastid-targeted proteins like SIG5 and GPT2 and VTC2 were transcriptionally upregulated in CML49- and SAP3-dependent manner but not preferentially translated after 10 min L→H (Figure 4 b-e, Suppl. Dataset 1). Nonetheless, we hypothesize that the SAP-CML49 co-regulated expression of SIG5 contributes to de-repression of chloroplast genes, especially since PSII D2 transcription is 1.4-fold increased after 60 min of L→H.

Discussion

Regulation of actively translated transcripts described herein adds an entirely new mechanism to retrograde communication in the cell. We identified components of the novel pathway of translation-dependent retrograde signaling (TraDeRS) that bifurcates into direct translation of chloroplast-targeted proteins. This process is completely uncoupled from nuclear transcription (Figure 5 I.). In addition, retrograde control of cytosolic translation affects synthesis of a suite of factors that in turn promote nuclear transcription. Both mechanisms enable the acclimation process

384 and are part of an anterograde response (Figure 5 II.). Collectively, the suite of chloroplast-targeted

385 and chloroplast-encoded proteins controlled by translation-dependent retrograde signaling includes

386 those involved in protein import and folding, chloroplast transcription, photosynthesis and

387 photoprotection.

Evidence for translation-dependent retrograde signaling (TraDeRS)

Translation-dependent regulation repressed global translation, while increasing translation of

stress-specific proteins within minutes of L→H-transfer (Figure 1 a, b). Our work demonstrates

that this process is rapidly and reversibly controlled under excess light (Figure 1 a). Most other

immediate short-term responses to excess light are chloroplast-located like non photochemical

quenching (43), malate valve stimulation (44) or activation of antioxidant metabolism (45). In

contrast to the chloroplast-localized processes, TraDeRS takes place in the cytosol and potentially

integrates multiple cues. Previously, we demonstrated that rapid transcriptional responses in the

nucleus affect a network of transcription factors within 20 s after the L→H-transfer (16, 46). This

included a H_2O_2 burst peaking at 30 min after the L→H (16). The decreased polysome association

of mRNA displayed a reciprocal behavior with the H_2O_2 burst after the L→H shift (16). The

maximal H_2O_2 accumulation after 30 min is followed by a return to basal levels after 60 min (16).

Parallel to the decreasing H_2O_2-content, general ribosome association of mRNA recovered after

one hour of H-light (Figure 1 a) (33).

MPK6 and AKIN10 take part of translation-dependent retrograde signaling.

Ribosome unloading and preferential loading of transcripts are decisive regulatory steps in

translational control and are mediated by translation initiation, elongation and termination factors

(47). The question is how does excess light in the chloroplast modulate translation? Protein

phosphorylation, thiol redox-related changes and acetylation are reported posttranslational

modifications (PTM) of involved translation factors and regulate translation initiation, elongation

and termination (16-19). Formation of the translational pre-initiation complex (PIC) is inhibited by

the General Control Non Derepressable-2 Kinase (GCN2) (48), and its activation has been shown

in to be linked to MAPK signalling pathways (49). Consequently, we investigated the involvement

of MAP kinases and showed that mutants lacking MPK6 exhibit an altered polysomal profile. The

mutant failed to realize the preferential ribosome loading of two of the markers of increased

translation, *SAP2* and *SAP3*, upon L→H-transfer (Suppl. Fig. 1).

The second rate-limiting mechanism of translation initiation is via the eIF4F complex formation and subsequent mRNA binding regulated via 4EBPs. In mammals 4EBPs are activated by the mammalian Target Of Rapamycin (mTOR) Kinase (50) mediating phosphorylation of the eukaryotic Initiation Factor 4E (eIF4E), disabling interaction with initiation factor eIF4G, and thus limiting eIF4F formation (51). Phosphorylation of the translation initiation factors eIF4E and eIFiso4E via SnRK1, an interactor of TOR, inhibited translation in a plant system (52). Herein we show that the lack of AKIN10, a subunit of the SnRK1, altered the polysomal profile compared to WT under control and stress conditions and interfered with the preferential loading of the marker, *SAP2*, onto ribosomes as well. A future question is what activates this pathway? ROS/Redox regulation are candidates, as are GCN and TOR-signaling via AKIN10 and MPK6 (Suppl Fig. 1, Suppl. Table 1) and DHAP. Of interest in this regard, we have previously reported that export of DHAP from the chloroplast via the triosephosphate/phosphate translocator was required for rapid phosphorylation of MAP Kinase 6 (MPK6) after 10 min L→H-transfer. In summary, the results show that both MPK6 and AKIN10 are required for translation-dependent retrograde signaling, most likely via a phosphorylation-dependent pathway.

Proteins affected by bifurcating translation-dependent signaling pathways

The set of preferentially loaded transcripts could be grouped into purely translationally regulated and both, translation- and transcription-regulated transcripts, with each group representing distinct regulatory bifurcation of acclimation responses. Transcripts related to photosynthesis, heat stress and H_2O_2 dominated the set of preferentially polysome-loaded and motif-carrying transcripts. Photosynthesis and chloroplast repair-related transcripts were highly enriched among the purely translationally regulated transcripts (Suppl. Dataset 1, PPLT). The 119 transcripts in the translation-alone regulatory cluster are 5-fold enriched in chloroplast-localized proteins according to SUBA4 (31). This set includes proteins critical for assembly, function and maintenance of the chloroplast. Minor and major light harvesting proteins like LHB1B1 (AT2G34430), LHCB3 (AT5G54270) and LHCB5 (AT4G10340), LHCA4 as a subunit of photosystem I (AT3G47470), cytochrome b_6f (AT2G26500) and NADP oxidoreductase (AT1G06690) as well as one small subunit of RuBisCO (AT5G38420). Additionally, PSI assembly factor PYG7/YCF37 (AT2G23670) and import chaperones like HSC70-1 (AT5G02500) and chloroplast vesiculation factors (AT2G25625). Our results strongly suggest the existence of a transcription-independent

regulation of photosynthesis repair mechanism that is under rapid translation-dependent control

and forms a direct retrograde circuit (Figure 5, I. Direct Translational Circuit).

Complementary to photosystem repair, translational control of nuclear transcription factors (HSFs,

DREBs) synchronizes the gene expression with the cellular demand for subsequent stress

acclimation. The identified HSF2A, DREB2A and MBF1c play major roles in transcriptional

control of the H-light acclimation processes (53, 54). Interestingly, transcripts for these translation

factors are already present during LL-conditions but not preferentially translated (Suppl.DataSet

1). Preferential translation after stress perception might serve as first activator of these transcription

factors that are known to be up-regulated by oxidative stresses in general, and high light stress, in

particular.

Regulatory elements controlling polysomal loading

The regulation of specific mRNA translation efficiencies by targeting of different protein classes

for increased and decreased translation must be inherently based on their individual sequence or

structural properties. To identify conserved common sequences that might serve as such regulatory

property, we conducted *de novo* motif search for conserved sequences in the set of translationally

regulated transcripts via the MEME-tool suite to identify 42 motifs. In this study we could *in vivo*

demonstrate the impact of the three identified motifs during high light stress on translation and

provide evidence suggesting that they serve as mRNP binding sites including their interacting RBP

(Figure 2, Suppl. Fig. 3). RNA-EMSA shifts with leaf extracts from L- and L→H-light leaves

indicate that M3 acts as an mRNP binding site (Suppl. Fig. 3). Conditional mRNP-binding to motifs

1-3 stimulated polysomal loading and could be shown for motifs M1>>M3>M2 in a protoplast

transfection assay, the latter for M2 not being significantly different to the control (Figure 2). The

motif M2 (TAGGGTTT) has previously been identified in the 5'-UTRs of a set of preferentially

polysome-associated transcripts in young seedlings treated with light for 30 min, making it a

promising target for subsequent analysis. Translation of reporters from transcripts carrying the

motifs *in vitro* occurred at higher rates than from constructs lacking the sequence (33). The motif

was also overrepresented in preferentially polysome-associated transcripts under hypoxic stress

(55), suggesting a role for motif 2 under various stresses, but its mode of action is so far not

reported.

The nature of the RBPs that recognized the motifs M1 and M2 was investigated by RNA-based
affinity pull downs and protein identification via MS provided first candidate RBPs the ATP
synthase subunit alpha and GAPC2, respectively. GAPDH functions as non-canonical RBP in
animal systems (56, 57) and this role was only recently discussed in plants as well (58). We
validated the GAPDH-motif interaction using the native M1 of SAP3 5'-UTR via ex vivo affinity
pull down and subsequent immunodetection and in vitro by RNA-EMSA. These results point to
light-dependent protein-RNA interaction that possibly could mediate changes in TE upon L➔H
shift. As protein amounts of GAPDH (C1/C2) were unaltered in L-light- or L➔H-light-treated
plants conditional binding to RNA could be mediated by PTM or the availability of co-factors or
substrates. This would link translational control by RBPs to the cellular metabolic state enabling
the cell to rapidly adjust protein synthesis according to metabolic requirements. This hypothesis
needs further validation in detailed analysis.

SAP2 and SAP3 contain motifs 1 and 2 and their proteins interact with CML49

Motifs 1 and 2 are both present in the UTRs of *SAP2* and *SAP3* from Arabidopsis and also in their
homologues in the monocot rice. The transcripts in rice showed the same polysomal association
during L➔H as in Arabidopsis. Interestingly, re-analyzing a dataset of Arabidopsis seedlings re-
illuminated after darkness for 10 min (23) for translationally regulated transcripts in the same
manner as performed for this study, revealed *SAP2* as one of only 5 transcripts overlapping with
the purely translationally regulated subgroup, PPLT (Suppl. Dataset 1).

The Stress Associated Proteins (SAPs) belong to a zinc-finger multigene family with a high level
of structural conservation and functional similarity among several plant species (34, 35). Despite
little knowledge on molecular function two regulatory roles have been attributed to SAPs, namely
ubiquitin ligase activity and redox sensor activity regulating gene expression under abiotic stress
(35, 39, 59, 60). Previous work demonstrated that AtSAP12 undergoes reversible redox-dependent
conformational changes induced by cellular redox changes during stress (39). This underlines the
possible role of SAPs as redox sensors during H-light acclimation. Furthermore, SAP5 interacts
with MBF1c during heat stress regulating heat acclimation responses via interaction with HSFs
and DREB transcription factors, ultimately influencing transcription (61). Given the structural
similarity between SAP2/3 and SAP5, the interaction of SAPs with MBF1c might co-regulate
specificity and interaction between HSFs, DREBs and MBF1c to synchronize transcriptional

demand, but remains subject to further study. Thus, retrograde signaling increases the abundance

of the protein and in parallel promotes its re-location to promoted transcript accumulation.

SAP2 and SAP3 interact with the Calmodulin-like-protein-49 (CML49). Based on the highly

conserved AN1 domain present in all members of the SAP protein family, an interaction of all

SAPs with CML49 is possible. Several CMLs display an electrophoretic mobility shift in the

presence of Ca^{2+}, indicating that they act as Ca^{2+} sensors (62, 63). Most CMLs lack known

functional domain impeding the prediction of biochemical functions. Nonetheless, CMLs are

important in signaling processes and are activated under stress conditions like heat, cold, drought

and heavy metals (64, 65). The Ca^{2+}-dependent binding of CaM7 to a Z-/G-box is involved in

regulating expression of light-responsive genes (66). H_2O_2 triggers an increase in cytosolic Ca^{2+}

by activating Ca^{2+} channels (67). Ca^{2+}¬-liganded CaM activates MPK8 which suppresses ROS

accumulation via transcriptional control of RbohD (68). *CML49* occurs in almost all tissues and

was not induced after 1 h of L→H-shift indicating that CML49 functions as constitutive signal

transmitter. Interestingly, the rice homologues of SAP2 and SAP3 also interact with CML49

homologues indicating a conserved regulatory mechanism potentially integrating ROS cues with

Ca^{2+} signaling (Suppl. Fig. 6) across multiple plant species.

SAP3 and CML49 co-regulate transcript accumulation

Both *sap3* and *cml49* knockouts display smaller sets of H-light-induced transcripts compared to

wild type plants. We propose that CML49 acts as general interactor together with the SAPs as

stress-specific partners. First, CML49 is constitutively expressed and SAP2 and SAP3 are

preferentially translated during H-light acclimation with the interaction of SAPs and CML49

resulting in relocation of the complex in the nucleus. Second, of the 215 transcripts downregulated

in *cml49* plants after 60 min of L→H-transfer a stress subset of 32 are downregulated in *sap3*.

Thus, the stress-responsive protein, SAP3, is providing specificity to what would be otherwise a

broader CML49 response (Figure 4 a). Cumulatively, the preferential translation of the SAPs and

their interaction with CML49 but also other TFs like DREB2A and HSF2A form a translation-

reliant signal pathway affecting nucleus transcription (Figure 5, II Translation-Reliant Nuclear

Circuit).

This translation-reliant nuclear circuit regulates two major clusters of genes. As noted above, the

majority of the transcripts regulated by the proposed CML49-SAP3 module code for stress

responsive transcripts. A subset which includes cytosolic *HSP70*, *HOP3*, *ROF2*, *HSFA2*, *HSP90.1*,
HSP20-like that increase after 1 h of L→H shift (Figure 4 a). Interestingly, *HSP90.1* and *ROF1*
interaction is needed for HSFA2-dependent heat memory and also regulates short HSPs (sHSPs)
(69). Of interest in this respect is our earlier report with Joanne Chory and colleagues of HSFs
translocation to the nucleus being redox-regulated (70). Taken together, we propose that this
mechanism forms a feed-forward loop that can amplify the transcriptional responses over time, the
translation-reliant retrograde stress signaling pathway (Figure 5, II. Translation-reliant Nuclear
Circuit).

A second subset of transcripts co-regulated by CML49 and SAP3 takes part in an acclimatory and
anterograde signaling pathway that collectively could contribute to chloroplast acclimation.
Among the transcripts were VTC2 encoding the GDP-L-galactose phosphorylase which catalyzes
the rate limiting enzymatic step in ascorbate synthesis (71). With respect to anterograde signaling,
plastidic sigma factor 5 (SIG5) is utilized by plastome-encoded RNA-polymerase (PEP) to
transcribe the plastidial psbD gene coding for the D2 protein of the reaction center of photosystem
II (5), which is known to be induced by H-light (72). SIG5 and D2 transcripts were increased in
response to the L→H-shifts, and SIG5 was reduced in the *cml49* and *sap3* knockouts, providing
evidence for an anterograde circuit altering chloroplast transcription in response to translational-
dependent retrograde signaling.

Conclusion

This work reveals a translation-dependent retrograde signaling pathway and identifies major
participants across key stages that regulate rapid stress acclimation responses after L→H-light
transfer. The translatome reorganization initiated by L→H involves a phosphorylation cascade,
that includes MPK6 and the AKIN10 subunit of SnRK1, which inhibit global translation and
preferential transcripts loading to the polysomes. This may serve as a very efficient local control
mechanism and possibly outcompetes transcriptional control in terms of time requirement needed
to initiate acclimation by just in time-provision of needed proteins to the plastids. The preferentially
loaded transcripts directly code for repair and photosynthesis-related proteins forming a direct
translational retrograde circuit (Figure 5). Translational efficiency is influenced by conserved
motifs in their 5'-UTR acting as RBP binding sites. The motifs, GAPC2 and ATP synthase subunit
alpha function in the direct circuit. In parallel, a translation-reliant nuclear circuit is activated to
increase nuclear expression of stress- and plastid-associated transcripts. In addition to known

transcription factors, SAP2 and SAP3 increase in response L$\rightarrow$H and interact with the Ca^{2+} sensor

CML49 to promote accumulation of stress-responsive transcripts and anterograde signaling via

SIG5, which functions in plastid transcription. The elucidation of multiple components of a

bifurcating translation-dependent retrograde signaling pathway allows for rapid tuning of protein

synthesis to account for the demand of the photosynthesizing cell.

Materials and Methods

Plant growth and treatments. *Arabidopsis thaliana* wild-type (ecotype Columbia) and mutants

(*cml49-1* (SALK_090873), *cml49-2* (SALK_035905), *sap3-2 (SALK_022340), sap3-3*

(SALK_136520C), *sap2-1* (SALK_008677), *sap2-2* (SAIL_1182_C11), *mpk6-3* (*73*), and *akin 10*

(GABI_579E09) were grown under short day conditions (10 h light with 80-100 µmol photons m^{-2}

^{2} s^{-1}, and 21°C/ 14 h dark, and 18°C, 50 % relative humidity) for three weeks prior to transfer to

8 µmol photons m^{-2} s^{-1} (L-light) for 10 days. Plants were transferred to 800 µmol photons m^{-2} s^{-1}

(H-light) for time periods as indicated, while control plants remained in L-light. Whole rosettes

were harvested under control and H-light conditions in the midday point of the light period to

exclude circadian effects and immediately frozen in liquid nitrogen and stored at -80°C.

Sucrose cushion for polysome isolation. Total ribosomes and ribosome-bound mRNA were

extracted and separated using sucrose gradient centrifugation as described (*74*). 3 g of pulverized

plant material was used with 6 mL polysomal extraction buffer (PEB; 0.2 M Tris, pH 9.0, 0.2 M

KCl, 25 mM EGTA, 35 mM MgCl, 1 % Brij, 1 % Triton X-100, 1 % Tween 20, 1 % Igepal CA630,

1% sodium deoxycholate (DOC), 1 % polyethylene-10-tridecylether (PTE), 5 mM DTT, 1 mM

PMSF, 100 µg/mL cycloheximide, 100 µg/mL chloramphenicol, 100 µg/mL lincomycin). The

suspension was passed twice through Miracloth^{TM} and subsequently cleared by centrifugation.

Extracts were loaded on a sucrose cushion (0.2 M Tris, pH 9.0, 0.2 M KCl, 0.025 M EGTA,

0.035 M MgCl, 1.75 M sucrose, 5 mM DTT, 50 µg/mL cycloheximide, 50 µg/mL

chloramphenicol, 50 µg/mL lincomycin) and centrifuged for 18 h at 100,000 xg in a Beckmann

SW71Ti rotor. The sediment was suspended in 100 µL RNase digestion buffer (20 mM Tris-HCl,

pH 8.0, 140 mM KCl, 35 mM MgCl$_2$, 50 µg/µL cycloheximide, 50 µg/µL chloramphenicol,

50 µg/µL lincomycin) and used immediately for RNase I digestion.

Polysomal mRNA and protein extraction. Guanidinium chloride-precipitated polysomal

fractions were used for RNA isolation and subsequent cDNA synthesis as described in (*75*). For

protein isolation guanidinium chloride-precipitated polysomal fractions were pooled, proteins
extracted with pH-buffered phenol, precipitated with ethanol/NH$_4$ acetate and used for analysis in
denaturing polyacrylamide gel electrophoresis.

RNase I digestion. RNase I digest was performed with 2,000 relative absorbance units OD$_{260}$ of
the in RNase digestion buffer suspended polysomes by adding 50 U of RNase If (New England
Biolabs) in 250 µL total volume by incubation at room temperature with constant rotation for 1 h.
In parallel same amounts of polysomes were treated with 5 µL SUPERase RNase inhibitor
(Thermo) and kept on ice as control sample. The digestion was stopped by adding 5 µL SUPERase
RNase inhibitor and the RNA precipitated by adding 1 volume of isopropanol, 300 mM NaOAc
pH 5.2 and 20 µg glycogen at -80 over-night. The supernatant was discarded after centrifugation
at 16,000 xg at 4 °C for 30 min and the sediment was washed twice with 75 % ethanol and
resuspended in RNA loading buffer for size exclusion electrophoresis.

Library sequencing and analysis of mRNAs. Total RNA was isolated using Tri Reagent (Sigma-
Aldrich) from extracts suspended in polysomal extraction buffer (see sucrose cushion for polysome
isolation, above). Briefly, 500 µL of extract was combined with 1 mL of TriReagent, followed
twice by extraction of the organic phase with 200 µL chloroform. RNA was precipitated with equal
volume 100% isopropanol and incubated overnight at -20°C. The RNA was recovered by
centrifugation and washed with 75% ethanol before resuspension in water. RNA quality was
assessed using a LabChip GXII (Perkin-Elmer). Total RNA was DNase-treated using TURBO
DNase (Invitrogen) following the manufacturer's protocol. Preparation of total RNAseq libraries
was carried out using Illumina TruSeq Stranded Total RNA with Ribo-Zero plant, scaled at half
reaction volumes.

Libraries of ribosome-protected fragments (RPFs) were prepared following RNase If digestion
(above) according to Juntawong et al. (2014) with minor modifications. RNA linker, rRNA
subtraction, and PCR oligonucleotides were ordered from Integrated DNA Technologies (IDT) and
are listed in Suppl. Table 2. Oligonucleotides for rRNA subtraction were adapted from those listed
in (*29*) and (*76*)

Total RNA and RPF sequencing libraries were quantified using the Qubit fluorometer (Invitrogen)
and size distribution was assessed using the LabChip GXII. Samples were pooled and sequenced

(75 bp single end) on the NextSeq500 at the Biomolecular Resource Facility (BRF) at the
Australian National University.

Quality control of sequencing data was performed with FastQC (v.0.11.5). Total RNA data had
adapters removed using scythe (v.0.991) with a prior of 0.01, and quality trimmed with sickle
(v.1.33) with a quality threshold of 20 and a minimum trimmed length of 20. Mapping to TAIR10
was performed with subjunc (v.1.4.5), reporting only reads with a single unambiguous mapping
location (*77*). RPF data was trimmed for adapter contamination and read quality using cutadapt
(v.1.9.1) with a minimum trimmed length of 20 bp and error rate of 0.1 (*78*). Trimmed reads were
mapped to TAIR10 using bowtie2 (v2.2.9) requiring end-to-end alignment and a substring length
of 10 (*79*). Total and RPF-mapped reads were summarised using featureCounts (*80*). Counts data
was converted to reads per kilobase million (RPKM) in R using edgeR (*81, 82*)

TE calculation and transcript clustering. Translation efficiency (TE) calculation followed
published protocols (*83*) as quotient of RPKM from RPF and RPKM from total RNA sequencing
for control and stress condition. The TE fold change (TE FC) was calculated as $TE_{stress}/TE_{control}$.
Cut-offs for DEGs were chosen according to upper and lower quantiles from a random distribution
analysis of all regulated polysome-associated and total transcripts (total mRNA: 1.5 and 0.75 and
polysomal: 1.1 and 0.75; 0.05 error margin). Clusters were derived for significantly transcripts
changed up and down in both polysomal and total RNA (cluster 1 and 4). Cluster 2 and 3 were
selected by filtering for only significantly up- and down-regulated in the polysomal set and not
significantly changed in total RNA and were termed preferential polysome-loaded transcripts
(PPLT) and preferential polysome-unloaded transcripts (PPUT).

Whole genome ATH1 microarray. Total RNA was isolated from 1 h L→H-light treated *sap3*,
cml49 and WT *Arabidopsis* plants using RNeasy mini Kit (Qiagen). Controls were kept in L-light.
The hybridization of the Affymetrix ATH1 genome Array was commissioned to KFB (Regensburg,
Germany). Each condition was hybridized three times from independent experiments for *sap3* and
WT and once for *cml49*. Microarray data were analysed with ROBIN (MPI Golm, Potsdam,
Germany) and normalized with RMA algorithm (*84*). Transcripts were considered as differentially
regulated if the fold change was >2 and the adjusted *p*-value <0.05 (*85, 86*).

Transcript profiling. RNA isolation and cDNA synthesis were performed as in (*75*). Quantitative
real-time PCR (qPCR) of total and polysomal transcripts was conducted as described in (*28*) with

the difference of using KAPA SYBR® FAST qPCR Kit (Kapa Biosystems, South Africa). *Actin2*

(At3g18780) and *Tubulin5* (At1g20010) were used as reference genes.

RNA electrophoretic mobility-shift assay (RNA-EMSA). 5'-biotinylated RNA and unmodified

RNA of Motif 3 (5'-AAAUCGCCGGAG-3' 5'-BioTEG-AAAUCGCCGGAG-3') were

synthesized by Eurofins (Hamburg, Germany) and used for LightShift™ Chemiluminescent RNA

EMSA Kit (Thermo Scientific). Total protein was extracted from L-light and 10 min L➔H-treated

plants using extraction buffer (25 mM Tris, pH 7.8, 75 mM NaCl, 15 mM EGTA, 15 mM

glycerophosphate, 15 mM 4-nitrophenyl pyrophosphate, 10 mM $MgCl_2$, 1 mM DTT, 1 mM NaF,

0.5 mM NA_3VO_4, 0.5 mM PMSF, 10 µg/mL leupeptin, 10 µg/mL aprotinin, 0.1 % Tween-20).

Prior to separation by non-denaturing polyacrylamide gel electrophoresis, RNAs were incubated

with extracts at room temperature for 30 min. Biotinylated RNA was detected by

chemiluminescence with streptavidin-HRP. For targeted EMSA AtGAPC1 and C2 were expressed

in *E.coli* BL21 (DE3) pLysS and purified as described in (*87*) with slight modifications. Buffers

were not supplemented with NAD^+ and purified protein was stored in 10 mM NaPi, pH 7.4.

AtGAPC1, AtGAPC2 and GAPDH from rabbit muscle (SIGMA Aldrich) were used for mobility

shift assays with 5'-biotinylated M1-RNA (5'-BioTEG-AUAUAUUCUCCGGAGAA-3'), 5'

biotinylated M2-RNA (5'-BioTEG-AUAUAUGAGAUUAGGGUUU-3') and 5'-biotinylated

SAP3 M1 (5'-Bio-ACCAGAUUUUUUCGCCGGAGUUUGUUUUGAUCCA-3').

Reporter gene construct and transient protoplast assay. For the reporter gene construct assay

solely ubiquitin 10 5'-UTR as control (CCATGGCTTGATCACGGTAGAGAGAATTG–

AGAGAAAGTTTTTAAGATTTTGAGAAATTGAAATTCTCG) or ubiquitin 10 5'-UTR with

motifs (M1: CCATGGCTTGATCACGGTAGAGAGAATTGAGAGAAAGTTTTTTTC–

TCCGGAGATTCTCCGGAGATTCTCCGGAGAAAGATTTTGAGAAATTGAAATTCTCGA

G; M2: CCATGGCTTGATCACGGTAGAGAGAATTGAGAGAAAGTTTTTAAACCC–

TAATCTCAAACCCTAATCTCAAACCCTAATCTCAAGATTTTGAGAAATTGAAATTCTC

GAG; M3: CCATGGCTTGATCACGGTAGAGAGAATTGAGAGAAAGTTTTTCTCCGG–

CGACTCCGGCGACTCCGGCGAAAGATTTTGAGAAATTGAAATTCTCGAGA) were

cloned into MS129 luciferase reporter (*88*), and 5 µg plasmid DNA was co-transfected with same

amount of low-rate constitutively expressed pRL-TK *Renilla* luciferase vector (Promega) into

mesophyll protoplasts isolated from *A. thaliana* leaves by the downsized polyethylene glycol-

mediated transfection method (Seidel et al., 2004). After 12 h of acclimation half of the transfected

protoplasts were transferred to 800 µmol photons·m^{-2}·s^{-1} for 10 min and subsequently flash frozen

in N$_{2\ liq}$ for cell extracts. Luciferase activity of cell extracts was measured with Sirius L Tube

Luminometer (Berthold) and the luciferase assay system (Promega).

RBP pulldown and identification. Affinity pulldown with motif-carrying biotinylated RNA for

detection and identification of potential protein-RNA interactors was conducted as described in

(*89*) with minor differences. In short: 500 µg (or 250 µg for pull down followed by

immunodetection of cytosolic GAPDH) of total protein from L-light and L→H-treated plants were

incubated for 30 min at RT with 1 µg biotinylated M1- or M2-RNA (3 µg for 5'-Bio-*SAP3*-M1),

1 µL aprotinin (25 µg/µL), 1 µL leupeptin (25 µg/µL), 2 µL RNasin (Promega), 250 µL TENT

buffer (0.02 M Tris-HCl, pH 8.0), 0.002 M EDTA (pH 8.0), 0.5 M NaCl, 1 % (v/v) Triton X-100)

and adjusted to 500 µL with TENT buffer. 50 µL Dynabeads (Thermo Fisher) were added and

incubated under slow shaking for 30 min at RT. After precipitation and three washing steps with

TENT buffer, Dynabeads were dissolved in 1x protein loading buffer and incubated for 10 min at

95 °C, precipitated and the supernatant loaded on SDS-PAGE for separation and band

identification via mass-spectrometry or immunodetection of GAPC1/C2 after blotting the gel-

separated proteins on nitrocellulose membrane by semi dry electro blotting technique.

Promotor and uORF analysis. The +50 to -1000-bp region upstream of the ATG start codon and

+500 downstream of the transcription stop site of the CML49 and SAP3 co-regulated genes were

analysed for common *cis*-elements using PlantPAN2.0 (*90*). For uORF analysis the web tool

uPEPperoni was used (*91*) and the eFP Browser 2.0 for general expression pattern (*92*).

mbSUS-analysis. Vector construction of the mating-based Split-Ubiquitin-System analysis was

performed with the Gateway® cloning system (InvitrogenTM). The coding sequences of SAP2,

SAP3 and CML49 were fused to attB recombination sites and cloned into the pDONRTM221-

vector by a BP-reaction. Subsequently, each entry clone was used in a LR-reaction with the

destination vectors pX-NubWTgate (Nub) and pMetYCgate (Cub). Controls were the transformed

diploid yeast with NubI- (pNubWtX-gate, positive control) and the NubG-vector (pNX35-Dest,

negative control) as prey. The resulting vectors were transformed via lithium acetate transformation

into *Saccharomyces cerevisiae* strains THY.AP5 and THY.AP4, respectively (*93*). Yeast cells were

grown in 20 mL selective Nub/Cub medium at 30°C/140 rpm for 5-7 days. To start the mating step,

THY.AP5 and THY.AP4 were transferred to YPD plates and grown for 1 d at 30°C. For the

methionine assays, mated yeast was cultivated in selective liquid diploid medium for 3 d (30°C,

140 rpm). OD600 of selected yeast cells were measured and diluted to OD600 of 0.1 and 1.0

respectively in sterile water. Then 6 µL drops of each dilution were transferred to agar plates

containing no (Met 0 µM) or 0.075 % (v/v) methionine (Met 50 µM). Subsequently, the yeast was

grown for 72 h at 30°C and results documented photographically (*93, 94*).

Subcellular localization and FRET measurements. The coding sequences of the genes of interest

were fused in frame to yellow or cyan fluorescent protein as indicated and fluorescence resonance

energy transfer (FRET)-experiments were performed as described by (*95*) with CLSM 780 (Zeiss).

More than 20 protoplasts were measured twice for each set of PEG-based transformation and the

experiment was repeated independently

Acknowledgements

The authors acknowledge support by the Deutsche Forschungsgemeinschaft (Di 346/19; SP1935) and especially thank Diep Ganguly for discussion of statistics and data analysis and Renate Scheibe for providing the pET16b construct for recombinant expression of GapC1/C2.

The authors declare that they have no competing interests.

Data availability

All relevant data is accessible in this manuscript or in the supplemental material.

Authors contributions

MM designed the research and performed the polysomal gradients, transcript profiling, microarrays, RPF-seq and total RNAseq bioinformatic data analysis, transient protoplast reporter gene assays, RNA-EMSA, protein affinity pull downs and wrote the paper. AS did the RPF-library preparation and sequencing and performed bioinformatic analyses. CW prepared polysomal gradients and transcript profiling. SS performed Y2H experiments and cellular localization and FRET. MW performed RBP affinity pull downs and gel shift assays on SAP3 motifs. AF measured FRET and transcript profiles of OsSAP and OsCML49. TS supervised and performed FRET measurements. BP supervised and discussed the results and wrote the paper. KJD designed the study, supervised and discussed the results, coordinated the group and wrote the paper.

Figure legends

Figure 1: Ribosome loading and ribosomal profiles. (a) Lowlight (L)-acclimated plants were maintained in L-light (0') or transferred to Highlight (H, 10') and then transferred back to L for the indicated time (recovery). Ribosome-loading efficiency was determined by comparing mRNA quantity in ribosomal footprints to total mRNA (mean ± SE from n=3 independent experiments). Comparison of RPKM of 10 min L→H relative to L-light control. Letters indicate significance of difference as determined with ANOVA post-hoc Tukey $p<0.5$. (b) Comparison of translational efficiency (TE) of individual mRNAs under LL and HL. Also indicated are the numbers of transcripts with significantly higher or lower TE than average between LL and 10 min HL. (c) Change in total mRNA and translation efficiency for same genes as in (b). Quantiles were defined by random distribution analysis and set as cut-offs (TE: 1.41 and 0.74; total mRNA: 1.5 and 0.75; 0.05 error margin) for regulation of translation vs. transcription of all significant RPKM reads (FDR significance of $p<0.05$). (d) Significant changes in polysomal and total mRNA given as fold changes under LL and HL (cut-offs poly: 1.1 and 0.75; total mRNA 1.75 and 0.7; 0.05 error margin; FDR significance of $p<0.05$) with clustering into groups by polysomal and not significantly total mRNA changed (PPLT; red and PPUT; steel blue), polysomal and total mRNA changed (Poly and Total; blue) and only total changed without significantly changed polysomal transcripts (buffered; salmon).

Figure 2: Identification and functional validation of three motifs in the 5'UTR of preferentially polysome-loaded transcripts. These motifs are hypothesized to confer preferential translation after 10 min L→H. (a) In 338 5'-UTRs of the preferentially loaded transcripts specific motifs were detected using the MEME algorithm against a randomized 5'-UTR set. Motifs 1 and 3 were present in 32% of those transcripts, and motif 2 in 13% with an additional 8% occurrence in their respective 3' UTR. (b) Schematics of constructs used in the reporter assays. The 5'-UTR of ubiquitin 10 was cloned upstream of the luciferase (LUC) reporter under control of the ubiquitin promotor. Triplet motifs were introduced into the 5'-UTR of ubiquitin mRNA. *A. thaliana* mesophyll protoplasts were co-transfected with one of each construct and with the pGL4::RLUC reference construct. (c) Relative LUC signal in lysed mesophyll protoplasts treated with H-light for 10' normalized to reference construct. The construct containing M1 showed a strong increase in LUC signal after L→H. Data are means ± SD from two independent experiments with 10 replicates, * and + indicate significant difference to ubiquitin control, two sided Student's t-test: $p<0.05$ and 0.1 , respectively.

Figure 3: Interaction of SAP2 or SAP3 with CML49 in the split-ubiquitin yeast-two-hybrid system and by Förster resonance energy transfer (FRET) of SAP2:CFP or SAP3:CFP with CML49:YFP in *A. thaliana* protoplasts. (a) Quantification of interaction strength between SAP2 or SAP3 and CML49 using a methionine-dependent split-ubiquitin-assay. Depicted are colonies after 72 h at 30 °C in different ODs (OD = 0.1 and 1.0). Growth was monitored either on Met 0 µM (no methionine, left colonies) and Met 50 µM (50 µM methionine, right colonies) medium plates. Diploid yeast was transformed with NubI- (pNubWtX-gate) as positive control and as negative control with the NubG-vector (pNX35-Dest) as prey (depicted are representative images of n=3 independent experiments). (b) *A. thaliana* mesophyll protoplasts were transfected with SAP2:CFP, SAP3:CFP or CML49:YFP vector DNA. Fluorescence images taken by CLSM for subcellular localization. Following transfection with SAP2:CFP or SAP3:CFP and CML49:YFP constructs, FRET efficiencies between CFP and YFP were quantified. FRET was also determined with CFP:YFP as positive control and 35S::CFP with 35S::YFP as negative control (for each set of transformation > 20 protoplast were measured twice with n=3 independent experiments).

Figure 4. Deregulated H-light response in *sap3* and *cml49* compared to WT. (a) Microarrays were hybridized with cDNA from WT, *cml49* and *sap3* after 60 min of L→H transfer. Assuming a concerted function of SAP2 and SAP3 together with CML49 in the H-light response, the data were filtered for co-regulated targets that are stronger deregulated in *cml49* than *sap3* background. The expression level corresponds to the colour intensity with yellow = -1 and blue = 5 ($\log_2$ fold). (b-f) Validation of transcript responses to H-light of putative CML49 and SAP3 targets. VTC2, HSP20-like, PS3, SIG5 and eEF1B were selected based on the transcriptome response, and their transcript levels quantified by qPCR following the L→H-transfer. Data are means ± S.D. of n=4 independent experiments, asterisks indicate significant difference, two sided Student's t-test p<0.05.

Figure 5: Schematics of translation-dependent retrograde signalling (TraDeRS). Rapid regulation of polysome loading and unloading after 10 min of L→H shift was partially disturbed in *mpk6* and *akin10* mutants indicating retrograde regulation of translation. Regulation involves transcription-independent changes of translational efficiency of majorly chloroplast targeted proteins involved in photosynthesis as well as chloroplast-associated assembly, repair and import of proteins, thus forming a direct feedback circuit regulated via translation (I. Direct Translational Circuit). The preferential loading of key transcription factors as well as *SAP2* and *SAP3* are part of the translation-reliant synchronization of the nucleus with the changed chloroplast demand (II. Translation-Reliant Nuclear Circuit). SAP2 and SAP3 interact with CML49 and this regulatory module affects transcript abundance of, e.g. VTC2 and SIG5, participating in anterograde signalling, and other major stress response genes like ANAC2, HSP101 and HOP3.

Supplementary materials

Suppl. Fig. 1: Preferential loading of selected transcripts after 10 min L➔H transfer and deregulation in *akin10*. (a-e) Five transcripts were selected from the transcriptome data set and subjected to qPCR analysis in wildtype (WT) and *akin10*. SAP2 and SAP3 showed significantly higher polysome association after L➔H-transfer for 10 min. (Data are means ± SE of n=3 independent experiments, * indicate significance difference relative to control, two sided Student's t-test: p <0.05, 0.1, respectively). (f-h) Absorbance (A254 nm) of sucrose density gradient of WT, *mpk6* and *akin10* treated with L-Light (control) and L➔H transfer for 10 min (depicted are representative profiles of n=3 replicates).

Suppl. Fig. 2: Distribution and secondary structure of motif-carrying 5'UTRs. (a) Motif 1, 2 and 3 occurrences in their respective transcripts as revealed by MEME-suit. (b) The secondary structure is given in Dot-Bracket notation and was calculated using the software *RNA Shapes Studio (96)*, with Motifs drawn at corresponding site(s) for motif 1 (green), 2 (yellow) and 3 (red), respectively. (c) Distribution of motif-carrying transcripts in translatomes are indicated as dots for transcripts carrying motif 1 (green), 2 (yellow) or 3 (red). (d) Relative ribosomal loading (HL/LL) of motif-carrying transcripts (M1 – M3) and all translationally regulated transcripts with (All Transcripts) and without motifs (All Other Transcripts).

Suppl. Fig. 3: Motifs are possible RBP binding sites. (a) 5'-biotinylated M3-RNA with and without unlabelled competitor M3-RNA was incubated with total protein extracts of L-light or 10 min L➔H shifted plants, respectively. Depicted is a representative blot. (b,c) mRNP 1-D acrylamide gel. Biotinylated M1 and M2 RNAs were coupled to strep-tagged beads and incubated with protein extracts from LL and 10 min L➔H treated plants. After washing, samples were eluted and loaded on one dimensional acrylamide gels. The indicated bands were cut out and analysed via MALDI. (d) Best Mascot scores for indicated bands from b & c. (e) RNA-EMSA was used to validate interaction of M1- and M2-RNA with Arabidopsis cytosolic GAPDH isoform C1 *in vitro*. (f) Immunodetection of AtGapC1/C2 after RNA-based affinity pull down with native M1 from *SAP3* 5'-UTR as described for M1 and M2 in b & c. Elution: eluted proteins associated with the RNA-functionalized beads, FT: Flow through, supernatant after affinity pull down, Input: Plant extract prior to affinity pull down. (g-i) RNA-EMSA performed with 5'-Bio *SAP3*-M1 and AtGapC2 (g), AtGapC1 (h) or GAPDH isolated from rabbit muscle (i).

Suppl. Fig. 4: Representation of motifs 1-3 from *A. thaliana* as Hidden Markow Models and comparison with related motifs in rice SAP8 and SAP9. OsSAP8 and OsSAP9 were identified as closest homologues of SAP2 and SAP3 from *A. thaliana*. Each motif could be identified in the 5'-and 3'-UTR of the rice homologues with slight variation.

Suppl. Fig. 5: Gene expression map and polysome loading of *OsSAP8, OsSAP9* and *OsCML49a*. (a) Transcript abundance of *SAP8*, (b) *SAP9* and (c) *CML49a* in different rice tissues retrieved from the Rice eFP Browser. *CML49b* was missing in group 1 data source of Rice eFP Browser. (d,e) qPCR-based quantification of *OsSAP8* and *OsSAP9* transcripts in total and polysomal RNA fractions after 10 min of L→H-transfer. Data sets were normalized to *Actin* and *α-Tubulin* and represent means ± SD of n=2 independent experiments, asterisks indicate significance relative to control, two sided Student's t-test: p <0.05.

Suppl. Fig. 6: Subcellular localization of rice OsSAP8, OsSAP9, OsCML49a and OsSAP49b in transfected Arabidopsis protoplasts and their molecular interaction shown by FRET. *A. thaliana* mesophyll protoplasts were transfected with OsSA8:YFP, OsSAP3:YFP, OsCML49b:CFP or OsCML49a:CFP vector DNA. Fluorescence images taken by CLSM for subcellular localization. Following transfection with constructs, FRET efficiencies between CFP and YFP were quantified.

Suppl. Table 1. Translation efficiency (TE) of selected transcripts in *mpk6* and *akin10* background. Target transcripts were quantified in total and polysomal mRNA via qPCR (Suppl. Fig. 1.). Two transcripts were selected for their increase in TE (*SAP2* and *SAP3*) and three for their decrease in TE (*ERF105*, *DREB2A*, *HSP70*). (TE was calculated from n=3 independent experiments, asterisks indicates significant difference p<0.05; ANOVA)

TE	TE increased		TE decreased		
Name	SAP2	SAP3	ERF105	DREB2A	HSP70
WT	6.1	2.6	0.4	0.2	0.07
mpk6	1.2*	1	0.3	0.4	0.26*
akin10	1.8*	0.6	1.1*	1.2*	0.58*

Suppl. Table 2. Oligonucleotides.

Target gene	ID number	Forward Sequence (5'–3')	Reverse Sequence (5' – 3')
HSP20-like	AT1G53540	GTTCGAAGGTTTCTTGACGC	TGTCTCCCTCCAATCCACTT
HSP70	AT3G12580	TCGACTGCTCAGACGACTAT	GAACCTAGCACGAGTGATGG
SAP2	AT1G51200	TGAGCCGATGACTGTTTCTG	TTGGTTTCGCAGCTTCTTCT
SAP3	AT2G27580	GGTGTGGTTTCGTTTTCTGC	GCTTTCACAATCGGATTCGC
VTC2	AT4G26850	GACTGCTGTGTTTGCCTTCA	CGGGCTCACTTCACCTAGAG
eEF1B	AT5G12110	ATCTCCACACAGAGCGAGGT	TTGCTAGCATTGGGAAAACC
SIG5	AT5G24120	TCCTCCTCGTGAGCAAGTTT	GGAGCCGAATTATGTTCCAA
PS3	AT3G47420	TTGGTTAACGGTCCTTACGC	CGTTACCGTAGCCAAAGCTC
OsSAP8	Os06G41010	CGATGTGGCAACTTGTACTG	TTAGCCTTGGCAATAGCATC
OsSAP9	Os07G07350	AAACAACTGTGGCTTCTTCG	CGTAAATGCCTTCTTCTCCA
OsActin	Os03G07181	AGGCCGTCCTCTCTCTGTAT	GGGGAGAGCATATCCTTCAT
Osα-tubulin	Os03G51600	TCTGAACGTGGATGTCAATG	GAGATCACTGGAGCGTAGGA
OsSAP8NotI		AAAAGCGGCCGCATGGAGCACAAGGAGACT	
OsSAP8EcoRI		TTTTGAATTCCTAAATTTTGTCAAGTTT	
OsSAP9NotI		AAAAGCGGCCGCATGGCACAAGAAAGCTGG	
OsSAP9EcoRI		TTTTGAATTCTCAGATCTTGTTGATCTT	
OsCML49NotI		AAAAGCGGCCGCATGGCCGGCTACCCGCCA	
OsCML49EcoRI		TTTTGAATTCTTATGCAATGAGAAAAGG	
rRNA depletion	rRNA1b	CAGCTGAGGCATCCTAACAGACCGGTAGACTTG	
rRNA depletion	rRNA21	AGTAAGCCCACCCCAAGATGAGTGCTCTC	
rRNA depletion	rRNA27	ACGACTCTCGGCAACGGATATCTCGGCTCTCGCATCGATGAAGAACG	
rRNA depletion	rRNA29	GGCCGAGGGCACGTCTGCCTGGGTGTCAC	
rRNA depletion	rRNA30	GACTGTGAAACTGCGAATGGCTCATTAA	
rRNA depletion	rRNA44	TAGCGACGTTGCTTTTTGATCCTTCG	
rRNA depletion	rRNA1c	GATAACCGTAGTAATTCTAGAG	
rRNA depletion	rRNA1	CATAAACGATGCCGACCAGGGATCAGCGG	
rRNA depletion	rRNA2	TTTATTAGATAAAAGGTCGACG	

Suppl. Table 3: Overview of *cis*-elements within the promoter regions of CML49 and SAP3 co-regulated genes.
The total number and type of *cis*-elements within the co-regulated subgroup was searched -1.000 bp upstream of the ATG region using PlantPAN2.0. The largest group of *cis*-elements belong to the WRKY-group, followed by MYB, MYB-related, SURE (sulfur-responsive element) and ABRE-like (early responsive to dehydration). Only few putative sites belong to the LS7 (salicylic acid-inducible element), C-box (bZIP factors), PHR1 (phosphate starvation response) and some represent cis-elements for light-dependent SORLIPs (Sequences Over-Represented in Light-Induced Promoters).

AGI	Name	MYB/ MYB-related	WBOX	SURE	PHR1	C-box	ABRElike	LS7	SORLIPs	Other
AT4G13800	unknown	29	37	35	4	2	20	4	17	24
AT5G52570	BETA-OHASE 2	24	36	29	6	2	27	4	30	26
AT5G49480	CP1	24	41	30	1	2	16	1	19	21
AT4G04020	FIB	22	39	18	5	3	19	3	26	16
AT5G58770	CPT4	35	35	22	3		23	3	30	29
AT3G56290	unknown	23	47	31	3	2	25	3	35	34
AT1G67360	REF	41	46	32	5	8	30	6	27	14
AT1G61800	GPT2	28	36	23	7	9	18	3	17	26
AT5G20150	SPX1	22	42	20	13	4	14	1	10	23
AT3G21890	BBX31	22	41	27	2	6	22	3	14	30
AT2G46830	CCA1	31	43	47	1	6	21	1	44	14
AT5G52640	HSP90.1	35	38	43	4	4	23	3	15	14
AT2G36800	DOGT1	32	48	44	5	6	28	3	30	14
AT5G48570	ROF2	24	49	30	2	4	22	1	15	14
AT1G52690	LEA7	27	44	27	5	8	32	4	28	25
AT2G24540	AFR	28	45	41	6	9	28	4	28	31
AT3G09350	Fes1A	34	41	40	3	9	26	9	25	21
AT4G21990	APR3	22	40	26	3	6	21	1	16	20
AT4G12400	HOP3	33	37	25	3	1	24	6	12	21
AT5G50800	ATSWEET13	26	43	32	4	7	24	5	30	19
AT2G37180	RD28	23	43	21	7	4	22	3	20	34
AT4G34550	PCH1-like	26	35	30	4	3	16	3	19	34
AT3G17609	HYH	27	38	27	6	5	24	3	28	30
AT5G64510	TIN1	34	44	34	4	12	27	9	30	30
AT1G01520	ASG4	33	34	45	3	2	29	7	38	26
AT1G32900	GBSS1	31	52	31	5	9	28	2	36	30
AT2G26150	HSFA2	20	60	46	0	4	19	3	30	34
AT5G12110	eEF1B	41	53	34	2	6	17	4	32	31
AT4G26850	VTC2	20	40	31	8	8	23	4	20	36
AT2G33380	RD20	29	42	26	6	7	38	3	22	29
AT3G21150	BBX32	26	41	39	5	6	17	5	30	23
AT3G09450	unknown	38	34	42	6	1	29	3	43	29
AT2G21320	BBX18	26	57	40	7	4	20	1	20	27
AT1G58520	RXW8	20	28	26	2	3	17	4	18	45
AT5G24120	SIG5	14	52	31	1	4	22	1	16	25
AT5G65380	MATE	24	47	31	4	8	32	7	33	29
AT3G53830	RCC1	24	40	35	6	4	31	2	18	27
AT3G17790	PAP17	37	44	34	7	8	28	6	31	29
AT1G54050	HSP20-like	44	40	30	6	5	15	3	38	31
AT1G62180	ATAPR2	29	53	24	5	3	26	3	27	28
AT3G21670	ATNPF6.4	22	44	27	3	1	25	2	21	37
AT3G12580	HSP70	20	42	24	4	1	18	2	11	32
AT2G46790	PRR9	22	43	26	6	3	19	1	44	33
AT1G77450	NAC032	31	45	35	4	6	33	8	35	30
AT2G15020	unknown	20	34	26	5	0	24	3	34	36
AT3G47420	PS3	30	42	10	7	2	21	5	18	27
AT5G63130	Phox/Bem1p	20	49	19	4	5	15	3	31	37
AT1G01720	ANAC2	29	39	50	2	6	22	2	36	23
AT4G26150	CGA1	26	34	27	5	2	15	2	14	36
AT1G74310	HSP101	20	35	33	5	2	13	1	17	34
	Total	1368	2112	1556	224	232	1148	173	1278	1368

Suppl. Table 4. uORF analysis of motif-carrying transcripts. The motif-carrying transcripts were screened for upstream open reading frames (uORF) coding for upstream peptides (uPEPs) using the online tool uPEPperoni.

Target gene	ID number	uPEPs Sequence (5'–3')	Position (start-end)
PHT3	AT2G17270	MRTLPTYTLFAFPFIAFL	(182, 235)
Methyltransferase superfamily protein	AT3G13440	MGLGRNLQTIRSSYKIRVF	(23, 79)
PDX1.2	AT3G16050	MEAHLVSFFCIRRRFAIPP	(14, 70)
RBL14	AT3G17611	MRVKRRSLWRICFRRDQAAAI	(8, 70)
Unknown protein	AT4G27010	MPDFELSASPICFVLLNPKTKTPRREKSSL	(53, 142)
HSFB1	AT4G36990	MEETKRNSDLLRSRVFLSGFYCWDWE	(267, 344)
SIG5	AT5G4088	MMMPSSSCFHFGLRFFISSIRDPPIRIDTQRSRVSTDMDMDKS	(159, 287)

References:

1. J. Barber, B. Andersson, Too much of a good thing: light can be bad for photosynthesis. *Trends in Biochemical Sciences* **17**, 61-66 (1992).
2. K. K. Niyogi, Safety valves for photosynthesis. *Current Opinion in Plant Biology* **3**, 455-460 (2000).
3. M. Baier, K. J. Dietz, Chloroplasts as source and target of cellular redox regulation: a discussion on chloroplast redox signals in the context of plant physiology. *J Exp Bot* **56**, 1449-1462 (2005).
4. K. Asada, THE WATER-WATER CYCLE IN CHLOROPLASTS: Scavenging of Active Oxygens and Dissipation of Excess Photons. *Annu Rev Plant Physiol Plant Mol Biol* **50**, 601-639 (1999).
5. A. Nagashima, The Multiple-Stress Responsive Plastid Sigma Factor, SIG5, Directs Activation of the psbD Blue Light-Responsive Promoter (BLRP) in Arabidopsis thaliana. *Plant and Cell Physiology* **45**, 357-368 (2004).
6. B. J. Pogson, N. S. Woo, B. Forster, I. D. Small, Plastid signalling to the nucleus and beyond. *Trends Plant Sci* **13**, 602-609 (2008).
7. J. Pfalz, T. Pfannschmidt, Essential nucleoid proteins in early chloroplast development. *Trends Plant Sci* **18**, 186-194 (2013).
8. T. Pfannschmidt, A. Nilsson, J. F. Allen, Photosynthetic control of chloroplast gene expression. *Nature* **397**, 625-628 (1999).
9. C. F. Beck, Signaling pathways from the chloroplast to the nucleus. *Planta* **222**, 743-756 (2005).
10. S. Koussevitzky, A. Nott, T. C. Mockler, F. Hong, G. Sachetto-Martins, M. Surpin, J. Lim, R. Mittler, J. Chory, Signals from chloroplasts converge to regulate nuclear gene expression. *Science* **316**, 715-719 (2007).
11. G. M. Estavillo, P. A. Crisp, W. Pornsiriwong, M. Wirtz, D. Collinge, C. Carrie, E. Giraud, J. Whelan, P. David, H. Javot, C. Brearley, R. Hell, E. Marin, B. J. Pogson, Evidence for a SAL1-PAP chloroplast retrograde pathway that functions in drought and high light signaling in Arabidopsis. *Plant Cell* **23**, 3992-4012 (2011).
12. Y. Xiao, T. Savchenko, E. E. Baidoo, W. E. Chehab, D. M. Hayden, V. Tolstikov, J. A. Corwin, D. J. Kliebenstein, J. D. Keasling, K. Dehesh, Retrograde signaling by the plastidial metabolite MEcPP regulates expression of nuclear stress-response genes. *Cell* **149**, 1525-1535 (2012).
13. S. W. Park, W. Li, A. Viehhauser, B. He, S. Kim, A. K. Nilsson, M. X. Andersson, J. D. Kittle, M. M. Ambavaram, S. Luan, A. R. Esker, D. Tholl, D. Cimini, M. Ellerstrom, G. Coaker, T. K. Mitchell, A. Pereira, K. J. Dietz, C. B. Lawrence, Cyclophilin 20-3 relays a 12-oxo-phytodienoic acid signal during stress responsive regulation of cellular redox homeostasis. *Proc Natl Acad Sci U S A* **110**, 9559-9564 (2013).
14. K. J. Dietz, Redox regulation of transcription factors in plant stress acclimation and development. *Antioxid Redox Signal* **21**, 1356-1372 (2014).
15. R. E. Häusler, L. Heinrichs, J. Schmitz, U. I. Flugge, How sugars might coordinate chloroplast and nuclear gene expression during acclimation to high light intensities. *Mol Plant* **7**, 1121-1137 (2014).
16. M. O. Vogel, M. Moore, K. Konig, P. Pecher, K. Alsharafa, J. Lee, K. J. Dietz, Fast retrograde signaling in response to high light involves metabolite export, MITOGEN-

ACTIVATED PROTEIN KINASE6, and AP2/ERF transcription factors in Arabidopsis. *Plant Cell* **26**, 1151-1165 (2014).

17. J. Schmitz, L. Heinrichs, F. Scossa, A. R. Fernie, M. L. Oelze, K. J. Dietz, M. Rothbart, B. Grimm, U. I. Flugge, R. E. Hausler, The essential role of sugar metabolism in the acclimation response of Arabidopsis thaliana to high light intensities. *Journal of Experimental Botany* **65**, 1619-1636 (2014).

18. P. A. Crisp, D. Ganguly, A. B. Smith, K. D. Murray, G. M. Estavillo, I. R. Searle, E. Ford, O. Bogdanovic, R. Lister, J. O. Borevitz, S. R. Eichten, B. J. Pogson, Rapid recovery gene downregulation during excess-light stress and recovery in Arabidopsis. *Plant Cell*, (2017).

19. G. Z. Wu, E. H. Meyer, A. S. Richter, M. Schuster, Q. Ling, M. A. Schottler, D. Walther, R. Zoschke, B. Grimm, R. P. Jarvis, R. Bock, Control of retrograde signalling by protein import and cytosolic folding stress. *Nat Plants* **5**, 525-538 (2019).

20. M. Moore, N. Gossmann, K. J. Dietz, Redox Regulation of Cytosolic Translation in Plants. *Trends Plant Sci* **21**, 388-397 (2016).

21. A. Lokdarshi, J. Guan, R. A. Urquidi-Camacho, S. K. Cho, P. W. Morgan, M. Leonard, M. Shimono, B. Day, A. G. von Arnim, Light Activates the Translational Regulatory Kinase GCN2 via Reactive Oxygen Species Emanating from the Chloroplast. *Plant Cell*, (2020).

22. M. Piques, W. X. Schulze, M. Hohne, B. Usadel, Y. Gibon, J. Rohwer, M. Stitt, Ribosome and transcript copy numbers, polysome occupancy and enzyme dynamics in Arabidopsis. *Mol Syst Biol* **5**, 314 (2009).

23. P. Juntawong, J. Bailey-Serres, Dynamic Light Regulation of Translation Status in Arabidopsis thaliana. *Front Plant Sci* **3**, 66 (2012).

24. J. Shin, A. Sanchez-Villarreal, A. M. Davis, S. X. Du, K. W. Berendzen, C. Koncz, Z. Ding, C. Li, S. J. Davis, The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. *Plant Cell Environ* **40**, 997-1008 (2017).

25. C. Robaglia, M. Thomas, C. Meyer, Sensing nutrient and energy status by SnRK1 and TOR kinases. *Curr Opin Plant Biol* **15**, 301-307 (2012).

26. M. Nietzsche, R. Landgraf, T. Tohge, F. Börnke, A protein–protein interaction network linking the energy-sensor kinase SnRK1 to multiple signaling pathways in Arabidopsis thaliana. *Current Plant Biology* **5**, 36-44 (2016).

27. E. Baena-Gonzalez, F. Rolland, J. M. Thevelein, J. Sheen, A central integrator of transcription networks in plant stress and energy signalling. *Nature* **448**, 938-942 (2007).

28. M. L. Oelze, M. O. Vogel, K. Alsharafa, U. Kahmann, A. Viehhauser, V. G. Maurino, K. J. Dietz, Efficient acclimation of the chloroplast antioxidant defence of Arabidopsis thaliana leaves in response to a 10- or 100-fold light increment and the possible involvement of retrograde signals. *J Exp Bot* **63**, 1297-1313 (2012).

29. P. Juntawong, T. Girke, J. Bazin, J. Bailey-Serres, Translational dynamics revealed by genome-wide profiling of ribosome footprints in Arabidopsis. *Proc Natl Acad Sci U S A* **111**, E203-212 (2014).

30. M. O. Vogel, M. Moore, K. Konig, P. Pecher, K. Alsharafa, J. Lee, K. J. Dietz, Fast Retrograde Signaling in Response to High Light Involves Metabolite Export, MITOGEN-ACTIVATED PROTEIN KINASE6, and AP2/ERF Transcription Factors in Arabidopsis. *Plant Cell* **26**, 1151-1165 (2014).

31. C. M. Hooper, I. R. Castleden, S. K. Tanz, N. Aryamanesh, A. H. Millar, SUBA4: the interactive data analysis centre for Arabidopsis subcellular protein locations. *Nucleic Acids Res* **45**, D1064-D1074 (2017).

32. T. L. Bailey, C. Elkan, Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology*, 28-36 (1994).

33. M. J. Liu, S. H. Wu, H. M. Chen, S. H. Wu, Widespread translational control contributes to the regulation of Arabidopsis photomorphogenesis. *Mol Syst Biol* **8**, 566 (2012).

34. S. Vij, A. K. Tyagi, Genome-wide analysis of the stress associated protein (SAP) gene family containing A20/AN1 zinc-finger(s) in rice and their phylogenetic relationship with Arabidopsis. *Mol Genet Genomics* **276**, 565-575 (2006).

35. F. Wang, R. A. Coe, S. Karki, S. Wanchana, V. Thakur, A. Henry, H. C. Lin, J. Huang, S. Peng, W. P. Quick, Overexpression of OsSAP16 Regulates Photosynthesis and the Expression of a Broad Range of Stress Response Genes in Rice (Oryza sativa L.). *PLoS One* **11**, e0157244 (2016).

36. A. Mukhopadhyay, S. Vij, A. K. Tyagi, Overexpression of a zinc-finger protein gene from rice confers tolerance to cold, dehydration, and salt stress in transgenic tobacco. *Proc Natl Acad Sci U S A* **101**, 6309-6314 (2004).

37. S. Yamasaki, A. Suzuki, Y. Yamano, H. Kawabe, D. Ueno, T. Demura, K. Kato, Identification of 5'-untranslated regions that function as effective translational enhancers in monocotyledonous plant cells using a novel method of genome-wide analysis. *Plant Biotechnol (Tokyo)* **35**, 365-373 (2018).

38. A. Nishizawa, Y. Yabuta, E. Yoshida, T. Maruta, K. Yoshimura, S. Shigeoka, Arabidopsis heat shock transcription factor A2 as a key regulator in response to several types of environmental stress. *Plant J* **48**, 535-547 (2006).

39. E. Stroher, X. J. Wang, N. Roloff, P. Klein, A. Husemann, K. J. Dietz, Redox-dependent regulation of the stress-induced zinc-finger protein SAP12 in Arabidopsis thaliana. *Mol Plant* **2**, 357-367 (2009).

40. F. Ferre, P. Clote, DiANNA: a web server for disulfide connectivity prediction. *Nucleic Acids Res* **33**, W230-232 (2005).

41. Y. Nishiyama, S. I. Allakhverdiev, N. Murata, Protein synthesis is the primary target of reactive oxygen species in the photoinhibition of photosystem II. *Physiol Plant* **142**, 35-46 (2011).

42. F. E. Belbin, Z. B. Noordally, S. J. Wetherill, K. A. Atkins, K. A. Franklin, A. N. Dodd, Integration of light and circadian signals that regulate chloroplast transcription by a nuclear-encoded sigma factor. *New Phytol*, (2016).

43. K. K. Niyogi, PHOTOPROTECTION REVISITED: Genetic and Molecular Approaches. *Annu Rev Plant Physiol Plant Mol Biol* **50**, 333-359 (1999).

44. I. Hebbelmann, J. Selinski, C. Wehmeyer, T. Goss, I. Voss, P. Mulo, S. Kangasjarvi, E. M. Aro, M. L. Oelze, K. J. Dietz, A. Nunes-Nesi, P. T. Do, A. R. Fernie, S. K. Talla, A. S. Raghavendra, V. Linke, R. Scheibe, Multiple strategies to prevent oxidative stress in Arabidopsis plants lacking the malate valve enzyme NADP-malate dehydrogenase. *J Exp Bot* **63**, 1445-1459 (2012).

45. M. L. Oelze, A. Kandlbinder, K. J. Dietz, Redox regulation and overreduction control in the photosynthesizing cell: complexity in redox regulatory networks. *Biochim Biophys Acta* **1780**, 1261-1272 (2008).

46. K. J. Dietz, M. O. Vogel, A. Viehhauser, AP2/EREBP transcription factors are part of gene regulatory networks and integrate metabolic, hormonal and environmental signals in stress acclimation and retrograde signalling. *Protoplasma* **245**, 3-14 (2010).

47. K. S. Browning, J. Bailey-Serres, Mechanism of cytoplasmic mRNA translation. *Arabidopsis Book* **13**, e0176 (2015).

48. Y. Zhang, Y. Wang, K. Kanyuka, M. A. Parry, S. J. Powers, N. G. Halford, GCN2-dependent phosphorylation of eukaryotic translation initiation factor-2alpha in Arabidopsis. *J Exp Bot* **59**, 3131-3141 (2008).

49. R. C. Wek, H. Y. Jiang, T. G. Anthony, Coping with stress: eIF2 kinases and translational control. *Biochemical Society Transactions* **34**, 7-11 (2006).

50. M. J. Clemens, A. Elia, S. J. Morley, Requirement for the eIF4E binding proteins for the synergistic down-regulation of protein synthesis by hypertonic conditions and mTOR inhibition. *PLoS One* **8**, e71138 (2013).

51. J. D. Richter, N. Sonenberg, Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature* **433**, 477-480 (2005).

52. A. N. Bruns, S. Li, G. Mohannath, D. M. Bisaro, Phosphorylation of Arabidopsis eIF4E and eIFiso4E by SnRK1 Inhibits Translation. *FEBS J*, (2019).

53. J. B. Rossel, I. W. Wilson, B. J. Pogson, Global changes in gene expression in response to high light in Arabidopsis. *Plant Physiol* **130**, 1109-1120 (2002).

54. N. Suzuki, H. Sejima, R. Tam, K. Schlauch, R. Mittler, Identification of the MBF1 heat-response regulon of Arabidopsis thaliana. *Plant J* **66**, 844-851 (2011).

55. C. Branco-Price, K. A. Kaiser, C. J. Jang, C. K. Larive, J. Bailey-Serres, Selective mRNA translation coordinates energetic and metabolic adjustments to cellular oxygen deprivation and reoxygenation in Arabidopsis thaliana. *Plant J* **56**, 743-755 (2008).

56. C. H. Chang, J. D. Curtis, L. B. Maggi, Jr., B. Faubert, A. V. Villarino, D. O'Sullivan, S. C. Huang, G. J. van der Windt, J. Blagih, J. Qiu, J. D. Weber, E. J. Pearce, R. G. Jones, E. L. Pearce, Posttranscriptional control of T cell effector function by aerobic glycolysis. *Cell* **153**, 1239-1251 (2013).

57. M. R. White, E. D. Garcin, The sweet side of RNA regulation: glyceraldehyde-3-phosphate dehydrogenase as a noncanonical RNA-binding protein. *Wiley Interdisc Rev RNA* **7**, 53-70 (2016).

58. C. Marondedze, L. Thomas, C. Gehring, K. S. Lilley, Changes in the Arabidopsis RNA-binding proteome reveal novel stress response mechanisms. *BMC Plant Biol* **19**, 139 (2019).

59. J. Giri, S. Vij, P. K. Dansana, A. K. Tyagi, Rice A20/AN1 zinc-finger containing stress-associated proteins (SAP1/11) and a receptor-like cytoplasmic kinase (OsRLCK253) interact via A20 zinc-finger and confer abiotic stress tolerance in transgenic Arabidopsis plants. *New Phytol* **191**, 721-732 (2011).

60. M. Kang, M. Fokar, H. Abdelmageed, R. D. Allen, Arabidopsis SAP5 functions as a positive regulator of stress responses and exhibits E3 ubiquitin ligase activity. *Plant Mol Biol* **75**, 451-466 (2011).

61. G. D. Kim, Y. H. Cho, S. D. Yoo, Regulatory functions of evolutionarily conserved AN1/A20-like Zinc finger family proteins in Arabidopsis stress responses under high temperature. *Biochem Biophys Res Commun* **457**, 213-220 (2015).

62. B. Vanderbeld, W. A. Snedden, Developmental and stimulus-induced expression patterns of Arabidopsis calmodulin-like genes CML37, CML38 and CML39. *Plant Mol Biol* **64**, 683-697 (2007).

63. S. Dobney, D. Chiasson, P. Lam, S. P. Smith, W. A. Snedden, The calmodulin-related calcium sensor CML42 plays a role in trichome branching. *J Biol Chem* **284**, 31647-31657 (2009).

64. H. C. Wu, D. L. Luo, F. Vignols, T. L. Jinn, Heat shock-induced biphasic Ca(2+) signature and OsCaM1-1 nuclear localization mediate downstream signalling in acquisition of thermotolerance in rice (Oryza sativa L.). *Plant Cell Environ* **35**, 1543-1557 (2012).

65. H. C. Wu, T. L. Jinn, Oscillation regulation of Ca2+ /calmodulin and heat-stress related genes in response to heat stress in rice (Oryza sativa L.). *Plant Signal Behav* **7**, 1056-1057 (2012).

66. R. Kushwaha, A. Singh, S. Chattopadhyay, Calmodulin7 plays an important role as transcriptional regulator in Arabidopsis seedling development. *Plant Cell* **20**, 1747-1759 (2008).

67. M. C. Rentel, M. R. Knight, Oxidative stress-induced calcium signaling in Arabidopsis. *Plant Physiol* **135**, 1471-1479 (2004).

68. F. Takahashi, T. Mizoguchi, R. Yoshida, K. Ichimura, K. Shinozaki, Calmodulin-dependent activation of MAP kinase for ROS homeostasis in Arabidopsis. *Mol Cell* **41**, 649-660 (2011).

69. D. Meiri, A. Breiman, Arabidopsis ROF1 (FKBP62) modulates thermotolerance by interacting with HSP90.1 and affecting the accumulation of HsfA2-regulated sHSPs. *Plant J* **59**, 387-399 (2009).

70. H. S. Jung, P. A. Crisp, G. M. Estavillo, B. Cole, F. Hong, T. C. Mockler, B. J. Pogson, J. Chory, Subset of heat-shock transcription factors required for the early response of Arabidopsis to excess light. *Proc Natl Acad Sci U S A* **110**, 14474-14479 (2013).

71. C. L. Linster, S. G. Clarke, L-Ascorbate biosynthesis in higher plants: the role of VTC2. *Trends Plant Sci* **13**, 567-573 (2008).

72. M. Mellenthin, U. Ellersiek, A. Borger, M. Baier, Expression of the Arabidopsis Sigma Factor SIG5 Is Photoreceptor and Photosynthesis Controlled. *Plants (Basel)* **3**, 359-391 (2014).

73. G. Bethke, T. Unthan, J. F. Uhrig, Y. Poschl, A. A. Gust, D. Scheel, J. Lee, Flg22 regulates the release of an ethylene response factor substrate from MAP kinase 6 in Arabidopsis thaliana via ethylene signaling. *Proceedings of the National Academy of Sciences* **106**, 8067-8072 (2009).

74. A. Mustroph, P. Juntawong, J. Bailey-Serres, Isolation of plant polysomal mRNA by differential centrifugation and ribosome immunopurification methods. *Methods Mol Biol* **553**, 109-126 (2009).

75. D. Wormuth, M. Baier, A. Kandlbinder, R. Scheibe, W. Hartung, K. J. Dietz, Regulation of gene expression by photosynthetic signals triggered through modified CO2 availability. *BMC Plant Biol* **6**, 15 (2006).

76. P. Chotewutmontri, N. Stiffler, K. P. Watkins, A. Barkan, Ribosome Profiling in Maize. *Methods Mol Biol* **1676**, 165-183 (2018).

77. Y. Liao, G. K. Smyth, W. Shi, The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic Acids Res* **41**, e108 (2013).

78. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10 (2011).

79. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359 (2012).

80. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

81. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010).

82. D. J. McCarthy, Y. Chen, G. K. Smyth, Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res* **40**, 4288-4297 (2012).

83. P. Juntawong, M. Hummel, J. Bazin, J. Bailey-Serres, Ribosome profiling: a tool for quantitative evaluation of dynamics in mRNA translation. *Methods Mol Biol* **1284**, 139-173 (2015).

84. R. A. Irizarry, B. M. Bolstad, F. Collin, L. M. Cope, B. Hobbs, T. P. Speed, Summaries of Affymetrix GeneChip probe level data. *Nucleic Acids Research* **31**, 15e-15 (2003).

85. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society* **57**, 289-300 (1995).

86. Y. Benjamini, Y. Hochberg, On the Adaptive Control of the False Discovery Rate in Multiple Testing With Independent Statistics. *Journal of Educational and Behavioral Statistics* **25**, 60-83 (2000).

87. S. Holtgrefe, J. Gohlke, J. Starmann, S. Druce, S. Klocke, B. Altmann, J. Wojtera, C. Lindermayr, R. Scheibe, Regulation of plant cytosolic glyceraldehyde 3-phosphate dehydrogenase isoforms by thiol modifications. *Physiol Plant* **133**, 211-228 (2008).

88. M. Sprenger-Haussels, B. Weisshaar, Transactivation properties of parsley proline-rich bZIP transcription factors. *The Plant Journal* **22**, 1-8 (2000).

89. A. Panda, J. Martindale, M. Gorospe, Affinity pulldown of biotinylated RNA for detection of protein-RNA complexes. *Bio-Protocol* **6**, (2016).

90. C. N. Chow, H. Q. Zheng, N. Y. Wu, C. H. Chien, H. D. Huang, T. Y. Lee, Y. F. Chiang-Hsieh, P. F. Hou, T. Y. Yang, W. C. Chang, PlantPAN 2.0: an update of plant promoter analysis navigator for reconstructing transcriptional regulatory networks in plants. *Nucleic Acids Res* **44**, D1154-1160 (2016).

91. A. Skarshewski, M. Stanton-Cook, T. Huber, S. Al Mansoori, R. Smith, S. A. Beatson, J. A. Rothnagel, uPEPperoni: an online tool for upstream open reading frame location and analysis of transcript conservation. *BMC Bioinformatics* **15**, 36 (2014).

92. D. Winter, B. Vinegar, H. Nahal, R. Ammar, G. V. Wilson, N. J. Provart, An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. *PLoS One* **2**, e718 (2007).

93. S. Lalonde, A. Sero, R. Pratelli, G. Pilot, J. Chen, M. I. Sardi, S. A. Parsa, D. Y. Kim, B. R. Acharya, E. V. Stein, H. C. Hu, F. Villiers, K. Takeda, Y. Yang, Y. S. Han, R. Schwacke, W. Chiang, N. Kato, D. Loque, S. M. Assmann, J. M. Kwak, J. I. Schroeder, S. Y. Rhee, W. B. Frommer, A membrane protein/signaling protein interaction network for Arabidopsis version AMPv2. *Front Physiol* **1**, 24 (2010).

94. P. Obrdlik, M. El-Bakkoury, T. Hamacher, C. Cappellaro, C. Vilarino, C. Fleischer, H. Ellerbrok, R. Kamuzinzi, V. Ledent, D. Blaudez, D. Sanders, J. L. Revuelta, E. Boles, B. Andre, W. B. Frommer, K+ channel interactions detected by a genetic system optimized for systematic studies of membrane protein interactions. *Proc Natl Acad Sci U S A* **101**, 12242-12247 (2004).

95. T. Seidel, D. Golldack, K. J. Dietz, Mapping of C-termini of V-ATPase subunits by in vivo-FRET measurements. *FEBS Lett* **579**, 4374-4382 (2005).

96. S. Janssen, R. Giegerich, The RNA shapes studio. *Bioinformatics* **31**, 423-425 (2015).

Figure 1: Moore et al.

Figure 2: Moore et al.

Figure 3: Moore et al.

Figure 4: Moore et al.

a

Gene ID	Target	WT	sap3 KO	cml49 KO	compartment
AT4G13800	unknown				membrane, plasma membrane
AT5G52570	BETA-OHASE 2				chloroplast, chloroplast membrane
AT5G49480	CP1				cytoplasm
AT4G04020	FIB				chloroplast
AT5G58770	CPT4				nucleus
AT3G56290	unknown				mitochondrium
AT1G67360	REF				cytoplasm, vacuola
AT1G61800	GPT2				chloroplast
AT5G20150	SPX1				cytoplasm, nucleus
AT3G21890	BBX31				cytoplasm, nucleus
AT2G46830	CCA1				nucleus
AT5G52640	HSP90.1				cytoplasm
AT2G36800	DOGT1				chloroplast, membrane
AT5G48570	ROF2				cytoplasm, vacuola
AT2G24540	AFR				cytoplasm
AT3G09350	Fes1A				cytoplasm, nucleus
AT4G21990	APR3				chloroplast
AT4G12400	HOP3				cytoplasm, nucleus
AT5G50800	ATSWEET13				membrane, plasma membrane
AT2G37180	RD28				membrane, plasma
AT4G34550	PCH1-like				chromosome, cytoplasm, nucleus
AT3G17609	HYH				nucleus
AT5G64510	TIN1				chloroplast, endoplasmic reticulum
AT1G01520	ASG4				nucleus
AT1G32900	GBSS1				chloroplast
AT2G26150	HSFA2				chloroplast, cytoplasm, nucleus
AT5G12110	eEF1B				cytoplasm
AT4G26850	VTC2				cytoplasm, nucleus
AT2G33380	RD20				chloroplast membrane, endoplasmic reticulum
AT3G21150	BBX32				cytoplasm, nucleus
AT3G09450	unknown				membrane
AT2G21320	BBX18				nucleus
AT1G58520	RXW8				chloroplast
AT5G24120	SIG 5				chloroplast, mitochondrium
AT5G65380	MATE				membrane
AT3G53830	RCC1				cytoplasm, nucleus
AT3G17790	PAP17				cell surface
AT1G54050	HSP20-like				cytoplasm
AT1G62180	ATAPR2				chloroplast
AT3G21670	ATNPF6.4				membrane
AT3G12580	HSP70				membrane
AT2G46790	PRR9				nucleus
AT1G77450	NAC032				nucleus
AT2G15020	unknown				mitochondrium
AT3G47420	PS3				membrane
AT5G63130	Phox/Bem1p				nucleus
AT1G01720	ANAC2				nucleus
AT4G26150	CGA1				nucleus
AT1G74310	HSP101				chloroplast stroma, cytoplasm, cytosol, nucleus

(log₂ fold change) -1 ——— 5

Figure 5: Moore et al.

Translation-Dependent Retrograde Signalling (TraDeRS)