Predicting the zoonotic capacity of mammals to transmit SARS-CoV-2

Ilya R. Fischhoff[1]*, Adrian A. Castellanos[1]*, João P.G.L.M. Rodrigues[2], Arvind Varsani[3,4],

Barbara A. Han[1]†

Affiliations:

[1] Cary Institute of Ecosystem Studies. Box AB Millbrook, NY 12545, USA

[2] Department of Structural Biology, Stanford University School of Medicine, Stanford, CA 94305,

USA

[3] The Biodesign Center for Fundamental and Applied Microbiomics, Center for Evolution and

Medicine, School of Life Sciences, Arizona State University, Tempe, AZ 85287, USA

[4] Structural Biology Research Unit, Department of Integrative Biomedical Sciences, University of

Cape Town, Rondebosch, 7700, Cape Town, South Africa

* contributed equally

† corresponding author

Email addresses:

Ilya Fischhoff: fischhoff@gmail.com

Adrian A. Castellanos: castellanosa@caryinstitute.org

João P.G.L.M. Rodrigues: joaor@stanford.edu

Arvind Varsani: arvind.varsani@asu.edu

Barbara A. Han: hanb@caryinstitute.org

Keywords: coronavirus; COVID-19; hosts; reservoirs; ecological traits; zoonotic; spillover;

spillback; susceptibility; machine learning; homology modelling; ACE2

Abstract

Back and forth transmission of SARS-CoV-2 between humans and animals may lead to wild
reservoirs of virus that can endanger efforts toward long-term control of COVID-19 in people,
and protecting vulnerable animal populations that are particularly susceptible to lethal disease.
Predicting high risk host species is key to targeting field surveillance and lab experiments that
validate host zoonotic potential. A major bottleneck to predicting animal hosts is the small
number of species with available molecular information about the structure of ACE2, a key
cellular receptor required for viral cell entry. We overcome this bottleneck by combining species'
ecological and biological traits with 3D modeling of virus and host cell protein interactions using
machine learning methods. This approach enables predictions about the zoonotic capacity of
SARS-CoV-2 for over 5,000 mammals — an order of magnitude more species than previously
possible. The high accuracy predictions achieved by this approach are strongly corroborated by
in vivo empirical studies. We identify numerous common mammal species whose predicted
zoonotic capacity and close proximity to humans may further enhance the risk of spillover and
spillback transmission of SARS-CoV-2. Our results reveal high priority areas of geographic
overlap between global COVID-19 hotspots and potential new mammal hosts of SARS-CoV-2.
With molecular sequence data available for only a small fraction of potential host species,
predictive modeling integrating data across multiple biological scales offers a conceptual
advance that may expand our predictive capacity for zoonotic viruses with similarly unknown
and potentially broad host ranges.

50
Introduction

The ongoing COVID-19 pandemic has surpassed 3.9 million deaths globally as of 25
June 2021 [1,2]. Like previous pandemics in recorded history, COVID-19 originated from the
spillover of a zoonotic pathogen, SARS-CoV-2, a betacoronavirus originating from an unknown
animal host [3–6]. The broad host range of SARS-CoV-2 is due in part to its use of a highly
conserved cell surface receptor to enter host cells, the angiotensin-converting enzyme 2
receptor (ACE2) [7] found in all major vertebrate groups [8].
58
The ubiquity of ACE2 coupled with the high prevalence of SARS-CoV-2 in the global
human population explains multiple observed *spillback* infections since the emergence of
SARS-CoV-2 in 2019 (see natural infections listed in Table 1 with references). In spillback
infection, human hosts transmit SARS-CoV-2 virus to cause infection in non-human animals. In
addition to threatening wildlife and domestic animals, repeated spillback infections may lead to
the establishment of new animal hosts from which SARS-CoV-2 can continue to pose a risk of
secondary spillover infection to humans through bridge hosts (e.g., [9]) or newly established
enzootic reservoirs. Indeed, this risk has already been realized in Denmark [10] and The
Netherlands, where SARS-CoV-2 spilled back from humans to farmed mink (*Neovison vison*)
with secondary spillover of a SARS-CoV-2 variant from mink back to humans [11]. A major
concern in such secondary spillover events is the appearance of a mutant strain [11,12]
affecting host range [13] or leading to increased transmissibility in humans [14,15] (but see
[16,17]). Preliminary evidence shows that the mink-derived variant exhibits moderately reduced
sensitivity to neutralizing antibodies [10], raising concerns that humans may eventually
experience infections from spillback variants, and that vaccines may become less efficient at
conferring immunity to these variants [18]. Conversely, human-derived variants pose spillback
risks to animals. For example, in contrast to previous infection trials [19], two new human
variants are now confirmed to have overcome the species barrier to infect lab mice (*Mus
musculus*) [20].
78
Spillback infections from humans to animals are already occurring worldwide. A variety
of pets, domesticated animals, zoo animals, and wildlife have also been documented as new
hosts of SARS-CoV-2 (Table 1). In addition to secondary spillover infections from mink farms,
SARS-CoV-2 has been found for the first time in wild and escaped mink in multiple states in the
United States, with viral sequences identical to SARS-CoV-2 in nearby farmed mink [21–23].
The global scale of human infections and the increasing range of known hosts observed for
SARS-CoV-2 demonstrate that SARS-CoV-2 has the capacity to establish novel enzootic
infection cycles in animals. In response, recent computational studies make predictions about
the susceptibility of particular animal species to SARS-CoV-2 [13,24–32]. These studies
compare known sequences of ACE2 orthologs across species (*sequence-based* studies), or
model the structure of the viral spike protein bound to ACE2 orthologs (*structure-based* studies).
These studies yield a wide range of predictions with varying degrees of agreement with
laboratory animal experiments (Figure 1).
92

93
94 **Table 1**. Species with confirmed suitability for SARS-CoV-2 infection from natural infections or *in vivo*
95 experiments. Asterisks reference species with infection status from preprints (not yet peer-reviewed).
96 Some species (e.g, dogs) with natural infection studies also have *in vivo* experimental studies.

Species	Susceptibility	Study type	Location	References
Cow (*Bos taurus*)	Yes	*In vivo* experiment	Lab	[33]
Dog (*Canis lupus familiaris*)	Yes	Natural infection	Multiple countries	[34–38]
African green monkey (*Chlorocebus aethiops*)	Yes	*In vivo* experiment	Lab	[39]
Big brown bat (*Eptesicus fuscus*)	No	*In vivo* experiment	Lab	[40]
Cat (*Felis catus*)	Yes	Natural infection	Multiple countries	[35,36,38,41]
Gorilla (*Gorilla gorilla*)	Yes	Natural infection	USA, Zoo	[42]
Crab-eating macaque (*Macaca fascicularis*)	Yes	*In vivo* experiment	Lab	[43]
Rhesus macaque (*Macaca mulatta*)	Yes	*In vivo* experiment	Lab	[44]
Golden hamster (*Mesocricetus auratus*)	Yes	*In vivo* experiment	Lab	[45]
House mouse (*Mus musculus*)	No	*In vivo* experiment	Lab	[19] (but see [20])
Ferret (*Mustela putorius furo*)	Yes	*In vivo* experiment	Lab	[37]
American mink (*Neovison vison*)	Yes	Natural infection	Multiple countries	[35,36,46]
Raccoon dog (*Nyctereutes procyonoides*)	Yes	*In vivo* experiment	Lab	[47]

European rabbit (*Oryctolagus cuniculus*)	Yes	*In vivo* experiment	Lab	[48]
Lion (*Panthera leo*)	Yes	Natural infection	Multiple countries, Zoos	[36,49]
Tiger (*Panthera tigris*)	Yes	Natural infection	USA and Sweden, Zoos	[35,36,49,50]
Deer mouse (*Peromyscus maniculatus*)*	Yes	*In vivo* experiment	Lab	[51,52]
Cougar (*Puma concolor*)	Yes	Natural infection	South Africa, Zoo	[36]
Egyptian fruit bat (*Rousettus aegyptiacus*)	Yes	*In vivo* experiment	Lab	[53]
Pig (*Sus scrofa*)	No	*In vivo* experiment	Lab	[37,53]
Northern treeshrew (*Tupaia belangeri*)	Yes	*In vivo* experiment	Lab	[54]
Snow leopard (*Uncia uncia*)	Yes	Natural infection	USA, Zoo	[55]
Bank vole (*Clethrionomys glareolus*)	Yes	*In vivo* experiment	Lab	[56]
Asian small-clawed otter (*Aonyx cinereus*)	Yes	Natural infection	USA, Zoo	[36,57]
White-tailed deer (Odocoileus virginianus)	Yes	In vivo experiment	Lab	[58]

97
98
99

Figure 1. A heatmap summarizing predicted susceptibility to SARS-CoV-2 for species with confirmed infection from *in vivo* experimental studies or from documented natural infections. Studies that make predictions about species susceptibility are shown on the x-axis, organized by method of prediction (those relying on ACE2 sequences, estimating binding strength using three dimensional structures, or laboratory experiments). Predictions about zoonotic capacity from this study are listed in the second to last column, with high and low categories determined by zoonotic capacity observed in *Felis catus*. Confirmed infections for species along the y-axis are summarized in [59] and are depicted as a series of filled or unfilled circles. Bolded species have been experimentally confirmed to transmit SARS-CoV-2 to naive conspecifics. Species predictions range from warmer colors (yellow: low susceptibility or zoonotic capacity for SARS-CoV-2) to cooler colors (purple: high susceptibility or zoonotic capacity). See

Supplementary Methods (https://doi.org/10.25390/caryinstitute.c.5293339) for detailed methods
about study categorization.

Sequence-based studies

Sequence-based studies predict host susceptibility based on amino acid sequence
similarity between human (hACE2) and non-human ACE2, and assume that a high degree of
similarity correlates with stronger viral binding, especially at amino acid residues where hACE2
interacts with the SARS-CoV-2 spike glycoprotein. For some species, such as rhesus
macaques [60], these qualitative predictions are borne out by *in vivo* studies (Figure 1), but
predictions from these methods do not consistently match real-world outcomes. For example,
sequence similarity predicted weak viral binding for minks and ferrets, which have all been
confirmed as highly susceptible, with minks capable of onward transmission [11,32,37] (Figure
1). These mismatches to experimental or real-world outcomes may arise in part because protein
three-dimensional structure, the main determinant of protein function, is incompletely
represented by 1D amino acid sequences [61,62]. As such, details about the interaction
between host ACE2 and the viral spike protein are not well captured by sequence-based
studies.

Structure-based studies

Modeling the three-dimensional structure of protein-protein complexes addresses some
of the limitations of sequence-based approaches, and has proven useful to predict how different
ACE2 orthologs bind to the SARS-CoV-2 viral spike protein receptor-binding domain (RBD)
[13,24]. These studies have also identified particular ACE2 amino acid residues essential for a
productive interaction with the viral RBD, thus improving predictive models of susceptibility
through structure-based inference [13]. These studies leverage known structures of the hACE2
receptor bound to the SARS-CoV-2 RBD and use powerful simulation methods to predict how
variation across different ACE2 orthologs affects binding with the viral RBD. While these
approaches successfully predicted strong binding for species that have been infected (e.g.
domestic cat, tiger, dog, and ferret) and weak binding for species in which experimental
infections failed (e.g. chicken, duck [37], mouse [19]), the results are also not consistently
supported by experiments. For instance, while guinea pig ACE2 scored favorably among
susceptible species in one of the studies [13], this ortholog was shown experimentally not to
bind to the SARS-CoV-2 RBD [63].
142
Although structural modeling has produced the most accurate results to date, all
currently available approaches for predicting the host range of SARS-CoV-2 are fundamentally
constrained by the availability and quality of ACE2 sequences across species. ACE2 is
ubiquitous across chordates, likely because of its role in several highly conserved physiological
pathways [64]. Because it is so highly conserved, the vast majority of mammal species (>6,000
species) are likely to have ACE2 receptors, but there are many fewer sequences available from
which to make predictions using existing modeling methods (~300 species). The functional
importance of the ACE2 receptor suggests that it has evolved in association with other intrinsic

organismal traits that are more easily observed and for which data are available for many more
species. These suites of correlated organismal traits may provide a robust statistical proxy that
can be leveraged to predict suitable hosts for SARS-CoV-2. Previous trait-based analyses
applied statistical (machine) learning techniques to accurately distinguish the zoonotic capacity
of various organisms [65–67], and predict likely hosts for particular groups of related viruses
[68,69], predictions which have subsequently been validated through independent laboratory
and field investigations (e.g., [70,71]).
158
Here, we combine molecular structural modeling of viral binding with machine learning of
species-level ecological and biological traits to predict species' zoonotic capacity for SARS-
CoV-2 virus across 5,400 mammal species, expanding our predictive capacity by an order of
magnitude (Figure 2). Crucially, this integrated approach enables predictions for the vast
majority of species whose ACE2 sequences are currently unavailable by leveraging information
from viral binding dynamics and biological traits of potential hosts. In our workflow (Figure 2), we
first carry out structural modeling to quantify the binding strength of SARS-CoV-2 RBD for
vertebrate species using published ACE2 amino acid sequences [72]. We then collate species
traits and train a machine learning model to predict the zoonotic capacity for 5,400 mammal
species. Zoonotic capacity (host susceptibility and capacity for onward transmission) was
approximated through a conservative threshold of binding strength applied to our structural
modeling results and reported by *in vivo* studies.
171
COVID-19 is, at this time, primarily a disease affecting humans, thus spillback infection
of SARS-CoV-2 from humans to animals is the most likely mode by which new host species will
become established. We therefore identify a subset of species for which the threat of spillback
infection appears greatest due to geographic overlaps and opportunities for contact with
humans in areas of high SARS-CoV-2 prevalence globally. Our predictions contribute to a
critical interdisciplinary and iterative process between computational modeling, field
surveillance, and laboratory experiments that is necessary for improving zoonotic risk
quantification, and to better inform next steps toward the prevention of enzootic SARS-CoV-2
transmission and spread. We demonstrate our approach using the SARS-CoV-2 sequence that
initially emerged in humans. These methods can be readily expanded to enable host range
predictions for new variants as their hACE2-RBD crystal structures become available.

183
Figure 2. A flowchart showing the progression of our workflow combining evidence from limited lab and
field studies with additional data types to predict zoonotic capacity across mammals through multi-scale
statistical modeling (gray boxes, steps 1-5). For all vertebrates with published ACE2 sequences, we
modelled the interface of species' ACE2 bound to the viral receptor binding domain using HADDOCK. We
then combined the HADDOCK scores, which approximate binding strength, with species' trait data and
trained machine learning models for both mammals and vertebrates (yellow boxes). Mammal species
predicted to have high zoonotic capacity were then compared to results of *in vivo* experiments and *in*
silico studies that applied various computational approaches. Based on predictions from our model, we
identified a subset of species with particularly high risk of spillback and secondary spillover potential to
prioritize additional lab validation and field surveillance (dashed line).

194
195

Methods

Protein sequence and alignment

We assembled a dataset of ACE2 NCBI GenBank accessions that are known human
ACE2 orthologs or have high similarity to known orthologs as determined using BLASTx [73].
Using the R package *rentrez* and the accession numbers, we downloaded ACE2 protein
sequences [74]. We supplemented these sequences by manually downloading four additional
sequences from the MEROPS database [75].

Structural Modeling of ACE2 orthologs bound to SARS-CoV-2 spike

The modeling of all 326 ACE2 orthologs bound to SARS-CoV-2 spike receptor binding
domain was carried out as described previously [13], with a few differences. Sequences of
ACE2 orthologs were aligned using MAFFT [76] and trimmed to the region resolved in the
template crystal structure of hACE2 bound to the SARS-CoV-2 spike (PDB ID: 6m0j, [77].

Ambiguous positions in each sequence, artifacts of the sequencing method, were replaced by
Glycine to minimize assumptions about the nature of the amino acid side-chain but still allow for
modeling. For each ortholog, we generated 10 homology models using MODELLER 9.24
[78,79], with restricted optimization (*fastest* schedule) and refinement (*very_fast* schedule)
settings, and selected a representative model based on the normalized DOPE score. These
representative models were then manually inspected and 27 were removed from further
analysis due to large insertions/deletions or to the presence of too many ambiguous amino
acids at the interface with spike. Each validated model was submitted for refinement to the
HADDOCK web server [80], which ran 50 independent short molecular dynamics simulations in
explicit solvent to optimize the interface between the two proteins . For each one of the animal
species in our study, we assigned an average and standard deviation of the scores of the 10
best refined models, ranked by their HADDOCK score -- a combination of van der Waals,
electrostatics, and desolvation energies. A lower (more negative) HADDOCK score predicts
stronger binding between the two proteins. We hereafter refer to predicted binding strength, or
simply binding strength, to indicate HADDOCK score. The HADDOCK server is freely available,
and we provide code to reproduce analyses or to aid in the application of this modeling
approach to other similar problems (https://zenodo.org/record/4517509).

Trait data collection and cleaning

We gathered ecological and life history trait data from AnAge [81], Amniote Life History
Database [82], and EltonTraits [83], among other databases (Supplementary Table 1; for details
on data processing, see Supplementary Methods with all supplementary data, figures, methods,
and tables available at https://doi.org/10.25390/caryinstitute.c.5293339). Using these data, we
also engineered additional traits that have shown importance in predicting host-pathogen
associations in other contexts. For example, as a measure of habitat breadth [84], we computed
for each species the percentage of ecoregions it occupies. To assess the influence of sampling
bias across species, we used the wosr R package [85] to count the number of studies returned
in a search in Web of Science for each species' Latin binomial and included this as a proxy for
sampling bias in our model.
236
Following the results of initial structural modeling (described above), we observed that
per-residue energy decomposition analysis of HADDOCK scores for 29 species indicated that
all species with strong predicted binding had in common a salt bridge between SARS-CoV-2
K417 and a negatively charged amino acid at position 30 in the ACE2 sequence [13]. Given the
apparent effect of amino acid 30 on overall binding strength, we constructed an additional
feature to denote whether amino acid 30 is negatively charged (and therefore more likely to
support strong binding) and included this feature as an additional trait in our models.

Modeling

245
Quantifying a threshold for zoonotic capacity using HADDOCK. While ACE2 binding is
necessary for viral entry into host cells, it is not sufficient for SARS-CoV-2 transmission. Multiple

in vivo experiments suggest that not all species that are capable of binding SARS-CoV-2 are
capable of transmitting active infection to other individuals (e.g., cattle, *Bos taurus* [33]; bank
voles, *Myodes glareolus [56]*). Viral replication, and infectious viral shedding that enables
onward transmission, are both required for a species to become a suitable bridge or reservoir
species for SARS-CoV-2. In order to constrain our predictions to species with the greatest
potential to perpetuate onward transmission, we trained our models on a conservative threshold
of binding strength (HADDOCK score = -129). This value is between the scores for two species:
the domestic cat (*Felis catus*), which is currently the species with weakest predicted binding with
confirmed conspecific transmission [86], and the pig (*Sus scrofa*), which shows the strongest
estimated binding for which experimental inoculation failed to cause detectable infection [37].
Binding strength was binarized according to this threshold, above which it is more likely that
both infection and onward transmission will occur following the results of multiple empirical
studies (Table 1). We note that there are species confirmed to be susceptible whose predicted
binding strength is weaker than cats, but conspecific transmission has not been confirmed in
these species. While it is likely that intraspecific transmission will be reported for additional
species as the pandemic continues, the binding strength selected for this analysis represents an
appropriately conservative threshold based on currently available evidence. For additional
modeling details, see Supplementary Methods.

Trait-based modeling to predict zoonotic capacity

We applied generalized boosted regression [87] to host trait data to predict species'
binding strength to SARS-CoV-2. We applied this approach initially to all of the vertebrate
species for which we estimated HADDOCK scores, but these models did not perform well. This
was likely due to extensive dissimilarities among traits describing different classes of organisms.
For instance, traits that are commonly measured for reptiles are different from those of interest
for birds or amphibians. Moreover, currently available ACE2 sequences are dominated by ray-
finned fishes and mammals.
274
Given that only mammals have so far been confirmed as both susceptible and capable
of onward transmission of SARS-CoV-2, we created a separate set of models to make zoonotic
capacity predictions for mammals only. For this mammal-only dataset, we gathered additional
species-level traits from PanTHERIA [88] and added a series of binary fields for taxonomic
order (based on [89]; Supplementary Table 2). We then applied boosted regression (BRT; gbm
package [90] in R version 4.0.0 [90,91]) to impute missing trait data for mammal species (e.g., [67];
see Supplementary Methods for details on imputation methods and results).
282
Many of the mammals for which we found the strongest evidence of zoonotic capacity
are domesticated to some degree (pets, farmed or traded animals, lab models) [11,37,53].
Relative to their ancestors or wild conspecifics, domesticated animals often have distinctive
traits [92] that are likely to influence the number of zoonoses found in these species [93]. To
account for trait variation due to domestication in certain species, we modeled mammals in two
ways. First, we incorporated a variable indicating whether the source populations from which
trait data were collected are wild or non-wild (e.g., farmed, pets, laboratory animals; non-wild

status confirmed by the Mammal Diversity Database [94]). Trait data collected from both wild
and non-wild individuals were considered to represent non-wild species for the purposes of this
model. In a second approach, we used only the wild species for model training and evaluation.
For both approaches, pre-imputation trait values were used for all non-wild mammals during
model training, evaluation, and prediction.
295
Boosted regression (BRT) is an ensemble machine learning approach that
accommodates non-random patterns of missing data, nonlinear relationships, and interacting
effects among predictors. In a BRT model, a sequence of regression models are fit by recursive
binary splits, with each additional regression modeling those instances that were poorly
accounted for by the previous regression iterations in the tree [87]. We applied grid search to
select optimal hyperparameters, and repeated model fitting 50 times using bootstrapped training
sets of 80% of labeled data. We measured performance by the area under the receiver
operating characteristic curve (AUC) for predictions made on the test dataset (remaining 20%),
corrected by comparing with null models created by target shuffling, which employed similar
bootstrapping (50 times). Detailed methods can be found in Supplementary Methods. We
discuss herein the results of model predictions about zoonotic capacity made by applying this
final model to all mammal species. We also report the mean and variation in predicted
probabilities across all 50 bootstrapped models in Supplementary File 1.
309
To visualize geographic patterns, we mapped the geographic ranges of mammal species
predicted within the 90th percentile of zoonotic capacity for SARS-CoV-2 using International
Union for the Conservation of Nature (IUCN) polygons of species distributions [95]. We subset
to the species found in human-associated habitats (e.g., urban areas, crop lands, heavily
degraded forests; based on IUCN 2020), and also masked their ranges to areas of high human
case counts (using SARS-CoV-2 case data from the COVID-19 Data Repository at Johns
Hopkins University [1]).
317
Additional methods and results of other uninformative model variations are also
described in Supplementary Methods and Supplementary Table 3 (e.g., a model in which
binding strength is modeled as a continuous rather than a threshold measure, a model
predicting the charge at amino acid 30, a model for all vertebrate species)
(https://doi.org/10.25390/caryinstitute.c.5293339). We provide code and data files for carrying
out boosted regression tree models
(https://github.com/HanLabDiseaseEcology/zoonotic_capacity). Details about how the species
susceptibility predictions from past studies were standardized into categories (low, medium,
high; Figure 1) are also available in Supplementary Methods.
327
328

Results

ACE2 host protein sequences and alignment

The ACE2 protein sequence alignment of the orthologs from 326 species spans eight
classes and 87 orders (https://zenodo.org/record/4517509). The majority of sequences
belonged to the classes Actinopterygii (22.1%), Aves (23.3%), and Mammalia (46.6%).
Sequence length ranged from 344 amino acids to 872 with a median length of 805.

Structural modeling of viral binding strength

We predicted binding strength for 299 vertebrates, including 142 mammals. These
binding strength scores represented six classes and 80 orders and ranged between -167.816
and -105.615. Across these six vertebrate classes, the strongest predicted binding between
ACE2 and SARS-CoV-2 (corresponding to the lowest mean HADDOCK scores) were in ray-
finned fishes (Actinopterygii; mean = -137.945) and mammals (Mammalia; mean = -129.193)
(Figure 3A). Four of these six classes included at least one species predicted to have stronger
binding than *Felis catus* (Figure 3B). Among well-represented mammalian orders (those
containing at least 10 species with binding strength predictions), Primates and Carnivora
showed predicted mean binding strengths that were stronger than domestic cats (Figure 3C).
345

346
347
Figure 3. Plots showing results from modeling species' ACE2 interaction with SARS-CoV-2 RBD using
HADDOCK to predict binding strength (measured as arbitrary units). HADDOCK scores that predict
stronger binding are more negative. The mean and standard deviation of the HADDOCK score for
vertebrate species **(A)** for which ACE2 orthologs are available. Binding strengths vary across vertebrate
classes **(B)** and across the five most speciose mammalian orders **(C)**. The "Other" category contains
species across multiple orders for which ACE2 sequences were available, each with fewer than 10
representative species in the order. The shaded regions of all panels represent predicted binding that is
as strong or stronger than (more negative values than) the domestic cat (*Felis catus*), which represents
our conservative zoonotic capacity threshold based on currently available empirical evidence.

Species predictions of zoonotic capacity from trait-based machine learning models

The best performing model was trained on a mammal-only dataset with trait imputation
and showed corrected test AUC of 0.72 (for results of all other model variations, see
Supplementary Table 3). We used this model to generate predictions of zoonotic capacity
among mammal species. Citation count, as a proxy for study effort, had ~1% relative
importance, suggesting that sampling bias across species had little influence on the model.

This zoonotic capacity model identified 540 species (out of 5400 total mammal species)
within the 90th percentile probability (0.826 or higher, compared to a total of 2,401 mammal
species with prediction scores above 0.5; see Supplementary File 1 for predictions on all 5,400
species, https://doi.org/10.25390/caryinstitute.c.5293339). The top 10% of species with the
highest predicted probabilities includes representatives from 13 orders. Most primates were
predicted to have high zoonotic capacity and collectively showed stronger viral binding
compared to other mammal groups (Figure 4). Additional orders with numerous species
predicted to have high zoonotic capacity (at least 75% of species above 0.5) include Hyracoidea
(hyraxes), Perissodactyla (odd-toed ungulates), Scandentia (treeshrews), Pilosa (sloths and
anteaters), Pholidota (pangolins), and non-cetacean Artiodactyla (even-toed ungulates) (Figure
4).

377
378
Figure 4. Ridgeline plots showing the distribution of predicted zoonotic capacity across mammals.
Predicted probabilities for zoonotic capacity across the x-axis range from 0 (likely not susceptible) to 1
(zoonotic capacity predicted to be the same or greater than *Felis catus*), with the vertical line representing
0.5. The y-axis depicts all mammalian orders represented by our predictions. Density curves represent
the distribution of the predictions, with those parts of the curve over 0.5 colored pink and lines
representing distribution quartiles. The predicted values for each order are shown as points below the

density curves. Points that were used to train the model are colored: orange represents species with
weaker predicted binding, blue represents species with stronger predicted binding. Selected family-level
distributions are shown in the Supplemental Figures 5-6
(https://doi.org/10.25390/caryinstitute.c.5293339).
389
390

Comparison of species predictions

Comparing species predictions across multiple computational approaches

Our model combined species traits with estimates of viral binding strength to predict
zoonotic capacity, which encompasses both susceptibility to SARS-CoV-2 and the probability of
onward transmission. Zoonotic capacity was defined as a threshold value based on the results
of experimental studies confirming intraspecific transmission among animals, and is therefore
more conservative than thresholds adopted by other studies (e.g., those based only on
estimates of viral binding strength, [30]). In addition, our modeling approach (machine learning)
and prediction targets (zoonotic capacity) differed compared to existing computational
approaches, which applied sequence-based or structure-based analyses constrained by the
small number of published ACE2 sequences. Despite these differences, comparing the species
predictions generated by multiple different approaches can be useful for gauging consensus,
and for comparing how species predictions change from one method to another.
404
Across approaches, there was general agreement in the predictions for primates as well
as for a select group of artiodactyls and carnivores (Figure 5). Our model results also agreed
with low susceptibility predictions made by several previous studies using sequence-based
approaches (e.g., in certain bats and rodents). In general, we note that structure-based models
predicted a smaller proportion of species to have low susceptibility compared to sequence-
based studies.

Figure 5. An alluvial plot comparing predictions of species susceptibility from multiple methods. Existing studies (listed in Supplementary Methods) are categorized as either sequence-based or structure-based. Predictions from our zoonotic capacity model result from combining structure-based modeling of viral binding with organismal traits using machine learning to distinguish species with zoonotic capacity above (1) or below (0) a conservative threshold value set by domestic cats (*Felis catus*). Colors represent unique mammalian orders, and the width of colored bands represent the relative number of species with that combination of predictions across methods. See Supplementary Methods (https://doi.org/10.25390/caryinstitute.c.5293339) for details on how species across multiple studies were assigned to categories (high, medium, low).

Comparing model predictions to in vivo *outcomes*

Our model predictions matched the results of several recently published *in vivo* studies on SARS-CoV-2 infection (Figure 1). For instance, experiments on deer mice (*Peromyscus maniculatus*; [51,52]) and raccoon dogs (*Nyctereutes procyonoides*; [47]) confirmed SARS-CoV-2 infection and transmission to naive conspecifics. Our model also estimated a high probability of zoonotic capacity of American mink for SARS-CoV-2 (*Neovison vison*, probability=0.83, 90th percentile), in which farmed individuals present severe infection from human spillback, and demonstrate the capacity to transmit to conspecifics as well as to humans [11,46]. Our model also correctly predicted relatively low zoonotic capacity for big brown bats (*Eptesicus fuscus*; [40]).

There were notable differences between our model results and the outcomes of some
experimental studies. For instance, our model estimated a moderately high probability of
zoonotic capacity for pigs (*Sus scrofa*, probability = 0.72, ~80th percentile). Similarly, some
computational and cell-based studies have also predicted strong viral binding in this species
[26,96], but *in vivo* studies report no detectable infection or onward transmission of SARS-CoV-
2 [37,53]. Similarly for cattle (*Bos taurus*), our model estimated a moderately high probability for
zoonotic capacity (0.72, ~80th percentile), and in a live animal experiment, cattle were
confirmed to be susceptible to infection but no onward transmission was observed to virus-naive
conspecifics [33].

444
Figure 6: Maps showing the global distribution of species with predicted capacity to transmit SARS-CoV-
2. **(A)** depicts global species richness of the top 10 percent of model-predicted zoonotic capacity.
Geographic ranges of this subset of species were filtered to those associated with human-dominated or
human-altered habitats **(B)**, and further filtered to show the subset of species that overlaps with areas of
high human SARS-CoV-2 positive case counts (over 100,000 cumulative cases as of 17 May 2021) **(C)**.
For a full list of model-predicted zoonotic capacity of species by country, see Supplementary File 2
(https://doi.org/10.25390/caryinstitute.c.5293339).
452
453

Discussion

We combined structure-based models of viral binding with species-level data on
biological and ecological traits to make predictions about the capacity of animal species to
become zoonotic hosts of SARS-CoV-2 (*zoonotic capacity*). This combined modeling approach
predicted zoonotic capacity with 72% accuracy, extending our predictive capacity beyond the
limited number of species for which ACE2 sequences are currently available. We identified
numerous mammal species whose predicted zoonotic capacity meets or exceeds the viral
susceptibility and transmissibility observed in experimental infections with SARS-CoV-2. In
addition to wide agreement with *in vivo* study results produced to date (Table 1), these model
predictions corroborate the predictions of previous studies generated using the limited number
of available ACE2 sequences (Figure 1). Below we discuss predictions of zoonotic capacity for
a number of ecologically and epidemiologically relevant categories of mammalian hosts.
466
Captive, farmed, or domesticated species. Given that the type and frequency of contact with
humans fundamentally underlies transmission risk, it is notable that our model predicted high
zoonotic capacity for multiple captive species that have also been confirmed as susceptible to
SARS-CoV-2 via experiments or natural infections. These include numerous carnivore species,
such as large cats from multiple zoos and pet dogs and cats. Our model also predicted high
SARS-CoV-2 zoonotic capacity for many farmed, domesticated, and live traded species. The
water buffalo (*Bubalus bubalis*), widely bred for dairy production and farming, had the highest
probability of zoonotic capacity among livestock (0.91). Model predictions in the 90th percentile
also included American mink (*Neovison vison*), red fox (*Vulpes vulpes*), sika deer (*Cervus
nippon*), white-lipped peccary (*Tayassu pecari*), nilgai (*Boselaphus tragocamelus*), and raccoon
dogs (*Nyctereutes procyonoides*), all of which are farmed, with the latter two considered
invasive species in some areas [97,98]. In addition to the risks of secondary spillover to humans
and the potential for large economic losses from culling infected animals [99], the escape of
farmed individuals into wild populations has implications for the spread and enzootic
establishment of SARS-CoV-2 [21]. These findings also have implications for vaccination
strategies, for instance, prioritizing people in regular contact with potential bridge species (e.g.,
veterinarians, abattoir-workers, farmers, etc).
484
Live traded or hunted wildlife species. The majority of the legally traded live mammals are
primates and carnivores [100], and model predictions included several species from these
groups. Our model predicted high zoonotic capacity in 20 out of 21 species in the primate genus
Macaca, which comprise the majority of all live-traded primates. Several live-traded carnivores
and pangolins were also assigned high zoonotic capacity, including the Asiatic black bear
(*Ursus thibetanus*), grey wolf (*Canis lupus*), and jaguar (*Panthera onca*), the Philippine pangolin
(*Manis culionensis*) and Sunda pangolin (*M. javanica*). Pangolins are notable because one of
the betacoronaviruses with the highest sequence similarity to SARS-CoV-2 was isolated from
Sunda pangolins [101,102]. Pangolin burrows are also known to be occupied by multiple other
animal species, including numerous bats [103].
495

496   Commonly hunted species in the top 10% of predictions include duiker (*Cephalophus*
497 *zebra*, West Africa), warty pig (*Sus celebes*, Southeast Asia), and two species of deer
498 (*Odocoileus hemionus* and *O. virginianus*) that are widespread across the Americas. The white-
499 tailed deer (*O. virginianus*) was recently confirmed to be capable of transmitting SARS-CoV-2 to
500 conspecifics via indirect contact (aerosolized virus particles) [58].
501
Bats. Similarly, bats are of special interest because of the high diversity of betacoronaviruses
found in *Rhinolophus spp.* and other bat species [104–107]. Our model identified 35 bat species
within the 90th percentile of zoonotic capacity for SARS-CoV-2. Within the genus *Rhinolophus*,
our model identified the large rufous horseshoe bat (*Rhinolophus rufus*), a known natural host
for bat betacoronaviruses [104] and a congener to three other horseshoe bats harboring
betacoronaviruses with high nucleotide sequence similarity to SARS-CoV-2 (~92-96%)
[6,108,109]. For these three species, our model assigned a range of probabilities for SARS-
CoV-2 zoonotic capacity (*Rhinolophus affinis* (0.58), *R. malayanus* (0.70), and *R. shameli*
(0.71)) and also predicted relatively high probabilities for two congeners, *Rhinolophus*
acuminatus (0.84) and *R. macrotis* (0.70). These predictions are in agreement with recent
experiments demonstrating efficient viral binding of SARS-CoV-2 RBD for *R. macrotis* [110] and
confirmation of SARS-CoV-2-neutralizing antibodies in field-caught *R. acuminatus* harboring a
closely related betacoronavirus [111].
515
Our model also identified 17 species in the genus *Pteropus* (flying foxes) with high
probabilities of zoonotic capacity for SARS-CoV-2. Some of these species are confirmed
reservoirs of other zoonotic viruses in Southeast Asia (e.g., henipaviruses in *P. lylei*, *P.*
vampyrus, *P. conspicillatus*, and *P. alecto*). While contact patterns between bats and humans
may be somewhat less direct compared with captive or farmed species, annual outbreaks
attributed to viral spillover transmission from bats illustrate a persistent epizootic risk to humans
[112–114] and confirm that gaps in systematic surveillance of zoonotic viruses, including
betacoronaviruses, remain an urgent priority (e.g., [115]).
524
Rodents. Our model identified 76 rodent species with high zoonotic capacity for SARS-CoV-2,
some of which thrive in human-altered settings. Among these, the deer mouse (*Peromyscus*
maniculatus) and the white-footed mouse (*P. leucopus*) showed high probabilities. These are
among the most well-studied mammals in North America, in part due to their status as zoonotic
reservoirs for multiple zoonotic pathogens and parasites [116–118]. Experimental infection, viral
shedding, and sustained intraspecific transmission of SARS-CoV-2 were recently confirmed for
P. maniculatus [51,52], but similar studies have not been conducted for *P. leucopus*, which is
widely distributed across the eastern United States and Mexico.
533
Our model predicted low zoonotic capacity for *Mus musculus* (0.11), corresponding with
in vivo experiments suggesting this species is not susceptible to infection by the initial human
variant of SARS-CoV-2[19], although notably, more recent experiments have confirmed the
susceptibility of *M. musculus* to two newer human-derived variants [20]. Also in the top 10%
were two rodent species considered to be human commensals whose geographic ranges are

expanding due to human activities: *Rattus argentiventer* (0.84) and *R. tiomanicus* (0.79)
(Supplementary File 1) [119–121]. Additional common rodent species with relatively high
probabilities of zoonotic capacity include domesticated guinea pigs (*Cavia porcellus*), gerbils
(*Gerbillus gerbillus*, *Meriones tristrami*), and several common mouse species (*Apodemus
peninsulae*, *A. flavicollis*, and *A. sylvaticus*), all of which are known reservoirs for other zoonotic
diseases [122–124]. It is notable that many of these rodent species are regularly preyed upon
by carnivore species, such as the red fox (*Vulpes vulpes*) or domestic cats (*Felis catus*) who
themselves were predicted to have high zoonotic capacity for SARS-CoV-2 by our model.
547
Species with large geographic ranges. With sufficient opportunity for infectious contact, the risk
of zoonotic spillback transmission increases with SARS-CoV-2 prevalence in human
populations. Among species with high model-predicted zoonotic capacity, there were several
relatively common species with very large geographic ranges or synanthropic tendencies that
overlap with global hotspots of COVID-19 in people (Figure 6, Supplementary File 2). Notable
species that are widely distributed across much of the northern hemisphere include the red fox
(*Vulpes vulpes*, ~50 countries), the European polecat (*Mustela putorius*), the raccoon dog
(*Nyctereutes procyonoides*), stoat (*Mustela erminea*) and wolf (*Canis lupus*). White-tailed deer
(*Odocoileus virginianus*) are among the most geographically widespread species across Latin
American countries with high SARS-CoV-2 prevalence. Globally, South and Southeast Asia had
the highest diversity of mammal species with high predicted zoonotic capacity for SARS-CoV-2
(~90 species). Notable examples in this region include both rodents and bats. For example,
Finlayson's squirrel (*Callosciurus finlaysonii*) is native to Mainland Southeast Asia, but
introductions via the pet trade in Europe have led to invasive populations in multiple countries
[125]. Hunting has been documented for numerous bat species with geographic ranges across
Southeast Asia (e.g., *Cheiromeles torquatus*, *Cynopterus brachyotis*, *Rousettus
amplexicaudatus*, *Macroglossus minimus*) [126,127], and there were multiple additional bat
species in the 90th percentile from Asia and Africa where bats are subject to hunting pressure
and from which other betacoronaviruses have been identified [107,128]. There were also
several wide-ranging species whose contact with humans are limited to specialized settings. For
instance, biologists and wildlife managers handle live individuals for research purposes,
including grizzly bear (*Ursus arctos*), polar bear (*Ursus maritimus*), and wolf (*Canis lupus*), all of
which are in the 89th percentile or above for predicted zoonotic capacity to SARS-CoV-2.
571
Other high priority mammal species. Species with more equivocal predictions about zoonotic
capacity that are in frequent contact with humans warrant further investigation. For instance,
while species such as horses (*Equus caballus*), goats (*Capra hircus*), and guinea pigs (*Cavia
porcellus*) are not in the top 10% of predicted zoonotic capacity, due to the nature of their
contact with humans they may experience greater risks of spillback infection, or pose a greater
risk to humans for secondary spillover infection compared to many wild species. Conversely,
while certain endangered or nearly extinct species are predicted to have relatively high zoonotic
capacity, they may have fewer opportunities for human contact. For species of conservation
concern, spillback transmission of SARS-CoV-2 from humans presents an important source of
risk [28,129], particularly for populations that are under active management, including *ex situ*

management such as captive breeding. These species include the scimitar-horned oryx (*Oryx*
dammah), addax (*Addax nasomaculatus*), some Antarctic fauna, and mountain gorillas (*Gorilla*
beringei) in which SARS-CoV-2 spillback infection may occur through close-proximity eco-
tourism activities [130,131]. Indeed, spillback transmission of SARS-CoV-2 has already been
confirmed in a closely related species, the Western lowland gorilla (*Gorilla gorilla*) in captivity
[132], leading to the vaccination of bonobos and orangutans with an experimental COVID-19
vaccine [133]. These species may benefit from focused risk mitigation efforts, such as those
enacted recently to protect endangered black-footed ferrets (*Mustela nigripes*) from potential
SARS-CoV-2 spillback [134].
591
All fifteen species of *Tupaia* treeshrews were predicted by our model to have medium to
high probability (ranging from 0.62 to 0.87). One species, *T. belangeri*, has been explored as a
potential lab model for several human infectious diseases including SARS-CoV-2 [135] but
relative to other treeshrews, our model assigned only medium probability for SARS-CoV-2
zoonotic capacity in this species (0.67). This result matches lab studies reporting asymptomatic
infection and low viral shedding in *T. belangeri* [54]. In contrast, the common treeshrew (*T. glis*)
was in the 94th percentile of zoonotic capacity (0.87 probability). These two species are
sympatric in parts of their range, exist in close proximity to humans, and also overlap
geographically with COVID-19 hotspots in Southeast Asia, suggesting the possibility of spillover
transmission among congeners if spillback transmission occurs from humans to these species.
602
Strengthening predictive capacity for zoonoses. While there was wide agreement between our
model predictions and empirical studies, examining biases and mismatches between
experimental results and model-generated predictions will focus research attention on
characterizing what factors underlie the disconnects between predicted and observed zoonotic
capacity. For instance, this study along with multiple other computational and experimental
studies predicted that pigs (*Sus scrofa*) would be susceptible to SARS-CoV-2 (Figure 1), but this
prediction has not been supported by results from whole animal inoculations [37,53].
610
Disconnects between real-world observations, *in vivo* experimental results, and *in silico*
predictions of zoonotic capacity may arise because host susceptibility and transmission capacity
are necessary but not sufficient for zoonotic risk to be realized in natural settings. These
processes are embedded in a broader ecological context that impacts host susceptibility, intra-
host infection dynamics (latency, recrudescence, tolerance), and viral persistence that
collectively determine where and when spillover will occur [136–139]. These processes also
depend strongly on the cellular environments in which cell entry and viral replication take place
(e.g., the presence of key proteases, [7]), and on host immunogenicity [139], factors which are
themselves influenced by the environment [140]. Insofar as data limitations preclude perfect
computational predictions of zoonotic capacity (e.g., limited ACE2 sequences, crystal structures,
or species trait data), laboratory experiments are also limited in assessing true zoonotic
capacity. For SARS-CoV-2 and other host-pathogen systems, animals that are readily infected
in the lab appear to be less susceptible in non-lab settings (ferrets in the lab vs. mixed results in
ferrets as pets [36,53,141]; rabbits in the lab vs. rabbits as pets [48,142]. Moreover, wildlife

hosts confirmed to shed multiple zoonotic viruses in natural settings (e.g., bats, [143]) can be
much less tractable for whole-animal laboratory investigations (for instance, requiring high
biosecurity containment and very limited sample sizes in unnatural settings). While laboratory
experiments are critical for understanding mechanisms of pathogenesis and disease, without
field surveillance and population-level studies they offer imperfect reflections of zoonotic
capacity in the natural world.
631

These examples illustrate that there is no single methodology sufficient to understand
and predict zoonotic transmission, for SARS-CoV-2 or any zoonotic pathogen. They also
demonstrate the need for improved coordination among theoretical and statistical models, lab
work, and field work to improve zoonotic predictive capacity [144], and to create new linkages to
underutilized data sources such as natural history collections, which are well-positioned to
augment basic knowledge gaps about the spatial and temporal extents of animal hosts and their
pathogens [145,146]. Integration of multiple methodologies and data streams across biological
scales offers avenues to more efficient iteration between computational predictions, laboratory
experiments, and targeted animal surveillance that will better link transmission mechanisms to
the broader conditions underpinning zoonotic disease emergence in nature.
642
643

Acknowledgments

We are grateful for discussions with Drs. Alexandre Bonvin, Dennis Bente, Susan
Hafenstein, Kathryn Hanley, Hyunwook Lee, Colin Parrish, and John Paul Schmidt about
various components of this project. This work was supported by the NSF EEID program (DEB
1717282), DARPA PREEMPT program (D18AC00031), CREATE-NEO, a member of the NIH
NIAID CREID program (1U01 AI151807-01), and the NVIDIA Corporation GPU grant program
(BAH); by the NSF Polar program (OPP 1935870, 1947040) (AV); and by NIH NIGMS
(R35GM122543) (JPGLMR).

652
653

654   **Competing interests**

655   The authors declare no competing interests.
656

References

1.   Dong E, Du H, Gardner L. 2020 An interactive web-based dashboard to track COVID-19 in
real time. *Lancet Infect. Dis.* **20**, 533–534. (doi:10.1016/S1473-3099(20)30120-1)

2.   WHO. 2021 WHO coronavirus disease (COVID-19) dashboard.

3.   Keele BF *et al.* 2006 Chimpanzee reservoirs of pandemic and nonpandemic HIV-1. *Science*
313, 523–526. (doi:10.1126/science.1126531)

4.   Gage KL, Kosoy MY. 2005 Natural history of plague: perspectives from more than a
century of research. *Annu. Rev. Entomol.* **50**, 505–528.
(doi:10.1146/annurev.ento.50.071803.130337)

5.   Taubenberger JK, Reid AH, Lourens RM, Wang R, Jin G, Fanning TG. 2005
Characterization of the 1918 influenza virus polymerase genes. *Nature* **437**, 889.
(doi:10.1038/nature04230)

6.   Zhou P *et al.* 2020 A pneumonia outbreak associated with a new coronavirus of probable
bat origin. *Nature* **579**, 270–273. (doi:10.1038/s41586-020-2012-7)

7.   Letko M, Marzi A, Munster V. 2020 Functional assessment of cell entry and receptor usage
for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol* **5**, 562–569.
(doi:10.1038/s41564-020-0688-y)

8.   Chou C-F *et al.* 2006 ACE2 orthologues in non-mammalian vertebrates (Danio, Gallus,
Fugu, Tetraodon and Xenopus). *Gene* **377**, 46–55. (doi:10.1016/j.gene.2006.03.010)

9.   Guth S, Visher E, Boots M, Brook CE. 2019 Host phylogenetic distance drives trends in
virus virulence and transmissibility across the animal-human interface. *Philos. Trans. R.*
Soc. Lond. B Biol. Sci. **374**, 20190296. (doi:10.1098/rstb.2019.0296)

10. WHO. 2020 SARS-CoV-2 mink-associated variant strain – Denmark.

11. Oude Munnink BB *et al.* 2020 Transmission of SARS-CoV-2 on mink farms between
humans and mink and back to humans. *Science* (doi:10.1126/science.abe5901)

12. Garry RF. 2021 Mutations arising in SARS-CoV-2 spike on sustained human-to-human
transmission and human-to-animal passage. *Virological*. See
https://virological.org/t/mutations-arising-in-sars-cov-2-spike-on-sustained-human-to-
human-transmission-and-human-to-animal-passage/578 (accessed on 28 January 2021).

13. Rodrigues JPGLM, Barrera-Vilarmau S, M C Teixeira J, Sorokina M, Seckel E, Kastritis PL,
Levitt M. 2020 Insights on cross-species transmission of SARS-CoV-2 from structural
modeling. *PLoS Comput. Biol.* **16**, e1008449. (doi:10.1371/journal.pcbi.1008449)

14. Davies NG *et al.* 2020 Estimated transmissibility and severity of novel SARS-CoV-2 Variant
of Concern 202012/01 in England. *medRxiv* , 2020.12.24.20248822.
(doi:10.1101/2020.12.24.20248822)

15. Volz E *et al.* 2021 Transmission of SARS-CoV-2 Lineage B.1.1.7 in England: Insights from
linking epidemiological and genetic data. *medRxiv* , 2020.12.30.20249034.
(doi:10.1101/2020.12.30.20249034)

16. Rambaut A *et al.* 2020 Preliminary genomic characterisation of an emergent SARS-CoV-2
lineage in the UK defined by a novel set of spike mutations. *Virological*. See
https://virological.org/t/preliminary-genomic-characterisation-of-an-emergent-sars-cov-2-
lineage-in-the-uk-defined-by-a-novel-set-of-spike-mutations/563 (accessed on 28 January
2021).

17. Tegally H *et al.* 2020 Emergence and rapid spread of a new severe acute respiratory
syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in
South Africa. *medRxiv* , 2020.12.21.20248640. (doi:10.1101/2020.12.21.20248640)

18. Van Egeren D *et al.* 2021 Risk of rapid evolutionary escape from biomedical interventions
targeting SARS-CoV-2 spike protein. *PLoS One* **16**, e0250780.
(doi:10.1371/journal.pone.0250780)

19. Bao L *et al.* 2020 The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. *Nature*
583, 830–833. (doi:10.1038/s41586-020-2312-y)

20. Montagutelli X *et al.* 2021 The B1.351 and P.1 variants extend SARS-CoV-2 host range to
mice. *bioRxiv*. , 2021.03.18.436013. (doi:10.1101/2021.03.18.436013)

21. DeLiberto T, Shriner S. 2020 ProMED.

22. ODA. 2020 Mink at affected Oregon farm negative for SARS-CoV-2, wildlife surveillance
continues. 23 December. See https://odanews.wpengine.com/mink-at-affected-oregon-
farm-negative-for-sars-cov-2-wildlife-surveillance-continues/.

23. Shriner S *et al.* 2021 SARS-CoV-2 Exposure in Escaped Mink, Utah, USA. *Emerging*
Infectious Disease journal **27**. (doi:10.3201/eid2703.204444)

24. Lam SD *et al.* 2020 SARS-CoV-2 spike protein predicted to form complexes with host
receptor protein orthologues from a broad range of mammals. *Sci. Rep.* **10**, 16471.
(doi:10.1038/s41598-020-71936-5)

25. Liu Z *et al.* 2020 Composition and divergence of coronavirus spike proteins and host ACE2
receptors predict potential intermediate hosts of SARS-CoV-2. *J. Med. Virol.* **92**, 595–601.
(doi:10.1002/jmv.25726)

26. Luan J, Jin X, Lu Y, Zhang L. 2020 SARS-CoV-2 spike protein favors ACE2 from Bovidae
and Cricetidae. *J. Med. Virol.*

27. Mathavarajah S, Stoddart AK, Gagnon GA, Dellaire G. 2020 Pandemic danger to the deep:
the risk of marine mammals contracting SARS-CoV-2 from wastewater. ,
2020.08.13.249904. (doi:10.1101/2020.08.13.249904)

28. Melin AD, Janiak MC, Marrone F 3rd, Arora PS, Higham JP. 2020 Comparative ACE2
variation and primate COVID-19 risk. *Commun Biol* **3**, 641. (doi:10.1038/s42003-020-
01370-w)

29. Kumar A, Pandey SN, Pareek V, Narayan RK, Faiq MA, Kumari C. 2020 Predicting
susceptibility for SARS-CoV-2 infection in domestic and wildlife animals using ACE2 protein
sequence homology. *Zoo Biol.* (doi:10.1002/zoo.21576)

30. Huang X, Zhang C, Pearce R, Omenn GS, Zhang Y. 2020 Identifying the Zoonotic Origin of
SARS-CoV-2 by Modeling the Binding Affinity between the Spike Receptor-Binding Domain
and Host ACE2. *J. Proteome Res.* **19**, 4844–4856. (doi:10.1021/acs.jproteome.0c00717)

31. Ahmed R, Hasan R, Siddiki AMAMZ, Islam MS. 2021 Host range projection of SARS-CoV-
2: South Asia perspective. *Infect. Genet. Evol.* **87**, 104670.
(doi:10.1016/j.meegid.2020.104670)

32. Damas J *et al.* 2020 Broad host range of SARS-CoV-2 predicted by comparative and
structural analysis of ACE2 in vertebrates. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 22311–
22322. (doi:10.1073/pnas.2010146117)

33. Ulrich L, Wernike K, Hoffmann D, Mettenleiter TC, Beer M. 2020 Experimental Infection of
Cattle with SARS-CoV-2. *Emerg. Infect. Dis.* **26**, 2979–2981. (doi:10.3201/eid2612.203799)

34. Sit THC *et al.* 2020 Infection of dogs with SARS-CoV-2. *Nature* **586**, 776–778.
(doi:10.1038/s41586-020-2334-5)

35. USDA. 2020 Cases of SARS-CoV-2 in animals in the United States. See
https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/sa_one_health/sars-cov-2-
animals-us (accessed on 12 January 2021).

36. OIE. 2021 Events in animals: OIE - World Organisation for Animal Health. See
https://www.oie.int/en/scientific-expertise/specific-information-and-
recommendations/questions-and-answers-on-2019novel-coronavirus/events-in-animals/
(accessed on 28 January 2021).

37. Shi J *et al.* 2020 Susceptibility of ferrets, cats, dogs, and other domesticated animals to
SARS-coronavirus 2. *Science* **368**, 1016–1020. (doi:10.1126/science.abb7015)

38. Hamer SA *et al.* 2021 SARS-CoV-2 Infections and Viral Isolations among Serially Tested
Cats and Dogs in Households with Infected Owners in Texas, USA. *Viruses* **13**.
(doi:10.3390/v13050938)

39. Woolsey C *et al.* 2020 Establishment of an African green monkey model for COVID-19.
bioRxiv , 2020.05.17.100289. (doi:10.1101/2020.05.17.100289)

40. Hall JS *et al.* 2020 Experimental challenge of a North American bat species, big brown bat
(Eptesicus fuscus), with SARS-CoV-2. *Transbound. Emerg. Dis.* (doi:10.1111/tbed.13949)

41. Zhang Q *et al.* 2020 SARS-CoV-2 neutralizing serum antibodies in cats: a serological
investigation. , 2020.04.01.021196. (doi:10.1101/2020.04.01.021196)

42. San Diego Zoo. 2021 Gorilla Troop at the San Diego Zoo Safari Park Test Positive for
COVID-19. See https://zoo.sandiegozoo.org/pressroom/news-releases/gorilla-troop-san-
diego-zoo-safari-park-test-positive-covid-19 (accessed on 28 January 2021).

43. Rockx B *et al.* 2020 Comparative pathogenesis of COVID-19, MERS, and SARS in a

nonhuman primate model. *Science* **368**, 1012–1015. (doi:10.1126/science.abb7314)

44. Munster VJ *et al.* 2020 Respiratory disease in rhesus macaques inoculated with SARS-
CoV-2. *Nature* **585**, 268–272. (doi:10.1038/s41586-020-2324-7)

45. Sia SF *et al.* 2020 Pathogenesis and transmission of SARS-CoV-2 in golden hamsters.
Nature **583**, 834–838. (doi:10.1038/s41586-020-2342-5)

46. Oreshkova N *et al.* 2020 SARS-CoV-2 infection in farmed minks, the Netherlands, April and
May 2020. *Euro Surveill.* **25**. (doi:10.2807/1560-7917.ES.2020.25.23.2001005)

47. Freuling CM *et al.* 2020 Susceptibility of Raccoon Dogs for Experimental SARS-CoV-2
Infection. *Emerg. Infect. Dis.* **26**, 2982–2985. (doi:10.3201/eid2612.203733)

48. Mykytyn AZ *et al.* 2021 Susceptibility of rabbits to SARS-CoV-2. *Emerg. Microbes Infect.*
10, 1–7. (doi:10.1080/22221751.2020.1868951)

49. Bartlett SL *et al.* 2021 SARS-COV-2 INFECTION AND LONGITUDINAL FECAL
SCREENING IN MALAYAN TIGERS (PANTHERA TIGRIS JACKSONI), AMUR TIGERS
(PANTHERA TIGRIS ALTAICA), AND AFRICAN LIONS (PANTHERA LEO KRUGERI) AT
THE BRONX ZOO, NEW YORK, USA. *J. Zoo Wildl. Med.* **51**, 733–744. (doi:10.1638/2020-
0171)

50. Wang L *et al.* 2020 Complete Genome Sequence of SARS-CoV-2 in a Tiger from a U.S.
Zoological Collection. *Microbiol Resour Announc* **9**. (doi:10.1128/MRA.00468-20)

51. Fagre A *et al.* 2021 SARS-CoV-2 infection, neuropathogenesis and transmission among
deer mice: Implications for spillback to New World rodents. *PLoS Pathog.* **17**, e1009585.
(doi:10.1371/journal.ppat.1009585)

52. Griffin BD *et al.* 2021 SARS-CoV-2 infection and transmission in the North American deer
mouse. *Nat. Commun.* **12**, 3612. (doi:10.1038/s41467-021-23848-9)

53. Schlottau K *et al.* 2020 SARS-CoV-2 in fruit bats, ferrets, pigs, and chickens: an
experimental transmission study. *Lancet Microbe* **1**, e218–e225. (doi:10.1016/S2666-
5247(20)30089-6)

54. Zhao Y *et al.* 2020 Susceptibility of tree shrew to SARS-CoV-2 infection. *Sci. Rep.* **10**,
16007. (doi:10.1038/s41598-020-72563-w)

55. Louisville Zoo. 2020 Louisville Zoo Female Snow Leopard Tests Positive for SARS-CoV-2.
See https://louisvillezoo.org/louisville-zoo-female-snow-leopard-tests-positive-for-sars-cov-
2-media-release/ (accessed on 28 January 2021).

56. Ulrich L, Michelitsch A, Halwe N, Wernike K, Hoffmann D, Beer M. 2021 Experimental
SARS-CoV-2 Infection of Bank Voles. *Emerg. Infect. Dis.* **27**, 1193–1195.
(doi:10.3201/eid2704.204945)

57. Georgia Aquarium. In press. Asian Small-Clawed Otters at Georgia Aquarium Test Positive
for COVID-19. See http://news.georgiaaquarium.org/stories/releases-20210418 (accessed
on 13 May 2021).

58. Palmer MV *et al.* 2021 Susceptibility of white-tailed deer (Odocoileus virginianus) to SARS-
CoV-2. *bioRxiv.* , 2021.01.13.426628. (doi:10.1101/2021.01.13.426628)

59. Gryseels S, De Bruyn L, Gyselings R, Calvignac-Spencer S, Leendertz FH, Leirs H. 2020
Risk of human-to-wildlife transmission of SARS-CoV-2. *Mamm. Rev.* **8**, e00373-17.
(doi:10.1111/mam.12225)

60. Deng W *et al.* 2020 Rhesus macaques can be effectively infected with SARS-CoV-2 via
ocular conjunctival route. *bioRxiv.* , 2020.03.13.990036. (doi:10.1101/2020.03.13.990036)

61. Rodrigues JPGLM *et al.* 2013 Defining the limits of homology modeling in information-
driven protein docking. *Proteins* **81**, 2119–2128. (doi:10.1002/prot.24382)

62. Sander C, Schneider R. 1991 Database of homology-derived protein structures and the
structural meaning of sequence alignment. *Proteins* **9**, 56–68.
(doi:10.1002/prot.340090107)

63. Li Y *et al.* 2020 SARS-CoV-2 and Three Related Coronaviruses Utilize Multiple ACE2
Orthologs and Are Potently Blocked by an Improved ACE2-Ig. *J. Virol.* **94**.
(doi:10.1128/JVI.01283-20)

64. Fournier D, Luft FC, Bader M, Ganten D, Andrade-Navarro MA. 2012 Emergence and
evolution of the renin-angiotensin-aldosterone system. *J. Mol. Med.* **90**, 495–508.
(doi:10.1007/s00109-012-0894-z)

65. Han BA, Schmidt JP, Bowden SE, Drake JM. 2015 Rodent reservoirs of future zoonotic
diseases. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 7039–7044. (doi:10.1073/pnas.1501598112)

66. Yang LH, Han BA. 2018 Data-driven predictions and novel hypotheses about zoonotic tick
vectors from the genus Ixodes. *BMC Ecol.* **18**, 7. (doi:10.1186/s12898-018-0163-2)

67. Han BA, O'Regan SM, Paul Schmidt J, Drake JM. 2020 Integrating data mining and
transmission theory in the ecology of infectious diseases. *Ecol. Lett.* **23**, 1178–1188.
(doi:10.1111/ele.13520)

68. Han BA, Schmidt JP, Alexander LW, Bowden SE, Hayman DTS, Drake JM. 2016
Undiscovered Bat Hosts of Filoviruses. *PLoS Negl. Trop. Dis.* **10**, e0004815.
(doi:10.1371/journal.pntd.0004815)

69. Han BA *et al.* 2019 Confronting data sparsity to identify potential sources of Zika virus
spillover infection among primates. *Epidemics* **27**, 59–65.
(doi:10.1016/j.epidem.2019.01.005)

70. Yang X-L *et al.* 2017 Genetically Diverse Filoviruses in Rousettus and Eonycteris spp. Bats,
China, 2009 and 2015. *Emerg. Infect. Dis.* **23**, 482–486. (doi:10.3201/eid2303.161119)

71. Goldstein T *et al.* 2018 The discovery of Bombali virus adds further support for bats as
hosts of ebolaviruses. *Nat Microbiol* **3**, 1084–1089. (doi:10.1038/s41564-018-0227-2)

72. Sorokina M, M C Teixeira J, Barrera-Vilarmau S, Paschke R, Papasotiriou I, Rodrigues
JPGLM, Kastritis PL. 2020 Structural models of human ACE2 variants with SARS-CoV-2
Spike protein for structure-based drug design. *Sci Data* **7**, 309. (doi:10.1038/s41597-020-

00652-6)

73. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990 Basic local alignment search
tool. *J. Mol. Biol.* **215**, 403–410. (doi:10.1016/S0022-2836(05)80360-2)

74. Winter D. 2017 rentrez: An R package for the NCBI eUtils API. *R J.* **9**, 520.
(doi:10.32614/rj-2017-058)

75. Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. 2018 The MEROPS
database of proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison
with peptidases in the PANTHER database. *Nucleic Acids Res.* **46**, D624–D632.
(doi:10.1093/nar/gkx1134)

76. Katoh K, Misawa K, Kuma K-I, Miyata T. 2002 MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066.
(doi:10.1093/nar/gkf436)

77. Lan J *et al.* 2020 Structure of the SARS-CoV-2 spike receptor-binding domain bound to the
ACE2 receptor. *Nature* **581**, 215–220. (doi:10.1038/s41586-020-2180-5)

78. Webb B, Sali A. 2016 Comparative Protein Structure Modeling Using MODELLER. *Curr.*
Protoc. Bioinformatics **54**, 5.6.1–5.6.37. (doi:10.1002/cpbi.3)

79. Sali A, Blundell TL. 1993 Comparative protein modelling by satisfaction of spatial restraints.
J. Mol. Biol. **234**, 779–815. (doi:10.1006/jmbi.1993.1626)

80. van Zundert GCP *et al.* 2016 The HADDOCK2.2 Web Server: User-Friendly Integrative
Modeling of Biomolecular Complexes. *J. Mol. Biol.* **428**, 720–725.
(doi:10.1016/j.jmb.2015.09.014)

81. de Magalhães JP, Costa J. 2009 A database of vertebrate longevity records and their
relation to other life-history traits. *J. Evol. Biol.* **22**, 1770–1774. (doi:10.1111/j.1420-
9101.2009.01783.x)

82. Myhrvold NP, Baldridge E, Chan B, Sivam D, Freeman DL, Ernest SKM. 2015 An amniote
life-history database to perform comparative analyses with birds, mammals, and reptiles:
Ecological ArchivesE096-269. *Ecology* **96**, 3109–3000. (doi:10.1890/15-0846r.1)

83. Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. 2014
EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. *Ecology*
95, 2027–2027. (doi:10.1890/13-1917.1)

84. Dallas T, Park AW, Drake JM. 2017 Predicting cryptic links in host-parasite networks. *PLoS*
Comput. Biol. **13**, e1005557. (doi:10.1371/journal.pcbi.1005557)

85. Baker C. 2018 *wosr: Clients to the 'Web of Science' and 'InCites' APIs*. See
https://CRAN.R-project.org/package=wosr.

86. Bosco-Lauth AM *et al.* 2020 Experimental infection of domestic dogs and cats with SARS-
CoV-2: Pathogenesis, transmission, and response to reexposure in cats. *Proc. Natl. Acad.*
Sci. U. S. A. **117**, 26382–26388. (doi:10.1073/pnas.2013102117)

87. Elith J, Leathwick JR, Hastie T. 2008 A working guide to boosted regression trees. *J. Anim.*
Ecol. **77**, 802–813.

88. Jones KE *et al.* 2009 PanTHERIA: a species-level database of life history, ecology, and
geography of extant and recently extinct mammals: Ecological Archives E090-184. *Ecology*
90, 2648–2648. (doi:10.1890/08-1494.1)

89. Wilson DE, Reeder DM. 2005 *Mammal Species of the World: A Taxonomic and Geographic*
Reference. Baltimore, MD: JHU Press. See
http://books.google.com/books?id=JgAMbNSt8ikC.

90. Greenwell B, Boehmke B, Cunningham J, Developers GBM. 2020 *Generalized Boosted*
Regression Models. Comprehensive R Archive Network (CRAN). See https://cran.r-
project.org/web/packages/gbm/index.html.

91. R Core Team. 2020 *R: A language and environment for statistical computing*. Vienna,
Austria. See http://www.R-project.org/.

92. Wilkins AS, Wrangham RW, Fitch WT. 2014 The 'domestication syndrome' in mammals: a
unified explanation based on neural crest cell behavior and genetics. *Genetics* **197**, 795–
808. (doi:10.1534/genetics.114.165423)

93. Cleaveland S, Laurenson MK, Taylor LH. 2001 Diseases of humans and their domestic
mammals: pathogen characteristics, host range and the risk of emergence. *Philos. Trans.*
R. Soc. Lond. B Biol. Sci. **356**, 991–999. (doi:10.1098/rstb.2001.0889)

94. Database MD. 2020 *Mammal Diversity Database*. (doi:10.5281/zenodo.4139818)

95. IUCN. 2020 The IUCN Red List of Threatened Species.

96. Liu Y *et al.* 2021 Functional and genetic analysis of viral receptor ACE2 orthologs reveals a
broad potential host range of SARS-CoV-2. *Proc. Natl. Acad. Sci. U. S. A.* **118**.
(doi:10.1073/pnas.2025373118)

97. Pitra C, Schwarz S, Fickel J. 2010 Going west—invasion genetics of the alien raccoon dog
Nyctereutes procnoides in Europe. *Eur. J. Wildl. Res.* **56**, 117–129. (doi:10.1007/s10344-
009-0283-2)

98. Milla R *et al.* 2018 Phylogenetic patterns and phenotypic profiles of the species of plants
and mammals farmed for food. *Nat Ecol Evol* **2**, 1808–1817. (doi:10.1038/s41559-018-
0690-4)

99. Kevany S. 2020 Danish Covid mink cull and future disease fears will kill fur trade, say
farmers. *The Guardian*, 6 November. See
http://www.theguardian.com/environment/2020/nov/06/danish-covid-mink-cull-and-future-
disease-fears-will-kill-fur-trade-say-farmers.

100.Can ÖE, D'Cruze N, Macdonald DW. 2019 Dealing in deadly pathogens: Taking stock of
the legal trade in live wildlife and potential risks to human health. *Glob Ecol Conserv* **17**,
e00515. (doi:10.1016/j.gecco.2018.e00515)

101.Lam TT-Y *et al.* 2020 Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins.

Nature **583**, 282–285. (doi:10.1038/s41586-020-2169-0)

102. Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. 2020 The proximal origin of
SARS-CoV-2. *Nat. Med.* **26**, 450–452. (doi:10.1038/s41591-020-0820-9)

103. Lehmann D, Halbwax ML, Makaga L, Whytock R, Ndindiwe Malata L, Bombenda Mouele
W, Momboua BR, Koumba Pambo AF, White LJT. 2020 Pangolins and bats living together
in underground burrows in Lopé National Park, Gabon. *Afr. J. Ecol.* (doi:10.1111/aje.12759)

104. Tsuda S *et al.* 2012 Genomic and serological detection of bat coronavirus from bats in the
Philippines. *Arch. Virol.* **157**, 2349–2355. (doi:10.1007/s00705-012-1410-z)

105. Olival KJ *et al.* 2020 Possibility for reverse zoonotic transmission of SARS-CoV-2 to free-
ranging wildlife: A case study of bats. *PLoS Pathog.* **16**, e1008758.
(doi:10.1371/journal.ppat.1008758)

106. Anthony SJ *et al.* 2013 Coronaviruses in bats from Mexico. *J. Gen. Virol.* **94**, 1028–1038.
(doi:10.1099/vir.0.049759-0)

107. Anthony SJ *et al.* 2017 Global patterns in coronavirus diversity. *Virus Evol* **3**, vex012.
(doi:10.1093/ve/vex012)

108. Zhou H *et al.* 2020 A Novel Bat Coronavirus Closely Related to SARS-CoV-2 Contains
Natural Insertions at the S1/S2 Cleavage Site of the Spike Protein. *Curr. Biol.* **30**, 2196–
2203.e3. (doi:10.1016/j.cub.2020.05.023)

109. Hul V *et al.* 2021 A novel SARS-CoV-2 related coronavirus in bats from Cambodia. *bioRxiv.*
, 2021.01.26.428212. (doi:10.1101/2021.01.26.428212)

110. Mou H *et al.* 2020 Mutations from bat ACE2 orthologs markedly enhance ACE2-Fc
neutralization of SARS-CoV-2. *bioRxiv.* , 2020.06.29.178459.
(doi:10.1101/2020.06.29.178459)

111. Wacharapluesadee S *et al.* 2021 Evidence for SARS-CoV-2 related coronaviruses
circulating in bats and pangolins in Southeast Asia. *Nat. Commun.* **12**, 972.
(doi:10.1038/s41467-021-21240-1)

112. Pulliam JRC *et al.* 2012 Agricultural intensification, priming for persistence and the
emergence of Nipah virus: a lethal bat-borne zoonosis. *J. R. Soc. Interface* **9**, 89–101.
(doi:10.1098/rsif.2011.0223)

113. Plowright RK *et al.* 2015 Ecological dynamics of emerging bat virus spillover. *Proceedings
of the Royal Society B.* **282**, 20142124. (doi:10.1098/rspb.2014.2124)

114. Kessler MK *et al.* 2018 Changing resource landscapes and spillover of henipaviruses. *Ann.
N. Y. Acad. Sci.* **1429**, 78–99. (doi:10.1111/nyas.13910)

115. Peel AJ, Field HE, Aravena MR, Edson D, McCallum H, Plowright RK, Prada D. 2020
Coronaviruses and Australian bats: a review in the midst of a pandemic. *Aust. J. Zool.*
(doi:10.1071/ZO20046)

116. Bordes F, Blasdell K, Morand S. 2015 Transmission ecology of rodent-borne diseases:

New frontiers. *Integr. Zool.* **10**, 424–435. (doi:10.1111/1749-4877.12149)

117. Ostfeld RS, Canham CD, Oggenfuss K, Winchcombe RJ, Keesing F. 2006 Climate, deer,
rodents, and acorns as determinants of variation in lyme-disease risk. *PLoS Biol.* **4**, e145.
(doi:10.1371/journal.pbio.0040145)

118. Machtinger ET, Williams SC. 2020 Practical Guide to Trapping Peromyscus leucopus
(Rodentia: Cricetidae) and Peromyscus maniculatus for Vector and Vector-Borne Pathogen
Surveillance and Ecology. *J. Insect Sci.* **20**. (doi:10.1093/jisesa/ieaa028)

119. Morand S *et al.* 2015 Global parasite and Rattus rodent invasions: The consequences for
rodent-borne diseases. *Integr. Zool.* **10**, 409–423. (doi:10.1111/1749-4877.12143)

120. Hamdan NES, Ng YL, Lee WB, Tan CS, Khan FAA, Chong YL. 2017 Rodent Species
Distribution and Hantavirus Seroprevalence in Residential and Forested areas of Sarawak,
Malaysia. *Trop Life Sci Res* **28**, 151–159. (doi:10.21315/tlsr2017.28.1.11)

121. Louys J, Herrera MB, Thomson VA, Wiewel AS, Donnellan SC, O'Connor S, Aplin K. 2020
Expanding population edge craniometrics and genetics provide insights into dispersal of
commensal rats through Nusa Tenggara, Indonesia. *Rec. Aust. Mus.* **72**, 287–302.
(doi:10.3853/j.2201-4349.72.2020.1730)

122. Tadin A *et al.* 2016 Molecular Survey of Zoonotic Agents in Rodents and Other Small
Mammals in Croatia. *Am. J. Trop. Med. Hyg.* **94**, 466–473. (doi:10.4269/ajtmh.15-0517)

123. Yousefi A, Eslami A, Mobedi I, Rahbari S, Ronaghi H. 2014 Helminth Infections of House
Mouse (Mus musulus) and Wood Mouse (Apodemus sylvaticus) from the Suburban Areas
of Hamadan City, Western Iran. *Iran. J. Parasitol.* **9**, 511–518.

124. Rahman MM, Yoon KB, Lim SJ, Jeon MG, Kim HJ, Kim HY, Cho JY, Chae HM, Park YC.
2017 Molecular detection by analysis of the 16S rRNA gene of fecal coliform bacteria from
the two Korean Apodemus species (Apodemus agrarius and A. peninsulae). *Genet. Mol.*
Res. **16**. (doi:10.4238/gmr16029510)

125. Bertolino S, Lurz PWW. 2013 Callosciurussquirrels: worldwide introductions, ecological
impacts and recommendations to prevent the establishment of new invasive populations:
Worldwide introductions ofCallosciurussquirrels. *Mamm. Rev.* **43**, 22–33.
(doi:10.1111/j.1365-2907.2011.00204.x)

126. Mildenstein T, Tanshi I, Racey PA. 2016 Exploitation of Bats for Bushmeat and Medicine. In
Bats in the Anthropocene: Conservation of Bats in a Changing World (eds CC Voigt, T
Kingston), pp. 325–375. Cham: Springer International Publishing. (doi:10.1007/978-3-319-
25220-9_12)

127. Ransaleleh TA, Nangoy MJ, Wahyuni I, Lomboan A, Koneri R, Saputro S, Pamungkas J,
Latinne A. 2020 Identification of bats on traditional market in dumoga district, North
Sulawesi. *IOP Conf. Ser.: Earth Environ. Sci.* **473**, 012067. (doi:10.1088/1755-
1315/473/1/012067)

128. Tampon NVT, Rabaya YMC, Malbog KMA, Burgos SC, Libre K Jr, Valila ASD, Achondo
MJMM, Onggo LS, Murao LAE. 2020 First molecular evidence for bat betacoronaviruses in

996   Mindanao. *Philipp J. Sci.* **149**, 91–94.

997 129.Logeot M, Mauroy A, Thiry E, De Regge N, Vervaeke M, Beck O, De Waele V, Van den
998   Berg T. 2021 Risk assessment of SARS-CoV-2 infection in free-ranging wild animals in
999   Belgium. *Transbound. Emerg. Dis.* (doi:10.1111/tbed.14131)

1000 130.Weber A, Kalema-Zikusoka G, Stevens NJ. 2020 Lack of Rule-Adherence During Mountain
1001   Gorilla Tourism Encounters in Bwindi Impenetrable National Park, Uganda, Places Gorillas
1002   at Risk From Human Disease. *Front Public Health* **8**, 1. (doi:10.3389/fpubh.2020.00001)

1003 131.Barbosa A *et al.* 2021 Risk assessment of SARS-CoV-2 in Antarctic wildlife. *Sci. Total*
1004   *Environ.* **755**, 143352. (doi:10.1016/j.scitotenv.2020.143352)

1005 132.Gibbons A. 2021 Captive gorillas test positive for coronavirus. *Science*
1006   (doi:10.1126/science.abg5458)

1007 133.Daly N. 2021 First great apes at U.S. zoo receive COVID-19 vaccine made for animals.
1008   *National Geographic* See https://www.nationalgeographic.com/animals/article/first-great-
1009   apes-at-us-zoo-receive-coronavirus-vaccine-made-for-animals.

1010 134.Aleccia J. 2020 'The Biggest Nemesis': Black-Footed Ferrets Get Experimental
1011   Coronavirus Vaccine. *Kaiser Health News*, 27 December. See
1012   https://www.cpr.org/2020/12/27/the-biggest-nemesis-black-footed-ferrets-get-experimental-
1013   coronavirus-vaccine/.

1014 135.Xu L *et al.* 2020 COVID-19-like symptoms observed in Chinese tree shrews infected with
1015   SARS-CoV-2. *Zool Res* **41**, 517–526. (doi:10.24272/j.issn.2095-8137.2020.053)

1016 136.Becker DJ, Washburne AD, Faust CL, Pulliam JRC, Mordecai EA, Lloyd-Smith JO,
1017   Plowright RK. 2019 Dynamic and integrative approaches to understanding pathogen
1018   spillover. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **374**, 20190014.
1019   (doi:10.1098/rstb.2019.0014)

1020 137.Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO.
1021   2017 Pathways to zoonotic spillover. *Nat. Rev. Microbiol.* (doi:10.1038/nrmicro.2017.45)

1022 138.Morris DH *et al.* 2020 The effect of temperature and humidity on the stability of SARS-CoV-
1023   2 and other enveloped viruses. *bioRxiv* (doi:10.1101/2020.10.16.341883)

1024 139.Bean AGD, Baker ML, Stewart CR, Cowled C, Deffrasnes C, Wang L-F, Lowenthal JW.
1025   2013 Studying immunity to zoonotic diseases in the natural host - keeping it real. *Nat. Rev.*
1026   *Immunol.* **13**, 851–861. (doi:10.1038/nri3551)

1027 140.Letko M, Seifert SN, Olival KJ, Plowright RK, Munster VJ. 2020 Bat-borne virus diversity,
1028   spillover and emergence. *Nat. Rev. Microbiol.* **18**, 461–471. (doi:10.1038/s41579-020-
1029   0394-z)

1030 141.Sawatzki K, Hill NJ, Puryear WB, Foss AD, Stone JJ, Runstadler JA. 2021 Host barriers to
1031   SARS-CoV-2 demonstrated by ferrets in a high-exposure domestic setting. *Proc. Natl.*
1032   *Acad. Sci. U. S. A.* **118**. (doi:10.1073/pnas.2025601118)

1033 142.Ruiz-Arrondo I, Portillo A, Palomar AM, Santibanez S, Santibanez P, Cervera C, Oteo JA.

2020 Detection of SARS-CoV-2 in pets living with COVID-19 owners diagnosed during the COVID-19 lockdown in Spain: A case of an asymptomatic cat with SARS-CoV-2 in Europe. *bioRxiv*. (doi:10.1101/2020.05.14.20101444)

143. Peel AJ *et al.* 2019 Synchronous shedding of multiple bat paramyxoviruses coincides with peak periods of Hendra virus spillover. *Emerg. Microbes Infect.* **8**, 1314–1323. (doi:10.1080/22221751.2019.1661217)

144. Restif O *et al.* 2012 Model-guided fieldwork: practical guidelines for multidisciplinary research on wildlife ecological and epidemiological dynamics. *Ecol. Lett.* (doi:10.1111/j.1461-0248.2012.01836.x)

145. Cook JA *et al.* 2020 Integrating Biodiversity Infrastructure into Pathogen Discovery and Mitigation of Emerging Infectious Diseases. *Bioscience* **70**, 531–534. (doi:10.1093/biosci/biaa064)

146. Thompson CW *et al.* 2021 Preserve a Voucher Specimen! The Critical Need for Integrating Natural History Collections in Infectious Disease Studies. *MBio* **12**. (doi:10.1128/mBio.02698-20)