Stress alters the activity of Leydig cells dependently on the diurnal time

Marija Lj Medar[1], Silvana A Andric[1], Tatjana S Kostic[1]

[1]*The University of Novi Sad, Faculty of Sciences Novi Sad, Department of Biology and Ecology,*

Laboratory for Chronobiology and Aging, Laboratory for Reproductive Endocrinology and

Signaling, Novi Sad, Serbia

* Correspondence:

Tatjana S Kostic

Dositeja Obradovica Sq 2,

Novi Sad, 21000

Serbia

tatjana.kostic@dbe.uns.ac.rs

Keywords: Stress, Leydig cell, Clock genes, Testosterone, Glucocorticoid receptor, Circadian rhythm

 Short title: Leydig cell stress response depends on daytime

Abstract

Background: The increasing amount of data points to the circadian timing system as an essential part of processes regulating androgen homeostasis. However, the relationship between stress response, timekeeping-, and steroidogenesis-related systems is unexplored.

Objective: The purpose of the study was to analyze the stress-response of the testosterone-producing Leydig cells depending on the stressful event's time.

Materials and methods: The study was designed to follow the effects of 3-hour immobilization (IMO) applied at different periods during the day. The IMO performed once (1xIMO) or repeated in 10 consecutive days (10xIMO). Principal-component-analysis (PCA) followed the expression study of the clock and steroidogenic-related genes in Leydig cells.

Results: Both types of IMO in all investigated periods increased corticosterone and decreased testosterone blood level. Transcriptional analysis revealed different sensitivity to IMO events depending on the circadian time. The majority of steroidogenesis-related genes (*Lhcgr*, *Cyp11a1*, *Cyp17a1*, *Hsd3b1/2*) were down-regulated in the inactive but unchanged or even up-regulated in the active phase of the day. Both types of IMO potentiated the expression of clock elements *Bmal1*/BMAL1, *Per1/*PER1 regardless of the day's stage and reduced *Rev-erba* in the inactive phase. The PCA confirmed a major shift, for both IMO-types, in the transcription of steroidogenesis and clock genes across the inactive/active phase. Further, the diurnal pattern of the glucocorticoid receptor (*Nr3c1*/GR) expression in Leydig cells was increased in the inactive phase due to 10xIMO. The observed time-dependent IMO-response of the Leydig cells correlated with different GR engagements.

Discussion: Stress- and the circadian-system coordinatively shape Leydig cell's physiology, assuming diverse GR engagement as a possible factor in mediating the diurnal effect of stress.

Conclusion: The Leydig cell's stress-response depends on the time of the stressful situation, emphasizing the importance of circadian activity in supporting androgen homeostasis and male fertility.

Introduction

The increase in evidence points to the circadian timing system as an essential part of processes regulating reproduction. The circadian timing system provides an organism with the sense of time of day to ensure that physiological and behavioral events coincide to give offspring [1]. Temporal coordination of different organs is mediated by the actions of the master clock in hypothalamic suprachiasmatic nuclei, together with hormonal and nervous systems employing multi-level signaling through negative and positive feedback loops. The content of hormonal signals is overwhelmingly conveyed in a rhythmic secretory pattern. In cells, including cells of the reproductive tract, the timing system consists of clock genes coding transcription factors that operate through positive (BMAL1, CLOCK, NPAS2, RORA,B) and negative (PER1-3, CRY1,2, REV-ERA,B, DEC) elements organized in transcription-translation regulatory feedback loops [2,3]. The rhythm in clock gene transcription is decoded into physiological signals through cell-specific transcriptional activity via circadian promoter elements such as E-boxes, D-boxes, and ROREs [4].

In males, testosterone production is a rhythmic process predominantly regulated by the rhythm of pulsatile LH secretion and clock elements [5,6]. Those cells rhythmically express clock genes (*Bmal1*, *Per1-3*, *Cry1,2*, *Reverba,b*, *Dec*, *Rora,b*) as well as major steroidogenic genes (*Star, Cyp11a1*, and *Cyp17a1*) followed by an oscillatory pattern of blood testosterone [7,8]. In humans and rats, testosterone reaches a peak at the beginning of the active phase [7–9]. Testosterone is a crucial hormone regulating male reproduction, including the development and maintenance of the male phenotype; consequently, potential changes in circadian rhythmicity of testosterone production/secretion could impact reproductive function.

It is known that stress disturbs male reproduction by decreasing blood testosterone [10–12] acting

on all three levels of the hypothalamic-pituitary-testicular axis [13]. The effect is achieved through

the neural and hormonal system, possibly associated with central and peripheral clocks [14].

In addition, some earlier studies suggest stress-response dependence on time of stress event [15,16],

pointing to the relationship between the circadian rhythm of life functions and the stress-

response. However, molecular events and signaling pathways responsible for such differences

are not described.

Keeping in mind the importance of sustainable circadian rhythm on homeostasis, including

maintaining reproductive ability, this study aims to explore the relationship between stress

response and circadian timekeeping in testosterone-producing Leydig cells. Precisely, this study

estimated the stress-response of the Leydig cells dependently on the time of the stressful event.

Materials and methods

Ethical Approval

All the experiments were carried out in the Laboratory for Chronobiology and Aging, the

Laboratory for Reproductive Endocrinology and Signaling, DBE, Faculty of Sciences at the

University of Novi Sad. All experiments were approved by the Ethical Committee on Animal

Care and Use of the University of Novi Sad (statement no. 01-201/3), operating under the rules

of the National Council for Animal Welfare and following statements of National Law for

Animal Welfare (copyright March 2009). All experiments were performed and conducted by

following the European Convention for the Protection of Vertebrate Animals used for

Experimental and other Scientific Purposes' (Council of Europe No 123, Strasbourg 1985) and

NIH Guide for the Care and Use of Laboratory Animals (NIH Publications, No. 80 23, revised,

7th ed., 1996). All experiments were in adherence to the ARRIVE guidelines.

Animals & stress model

Adult (age, 3 mo; body wt, 250–270 g) male *Wistar* rats, bred and raised in the Animal Facility

of the Department of Biology and Ecology (Faculty of Sciences, University of Novi Sad), were

used for the experiments. The rats were raised in controlled environmental conditions (22 ± 2°C;

14h light:10h dark; lights on at 6:00 AM) with food and water ad libitum. The chosen

experimental model of stress was psychophysical stress induced by 3h immobilization (IMO).

IMO was performed as described previously [10,17]. Briefly, rats were bound in the supine position

for 3h by fixing their limbs to the wooden board with thread. The head motion was not limited.

At the end of experiments, rats were quickly decapitated without anesthesia. All activities during

the dark phase were performed under the red light.

The two types of experiments were performed depending on number od IMO session: (1) to

analyze circadian sensitivity on acute IMO (1xIMO), rats were subject to IMO in different

periods during 24h (ZT0-3, ZT8-11, ZT14-17, and ZT20-23; ZT0 is a time when the light turned

on); (2) to examine the effect of repeated IMO, rats were immobilized each day at the same time

(ZT0-3, ZT8-11, and ZT20-23) for 10 consecutive days (10xIMO). Corresponding groups of

freely moving rats served as controls in both IMO experiments.

For both tipes of IMO experiments, rats for control and experimental groups were randomly

divided into 4 or 3 time points consisted each from 5-6 rats per time points. No masking was

used during group allocation, data collection or data analysis. Previous results from our group

obtained on the rat model showed that the size of the group of 5-6 rats was sufficient to calculate

hormone parameters of the circadian rhythm [7,8]. The sample size was checked also by Power

121   Analysis using G Power software (http://core.ecu.edu/psyc/wuenschk/Power.htm) according to

122   previous results. Experiments were repeated two times.

Preparation of purified Leydig cells

The testes were isolated, decapsulated, and the main blood vessel was removed as it was

previously described (Kostic et al., 2010; Baburski et al., 2016, 2015). Decapsulated testes were

placed in plastic tubes (2 testes per tube) containing collagenase solution (0.25 mg

collagenase/mL-1.5 %-BSA-20 mM HEPES-M199, Sigma, St Louise, Missouri) and incubated

in a shacking-water bath (15 min/34 °C/120 cycles/min). The reaction was stopped by adding a

sufficient amount of cold M199 medium, and seminiferous tubules were removed by filtration.

The interstitial cells' suspension was centrifuged (160xg/5 min), and the cell pellet was

resuspended in DMEM-F12 medium (Sigma, St Louise, Missouri) 8 mL/rat. The resuspended

cells ($35-40x10^6$) were moved on the Percoll gradients (Sigma, St Louise, Missouri) and

centrifuged 1100xg/28 min. After centrifugation, Leydig cells were harvested, washed with a

sufficient amount of the M199-0.1% BSA, centrifuged (200xg/5 min), and resuspended in

DMEM-F12 medium (5 mL/animal). In line with the procedure and according to the HSD3B

staining [20] presences of Leydig cells in the culture was greater than 90%; according to the Trypan

blue exclusion test [7], viability of the cells was greater than 90%. The Leydig cell suspension was

centrifuged, and the pellet was stored at -80 °C for RNA and protein analysis.

RNA isolation and cDNA synthesis

Total RNA from purified Leydig cells was isolated using the using Trizol (TermoFisher

Scientific, Waltham, USA) or Rneasy kit reagent following the protocol recommended by the

manufacturer (QIAGEN, Valencia, California). The amount of total RNA's and purity were

assesed by spectrophotometric measurement (NanoDrop, Shimadzu). After DNase I treatment,

cDNA was synthesized using the Applied Biosystems kit (Termo Fisher Scientific,

Massachusetts, USA) according to the manufacturer's instructions. Analyses of appropriate and

relevant positive and negative controls were performed for both DNase treatment and cDNA

synthesis.

Real-time PCR and relative quantification

Relative gene expression was done by RQ-PCR using SYBR Green-based chemistry (Applied

Biosystems, Foster City, California) in the presence of a 5 μl aliquot of the reverse transcription

reaction product (25 ng RNA calculated on the starting RNA) and specific primers. The primer

sequences used for RQ-PCR analysis and GenBank accession codes for full gene sequences

(www.ncbi.nlm.nih.gov/sites/entrez) are provided in Suppl. Table 1. *Gapdh* was also measured

in the same samples and used to correct variations in RNA content. The relative quantification of

each gene was performed in duplicate.

Protein extraction and western blot analysis

Platted cells were lysed by buffer (20 mM HEPES, 10 mM EDTA, 2.5 mM $MgCl_2$, 1 mM DTT,

40 mM β-glicerophosphate, 1% NP-40, 2 μM leupeptin, 1 μM aprotinin, 0.5 mM AEBSF,

phosphatase inhibitor cocktail (PhosSTOP, Roche, Basel, Switzerland) and the lysates were

uniformed by Bradford method. Samples were mixed with SDS protein gel loading solution

(Quality Biological, Gaithersburg, USA), boiled for 5 min, and separated by one-dimensional

SDS-PAGE electrophoresis (Bio-rad, Berkeley, USA).

Transfer of the proteins to the PVDF membrane was performed by electroblotting overnight at

4°C/40A, and the efficacy of transfer was checked by staining and distaining the gels. The

membranes were blocked by 3% BSA-1xTBS for 2h/room temperature, incubation with primary

antibody was done overnight at 4 °C, and 0.1% Tween-1xTBS was used for washing membranes

incubation with secondary antibody was done for 1h/room temperature [6]. Immunoreactive bands

were detected using luminol reaction and MyECL imager (www.Fischer.sci)/films and

densitometric measurements were performed by Image J program version 1.32

(https://rsbweb.nich.gov//ij/download.html). Anti-BMAL1 purchased from Abcam (Cambridge,

UK, #ab49421, dilution 1:800); anti-PER1 purchased from Abcam (Cambridge, UK, #ab3443,

dilution 1:500); anti-GR purchased from Thermo Fisher Scientific (Waltham, Massachusetts,

USA, #MA1-510, dilution 1:1000); anti-ACT purchased from Santa Cruz Biotechnology

(Heidelberg, Germany, #sc-8432, dilution 1:800). All antibodies used in this study are validated

and previously published by our group [6,7,21].

Hormone and glucose measurements

Androgen level was measured by RIA and it was referred to testosterone + dihydro-testosterone

(T + DHT) because the anti-testosterone serum No. 250 showed 100% cross-reactivity with DHT

[18,19]. All samples were measured in duplicate in one assay (sensitivity: 6 pg per tube; intraassay

coefficient of variation 5–8%). For serum corticosterone levels [22], all samples were measured in

duplicate, in one assay by the corticosterone EIA Kit (Caymanchem, Michigan, USA) with 30

pg/ml as the lowest standard significantly different from blank. The level of glucose in serum

was estimated by using BAYER Contour plus Glucose Monitoring System (Bayer, Basel,

Switzerland; measuring range 0.6-33.3 mmol/L). Technically, all serum samples obtained

individually, were measured in duplicate.

Statistical and rhythm analysis

The results represent group means ± SEM values of two in vivo experiments. Rhythm

parameters (p, MESOR, amplitude, and acrophase) were obtained by cosinor.lm() function from

the "cosinor" package [23] in the R environment and by using online Cosinor.Online software fitted

to 24 h period (http://www.circadian.org/softwar.html) [24]. Statistical significance (p < 0.05)

between the treated group and control group within the same time point and between the groups

including all time points (p < 0.05) were analyzed using Mann–Whitney test and two-way

ANOVA, respectively.

Principal component analysis (PCA) was done with dudi.PCA() function implemented in "ade4"

package [25], on scaled and centered data matrix, within the R environment. We decided to retain

the first two PC based on eigen values and cumulative variation. In support of such a decision,

we performed Horn's parallel analysis for a PCA with the "paran" package, to adjust for finite

sample bias in retaining components [26]. Biplots visualization were done with "factoextra"

package [27]. The effects of 1xIMO and 10xIMO on gene expression were presented by using The

Clustered Image Map (CIM) analysis (https://discover.nci.nih.gov/ cimminer / oneMatrix.do).

Results

Time-dependent effect of IMO on hormone levels.

To examine whether the circadian time of stress event can provoke different responses, rats were

immobilized for 3h, in different periods during 24h, once (1xIMO) or repeatedly for 10 days

(10xIMO), and responses were analyzed immediately after the IMO session.

As expected [21], corticosterone, glucose, and testosterone show diurnal pattern in the serum of

control rats (Fig. 1A&B&C; the rhythm parameters in Suppl. Table 2). The peak of these

oscillations occurs around ZT11 for all three. 1xIMO effectively increased corticosterone (Fig.

1A) and glucose (Fig. 1B) levels in all investigated ZT, although glucose levels were increased

in the beginning of inactive and at the end of active phase (ZT3 and ZT23). On the contrary,

1xIMO reduced testosterone levels in serum in all investigated ZTs (Fig. 1C).

Analysis of the relationship between blood testosterone and corticosterone (T/C) could point to

the ability of the Leydig cells to produce testosterone in stress conditions. In that respect, the

relative T/C was calculated on ZT3 in control group (ZT3=1). Obtained results depicts circadian

fluctuations in blood testosterone and corticosterone; the peak of fluctuations observed in the

control group was at ZT11 (T/C ZT11=1.6). Nevertheless, 1xIMO reduced T/C approximately

38-fold, depending on time of IMO event. The lowest T/C value was observed in ZT3 (T/CZT3

= 0.02), while the highest was observed in ZT11 (T/CZT11 = 0.03) (Fig. 1D).

The repetition of stress events over ten days always at the same time, 10xIMO, increased blood

corticosterone (Fig. 1G), and glucose (Fig. 1H). However, increments of corticosterone were

one-third lower than after 1xIMO at all tested times (Fig. 1G). In addition, the repeated IMO

profoundly reduced testosterone (Fig.1I), which resulted in a substantially decreased T/C ratio

(Fig. 1J). Described patterns were observed in all examined ZTs.

Since LH is the principal regulator of Leydig cell endocrine function, the transcriptional pattern

of the genes encoding pituitary LH subunits (*Cga*, *Lhb*) were studied. Both genes showed a

diurnal transcriptional model in control conditions (Fig.1 E, F; please see rhythm parameters in

Suppl.Table2). 1xIMO significantly decreased *Cga* at ZT11 and ZT17 (Fig. 1E) and *Lhb* at

ZT11, ZT17, and ZT23 (Fig. 1F). The 10xIMO turn-down transcription of *Cga* (Fig. 1K)

and *Lhb* (Fig. 1L) at ZT11.

Figure 1. Time-dependent effect of IMO on hormone levels and pituitaries gene levels.
Wistar rats were subject to 1xIMO for three hours in four-time points (ZT0-3, ZT8-11, ZT14-17, ZT20-24) or to 10xIMO in three-time points (ZT0-3, ZT8-11, ZT20-24) during the 24h. Serum was collected for determination of corticosterone, glucose, and testosterone levels as well as testosterone/corticosterone (T/C) ratio (1xIMO: A-D; 10xIMO: G-J). Pituitaries were collected, RNA isolated for RT followed by RQ-PCR analysis of genes encoding the LH subunits (1xIMO: E, F; 10xIMO: K, L). Data points are group mean ± SEM values, (n = 10-12 for A, B, C, G, H, I, J, and n=5-6 for E, F, K, L). For possible rhythm prediction please see Suppl. Table 2. *Statistical significance between control and IMO groups for the same time point (p < 0.05).

The time-dependent effects of IMO on the steroidogenesis-related genes expression in Leydig cells

The endocrine function of the Leydig cell entirely depends on the expression pattern of the steroidogenic elements. Therefore, to detect a possible different time-dependent stress effect on the gene expression essential for the Leydig cells' endocrine function, we perform transcriptional analysis in Leydig cells originating from control and IMO rats in various periods during 24h.

The *Lhcgr*, encoding LH receptor, did not show a diurnal pattern of transcription, (Fig. 2A).

1xIMO significantly decreased the *Lhcgr* transcription in the light phase, while the dark phase

was without effect (Fig. 2 A).

Key steroidogenic genes, *Star*, *Cyp11a1*, *Cyp17a4*, exhibit a diurnal transcriptional pattern

which reach a maximum approximately in the middle of the light phase (Fig. 2B, C, E,

Suppl.Table2), a few hours earlier than the peak of blood testosterone. 1xIMO changed the

expression pattern of the *Star, Cyp11a1*, and *Cyp17a4,* underlying a pronounced reduction in the

inactive (light) phase (Fig. 2B, C, E). Although, results did not support diurnal pattern in the

transcription of *Hsd3b1/2* and *Hsd17b4,* 1xIMO significantly reduced transcription of both genes

during the inactive (light) phase (Fig. 2D, F).

The time-dependent effects of stress response were also seen in 10xIMO conditions (Fig. 2 G-L).

The *Lhcgr* (Fig. 2G), *Star* (Fig. 2H), *Cyp11a1* (Fig. 2I), *Hsd3b1/2* (Fig. 2J), and *Cyp17a4* (Fig.

2K) were down-regulated in inactive without significant effect in active period. In addition,

the *Hsd17b4* (Fig. 2L) was down-regulated by 10xIMO in all three examined ZTs.

Thus, the results showed different expression of steroidogenic genes over 24 h, including

different IMO effects on their expression: in control samples, inactive and active phases tend to

separate across the first two PC for 67.7% of the total dataset of steroidogenesis-related genes

variability (Fig. 2M; variable loadings are shown in Suppl. Table 3); in 1xIMO and 10xIMO PC

accounts for 74.5% (Fig. 2N, variable loadings are shown in Suppl. Table 4) and 94.4% (Fig. 2O,

variable loadings are shown in Suppl. Table 5) data variability, respectively. In 1xIMO

conditions, inactive and active phases are strongly driven by distinct expression profiles

of *Lhgcr, Hsd3b1/2, Cyp11a1*, and *Hsd17b4* (Fig. 2N). Similarly, in 10xIMO conditions the

transcription of *Lhgcr, Hsd3b1/2, Cyp11a1* and *Star* were less changed in the active than in the

inactive phase (Fig. 2O). Such gene expression profiles point to the robustness of steroidogenesis-related genes (light and dark phase, respectively) under different IMO treatments, which may pinpoint them as a landmark for stress conditions.

Figure 2. The time-dependent effects of IMO on the expression of the steroidogenesis-related genes in Leydig cells. *Wistar* rats were subject to 1xIMO for three hours in four time points (ZT0-3, ZT8-11, ZT14-17, ZT20-24) or to 10xIMO in three time points (ZT0-3, ZT8-11, ZT20-24) during the 24h. Leydig cells were purified, RNA extracted and used for RT followed by RQ-PCR analysis of steroidogenesis-related genes expression (1xIMO: A-F; 10xIMO: G-L). Data points are group mean ± SEM values, (n = 5-6). For possible rhythm prediction please see Suppl. Table 2. *Statistical significance between control and IMO groups for the same time point ($p < 0.05$). PCA of steroidogenic-related genes depending on active/inactive phase in control (M), 1xIMO (N) and 10xIMO (O) conditions; Dim1 and Dim2 represents the first two

286   PC and % of retained variation. cos2 estimates the qualitative representation of variables. In M,
287   N and O datasets were prepared as a value that deviates from control at a given time
288   (Supplemental table 3, 4 and 5).

290   The time-dependent effects of IMO on the expression of the clock genes in Leydig cells

The effect of IMO on the peripheral clock in Leydig cells was estimated by transcription analysis

of canonical clock genes (Fig. 3; Suppl.Table2).

The majority of examined genes (*Bmal1*, *Per1/2*, *Cry1/2*, *Rev-erba/b*, *Rob*) had a diurnal

expression in rat Leydig cells (Fig. 3; Suppl.Table2). On the contrary, *Clock*, *Npas*,

and *Rora* transcripts did not follow any significant pattern (Fig. 3 B, C, J; Supplement Table

2). *Bmal1* transcript was the biggest at the beginning (Fig. 3A) while *Per1/2* (Fig. 3D, E)*,*

Cry2 (Fig. 3G)*, Rev-erba/b* (Fig. 3H, I)*, and Rorb* (Fig. 3K) at the end of the light phase,

followed by *Cry1* (Fig. 3F) rise in the dark phase. Generally, 1xIMO changed transcription of

clock genes in Leydig cells: *Bmal1* increased in all IMO-times (Fig. 3A), and *Clock* (Fig. 3B)

decreased immediately after morning IMO (ZT3) while its counterpart *Npas2* was unaffected by

IMO situations (Fig. 3C). However, clock negative regulators in the circadian clock's primary

loop were enhanced: *Per1* after all IMO sessions regardless of the time of occurrence (Fig. 3D),

and *Per2* at ZT3 and ZT23*, Cry1* at ZT3, ZT11 and ZT23*, and Cry2* at ZT23 (Fig. 3E, F, G).

Additionally, IMO affected the transcription of genes in the secondary loop by decreased

expression of *Rev-erba/b* during the inactive phase (Fig. 3H, I) and *Rorb* at ZT17 and ZT23 (Fig.

3K). Changed protein levels of BMAL1 and PER1 due to IMO were confirmed by western blot

analysis. As expected, BMAL1 and PER1 showed circadian profiles in the control samples with

an increased level of BMAL1 in the light in relation to the dark phase and vice versa situation for

its negative regulator PER1 (Fig. 3L, M). IMO increased BMAL1 (Fig. 3L) and PER1 in all

examined time points (Fig. 3M).

10xIMO altered clock genes expression in a similar way as 1xIMO (Fig. 3N-W):

the *Bmal1* increased and (Fig. 3N) *Per1* increased in all tested IMO-periods (Fig. 3Q) while *Rev-*

erba decreased by IMO-sessions in light phase (Fig. 3U). 10xIMO efficiently increased BMAL1

(Fig. 3Z) and PER1 (Fig. 3Y) after all analyzed time-points.

Likewise previous, a PCA was run to convince into IMO effects on diurnal fluctuations of clock

genes expression in Leydig cells. As expected, the diurnal phase transition is characterized by a

distinct expression pattern of clock genes, *Rev-erbb/a, Rora/b, Cry2* in the inactive (light) phase,

and *Cry1* during the active (dark) phase (Fig. 3AA, variable loadings are shown in Suppl. Table

6). The *Bmal1* and *Per1/2* had opposite directions (Fig. 3A, variable loadings are shown in

Suppl. Table 6). Additionally, PCA confirmed that light and dark phases have different

expression patterns under 1xIMO or 10xIMO conditions. In 1xIMO conditions, a pronounced

cluster of genes in active (*Rorb, Cry2, Rev-erbb, Per1)*, opposes *Bmal1* in inactive phase and

contributes to the separation between phases (Fig. 3AB, variable loadings are shown in Suppl.

Table 7). Interestingly, *Clock*, *Npas2*, *Per2* but also *Rev-erba* are minorly contributing to the

separation between phases (Fig. 3AB, variable loadings are shown in Suppl. Table 7).

Additionally, 10xIMO treatment drives a robust difference in expression of *Rev-erba* and *Rora*

during the active and *Rev-erbb* and *Cry2*, during inactive phase (Fig. 3AC, variable loadings are

shown in Suppl. Table 8). Also, the light and dark phase shows the most distinctive change/shift

across all analyzed data (94.0% variability), which implies that 10xIMO is strongly affecting the

clock genes expression profiles (Fig. 3AC, variable loadings are shown in Suppl. Table 8).

Figure 3. The time-dependent effects of IMO on the expression of the clock-related genes in Leydig cells. Wistar rats were subject to 1xIMO for three hours in four-time points (ZT0-3, ZT8-11, ZT14-17, ZT20-24) or to 10xIMO in three-time points (ZT0-3, ZT8-11, ZT20-24) during the

24h. Leydig cells were purified, RNA extracted, and used for RT followed by RQ-PCR analysis
of steroidogenesis-related genes expression (1xIMO: A-K; 10xIMO: N-W). Data points are
group mean ± SEM values, (n = 5-6). For possible rhythm prediction please see Suppl. Table 2.
The protein level of BMAL1 (1xIMO: L; 10xIMO: Z) and PER1 (1xIMO: M; 10xIMO: Y)
were determined by western blot. Representative blots are shown. Bars above blots represent
mean ± SEM values from pooled data using scanning densitometry values normalized on ACT.
*Statistical significance between Control and IMO group at the same time point (p < 0.05).
Expression pattern of clock genes depending on active/inactive phase in Leydig cells from
control (AA), 1xIMO (AB) and 10xIMO (AC) rats analized by PCA; Dim1 and Dim2 represents
the first two PC and % of retained variation. cos2 estimates the qualitative representation of
variables. In AA, AB and AC datasets were prepared as a value that deviates from control at a
given time (Supplemental table 6,7 and 8).
347

The time-dependent effects of IMO on the *Nr3c1*/GR expression in Leydig cells

Glucocorticoids are the primary hormonal mediators of stress with a large regulatory impact on

Leydig cell`s steroidogenesis as well as survival [10]. In that respect, the time-dependent *Nr3c1*/GR

expression was monitored in Leydig cells from control and 1xIMO as well as 10xIMO rats. In

Leydig cells from control rats, the diurnal expression pattern of *Nr3c1* and GR was observed

(Fig. 4) with increased expression at the end of the light and the beginning of the dark phase.

That 24h expression pattern of GR correlates with blood corticosterone fluctuation (R=0.885),

allowing efficient regulation of target genes in Leydig cells. However, 1xIMO in the rest period

reduced *Nr3c1* transcript (Fig. 4A) without significant effect on GR (Fig. 4B). Still, a decrease in

Nr3c1 was not detected in the case of 1xIMO in the active period. On the contrary, IMO

repetition (10xIMO), in rest period, elevated *Nr3c1* (Fig. 4C) and GR (Fig. 4D) in Leydig cells.

Again, the effect of stress was less pronounced in the active period.

Since several steroidogenesis-related and clock genes showed different expression patterns

regarding the time and type of IMO session, we perform a correlation study to assess the possible

association with GR-signaling. The Cluster Image Map (Fig. 4G, H) represents the correlation

between the GR-signaling (presented as a ratio of the blood corticosterone/GR in Leydig cells)

and the transcriptional level of several steroidogenesis-related (Fig. 4G) and clock genes (Fig.

4H) in Leydig cells. The map presents a relationship of GR-signaling and these two groups of

genes in inactive (-) and active (+) phase of the day in the control, 1xIMO or 10xIMO

conditions.

In the inactive phase of the day in the control and 1xIMO groups, the analysis showed a negative

correlation between C/GR and *Lhcgr*, *Star*, *Cyp11a1*, and *Hsd3b* (Fig. 4G). A positive

correlation in the control and 1xIMO group's inactive phase was observed between C/GR

and *Hsd17b* (Fig. 4G). In the 10xIMO condition, R values for all investigated genes vary

between -0.4 and 0.4 (Fig. 4G). Clock genes, such are *Bmal1*, *Npas2*, *Per1,2,* and

Cry1,2, positively correlate with C/GR in conditions when 1xIMO occurred in both inactive and

active phase. Furthermore, a positive correlation was observed for *Rora* and *Rorb* in the inactive

phase, and negative in the active phase of the day (Fig. 4H). However, the opposite was

registered for *Rev-ereba*, i.e.*,* negative in the inactive and positive in the active phase (Fig. 4H).

In the 10xIMO conditions, regardless of the phase of the day, a negative correlation was

observed with *Clock, Npas2, Per1,2*. The *Cry1,2*, and *Rev-erbb,* negatively correlate with C/GR

in the inactive period, while *Rev-erba, Rora*, and *Rorb* positively (Fig. 4H).

Figure 4. The time-dependent effects of IMO on the expression of the glucocorticoid receptor gene and protein level in Leydig cells. Wistar rats were subject to 1xIMO for three hours in four-time points (ZT0-3, ZT8-11, ZT14-17, ZT20-24) or to 10xIMO in three-time points (ZT0-3, ZT8-11, ZT20-24) during the 24h. Leydig cells were purified, RNA extracted, and used for RT followed by RQ-PCR analysis of *Nr3c1* gene expression (1xIMO: A; 10xIMO: C;). Data points are group mean ± SEM values, (n = 5-6). For possible rhythm prediction please see Suppl. Table 2. The protein level of GR (1xIMO: B; 10xIMO: D;) was determined by western blot. Representative blots are shown. Bars above blots represent mean ± SEM values from pooled data using scanning densitometry values normalized on ACTB. *Statistical significance between Control and IMO group at the same time point (p < 0.05). For The Clustered Image Map datasets show R values from correlation analysis between GR-signaling (presented as a blood corticosterone/GR ratio) and transcription of steroidogenesis-related genes (G) or clock genes (H) during inactive (-) and active (+) phase of the day at different experimental conditions. The red line represents the genes positively correlated with GR-signaling. The blue line corresponds to the genes that were negatively correlated with the GR-signaling.

Discussion

Stress is challenged by any stimulus that disrupts or threatens homeostatic equilibrium. The stress response engages multiple coordinated processes to restore homeostasis and preserve life

[14]. However, reproductive functions are suppressed under various stress conditions depending on

genetic predisposition to stress, type, and duration of stress.

Besides that, results from the present study strongly suggest diurnal time as the critical element

in determining the stress response. Namely, our previous work pointed to diurnal expression of

Leydig cell`s clock genes (*Bmal1*, *Per1/2/3*, *Cry1/2*, *Rev-erba/b*, *Rorb*, *Dec1/2*, *Dbp,* and *E4bp4*)

but also genes responsible for testosterone production (*Star*, *Cyp11a1,* and *Cyp17a1*) [7,8].

However, the expression pattern of these genes changed due to stress and varied depending on

the time of the stressful event. In the inactive (light) phase, the critical steroidogenesis-related

genes (*Lhcgr, Cyp11a1*, *Cyp17a1, Hsd3b1/2)* were down-regulated, but in the active (dark)

phase of the day, they were unchanged or even up-regulated. The PCA confirmed significant

separation of effects on steroidogenesis-related genes depending on the time of the stress in both

types of stress used.

It is well known that all steroidogenic events in Leydig cells are initiated and depend on LH-

receptor activation, including expression of proteins involved in steroidogenesis such

are *Star*/StAR, *Cyp11a1*/CYP11A, *Hsd3b*/HSD3B, *Cyp17a*/CYP17A,   and *Hsd17b*/HSD17B

(Payne and Hales, 2004). However, results indicate severe inhibitory IMO effects on *Lhcgr*

transcription in inactive but not in the active period. This effect did not depend on the type of

stress applied. Since LH-receptor signaling is crucial for steroidogenic genes transcription and

steroidogenesis activation [28,29] decreased expression of *Lhcgr* due to IMO in the rest period

could make a difference in respect to IMO in the active period. Also, IMO decreases

transcription of pituitary genes encoding LH subunits, *Cga* during the rest and at the beginning

of the active period together with *Lhb* in both types of stress. Changes of both genes likely have

consequences for the quantity of pulsatile LH secretion and impaired Leydig cells stimulation.

Nevertheless, the IMO is changing the diurnal organization of clocks in the Leydig cells. IMO in

both conditions applied potentiates the expression of *Bmal1*/BMAL1 and *Per1/2*/PER1 and in

the inactive period decreased *Rev-erba*, suggesting that stressful stimuli can entrain the clock in

the Leydig cells. Further, BMAL1 is cyclically expressed in the rat [6,8] and mouse [30] Leydig cells,

stimulating *StAR* expression [31] therefore contributing to the cyclic expression of StAR protein.

Since IMO in the dark phase promotes BMAL1, this may mitigate the effects of stress on

steroidogenesis during the night. Also, the promoters of steroidogenic genes such are

Star, *Cyp11a1*, and *Cyp17a1* [30] as well as promoter of steroidogenic regulators such as *Nur77*

contain clock-controlled RORE and E-box, suggesting BMAL1, PERs, and REV-ERBs as

potential regulators of rhythmic expression of steroidogenic genes. Further studies are necessary

to confirm this hypothesis.

The clock system interacts with many different signals to produce an integrated output over the

24h cycle, entraining other cyclic activities in the cell [32]. It is well known that rhythmic

corticosterone secretion is an important timing signal for the coordination of the peripheral clock

[33]. Both circadian and stress-mediated aspects play a crucial role in determining the level of

corticosterone secretion and mediating the effects on target cells. Corticosterone acts as a

peripheral clock synchronizer [34,35] by regulation of clock genes expression through genomic

actions mediated by activation of GR [36]. Actually, it was found that *Per1* and *Per2* contain

glucocorticoid-responsive elements. Further, *Rev-erba* and *Rora* are negatively regulated by

glucocorticoids [37,38]. Results from this study showed altered *Nr3c1*/GR expression profile in

Leydig cells due to IMO. Thus, results suggest that the diurnal effect of stress is mediated,

among others, by the circadian dynamics of corticosterone signaling.

Indeed, correlation study suggested a positive relationship between GR signaling (C/GR) and

transcriptional activity of the clock *Bmal1*, *Per1,2*, *Cry1,2*, and *Rora,b* and negative with

steroidogenesis-related *Lhcgr*, *Star*, *Cyp11a1*, and *Hsd3b* when 1xIMO occurred in the inactive

phase of the day. However, when IMO repeatedly occurred in the inactive period, a negative

relationship with *Clock, Npas2, Per1,2*, *Cry1,2*, and *Rev-erbb* suggested different regulation in

1xIMO and repeated IMO conditions. Additionally, the diurnal pattern of the *Nr3c1*/GR

expression in Leydig cells was changed ie increased in the 10xIMO, which contributes to the

different effect observed. However, many other signaling pathways directed by hormones,

especially LH, PRL, adrenaline, involved in highly multi-regulated processes such as gene

expression, could be affected by IMO and vary the circadian rhythm.

In summary, the results suggest different sensitivity to stress depending on the diurnal time of the

stressful event. Transcriptions of the clock and steroidogenic genes are much more affected by

stress in the rest than in the active phase, pointed to the relevance of the stress event's timing.

Both types of stress (1xIMO and 10xIMO) potentiates expression of clock elements that act over

E-box (BMAL1 and PER1) regardless of the phase of the day and decreased elements which

exert their action via RORE-box (*Rev-erba,b*) in rest period, suggesting a possible effect on gene

expression not only on the clock system but also outside it.

The obtained results pointed to stress and the circadian system's coordinated action in shaping

male reproductive physiology, emphasizing the importance of the Leydig cells' rhythmic activity

underpinning male fertility.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

M.LJ.M.— data acquisition; data analysis and interpretation; figures drafting; revising manuscript critically for important intellectual content; S.A.A.—acquisition of the data; analysis and interpretation of the data; revising manuscript critically for important intellectual content; T.S.K.—the conception and design of the research; acquisition of the data; analysis and interpretation of the data; drafting the manuscript; All authors—approved the final version of the manuscript; agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding

This research was supported by the Ministry of Education, Science and Technological Development Republic of Serbia grant no. 173057 and CeRES grant, and the Autonomic Province of Vojvodina, Serbia, grants no. 3822.

Acknowledgments

We are very grateful to Professor dr Gordon Niswender (Colorado State University) for supplying antibodies for radioimmunoassay analysis. Also we thank dr Aleksandar Baburski for technical assistance including data analysis as well as dr Milan Župunski for technical help in drafting figures.

References

1. Kennaway, D. J. The role of circadian rhythmicity in reproduction. *Hum Reprod Update* **11**,

    91–101 (2005).

2. Albrecht, U. Invited Review: Regulation of mammalian circadian clock genes. *Journal of*

    Applied Physiology **92**, 1348–1355 (2002).

3. Takahashi, J. S. Transcriptional architecture of the mammalian circadian clock. *Nat Rev*

    Genet **18**, 164–179 (2017).

4. Ukai, H. & Ueda, H. R. Systems Biology of Mammalian Circadian Clocks. *Annu. Rev.*

    Physiol. **72**, 579–603 (2010).

5. Alvarez, J. D. *et al.* The circadian clock protein BMAL1 is necessary for fertility and proper

testosterone production in mice. *J Biol Rhythms* **23**, 26–36 (2008).

6. Baburski, A. Z., Andric, S. A. & Kostic, T. S. Luteinizing hormone signaling is involved in

synchronization of Leydig cell's clock and is crucial for rhythm robustness of testosterone

production†. *Biol Reprod* **100**, 1406–1415 (2019).

7. Baburski, A. Z. *et al.* Circadian rhythm of the Leydig cells endocrine function is attenuated

during aging. *Experimental Gerontology* **73**, 5–13 (2016).

8. Baburski, A. Z. *et al.* Melatonin replacement restores the circadian behavior in adult rat

Leydig cells after pinealectomy. *Molecular and Cellular Endocrinology* **413**, 26–35 (2015).

9. Brambilla, D. J., Matsumoto, A. M., Araujo, A. B. & McKinlay, J. B. The effect of diurnal

variation on clinical measurement of serum testosterone and other sex hormone levels in

men. *J Clin Endocrinol Metab* **94**, 907–913 (2009).

10. Andric, S. A. *et al.* The opposite roles of glucocorticoid and α1-adrenergic receptors in stress

triggered apoptosis of rat Leydig cells. *American Journal of Physiology-Endocrinology and*

Metabolism **304**, E51–E59 (2012).

11. Ge, R.-S. *et al.* Stimulation of testosterone production in rat Leydig cells by aldosterone is

mineralocorticoid receptor mediated. *Molecular and Cellular Endocrinology* **243**, 35–42

(2005).

12. Marić, D., Kostić, T. & Kovačević, R. Effects of acute and chronic immobilization stress on

rat Leydig cell steroidogenesis. *The Journal of Steroid Biochemistry and Molecular Biology*

58, 351–355 (1996).

13. Rivier, C. & Rivest, S. Effect of Stress on the Activity of the Hypothalamic-Pituitary-

Gonadal Axis: Peripheral and Central Mechanisms. *Biol Reprod* **45**, 523–532 (1991).

14. Russell, G. & Lightman, S. The human stress response. *Nat Rev Endocrinol* **15**, 525–534

(2019).

15. Gibbs, F. P. Circadian variation of ether-induced corticosterone secretion in the rat. *Am J*

Physiol **219**, 288–292 (1970).

16. Zimmermann, E. & Critchlow, V. Effects of Diurnal Variation in Plasma Corticosterone

Levels on Adrenocortical Response to Stress. *Proceedings of the Society for Experimental*

Biology and Medicine **125**, 658–663 (1967).

17. Kvetnansky, R. & Mikulaj, L. Adrenal and Urinary Catecholamines in Rats During

Adaptation to Repeated Immobilization Stress. *Endocrinology* **87**, 738–743 (1970).

18. Andric, S. A., Janjic, M. M., Stojkov, N. J. & Kostic, T. S. Protein kinase G-mediated

stimulation of basal Leydig cell steroidogenesis. *American Journal of Physiology-*

Endocrinology and Metabolism **293**, E1399–E1408 (2007).

19. Kostic, T. S., Stojkov, N. J., Janjic, M. M. & Andric, S. A. Structural complexity of the testis

and PKG I / StAR interaction regulate the Leydig cell adaptive response to repeated

immobilization stress. *International Journal of Andrology* **33**, 717–729 (2010).

20. Payne, A. H., Downing, J. R. & Wong, K. L. Luteinizing hormone receptors and testosterone

synthesis in two distinct populations of Leydig cells. *Endocrinology* **106**, 1424–1429 (1980).

21. Medar, M. L., Andric, S. A. & Kostic, T. S. Stress-induced glucocorticoids alter the Leydig

cells' timing and steroidogenesis-related systems. *Molecular and Cellular Endocrinology*

538, 111469 (2021).

22. Stojkov, N. J. *et al.* Repeated immobilization stress disturbed steroidogenic machinery and

stimulated the expression of cAMP signaling elements and adrenergic receptors in Leydig

cells. *American Journal of Physiology-Endocrinology and Metabolism* **302**, E1239–E1251

(2012).

23. Sachs, M. Tools for estimating and predicting the cosinor model. R package version 1.1.

(2014).

24. Molcan, L. Time distributed data analysis by Cosinor.Online application. *bioRxiv* 805960

(2019) doi:10.1101/805960.

25. Dray, S. & Dufour, A.-B. The **ade4** Package: Implementing the Duality Diagram for

Ecologists. *J. Stat. Soft.* **22**, (2007).

26. Dinno, A. Tools for estimating and predicting the cosinor model. R package version 1.1.

(2018).

27. Kassambara, A. & Mundt, A. Extract and visualize the results of multivariate data analyses.

R package version 1.0.7. (2020) doi:https://CRAN.R-project.org/package=factoextra.

28. Dufau, M. L. The Luteinizing Hormone Receptor. *Annual Review of Physiology* **60**, 461–496

(1998).

29. Zirkin, B. R. & Papadopoulos, V. Leydig cells: formation, function, and regulation†. *Biology*

of Reproduction **99**, 101–111 (2018).

30. Chen, H. *et al.* Circadian clock and steroidogenic-related gene expression profiles in mouse

Leydig cells following dexamethasone stimulation. *Biochem Biophys Res Commun* **483**,

294–300 (2017).

31. Alvarez, J. D. *et al.* The Circadian Clock Protein BMAL1 Is Necessary for Fertility and

Proper Testosterone Production in Mice: *Journal of Biological Rhythms* (2008)

doi:10.1177/0748730407311254.

32. Ruginsk, S. G. *et al.* Glucocorticoid-Mediated Regulation of Circadian Rhythms: Interface

with Energy Homeostasis and Reproduction. in *Corticosteroids* (ed. Al-kaf, A. G.) (InTech,

2018). doi:10.5772/intechopen.73599.

33. Koch, C. E., Leinweber, B., Drengberg, B. C., Blaum, C. & Oster, H. Interaction between

circadian rhythms and stress. *Neurobiology of Stress* **6**, 57–67 (2017).

34. Kino, T. Circadian Rhythms of Glucocorticoid Hormone Actions in Target Tissues: Potential

Clinical Implications. *Sci. Signal.* **5**, pt4–pt4 (2012).

35. Oishi, K. *et al.* Genome-wide Expression Analysis Reveals 100 Adrenal Gland-dependent

Circadian Genes in the Mouse Liver. *DNA Research* **12**, 191–202 (2005).

36. Balsalobre, A. *et al.* Resetting of Circadian Time in Peripheral Tissues by Glucocorticoid

Signaling. *Science* **289**, 2344–2347 (2000).

37. Surjit, M. *et al.* Widespread Negative Response Elements Mediate Direct Repression by

Agonist- Liganded Glucocorticoid Receptor. *Cell* **145**, 224–241 (2011).

38. Torra, I. P. *et al.* Circadian and Glucocorticoid Regulation of Rev-erbα Expression in Liver*.

Endocrinology **141**, 3799–3806 (2000).