1

Messages From the Small Intestine Carried by Extracellular Vesicles

in Prediabetes: a Proteomic Portrait

Author(s) Names

Inês Ferreira[1,2,3#], Rita Machado de Oliveira[1#], Ana Sofia Carvalho[1], Akiko Teshima[1], Hans

Christian Beck[4], Rune Matthiesen[1], Bruno Costa-Silva[3,*,#], Maria Paula Macedo[1,5,6,*,#]

Author(s) Address Information

1. *Chronic Diseases Research Centre, CEDOC, NOVA Medical School, (NMS/FCM), Lisbon*
1169-056, Portugal
2. *Bioengineering – Cell Therapies and Regenerative Medicine PhD Program, Instituto*
Superior Técnico, University of Lisbon, Lisbon 1049-001, Portugal
3. *Champalimaud Research, Champalimaud Centre for the Unknown, 1400-038 Lisbon,*
Portugal
4. *Centre for Clinical Proteomics, Department of Clinical Biochemistry and Pharmacology,*
Odense University Hospital, Denmark
5. *APDP-ERC Portuguese Diabetes Association Education and Research Centre, Lisbon*
1250-189, Portugal
6. *Departament of Medical Sciences, University of Aveiro, Aveiro 3810-193, Portugal*
24

Symbol

[#] Equal contribution

[*] Corresponding author(s)

E-mail: bruno.costa-silva@research.fchampalimaud.org (Costa-Silva B),

paula.macedo@nms.unl.pt (Macedo MP)

CEDOC, NOVA Medical School / Faculdade de Ciências Medicas (NMS / FCM),

Universidade Nova de Lisboa,

Campo Mártires da Pátria, 130, 1169-056 Lisboa, Portugal

Phone: (+351) 21 880 3017

Fax: (+351) 21 880 3028

36
37

Abstract

Extracellular vesicles (EVs) mediate communication in physiological and pathological conditions. In the pathogenesis of type 2 diabetes, inter-organ communication plays an important role in its progress and metabolic surgery leads to its remission. Moreover, gut dysbiosis is emerging as a diabetogenic factor. However, it remains unclear how gut senses metabolic alterations and whether this is transmitted to other tissues via EVs. Using a diet induced-prediabetic mouse model, we observed that protein packaging in gut derived EVs (GDE), specifically small intestine, is increased in prediabetes. Proteins related to lipid metabolism and to oxidative stress management were more abundant in prediabetic GDE compared to healthy controls. On the other hand, proteins related to glycolytic activity, as well as those responsible for the degradation of polyubiquitinated composites, were depleted in prediabetic GDE. Together, our findings show that protein packaging in GDE is markedly modified during prediabetes pathogenesis. Thus, suggesting that prediabetic alterations in small intestine are translated into modified GDE proteome, which are dispersed into the circulation where they can interact with and influence the metabolic status of other tissues. This study highlights the importance of the small intestine as a tissue that propagates prediabetic metabolic dysfunction throughout the body and the importance of GDE as the messenger.

KEYWORDS: Small Instestines; Exosome; Type 2 diabetes; Lipids; Diet

Introduction

Type 2 diabetes (T2D) is the most common chronic metabolic disease characterized by high blood glucose and it accounts for 95% of diabetes cases in adulthood [1]. The escalating incidence of T2D as a consequence of sedentary lifestyle and over-nutrition, imposes a substantial burden to health care systems worldwide. Among the therapeutic interventions, metabolic surgery which targets the small intestine, is the most effective procedure in diabetes remission [2;3]. Prediabetes is a high-risk state for T2D, and it is defined by glycemic levels higher than normal, but lower than diabetic limits [4]. This stage can be seen as a fork in the road, if left untreated, the likelihood for progression to diabetes is high. Importantly, with timely diagnosis and with patient's compliance to lifestyle intervention, prediabetes can be reversed [5]. Obesity is a potent risk factor for diabetes. Moreover, a diet rich in fat and sugar, with large intake of finely processed grains and starchy carbohydrates is directly associated with this disease [6].

Blood glucose homeostasis requires a constant communication between insulin-secreting and insulin-sensitive organs. Failure in the coordination between these organs can lead to a rise in blood glucose levels and to prediabetes. Recently, extracellular vesicles (EVs) emerged as an important component of organismal communication and have been implicated in the pathophysiology of diabetes [7]. These small vesicles are able to transfer proteins and nucleic acids from a releasing cell to a distant target cell, thus modulating gene and protein expression and leading to functional changes [8-9]. Interestingly, visceral adipose tissue derived EVs trigger systemic inflammation, glucose intolerance and insulin resistance; hallmarks of prediabetes [10]. In obese patients, adipose tissue-EVs were able to modulate insulin responses in hepatocytes and muscle cells [11].

The intestinal system is daily exposed to food, millions of pathogens, and high concentrations of foreign antigens. In the case of the gut EVs, the composition and effects of EVs released by the resident microbiota have been explored [12-13]. For instance, these EVs are released into the gut mucosa and participate in local innate responses to invading bacteria through microbicidal activity [14] and mediate gut permeability through the regulation of tight junctions [15]. While the contribution of microbiota is crucial to maintain intestinal homeostasis, the epithelial cells are a critical component of the intestinal defense system forming a selective permeable barrier, allowing the absorption of nutrients, electrolytes and water, and blocking ingress of toxins and antigens. Epithelial cells in the gut were described to secrete EVs and up today they were described to play a role as antigen-carrying structures [16]. Whether EVs derived from the small intestine play a primordial role as a diabetogenic trigger needs to be elucidated. In fact, the relevance of the gastrointestinal tract in diabetes is elucidated by the fact that metabolic surgery improves glucose homeostasis more effectively than any known pharmaceutical or behavioral approach, causing durable remission in many patients with T2D [17]. Attempts to elucidate the

exact mechanism by which gastro-intestinal surgery reverses T2D have implicated changes in the
gut hormones, bile acid metabolism, intestinal nutrient sensing and metabolism, gut microbiota,
and other factors [18]. Gut dysbiosis recently emerged as a possible player in the development of
T2D [19]. Despite the literature on the impact of microbiota in T2D onset and development, the
role of the small intestine EVs, independent of EVs derived from the biota is still elusive. While
gut derived EVs (GDE) are considered to play an important role in intestinal diseases [20] it is
not known how GDE protein content is modulated by a western diet. Therefore, it is timely to
understand the importance of the small intestine as a metabolically active organ in prediabetes
and the means by which it communicates with other tissues via EVs. Here we aim at
characterizing the differences in the GDE proteome of diet-induced prediabetic mice compared
to that of healthy mice fed standard chow.

121
122

Results

Description of the prediabetes mouse model

First, we established a mouse model that recapitulates many features of prediabetes [21]. High fat
and sugar feeding leads to obesity, hyperinsulinemia and glucose intolerance. Therefore, we used
an hypercaloric diet (HD) which composition is 58% fat and 18 % sucrose, in contrast with
normal-chow diet (NCD) that is 11% fat with no added sugar. After 12 weeks of diet, HD mice
demonstrate significantly increased body weight (Figure 1A, 1B) and an abnormal intrahepatic
fat accumulation – a hallmark of liver steatosis (Figure 1C). They also developed glucose
intolerance, evaluated by a glucose tolerance test in the previous week of sacrifice and
correspondent area under the curve for each group (Figure 1D, 1E) (**Figure 1**).
133

Characterization of intestine derived extracellular vesicles

Once the prediabetic phenotype was validated in the diet-induced obese animals, we evaluated
the presence of EVs in small intestine. We stained cross sections for CD81, a commonly used
EV-marker, and the presence of this protein was detected in similar levels in the lumen of the
intestine in both conditions (NCD and HD fed mice - Figure 1F). Moreover, western blot was
performed to confirm the enrichment of CD63, a commonly used EV marker, in GDE from NCD
and HD when compared with the tissue (Figure 1G). Next, GDE were characterized by
nanoparticle tracking analysis and total amount of protein was quantified by BCA. The mean size
of the particles, in both conditions, was around 100 nm (NCD=152.8±67.5; HD=128.6±50.7)
(Figure 1H). While the number of EVs is decreased in HD animals (Figure 1I), protein cargo was
higher in GDE isolated from HD-fed mice (HD-GDE) than in NCD-fed (NCD-GDE) mice
(Figure 1J). Interestingly, the amount of protein per GDE was higher in prediabetic mice than in
controls (Figure 1K) (**Figure 1**).

Proteomics Study

EVs protein cargo provides valuable information regarding the cellular environment where they are produced [22]. We proposed that a diet induced dysmetabolic environment modulates the protein composition of GDE. Thus, to identify the impact of the diet-induced prediabetic gut milieu on GDE protein cargo, we isolated GDE (Supplementary Figure1) from mice under HD and respective controls and analysed it by mass-spectrometry (MS) (Supplementary Figure 2A). Potential contribution of gut bacteria EVs was ruled out by mass spectrometry analysis, which showed absence of bacterial proteins (Supplementary Figure 2B), and culture of gut conditioned media in antibiotics-free agar, which did not display bacterial growth (Supplementary Figure 2C).

Bioinformatics analysis of the MS identified proteins

After filtering the identified proteins for reverse proteins used for false discovery rate and possible contaminating proteins (e.g. keratin and bovine proteins from cell culture media), we identified a total of 2467 proteins after removing isoforms from the same gene. Intensity based absolute quantitation (iBAQ) was calculated [23]. Excluding proteins with missing quantification values resulted in 2156 proteins (Supplementary file). The raw data was then normalized (Figure 2A). Subsequently, we distinguished between identified and regulated proteins. Regulated proteins are those at least two-fold regulated and with an adj. p-value < 0.05. After applying the aforementioned criteria to our data set, we obtained a list of robustly regulated 558 proteins, which shows that diet-induced prediabetes alters the protein composition of GDEs (Figure 2A). All data, before applying the filtering criteria, is represented in a volcano plot that describes the level of significance and magnitude of changes observed, comparing the HD with the NCD group (Figure 2B). The dotted line in the plot represents the threshold of adj.p-value<0.05, and from that line above each individual dot represents a protein within the 558 regulated ones.

Gene Ontology

To gain insight into the biological role of the 588 regulated proteins, we performed Gene Ontology and Kyoto Encyclopedia for Genes and Genomes (KEGG) enrichment analysis, which revealed that the vast majority of proteins were related with metabolism. In terms of molecular function, enzyme activity (including isomerases, hydrolases and oxidoreductase) shows up in the top of list (Figure 3A). Most of the enriched pathways for gene ontology biological process emphasized lipid metabolism, highlighting fatty acids (FA) β-oxidation. Curiously, glycolysis-related and proteolysis in cellular protein catabolic processes-related proteins are mostly down regulated (Figure 3B). Considering KEGG pathways enrichment analysis, the affected proteins belong primarily to FA, carbohydrate and amino acid metabolism. Interestingly, alcoholism-related proteins show up with great significance (Figure 3C). In terms of down regulated proteins, based on the KEGG analysis the proteasome and focal adhesions proteins are markedly down regulated. Finally, to assess sub-cellular localization of the GDE protein cargo regulated by diet

185 we used gene ontology cellular component analysis and found that the majority of altered proteins
186 are mitochondrial, followed by the cytosolic ones. Once again, proteasome related proteins were
187 down regulated (Figure 3D).
188
Cluster analysis for GDE regulated proteins
For a more detailed characterization of GDE proteome, we looked into the subset of proteins that
were overly altered among groups. For that, we established an adj.p-value<0.0001 as cut-off
(Figure 4A). The proteins with that significance are represented in the heatmap (Figure 4B). To
gain insight into protein-protein interaction of GDE identified protein cargo, we used the STRING
software to link some of the altered proteins and to display the level of relationship between them.
The proteins that demonstrate higher level of interaction are the ones related with the proteasome,
including both α subunits (PSMA) and β (PSMB). Actin related proteins, which function is to
maintain cellular integrity, also display relevant interaction. Moreover, network analysis
highlights strong tendency towards metabolic-related proteins (Supplementary Figure 2). The fold
change of all altered proteins varied up to 2-fold when comparing HD with NCD GDE. Within
metabolic pathways, the altered proteins are mainly involved in nutrient metabolism, such as
glucose, FAs and amino acids. In addition, cell integrity, focal adhesion and proteasome-related
proteins are all down regulated (Figure 4C and Supplementary Figure 4). Key GDE proteins up
and down regulated by HD are also represented in their subcellular location (**Figure 5**).
Noteworthy to mention, proteins related with carbohydrate metabolism, which ensures the supply
of energy to the cells (PFKL, ALDOA, ALDOB, PGAM2, ENO1, PGM5), with the tricarboxylic
acid (TCA) cycle (DLST, IDH1). cholesterol homeostasis (ACAT1, ACAT2) and proteasome
(PSMA1, PSMA2, PSMA3, PSMA4, PSMA5, PSMA6, PSMA7, PSMB1, PSMB2, PSMB3,
PSMB8, PSMB10) are down regulated by HD. On the other hand, proteins related with lipid
homeostasis (ACOT1 ACOT2, ACOT3, ACOT5, ACOT6), FA mitochondrial β-oxidation
(HDHA, ACADVL, ACADL, ECHS1, ACAA2), very-long chain FA peroxisomal β-oxidation
(ECH1, ACOX1), amino acids metabolism (GSTA1, GSTA2) and antioxidant defence (ALDH2,
ALDH1B1, PRDX6, GSTA4, GSTM2) are majorly upregulated. **Table 1** describes all the
proteins up and down regulated according to their biological function.
214
215

Discussion
The relationship between diet composition and the incidence of metabolic disorders is a public
health concern. One of the first organs and by far the largest one to contact with food and its
derivates is the gut, as a consequence it is where the first organismal response should be found
[24]. Here, we report that GDE isolated from animals fed an HD, mimicking the western one,
have a distinct proteome that strongly reflects the intestinal dysmetabolic milieu. We found

numerous pathways associated with macronutrients altered in GDE isolated from HD-fed mice;
of great relevance the ones related with lipidic and carbohydrate metabolic processes.
224

From the enrichment analysis we observed that both glycolysis-gluconeogenesis and pyruvate
pathways are within the most affected ones (Figure 3B). Rate limiting enzymes for glycolysis
were found to be less abundant, namely hexokinase (HK) and phosphofructokinase (PFKL), thus
suggesting changes in glucose utilization and lactate production. In accordance, in diabetic rats,
phosphorylation of glucose is inhibited in tissues such as the heart, skeletal muscle and liver [25;
26; 27; 28]. Interestingly, in hearts from diabetic rats, PFK and HK down regulation is reverted
by an oral anti-diabetic drug - metformin [29]. We suggest that the observed decreased levels of
PFKL in prediabetic GDEs is reflecting a similar down regulation in the small intestine of
prediabetic mice and is an early event in the progression of the disease [30]. Down in the
glycolysis pathway, pyruvate dehydrogenase A1 (PDHA1), which belongs to the PDH complex,
was upregulated in HD GDEs, indicating a possible compensation in the pathway given the
importance of acetyl-CoA for the TCA cycle and for many other proteins and lipidic biochemical
reactions. PDH is fundamental for metabolic flexibility, the ability of switching from
carbohydrates to lipid fuel as energy sources, and the inability of cells to do so is an important
hallmark of T2D [31]. Our results suggest that in the early stages of the disease gut cells tackle
the insult of HD feeding by changing the source of fuel. The decrease in glycolysis might be
explained by the low amount of fibers provided by the diet; since the percentage of fibers in HD
is considerably reduced and in part replaced by sucrose. Likewise, fibers have a beneficial effect
on NCD mice, by improving intestinal integrity, enhancing energy expenditure and reducing
inflammation [32]. In fact, because HD has increased sucrose, we envisage that sucrose is being
directed to an alternative pathway, such as de novo lipogenesis, known to also occur in the gut
[33].
247

In accordance with the high lipidic content present in the HD, we observed that lipid metabolism
(biosynthesis of unsaturated FA and FA elongation) is vastly altered. The excess of fat in the diet
is known to cause increased fatty acids uptake, which can be either metabolized for the production
of energy or, when in excess, will be stored. Several proteins involved in mitochondrial β-
oxidation (HADHA, ACADVL, ACADL, ECH1, ECHS1), which generates acetyl-CoA to feed
the TCA cycle, and peroxisome β-oxidation (ECH1, ACOX1), major fatty acids degradative
pathway, are upregulated in GDE by HD, suggesting that fatty acids are favoured as energy source
over glucose (Figure 5). In addition, over-activation of lipid metabolism related pathways is a
common feature of obesity and T2D, leading to a state of hyperglycemia [34; 35; 36]. Moreover,
a family of acyl-CoA thioesterases (ACOTs) were particularly upregulated; these are
intermediaries in the decision whether fatty acids are directed to the TCA cycle or stored, (Figure
5, Supplementary Figure 4) [37; 38]. Altogether, this suggests that HD shifts the burden of energy

provision from carbohydrates toward fat derived from diet, since ACSS2 (a protein that conducts fatty acids to the TCA cycle) is down regulated and PDHA1 (which provides a link between glycolysis, TCA cycle and lipogenesis) is upregulated in order to ensure the energetic need of the cell. Based on recent literature, obesity-induced activation of ACOTs directs fatty acids towards triglycerides synthesis for incorporation into chylomicrons or VLDL particles [39; 40]. Noteworthy, some studies consider activation of ACOTs protective against fatty acids over-supply, slowing the flux of fatty acids to downstream metabolic pathways [41; 42; 43]. Whether this is the case in the gut, needs to be further explored. Additionally, intestinal cells in the HD model appear to prioritize production of energy from fatty acids, instead of carbohydrates. This is shown by the upregulation of ATP-citrate lyase (ACLY), an enzyme responsible for the conversion of citrate to acetyl-CoA – the first step in DNL. Overall, the major enzymes found altered within the gut EVs are very much responsible for regulating acetyl-CoA levels in the cells, given that this molecule is a fundamental node in metabolism, in the future it would be of major interest to quantify it in enterocytes.

Increased fatty acids oxidation promotes mitochondrial dysfunction, as well as fatty acids peroxidation and glucose dysmetabolism, altogether leading to reactive oxygen species (ROS) accumulation to pernicious levels [44]. Oxidative stress is implicated in various diseases including diabetes [45]. We observed several proteins involved in antioxidant defense mechanisms, such as removal of free radicals (PRDX6) and elimination of xenobiotic compounds and products of lipids peroxidation (GSTA4, GSTM2) upregulated in HD-GDE, suggesting an activation of an antioxidant response in the gut cells (Figure 5). Consistent with our model, the presence of numerous isoforms of the aldehyde dehydrogenase (ALDH) family was upregulated in HD-derived GDE. ALDHs are a family of oxidizing enzymes responsible for cellular detoxification by oxidation of biogenic and xenogeneic aldehydes that accumulate through metabolism [46]. In particular, we highlight ALDH1B1 due to its ability to metabolize acetaldehyde and maintain glucose homeostasis [46]. Of interest, ALDH2 has been explored as a pharmacological target, due to its role in cellular damage defense against oxidative stress induced by pathological conditions, for instances high glucose [47].

Besides glucose and fatty acids, amino acids, in particular lysine, also play important roles in providing energy to gut epithelial cells (Figure 3C) [48]. Lysine, which pathway we see affected in HD-GDE, mediates protein biosynthesis, such as of carnitine, a key factor in fatty acid metabolism [49]. Interestingly, carnitine deficiency due to poor lysine diet induces triglycerides accumulation [50]. Evidence that the HD not only induces obesity, but also a general pathological state arises from the massive alteration of proteasome-related proteins. The proteasome is recognized as an important target in many diseases [51]. Our data showed down regulation of proteasome proteins, which is consistent with decreased proteasome function reported in a broad

298 array of chronic diseases (Figure 3C; 5) and defective degradation of damaged proteins, a
299 hallmark of neurodegenerative diseases [52]. One hypothesis is that the decreased content of
300 proteasome proteins in prediabetic GDE mirror in the cell of origin. Alternatively, the GDE
301 producing cells can be consuming proteasome to counteract the deleterious effects of oxidative
302 stress, resulting in lower packaging of proteasome proteins in GDEs.
303
The present study is the first comprehensive analysis of GDE protein content. We found that diet
composition strongly modulates GDE proteomic profile. Several lines of evidence support a
model in which the metabolic alterations induced by the HD are translated in a similar manner in
alterations in GDE protein cargo. These results bring light into the important modification taking
place in the small intestine and how it translates into GDE content. We propose that future studies
should focus on whether GDE can act in local and distant cellular targets and mediate prediabetes.
We also propose that the analysis of GDE in biofluids may represent a strategy to non-invasively
detect metabolic modifications in the gut linked with prediabetes and potentially be employed
early diagnose and follow-up prediabetes patients.
313
314

Material and methods

Animals

Male C57Bl/6J mice were housed in a temperature-controlled room, in a 12-hour light/dark cycle.
For induction of prediabetic phenotype, C57BL/6J mice started on a high-fat diet at 6 weeks of
age, with free access to food and water, for 12 weeks. Hypercaloric diet (HD) (OpenSource Diet,
D12331) composition is 16.5% protein, 25.5% carbohydrate, and 58% fat with 13% addition of
sucrose. The control normal chow diet (NCD) (Special Diets Services, RM3) composition is
26.51% proteins, 62.14% carbohydrates and 11.35% fat and no sucrose. At 18 weeks of age, 12
weeks of diet, mice were sacrificed. Mice were anesthetized using 2% isoflurane. Mice were
sacrificed under 12 hours of fasting, with a previous feeding period of 2 hours, with the previous
8 hours of fasting. Blood, liver and small intestine were collected from prediabetic mice and
respective controls. All animals were treated according with National and European Union
Directive for Protection of Vertebrates.
328

Glucose Tolerance Test

330 At week 11 of diet GTT was performed. Mice were fasted overnight and weighed in the morning.
331 Mice were then given an intra-peritoneal injection of a glucose solution (20% m/v – Sigma
332 Aldrich) at 2g/kg body weight. Serum blood glucose levels were measured at 15, 30, 60, 90 and
333 120 min after injection. Blood glucose levels we measured using OneTouch Ultra glucose meter
334 (LifeScan Inc).

Small Intestine Histology

After sacrifice, liver and small intestine were washed in PBS and fixed for histology and a portion of small intestine was snap frozen in liquid nitrogen for later analysis by western blot. The clean gut was placed in 2% Paraformaldehyde with 20% Sucrose, overnight at 4ºC; washed three times in Phosphate Buffered Saline (PBS), 10 min each; transferred to 30% sucrose for 2 hours at 4ºC; left overnight at 4ºC in Optimal Cutting Temperature compound (OCT) with 30% Sucrose (1:1). In the following day, tissues were embedded in 20% Sucrose (1:4) + (OCT) (3:4) in the presence of dry ice and immediately stored -80ºC. Consequently, tissues were sliced into 6 μm thin sections in the cryostat and stored at -80ºC for following applications. Gut samples were used for CD81 staining and liver samples for Hematoxylin-Eosin staining.

Exosomes Isolation and Nanoparticle Tracking Analysis

After sacrifice the small intestine was removed, washed with PBS and cultured overnight, at 37ºC and 5%CO2, in RPMI medium (Sigma) supplemented with 1% Penicillin/Streptomycin and 10% Fetal Bovine Serum EV-free, to avoid contaminations from serum EVs. In the following day medium was collected. Conditioned medium was submitted to two initial centrifugations (10 min, 500 g and 20 min, 3 000 g) to remove any suspended or dead cells in the medium. To remove large EVs, media was centrifuged (20 min, 12 000 g) and the pellet was discarded. The supernatant enriched in small EVs was again centrifuged (2 h 20 min, 100 000 g), and the EVs enriched pellet was collected. For sucrose cushion purification, this pellet was resuspended in 14 mL filtered phosphate buffered saline (PBS, Corning 15313581, NY, US) and added the top of 4 mL sucrose solution (D2O containing 1.2 g of protease-free sucrose and 96 mg of Tris base adjusted to pH 7.4). A new ultracentrifugation was performed (1 h 10 min, 100 000 g), after which 4 mL of the sucrose fraction was collected using a 18 G needle placed at the bottom of the ultracentrifugation tube (away from the pellet). Finally, 16 mL of PBS were added to the collected sucrose/EVs solution and an overnight (16 h, 100 000 g) ultracentrifugation was performed. The pellet containing the isolated EVs was resuspended in filtered PBS. The concentration and size of EVs were analyzed in NanoSight NS300 (NS3000) system equipped with a blue laser (405 nm), according to the manufacturer's instructions. EVs were diluted 1:1000 in filtered sterile PBS. Each sample analysis was conducted for 90 seconds and measured 5 times using Nanosight automatic analysis settings. The software calculated the size distribution in nanometers and the concentration in number of particles per mL.

Western Blotting

Protein extracts for western blot analysis were obtained using cold lysis buffer (20 mM Tris- HCl pH 7.4, 5 mM EDTA pH 8.0, 1% Triton-X 100, 2 mM Na3VO4, 100 mM NaF, 10 mM Na4P2O7)

in the presence of protease inhibitors (completeTM, Mini, EDTA-free Protein inhibitor cocktail

tablets, Roche, Sigma). Gut tissue and GDE were homogenized in the lysis buffer and sonicated

three time for 30 seconds each at 10 μm amplitude, incubated on ice between each sonication

step. Lysates were centrifuged at 18.000 g for 10 min at 4°C. Soluble fraction was collected and

protein concentration was determined using the PierceTM BCA Protein Assay kit (Thermo

Fisher). For protein expression analysis, equal amounts of total protein (20 μg) were mixed with

sample buffer (250 mM Tris-HCl pH 6.8, 8% SDS, 40% glycerol, 8% β-mercaptoethanol,

bromophenol), before being denaturated at 95ºC for 10 min. Samples were then resolved by an

8% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Proteins were

subsequently electrotransferred to a polyvinylidene fluoride (PVDF) membrane (Immobilon-P

Membrane, PVDF, Millipore) for 20 min at 25 mA and 2,5 mV in a Trans-Blot Turbo TM

Transfer System (BioRad). Membranes were blocked for 1 hour at RT with 3% BSA (Thermo

Fisher) in tris-buffered saline with 0,1% Tween (TBS-T) (blocking solution). Membranes were

incubated overnight at 4ºC, with CD63 antibody (Santa Cruz Biotechnology) prepared in 3% BSA

in TBS-T, diluted 1:1000. Membranes were washed three times with TBS-T and incubated for 1

hour at RT with the secondary antibody anti-rabbit antibody diluted 1:5000 (Santa Cruz

Biotechnology). Blots were developed with ECL (ECL Prime, GE Haelthcare) according to

manufacturer's instructions. Chemidoc Touch (Biorad) was used to detect chemilumunescence.

Band intensities were quantified usinf ImageLab software and normalized by β-actin as a loading

control.

Nano-LC-MSMS analysis

Peptide samples were analysed by nano-LC-MSMS (Dionex RSLCnano 3000) coupled to a Q-

Exactive Orbitrap mass spectrometer (Thermo Scientific). Briefly, 5 μL of sample was loaded

onto a custom made fused capillary pre-column (2 cm length, 360 μm OD, 75 μm ID) with a flow

of 5 μL per min for 7 min. Trapped peptides were separated on a custom made fused capillary

column (20 cm length, 360 μm outer diameter, 75 μm inner diameter) packed with ReproSil Pur

C18 3-μm resin (Dr. Maish, Ammerbuch- Entringen, Germany) with a flow of 300 nL per minute

using a linear gradient from 92 % A (0.1% formic acid) to 28 % B (0,1% formic acid in 100

acetonitrile) over 93 min followed by a linear gradient from 28 % B to 35 % B over 20 min at a

flowrate of 300 nl per minute. Mass spectra were acquired in positive ion mode applying

automatic data-dependent switch between one Orbitrap survey MS scan in the mass range of 400

to 1200 m/z followed by HCD fragmentation and Orbitrap detection of the 15 most intense ions

observed in the MS scan. AGC target value in the Orbitrap for MS and MSMS scans were

1,000,000 ions at a resolution of 70,000 at m/z 200 for MS scans, and 25,000 ions at m/z 200 for

MSMS scans. Peptide fragmentation in the HCD cell was performed at normalized collision

energy of 31 eV. Ion selection threshold was set to 5,000 ions for MSMS analysis, and maximum

injection time was 100 ms for MS scans and 200 ms for MSMS scans. Selected sequenced ions
were dynamically excluded for 60 seconds.
410

MSMS analysis

Mass accuracy was set to 5 ppm on the peptide level and 10 Da on the fragment ions. A maximum
of four missed cleavages was used. Carbamidomethyl was set as a fixed modification. M
oxidation, N-terminal protein Acetyl, Q Deamidation and N Deamidation were set as variable
modifications. The MSMS data was searched against all reviewed mouse proteins from UniProt
with concatenation of all the sequences in reverse maintaining only lysine and arginine in place.
The data was searched and quantified with both MaxQuant [53] and VEMS [54].
418

Statistical Analysis

420 Data are presented as means ± SEM. GTT curves, bar plots, volcano plots, heatmap and linear
421 regression analyses were done using GraphPad Prism 8 (GraphPad Software). Differences
422 significance was calculated through unpaired student's t tests with Welch's correction; one-way
423 ANOVA followed by Tukey-Kramer multiple comparison tests. Differences were accepted as
424 statistically significant at $p < 0.05$.
425

Author's contributions

IF and RMO carried out the animal model of disease, western blotting and drafted the manuscript.
IF did the isolation of the extracellular vesicles and characterization by NTA; gut histology and
performed bioinformatics analysis, together with MS, for graphical representation. ASC prepared
GDE for MS analysis and RM was responsible for the mass spectrometry analysis and
bioinformatics. Mass spectrometry was performed by HCB. MPM and BCS conceived the study,
participated in its design and coordination and edited the manuscript. All authors read and
approved the final manuscript.
435

Competing interests

437 The authors have declared no competing interests, except for MPM who serves on advisory board
438 for Gilead Sciences.
439

Acknowledgements

This work was supported by the Fundação para a Ciência e Tecnologia (Grant No. PTDC/MEC-
MET/29314/2017) and Programa Gilead GÉNESE - Edição de 2019 awarded to MPM, RMO and
BCS. IF is a recipient of Fundação para a Ciência e Tecnologia fellowship (PD/BD/114044/2015).
AT is a recipient a European Foundation for the Study od Diabetes and Japonese Diabetes Society
Reciprocal Travel Research Fellowship Programme.

References

[1] IDF. IDF Diabetes Atlas Ninth. Dunia: IDF. 2019. 168 p.

[2] Rubino F, Nathan DM, Eckel RH, Schauer PR, Alberti KGMM, Zimmet PZ, et al.
Metabolic surgery in the treatment algorithm for type 2 diabetes: A joint statement by
international diabetes organizations. Diabetes Care. 2016;39(6):861–77.

[3] Aminian A, Vidal J, Salminen P, Still CD, Hanipah ZN, Sharma G, et al. Late relapse of
diabetes after bariatric surgery: Not rare, but not a failure. Diabetes Care. 2020;43(3):534–40.

[4] Petersen MC, Shulman GI. Mechanisms of insulin action and insulin resistance. Physiol
Rev. 2018;98(4):2133–223.

[5] Tabák AG, Herder C, Rathmann W, Brunner EJ, Kivimäki M. Prediabetes: A high-risk state

for diabetes development. Lancet [Internet]. 2012;379(9833):2279–90

[6] Hamdy O, Tasabehji MW, Elseaidy T, Tomah S, Ashrafzadeh S, Mottalib A. Fat Versus
Carbohydrate-Based Energy-Restricted Diets for Weight Loss in Patients With Type 2 Diabetes.
Curr Diab Rep. 2018;18(12).

[7] Ban LA, Shackel NA, McLennan S V. Extracellular vesicles: A new frontier in biomarker
discovery for non-alcoholic fatty liver disease. Int J Mol Sci. 2016;17(3).

[8] Zhang J, Li S, Li L, Li M, Guo C, Yao J, et al. Exosome and exosomal microRNA:
Trafficking, sorting, and function. Genomics, Proteomics Bioinforma. 2015;13(1).

[9] Théry C, Zitvogel L, Amigorena S. Exosomes: composition, biogenesis and function. Nat
Rev Immunol. 2002;2(8):569–79.

[10] Deng Z, Poliakov A, Hardy RW, Clements R, Liu C, Liu Y, et al. Adipose Tissue
Exosome-Like Vesicles Mediate Activation of Macrophage-Induced Insulin Resistance.
2009;58 (November).

[11] Kranendonk MEG, Visseren FLJ, Van Herwaarden JA, Nolte-'t Hoen ENM, De Jager W,
Wauben MHM, et al. Effect of extracellular vesicles of human adipose tissue on insulin
signaling in liver and muscle cells. Obesity. 2014;22(10):2216–23.

[12] Chang X, Wang SL, Zhao SB, Shi YH, Pan P, Gu L, et al. Extracellular vesicles with
possible roles in gut intestinal tract homeostasis and IBD. Mediators Inflamm. 2020;2020.

[13] Smythies LE, Smythies JR. Exosomes in the gut. Front Immunol. 2014;5(MAR):2–5.

[14] Hu G, Gong AY, Roth AL, Huang BQ, Ward HD, Zhu G, et al. Release of Luminal
Exosomes Contributes to TLR4-Mediated Epithelial Antimicrobial Defense. PLoS Pathog.
2013;9(4).

[15] Chelakkot C, Choi Y, Kim DK, Park HT, Ghim J, Kwon Y, et al. Akkermansia
muciniphila-derived extracellular vesicles influence gut permeability through the regulation of
tight junctions. Exp Mol Med [Internet]. 2018;50(2):e450-11.

[16] Mallegol J, Van Niel G, Heyman M. Phenotypic and functional characterization of
intestinal epithelial exosomes. Blood Cells, Mol Dis. 2005;35(1):11–6.

[17] Evers SS, Sandoval DA, Seeley RJ. The Physiology and Molecular Underpinnings of the
Effects of Bariatric Surgery on Obesity and Diabetes. Annu Rev Physiol. 2017;79(1):313–34.

[18] Singh AK, Singh R, Kota SK. Bariatric surgery and diabetes remission: Who would have
thought it? Indian J Endocrinol Metab. 2015;19(5):563–76.

[19] David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet
rapidly and reproducibly alters the human gut microbiome. Nature [Internet].
2014;505(7484):559–63.

[20] Burcelin R, Serino M, Chabo C, Blasco-Baque V, Amar J. Gut microbiota and diabetes:
From pathogenesis to therapeutic perspective. Acta Diabetol. 2011;48(4):257–73.

[21] Kowalski GM, Bruce CR. The regulation of glucose metabolism: Implications and
considerations for the assessment of glucose homeostasis in rodents. Am J Physiol - Endocrinol
Metab. 2014;307(10):E859–71.

[22] Kalra H, Drummen GPC, Mathivanan S. Focus on extracellular vesicles: Introducing the
next small big thing. Int J Mol Sci. 2016;17(2).

[23] Schwanhüusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, et al. Global
quantification of mammalian gene expression control. Nature. 2011;473(7347):337–42.

[24] Abumrad NA, Davidson NO. Role of the gut in lipid homeostasis. Physiol Rev.
2012;92(3):1061–85.

[25] Amir, Weber, Beard, Bomyea T. NIH Public Access. Bone [Internet]. 2008;23(1):1–7.

[26] Kipnis DM, Cori CF. Studies of Tissue Permeability. J Biol Chem [Internet].
1960;235(11):3070–5.

[27] Chernick SS, Chaikoff IL, Abraham S. Localization of initial block in glucose metabolism
in diabetic liver slices. J Biol Chem. 1951;193(2):793-802.
506
[28] Renold AE, Hastings AB, Nesbett FB. Studies on carbohydrate metabolism in rat liver
slices. III. Utilization of glucose and fructose by liver from normal and diabetic animals. J Biol
Chem. 1954;209(2):687-696.

[29] Da Silva D, Ausina P, Alencar EM, Coelho WS, Zancan P, Sola-Penna M. Metformin
reverses hexokinase and phosphofructokinase downregulation and intracellular distribution in
the heart of diabetic mice. IUBMB Life. 2012;64(9):766–74.

[30] Randle PJ, Garland PB, Hales CN, Newsholme EA. The Glucose Fatty-Acid Cycle Its
Role in Insulin Sensitivity and the Metabolic Disturbances of Diabetes Mellitus. Lancet.
1963;281(7285):785–9.

[31] Kelley DE, Mandarino LJ. Fuel selection in human skeletal muscle in insulin resistance: A
reexamination. Diabetes. 2000;49(5):677–83.

[32] Jangra S, Raja SK, Sharma RK, Pothuraju R, Mohanty AK. Ameliorative effect of
fermentable fibres on adiposity and insulin resistance in C57BL/6 mice fed a high-fat and
sucrose diet. Food Funct. 2019;10(6):3696–705.

[33] Hoffman S, Alvares D, Adeli K. Intestinal lipogenesis: How carbs turn on triglyceride
production in the gut. Curr Opin Clin Nutr Metab Care. 2019;22(4):284–8.

[34] Deng WJ, Nie S, Dai J, Wu JR, Zeng R. Proteome, phosphoproteome, and
hydroxyproteome of liver mitochondria in diabetic rats at early pathogenic stages. Mol Cell
Proteomics. 2010;9(1):100–16.

[35] Kanuri BN, Rebello SC, Pathak P, Agarwal H, Kanshana JS, Awasthi D, et al. Glucose and
lipid metabolism alterations in liver and adipose tissue pre-dispose p47phox knockout mice to
systemic insulin resistance. Free Radic Res [Internet]. 2018;52(5):568–82.

[36] Knebel B, Hartwig S, Haas J, Lehr S, Goeddeke S, Susanto F, et al. Peroxisomes
compensate hepatic lipid overflow in mice with fatty liver. Biochim Biophys Acta - Mol Cell
Biol Lipids [Internet]. 2015;1851(7):965–76.

[37] Wang F, Wu J, Qiu Z, Ge X, Liu X, Zhang C, et al. ACOT1 expression is associated with
poor prognosis in gastric adenocarcinoma. Hum Pathol [Internet]. 2018;77:35–44.

[38] Steensels S, Qiao J, Zhang Y, Maner-Smith KM, Kika N, Holman CD, et al. Acot9 traffics
mitochondrial short-chain fatty acids towards de novo lipogenesis and glucose production in the
liver. Hepatology. 2020;0–3.

[39] Alves-Bezerra M, Li Y, Acuña M, Ivanova AA, Corey KE, Ortlund EA, et al. Thioesterase
Superfamily Member 2 Promotes Hepatic VLDL Secretion by Channeling Fatty Acids Into
Triglyceride Biosynthesis. Hepatology. 2019;70(2):496–510.

[40] Ersoy BA, Maner-Smith KM, Li Y, Alpertunga I, Cohen DE. Thioesterase-mediated
control of cellular calcium homeostasis enables hepatic ER stress. J Clin Invest.
2018;128(1):141–56.

[41] Yamaguchi K, Yang L, McCall S, Huang J, Xing XY, Pandey SK, et al. Inhibiting
triglyceride synthesis improves hepatic steatosis but exacerbates liver damage and fibrosis in
obese mice with nonalcoholic steatohepatitis. Hepatology. 2007;45(6):1366–74.

[42] Fujita M, Momose A, Ohtomo T, Nishinosono A, Tanonaka K, Toyoda H, et al.
Upregulation of fatty acyl-CoA thioesterases in the heart and skeletal muscle of rats fed a high-
fat diet. Biol Pharm Bull. 2011;34(1):87–91.

[43] Franklin MP, Sathyanarayan A, Mashek DG. Acyl-CoA thioesterase 1 (ACOT1) regulates
PPARα to couple fatty acid flux with oxidative capacity during fasting. Diabetes.
2017;66(8):2112–23.

[44] Volpe CMO, Villar-Delfino PH, Dos Anjos PMF, Nogueira-Machado JA. Cellular death,
reactive oxygen species (ROS) and diabetic complications review-Article. Cell Death Dis.
2018;9(2).

[45] Luo W, Xu Q, Wang Q, Wu H, Hua J. Effect of modulation of PPAR-γ activity on Kupffer
cells M1/M2 polarization in the development of non-alcoholic fatty liver disease. Sci Rep.
2017;7(March):1–13.

[46] Yang CK, Wang XK, Liao XW, Han CY, Yu TD, Qin W, et al. Aldehyde dehydrogenase 1
(ALDH1) isoform expression and potential clinical implications in hepatocellular carcinoma.
PLoS One. 2017;12(8):1–13.

[47] Chen CH, Ferreira JCB, Gross ER, Mochly-Rosen D. Targeting aldehyde dehydrogenase 2:
New therapeutic opportunities. Physiol Rev. 2014;94(1):1–34.

[48] Van Goudoever JB, Stoll B, Henry JF, Burrin DG, Reeds PJ. Adaptive regulation of
intestinal lysine metabolism. Proc Natl Acad Sci U S A. 2000;97(21):11620–5.

[49] Azevedo C, Saiardi A. Why always lysine? The ongoing tale of one of the most modified
amino acids. Adv Biol Regul [Internet]. 2016;60:144–50.

[50] Khan L, Bamji MS. Tissue Carnitine Deficiency due to Dietary Lysine Deficiency:
Triglyceride Accumulation and Concomitant Impairment in Fatty Acid Oxidation. J Nutr.
1979;109(1):24–31.

[51] Liu H, Yu S, Xu W, Xu J. Enhancement of 26S proteasome functionality connects
oxidative stress and vascular endothelial inflammatory response in diabetes. Arterioscler Throm
Vasc Biol. 2012; 32(9):2131-2140.

[52] Rousseau A, Bertolotti A. Regulation of proteasome assembly and activity in health and
disease. Nat Rev Mol Cell Biol [Internet]. 2018;19(11):697–712.

[53] Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-
range mass accuracies and proteome-wide protein quantification. Nat Biotechnol.
2008;26(12):1367–72.

[54] Carvalho AS, Ribeiro H, Voabil P, Penque D, Jensen ON, Molina H, et al. Global mass
spectrometry and transcriptomics array based drug profiling provides novel insight into
glucosamine induced endoplasmic reticulum stress. Mol Cell Proteomics. 2014;13(12):3294–
307.

Figure legends

Figure 1 High fat diet induces prediabetes in mice and increases protein cargo in gut derived EV

A. Diet plan schematic representation. **B**. Body weight (n=20 per group). **C**. Representative Hematoxylin-Eosin images of liver histological sections of NCD and HD mice. White areas indicated by black arrows are lipid droplets, revealing steatosis in the liver of mice subjected to HD. **D.** Plasma glucose profile (md/dl) along a glucose tolerance test (GTT) at different time points after glucose bolus (0, 15, 30, 60, 90, 120 minutes) at week 17 of age. **E**. Mean area under the curve of the GTT. Results are expressed as mean $\pm$ SEM; n=20 per group. Unpaired t test with Welch's correction for B and D; ****$p<0.0001$. **F**. Gut cross sections, EVs are labelled in green (CD81) and nuclei in blue (DAPI). NCD gut tissue in the left and HD gut tissue in the right. CD81 quantification in the gut. **G**. Immunoblotting of gut tissue and gut derived EV (GDE) isolated from NCD and HD mice. Protein extracts prepared from gut (10 μg) and from GDE (15 μg) were analyzed by western blotting using antibodies against EV marker CD63 and β-actin. **H**. Size and concentration distribution of GDE determined by nanoparticle tracking analysis (NTA). A representative NTA graph of GDE of mice fed with NCD (NCD-EV) (left) and HD (HD-EV) (right). **I**. Analysis of the number of particles per mL of sample. **J.** Protein quantification in GDE in mg per mL of sample, obtained by BCA. **K.** Protein content per GDE, represented in mg of protein per particle. Results are expressed as mean $\pm$ SEM; n=17 for NCD-EV and n=25 for HD-EV. Unpaired t test with Welch's correction (F, I, J, K); ***$p<0.0005$, **$p<0.005$.

Figure 2 Study flow of proteomic approach for characterization of gut derived EV protein cargos

A. Strategy used for selection of proteins to be analyzed with confidence. **B**. Volcano plot representing the fold change and adj.p-value in the gut-derived EV (GDE) isolated from prediabetic mice and control groups. x-axis represents fold change, and y-axis adj.p-value. Dotted line sets the threshold of adj.p-value<0.05. In grey are the non-regulated proteins, in black the 558 regulated proteins.

Figure 3 Characterization of prediabetic gut derived EV by GO and KEGG enrichment analysis directly against mouse annotation, according to GO terms

A. Molecular function analysis of the GDE identified proteins. GO enrichment analysis according to Molecular Function. **B**. Analysis of the main biological regulated by the proteins identified in the GDE. GO enrichment analysis according to Biological Process. **C**. KEGG enrichment analysis for KEGG pathway. **D**. GO enrichment analysis according to Cellular Compartment. On the left panels, changes are displayed as the number of proteins with increased (red) or decreased (blue) levels (horizontal axis). On the right panels, the horizontal axis indicates the significance -log10 (p-value) of the functional association, which is dependent on the number of submitted proteins in the class, the number of proteins annotated in the class, the total number of submitted proteins and the total number of mouse proteins.

Figure 4 Cluster analysis of the top50 altered proteins of gut derived EV in
prediabetic mice
A. Volcano plot representing the fold change and adj.p-value of gut-derived EV (GDE) proteins isolated
from prediabetic mice and control groups. x-axis represents fold change, and y-axis adj.p-value. Dotted line
sets the threshold of adj.p-value<0.0001, highlighting the most altered proteins in GDE from HD comparing
with GDE from NCD. In blue are the most down regulated proteins and in red the most upregulated. **B**.
Heatmap representing mean-centered abundance of the most altered proteins decorated in the previous
volcano plot. **C**. Violin plot demonstrating the fold changes of the most altered proteins and correspondent
cellular functions. Cellular metabolic process (GO) represented in green.
638

Figure 5 Overview of prediabetes impact on gut derived EV proteome
The identified proteins with higher differential expression are represented in the image according to their
function and estimated subcellular localization. Red color for proteins upregulated and blue color for
proteins down regulated in small intestine derived EV of mice fed with HD compared with controls, mice
fed with NCD. ROS, reactive oxygen species; TCA, tricarboxylic acid cycle. Image done in BioRender.

Tables

Table 1 Biological function of the main up and down regulated proteins in HD-GDE
compared with NCD-GDE by proteomics

Supplementary material

Supplementary Figure 1 Study flow of proteomic approach for characterization of gut
derived EV protein cargos
A. Schematic representation of the detailed protocol used for isolation of GDE. Image done in
BioRender.
B. Schematic representation of the experimental process of analysis of gut derived EV (GDE).
Image done in BioRender. **C**. Bar plot indicating the number of proteins uniquely identified from
different species when searched against all proteins from all species in UniProt. **D**. Picture of LB-
agar (without antibiotics) petri dish after 24 after hours of inoculation with media where guts were
deposit overnight at 37ºC prior to EVs isolation.
658

Supplementary Figure 2 Protein-protein interaction network
The network was built using the STRING online software with a medium confidence level (0.4).
Each circle represents one protein. The line thickness indicates the strength of the evidence, with

thicker connections indicating higher confidence in the protein-protein interaction. Green circles

represent the proteins belonging to cellular metabolic process according to GO enrichment

analysis for biological processes.

Supplementary Figure 3 Individual protein alteration levels according to its biological

function

Data plots demonstrating the log fold changes of each protein in the image in accordance to its

related pathway.

Figure 1

Figure 2

A

B

Figure 3

A

Molecular Function

B

Biological Process

C

KEGG Pathways

D

Celullar Compartment

Figure 4

A

C

B

Figure 5

Table 1

Biological Function	Up	Down
Lipid Metabolism		
Mitochondrial ß-oxidation	ACADVL; ACADL	
Lipid Homeostasis	ACOT1; ACOT2; ACOT3; ACOT5; ACOT6	
Glucose Metabolism		
Glycolysis		ENO1; PGM5
Conversion of pyruvate to acetyl-CoA	PDHA1	
Amino Acid Metabolism		
C-terminal amino acid release	CPA1	
Protein degradation		VCP
Peptide metabolism		DNPEP
Purine Metabolism		XDH
TCA Cycle		IDH1
Oxidative phosphorylation	ETFA; ETFB	
Cell Integrity		ACTA1; ACTA2; ACTB; ACTC1; ACTG1; ACTG2; CORO1C; FLNA
Nucleosome	HIST1H4A; HIST3H2BB; HIST3H2BA; HIST2H2BE; HIST1H2BF	
Endocytosis		
Caveolar endocytosis	CAVIN1	
Clathrin-dependent endocytosis	AP2B1	
Focal adhesion		ILK; CDH17
Protease inhibitor	SERPINA1A; SERPINA1C	
Antioxidant Defense	PRDX6; GSTA4; GSTA1; GSTA2; ALDH2; ALDH1B1	TXN; SUOX
Proteasome		PSMA1; PSMA2; PSMA3; PSMA4; PSMA5; PSMA6; PSMA7; PSMB1; PSMB2; PSMB3; PSMB8; PSMB10