1 **Parallel Characterization of *cis*-Regulatory Elements for Multiple Genes Using**
2 **CRISPRpath**
3

4 **Authors:** Xingjie Ren[1], Mengchi Wang[2], Bingkun Li[1], Kirsty Jamieson[1], Lina Zheng[2], Ian R.
5 Jones[1], Bin Li[3], Maya Asami Takagi[1], Jerry Lee[1], Lenka Maliskova[1], Tsz Wai Tam[1], Miao Yu[3],
6 Rong Hu[3], Lindsay Lee[4], Armen Abnousi[4], Gang Li[5], Yun Li[6,7,8], Ming Hu[4], Bing Ren[3,9,10], Wei
7 Wang[2,11], Yin Shen[1,12,#]
8

Affiliation:
[1]Institute for Human Genetics, University of California, San Francisco, San Francisco, CA, USA.
[2]Bioinformatics and Systems Biology Graduate Program, University of California, San Diego,
La Jolla, CA, USA.
[3]Ludwig Institute for Cancer Research, La Jolla, California, USA.
[4]Department of Quantitative Health Sciences, Lerner Research Institute, Cleveland Clinic
Foundation, Cleveland, OH, USA.
[5]Department of Statistics and Operations Research, University of North Carolina, Chapel Hill,
NC, USA.
[6]Department of Biostatistics, University of North Carolina, Chapel Hill, NC, USA.
[7]Department of Genetics, University of North Carolina, Chapel Hill, NC, USA.
[8]Department of Computer Science, University of North Carolina, Chapel Hill, NC, USA.
[9]Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla,
CA, USA.
[10]Moores Cancer Center, University of California, San Diego, La Jolla, California, USA.
[11]Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA,
USA.
[12]Department of Neurology, University of California, San Francisco, San Francisco, CA, USA.
27
28 [#]Corresponding Author: yin.shen@ucsf.edu
29

Abstract

Current pooled CRISPR screens for *cis*-regulatory elements (CREs) can only accommodate one gene based on its expression level. Here, we describe CRISPRpath, a scalable screening strategy for parallelly characterizing CREs of genes linked to the same biological pathway and converging phenotypes. We demonstrate the ability of CRISPRpath for simultaneously identifying functional enhancers of six genes in the 6-thioguanine-induced DNA mismatch repair pathway using both CRISPR interference (CRISPRi) and CRISPR nuclease (CRISPRn) approaches. 60% of the identified enhancers are known promoters with distinct epigenomic features compared to other active promoters, including increased chromatin accessibility and interactivity. Furthermore, by imposing different levels of selection pressure, CRISPRpath can distinguish enhancers exerting strong impact on gene expression from those exerting weak impact. Our results offer a nuanced view of *cis*-regulation and demonstrate that CRISPRpath can be leveraged for understanding the complex gene regulatory program beyond transcriptional output at scale.

Main

Cis-regulatory elements (CREs) are key regulators for spatial-temporal control of gene expression. Mutations in CREs can contribute to complex diseases by modulating gene expression over long genomic distances[1-3]. Thus, functionally characterizing CREs can provide important insight into gene regulation mechanisms and enable us to better interpret non-coding genetic variants associated with diseases. Despite the fact that tremendous numbers of candidate CREs have been mapped by biochemical signature[4], our knowledge of whether, how, and how much these putative CREs are functional on gene expression remain scarce in the human genome. Pooled CRISPR screens have been developed for testing CREs in their native chromatin context by monitoring the transcriptional levels for the gene of interest[5-11]. Although results from these studies have made significant contributions to the annotation of functional DNA elements, challenges remain in pooled CRISPR screens of CREs. First, CRISPR screens for enhancers based on gene expression levels largely depend on generating reporter knock-in cell lines[7] or using FlowFISH signals[8]. These procedures, involving generation of reporter lines and selection of cells with positive hits by flow cytometry, are time-consuming and difficult to scale up to multiple genes in the same experiment. Second, the approaches of using gene

expression as the screening phenotype[9, 10] fail to connect the functions of DNA elements from

transcriptional regulation at the molecular level to interpretable cellular and physiological

functions. Third, in cases of CRE screens using phenotypes such as cell proliferation and

survival[11, 12], they fail to quantify the effect sizes of enhancers on transcriptional output.

To address these limitations, we developed CRISPRpath, a pooled CRISPR screening

approach to simultaneously characterizing CREs for multiple target genes involved in the same

biological pathway. CRISPRpath allows us to screen functional DNA elements based on

phenotypes associated with well-defined biological pathways. We demonstrate the capacity of

CRISPRpath by performing CRISPR interference (CRISPRi) and nuclease (CRISPRn) screens

for six genes in human induced pluripotent stem cells (iPSCs), and reveal different strengths of

enhancer functions by imposing varying levels of selection pressure on the cells.

Results

Leveraging CRISPRpath for parallel characterization of CREs for multiple genes in

iPSCs.

To characterize candidate *cis*-regulatory elements (CREs) for multiple genes within the same

pooled CRISPR screening, we designed and applied CRISPRpath to six genomic loci

containing six genes (*HPRT1, MSH2, MSH6, MLH1, PMS2, PCNA*) involved in the 6-

thioguanine (6TG)-induced mismatch repair (MMR) (**Fig. 1a**). The MMR pathway is highly

conserved and essential for the maintenance of genome stability[13]. The MMR pathway

recognizes DNA mismatches caused by 6TG treatment and induces cell apoptosis[14, 15]. On the

other hand, cells with a malfunctioning MMR pathway, due to aberrant expression levels of 6TG

metabolism genes or MMR genes, may survive during 6TG treatment. Employing the properties

of the MMR pathway, we used cell survival for selecting cells with the reduced expression of

MMR genes due to defects in enhancer activities (**Fig. 1b**). To design the screening library, we

first identified open chromatin regions by performing Assay for Transposase Accessible

Chromatin using sequencing (ATAC-Seq) in WTC11 iPSCs. We included all open chromatin

regions defined by ATAC-seq peaks located 1Mb upstream and 1Mb downstream of each of

the six genes (spanning a total of 10.6 Mb genomic regions) as candidate CREs for functional

characterization (**Supplementary Fig. 1a, b, Supplementary Table 1**). We then designed a

sgRNA library with 32,383 distal sgRNAs targeting 294 distal ATAC-seq peaks, 2,755 proximal

sgRNAs targeting 81 ATAC-seq peaks overlapped with transcription start site (TSS) and coding

regions of the six genes, and 625 non-targeting sgRNAs with genomic sequences in the same

genomic loci but are not followed by PAM sequences (**Supplementary Fig. 1c, Supplementary**

Table 2). In total, we included 35,763 sgRNAs in the library with an average of 110 sgRNA per

ATAC-seq peak (**Supplementary Fig. 1d, Supplementary Fig. 2a, b**). We generated a

lentiviral library expressing these sgRNAs and transduced this library into two engineered

WTC11 iPSC lines, one expressing doxycycline-inducible dCas9-KRAB (CRISPRi) and the

other doxycycline-inducible Cas9 (CRISPRn)[16], both at a multiplicity of infection (MOI) of 0.5

(**Fig. 1b**).

To carry out the screening, we pre-determined the minimal lethal concentration of 6TG at 80

ng/mL for CRISPRi and CRISPRn iPSC lines (see **Methods** for more details), and applied three

different 6TG concentrations (1X: 80 ng/mL, 2X: 160 ng/mL, 3X: 240 ng/mL) in both CRISPRi

and CRISPRn screens. We extracted and sequenced DNA samples from the survival cells

seven days after 6TG treatment to determine enriched sgRNAs by comparing the results to that

of the control cells taken after sgRNA library infection before the 6TG treatment (**Fig. 1b**). To

avoid confounding signals generated by off-target effects of low-quality sgRNAs[17], we only used

sgRNAs with high specificity (defined as specificity score > 0.2[18], and without any off-target sites

with sequence similarity of ≤2 mismatches) for data analysis. This led to the use of a total of

12,702 high-quality sgRNAs with an average of 38 sgRNAs per ATAC-seq peak for analysis

(**Supplementary Fig. 1e, f, Supplementary Fig. 2c, d**). We performed each screen in two

biological replicates with each pair of replicates exhibiting high reproducibility (**Supplementary**

Fig. 2e). We compared the abundance of each sgRNA between the 6TG treated population and

the control population using a negative binomial model, and computed the fold change and P

value to quantify the effect size and the significance of enrichment of each sgRNA. We used

the 5% percentile of the P values from non-targeting control sgRNAs as the empirical

significance threshold to achieve a false discovery rate of 5%. sgRNAs with P value less than

the empirical significance threshold and with fold change > 2 were defined as enriched.

(**Supplementary Fig. 3)**. As expected, sgRNA targeting TSS and coding region were identified

as positive hits from both CRISPRi and CRISPRn screens exhibiting greater fold change in

CRISPRn screens compared to the CRISPRi screens (**Fig. 1e, Supplementary Fig. 4a**). We also observed enrichment of sgRNA bias towards coding regions over TSS regions for CRISPRn screen (**Supplementary Fig. 4b**). These results are consistent with CRISPRi functioning best near TSS by inhibiting transcription, and CRISPRn can disrupt gene function by generating indels downstream of TSS[19, 20].

Further sgRNA fold-change ranking analysis revealed strong positive correlation between the screens with 2X and 3X 6TG treatment for both CRISPRi and CRISPRn screen (Spearman correlation, CRISPRi = 0.97, CRISPRn = 0.84) (**Fig. 1c**) with the correlations for proximal sgRNAs being higher than for distal sgRNAs (**Supplementary Fig. 4c**). On the contrary, results from the 1X screen correlated poorly with either 2X or 3X screens (**Fig. 1c**), suggesting more substantial selection pressure (2X and 3X) can reduce background noise in CRISPRpath screens. Thus, we used sgRNAs enriched from 2X and 3X screens data for identifying active enhancers in the following section (**Fig. 1d**).

CRISPRi is more efficient than CRISPRn in pooled CRISPR screens of CREs.

Performing CRISPRpath with CRISPRi and CRISPRn in the same genetic background with an identical sgRNA library offers a unique opportunity for comparing the efficacies of CRISPRi and CRISPRn in pooled CRISPR screens of CREs. We noticed that CRISPRn screens recovered fewer enriched distal sgRNAs than CRISPRi screens (**Fig. 1f**). This is possibly due to the fact that CRISPRi-mediated heterochromatin formation can more effectively perturb CREs compared to CRISPRn-mediated genetics perturbations. We then called an candidate element as an enhancer if there are at least 3 enriched sgRNAs in that CRE. Based on this criterion, we identified 62 and 33 enhancers from the 2X and 3X CRISPRi screen, respectively, and 19 enhancers from the 2X CRIPSRn screen. (**Fig. 1g, Supplementary Table 3**). However, no enhancer was identified from the 3X CRISPRn screen, indicating either the CRISPRn induced mutations did not lead to any strong effect on gene expression to make the cells survive the 3X 6TG treatment or there are insufficient numbers of sgRNAs exhibiting deleterious effects on the tested DNA elements to satisfy our criterion of calling functional enhancers. In total, 66 unique enhancers were identified for the six target genes with CRISPRpath under different 6TG treatments (**Fig. 1g**). Together, we demonstrate CRISPRpath can simultaneously identify

enhancers for multiple target genes with CRISPRi outperforming CRISPRn. For the following

analysis, we focused on the 63 enhancers identified from the 2X and 3X CRISPRi screens (**Fig.**

1g).

Genomic feature of CRISPRpath identified enhancers.

To determine the genomic feature of the enhancers, we plotted all the tested elements by their

genomic locations and enrichment scores (average of $\log_2$(fold change) of enriched sgRNAs of

each element) (**Fig. 2a**). Not surprisingly, our data suggest that each gene can be regulated by

multiple enhancers with the identified functional enhancers having no position bias relative to

the TSS. The average distance between an enhancer and its paired TSS is about 530 Kb (**Fig.**

2b) with an average of 10 interval genes between an identified enhancer and its target gene

pairs (**Fig. 2c**). Interestingly, we observed a weak negative correlation between the enhancer

enrichment score and the distance between an enhancer and its paired TSS (**Fig. 2d**, Pearson

correlation, ρ = -0.36, *P* = 0.01), suggesting enhancers near to TSS tend to have higher

regulatory activity compared to enhancers further away from their target genes. It is worth

noting, the relative positions for the enriched sgRNAs exhibited no preference relative to ATAC-

seq peaks (**Fig. 2e, Supplementary Fig. 5a**) and no preference for the strand on which the

sgRNAs were designed (**Supplementary Fig. 5b**), consistent with our knowledge that CRISPRi

mediated heterochromatin spreads over hundreds of base pairs in distance[21].

Previous studies have revealed that promoters can function as enhancers[7, 22]. Indeed, 60% (38

out of 63) of the functional enhancers identified in CRISPRi screens overlapped with annotated

promoters, providing an excellent opportunity to further explore the genomic features of these

enhancer-like promoters. To validate whether these promoters function as bona fide enhancers,

we targeted three enhancer-like promoters with CRISPRi. We confirmed significant

downregulation of their target genes including *MSH6*, *MSH2*, and *PCNA* (**Fig. 3a-c**). In contrast,

shRNAs against the transcripts from these promoters (*SOC5*, *FOXN2*, and *TMEM230*) only led

to a significant downregulation of its own transcripts and did not affect their target gene

expression (**Fig. 3a-c**). These results confirm that these promoter sequences identified by

CRISPRpath can function as enhancers. Although it has been shown that enhancer-like

protomors are enriched with active chromatin marks and physically close to target genes[7], it is

not clear whether enhancer-like promoters have unqiue genome features that can differentiate
them from other regular active promoters. To this end, we compared chromatin accessibility,
occupancy of histone 3 lysine 4 trimethylation (H3K4me3), histone 3 lysine 27 acetylation
(H3K27ac) and CTCF, transcription, and chromatin interactivity levels between enhancer-like
promoters and all other active promoters that did not show enhancer activity in our CRISPRi
screens. We show that enhancer-like promoters exhibit higher chromatin accessibility, higher
level of transcription, stronger H3K4me3 and H3K27ac signals than those at other active
promoters (**Fig. 3d**). On the other hand, we did not observe a significant difference for CTCF
binding signals between enhancer-like promoters and control promoters (**Fig. 3d**). Furthermore,
by evaluating chromatin interaction data using H3K4me3 Proximity Ligation-Assisted ChIP-seq
(PLAC-seq), we show enhancer-like promoters have significantly more and stronger
interactions compared to control promoters (**Fig. 3e**).

CRISPRpath is capable of distinguishing enhancers with distinct effect sizes.
Gene expression often is a result of combinatorial regulatory effects from multiple *cis*-regulatory
elements[11, 23]. Understanding how individual enhancers contribute to gene expression in a
quantitative manner is an important first step in dissecting how enhancers orchestrate precise
transcriptional control. We seek a new strategy to differentiate enhancers based on their effect
sizes on gene expression using CRISPRpath. We hypothesized that cells with drastic down-
regulation of MMR genes have a fitness advantage under higher 6TG concentration than cells
with modest down-regulation of MMR genes. Consistent with this hypothesis, proximal sgRNAs
exhibit larger fold changes than distal sgRNAs (**Fig. 1e**) because perturbing proximal regions
has more profound effects on gene down-regulation than perturbing distal regulatory regions.
Based on these observations, we hypothesize that enhancers identified under different
selection pressure represent distinct regulatory strengths on transcriptional activation. We
noticed that enhancers identified under strong selection pressure (3X) have higher enrichment
scores compared to enhancers uniquely identified under weak selection pressure (2X), with the
TSS regions manifesting the highest enrichment scores (**Fig. 4a**). Thus, enhancers identified in
the 3X screen are strong enhancers (n=33), while enhancers uniquely identified in the 2X screen
are weak enhancers (n=30) (**Fig. 1g**). To confirm the quantitative effect of enhancers on target
gene expression, we tested 11 strong and 10 weak enhancers using CRISPRi followed by RT-

qPCR measurement of the corresponding target gene expression (**Fig. 4b, Supplementary**
Fig. 6a). We show perturbations of strong enhancers led to significantly more down-regulation
of target gene expression (mean down-regulation of target gene by 21%) than perturbations of
weak enhancers (mean down-regulation of target gene by 6%), with the perturbations of TSS
regions achieving the strongest down-regulation of target genes, by an average of 68%
reduction in gene expression (**Fig. 4b, Supplementary Fig. 6a, b**). These quantitative effects
on target gene expression are consistent with the enrichment scores from our CRISPRpath
screens (**Supplementary Fig. 6c**) and demonstrate the capacity of distinguishing enhancers
with different effect sizes by imposing different levels of selection pressures.

We further explored chromatin features of strong and weak enhancers by analyzing chromatin
accessibility, H3K4me3, H3K27ac, and CTCF binding signals in these regions. At individual
chromatin mark level, while CRISPRpath-identified enhancers were more accessible, and
enriched with active chromatin marks, such as H3K4me3 and H3K27ac, and CTCF binding
compared to negative elements or random elements (**Fig. 4c**), we did not observe significant
differences between strong and weak enhancers in the chromatin features we individually
examined. However, strong enhancers tend to have more active chromatin signatures than
weak enhancers (**Fig. 4d**), suggesting combined signatures of active chromatin can be a better
indicator of enhancer strength. Strong enhancers tend to have higher distance normalized
PLAC-seq contact frequencies with their target promoters than weak enhancers, though not
statistically significant, possibly due to the small sample size in this study (**Fig. 4e**). We obtained
similar results by expanding this analysis for characterized enhancers in K562 cells and mouse
embryonic stem cells[6, 8] (**Supplementary Fig. 7**), which reinforces the idea that enhancers with
larger effects on gene expression tend to have higher chromatin interactions with their cognate
promoters. To explore the possible mechanisms that drive enhancer activities in a quantitative
manner, we evaluated potential transcription factors (TFs) binding motifs in strong and weak
enhancer sequences. Both strong and weak enhancers are enriched with CTCF binding motif
(**Fig. 4f**). Indeed, most of strong and weak enhancers are bound by CTCF (**Fig. 4d**), consistent
with the notion that CTCF-mediated chromatin loops are essential for gene activation[24].
Furthermore, strong enhancers and weak enhancers have differential enrichment with TFs
binding motifs. For example, the binding motifs for SP/KLF family[25] and E2F family[26, 27] appear

more frequently in strong enhancers compared to weak enhancers, suggesting these strong

enhancers could be major docking sites for master regulators in iPSCs (**Fig. 4f**).

Discussion

CRISPR-mediated high throughput screening using bulk cells allows for the functional

characterization of regulatory elements in their native genomic context. However, current

approaches are limited to validating a small number of regulatory elements for a single gene[5,]

[7, 9, 12, 28, 29]. To overcome this bottleneck, we developed CRISPRpath, a strategy for functional

characterization of enhancers for multiple genes simultaneously by leveraging the genes

involved in the same biological pathway so that the effects can be measured via a define

phenotype. For example, alpha-toxin resistance phenotype can be used to identify CREs for 17

genes in glycosylphosphatidylinositol (GPI)-anchor synthesis pathway[30]. CRISPRpath can also

be leveraged to identify CREs for protein folding regulators that contribute to the endoplasmic

reticulum stress-response pathway[31] using UPRE reporter in mammalian cells. Compared to

the existing pooled CRISPR screens of CREs[5, 7, 8, 10-12, 28, 29, 32], CRISPRpath is scalable with

additional benefits of connecting DNA elements to cellular function, beyond the most standard

molecular phenotype of gene expression.

Promoters can function as enhancers more widespread than expected, with more than half of

the enhancers identified for MMR genes in our study being previously annotated promoters.

This is consistent with previous reports that enhancer-like promoters are more prevalent for

ubiquitously expressed genes[33]. Enhancer-like promoters are more accessible compared to

other promoters, possibly because these regions are required to be more open to accommodate

additional transcriptional machinery such as TF for activating target gene expression besides

their own transcription[34]. Enhancer-like promoters also exhibit significantly higher levels of

chromatin interactions with distal regions compared to other active promoters. This observation

can be explained by the fact that enhancer-like promoters will not only form chromatin loops

with their distal target genes, but also with CREs for controlling the expression of their own

genes.

Genomic studies of chromatin marks have revealed hundreds of thousands candidate CREs in

the human genome but with very little quantitative information regarding how CREs contribute

to gene regulation[35, 36]. Using CRISPRpath, we can systematically classify enhancers based on

their effect sizes on transcription. Identifying and charaterizing the effect size for each individual

enhancer is the critical first step to future studies of their combinatory effects on target gene

expression. Interestingly, strong and weak enhancers can not be distinguished by individual

epigenetic marks we examined. One possible explanation for this observation is that chromatin

features only mark enhancer's identity but do not quantify enhancer activity. On the other hand,

the strong and weak enhancers we identified may regulate other genes differently from

regulating the MMR gene. Interestingly, strong enhancers tend to harbor more than one active

chromatin signature, which indicates that enhancer activities are regulated by multiple

epigenetic factors, for example, TF mediated transcriptional regulation. Differential TFs binding

motifs observed within strong and weak enhancers suggest that enhancer strength is modulated

by TF binding. Future studies that further integrating TF binding datasets with functional data of

enhancers will shed light on the molecular mechanisms that drives enhancers' effect sizes on

gene regulation.

Acknowledgements

This work was supported by the National Institutes of Health grants UM1HG009402 (to Y.S.).

Data availability

The CRISPRpath screen datasets used in this study are available at the ENCODE portal

(www.encodeproject.org, accession number: ENCSR617AZY (sgRNA plasmid library),

ENCSR427OPP (CRISPRi control), ENCSR900AXT, ENCSR254SJU (CRISPRi 2X),

ENCSR793DSE (CRISPRi 3X), ENCSR250ZWC (CRISPRn control), ENCSR117YGQ

(CRISPRn 1X), ENCSR071ZGB (CRISPRn 2X), ENCSR482PHH (CRISPRn 3X)). WTC11

iPSCs H3K4me3 PLAC-seq datasets are available at the 4DN data portal

(data.4dnucleome.org, accession number: 4DNESIZ5TTHO and 4DNESDRL4ZKM). ATAC-

seq, ChIP-seq, CUT&Tag, and RNA-seq datasets in WTC11 iPSCs are available at the Gene

Expression Omnibus under the accession number GSE166839.

Data can be visualized on the WashU Epigenome Browser using the session at the following link (https://epigenomegateway.wustl.edu/browser/?genome=hg38&sessionFile=https://shen-xren.s3-us-west-1.amazonaws.com/CRISPRpath/eg-session-QRXJ0218-4d710b60-6ea7-11eb-8d8d-03c7189570c0.json). Tracks include ATAC-seq, H3K27ac, H3K4me3, and CTCF signals, and the identified enhancers from CRISPRi 2X and 3X screens. RefGene 38 genes are also displayed.

The plasmids generated in this study are available from Addgene (#167186, #167188, # 167189).

Code availability

The computer code used for analyzing CRISPRpath datasets is available at https://github.com/MichaelMW/crispy. A copy of the custom code used for data analysis and figure generation in this study is available upon request.

Competing interests

B.R. is co-founder and shareholder of Arima Genomics and Epigenome Technologies. The other authors declare that they have no competing interests.

Author contributions

X.R. and Y.S. designed the study. X.R. and B.L. designed the sgRNA library under the supervision of Y.S. and B.R. X.R., K.J., I.R.J., M.A.T., J.L., L.M., and T.W.T. performed the experiments. M.W. designed the CRISPY under the supervision of W.W. X.R., B.L., L.Z., G.L., and Y.L. performed data analysis. M.Y. and R.H. constructed the H3K4me3 PLAC-seq libraries. L.L., A.A. and M.H. analyzed PLAC-seq and HiChIP data. X.R. and Y.S. prepared the manuscript with input from all other authors.

Figure 1. CRISPRpath for identifying enhancers of multiple genes.

(**a**) Six genes (*HPRT1*, *MSH2*, *MSH6*, *MLH1*, *PMS2* and *PCNA*) in the 6TG-induced mismatch repair process were used for CRISPRpath screen in this study. (**b**) Schematic of the CRISPRpath screening strategy with 6TG treatment in iPSCs. Cell survival was used as readout for the screen. (**c**) Spearman correlation analysis of sgRNA ranking based on fold change for CRISPRpath screens with different 6TG concentrations (1X, 2X and 3X). (**d**) Venn diagram shows the overlapping enriched sgRNAs identified from the screens with 2X and 3X 6TG treatments. (**e**) Box plots show the fold change of the enriched distal and proximal sgRNAs from 2X, 3X CRISPRi and CRISPRn screens. Star indicates no enriched distal sgRNA was identified from 3X CRISPRn screen. Boxplots indicate the median, interquartile range (IQR), Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. (**f**) Bar plot shows the number of enriched distal and proximal sgRNAs from 2X, 3X CRISPRi and CRISPRn screens. Star indicates no enriched distal sgRNA was identified from 3X CRISPRn screen. (**g**) Venn diagram shows the identified enhancers from each CRISPRpath screen.

Figure 2. Genomic features of identified enhancers from CRISPRpath using CRISPRi.

(**a**) Genomic locations of identified enhancers relative to TSS. Circles indicate enhancers identified from the CRISPRi 3X screen (red), enhancers uniquely identified from the CRISPRi 2X screen (blue), tested CREs that are not identified as enhancers (grey). Purple lines label the location of each target gene. (**b**) Histogram shows the distance distribution between identified enhancers and their paired TSS. (**c**) Histogram shows the number of interval genes between enhancers and their target gene TSS. Mean is indicated with an orange dashed line and only the enhancers for *MLH1*, *PMS2*, *PCNA*, *HPRT1* are included in **b** and **c**. (**d**) A weak negative correlation is observed between enrichment score and genomic distance between enhancers and their target genes (Pearson correlation, r = -0.36, *P* = 0.01). Black circles indicate promoters. The red and blue circles are enhancers showing in **a**. (**e**) Density plot shows no significant difference (two-tailed two-sample Kolmogorov–Smirnov test) for the distribution of all distal sgRNAs (gray), enriched distal sgRNAs from 2X (blue) and 3X screen (red) CRISPRi screens.

Figure 3. Enhancer-like promoters act as functional enhancers.

(**a**, **b**, **c**) Three examples of promoters function as enhancers. CRISPRi silencing of the
promoter region of *SOCS5*, *FOXN2* and *TMEM230* results in significant downregulation of
MSH6, *MSH2* and *PCNA*, respectively. shRNA knockdown of *SOCS5, FOXN2* and *TMEM230*
can only downregulate *SOCS5, FOXN2* and *TMEM230* expression. Three independent
replicates per condition and two independent sgRNAs or shRNAs per replicate were used for
each experiment. *P* values are from two-tailed two-sample t-test. (**d**) Average signal enrichment
of ATAC-seq, gene transcription, H3K4me3, H3K27ac and CTCF binding for enhancer-like
promoters (n = 38) and control promoters (n = 47). *P* values are from Wilcoxon test. Boxplots
indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. (**e**) Number of H3K4me3
mediated chromatin interactions and cumulative interaction score for enhancer-like promoters
(n = 31) and control promoters (n = 43). Boxplots indicate median, IQR, Q1 − 1.5 × IQR and Q3
+ 1.5 × IQR. *P* values are calculated from Wilcoxon test.

Figure 4. CRISPRpath can distinguish weak and strong enhancers by imposing different
selection pressures.

(**a**) Box plots show the enrichment score of the tested elements. TSS regions (black circles)
show highest enrichment scores. Enhancers uniquely identified from the lower selection
pressure (CRISPRi 2X, blue circles) exhibit lower enrichment scores compared to the
enhancers identified from the higher selection pressure (CRISPRi 3X, red circles). *P* values are
from Wilcoxon test. (**b**) Box plots show the CRISPRi perturbation at enhancers induced various
degrees of transcriptional repression of target genes measured with RT-qPCR. Each dot
represents the average value from three biological replicates. CRISPRi targeting TSS regions
(dark gray) achieved the highest transcriptional repression. CRISPRi targeting strong
enhancers (pink) leads a more substantial transcription silencing on target gene compared to
CRISPRi targeting weak enhancers (cyan). *P* values are from Wilcoxon test. (**c**) Enrichment
analysis of ATAC-seq, H3K27ac, H3K4me3, CTCF binding signals for strong (n = 33) and weak
(n = 30) enhancers. Boxplots indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. *P*
values for the difference between strong and weak enhancers are from Wilcoxon test; see
Supplementary Table 7 for *P* values of all pairwise comparisons. (**d**) Intersection of genomic
features for weak enhancers (blue bar) and strong enhancers (red bar). (**e**) Distance normalized
H3K4me3 PLAC-seq contact frequency for strong (n = 23) and weak (n = 21) enhancers. Only

the enhancers for *MLH1*, *PMS2*, *PCNA*, *HPRT1* are included (see Methods). Boxplots indicate

the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. *P* value is from Wilcoxon test. (**f**) Heatmap

shows the frequency of transcription factor motifs found in strong and weak enhancers.

Supplementary Figure 1. Features of sgRNA library for CRISPRpath screen.

(**a**) Bar graph shows the number of distal ATAC-seq peaks used as candidate CREs for six

target genes. (**b**) Histogram shows size distribution of distal ATAC-seq peaks. The average size

is 488 bp (blue dash line). (**c**, **e**) The composition of the sgRNA library. In total, 35,763 sgRNAs

were included in the library (**c**), and 12,702 sgRNAs are high quality sgRNAs (**e**). (**d**, **f**)

Distribution of the number of sgRNAs per distal ATAC-seq peak. Average numbers of sgRNA

per ATAC-seq peak are indicated with blue dash lines.

Supplementary Figure 2. Quality of the sgRNA library and CRISPRpath screen libraries.

(**a**) Distribution of sgRNA oligo read counts in the sgRNA library. (**b**) Cumulative frequency of

sgRNAs in the sgRNA library. (**c**) Distribution of high quality sgRNAs read counts in the sgRNA

library. (**d**) Cumulative frequency of high quality sgRNAs in the sgRNA library. The constructed

sgRNA plasmid library recorvered all the designed sgRNAs with the copy number difference

less than five fold for at least 97% designed sgRNAs. (**e**) PCA analysis shows the high

reproducibility of the CRISPRpath screen libraries between biological replicates.

Supplementary Figure 3. *P* value cutoff used for identifying enriched sgRNAs from each

screens.

(**a-f**) Distribution of *P* value for tested distal, proximal and non-targeting control sgRNA groups.

Orange dash lines indicate 5% percentile of the *P* values from non-targeting control sgRNAs to

achieve a false discovery rate of 5%.

Supplementary Figure 4. Enriched proximal sgRNAs and sgRNA ranking analysis.

(**a**) Number and fold change of the enriched proximal sgRNAs for the six target genes from

CRISPRi and CRISPRn screens. The color indicates fold changes, and the size of circle

indicates the number of enriched sgRNAs. (**b**) Enrichment analysis shows the enriched proximal

sgRNAs bias towards to the TSS region for CRISPRi screens and the protein coding region

(CDS) for CRISPRn screens. Color represents the number of enriched sgRNAs. (**c**) Spearman

correlation analysis of the distal and proximal sgRNAs ranking shows proximal sgRNAs

exhibiting higher correlation between each screen compared to distal sgRNAs.

Supplementary Figure 5. Enriched sgRNAs identified from CRISPRi screens exhibit no

position and strand preference.

(**a**) Enriched sgRNAs from CRISPRi 2X (red dots, n=448) and CRISPRi 3X (red dots, n=260)

screens showed similar distributions across candidate CREs. (**b**) Odds ratio analysis of the fold

change of enriched sgRNAs shows enriched sgRNAs have no strand preference. Enhancers

with enriched sgRNAs only targeting one strand were exclude for the analysis. Odds ratio was

calculated for each element with the equation of ave($\log_2$(fold change of sgRNA targeting plus

strand)) / ave($\log_2$(fold change of sgRNA targeting minus strand)). Violin plots show the

distributions of odds ratio values within each screen, and boxplots indicate the median, IQR,

$Q1 - 1.5 \times IQR$ and $Q3 + 1.5 \times IQR$.

Supplementary Figure 6. Validation of CIRSPRpath identified enhancers.

(**a**) Validation of the strong (black) and weak (grey) enhancers with CRISPRi followed by RT-

qPCR. Three independent replicates per condition. The significance was calculated with two-

tailed two-sample *t*-test. Data are mean and s.d. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. (**b**)

CRISPRi-mediated transcriptional repression of six target genes by targeting TSS of each gene.

Three independent replicates per condition. The significance was calculated with two-tailed two-

sample *t*-test. Data are mean and s.d. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. (**c**) Pearson

correlation analysis reveals element enrichment score from CRISPRpath screens correlates

with element effect size on transcription from CRISPRi (Pearson correlation, PCC = -0.68, $P = $

3.2×10^{-5}).

Supplementary Figure 7. Chromatin contract frequency analysis for the enhancers in

K562 cells and mESCs.

(**a**) Distance normalized H3K27ac HiChIP contact frequency for strong (n = 34) and weak (n =

82) enhancers identified with crisprQTL mapping in K562 cells. (**b**) Distance normalized

H3K27ac HiChIP contact frequency for strong (n = 2) and weak (n = 20) enhancers identified

with CRISPRi-FlowFISH screen in K562 cells. (**c**) Distance normalized H3K4me3 PLAC-seq contact frequency for strong (n = 10) and weak (n = 3) enhancers identified in mouse embryonic stem cells. Boxplots indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. *P* values are from Wilcoxon test.

Supplementary Table 1. ATAC-seq peak regions used to design the sgRNA library.

Supplementary Table 2. List of sgRNA sequences used for CRISPRpath screen.

Supplementary Table 3. Enhancers identified from CRISPRn 2X, CRISPRi 2X and CRISPRi 3X screens.

Supplementary Table 4. List of primers used for RT-qPCR.

Supplementary Table 5. List of shRNA sequences used for RNA interference experiments.

Supplementary Table 6. List of sgRNA sequences used for CRISPRi-mediated enhancer validation experiments.

Supplementary Table 7. Pairwise comparisons of data in Figure 4c.

 Methods

Cell culture

Doxycycline inducible CRISPRi and CRISPRn WTC11 iPSC lines were purchased from

Gladstone Stem Cell Core. Both CRISPRi and CRISPRn WTC11 iPS cells were cultured on

Matrigel-coated (Corning, 354277) plates with Essential 8™ Medium (Life Technologies,

A1517001). iPSCs were passaged using Accutase (STEMCELL Technologies, 07922) and 10

µM ROCK inhibitor Y-27632 (STEMCELL Technologies, 72302). HEK293T cells were cultured

in Dulbecco's modified Eagle's medium (Gibco, 11995065) with 10% fetal bovine serum

(CPSSerum, FBS-500). HEK293T cells were passaged with Trypsin-EDTA (Gibco, 25200072).

All the cells were grown with 5% CO_2 at 37°C and verified mycoplasma free using the MycoAlert

Mycoplasma Detection Kit (Lonza, LT07-218).

sgRNA library design

CRISPRpath sgRNA library was designed to screen cis-regulatory elements for *HPRT1*, *MSH2*,

MSH6, *MLH1*, *PMS2* and *PCNA*. ATAC-seq peaks within the region of 1 Mb upstream and 1

Mb downstream of each target gene including TSS and coding regions were selected as

targeting regions for the sgRNA library design (**Supplementary Table 1**). We generated a

genome-wide sgRNA database containing all the available unique sgRNAs, each followed by a

'NGG' PAM sequence. All the designed unique sgRNAs in the target regions were added in the

sgRNA library, excluding sgRNAs containing AATAAA, AAAAA, TTTTT or TTTTTT sequences.

Unique 20-bp sequences in the target regions that were not followed by the 'NGG' or 'NAG'

PAM sequences were taken as non-targeting control sgRNAs, excluding non-targeting sgRNAs

containing TTT, TTNTT, AATAAA, AAAAA, TTTTT or TTTTTT sequences. Then, a guanine

nucleotide was added to all the sgRNAs if the sequence did not start with G to increase

efficiency of transcription from U6 promoter. Final sgRNA oligos adhered to the following

template: 5'-ATATCTTGTGGAAAGGACGAAACACC-[20- or 21-bp sgRNA sequence]-

GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGC-3'. In total, 35,763 sgRNAs were

included in the library (**Supplementary Fig. 1 and Supplementary Table 2**). We retrieved

specificity score and off-target site for each sgRNA from GuideScan[18] (www.guidescan.com)

and assigned the specificity score of sgRNAs not existed in the GuideScan database to 0. The

high-quality sgRNAs were filtered with specificity score >0.2 and without perfectly matched or

1-2 mismatches off-target sites.

Oligo synthesis and library cloning

sgRNA library oligos were synthesized by TWIST BIOSCIENCE and amplified with the forward

primer 5'- TCGATTTCTTGGCTTTATATATCTTGTGGAAAGGACGAAACAC-3' and the

reverse primer

5'-AACGGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC-3'. We replaced the

Cas9 sequence in lentiCRISPR v2 plasmid (Addgene, 52961) with blasticidin S deaminase

sequence to construct the lentiCRISPR-v2-Blast-Puro plasmid (Addgene, 167186). The PCR

products were purified via gel excision and column purification (Promega, A9282), and then

inserted into the BsmBI-digested lentiCRISPR-v2-Blast-Puro vector by Gibson assembly (New

England Biolabs, E2621L). The assembled products were transformed into NEB 5-alpha

electrocompetent *E. coli* cells (New England Biolabs, C2989K) by electroporation. About 40

million independent bacterial colonies were cultured, and sgRNA library plasmids were

extracted with the Qiagen EndoFree Plasmid Mega Kit (Qiagen, 12381). The recovery rate and

distribution of the sgRNA library were checked with next generation sequencing

(**Supplementary Fig. 2a-d**).

Lentivirus production and titration

To make the lentiviral library, 5 μg of sgRNA plasmid library was co-transfected with 3 μg of

psPAX (Addgene, 12260) and 1 μg of pMD2.G (Addgene, 12259) lentivirus packaging plasmids

into 8 million HEK293T cells in a 10-cm dish with PolyJet (SignaGen Laboratories, SL100688).

For each individual sgRNA, 3.75 μg of sgRNA plasmid was co-transfected with 2.25 μg of

psPAX (Addgene, 12260) and 0.75 μg of pMD2.G (Addgene, 12259) plasmids into 4 million

HEK293T cells in a T25 flask with PolyJet (SignaGen Laboratories, SL100688). Media was

replaced 12 h after transfection, and harvested every 24 h for a total of three harvests.

Harvested media containing the desired virus were filtered through Millex-HV 0.45-μm PVDF

filters (Millipore, SLHV033RS) and further concentrated with 100,000 NMWL Amicon Ultra-15

centrifugal filter units (Amicon, UFC910008).

The titer of lentivirus was determined by transducing 500,000 cells with varying amount (0, 0.5,
1.0, 2.0, 4.0 and 8.0 µL) of concentrated virus and polybrene (Millipore, TR-1003-G, 8 µg/mL,).
Viral transduction was performed by centrifuging the lentivirus and cell combination at 1000
RCF for 90 min at 37°C. 3 to 4 h later, virus containing media was replaced with fresh media.
24 h after the transduction, transduced cells were dissociated with Accutase, and seeded as
duplicates. One replicate was treated with blasticidin (Gibco, A1113903, 4 µg/mL), and the other
replicate was not treated with blasticidin. Four days later, the blasticidin resistant cells and
control cells were counted to calculate the ratio of infected cells and the viral titer.

Determining 6TG concentration via killing curve titration
Both CRISPRi and CRISPRn WTC11 iPSCs were used to determine the minimal lethal
concentration of 6TG. Cells were seeded in 24 well plates. When the cells reached around 50%
confluence (Day 0), they were treated with 6TG concentrations of 0 (control), 20, 40, 60, 80,
100, 120, 140, and 160 ng/mL. Two wells were allocated for each condition. The cells were
examined daily and cultured for 7 days. The media was replaced daily with the specified 6TG
concentration. After 3 days, wells with 6TG concentration greater than or equal to 100 ng/mL
had no surviving cells. On Day 4 of treatment, the wells with 80 ng/mL 6TG treatment had no
surviving cells. On the last day of treatment, the wells with 40 and 60 ng/mL treatments had
very few surviving cells, while the 20 ng/mL treatment had many surviving cells. Based on these
results, we set 80 ng/mL as the minimal lethal concentration for 6TG.

CRISPRpath screening and sequencing library preparation
CRISPRpath screens were carried out with 72 million doxycycline inducible CRISPRi or
CRISPRn iPSCs in biological replicate. The cells for lentiviral transduction were seeded into 6
well plates with 1 million cells per well, and the lentiviral library (MOI = 0.5) was transduced into
the iPSCs with 8 µg/mL Polybrene (Millipore, TR-1003-G) and spun at 1000 RCF at 37°C for
90 min. The transduced cells were treated with doxycycline (Sigma, D9891, 2 µM) and
blasticidin (Gibco, A1113903, 4 µg/mL) for 4 days. After this doxycycline and blasticidin
treatment, 10 million cells were reserved as a control population, and 100 million cells were
used for CRISPRpath screen with doxycycline and 6TG (Sigma, A4660) treatment for 7 days.
Finally, survival cells were collected from the 6TG treated population.

The genomic DNA was extracted from each sample via cell lysis and digestion (100 mM Tris-

HCl pH8.5, 5 mM EDTA, 200 mM NaCl, 0.2% SDS, 100 µg/mL proteinase k), phenol:chloroform

(Thermo Scientific, 17908) extraction and isopropanol (Fisher Scientific, BP2618500)

precipitation. To amplify the sgRNA sequences from each sample, thirty-two 50 µl PCR

reactions were performed using 500 ng genomic DNA for each reaction and NEBNext® High-

Fidelity 2X PCR Master Mix (New England Biolabs, M0541S). The purified libraries were

sequenced on the NovaSeq 6000 with 150-bp paired-end sequencing. The detail protocol is

available on ENCODE portal (https://www.encodeproject.org/documents/2e6451a9-3b98-

4d95-922e-a3d8d2100ddf/).

CRISPRpath data analysis

The sequence files were down-sampled to the same amount of total reads, and then mapped

to the sgRNA library with the requirement of exact match of designed sgRNA sequences in the

following patten 5'-CCG-[N19 or N20]-GTT-3'. Only the highly specific sgRNAs (specificity score

>0.2, without perfectly matched or 1-2 mismatches off-target sites) were used for downstream

data analysis. The sgRNA enrichment for each screen was calculated by comparing 6TG

treated samples with the associated control samples with edgeR and TMM normalization. We

first used edgeR[37] to calculate P value based on negative binomial model for both targeting

sgRNAs and non-targeting control sgRNAs. To achieve empirical false discovery rate less than

5%, we then selected a P value cutoff corresponding to the 5% percentile of P values from non-

targeting control sgRNAs. Finally, we defined enriched sgRNAs with P value less than the

selected P value cutoff, and fold change >2. The ATAC-seq peaks were identified as functional

enhancers for the six MMR genes by having at least 3 significant enriched sgRNAs. Analysis

scripts are available at https://github.com/MichaelMW/crispy.

Analysis of genomic feature and chromatin signature of identified enhancers

Genomic distances between enhancer and TSS pairs were calculated based on the distance

from the center of enhancers to the transcription start sites of the target genes. The number of

interval genes is the number of all the RefSeq annotated genes between each enhancer and

paired target gene. The signal of chromatin signatures, including ATAC-seq, H3K27ac,

H3K4me3, CTCF binding and RNA-seq, were calculated by deeptools (v3.4.3)[38]. The enhancer-
like promoters are the enhancers overlap with the region 500 bp upstream and downstream of
a RefSeq annotated TSS.

Validation of identified enhancers using CRISPRi

We cloned lentiCRISPR-v2-HygR-EGFP (Addgene, 167188) and lentiCRISPR-v2-HygR-
mCherry (Addgene, 167189) vectors by replacing the Cas9 and puromycin N-acetyltransferase
sequences in lentiCRISPR v2 plasmid (Addgene, 52961) with hygromycin B
phosphotransferase and EGFP or mCherry sequences. To validate the identified enhancers,
individual sgRNAs targeting identified enhancers were cloned into the lentiCRISPR-v2-HygR-
GFP or lentiCRISPR-v2-HygR-mCherry vector. The doxycycline inducible CRISPRi WTC11
iPSCs were infected with the lentivirus expressing sgRNAs for three replicates per sgRNA. The
sgRNA infected cells were grown with hygromycin (Gibco, 10687010, 150 µg/mL) and
doxycycline (Sigma, D9891, 2 µM) containing media. Seven days later, the cells were collected
and total RNA was extracted from the cells using the Qiagen RNeasy® Plus Kit (Qiagen,
74134). One µg of RNA was then used to synthesize cDNA using the Bio-RAD iScript cDNA
Synthesis Kit (Bio-RAD, 1708840). qPCR reactions for targeted genes were performed with the
Luminaris HiGreen qPCR Master Mix (Thermo Scientific, K0993) on the Roche LightCycler 96
System. The qPCR primers are listed in **Supplementary Table 4** and the sgRNA sequences
are listed in **Supplementary Table 6**. For each tested element in **Fig. 3a-c** and **Supplementary**
Fig. 6b, we performed CRISPRi experiments with two independent sgRNAs and used the
results from the sgRNA with stronger transcriptional repression in **Fig. 4b**.

shRNA mediated RNA interference

shRNAs were designed by using DSIR tool (http://biodev.extra.cea.fr/DSIR/DSIR.html)
targeting *SOCS5*, *FOXN2* and *TMEM230*. The sequences of shRNAs are listed in
Supplementary Table 5. The shRNAs were cloned into lentiCRISPR-v2-HygR-mCherry vector
under the control of human U6 promoter and packaged into lentivirus for cell transduction. The
WTC11 iPSCs transduced with shRNA lentivirus were treated with hygromycin (Gibco,
10687010, 150 µg/mL) for 7 days and then collected for RNA extraction and qPCR.

ATAC-seq

ATAC-seq was carried out using the Nextera DNA Library Prep Kit (Illumina, FC-121-1030) as previously described[39]. The detailed protocol is available on the ENCODE portal (https://www.encodeproject.org/documents/0317894c-5a42-4f03-b865-c2a2d08708ef/).

Briefly, each library started with 100,000 fresh iPSCs, and the cells were incubated with ice cold nuclei extraction buffer (10 mM TrisHCl pH 7.5, 10 mM NaCl, 3 mM MgCl2, 0.1% Igepal CA630, and 1x protease inhibitor) for 5 min on ice, then centrifuged at 500 RCF for 5 min. 50,000 resulting nuclei were treated with tagmentation buffer (25 µL Buffer TD with 50,000 nuclei, 22.5 µL water, 2.5 µL TDE1) for 30 min at 37°C. The transposed DNA was purified using Qiagen MinElute PCR purification kit (Qiagen, 28006), amplified using Nextera primers, then size-selected for fragments between 150 and 1000 bp using SPRISelect beads (Beckman Coulter, B23319). Libraries were sent for single-end sequencing on the HiSeq 4000 (50 bp single-end reads). Reads were mapped to hg38/GRCh38 and processed using the ENCODE pipeline (https://github.com/kundajelab/atac_dnase_pipelines, V1.8.0), which ran on the default settings. The ATAC-seq peaks were filtered with FDR cutoff of 0.1%, and adjacent peaks were merged if they are less than 1 kb apart.

RNA-seq

RNA was extracted from fresh cells using the RNeasy Plus Mini Kit (Qiagen, 74134). Approximately 1000 ng of extracted RNA was used to prepare libraries for sequencing using the TruSeq Stranded mRNA Library Prep Kit (Illumina, 20020594). Libraries were sent for paired-end sequencing on the NovaSeq 6000 (100 bp paired-end reads). Reads were aligned to hg38/GRCh38 using STAR 2.7.0f[40] with the standard ENCODE settings, and transcript quantification was performed in a strand-specific manner using RSEM 1.3.1[41] with the annotation from GENCODE v32. Only the first read was used, and all reads were trimmed to 51bp using TrimGalore 0.4.5 running the following options: -q 20 --length 20 -- stringency 3 -- trim-n. The edgeR package in R (3.20.9)[37] was used to calculate TMM-normalized FPKM values for each gene based on the expected counts and gene lengths for each library. The mean gene expression across all replicates was used for analysis.

ChIP-seq

ChIP-seq libraries were constructed from 2 million WTC11 iPSCs. Cells were crosslinked in 1%

formaldehyde at room temperature for 20 min and then quenched with 2.5 M glycine at room

temperature for 5 min. Fixed cells were lysed and chromatin was sonicated by Covaris with the

following parameters: cycle 2, 105 W, 200 cycles per burst time for 30 min. Input chromatin was

removed and stored at -20°C for later processing. Magnetic beads (Invitrogen, Dynabeads

Protein A, 10001D) were preincubated with H3K27ac antibody (Active Motif, 39133, Lot

22618011) for 2 hours at 4°C before being added to sheared chromatin. Samples were

incubated overnight at 4°C. Beads were washed 3 times and chromatin was then eluted.

Samples were incubated at 65°C overnight to reverse the crosslinking. DNA was treated with

RNase A for 1 hr at 37°C and Proteinase K (New England Biolabs, 8107) for 1 hr at 55°C. DNA

was purified by phenol-chloroform extraction and ethanol precipitation. Libraries were prepared

using Tru-seq adapters and size-selected using SPRIselect beads prior to amplification and

paired-end sequencing. Libraries were sent for paired-end sequencing on the NovaSeq 6000

(150 bp paired-end reads). Sequencing reads were trimmed to 50 bp and mapped to hg38 using

bowtie2 with the following options: --local --very-sensitive-local --no-unal --no-mixed --no-

discordant --phred33 -I 10 -X 700. Picard Tools was used to remove blacklisted regions and

duplicate reads and MACS2 was used to call peaks on merged replicates at an FDR cutoff of

1%.

CUT&Tag

CUT&Tag libraries were constructed from 150,000 WTC11 iPSCs according to previously

described methods[42]. Cells were lysed in nuclei extraction buffer (20 mM HEPES-KOH pH 7.9,

10 mM MgCl2, 0.1% Triton X-100, 20% glycerol and 1x protease inhibitor) on ice for 10 min.

The samples were spun and resuspended in 100 µl nuclei extraction buffer. Meanwhile, 10 µl

of BioMag Plus Concanavalin A (Bangs Laboratories, BP531) were equilibrated in binding buffer

(1x PBS, 1 mM $CaCl_2$, 1 mM $MgCl_2$ and 1 mM $MnCl_2$). The equilibrated beads were added to

the samples and incubated with rotation for 15 min at 4°C. Nuclei-bound beads were washed

with Buffer 1 (20 mM HEPES-KOH pH 7.9, 150 mM NaCl, 2 mM EDTA, 0.5 mM spermidine,

0.1% BSA and 1x protease inhibitor) and Buffer 2 (20 mM HEPES-KOH pH 7.9, 150 mM NaCl,

0.5 mM spermidine, 0.1% BSA and 1x protease inhibitor). After washing nuclei-bound beads

were resuspended in 50 µl Buffer 2 with 0.5 µl antibody (H3K4me3 from Millipore, 04-745, Lot

3543820 and CTCF from Millipore, 07-729, Lot 3059608) and incubated with rotation overnight

at 4°C. Samples were washed twice with Buffer 2 and resuspended in 50 µl Buffer 2 with

antibody (antibodies-online Inc., Guinea Pig anti-Rabbit IgG, ABIN101961, Lot 42323) and

incubated for 1 hr at room temperature with rotation. Samples were washed again with Buffer 2

and resuspended in 100 µl Buffer 3 (20 mM HEPES-KOH pH 7.9, 300 mM NaCl, 0.5 mM

Spermidine, 0.1% BSA and 1x proteinase inhibitor) containing 0.04 µM pA-Tn5. Samples were

incubated for 1 hr at room temperature, washed three times with Buffer 3 and resuspended in

tagmentation buffer (20 mM HEPES-KOH pH 7.9, 300 mM NaCl, 0.5 mM Spermidine, 10 mM

$MgCl_2$, 0.1% BSA and 1x proteinase inhibitor). Samples were incubated for 1 hr at 37°C.

Samples were treated with Proteinase K (New England Biolabs, 8107) for 1 hr at 50°C. DNA

was purified by phenol-chloroform extraction and ethanol precipitation. Libraries were prepared

using Tru-seq adapters and size-selected using SPRIselect beads prior to amplification and

paired-end sequencing. Libraries were sent for paired-end sequencing on the Mini-seq (37 bp

paired-end reads, H3K4me3 libraries) or NovaSeq 6000 (150 bp paired-end reads, CTCF

libraries). Sequencing reads (CTCF libraries were trimmed to 50 bp) were mapped to hg38

using bowtie2 with the following options: --local --very-sensitive-local --no-unal --no-mixed --no-

discordant --phred33 -I 10 -X 700. Picard Tools was used to remove blacklisted regions and

duplicate reads and SEACR[43] was used to call peaks on merged replicates.

H3K4me3 PLAC-seq

H3K4me3 PLAC-seq data in WTC11 cells were generated as previously described[44] in

biological replicates (clone 6 and clone 28) (https://data.4dnucleome.org/experiment-set-

replicates/4DNESDRL4ZKM/   and   https://data.4dnucleome.org/experiment-set-

replicates/4DNESIZ5TTHO/). We combined the two biological replicates, and applied the MAPS

pipeline[45] to identify significant long-range chromatin interactions at 5 kb bin resolution for the

genomic distance 10 kb ~ 1 Mb. The reference genome is GRCh38. In addition, for each 5 kb

bin pair anchored at H3K4me3 peaks, the MAPS pipeline outputs the normalized contact

frequency, which adjusts for the biases from effective fragment length, GC content, sequence

mappability, H3K4me3 enrichment level and 1D genomic distance effect.

Comparison between strong enhancers and weak enhancers using H3K4me3 PLAC-seq data

For 4 genes *HPRT1*, *MLH1*, *PMS2* and *PCNA*, there are 23 enhancer-promoter pairs between strong enhancers and their target genes, and 21 enhancer-promoter pairs between weak enhancers and their target genes. We mapped each enhancer and promoter of target gene into 5 kb bins, and obtained the distance normalized H3K4me3 PLAC-seq contact frequency for 5 kb bin pairs containing the enhancer-promoter pairs. Since *MSH2* and *MSH6* are located within 407 kb linear genomic distance with each other and we can't assign enhancers to either gene reliably, enhancers identified near *MSH2* and *MSH6* were excluded from this analysis.

Comparison between enhancer-like promoters and control promoters using H3K4me3 PLAC-seq data

For this analysis, control promoters are active promoter regions with annotated ATAC-seq peaks and tested negative as enhancers for the for MMR genes. We mapped each promoter into a 5 kb bin that was used in the PLAC-seq analysis. We only choose the bins with one annotated active promoter, which gave us 31 enhancer-like promoters and 43 control promoters in this analysis. We counted the number of significant H3K4me3 PLAC-seq interactions anchored at the 5 kb bins with these promoter sequences. In addition, as described in our previous study[23], for promoters with at least one significant interaction, we calculated the summation of –log10 FDR of significant interactions, which is a measure of the overall interaction strength.

Chromatin contact frequency comparison between strong enhancers and weak enhancers in K562 cells and mESCs

For the chromatin contact frequency comparison of enhancers in K562 cells and mouse embryonic stem cells (mESCs), we downloaded the identified enhancers from each publication[6, 8], and defined strong enhancer with cutoff of 50% ≤ transcriptional contribution ≤ 100%, weak enhancer with cutoff of 0% < transcriptional contribution ≤ 20%. H3K27ac HiChIP data in K562 cells[46] and H3K4me3 PLAC-seq data in mESCs[45] were used for comparison. The comparisons were performed in 10 kb resolution.

Motif scan and Transcription factor identification

The fasta files were first generated in the hg38 genome for the identified strong enhancers and

weak enhancers separately. For each strong enhancer and weak enhancer, the FIMO software

(version 5.1.0)[47] with human motif database HOCOMOCO (v11 FULL)[48] was used to scan the

motifs. All the FIMO motif scans were in default settings. We then filtered the transcription

factors (TFs) in each strong and weak enhancer loci by FDR cutoff of 0.05 and p-value cutoff of

0.0001 and gene expressions cutoff of FPKM > 1. By taking the TFs with TF motif appeared in

more than 80% enhancers, 47 TFs were considered as commonly appearing in the strong

enhancers, and 35 TFs were in the weak enhancers.

References

1. Soldner, F. et al. Parkinson-associated risk variant in distal enhancer of alpha-synuclein modulates target gene expression. *Nature* **533**, 95-99 (2016).
2. Visel, A., Rubin, E.M. & Pennacchio, L.A. Genomic views of distant-acting enhancers. *Nature* **461**, 199-205 (2009).
3. Long, H.K. et al. Loss of Extreme Long-Range Enhancers in Human Neural Crest Drives a Craniofacial Disorder. *Cell Stem Cell* **27**, 765-783 e714 (2020).
4. Consortium, E.P. et al. Expanded encyclopaedias of DNA elements in the human and mouse genomes. *Nature* **583**, 699-710 (2020).
5. Sanjana, N.E. et al. High-resolution interrogation of functional elements in the noncoding genome. *Science* **353**, 1545-1549 (2016).
6. Gasperini, M. et al. A Genome-wide Framework for Mapping Gene Regulation via Cellular Genetic Screens. *Cell* **176**, 377-390 e319 (2019).
7. Diao, Y. et al. A tiling-deletion-based genetic screen for cis-regulatory element identification in mammalian cells. *Nat Methods* **14**, 629-635 (2017).
8. Fulco, C.P. et al. Activity-by-contact model of enhancer-promoter regulation from thousands of CRISPR perturbations. *Nat Genet* **51**, 1664-1669 (2019).
9. Simeonov, D.R. et al. Discovery of stimulation-responsive immune enhancers with CRISPR activation. *Nature* **549**, 111-115 (2017).
10. Klann, T.S. et al. CRISPR-Cas9 epigenome editing enables high-throughput screening for functional regulatory elements in the human genome. *Nat Biotechnol* **35**, 561-568 (2017).
11. Fulco, C.P. et al. Systematic mapping of functional enhancer-promoter connections with CRISPR interference. *Science* **354**, 769-773 (2016).
12. Gasperini, M. et al. CRISPR/Cas9-Mediated Scanning for Regulatory Elements Required for HPRT1 Expression via Thousands of Large, Programmed Genomic Deletions. *Am J Hum Genet* **101**, 192-205 (2017).
13. Pecina-Slaus, N., Kafka, A., Salamon, I. & Bukovac, A. Mismatch Repair Pathway, Genome Stability and Cancer. *Front Mol Biosci* **7**, 122 (2020).
14. Yan, T., Berry, S.E., Desai, A.B. & Kinsella, T.J. DNA mismatch repair (MMR) mediates 6-thioguanine genotoxicity by introducing single-strand breaks to signal a G2-M arrest in MMR-proficient RKO cells. *Clin Cancer Res* **9**, 2327-2334 (2003).
15. Li, G.M. Mechanisms and functions of DNA mismatch repair. *Cell Res* **18**, 85-98 (2008).
16. Mandegar, M.A. et al. CRISPR Interference Efficiently Induces Specific and Reversible Gene Silencing in Human iPSCs. *Cell Stem Cell* **18**, 541-553 (2016).
17. Tycko, J. et al. Mitigation of off-target toxicity in CRISPR-Cas9 screens for essential non-coding elements. *Nat Commun* **10**, 4063 (2019).
18. Perez, A.R. et al. GuideScan software for improved single and paired CRISPR guide RNA design. *Nat Biotechnol* **35**, 347-349 (2017).
19. Radzisheuskaya, A., Shlyueva, D., Muller, I. & Helin, K. Optimizing sgRNA position markedly improves the efficiency of CRISPR/dCas9-mediated transcriptional repression. *Nucleic Acids Res* **44**, e141 (2016).
20. Rosenbluh, J. et al. Complementary information derived from CRISPR Cas9 mediated gene deletion and suppression. *Nat Commun* **8**, 15403 (2017).
21. Li, K. et al. Interrogation of enhancer function by enhancer-targeting CRISPR epigenetic editing. *Nat Commun* **11**, 485 (2020).

22. Engreitz, J.M. et al. Local regulation of gene expression by lncRNA promoters,
transcription and splicing. *Nature* **539**, 452-455 (2016).
23. Song, M. et al. Cell-type-specific 3D epigenomes in the developing human cortex. *Nature*
587, 644-649 (2020).
24. Kubo, N. et al. Promoter-proximal CTCF binding promotes distal enhancer-dependent
gene activation. *Nat Struct Mol Biol* (2021).
25. Fernandez-Zapico, M.E. et al. A functional family-wide screening of SP/KLF proteins
identifies a subset of suppressors of KRAS-mediated cell growth. *Biochem J* **435**, 529-
537 (2011).
26. Ren, B. et al. E2F integrates cell cycle progression with DNA repair, replication, and
G(2)/M checkpoints. *Genes Dev* **16**, 245-256 (2002).
27. Polager, S., Kalma, Y., Berkovich, E. & Ginsberg, D. E2Fs up-regulate expression of
genes involved in DNA replication, DNA repair and mitosis. *Oncogene* **21**, 437-446
(2002).
28. Canver, M.C. et al. BCL11A enhancer dissection by Cas9-mediated in situ saturating
mutagenesis. *Nature* **527**, 192-197 (2015).
29. Rajagopal, N. et al. High-throughput mapping of regulatory DNA. *Nat Biotechnol* **34**, 167-
174 (2016).
30. Koike-Yusa, H., Li, Y., Tan, E.P., Velasco-Herrera Mdel, C. & Yusa, K. Genome-wide
recessive genetic screening in mammalian cells with a lentiviral CRISPR-guide RNA
library. *Nat Biotechnol* **32**, 267-273 (2014).
31. Adamson, B. et al. A Multiplexed Single-Cell CRISPR Screening Platform Enables
Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882 e1821
(2016).
32. Diao, Y. et al. A new class of temporarily phenotypic enhancers identified by
CRISPR/Cas9-mediated genetic screening. *Genome Res* **26**, 397-405 (2016).
33. Zabidi, M.A. et al. Enhancer-core-promoter specificity separates developmental and
housekeeping gene regulation. *Nature* **518**, 556-559 (2015).
34. Dao, L.T.M. & Spicuglia, S. Transcriptional regulation by promoters with enhancer
function. *Transcription* **9**, 307-314 (2018).
35. Mikhaylichenko, O. et al. The degree of enhancer or promoter activity is reflected by the
levels and directionality of eRNA transcription. *Genes Dev* **32**, 42-57 (2018).
36. Thomas, H.F. et al. Temporal dissection of an enhancer cluster reveals distinct temporal
and functional contributions of individual elements. *Mol Cell* (2021).
37. Robinson, M.D., McCarthy, D.J. & Smyth, G.K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
140 (2010).
38. Ramirez, F. et al. deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **44**, W160-165 (2016).
39. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. & Greenleaf, W.J. Transposition
of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-
binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218 (2013).
40. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
41. Li, B. & Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with
or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

862 42. Kaya-Okur, H.S. et al. CUT&Tag for efficient epigenomic profiling of small samples and
863   single cells. *Nat Commun* **10**, 1930 (2019).
864 43. Meers, M.P., Tenenbaum, D. & Henikoff, S. Peak calling by Sparse Enrichment Analysis
865   for CUT&RUN chromatin profiling. *Epigenetics Chromatin* **12**, 42 (2019).
866 44. Fang, R. et al. Mapping of long-range chromatin interactions by proximity ligation-
867   assisted ChIP-seq. *Cell Res* **26**, 1345-1348 (2016).
868 45. Juric, I. et al. MAPS: Model-based analysis of long-range chromatin interactions from
869   PLAC-seq and HiChIP experiments. *PLoS Comput Biol* **15**, e1006982 (2019).
870 46. Mumbach, M.R. et al. Enhancer connectome in primary human cells identifies target
871   genes of disease-associated DNA elements. *Nat Genet* **49**, 1602-1612 (2017).
872 47. Grant, C.E., Bailey, T.L. & Noble, W.S. FIMO: scanning for occurrences of a given motif.
873   *Bioinformatics* **27**, 1017-1018 (2011).
874 48. Kulakovskiy, I.V. et al. HOCOMOCO: towards a complete collection of transcription factor
875   binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids*
876   *Res* **46**, D252-D259 (2018).
877

Figure 1

Figure 1. CRISPRpath for identifying enhancers of multiple genes.
(**a**) Six genes (*HPRT1*, *MSH2*, *MSH6*, *MLH1*, *PMS2* and *PCNA*) in the 6TG-induced mismatch repair process were used for CRISPRpath screen in this study. (**b**) Schematic of the CRISPRpath screening strategy with 6TG treatment in iPSCs. Cell survival was used as readout for the screen. (**c**) Spearman correlation analysis of sgRNA ranking based on fold change for CRISPRpath screens with different 6TG concentrations (1X, 2X and 3X). (**d**) Venn diagram shows the overlapping enriched sgRNAs identified from the screens with 2X and 3X 6TG treatments. (**e**) Box plots show the fold change of the enriched distal and proximal sgRNAs from 2X, 3X CRISPRi and CRISPRn screens. Star indicates no enriched distal sgRNA was identified from 3X CRISPRn screen. Boxplots indicate the median, interquartile range (IQR), Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. (**f**) Bar plot shows the number of enriched distal and proximal sgRNAs from 2X, 3X CRISPRi and CRISPRn screens. Star indicates no enriched distal sgRNA was identified from 3X CRISPRn screen. (**g**) Venn diagram shows the identified enhancers from each CRISPRpath screen.

Figure 2

Figure 2. Genomic features of identified enhancers from CRISPRpath using CRISPRi.
(**a**) Genomic locations of identified enhancers relative to TSS. Circles indicate enhancers identified from the CRISPRi 3X screen (red), enhancers uniquely identified from the CRISPRi 2X screen (blue), tested CREs that are not identified as enhancers (grey). Purple lines label the location of each target gene. (b) Histogram shows the distance distribution between identified enhancers and their paired TSS. (**c**) Histogram shows the number of interval genes between enhancers and their target gene TSS. Mean is indicated with an orange dashed line and only the enhancers for *MLH1*, *PMS2*, *PCNA*, *HPRT1* are included in **b** and c. (**d**) A weak negative correlation is observed between enrichment score and genomic distance between enhancers and their target genes (Pearson correlation, r = -0.36, *P* = 0.01). Black circles indicate promoters. The red and blue circles are enhancers showing in **a**. (**e**) Density plot shows no significant difference (two-tailed two-sample Kolmogorov–Smirnov test) for the distribution of all distal sgRNAs (gray), enriched distal sgRNAs from 2X (blue) and 3X screen (red) CRISPRi screens.

Figure 3

Figure 3. Enhancer-like promoters act as functional enhancers.
(**a, b, c**) Three examples of promoters function as enhancers. CRISPRi silencing of the promoter region of *SOCS5*, *FOXN2* and *TMEM230* results in significant downregulation of *MSH6*, *MSH2* and *PCNA*, respectively. shRNA knockdown of *SOCS5*, *FOXN2* and *TMEM230* can only downregulate *SOCS5*, *FOXN2* and *TMEM230* expression. Three independent replicates per condition and two independent sgRNAs or shRNAs per replicate were used for each experiment. *P* values are from two-tailed two-sample t-test. (**d**) Average signal enrichment of ATAC-seq, gene transcription, H3K4me3, H3K27ac and CTCF binding for enhancer-like promoters (n = 38) and control promoters (n = 47). *P* values are from Wilcoxon test. Boxplots indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. (**e**) Number of H3K4me3 mediated chromatin interactions and cumulative interaction score for enhancer-like promoters (n = 31) and control promoters (n = 43). Boxplots indicate median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. *P* values are calculated from Wilcoxon test.

Figure 4

Figure 4. CRISPRpath can distinguish weak and strong enhancers by imposing different selection pressures.
(**a**) Box plots show the enrichment score of the tested elements. TSS regions (black circles) show highest enrichment scores. Enhancers uniquely identified from the lower selection pressure (CRISPRi 2X, blue circles) exhibit lower enrichment scores compared to the enhancers identified from the higher selection pressure (CRISPRi 3X, red circles). *P* values are from Wilcoxon test. (**b**) Box plots show the CRISPRi perturbation at enhancers induced various degrees of transcriptional repression of target genes measured with RT-qPCR. Each dot represents the average value from three biological replicates. CRISPRi targeting TSS regions (dark gray) achieved the highest transcriptional repression. CRISPRi targeting strong enhancers (pink) leads a more substantial transcription silencing on target gene compared to CRISPRi targeting weak enhancers (cyan). *P* values are from Wilcoxon test. (**c**) Enrichment analysis of ATAC-seq, H3K27ac, H3K4me3, CTCF binding signals for strong (n = 33) and weak (n = 30) enhancers. Boxplots indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. *P* values for the difference between strong and weak enhancers are from Wilcoxon test; see Supplementary Table 7 for *P* values of all pairwise comparisons. (**d**) Intersection of genomic features for weak enhancers (blue bar) and strong enhancers (red bar). (**e**) Distance normalized H3K4me3 PLAC-seq contact frequency for strong (n = 23) and weak (n = 21) enhancers. Only the enhancers for *MLH1*, *PMS2*, *PCNA*, *HPRT1* are included (see Methods). Boxplots indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. *P* value is from Wilcoxon test. (**f**) Heatmap shows the frequency of transcription factor motifs found in strong and weak enhancers.

Supplementary Figure 1

Supplementary Figure 1. Features of sgRNA library for CRISPRpath screen.
(**a**) Bar graph shows the number of distal ATAC-seq peaks used as candidate CREs for six target genes. (**b**) Histogram shows size distribution of distal ATAC-seq peaks. The average size is 488 bp (blue dash line). (**c, e**) The composition of the sgRNA library. In total, 35,763 sgRNAs were included in the library (c), and 12,702 sgRNAs are high quality sgRNAs (**e**). (**d, f**) Distribution of the number of sgRNAs per distal ATAC-seq peak. Average numbers of sgRNA per ATAC-seq peak are indicated with blue dash lines.

Supplementary Figure 2

Supplementary Figure 2. Quality of the sgRNA library and CRISPRpath screen libraries.
(**a**) Distribution of sgRNA oligo read counts in the sgRNA library. (**b**) Cumulative frequency of sgRNAs in the sgRNA library. (**c**) Distribution of high quality sgRNAs read counts in the sgRNA library. (**d**) Cumulative frequency of high quality sgRNAs in the sgRNA library. The constructed sgRNA plasmid library recorvered all the designed sgRNAs with the copy number difference less than five fold for at least 97% designed sgRNAs. (**e**) PCA analysis shows the high reproducibility of the CRISPRpath screen libraries between biological replicates.

Supplementary Figure 3

Supplementary Figure 3. P value cutoff used for identifying enriched sgRNAs from each screens.
(**a-f**) Distribution of P value for tested distal, proximal and non-targeting control sgRNA groups. Orange dash lines indicate 5% percentile of the *P* values from non-targeting control sgRNAs to achieve a false discovery rate of 5%.

Supplementary Figure 4

Supplementary Figure 4. Enriched proximal sgRNAs and sgRNA ranking analysis.
(**a**) Number and fold change of the enriched proximal sgRNAs for the six target genes from CRISPRi and CRISPRn screens. The color indicates fold changes, and the size of circle indicates the number of enriched sgRNAs. (**b**) Enrichment analysis shows the enriched proximal sgRNAs bias towards to the TSS region for CRISPRi screens and the protein coding region (CDS) for CRISPRn screens. Color represents the number of enriched sgRNAs. (**c**) Spearman correlation analysis of the distal and proximal sgRNAs ranking shows proximal sgRNAs exhibiting higher correlation between each screen compared to distal sgRNAs.

Supplementary Figure 5

Supplementary Figure 5. Enriched sgRNAs identified from CRISPRi screens exhibit no position and strand preference.

(**a**) Enriched sgRNAs from CRISPRi 2X (red dots, n=448) and CRISPRi 3X (red dots, n=260) screens showed similar distributions across candidate CREs. (**b**) Odds ratio analysis of the fold change of enriched sgRNAs shows enriched sgRNAs have no strand preference. Enhancers with enriched sgRNAs only targeting one strand were exclude for the analysis. Odds ratio was calculated for each element with the equation of ave(log2(fold change of sgRNA targeting plus strand)) / ave(log2(fold change of sgRNA targeting minus strand)). Violin plots show the distributions of odds ratio values within each screen, and boxplots indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR.

Supplementary Figure 6. Validation of CIRSPRpath identified enhancers.
(**a**) Validation of the strong (black) and weak (grey) enhancers with CRISPRi followed by RT-qPCR. Three independent replicates per condition. The significance was calculated with two-tailed two-sample t-test. Data are mean and s.d. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. (**b**) CRISPRi-mediated transcriptional repression of six target genes by targeting TSS of each gene. Three independent replicates per condition. The significance was calculated with two-tailed two-sample t-test. Data are mean and s.d. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. (**c**) Pearson correlation analysis reveals element enrichment score from CRISPRpath screens correlates with element effect size on transcription from CRISPRi (Pearson correlation, PCC = -0.68, $P = 3.2 \times 10\text{-}5$).

Supplementary Figure 7

Supplementary Figure 7. Chromatin contract frequency analysis for the enhancers in K562 cells and mESCs.
(**a**) Distance normalized H3K27ac HiChIP contact frequency for strong (n = 34) and weak (n = 82) enhancers identified with crisprQTL mapping in K562 cells. (**b**) Distance normalized H3K27ac HiChIP contact frequency for strong (n = 2) and weak (n = 20) enhancers identified with CRISPRi-FlowFISH screen in K562 cells. (**c**) Distance normalized H3K4me3 PLAC-seq contact frequency for strong (n = 10) and weak (n = 3) enhancers identified in mouse embryonic stem cells. Boxplots indicate the median, IQR, Q1 − 1.5 × IQR and Q3 + 1.5 × IQR. P values are from Wilcoxon test.