1 TITLE

2 Training-induced bioenergetic improvement in human skeletal muscle is associated with non-

3 stoichiometric changes in the mitochondrial proteome without reorganization of respiratory

4 chain content

5 AUTHORS

6 Cesare Granata[1,2,11,*], Nikeisha J. Caruana[2,3,11], Javier Botella[2], Nicholas A. Jamnick[2,4], Kevin

7 Huynh[5], Jujiao Kuang[2], Hans A. Janssen[2], Boris Reljic[3,6], Natalie A. Mellett[5], Adrienne

8 Laskowski[1,7], Tegan L. Stait[7], Ann E. Frazier[7,8], Melinda T. Coughlan[1,9], Peter J. Meikle[5],

9 David R. Thorburn[7,8,10], David A. Stroud[3,12,*] and David J. Bishop[2,12,13,*]

10 AFFILIATIONS

[1]Department of Diabetes, Central Clinical School, Monash University, Melbourne, VIC,

3004, Australia

[2]Institute for Health and Sport (IHES), Victoria University, Melbourne, VIC, 3011, Australia

[3]Department of Biochemistry and Pharmacology and Bio21 Molecular Science and

Biotechnology Institute, The University of Melbourne, Parkville, VIC, 3010, Australia

[4]Metabolic Research Unit, School of Medicine and Institute for Mental and Physical Health

and Clinical Translation (iMPACT), Deakin University, Geelong, VIC, Australia

[5]Metabolomics Laboratory, Baker Heart & Diabetes Institute, Melbourne, VIC, 3004,

Australia

[6]Current address: Department of Biochemistry and Molecular Biology, Monash Biomedicine

Discovery Institute, Monash University, 3800, Melbourne, Australia.

[7]Murdoch Children's Research Institute, Royal Children's Hospital, Melbourne, VIC, 3052,

Australia

[8]Department of Paediatrics, The University of Melbourne, Melbourne, VIC, 3052, Australia

[9]Baker Heart & Diabetes Institute, Melbourne, VIC, 3004, Australia

[10]Victorian Clinical Genetics Services, Royal Children's Hospital, Melbourne, VIC, 3052,

Australia

[11]Co-first author

[12]Co-senior author

[13]Lead contact

*Correspondence: cesare.granata@monash.edu - cesare.granata@ddz.de (C.G.),

david.stroud@unimelb.edu.au (D.A.S), david.bishop@vu.edu.au (D.J.B)

SUMMARY

Mitochondrial defects are implicated in multiple diseases and aging. Exercise training is an

accessible and inexpensive therapeutic intervention improving mitochondrial bioenergetics

and quality of life. By combining a multi-omics approach with biochemical and *in silico*

normalization, we removed the bias arising from the training-induced increase in human

skeletal muscle mitochondrial content to unearth an intricate and previously undemonstrated

network of differentially prioritized mitochondrial adaptations. We show that changes in

hundreds of transcripts, proteins, and lipids are not stoichiometrically linked to the increase in

mitochondrial content. We demonstrate that enhancing electron flow to oxidative

phosphorylation (OXPHOS) is more important to improve ATP generation than increasing

the abundance of the OXPHOS machinery, and that training-induced supercomplex

formation does not confer enhancements in mitochondrial bioenergetics. Our study provides a

new analytical approach allowing unbiased and in-depth investigations of training-induced

mitochondrial adaptations, challenging our current understanding and calling for careful

reinterpretation of previous findings.

INTRODUCTION

Mitochondria are the main site of energy conversion in the cell and have critical roles in other important biological processes (Spinelli and Haigis, 2018). Owing to this, defects in mitochondria have been implicated in multiple diseases, medical conditions, and aging (Gorman et al., 2016; Lin and Beal, 2006; López-Otín et al., 2013; Nunnari and Suomalainen, 2012). The development of interventions to improve the content and function of mitochondria is therefore essential to enhance quality of life and to extend life expectancy.

Exercise training is one of the most widely accessible interventions or "therapies" to stimulate mitochondrial biogenesis (Holloszy, 1967; Hoppeler et al., 1973). As arguably the most extensive and also natural perturbation, exercise is also an excellent experimental model for understanding both the complex nature of mitochondrial biogenesis and the plasticity of the skeletal muscle mitochondrial proteome in humans (Neufer et al., 2015). It is well accepted that exercise training induces an increase in mitochondrial content and respiratory function in human skeletal muscle (Granata et al., 2018b). However, our knowledge of mitochondrial adaptations is limited to a small fraction of the mitochondrial proteome (Granata et al., 2018a), and is complicated by large training-induced changes in mitochondrial content, which limit the ability to interpret adaptations in individual proteins independent of the overall change in mitochondrial content. To account for this, most studies to date have either performed analyses in mitochondrial isolates obtained after differential centrifugation or have employed a crude normalization strategy involving the use of markers of mitochondrial content, such as citrate synthase (CS) activity (Granata et al., 2018b; Groennebaek et al., 2020). Nonetheless, due to limitations in both approaches (Franko et al., 2013; Groennebaek et al., 2020), there remain major gaps in our understanding of the magnitude, timing, and direction of change in individual mitochondrial proteins and related functional pathways, independent of the overall change in mitochondrial content.

A key biological function of mitochondria is energy conversion via oxidative

phosphorylation (OXPHOS), which is carried out by the four multi-protein complexes

(complexes I to IV [CI-CIV]) of the electron transport chain (ETC) and F_0F_1-ATP synthase

(or CV) (Pesta and Gnaiger, 2012). Assembly of the OXPHOS complexes is an intricate

process requiring the coordination of two genomes (nuclear and mitochondrial) and non-

subunit proteins known as assembly factors (Signes and Fernandez-Vizarra, 2018). Only two

studies have investigated training-induced changes in OXPHOS complexes in humans

(Greggio et al., 2017; Wyckelsma et al., 2017), reporting that exercise differentially

modulates the assembly and composition of respiratory chain complexes - including their

high molecular weight supercomplex (SC) assemblies. Furthermore, no study has

investigated if training-induced changes in chaperones, assembly factors, and single

OXPHOS subunits occur in conjunction with, or precede, changes in the complete, functional

complexes. Many other processes support assembly and maintenance of the OXPHOS

complexes, including mitochondrial protein import and assembly pathways, mitochondrial

DNA transcription and translation, metabolite carriers, and pathways producing critical

cofactors such as coenzyme Q and iron-sulphur proteins (Fe-S) (Signes and Fernandez-

Vizarra, 2018). In turn, these processes are reliant on pathways supporting mitochondrial

biogenesis more broadly, such as lipid biogenesis and maintenance of mitochondrial

morphology. Whether these processes and pathways are activated by exercise training and at

which specific times remains unknown.

Here, we employed a 3-phase training intervention, combined with a multi-omics approach

(transcriptomics, proteomics, and lipidomics) and a newly developed normalization strategy

that removed the bias induced by large alterations in mitochondrial content, to investigate

mitochondrial adaptations in human skeletal muscle. Our findings revealed an intricate and

timely remodeling in the mitochondrial transcriptome, proteome, and lipidome, independent

of changes in overall mitochondrial content. Post-normalization, we identified 185

differentially expressed mitochondrial proteins, representing multiple protein functional

classes, as well as bioenergetic and metabolic pathways, that followed distinct patterns of

adaptation. In addition, we provide evidence that training-induced changes in the abundance

and/or organization of SCs do not contribute to improvements in mitochondrial bioenergetics.

Our study provides a new analytical approach that expands our fundamental understanding of

training-induced mitochondrial biogenesis. It challenges our current knowledge and calls for

careful reinterpretation of previous findings, while leading to new hypotheses on how

training alters mitochondrial protein content and function, thus providing an important

resource with implications for both health and disease.

RESULTS AND DISCUSSION

Increases in mitochondrial content underlie changes in markers of mitochondrial

biogenesis and mitochondrial respiratory function following training. The traditional view

of mitochondrial biogenesis suggests that changes in individual mitochondrial proteins are

stoichiometrically linked to changes in mitochondrial content. Although this has been

questioned previously (Ørtenblad, 2018), it remains an unresolved matter. To challenge this

paradigm, we employed an experimental model in which ten men were sequentially exposed

to different volumes (normal- [NVT], high- [HVT] and reduced- [RVT] volume training) of

high-intensity interval training (HIIT) with biopsies taken at baseline (BL) and post-NVT

(PN), post-HVT (PH) and post-RVT (PR) (Figure 1A). This design was chosen as we have

previously demonstrated that changes (increases or decreases) in training volume lead to

changes in mitochondrial content and markers of mitochondrial biogenesis (Bishop et al.,

2019; Bishop et al., 2014; Granata et al., 2018b; Granata et al., 2016a). Weekly training

volumes for the three phases are shown in Figure 1B; changes in physiological and

performance parameters are presented in Table 1.

Figure 1. RNA-seq highlights a complex network of transcriptional responses underpinning the training-induced increases in markers of mitochondrial content in human vastus lateralis muscle

(A) Study design; colored boxes indicate a training phase: NVT (normal-volume training), HVT (high-volume training), and RVT (reduced-volume training), whereas open boxes indicate a testing session and resting skeletal muscle biopsy: baseline (BL), post-NVT (PN), post-HVT (PH), and post-RVT (PR)

(B) Weekly training volume during the NVT, HVT, and RVT training phase; training volume was calculated by multiplying the absolute exercise intensity in Watts by the effective duration of exercise training in minutes (excluding the warm up and the rest periods between intervals) by the total number of training sessions in each phase.

(C) Mitochondrial yield per mg of tissue achieved during mitochondria isolation of human vastus lateralis biopsy samples (biochemical enrichment of mitochondrial protein).

(D) Citrate synthase (CS) enzyme activity assessed in whole-tissue (vastus lateralis) homogenates.

(E) Heatmap of differentially expressed transcripts between training phases determined with an adjusted $P < 0.05$ (Benjamini Hochberg). Row clustering determined by unsupervised hierarchical cluster analysis.

(F) Profile plots showing relative scaled transcripts of subunits of the five oxidative phosphorylation (OXPHOS) complexes (CI to CV), mitochondrial ribosomes (mtribo), OXPHOS assembly, TCA cycle, SLC25As and mitochondrial contact site and cristae organising system (MICOS) complex. Transcripts grouped according to known literature (see methods).

(G), Biological process (BP) gene ontology of all differentially expressed transcripts as in (E); the five terms with the highest combined score, as determined by *Enrichr* (see methods), are displayed.

HIIT: high-intensity interval training; MJ: megajoules; CI-V: complex I to V; IB: immunoblotting; ▼,◊,○, and ● represent individual values; bars represent mean values. Datasets (B), (C), and (D) were analyzed by repeated measures one-way ANOVA followed by Tukey's post hoc testing; n = 10; significance: $P < 0.05$. Dataset (E), (F), and (G) n = 5.

Table 1. Participants' physiological and endurance performance measurements before and after each training phase

Measurement	BL	PN	PH	PR
Age	22.3 ± 5.2			
Height	178.1 ± 10.7			
Body mass	78.8 ± 12.1	79.0 ± 12.9	78.0 ± 12.5	$77.9 \pm 11.8^{\#}$
BMI	24.8 ± 2.9	24.9 ± 3.2	24.6 ± 3.0	$24.5 \pm 2.9^{\#}$
$\dot{V}O_{2Peak}$	46.7 ± 8.2	47.7 ± 8.8	$50.5 \pm 6.0^{*\#}$	$50.0 \pm 6.0^{*}$
$\dot{W}_{LT}$	174.1 ± 30.7	182.5 ± 33.0	$222.2 \pm 36.2^{*\#}$	$219.6 \pm 31.4^{*\#}$
$\dot{W}_{peak}$	231.1 ± 43.2	233.2 ± 43.5	$266.3 \pm 42.4^{*\#}$	$264.5 \pm 40.4^{*\#}$
20k-TT time	2321.0 ± 168.1	2309.1 ± 257.4	$2094.6 \pm 130.8^{*\#}$	$2030.1 \pm 140.5^{*\#}$

BL: baseline; PN: post normal-volume training; PH: post high-volume training; PR: post
reduced-volume training; BMI: body mass index [kg m^{-2}]; $\dot{V}O_{2Peak}$: peak oxygen uptake [mL
min^{-1} kg^{-1}]; $\dot{W}_{LT}$: power at the lactate threshold [Watts]; $\dot{W}_{peak}$: peak power output [Watts];
20k-TT: 20-km time trial [seconds]. All values are mean ± SD; n = 10 for all analyses; all
datasets analysed by repeated measures one-way ANOVA followed by Tukey's post hoc
testing. Significantly different ($P < 0.05$) * vs. BL, # vs. PN.

We first verified that changes in training volume induced changes in the yield of

mitochondrial proteins isolated from muscle biopsies (Figure 1C), and citrate synthase (CS)

activity (Figure 1D) - a commonly used biomarker of mitochondrial content (Larsen et al.,

2012), as previously reported (Barnard et al., 1970; Egan et al., 2011; Granata et al., 2018b;

Granata et al., 2016a; Holloszy, 1967; Morgan et al., 1971). We then confirmed that classic

markers of mitochondrial biogenesis, such as the protein content of OXPHOS subunits

assessed in whole-muscle lysates (Figure S1A, upper panels), as well as measurements of

mitochondrial respiratory function, such as mitochondrial respiration in permeabilized fibers

and ETC enzyme activity (Figure S1B, and C, respectively, all upper panels) changed, for the

most part, as a consequence of changes in training volume. RVT resulted in no change in any

of these markers, suggesting that the training stimulus was sufficient to prevent a decrease in

these parameters. Subsequently, to investigate the underpinnings of training-induced changes

in the above markers, we normalized these values by CS activity, a widely-used

normalization strategy to account for differences in mitochondrial content (Granata et al.,

2018b; Granata et al., 2016a; Larsen et al., 2012; Meinild Lundby et al., 2018). This

demonstrated that, for the most part, absolute changes in markers of mitochondrial biogenesis

and respiratory function were stoichiometrically associated with the overall increase in

mitochondrial content (Figure S1A, B, and C, all lower panels).

Disentangling changes in the mitochondrial proteome from the general increase in

mitochondrial content observed post-training. The above findings seem to indicate that

altering training volume results in a stoichiometric relationship between training-induced

changes in mitochondrial content and markers of mitochondrial biogenesis and mitochondrial

respiratory function (Granata et al., 2018b). However, whether there is also a fixed

stoichiometry between changes in mitochondrial content and individual mitochondrial

proteins, protein functional classes and metabolic pathways, remains undetermined. To test

this, we first employed RNA sequencing (RNA-seq) based transcriptomics (Table S1). We

identified 1206 differentially expressed transcripts (Figure 1E, Table S2), and we observed

the expected increase in gene transcripts encoding subunits of the OXPHOS complexes

(Figure 1F, upper panels, compare with Figure S1A, upper panels) and enzymes of the

tricarboxylic acid (TCA) cycle (Figure 1F, central lower panel), consistent with previous

studies in humans (Pillon et al., 2020). However, transcripts for genes encoding other

mitochondrial proteins did not all follow the same trend (Figure 1F, lower panels). Moreover,

gene ontology enrichment analysis of all differentially expressed transcripts did not identify

significant alterations of pathways involved in mitochondrial respiratory function (Figure

1G), despite the increase in mitochondrial respiration and ETC enzyme activity reported

above (Figure S1B and C, respectively). This reveals a greater complexity in the

transcriptional responses to exercise than previously hypothesized; i.e., not all mitochondria-

related gene transcripts changed in the same direction and with a similar magnitude following

different training phases (Figure 1F).

As transcriptomics suggested there may not be a fixed stoichiometry between training-

induced changes in mitochondrial content and individual mitochondrial proteins, we sought

to support these findings with quantitative proteomics. To negate the general increase in

mitochondrial content following exercise training (Figure 1D) and to avoid this inherent bias

on our proteomics measurements, we performed label-free quantitative (LFQ) proteomics on

equal amounts of mitochondria-enriched fractions (biochemical enrichment of mitochondrial

proteins using differential centrifugation). We quantified 1,411 proteins - 726 of which were

annotated as mitochondrial based on either the Mitocarta2.0 (Calvo et al., 2016) or the

Integrated Mitochondrial Protein Index (IMPI; Known and Predicted Mitochondrial) (Smith

and Robinson, 2019) databases (Table S3). Considering only the high-confidence IMPI

(Known Mitochondrial) dataset, mitochondrial proteins represented 41% of the total number

of proteins identified in our mitochondria-enriched fraction (584 annotated mitochondrial

proteins), contributing to 32% of the overall protein abundance based on raw intensity data

for each protein (Table S3). This demonstrated the presence of a significant proportion of co-

isolating non-mitochondrial proteins, with enrichment analysis indicating these to be

predominantly of myofibrillar origin (Figure S2A), consistent with most mitochondria in

skeletal muscle being tightly associated with myofibrils (Egan et al., 2011). Surprisingly, we

still observed a relative increase in the abundance of mitochondrial proteins within isolates

following training (Figure 2A, upper panel, magenta profiles), an artefact that likely stems

from differences in mitochondrial protein enrichment (MPE) - the contribution of

mitochondrial protein intensities in biochemically isolated mitochondria relative to all protein

intensities - at different time points. Indeed, the MPE for the 4 different time points in the

present study was significantly different (Figure 2A, middle panel) and corresponded to

24.5% at baseline (BL), 31.5% post-NVT (PN), 39.3% post-HVT (PH), and 34.5% post-RVT

(PR) (Table S3) - a pattern consistent with our assessment of mitochondrial content (compare

Figure 2A, middle panel with Figure 1C and D).

Figure 2. Correcting for training-induced changes in mitochondrial content reveals differentially expressed protein clusters with divergent prioritization in response to training

230   (A) Training-induced differences in the mitochondrial protein content of isolated

231   mitochondrial (IM) fractions obtained from vastus lateralis muscle biopsies. Upper panel:

profile plot of all non-normalized intensity values displaying "known mitochondrial" (IMPI "Known Mitochondrial") (magenta) and non-mitochondrial (salmon) proteins co-precipitating during mitochondrial isolation. Proteins identified in less than 70% of samples were removed. Middle panel: mitochondrial protein enrichment (MPE) obtained by adding the raw intensity values from LC-MS/MS analysis of all "known mitochondrial" proteins (IMPI "Known Mitochondrial") at the four different time points. ▼,◊,○, and ● represent individual values; bars represent mean values; analyzed by repeated measures one-way ANOVA followed by Tukey's post hoc testing. Significance: $P < 0.05$. Lower panel: same as upper panel but following removal of non-mitochondrial proteins and statistical correction (BESt-) normalization.

(B) Multidimensional scaling analysis showing segregation of samples obtained from IM fractions of BESt-normalized proteomics values.

(C) Scaled profile plots showing the relative abundance of subunits of OXPHOS complexes and mtribo of the non-normalized intensity values of "known mitochondrial" proteins (IMPI "Known Mitochondrial") following valid value filtering (left panels), and values obtained after BESt-normalization (right panels).

(D) Venn diagram representing the overlap in differentially expressed proteins between the present study and previous literature utilizing young healthy humans (Egan et al., 2011; Holloway et al., 2009; Hostrup et al., 2018). The bubble size for each study is proportional to the number of differentially expressed proteins identified (this was not possible for the overlapping size).

(E) Heatmap of BESt-normalized differentially expressed mitochondrial proteins (IMPI "Known Mitochondrial"), determined with an adjusted $P < 0.01$ (Benjamini Hochberg). Row clustering determined by unsupervised hierarchical cluster analysis.

(F) Venn diagram representation of the gene ontology of the six clusters identified in (E) obtained from enrichment analysis (using the ClueGO [v2.5.6] application in Cytoscape [v3.7.1]; see methods) of all proteins belonging to each cluster performed separately (Table S5).

(G) Venn diagram representations of the number of BESt-normalized differentially expressed mitochondrial proteins (IMPI "Known Mitochondrial") between each of the time points. Differentially expressed proteins for each time point comparison were identified by linear modelling with an adjusted P value < 0.01 using the Benjamini Hochberg method, as shown in Table S7.

BL: baseline; PN: post-NVT; PH: post-HVT; PR: post-RVT. n = 5 for **a**, **b**, and **c**, n = 10 for all other analyses.

The above findings highlight that the standard approach of relying on biochemical

enrichment to eliminate the bias arising from different MPEs across time points is insufficient

and that an additional strategy is needed. We therefore complemented our biochemical

enrichment approach with a novel statistical correction strategy consisting of *i*) removing

proteins that were identified in less than 70% of samples, *ii*) removing all except high-

confidence mitochondrial proteins (IMPI Known Mitochondrial), and *iii*) undertaking

variance stabilizing normalization (VSN) (see methods for more detail). This resulted in the

retention of 498 mitochondrial proteins normalized across the dataset with high confidence

(Table S4). Unlike CS activity, which is a single value for each sample, our normalization

strategy utilizes the trends of hundreds of proteins and reduces the confounding effect of

potential outliers. This is extremely important, given that previous studies have shown a > 5-

fold difference in training-induced synthesis rate between fast and slow synthesized

mitochondrial proteins in human skeletal muscle (Groennebaek et al., 2020; Scalzo et al.,

2014; Shankaran et al., 2016). Plotting the resulting data revealed that the general increase in

mitochondrial proteins with different training volumes had been compensated for, indicating

that our <u>b</u>iochemical <u>e</u>nrichment and <u>st</u>atistical correction (BESt-normalization) was

successful (Figure 2A, lower panel). In further support of our normalization strategy,

multidimensional scaling analysis showed good segregation of samples corresponding to the

different time points (Figure 2B).

To assess the power of our normalization strategy, we compared results from non-normalized

and BESt-normalized proteomics values. Whereas analysis of non-normalized intensity

values revealed a lack of change in the abundance of OXPHOS and mitochondrial ribosomal

subunits post-NVT, followed by a marked increase post-HVT, and a decrease post-RVT

(Figure 2C, left panels), results from BESt-normalized values showcased a very different

picture, as the content of OXPHOS subunits was markedly decreased (deprioritized) post-

NVT, and showed discreet and specific changes (mainly modest increases) thereafter (Figure

2C, right panels; results discussed in more detail later), whereas mitochondrial ribosomal

subunits increased to a greater extent than mitochondrial content (prioritized) (Figure 2C,

right panels). These findings demonstrate the power of our BESt-normalization strategy

compared to classical approaches, as it enables the first unbiased comparison of the changes

in individual mitochondrial proteins relative to the overall increase in mitochondrial content

following training, providing clear evidence that these changes are not stoichiometrically

linked. As a means of comparison, we also employed the classical approach of normalizing

the abundances of specific proteins derived from SDS-PAGE and immunoblot analysis on

isolated mitochondrial fractions by CS activity; this demonstrates that while this widely used

approach partially corrects for changes in MPE it fails to recapitulate most of the findings

observed with our BESt-normalization strategy (Figure S2B).

From this point onwards, unless specified otherwise, post BESt-normalization proteomics

results will be presented, enabling the investigation of training-induced changes in

mitochondrial proteins without the confounding influence of changes in mitochondrial

content.

Discovery of novel mitochondrial proteins and functional classes and metabolic pathways

differentially affected by exercise training. We next interrogated our BESt-normalized

mitochondrial proteome to gain new insights into non-stoichiometric changes in individual

mitochondrial proteins following three training phases designed to induce different effects on

mitochondrial biogenesis. Using an ANOVA, we identified 185 mitochondrial proteins that

were differentially expressed across the three training phases (Table S5). This is ~10-fold

greater than the number of proteins reported to change post-training in previous human

skeletal muscle studies (Egan et al., 2011; Holloway et al., 2009; Hostrup et al., 2018;

Robinson et al., 2017) (Figure 2D). Unsupervised hierarchical clustering of all differentially

expressed mitochondrial proteins revealed six clusters (Figure 2E) with distinct patterns of

change in response to training (Figure S2C), which also confirmed the absence of a fixed

stoichiometry between training-induced changes in mitochondrial content and mitochondrial

protein functional classes.

Enrichment analysis of differentially expressed proteins within each of the six clusters

revealed them to be enriched in, but not always exclusively contain, proteins involved in fatty

acid β-oxidation (FAO), mitochondrial translation, the TCA cycle, OXPHOS, metabolism of

amino acids, and cristae formation, respectively (Figure 2F; Table S5). Because not all

proteins belonging to each individual Reactome pathway segregated within one cluster

(Figure S2D), we compared our 185 differentially expressed proteins with proteins belonging

to the six aforementioned functional classes (Reactome R-HSA) and other related pathways;

matching proteins (Table S6) were used to generate the individual pathway heatmaps

presented in Figure 3 and to discuss our findings. In addition, because Reactome, like other

protein databases, only provides an extensive but not exhaustive list of proteins involved in

each pathway, we complemented our heatmaps with other proteins identified through

literature (see methods for more information on how pathways in Table S6 and Figure 3 were

generated).

To pinpoint the origin of training-induced changes in individual mitochondrial proteins, we

performed unpaired t-tests between group pairs following permutation-based false discovery

rate (FDR) correction. Using a conservative approach, we matched the differentially

expressed proteins identified by each t-test comparison with the 185 differentially expressed

proteins identified by the ANOVA (Table S7). The majority of changes occurred during the

initial NVT phase (Figure 2G; Table S7), where 102 proteins were differentially expressed,

suggesting that the greatest number of mitochondrial adaptations took place within the first

few training sessions and these became harder to obtain as the training progressed (Figure 2E,

note the strongest changes are found in the "PN *vs.* BL" column). Our study also confirms

that the training stimulus during the RVT phase was sufficient to prevent a loss of adaptation

(Figure 2G; Table S7).

Figure 3. Overview of training-induced changes in mitochondrial protein functional

classes and metabolic pathways

Protein functional classes and/or metabolic pathways (defined in Table S6) as determined in

Table S7. Row clustering determined by unsupervised hierarchical cluster analysis. BL:

baseline; PN: post-NVT; PH: post-HVT; PR: post-RVT; ROS: reactive oxygen species;

FAO: fatty acid β-oxidation; CoQ: Coenzyme Q; TCA: tricarboxylic acid cycle; OXPHOS:

oxidative phosphorylation.

Proteins important for the translation of mitochondrial proteins are sensitive to training

volume. We next examined the six over-represented pathways highlighted by our enrichment

analysis, beginning with the pathway related to mitochondrial protein translation (Figure 3),

which mostly segregated within cluster-2 (Figure 2F, and S2D; Table S5). Specifically, we

identified proteins from both the small 28S (MRPS5, MRPS22, MRPS27) and large 39S

(MRPL4, MRPL18) subunits of the mitochondrial ribosome, as well as proteins involved in

translation initiation (MTIF2) and elongation (GFM1, PTCD3, TSFM, TUSM) (Figure 3).

Whereas only four proteins were upregulated post-NVT, eight of the 10 aforementioned

proteins were upregulated post-HVT, with six remaining elevated post-RVT (Table S7). This

is the first study to directly demonstrate training-induced changes in these or related

mitochondrial translation proteins in young, healthy humans, as previous research reported

training-induced increases in older (65 to 80 years), but not younger (18 to 30 years),

individuals (Robinson et al., 2017). Our findings also indicate that changes in proteins

involved in mitochondrial protein translation are sensitive to training volume, consistent with

a study observing a positive association between these proteins and physical activity levels in

healthy individuals (20 to 87 years) (Ubaida-Mohien et al., 2019).

Mitochondria prioritize the TCA cycle and FAO in response to NVT. Proteins involved in

the TCA cycle pathway grouped mainly within cluster-3 and, to a lesser extent, cluster-1

(Figure 2F and S2D; Table S5). This pathway was characterized by a large and concerted

increase post-NVT (12 of 16 enzymes were significantly increased), with many of these

proteins remaining upregulated compared to BL both post-HVT and post-RVT (Figure 3,

Table S7). Of note, SDHA and SDHB, subunits of CII - the only TCA cycle enzyme that also

participates in the ETC - clustered with the majority of OXPHOS subunits (discussed below)

and decreased during NVT. While training-induced changes in some TCA cycle proteins

have been reported (Egan et al., 2011; Hostrup et al., 2018), this is the first study to reveal

training-induced increases in ACO2, SUCLG1, SUCLG2, SUCLA2, and FAHD1, and that

these enzymes adapted mostly in a coordinated fashion.

Proteins involved in FAO segregated mainly within cluster-1 and cluster-3 (Figure 2F and

S2D; Table S5). Our data demonstrated a large and concerted increase in these proteins post-

NVT, with minimal (HVT) and no (RVT) changes thereafter (Figure 3; Table S7). Proteins

increased post-NVT included chain shortening enzymes involved in FAO (ACAA2, ACSF2,

ECHS1, HADH, HSD17B10, MECR, MLYCD) and enzymes required to convert unsaturated

fatty acids into intermediates of FAO (DECR1 and HSD17B8). Similar increases were also

observed for ETFA and ETFB, which transfer electrons produced during FAO to the

OXPHOS system, further highlighting the early increase in FAO. HMGCL and ACAT1, two

enzymes involved in the ketogenesis pathway, were also increased, possibly suggesting that

excess acetyl-CoA generated by enhanced FAO may be fed into ketogenesis. Taken together,

these findings indicate that the synthesis of proteins involved in both the TCA cycle and FAO

pathways is prioritized in the early stages of an endurance training intervention; these

increases may have contributed, at least in part, to the training-induced increase in

mitochondrial respiration (Figure S1B, upper panels).

Lipidomics highlights a divergent response to training of different lipid classes. To

highlight specific changes in fatty acid composition, LC-MS/MS based comparative

lipidomics was performed using the same mitochondrial isolates used for proteomics. A total

of 779 species representing 30 lipid classes (Table S8) were quantified. Indeed, changes in

the mitochondria-specific cardiolipins (CLs), the biomarker most strongly associated with

mitochondrial content in human skeletal muscle (Larsen et al., 2012), mirrored changes in

training volume and in non-normalized mitochondrial proteins (compare Figure 4A, left panel

with Figure 2A, upper panel; Table S8). Therefore, to ensure that our lipidomics data were

corrected for changes in MPE in our fractions, we applied a similar approach to the one used

for proteomics where lipidomic results were normalized according to the trend observed for

the CL class (see methods). Post-normalization profile plots of CL species indicated that,

despite changes in individual CL species, the general increase in mitochondrial content with

different training volumes had been compensated for by our normalization strategy (Figure

4A; right panel), and the lipid profiles could be used to readily segregate samples across

different time points (Figure 4B). From this point on we will present post-normalization

lipidomics results, unless specified otherwise.

We identified 182 differentially expressed lipid species (Figure 4C; Table S9). Similar to

what we observed in our proteomics dataset, and consistent with previous research in humans

(Dube et al., 2011; Shepherd et al., 2017), our study highlighted the divergent response to

training of different lipid classes and species relative to overall changes in mitochondrial

content (Figure 4D). However, in contrast to proteomics, where the greatest number of

changes were observed early (during NVT) and no changes were observed during RVT, our

lipidomics assessment showed the most significant changes to occur during the latter two

stages (Figure S3). This suggests a delayed response to changes in lipid content compared to

proteins, and may reflect overall mitochondrial biogenesis as training-induced changes in

lipid species were mainly related to mitochondrial membrane-based lipids (CLs,

phosphatidylethanolamines and phosphatidylcholines) and the triglycerides (TGs) (Figure

S3). Contrary to previous training studies in obese humans (Bruce et al., 2006; Dube et al.,

2011; Shepherd et al., 2017), we did not observe reductions in muscle ceramides [with the

exception of Cer(d18:1/18:0) during HVT] or diacylglycerol content in our healthy, young

men following training. While this could indicate a different response of lipid species in

mitochondria isolates (our study) compared to whole-muscle lysates (Bruce et al., 2006;

Dube et al., 2011; Shepherd et al., 2017), as previously suggested (Perreault et al., 2018), it

could also highlight adaptation differences between obese and healthy individuals.

Of relevance to the increase in FAO reported above, we observed a post-NVT decrease in

TGs (Figure 4D), a class of lipids used as an important source of energy production during

exercise (Goodpaster and Kelley, 1998); however, likely due to the high number of multiple

comparisons in our bioinformatic analysis (n = 779), no significant changes in single TG

species were detected (Figure S3). The decrease in TGs in our study is consistent with an

increase in fatty acid turnover, and is corroborated by the post-NVT increase in FAO

enzymes (Figure 3). Finally, changes in TGs (Figure 4D), which are not found in

mitochondria but rather in lipid droplets (Walther and Farese Jr, 2012), mirrored changes in

PLIN5 (Figure 3). Since PLIN5 is a protein that tethers mitochondria to lipid droplets and

regulates the release of lipids for FAO (Benador et al., 2019; Wang et al., 2011), we

hypothesize that training-induced changes in TGs may reflect increased association of lipid

droplets with mitochondria. While few studies have evaluated the effects of different training

interventions on skeletal muscle lipid species, our results add to the growing evidence that

changes with training are specific to the exercise prescription and that training-induced

changes in individual lipid species and lipid classes are not stoichiometrically linked.

Figure 4. Lipidomics highlights a divergent response to training of different lipid classes

(A) All-lipids profile plot of non-normalized (left panel) and cardiolipin (CL) class-normalized (right panel) intensity values from mitochondrial isolates, displaying all individual CL species (gold) and all other lipid species (grey).

(B) Multidimensional scaling analysis showing segregation of samples obtained from IM fractions of lipidomics values normalized by the entire CL class.

(C) Heatmap of differentially expressed lipids after normalization by the entire CL class

between training phases determined with an adjusted $P < 0.05$ (Benjamini Hochberg). Row

clustering determined by unsupervised hierarchical cluster analysis.

(D) Profile plots of the scaled concentration mean of different lipid classes. For each profile

plot, all the individual lipid species identified within the specific class are represented in

color over the entire lipidome (grey) at each time point. The intensity values for all lipid

species were normalized by the entire CL class to adjust for changes in mitochondrial

content.

BL: baseline; PN: post-NVT; PH: post-HVT; PR: post-RVT. n = 10 for all analyses.

A training-induced increase in OXPHOS subunits is not a priority to enhance

mitochondrial respiration. We next examined the OXPHOS pathway, which segregated

almost entirely in cluster-4 (Figure 2F and S2D; Table S5). Our analysis provided excellent

coverage of OXPHOS proteins and identified 78 OXPHOS subunits. We report training-

induced changes in 26 OXPHOS subunits (Figure 3), equivalent to 33% of all subunits

identified, most for the first time. Differentially expressed OXPHOS proteins included

subunits from all six of the modules required for the assembly of CI, both catalytic subunits

of CII, as well as subunits from CIII, one of the three core subunits of CIV, and proteins

found in both functional domains of CV. Specifically, we observed a large and coordinated

(23 subunits) decrease post-NVT, with no changes during HVT and/or RVT (Figure 3; Table

S7). Changes in SLC25A4 (ANT1), a protein catalyzing the exchange of cytosolic ADP and

mitochondrial ATP across the mitochondrial inner membrane (Palmieri, 2013), mirrored

these changes (Table S7). The post-NVT decrease in OXPHOS subunits indicates that during

times of increased mitochondrial biogenesis, synthesis of proteins involved in OXPHOS

pathways is deprioritized. Moreover, our results demonstrated a large post-NVT increase in

TCA cycle and FAO related enzymes - the major providers of reducing equivalents (NADH

and $FADH_2$) to the OXPHOS system (Wang and Hekimi, 2016); this suggests that

enhancement of these two metabolic pathways may be more important to improve

mitochondrial respiration following exercise training (Figure S1B, upper panels) than an

increase in OXPHOS subunits.

An increase in reducing equivalents coupled with the de-prioritization of the OXPHOS

machinery could lead to increased reactive oxygen species (ROS) generation. Here we report

an immediate post-NVT increase in the abundance of enzymes involved in protection from

oxidative stress, such as PRDX5 (the most increased protein), TXN2, and GPX1 (Powers et

al., 2020) (Figure 3; Table S7). Moreover, both SOD2, the mitochondrial ROS scavenger

(Powers et al., 2020), and PRDX5 were increased post-HVT and post-RVT. These findings

suggest that training-induced synthesis of proteins involved in the protection from oxidative

stress was emphasized early during the training intervention and maintained throughout,

consistent with the notion that exercise training provides protection against oxidative stress

(LaHue et al., 2016; Powers et al., 2020).

The coordinated assembly of OXPHOS complexes involves various assembly factors,

chaperones, and protein translocation components, and has been suggested to be in the order

of days (McKenzie et al., 2007). Whereas increases in OXPHOS assembly factors occurred

mostly post-HVT (Figure 3; Table S7), changes in subunits of the mitochondrial ribosome

(Figure 2C, lower right panel), which synthesize the 13 OXPHOS subunits encoded by

mitochondrial DNA, and in several chaperone proteins regulating import and folding of

proteins destined for the mitochondrial matrix, took place earlier (post-NVT; Figure 3; Table

S7). Chaperone proteins upregulated post-NVT included HSPA9 (mtHsp70), HSPD1

(mtHsp60), HSPE1 (mtHsp10), and TRAP1 (mtHsp75), whereby TIMM44 was increased

during HVT (Figure 3; Table S7). All five proteins remained elevated compared to BL both

post-HVT and post-RVT. Except for HSPE1 (Egan et al., 2011), none of the above

chaperones have previously been reported to respond to exercise. Owing to the notion that all

of these chaperones are known to interact (Voos, 2013), our data indicate a coordinated

increase in the protein quality control system in response to exercise training. While we

detected many subunits of the translocases of the outer and inner membrane (TOM and TIM,

respectively; Table S3), which are required for importing nuclear-encoded subunits

(Wiedemann and Pfanner, 2017), most did not change significantly with training, suggesting

that their levels remain stable relative to changes in mitochondrial content.

Training-induced changes in mitochondrial content, but not changes in the abundance

and/or organization of SCs, mediate improvements in mitochondrial respiration. A

previous study reported that training increases the formation of ETC SCs - high molecular

weight assemblies comprised of CI, CIII, and CIV (Schägger and Pfeiffer, 2000) - and

suggested this to contribute to improvements in mitochondrial respiration (Greggio et al.,

2017). Although SCs were originally proposed to support enhanced ETC function (Lapuente-

Brun et al., 2013), their role has become controversial, with multiple groups reporting that

they confer no bioenergetic advantage (Hirst, 2018). To provide further insight, we

performed BN-PAGE analysis of ETC SCs (Figure 5A). We observed increases in SC

abundance with increasing training volumes, achieving significance post-HVT (Figure 5B,

upper panels); however, these increases were no longer significant upon normalization by

MPE (Figure 5B, middle panels).

We next assessed whether exercise training affects the distribution of ETC complexes into

SCs. In contrast to the only previous report showing a post-training redistribution of CIII and

CIV (but not CI) into SCs in elderly individuals (Greggio et al., 2017), our analysis revealed

no significant changes throughout the intervention (Figure 5B, lower panels). This

demonstrates that in young healthy men, endurance training does not alter the ETC complex

distribution within major and minor SC assemblies. While the contrasting findings may relate

to differences in the participants' average age (66 *vs.* 22 y) and baseline $\dot{V}O_{2max}$ (1.9 *vs.* 3.6 L

min^{-1}), as well as the training interventions (48 sessions of moderate-intensity training in 16

weeks *vs.* 52 sessions of HIIT in 9 weeks), our results are consistent with the concept that SC

assemblies are a structural feature of the ETC rather than a phenomenon conferring enhanced

bioenergetics (Hirst, 2018). In line with this, our data indicated that the greatest changes in

maximal mitochondrial respiration ([ETF+CI+II]$_P$) occurred in parallel with the greatest

changes in SC abundance (compare Figure 5B with 5C, both upper panels). However,

following normalization by MPE, both of these adaptations were no longer significant

(compare Figure 5B, middle panels with Figure 5C, lower panel). Taken together, our

findings do not support the hypothesis that training-induced changes in the abundance or

organization of SCs contribute to improvements in mitochondrial respiration. Conversely, our

data indicate that training-induced changes in both mitochondrial respiration and SC

formation could largely be attributed to increases in mitochondrial content.

Given the importance of CLs for the assembly and stability of mitochondrial membrane

protein complexes and ETC SCs (Gebert et al., 2009; Pfeiffer et al., 2003), we again

interrogated our normalized mitochondrial lipidome. Our analysis demonstrated an extensive

remodeling of CL composition (Figure 5D); notably, following NVT and HVT, we observed

a decrease in the abundance of tetra-linoleoyl CL (18:2_18:2_18:2_18:2), the dominant form

of CLs found in skeletal muscle (Ikon and Ryan, 2017). Accompanying this were

concomitant increases in CLs containing oleic acid (18:1) and to a lesser extent palmitoleic

acid (16:1) acyl chains. This may reflect a lack in relative availability of dietary linoleic acid

due to increased mitochondrial biogenesis, since the nature of CL remodeling catalyzed by

TAZ, which was unaltered in our dataset, appears to be controlled predominantly by lipid

availability (Malhotra et al., 2009). This compensation in CL content is in line with our

observation that increased training volume does not lead to changes in ETC SC assembly

(Figure 5B middle and lower panels).

Figure 5. Training-induced changes in mitochondrial content, but not changes in the

abundance and/or organization of SCs, mediate improvements in mitochondrial

respiration

(A) Representative BN-PAGE blots of isolated mitochondria (IM) fractions from human

vastus lateralis muscles (images from baseline [BL] IM fractions). Band 1 (SCs): mature

supercomplexes (SCs) consisting of complex $I+III_n+IV_n$; band 2 ($CIII_2+IV$): a SC consisting

of CIII and CIV; band 3 (LMW CI int): low molecular weight intermediate of CI (band not

present in all samples); band 4 ($CIII_2$): CIII dimer; band 5 (CIV_2): CIV dimer; band 6 (CIV):

CIV monomer.

(B) Top panels: protein content of SCs of the electron transport chain (ETC) by BN-PAGE in

IM fractions from human vastus lateralis muscle biopsies (antibodies probed on separate

membranes); middle panels: values from top panels normalized by mitochondrial protein

enrichment (MPE); lower panels: distribution of ETC complexes into SCs from images

obtained in b at each time point.

(C) Top panel: maximal mass-specific mitochondrial respiration (MR) in permeabilized

human vastus lateralis muscle fibers with convergent electron input through ETF+CI+CII

($[ETF+CI+II]_P$); lower panel: values of mitochondrial- (mt-) specific MR, obtained after

normalizing values from the top panel by MPE ($[ETF+CI+II]_P/MPE$). Results for the entire

substrate-uncoupler-inhibitor-titration (SUIT) protocol are shown in Figure S1B.

(D) Training-induced changes in all individual CL species that were differentially expressed

(as determined in Figure 4C, with an adjusted $P < 0.05$ [Benjamini Hochberg]).

PN: post-NVT; PH: post-HVT; PR: post-RVT; IB: immunoblotting. ▼,◊,○, and ● represent

individual values; bars represent mean values; error bars in c represent SD; n = 10 for all

analyses; datasets (B) and (C) were analyzed by repeated measures one-way ANOVA

followed by Tukey's post hoc testing, except for the lower right panel of (B), which was

analyzed by Friedman test followed by Dunn's post hoc testing, as not normally distributed.

Significance: $P < 0.05$.

Coenzyme Q biosynthesis and cristae formation are not prioritized during high-volume

endurance training. Despite enrichment analysis suggesting cluster-5 to be enriched in

metabolism of amino acids and derivatives, upon further inspection we identified this cluster

to predominantly contain proteins involved in the biosynthesis and function of coenzyme Q

(CoQ or ubiquinone; Figure 2F and S2D; Table S5) - an enzyme with an essential role as an

electron transfer lipid in OXPHOS processes (Wang and Hekimi, 2019). We identified

changes in four constituents (COQ3, COQ5, COQ7, COQ9) of the CoQ synthome (Figure 3;

Table S6), a multi-subunit complex necessary for the biosynthesis of CoQ (Stefely and

Pagliarini, 2017). We report no changes post-NVT, consistent with previous research

investigating the effects of short-duration (< 3 weeks) exercise training on CoQ content in

human skeletal muscle (Svensson et al., 1999). However, post-HVT we report a concerted

reduction in three of these proteins, which also persisted post-RVT (Table S7). This indicates

that CoQ biosynthesis is not prioritized in response to high-volume endurance training,

suggesting that enhancing the CoQ pool may not be necessary to support an increase in

mitochondrial respiration in human skeletal muscle of young healthy individuals.

CoQ is an electron carrier shuttling electrons derived from the TCA cycle (via CI and CII),

pyrimidine biosynthesis (via DHODH), glycolysis (via GPD2), and FAO (via EFTDH) to

Complex III (Wang and Hekimi, 2016). Moreover, other enzymes located near or on the

inner mitochondrial membrane feeding electrons to CoQ, such as IVD (Finocchiaro et al.,

1987), DLD (Xia et al., 2001), and L2HGDH (Oldham et al., 2015), were also identified in

our study (Table S6). Similar to the decrease reported above in CI and CII subunits post-

NVT, we report decreases in four of these enzymes during HVT, with three of them

remaining downregulated both post-HVT and post-RVT; conversely, IVD was increased

during NVT and DHODH was elevated post-HVT (Figure 3; Table S7). The lack of a clear

correlation between the content of the above proteins and that of CoQ may be ascribed to

changing preferences for energy utilization over the course of the training intervention, as

also indicated by the differential prioritization of the various metabolic pathways described

above.

The mitochondrial contact site and cristae organizing system (MICOS) is pivotal for the

formation of cristae junctions, providing the extended membrane surface hosting OXPHOS

complexes (Huynen et al., 2016). The MICOS associates with the outer membrane sorting

and assembly machinery (SAM) to yield the mitochondrial intermembrane space bridging

complex (MIB), thus linking the mitochondrial inner and outer membranes (Huynen et al.,

2016). Eight proteins involved in cristae formation were differentially expressed in our study:

APOO (MIC26), APOOL (MIC27), CHCHD3 (MIC19), and MIC60 from the MICOS

complex; MTX1 and MTX2 from the SAM complex; TMEM11, a protein associating with

multiple MICOS subunits for cristae biogenesis (Guarani et al., 2015); and OMA1, a

mitochondrial protease regulating OPA1 (Ehses et al., 2009) - a key protein implicated in

cristae remodeling (Frezza et al., 2007) (Figure 3; Table S6). Seven of these proteins, which

almost entirely grouped within cluster-6 (Figure 2F and S2D; Table S5), were significantly

decreased post-HVT, with four being decreased post-RVT (Table S7). To our knowledge,

this is the first study to report a training-induced de-prioritization of these proteins. This is

consistent with the minimal changes observed in cristae density following short-term (< 3

months) exercise training in obese individuals (Nielsen et al., 2017). Thus, our results support

the notion that cristae remodeling is not a priority during short-term training and that changes

in other mitochondrial proteins are more important to adapt to the higher metabolic demand

of exercise. Despite MICOS-MIB subunits being known to interact with CLs (Weber et al.,

2013), we report no correlation between the levels of MICOS-MIB subunits and changes in

CLs content following different training volumes.

CONCLUSION

The present study utilized the power of a multi-omics approach, and a novel *in silico*

normalization strategy eliminating the bias introduced by changes in mitochondrial content,

to unearth an intricate and previously undemonstrated network of differentially prioritized

mitochondrial adaptations to exercise training in men. We identified 185 differentially

expressed mitochondrial proteins, an ~10-fold increase compared to previous studies. This

increased number of proteins was largely related to our normalization strategy, and does not

seem to depend on the large training volume utilized (120 differentially expressed proteins

would have been identified after the modest training volume of the NVT phase alone), nor is

the result of comparing multiple time points (118 differentially expressed proteins would

have been identified by sampling muscle only at the beginning and end of the study). This

increased depth of analysis enabled us to identify the time-dependent and complex

remodeling of the mitochondrial proteome following endurance exercise. We demonstrate

that training-induced changes in individual proteins, protein functional classes, and metabolic

pathways follow markedly different patterns of adaptation, which are not stoichiometrically

linked to the overall training-induced increase in mitochondrial content. Moreover, we show

that the lack of stoichiometry and the differential prioritization extended to all three levels

investigated (i.e., the transcriptome, proteome, and lipidome) and between levels, as

demonstrated by the delayed adaptive response of lipids compared to proteins.

A striking and unexpected finding was the early (post-NVT) deprioritization of OXPHOS

subunit formation, which indicates that the overall training-induced increase in biogenesis

and the proliferation of mitochondria is greater than the increase in specific components of

the OXPHOS machinery. This become even more striking when considering the early

increase in mitochondrial respiration, as well as proteins involved in the TCA cycle and FAO

processes - two of the main suppliers of reducing equivalents to the OXPHOS system. This

could not be explained by changes in the abundance and/or organization of SCs, as

previously hypothesized (Lapuente-Brun et al., 2013), as our findings showed no change in

these parameters, adding to the growing evidence that SC formation does not confer

enhancements in mitochondrial bioenergetics (Hirst, 2018). This is also unlikely to be related

to the lengthy time requirements necessary to assemble large multi-protein complexes

(McKenzie et al., 2007), as the relative amount of OXPHOS subunits remained reduced

throughout the intervention (lasting several weeks), whereas mitochondrial respiration, as

well as TCA cycle and FAO process, remained upregulated throughout. Instead, these

increases were likely supported by the inherent reserve capacity of mitochondria (i.e., their

ability to respond to sudden increases in energy requirements) (Desler et al., 2012), and

demonstrate that enhancing electron flow to OXPHOS is more important to increase ATP

production than an increase in the components of the OXPHOS machinery. The relative

decrease in proteins important for both CoQ biosynthesis and cristae formation further

substantiates these findings.

Finally, the above findings were the result of changes in only one of the several exercise

prescription programming variables (i.e., training volume). Future research should investigate

the effects of manipulating other programming variables (e.g., exercise intensity, frequency,

recovery between sessions), the type of exercise (e.g., running, swimming, resistance

training), and different populations (e.g., differences in age, sex, health status) utilizing a

similar approach. These interventions are likely to induce adaptations that are specific and

presumably different from the one presented here. Because of the well-documented

therapeutic benefits of exercise training (Booth et al., 2000; Pedersen and Saltin, 2006), the

knowledge generated by our novel findings, which are readily available in supplemental

tables, together with the results from future research, could be mined by exercise and health

professionals to design focused and personalized training interventions.

Acknowledgments. We thank the participants for their time and commitment to this study.

The authors would like to acknowledge all members of the Stroud and Bishop labs for input

into interpretation and presentation of data. We thank the Bio21 Mass Spectrometry and

Proteomics Facility (MMSPF) for provision of instrumentation, training, and technical

support. We acknowledge the use of the services and facilities of the Australian Genome

Research Facility (AGRF). This study was funded by grants from the ANZ-MASON

Foundation (to D.J.B), the Australian Research Council (Discovery Project DP140104165 to

D.J.B), the Australian National Health and Medical Research Council (NHMRC Project

Grant 1140906 to D.A.S.; NHMRC Fellowships 1140851 to D.A.S. and 1155244 to D.R.T.),

JDRF Australia (JDRF Career Development to M.T.C.), and the Australian Research Council

Special Research Initiative in Type 1 Juvenile Diabetes (to M.T.C.). We acknowledge the

support of the Mito Foundation for the provision of instrumentation and the Victorian

Government's Operational Infrastructure Support Program.

Author contributions. C.G., D.A.S., and D.J.B. conceptualized the study. C.G., D.A.S., and

D.J.B. devised the study methodology. C.G., N.A.J., and H.A.J. performed study validation.

C.G., N.A.J, H.A.J. delivered the training and performed sample collection. C.G., J.B., J.K.,

B.R., A.L., T.L.S., and A.E.F. performed biochemical analyses. C.G., B.R., and D.A.S.,

performed proteomic analysis. K.H., and N.A.M. performed lipidomic analysis. RNA-seq

analysis was performed at the Australian Genome Research Facility (AGRF). C.G., N.J.C.,

and D.A.S. performed statistical and bioinformatic analysis. C.G., N.J.C., and D.A.S.

delivered the visualization. C.G., D.A.S., and D.J.B. wrote the manuscript. M.T.C., P.J.M.,

D.R.T., D.A.S, and D.J.B. provided supervision. D.R.T., M.T.C., D.A.S, and D.J.B. funded

the research. C.G., D.A.S., and D.J.B. have primary responsibility for final content. Data

collection took place at Victoria University. Muscle analysis took place at Victoria

University, Monash University, Murdoch Children's Research Institute, the Baker Heart &

Declaration of interests. The authors declare no conflict of interest.

STAR METHODS

RESOURCE AVAILABILITY

Lead Contact and Materials Availability. Further information and requests for resources

and reagents should be directed to and will be fulfilled by the lead contacts, David Stroud

(david.stroud@unimelb.edu.au) and David Bishop (david.bishop@vu.edu.au). This study did

not generate new unique reagents.

Data and Code Availability. The mass spectrometry proteomics data will be deposited to the

ProteomeXchange Consortium via the PRIDE partner repository upon publication. The

transcriptomic data will be deposited to NCBI upon publication. Lipidomics data will be

deposited to the Metabolomics Workbench Data Repository. The R scripts used for all omics

analyses described above are deposited on GitHub at

https://github.com/XercisOmics/EnduranceMitoMultiOmics.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Ten healthy men volunteered to take part in this study (physiological and performance

parameters are presented in Table 1). Potential participants were deemed suitable if aged 18-

35 y, were moderately-trained (i.e., less than 4 h per week of unstructured aerobic activity for

half a year prior to the study), not regularly engaged in cycling-based sports, and were non-

smokers and medication free prior to and during the study. Participants underwent a medical

screening to exclude conditions that may have precluded their participation (e.g.,

cardiovascular, musculoskeletal and/or metabolic problems), and were informed of the study

requirements, risks, and benefits, before giving written informed consent. Approval for the

study's procedures, which conformed to the standards set by the latest revision of the

Declaration of Helsinki, was granted by the Victoria University Human Research Ethics

Committee (HRE15-126).

METHOD DETAILS

Study design. The study consisted of three consecutive training phases: the normal- (NVT),

high- (HVT), and reduced- (RVT) training volume phase (Figure 1A). Each training phase

was preceded (and followed) by performance testing, which included a 20-km cycling time

trial (20k-TT), a graded exercise test (GXT) (participants were previously familiarized with

both tests), and a resting muscle biopsy. Overall study duration was ~9 weeks.

Testing procedures. Participants were required to avoid any vigorous exercise for the 48 h

preceding each performance test (72 h for the skeletal muscle biopsy), from alcohol and any

exercise for 24 h before testing, and from food and caffeine consumption for the 2 h

preceding each test. Similar tests were performed at the same time of the day throughout the

study to avoid variations caused by changes in circadian rhythm.

GXT. A graded exercise test was performed on an electronically-braked cycle ergometer

(Lode Excalibur v2.0, Groningen, The Netherlands) to determine peak oxygen uptake

($\dot{V}O_{2Peak}$), peak power output ($\dot{W}_{Peak}$), the power attained at the lactate threshold ($\dot{W}_{LT}$) using

the modified D_{Max} method (Bishop et al., 2000), and the training intensity for each training

phase. The test consisted of consecutive 4-min stages at constant power output; the test

starting intensity (range: 45-77 W) and the intensity increase of each stage (range: 17-28 W)

were chosen so as to obtain at least 8 time points for the determination of the $\dot{W}_{LT}$ (Jamnick et

al., 2018) and were based on participants' fitness levels. An identical protocol was used at all

4 time points for each participant. Prior to the test, and in the last 30 s of each stage, venous

blood samples were taken for measurement of blood lactate concentration ([La$^-$]). Participants

were instructed to keep a cadence > 60 rpm and were only allowed access to cadence. The

test was stopped when a participant reached volitional exhaustion or cadence dropped below

60 rpm for over 10 s. The $\dot{W}_{Peak}$ was determined as the power of the last completed stage plus

an additional 25% of the stage increase wattage for every additional minute completed. At the

end of each GXT, after a 5-min recovery, a verification exhaustive bout was performed at an

intensity equivalent to $\dot{V}O_{2Peak}$ to confirm the highest measured $\dot{V}O_{2Peak}$ (Jamnick et al.,

2018).

20k-TT. Cycling time trials were performed on an electronically-braked cycle ergometer

(Velotron, RacerMate, Seattle, WA, USA) after a 6-min cycling warm-up (4 min at 66% of

$\dot{W}_{LT}$ followed by 2 min at $\dot{W}_{LT}$), and 2 min of rest. Participants were only allowed access to

cadence and completed distance.

Gas Analysis during the GXT. During the GXT, expired air was continuously analyzed for O_2

and CO_2 concentrations via a gas analyzer (Moxus 2010, AEI Technologies, Pittsburgh, PA,

USA), which was calibrated immediately before each test. $\dot{V}O_2$ values were recorded every

15 s and the average of the two highest consecutive 15-s values was recorded as a

participant's $\dot{V}O_{2Peak}$.

Venous blood sampling. Venous blood samples (~1 mL) were collected during the GXT from

a cannula inserted in the antecubital vain for the determination of venous blood [La^-] using a

blood-lactate analyzer (2300 STAT Plus; YSI, Yellow Spring, OH, USA).

Muscle biopsies. A biopsy needle with suction under local anesthesia (1% xylocaine) was

used to obtain vastus lateralis muscle biopsies at rest at the following four time points: BL,

PN, PH and PR. After being cleaned of excess blood, connective and fat tissue muscle

biopsies were divided as follows: ~10 mg was immediately immersed in ~2 mL of ice-cold

BIOPS for measurements of mitochondrial respiration, whereas the remainder was promptly

frozen in liquid nitrogen and stored at -80°C for follow-up analyses.

Training intervention. All training sessions were performed on an electronically braked

cycle ergometer (Velotron, RacerMate, USA), following an 8-min warm up (see 20k-TT) and

consisted of HIIT (2:1 work-to-rest ratio). Training intensity was set relative to $\dot{W}_{LT}$ (rather

than $\dot{W}_{Peak}$) so as to induce similar metabolic and cardiac stresses amongst participants of

differing fitness levels (Baldwin et al., 2000). Exercise intensity was maintained between

$\dot{W}_{LT}$ and $\dot{W}_{Peak}$ throughout the entire study so that training volume was the only manipulated

variable between the 3 phases.

NVT phase. This consisted of 6 HIIT sessions within 2 weeks of 5 to 7 4-min cycling

intervals interspersed with a 2-min recovery at 60 W. Exercise intensities were defined as

$[\dot{W}_{LT} + x(\dot{W}_{Peak}-\dot{W}_{LT})]$, with x increasing from 0.5 to 0.7 throughout the phase.

HVT phase. Participants performed HIIT twice a day for 20 consecutive days; training

sessions consisted of either 7 to 10 4-min intervals interspersed with a 2-min recovery at 60

W at intensities ranging from $[\dot{W}_{LT} + 0.5(\dot{W}_{Peak}-\dot{W}_{LT})]$ to $[\dot{W}_{LT} + 0.8(\dot{W}_{Peak}-\dot{W}_{LT})]$, or 15 to 20

2-min intervals at intensities ranging from $[\dot{W}_{LT} + 0.5(\dot{W}_{Peak}-\dot{W}_{LT})]$ to $[\dot{W}_{LT} + 0.95(\dot{W}_{Peak}-$

$\dot{W}_{LT})]$, interspersed with a 1-min recovery at 60 W. Single session duration increased from

~45 min to 60 min.

RVT phase. The RVT phase consisted of 6 HIIT sessions in 6 days; participants performed

10, 9, 8, 7, 6, and 4, 4-min intervals interspersed with a 2-min recovery at 60 W, at an

intensity of $[\dot{W}_{LT} + x(\dot{W}_{Peak}- \dot{W}_{LT})]$, with x increasing from 0.5 to 0.7 throughout the phase.

Physical activity and nutritional control. Physical activity and dietary patterns were

maintained throughout the study and were monitored with the use of food and physical

activity recall diaries. The last 3 meals prior to each performance test undertaken during

baseline testing were recorded by each participant and were replicated thereafter before the

same type of test. To control for dietary effects on muscle metabolism, participants were

provided with a standardized dinner (55 kJ kg^{-1} body mass (BM), providing 2.1 g

carbohydrate (CHO) kg^{-1} BM, 0.3 g fat kg^{-1} BM, and 0.6 g protein kg^{-1} BM) and breakfast

(41 kJ kg^{-1} BM, providing 1.8 g CHO kg^{-1} BM, 0.2 g fat kg^{-1} BM, and 0.3 g protein kg^{-1} BM),

to be consumed 15 and 3 h prior to the muscle biopsy, respectively.

Muscle analyses

Enzymatic activity. Enzyme activities were determined spectrophotometrically in post-600g

supernatants of skeletal muscle homogenates according to the method described for

respiratory chain complexes I-IV and citrate synthase (Frazier et al., 2020). Briefly, ~20mg of

skeletal muscle sample was homogenized in sucrose/mannitol containing buffer using a

glass/glass homogenizer and spun for 10 min at 600 g and 4 °C. The supernatant was then

subjected to two freeze/thaw cycles and stored at -80 °C until measurement of respiratory

chain enzymes. CI was assayed as rotenone-sensitive NADH:CoQ1 oxidoreductase by

monitoring the decrease in absorbance due to NADH oxidation at 340 nm. For CII, activity

was measured as succinate:CoQ1 oxidoreductase by measuring CoQ1 reduction at 280 nm.

CIII was assayed as decylbenzylquinol; cytochrome *c* oxidoreductase by following the

increase in absorbance resulting from cytochrome *c* reduction at 550 nm. CIV was measured

as cytochrome *c* oxidase by following the decrease in absorbance resulting from cytochrome

c oxidation at 550 nm. To assay the CS catalyzed production of coenzyme A (CoA.SH) from

oxaloacetate, the generation of free sulfhydryl groups was monitored using the thiol reagent

5,5'-dithio-bis-(2-nitobenzoic acid) (DTNB), which reacts spontaneously with the sulfhydryl

groups to produce 5-thio-2-nitrobenzoate anions. CS specific activity was measured by

following the increase in absorbance resulting from the formation of 5-thio-2 nitrobenzoate

anions at 412nm. Following the enzyme measurements, the amount of protein in each sample

was determined using a bicinchoninic acid assay and activity calculated as initial rates

(complexes I, II and citrate synthase) or as first-order rate constants (complexes III and IV).

Preparation of whole-muscle lysates for SDS-PAGE assessment of ETC subunits. Frozen

skeletal muscle samples (~10 mg) were homogenized in a TissueLyser for 2x 2 min at

maximum speed in ice-cold lysis buffer (1:20 w/v) containing 50 mM Tris-HCl, 150 mM

NaCl, 1 mM EDTA, 1% NP-40 and a phosphatase/protease inhibitor (5872, Cell Signaling

Technology, Danvers, MA, USA). Homogenates were rotated end-over-end at 4 °C for 1 h

and protein concentration was determined in triplicate using a commercial colorimetric assay

(Bio-Rad Protein Assay kit-II, Australia).

Mitochondrial isolation for SDS- and BN-PAGE assessment of ETC subunits, complexes and

SCs. Frozen skeletal muscle samples (~30 mg) were homogenized with 2x 20 strokes in a

Potter-Elvehjem tissue grinder attached to a rotating drill (~1000 rpm) in 5 mL solution A (1

mM EDTA, 220 mM mannitol, 20 mM HEPES-KOH [pH=7.6], 70 mM sucrose, 2 mg/mL

BSA, 0.5 mM PMSF) and spun at 800 g for 5 min at 4°C. The supernatant was collected,

whereas the pellet was re-homogenized as above in 5 mL of solution A to maximize

extraction. The two supernatants were mixed and further spun at 800 g for 5 min at 4°C. The

ensuing supernatant was then spun at 10,000 g for 20 min at 4°C, and the pellet was

resuspended in 200 µL of sucrose buffer (0.5 M sucrose, 10 mM HEPES-KOH [pH=7.6], 0.5

mM PMSF). Protein concentration was determined by the bicinchoninic acid method

according to the manufacturer's instructions (BCA Protein Assay Kit, Pierce-Thermo Fisher

Scientific, Melbourne, Australia). This measurement was used to generate Figure 1C.

SDS-PAGE. Both whole-muscle lysates (7.5 μg) and mitochondrial isolates (5 μg) were

separated by electrophoresis using 12 or 15% SDS-PAGE gels, as previously described

(Granata et al., 2016b), and blotted with a total OXPHOS (ab110411, Abcam, Cambridge,

MA, USA) or with a single CI (ab110242, Abcam, Cambridge, MA, USA) antibody where

separation with CIV was not optimal using the total OXPHOS antibody.

BN-PAGE. Mitochondrial isolates (6-15 μg) were separated by electrophoresis using 3-12%

NativePAGE gels (Life Technologies Australia, Mulgrave, Australia) as previously described

(Jha et al., 2016). A 4 g/g digitonin/protein ratio was used for assessment of SCs. The

following primary antibodies were used: NADH:ubiquinone oxidoreductase subunit A9

(NDUFA9; ab14713), ubiquinol-cytochrome *c* reductase core protein 2 (UQCRC2; ab14745),

cytochrome *c* oxidase subunit IV (COX IV; ab14744) (all Abcam, Cambridge, MA, USA).

For both SDS- and BN-PAGE, protein bands were visualized using a Bio-Rad ChemiDoc

imaging system and bands were quantified using Bio-Rad Image Lab 5.0 software (Bio-Rad

laboratories, Gladesville, NSW, Australia). An internal standard (made of a mixture of all

samples) was loaded in each SDS- and BN-PAGE gel, and each lane was normalized to this

value, to reduce gel-to-gel variability.

Fiber preparation for high-resolution respirometry. Fresh muscle fibers were mechanically

separated in ice-cold BIOPS (in mM: 2.77 CaK$_2$EGTA, 7.23 K$_2$EGTA, 5.77 Na$_2$ATP, 6.56

MgCl$_2$, 20 taurine, 50 MES, 15 Na$_2$phosphocreatine, 20 imidazole and 0.5 dithiothreitol

adjusted to pH 7.1 (Pesta and Gnaiger, 2012), followed by permeabilization by gentle

agitation for 30 min at 4°C in BIOPS containing 50 μg/mL of saponin, and 3 5-min washes in

MiR05 (in mM, unless specified: 0.5 EGTA, 3 MgCl$_2$, 60 K-lactobionate, 20 taurine, 10

KH_2PO_4, 20 HEPES, 110 sucrose and 1 g/L BSA essentially fatty acid-free, pH 7.1) (Pesta

and Gnaiger, 2012). Mitochondrial respiration was measured in duplicate (from 2-3 mg wet

weight of muscle fibers) in MiR05 at 37°C using the high-resolution Oxygraph-2k (Oroboros,

Innsbruck, Austria). To avoid potential oxygen diffusion limitation, oxygen concentration

was maintained between 270 and 480 nmol mL^{-1} by re-oxygenation via direct syringe

injection of O_2.

Mitochondrial respiration protocol. The substrate-uncoupler-inhibitor titration (SUIT)

protocol (Pesta and Gnaiger, 2012) used was as follows: 0.2 mM octanoylcarnitine and 2 mM

malate ([ETF]$_L$: leak respiration state [L] in the absence of adenylates and limitation of flux

by electron input through electron transfer flavoprotein [ETF]); 3 mM $MgCl_2$ and 5 mM ADP

([ETF]$_P$: maximal OXPHOS capacity [P] with saturating levels of ADP and limitation of flux

by electron input through ETF); 5 mM pyruvate ([ETF+CI]$_P$: P with saturating levels of ADP

and limitation of flux by convergent electron input through ETF + CI); 10 mM succinate

([ETF+CI+II]$_P$: P with saturating levels of ADP and limitation of flux by convergent electron

input through ETF + CI + CII); 10 μM cytochrome *c* (outer mitochondrial membrane

integrity test); 0.75-1.5 μM carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone (FCCP)

via stepwise titration ([ETF+C+II]$_E$, maximal electron transport chain capacity [E] with

saturating levels of ADP and limitation of flux by convergent electron input through ETF +

CI + CII); 0.5 μM rotenone ([CII]$_E$: E with saturating levels of ADP and limitation of flux by

electron input through CII); 5 μM antimycin A (residual non-mitochondrial oxygen

consumption [ROX]). Data are presented as mass-specific mitochondrial respiration [pmol O_2

s^{-1} mg^{-1} wet weight] and as mitochondrial-specific respiration [pmol O_2 s^{-1} mg^{-1} wet

weight/CS activity].

RNA-seq analysis. Approximately 10 to 15 mg of frozen muscle was used to isolate RNA

using the RNeasy Mini Kit (Qiagen, Canada) according to the manufacturer's instructions.

Samples were homogenized using the TissueLyser II (Qiagen, Canada). RNA concentration

and purity were determined spectrophotometrically (NanoDrop 2000, Thermo Fisher

Scientific, Wilmington, DE, USA) at 260 and 280 nm. RNA integrity was assessed using an

Agilent Bioanalyzer according to manufacturer's instructions. The RNA was stored at -80 °C.

Sequencing and assembly of RNA-seq. This analysis was conducted on n = 5 participants (20

samples in total); samples were sequenced (100 base pair, single reads) on the Illumina

NovaSeq 6000 platform at the Australian Genome Research Facility (AGRF). Transcriptome

assembly was completed at AGRF with reads screened for presence of any Illumina

adapter/overrepresented sequences and cross-species contamination. Per base sequence

quality for all samples was >96% bases above Q30. Cleaned sequence reads were aligned

against the *Homo sapiens* genome (Build version HG38). The STAR aligner (v2.5.3a) was

used to map reads to the genomic sequences. Count of read mapping to each known gene

were summarized to provide the matrix used for further analysis.

Bioinformatic analysis of RNA-seq data. For downstream RNA-seq analysis the R package

limma (Ritchie et al., 2015) was used to conduct the differential expression analysis from

count data. Count data was normalized using calcNormFactors in the *edgeR* package in R.

Differential expression analysis was performed between each subsequent biopsy to show

comparative changes between each of the training volumes, though all comparisons were

accounted for within the analysis. The resulting differential expression values were filtered

for an adjusted *P* value < 0.05 using the Benjamini Hochberg method. Heatmaps were

visualized using hierarchical clustering using the "average" method. Categorical columns for

the generation of Figure 1F were determined by identifying the transcripts to known

descriptions in the literature; MICOS (Eramo et al., 2020), TCA cycle (Martínez-Reyes and

Chandel, 2020), SLC25A (Palmieri, 2013), Assembly Factors (Hock et al., 2020). All

differentially expressed transcripts were run through Enrichr

[https://maayanlab.cloud/Enrichr/ (Chen et al., 2013; Kuleshov et al., 2016)] and the top

biological processes were identified with a combined score sorting for Figure 1G.

Mitochondrial isolation for proteomics and lipidomics assessment. Frozen skeletal muscle

samples (~30 mg) were homogenized with 2x 20 strokes in a Potter-Elvehjem tissue grinder

attached to a rotating drill (~1000 rpm) in 3 mL solution B (1 mM EDTA, 220 mM mannitol,

20 mM HEPES-KOH [pH=7.6], 70 mM sucrose, 0.5 mM PMSF) and spun at 1000 g for 5

min at 4°C. The supernatant was further spun at 12000 g for 10 min at 4°C, and the ensuing

pellet was resuspended in 200 μL of sucrose buffer (0.5 M sucrose, 10 mM HEPES-KOH

[pH=7.6]). Protein concentration was determined by the bicinchoninic acid method according

to the manufacturer's instructions (BCA Protein Assay Kit, Pierce-Thermo Fisher Scientific,

Melbourne, Australia). Two IM fractions per sample (50 μg each) were spun at 12000 g for

10 min at 4°C; after removal of the supernatant, the pellets were stored frozen at -80°C for

subsequent proteomics and lipidomics analysis.

Proteomics. IM fractions were prepared for proteomics analysis as previously described

(Stroud et al., 2016) with minor modifications. Briefly, 50 μg of frozen mitochondrial isolates

were solubilized in 20 μL of 8 M urea, 40 mM chloroacetamide, 10 mM tris(2-

carboxyethyl)phosphine, 100 mM Tris, pH 8.1; this was followed by 15 min of sonication in

a water bath sonicator and 30 min of shaking (1500 rpm, at 37°C). The urea concentration

was reduced to 2 M with H_2O prior to protein digestion with trypsin (Promega, Alexandria,

NSW, Australia) at a 1:60 trypsin:protein ratio and subsequent overnight digestion at 37°C.

The next day, samples were acidified with trifluoroacetic acid (1% [v/v] final concentration)

and centrifuged for 5 min at 20,100 g at RT. The supernatants were desalted on pre-activated

(100% acetonitrile [ACN]) and pre-equilibrated (0.1% TFA, 2% ACN) Empore™ SPE Disks

(matrix active group polystyrene-divinylbenzene [SDB-XC] Merck, Bayswater, VIC,

Australia) stage tips (Kulak et al., 2014) made in-house, before being washed (0.1% TFA, 2%

ACN) and eluted in 0.1% TFA, 80% ACN. Samples were concentrated under vacuum and

reconstituted in 0.1% TFA, 2% ACN. After 15 min sonication and subsequent vortexing

samples were centrifuged at 20,100 g at RT before estimation of peptide concentration

(Direct Detect, Merck). Approximately 600-800 ng of peptides were analyzed on a Thermo Q

Exactive™ Plus mass spectrometer coupled to an Ultimate 3000 HPLC (both Thermo Fisher

Scientific, Mulgrave, VIC, Australia). Peptides were first loaded onto a trap column (Dionex-

C18, 100 Å, 75 μm x 2 cm; Thermo Fisher Scientific) at an isocratic flow of 5 μL min^{-1} of

2% (v/v) ACN containing 0.1 % (v/v) formic acid (FA) for 5 min before switching the trap-

column in line with the analytical column (Dionex-C18, 100 Å, 75 μm x 50 cm; Thermo

Fisher Scientific). The separation of peptides was performed at 300 nL min^{-1} using a

nonlinear ACN gradient of buffer A (2% ACN, 0.1% FA) and buffer B (80% ACN, 0.1%

FA) over 125 min including void and equilibration. Data were collected in positive mode

using Data Dependent Acquisition using m/z 375–1400 as MS scan range, HCD for MS/MS

of the 15 most intense ions with $z \geq 2$. Other instrument parameters were: MS1 scan at 70,000

resolution (at 200 m/z), MS maximum injection time 50 ms, AGC target 3e6, stepped

normalized collision energy of 27, 30, 32, isolation window of 1.6 m/z, MS/MS resolution

17,500, MS/MS AGC target of 5e4, MS/MS maximum injection time 50 ms, minimum

intensity was set at 2e3 and dynamic exclusion was set to 30 s.

Raw files were analyzed using the MaxQuant platform (Cox and Mann, 2008) version

1.6.1.0, searching against the Uniprot human database containing reviewed, canonical

variants in FASTA format (June 2018) and a database containing common contaminants by

the Andromeda search engine (Cox et al., 2011). Default search parameters for a label free

quantification (LFQ) experiment were used with modifications. In brief, "Label free

quantification" was set to "LFQ" using a minimum ratio count of 2. Cysteine

carbamidomethylation was used as a fixed modification, and N-terminal acetylation and

methionine oxidation were used as variable modifications. False discovery rates of 1% for

proteins and peptides were applied by searching a reverse database, and 'match from and to',

'match between runs' options were enabled with a match time window of 0.7 min. Unique

and razor peptides with a minimum ratio count of 2 were used for quantification.

Bioinformatic analysis of proteomics data. The R package *limma* (Ritchie et al., 2015) was

used to conduct the differential expression analysis of MaxQuant LFQ intensities (extracted

from proteinGroups.txt) after first performing normalization using variance stabilizing

normalization (VSN) as found in the *limma* package. Identifications labelled by MaxQuant as

'only identified by site', 'reverse' and 'potential contaminant' were removed. Proteins having

less than 70% valid values were removed and remaining missing data was imputed using

QRILC method from the *imputeLCMD* package in R (Lazar). Differential expression

analysis was performed between BL and then each subsequent time point to show

comparative changes between each of the training volumes, though all comparisons were

accounted for within the analysis. Linear modelling was determined using eBayes in the

limma package. The resulting differential expression values were filtered for an adjusted *P*

value < 0.01 using the Benjamini Hochberg method. For the mitochondrial normalization, all

"Known Mitochondrial" proteins identified using the Integrated Mitochondrial Protein Index

(IMPI) (Smith and Robinson, 2019) were subset from the rest of the dataset, and then

followed the normalization and statistical validation as described above. Heatmaps were

produced using hierarchical clustering using the "complete" method. Gene ontology of the

clusters was determined by taking the proteins identified in the cluster and performing an

enrichment analysis using the ClueGO (v2.5.6) application in Cytoscape (v3.7.1) using

default settings except for a GO tree interval of 3 to 5, with only the Biological Processes

Ontology switched on (Figure 2F and Figure S2A). Full tables of identified proteins and post-

normalization differentially expressed proteins in their respective clusters and their

annotations can be found in Table S3 to 7.

The heatmaps presented in Figure 3 and discussed in the Results section were generated

according to Reactome pathways and/or literature searches (Table S6). Proteins involved in

two or more pathways were either presented in both pathways, or were assigned to the

pathway involving the protein's primary function and/or most closely matching their training-

induced changes. Specifically, "mitochondrial translation" was based on Reactome pathway

R-HSA:5368287 (Mitochondrial translation). "TCA cycle" was based on Reactome pathway

"R-HSA:71403 (Citric acid cycle [TCA cycle]); NNT was removed because its main

physiological function is the generation of NADPH (Kampjut and Sazanov, 2019). "FAO"

was based on Reactome pathway R-HSA:556833 (Metabolism of Lipids); ETFA (Watmough

and Frerman, 2010), ETFB (Watmough and Frerman, 2010), HSD17B10 (Schulz, 1991), and

PLIN5 (Benador et al., 2019) were added based on their involvement with FAO and lipid

metabolism; GPD2 (Mráček et al., 2013; Wang and Hekimi, 2016), GPX1 [R-HSA-3299685

and (Powers et al., 2020)], and GPX4 (Powers et al., 2020) where not presented within this

pathway as their primary function suggested their inclusion in different pathways of Figure 3;

SAR1B was removed because its main function is the regulation of vesicle budding (Takai et

al., 2001). "OXPHOS" was based on Reactome pathway R-HSA:1632000 (Respiratory

electron transport, ATP synthesis by chemiosmotic coupling, and heat production by

uncoupling proteins); ETFA (Watmough and Frerman, 2010), ETFB (Watmough and

Frerman, 2010), ETFDH (Wang and Hekimi, 2016), TRAP1 (Voos, 2013), and CYCS (R-

HSA-3299685) were not presented within this pathway as their primary function and/or

training-induced changes suggested their inclusion in different pathways of Figure 3 was

more appropriate; based on the main function of its constituent proteins the "OXPHOS"

pathway was then subdivided in 2 subgroups: "OXPHOS - subunits", to which MP68

(Fujikawa et al., 2014) was added, and "OXPHOS - Assembly factors" to which ATPAF1

(Signes and Fernandez-Vizarra, 2018), BCS1L (Signes and Fernandez-Vizarra, 2018), and

COA3 (Signes and Fernandez-Vizarra, 2018) were added. "Response to ROS" was based on

Reactome pathway: R-HSA-3299685 (Detoxification of Reactive Oxygen Species); GPX4

was added based on its ability to reduce H_2O_2 (Powers et al., 2020). "Protein Folding" was

based on literature searches: TRAP1, HSPA9, HSPD1, HSPE1, and TIMM44 were all added

based on their involvement in protein folding and quality control mechanisms (Voos, 2013).

"CoQ biosynthesis" was based on Reactome pathway R-HSA-2142789 (Ubiquinol

biosynthesis). "CoQ e- donors" was based on literature searches: DHODH (Wang and

Hekimi, 2016), ETFDH (Wang and Hekimi, 2016), GPD2 (Wang and Hekimi, 2016), IVD

(Finocchiaro et al., 1987), DLD (Xia et al., 2001) and L2HGDH (Oldham et al., 2015), were

all added based on their function to feed electrons into the respiratory chain via CoQ. "Cristae

formation" was based on Reactome pathway R-HSA-8949613 (Cristae formation); OMA1

(Ehses et al., 2009; Frezza et al., 2007) was added based on its involvement in cristae

formation, whereas ATP5A1, ATP5J, ATP5B, ATP5H, and ATP5O (all based on R-

HSA:163200), as well as HSPA9 (Voos, 2013) were not presented within this pathway as

their primary function suggested their inclusion in different pathways of Figure 3.

Lipid extraction for lipidomics. Mitochondrial isolates were extracted using a modified

single-phase chloroform/methanol extraction as described previously (Weir et al., 2013). In

brief, 20 volumes of chloroform:methanol (2:1) were added to the sample along with a series

of internal standards. Samples were vortexed and centrifuged on a rotary mixer for 10 min.

Following sonication on a sonicator bath for 30 min, samples were rested for 20 min prior to

centrifugation at 13,000 g for 10 min. Supernatants were transferred into a 96 well plated,

dried down and reconstituted in 50 µL H_2O saturated butanol and sonicated for 10 min. After

the addition of 50 µL of methanol with 10 mM ammonium formate, samples were

centrifuged at 4000 rpm on a plate centrifuge and transferred into glass vials with inserts for

mass spectrometry analysis.

Targeted lipidomics analysis. LC-MS/MS was performed according to previously published

methods, with slight modification for tissue samples (Huynh et al., 2019). Sample extracts

were analyzed using either (i) an AB Sciex Qtrap 4000 mass spectrometer coupled to an

Agilent 1200 HPLC system for CL assessment, as described preciously (Tan et al., 2020) or

(ii) an Agilent 6490 QQQ mass spectrometer coupled with an Agilent 1290 series HPLC

system for assessment of all other lipid species (Huynh et al., 2019). Lipids run on the

Agilent 6490 were measured using scheduled multiple reaction monitoring with the following

conditions: isolation widths for Q1 and Q3 were set to "unit" resolution (0.7 amu), gas

temperature 150°C, nebulizer 20 psi, sheath gas temperature 200°C, gas flow rate 17 L/min,

capillary voltage 3500 V and sheath gas flow 10 L/min. The list of MRMs used and the

chromatographic conditions were described previously (Huynh et al., 2019).

Bioinformatic analysis of lipidomics data. For lipidomics we identified that the increase in

CLs in the un-normalized data was due to a training-induced increase in MPE in our

mitochondrial isolates. To eliminate this bias, all lipid species were normalized by total CL

amount (Table S8). Lipid species were log transformed before undergoing differential

expression analysis in *limma* (Ritchie et al., 2015), with linear models using the eBayes

function. The resulting differential expression values were filtered for an adjusted P value <

0.05 using Benjamini Hochberg method (Table S9). Heatmaps were visualized using

hierarchical clustering using the "complete" method. Profile plots were determined by taking

the mean of the z-score for all participants' samples for each lipid species at each time point

(Figure 4D). Forest plots are depicted using logFC results from *Limma* analysis (Figure S3).

Higher classes were determined through descriptions of the literature as assigned by the Lipid

Metabolites and Pathways Strategy (Lipid MAPS) (Fahy et al., 2011).

QUANTIFICATION AND STATISTICAL ANALYSIS

All values are reported as means ± SD, unless otherwise specified. For non-omics analyses:

outliers were first removed using the ROUT method set at Q = 1% (Motulsky and Brown,

2006). Normally distributed datasets (Shapiro-Wilk test $P > 0.05$) were analyzed by a

repeated measures one-way ANOVA followed by Tukey's correction post hoc testing. Non-

normally distributed datasets (Shapiro-Wilk test $P < 0.05$) were transformed using, in order,

log(Y), 1/Y, √Y, until normality was met, before being analyzed as above; datasets that

remained non-normal following 3 independent transformation attempts were analyzed using

the non-parametric Friedman test on the raw data followed by Dunn's correction post hoc

testing. The level of statistical significance was set a priori at $P < 0.05$. GraphPad Prism (v.

8.4.2) was used for all statistical non-omics analyses. Differential expression for all omics

was completed in R (v.3.6.3), using the packages, limma and edgeR to determine

significantly enriched results between the groups. Significance was determined by an

adjusted p value < 0.05 for the transcriptomics and lipidomics and 0.01 for the proteomics.

Statistical details can be found in the associated figure legends.

REFERENCES

Baldwin, J., Snow, R.J., and Febbraio, M.A. (2000). Effect of training status and relative
exercise intensity on physiological responses in men. Med. Sci. Sports Exerc. *32*, 1648-1654.

Barnard, R.J., Edgerton, V.R., and Peter, J. (1970). Effect of exercise on skeletal muscle. I.
Biochemical and histochemical properties. J. Appl. Physiol. *28*, 762-766.

Benador, I.Y., Veliova, M., Liesa, M., and Shirihai, O.S. (2019). Mitochondria bound to lipid
droplets: where mitochondrial dynamics regulate lipid storage and utilization. Cell Metab. *29*,
827-835.

Bishop, D.J., Botella, J., Genders, A.J., Lee, M.J.-C., Saner, N.J., Kuang, J., Yan, X., and
Granata, C. (2019). High-Intensity Exercise and Mitochondrial Biogenesis: Current
Controversies and Future Research Directions. Physiology *34*, 56-70.

Bishop, D.J., Granata, C., and Eynon, N. (2014). Can we optimise the exercise training
prescription to maximise improvements in mitochondria function and content? Biochim.
Biophys. Acta, Gen. Subj. *1840*, 1266-1275.

Bishop, D.J., Jenkins, D.G., McEniery, M., and Carey, M.F. (2000). Relationship between
plasma lactate parameters and muscle characteristics in female cyclists. Med. Sci. Sports
Exerc. *32*, 1088-1093.

Booth, F.W., Gordon, S.E., Carlson, C.J., and Hamilton, M.T. (2000). Waging war on
modern chronic diseases: primary prevention through exercise biology. J. Appl. Physiol. *88*,
774-787.

Bruce, C.R., Thrush, A.B., Mertz, V.A., Bezaire, V., Chabowski, A., Heigenhauser, G.J.F.,
and Dyck, D.J. (2006). Endurance training in obese humans improves glucose tolerance and
mitochondrial fatty acid oxidation and alters muscle lipid content. American Journal of
Physiology - Endocrinology and Metabolism *291*, E99-E107.

Calvo, S.E., Clauser, K.R., and Mootha, V.K. (2016). MitoCarta2.0: an updated inventory of
mammalian mitochondrial proteins. Nucleic Acids Res. *44*, D1251-1257.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and
Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. BMC Bioinformatics *14*, 128.

Cox, J., and Mann, M. (2008). MaxQuant enables high peptide identification rates,
individualized ppb-range mass accuracies and proteome-wide protein quantification. Nat.
Biotechnol. *26*, 1367-1372.

Cox, J., Neuhauser, N., Michalski, A., Scheltema, R.A., Olsen, J.V., and Mann, M. (2011).
Andromeda: a peptide search engine integrated into the MaxQuant environment. J. Proteome
Res. *10*, 1794-1805.

Desler, C., Hansen, T.L., Frederiksen, J.B., Marcker, M.L., Singh, K.K., and Juel Rasmussen,
L. (2012). Is there a link between mitochondrial reserve respiratory capacity and aging?
Journal of aging research *2012*.

Dube, J., Amati, F., Toledo, F., Stefanovic-Racic, M., Rossi, A., Coen, P., and Goodpaster, B.
(2011). Effects of weight loss and exercise on insulin resistance, and intramyocellular
triacylglycerol, diacylglycerol and ceramide. Diabetologia *54*, 1147-1156.

Egan, B., Dowling, P., O'Connor, P.L., Henry, M., Meleady, P., Zierath, J.R., and O'Gorman,
D.J. (2011). 2-D DIGE analysis of the mitochondrial proteome from human skeletal muscle
reveals time course-dependent remodelling in response to 14 consecutive days of endurance
exercise training. Proteomics *11*, 1413-1428.

Ehses, S., Raschke, I., Mancuso, G., Bernacchia, A., Geimer, S., Tondera, D., Martinou, J.-
C., Westermann, B., Rugarli, E.I., and Langer, T. (2009). Regulation of OPA1 processing and
mitochondrial fusion by m-AAA protease isoenzymes and OMA1. J. Cell Biol. *187*, 1023-
1036.

Eramo, M.J., Lisnyak, V., Formosa, L.E., and Ryan, M.T. (2020). The 'mitochondrial contact
site and cristae organising system'(MICOS) in health and human disease. The Journal of
Biochemistry *167*, 243-255.

Fahy, E., Cotter, D., Sud, M., and Subramaniam, S. (2011). Lipid classification, structures
and tools. Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids *1811*,
637-647.

Finocchiaro, G., Ito, M., and Tanaka, K. (1987). Purification and properties of short chain
acyl-CoA, medium chain acyl-CoA, and isovaleryl-CoA dehydrogenases from human liver. J.
Biol. Chem. *262*, 7982-7989.

Franko, A., Baris, O.R., Bergschneider, E., Von Toerne, C., Hauck, S.M., Aichler, M.,
Walch, A.K., Wurst, W., Wiesner, R.J., and Johnston, I.C. (2013). Efficient isolation of pure
and functional mitochondria from mouse tissues using automated tissue disruption and
enrichment with anti-TOM22 magnetic beads. PLoS One *8*, e82392.

Frazier, A.E., Vincent, A.E., Turnbull, D.M., Thorburn, D.R., and Taylor, R.W. (2020).
Assessment of mitochondrial respiratory chain enzymes in cells and tissues. In Methods Cell
Biol. (Elsevier), pp. 121-156.

Frezza, C., Cipolat, S., and Scorrano, L. (2007). Organelle isolation: functional mitochondria
from mouse liver, muscle and cultured filroblasts. Nat. Protoc. *2*, 287.

Fujikawa, M., Ohsakaya, S., Sugawara, K., and Yoshida, M. (2014). Population of ATP
synthase molecules in mitochondria is limited by available 6.8-kDa proteolipid protein
(MLQ). Genes Cells *19*, 153-160.

Gebert, N., Joshi, A.S., Kutik, S., Becker, T., McKenzie, M., Guan, X.L., Mooga, V.P.,
Stroud, D.A., Kulkarni, G., and Wenk, M.R. (2009). Mitochondrial cardiolipin involved in
outer-membrane protein biogenesis: implications for Barth syndrome. Curr. Biol. *19*, 2133-
2139.

Goodpaster, B.H., and Kelley, D.E. (1998). Role of muscle in triglyceride metabolism. Curr.
Opin. Lipidol. *9*, 231-236.

Gorman, G.S., Chinnery, P.F., DiMauro, S., Hirano, M., Koga, Y., McFarland, R.,
Suomalainen, A., Thorburn, D.R., Zeviani, M., and Turnbull, D.M. (2016). Mitochondrial
diseases. Nature reviews Disease primers *2*, 1-22.

Granata, C., Jamnick, N.A., and Bishop, D.J. (2018a). Principles of exercise prescription, and
how they influence exercise-induced changes of transcription factors and other regulators of
mitochondrial biogenesis. Sports Med. *48*, 1541–1559.

Granata, C., Jamnick, N.A., and Bishop, D.J. (2018b). Training-induced changes in
mitochondrial content and respiratory function in human skeletal muscle. Sports Med. *48*,
1809–1828.

Granata, C., Oliveira, R.S.F., Little, J.P., Renner, K., and Bishop, D.J. (2016a). Mitochondrial
adaptations to high-volume exercise training are rapidly reversed after a reduction in training
volume in human skeletal muscle. FASEB J. *30*, 3413-3423.

Granata, C., Oliveira, R.S.F., Little, J.P., Renner, K., and Bishop, D.J. (2016b). Training
intensity modulates changes in PGC-1α and p53 protein content and mitochondrial
respiration, but not markers of mitochondrial content in human skeletal muscle. FASEB J. *30*,
959-970.

Greggio, C., Jha, P., Kulkarni, S.S., Lagarrigue, S., Broskey, N.T., Boutant, M., Wang, X.,
Alonso, S.C., Ofori, E., and Auwerx, J. (2017). Enhanced respiratory chain supercomplex
formation in response to exercise in human skeletal muscle. Cell Metab. *25*, 301-311.

Groennebaek, T., Nielsen, J., Jespersen, N.R., Bøtker, H.E., de Paoli, F.V., Miller, B.F., and
Vissing, K. (2020). Utilization of biomarkers as predictors of skeletal muscle mitochondrial
content after physiological intervention and in clinical settings. American Journal Of
Physiology-Endocrinology And Metabolism *318*, E886-E889.

Guarani, V., McNeill, E.M., Paulo, J.A., Huttlin, E.L., Fröhlich, F., Gygi, S.P., Van Vactor,
D., and Harper, J.W. (2015). QIL1 is a novel mitochondrial protein required for MICOS
complex stability and cristae morphology. Elife *4*, e06265.

Hirst, J. (2018). Open questions: respiratory chain supercomplexes—why are they there and
what do they do? BMC Biol. *16*, 111.

Hock, D.H., Robinson, D.R., and Stroud, D.A. (2020). Blackout in the powerhouse: clinical
phenotypes associated with defects in the assembly of OXPHOS complexes and the
mitoribosome. Biochem. J. *477*, 4085-4132.

Holloszy, J.O. (1967). Biochemical adaptations in muscle. Effects of exercise on
mitochondrial oxygen uptake and respiratory enzyme activity in skeletal muscle. J. Biol.
Chem. *242*, 2278-2282.

Holloway, K.V., O'Gorman, M., Woods, P., Morton, J.P., Evans, L., Cable, N.T., Goldspink,
D.F., and Burniston, J.G. (2009). Proteomic investigation of changes in human vastus
lateralis muscle in response to interval-exercise training. Proteomics *9*, 5155-5174.

Hoppeler, H., Lüthi, P., Claassen, H., Weibel, E.R., and Howald, H. (1973). The
ultrastructure of the normal human skeletal muscle. Pflügers Archiv *344*, 217-232.

Hostrup, M., Onslev, J., Jacobson, G.A., Wilson, R., and Bangsbo, J. (2018). Chronic β2-
adrenoceptor agonist treatment alters muscle proteome and functional adaptations induced by
high intensity training in young men. The Journal of physiology *596*, 231-252.

Huynen, M.A., Mühlmeister, M., Gotthardt, K., Guerrero-Castillo, S., and Brandt, U. (2016).
Evolution and structural organization of the mitochondrial contact site (MICOS) complex and
the mitochondrial intermembrane space bridging (MIB) complex. Biochimica et Biophysica
Acta (BBA)-Molecular Cell Research *1863*, 91-101.

Huynh, K., Barlow, C.K., Jayawardana, K.S., Weir, J.M., Mellett, N.A., Cinel, M., Magliano,
D.J., Shaw, J.E., Drew, B.G., and Meikle, P.J. (2019). High-throughput plasma lipidomics:
detailed mapping of the associations with cardiometabolic risk factors. Cell chemical biology
26, 71-84. e74.

Ikon, N., and Ryan, R.O. (2017). Cardiolipin and mitochondrial cristae organization.
Biochimica et Biophysica Acta (BBA)-Biomembranes *1859*, 1156-1163.

Jamnick, N.A., Botella, J., Pyne, D.B., and Bishop, D.J. (2018). Manipulating graded
exercise test variables affects the validity of the lactate threshold and V̇ O2peak. PLoS One
13, e0199794.

Jha, P., Wang, X., and Auwerx, J. (2016). Analysis of Mitochondrial Respiratory Chain
Supercomplexes Using Blue Native Polyacrylamide Gel Electrophoresis (BN-PAGE). Curr.
Protoc. Mouse Biol., 1-14.

Kampjut, D., and Sazanov, L.A. (2019). Structure and mechanism of mitochondrial proton-
translocating transhydrogenase. Nature *573*, 291-295.

Kulak, N.A., Pichler, G., Paron, I., Nagaraj, N., and Mann, M. (2014). Minimal, encapsulated
proteomic-sample processing applied to copy-number estimation in eukaryotic cells. Nature
methods *11*, 319.

Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev,
S., Jenkins, S.L., Jagodnik, K.M., and Lachmann, A. (2016). Enrichr: a comprehensive gene
set enrichment analysis web server 2016 update. Nucleic Acids Res. *44*, W90-W97.

LaHue, S.C., Comella, C.L., and Tanner, C.M. (2016). The best medicine? The influence of
physical activity and inactivity on Parkinson's disease. Mov. Disord. *31*, 1444-1454.

Lapuente-Brun, E., Moreno-Loshuertos, R., Acín-Pérez, R., Latorre-Pellicer, A., Colás, C.,
Balsa, E., Perales-Clemente, E., Quirós, P.M., Calvo, E., and Rodríguez-Hernández, M.
(2013). Supercomplex assembly determines electron flux in the mitochondrial electron
transport chain. Science *340*, 1567-1570.

Larsen, S., Nielsen, J., Hansen, C.N., Nielsen, L.B., Wibrand, F., Stride, N., Schroder, H.D.,
Boushel, R., Helge, J.W., Dela, F., et al. (2012). Biomarkers of mitochondrial content in
skeletal muscle of healthy young human subjects. J. Physiol. *590*, 3349-3360.

Lazar, C. imputeLCMD: A Collection of Methods for Left-Censored Missing Data
Imputation. R package, version 2.0.

Lin, M.T., and Beal, M.F. (2006). Mitochondrial dysfunction and oxidative stress in
neurodegenerative diseases. Nature *443*, 787-795.

López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2013). The
hallmarks of aging. Cell *153*, 1194-1217.

Malhotra, A., Xu, Y., Ren, M., and Schlame, M. (2009). Formation of molecular species of
mitochondrial cardiolipin. 1. A novel transacylation mechanism to shuttle fatty acids between
sn-1 and sn-2 positions of multiple phospholipid species. Biochimica et Biophysica Acta
(BBA)-Molecular and Cell Biology of Lipids *1791*, 314-320.

Martínez-Reyes, I., and Chandel, N.S. (2020). Mitochondrial TCA cycle metabolites control
physiology and disease. Nature communications *11*, 1-11.

McKenzie, M., Lazarou, M., Thorburn, D.R., and Ryan, M.T. (2007). Analysis of
mitochondrial subunit assembly into respiratory chain complexes using Blue Native
polyacrylamide gel electrophoresis. Anal. Biochem. *364*, 128-137.

Meinild Lundby, A.K., Jacobs, R.A., Gehrig, S., de Leur, J., Hauser, M., Bonne, T.C., Flück,
D., Dandanell, S., Kirk, N., Kaech, A., et al. (2018). Exercise training increases skeletal
muscle mitochondrial volume density by enlargement of existing mitochondria and not de
novo biogenesis. Acta Physiol *222*, e12905.

Morgan, T., Cobb, L., Short, F., Ross, R., and Gunn, D. (1971). Effects of long-term exercise
on human muscle mitochondria. In Muscle metabolism during exercise (Springer), pp. 87-95.

Motulsky, H.J., and Brown, R.E. (2006). Detecting outliers when fitting data with nonlinear
regression–a new method based on robust nonlinear regression and the false discovery rate.
BMC Bioinformatics *7*, 123.

Mráček, T., Drahota, Z., and Houštěk, J. (2013). The function and the role of the
mitochondrial glycerol-3-phosphate dehydrogenase in mammalian tissues. Biochimica et
Biophysica Acta (BBA)-Bioenergetics *1827*, 401-410.

Neufer, P.D., Bamman, M.M., Muoio, D.M., Bouchard, C., Cooper, D.M., Goodpaster, B.H.,
Booth, F.W., Kohrt, W.M., Gerszten, R.E., Mattson, M.P., et al. (2015). Understanding the
cellular and molecular mechanisms of physical activity-induced health benefits. Cell Metab.
22, 4-11.

Nielsen, J., Gejl, K.D., Hey-Mogensen, M., Holmberg, H.C., Suetta, C., Krustrup, P.,
Elemans, C.P., and Ørtenblad, N. (2017). Plasticity in mitochondrial cristae density allows
metabolic capacity modulation in human skeletal muscle. J. Physiol. *595*, 2839-2847.

Nunnari, J., and Suomalainen, A. (2012). Mitochondria: in sickness and in health. Cell *148*,
1145-1159.

Oldham, W.M., Clish, C.B., Yang, Y., and Loscalzo, J. (2015). Hypoxia-mediated increases
in L-2-hydroxyglutarate coordinate the metabolic response to reductive stress. Cell Metab.
22, 291-303.

Ørtenblad, N. (2018). Mitochondrial increase in volume density with exercise training: More,
larger or better? Acta Physiologica *222*, e12976.

Palmieri, F. (2013). The mitochondrial transporter family SLC25: identification, properties
and physiopathology. Mol. Aspects Med. *34*, 465-484.

Pedersen, B.K., and Saltin, B. (2006). Evidence for prescribing exercise as therapy in chronic
disease. Scand. J. Med. Sci. Sports *16*, 3-63.

Perreault, L., Newsom, S.A., Strauss, A., Kerege, A., Kahn, D.E., Harrison, K.A., Snell-
Bergeon, J.K., Nemkov, T., D'Alessandro, A., and Jackman, M.R. (2018). Intracellular
localization of diacylglycerols and sphingolipids influences insulin sensitivity and
mitochondrial function in human skeletal muscle. JCI insight *3*.

Pesta, D., and Gnaiger, E. (2012). High-resolution respirometry: OXPHOS protocols for
human cells and permeabilized fibers from small biopsies of human muscle. In Mitochondrial
Bioenergetics: Methods and Protocols. C.M. Palmeira, and A.J. Moreno, eds. (Springer
Science+Business Media), pp. 25-58.

Pfeiffer, K., Gohil, V., Stuart, R.A., Hunte, C., Brandt, U., Greenberg, M.L., and Schägger,
H. (2003). Cardiolipin stabilizes respiratory chain supercomplexes. J. Biol. Chem. *278*,
52873-52880.

Pillon, N.J., Gabriel, B.M., Dollet, L., Smith, J.A., Puig, L.S., Botella, J., Bishop, D.J.,
Krook, A., and Zierath, J.R. (2020). Transcriptomic profiling of skeletal muscle adaptations
to exercise and inactivity. Nature communications *11*, 1-15.

Powers, S.K., Deminice, R., Ozdemir, M., Yoshihara, T., Bomkamp, M.P., and Hyatt, H.
(2020). Exercise-induced oxidative stress: Friend or foe? Journal of sport and health science.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015).
limma powers differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res. *43*, e47-e47.

Robinson, M.M., Dasari, S., Konopka, A.R., Johnson, M.L., Manjunatha, S., Esponda, R.R.,
Carter, R.E., Lanza, I.R., and Nair, K.S. (2017). Enhanced protein translation underlies
improved metabolic and physical adaptations to different exercise training modes in young
and old humans. Cell Metab. *25*, 581-592.

Scalzo, R.L., Peltonen, G.L., Binns, S.E., Shankaran, M., Giordano, G.R., Hartley, D.A.,
Klochak, A.L., Lonac, M.C., Paris, H.L.R., Szallar, S.E., et al. (2014). Greater muscle protein
synthesis and mitochondrial biogenesis in males compared with females during sprint interval
training. FASEB J. *28*, 2705-2714.

Schägger, H., and Pfeiffer, K. (2000). Supercomplexes in the respiratory chains of yeast and
mammalian mitochondria. The EMBO journal *19*, 1777-1783.

Schulz, H. (1991). Beta oxidation of fatty acids. Biochimica et Biophysica Acta (BBA)-
Lipids and Lipid Metabolism *1081*, 109-120.

Shankaran, M., King, C.L., Angel, T.E., Holmes, W.E., Li, K.W., Colangelo, M., Price, J.C.,
Turner, S.M., Bell, C., and Hamilton, K.L. (2016). Circulating protein synthesis rates reveal
skeletal muscle proteome dynamics. The Journal of clinical investigation *126*, 288-302.

Shepherd, S., Cocks, M., Meikle, P., Mellett, N., Ranasinghe, A., Barker, T., Wagenmakers,
A., and Shaw, C. (2017). Lipid droplet remodelling and reduced muscle ceramides following
sprint interval and moderate-intensity continuous exercise training in obese males. Int. J.
Obes. *41*, 1745.

Signes, A., and Fernandez-Vizarra, E. (2018). Assembly of mammalian oxidative
phosphorylation complexes I–V and supercomplexes. Essays Biochem. *62*, 255-270.

Smith, A.C., and Robinson, A.J. (2019). MitoMiner v4. 0: an updated database of
mitochondrial localization evidence, phenotypes and diseases. Nucleic Acids Res. *47*, D1225-
D1228.

Spinelli, J.B., and Haigis, M.C. (2018). The multifaceted contributions of mitochondria to
cellular metabolism. Nat. Cell Biol. *20*, 745-754.

Stefely, J.A., and Pagliarini, D.J. (2017). Biochemistry of mitochondrial coenzyme Q
biosynthesis. Trends Biochem. Sci. *42*, 824-843.

Stroud, D.A., Surgenor, E.E., Formosa, L.E., Reljic, B., Frazier, A.E., Dibley, M.G.,
Osellame, L.D., Stait, T., Beilharz, T.H., and Thorburn, D.R. (2016). Accessory subunits are
integral for assembly and function of human mitochondrial complex I. Nature *538*, 123.

Svensson, M., Malm, C., Tonkonogi, M., Ekblom, B., Sjödin, B., and Sahlin, K. (1999).
Effect of Q10 supplementation on tissue Q10 levels and adenine nucleotide catabolism during
high-intensity exercise. Int. J. Sport Nutr. Exerc. Metab. *9*, 166-180.

Takai, Y., Sasaki, T., and Matozaki, T. (2001). Small GTP-binding proteins. Physiol. Rev.
81, 153-208.

Tan, S.M., Ziemann, M., Thallas-Bonke, V., Snelson, M., Kumar, V., Laskowski, A.,
Nguyen, T.-V., Huynh, K., Clarke, M.V., and Libianto, R. (2020). Complement C5a Induces
Renal Injury in Diabetic Kidney Disease by Disrupting Mitochondrial Metabolic Agility.
Diabetes *69*, 83-98.

Ubaida-Mohien, C., Gonzalez-Freire, M., Lyashkov, A., Moaddel, R., Chia, C.W.,
Simonsick, E.M., Sen, R., and Ferrucci, L. (2019). Physical activity associated proteomics of
skeletal muscle: being physically active in daily life may protect skeletal muscle from aging.
Front. Physiol. *10*, 312.

Voos, W. (2013). Chaperone–protease networks in mitochondrial protein homeostasis.
Biochimica et Biophysica Acta (BBA)-Molecular Cell Research *1833*, 388-399.

Walther, T.C., and Farese Jr, R.V. (2012). Lipid droplets and cellular lipid metabolism.
Annu. Rev. Biochem. *81*, 687-714.

Wang, H., Sreenivasan, U., Hu, H., Saladino, A., Polster, B.M., Lund, L.M., Gong, D.-w.,
Stanley, W.C., and Sztalryd, C. (2011). Perilipin 5, a lipid droplet-associated protein,
provides physical and metabolic linkage to mitochondria. J. Lipid Res. *52*, 2159-2168.

Wang, Y., and Hekimi, S. (2016). Understanding ubiquinone. Trends Cell Biol. *26*, 367-378.

Wang, Y., and Hekimi, S. (2019). The complexity of making ubiquinone. Trends Endocrinol.
Metab. *30*, 929-943.

Watmough, N.J., and Frerman, F.E. (2010). The electron transfer flavoprotein: ubiquinone
oxidoreductases. Biochimica et Biophysica Acta (BBA)-Bioenergetics *1797*, 1910-1916.

Weber, T.A., Koob, S., Heide, H., Wittig, I., Head, B., van der Bliek, A., Brandt, U.,
Mittelbronn, M., and Reichert, A.S. (2013). APOOL is a cardiolipin-binding constituent of
the Mitofilin/MINOS protein complex determining cristae morphology in mammalian
mitochondria. PLoS One *8*, e63683.

Weir, J.M., Wong, G., Barlow, C.K., Greeve, M.A., Kowalczyk, A., Almasy, L., Comuzzie,
A.G., Mahaney, M.C., Jowett, J.B., and Shaw, J. (2013). Plasma lipid profiling in a large
population-based cohort. J. Lipid Res. *54*, 2898-2908.

Wiedemann, N., and Pfanner, N. (2017). Mitochondrial machineries for protein import and
assembly. Annu. Rev. Biochem. *86*, 685-714.

Wyckelsma, V.L., Levinger, I., McKenna, M.J., Formosa, L.E., Ryan, M.T., Petersen, A.C.,
Anderson, M.J., and Murphy, R.M. (2017). Preservation of skeletal muscle mitochondrial
content in older adults: relationship between mitochondria, fibre type and high-intensity
exercise training. The Journal of physiology *595*, 3345-3359.

Xia, L., Björnstedt, M., Nordman, T., Eriksson, L.C., and Olsson, J.M. (2001). Reduction of
ubiquinone by lipoamide dehydrogenase: An antioxidant regenerating pathway. Eur. J.
Biochem. *268*, 1486-1490.

1368