TITLE

Impacts of an urban sanitation intervention on fecal indicators and the prevalence of human fecal

contamination in Mozambique

AUTHORS AND AFFILIATIONS

David A. Holcomb[a,b] , Jackie Knee[c,d] , Drew Capone[a,c], Trent Sumner[c], Zaida Adriano[e], Rassul Nalá[f],

Oliver Cumming[d], Joe Brown[a], Jill R. Stewart[a,*]

[a] Department of Environmental Sciences and Engineering, Gillings School of Global Public Health,

University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States

[b] Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina at

Chapel Hill, Chapel Hill, North Carolina, United States

[c] School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, Georgia,

United States

[d] Department of Disease Control, London School of Hygiene & Tropical Medicine, London, United

Kingdom

[e] We Consult, Maputo, Mozambique

[f] Instituto Nacional de Saúde, Ministério da Saúde, Maputo, Mozambique

CORRESPONDING AUTHOR

* E-mail: jill.stewart@unc.edu; phone: (919) 966-7553

ABSTRACT

Fecal source tracking (FST) may be useful to assess pathways of fecal contamination in domestic environments and to estimate the impacts of water, sanitation, and hygiene (WASH) interventions in low-income settings. We measured two non-specific and two human-associated fecal indicators in water, soil, and surfaces before and after a shared latrine intervention from low-income households in Maputo, Mozambique participating in the Maputo Sanitation (MapSan) trial. Up to a quarter of households were impacted by human fecal contamination, but trends were unaffected by improvements to shared sanitation facilities. The intervention reduced *E. coli* gene concentrations in soil but did not impact culturable *E. coli* or the prevalence of human FST markers in a difference-in-differences analysis. Using a novel Bayesian hierarchical modeling approach to account for human marker diagnostic sensitivity and specificity, we revealed a high amount of uncertainty associated with human FST measurements and intervention effect estimates. The field of microbial source tracking would benefit from adding measures of diagnostic accuracy to better interpret findings, particularly when FST analyses convey insufficient information for robust inference. With improved measures, FST could help identify dominant pathways of human and animal fecal contamination in communities and guide implementation of effective interventions to safeguard health.

KEYWORDS

Diagnostic accuracy; water, sanitation, and hygiene; shared sanitation; microbial source tracking; fecal indicator; qPCR; Bayesian hierarchical model

44   SYNOPSIS

45   An urban sanitation intervention had minimal and highly uncertain effects on human fecal

46   contamination after accounting for fecal indicator sensitivity and specificity.

48   TOC GRAPHIC/ABSTRACT ART

Introduction

Water, sanitation, and hygiene (WASH) interventions aim to improve health by preventing exposure to enteric pathogens, which are introduced to the environment in the feces of infected human and animal hosts.[1] Environmental pathways of pathogen exposure include contaminated environmental compartments like water, soil, and surfaces, as well as hands, flies, food, and fomites that have been in contact with contaminated environments.[2–4] Recent evaluations of a range of WASH interventions found inconsistent and largely negligible impacts on child diarrhea, growth, and enteric infection.[5–12] Notably, combined interventions did not provide greater protection than their constituent interventions alone, suggesting that key sources of pathogens and pathways of exposure are inadequately addressed by conventional WASH strategies.[6,7,9,13–15]

Characterizing fecal contamination in potential exposure pathways may help explain why specific interventions do or do not improve health by identifying which pathways the intervention interrupts and which remain unaffected. Fecal contamination is typically assessed by measuring fecal indicator organisms, microbes abundant in feces used to infer the presence of fecal contamination and therefore the likely presence of enteric pathogens, which are challenging to measure directly due to their diversity and low environmental concentrations.[15,16] Indicator organisms can also be used for fecal source tracking (FST) by targeting microbes specific to the feces of a particular host. Animals are important sources of fecal contamination in both domestic and public environments but traditional efforts have focused on preventing exposure to human feces; differentiating between human and various animal feces could inform more appropriate intervention approaches.[4,17–22]

Fecal indicator approaches have increasingly been applied to domestic environments in

low-income settings with high burdens of enteric disease.[3,15–18,23–26] Occurrence of non-specific

indicators like *Escherichia coli* is challenging to interpret in these settings due to elevated and

highly variable ambient concentrations, possibly from naturalized sources, which are typically

assessed in limited numbers of (cross-sectional) observations from each environmental

compartment.[16,27–30] Other than ruminant FST markers, host-associated fecal indicators have

demonstrated poor diagnostic accuracy in domestic settings.[17,31,32,26,16] The use of multiple FST

markers has been proposed to help address the limited accuracy of individual indicators.[33,34]

Several studies have calculated the conditional probability of contamination by a specific fecal

source given the detection of one or more source-associated indicators in one sample.[31,34–36] Such

analyses provide valuable intuition about the uncertainty associated with individual

measurements, which can be particularly important in decision-making contexts like beach

closures. To our knowledge, diagnostic performance has not been similarly accounted for when

FST has been used to infer patterns and predictors of source-specific fecal contamination in

domestic environments, likely overstating the confidence of such estimates.[4,17,18,26,37–39]

In this study, we analyze two non-specific and two human-associated fecal indicators in

water, soil, and surfaces from low-income households in Maputo, Mozambique before and after

a shared sanitation intervention. We explore the conditional probability of human fecal

contamination in individual samples under different prevalence and indicator detection scenarios

and develop a Bayesian hierarchical modeling approach that accounts for the diagnostic accuracy

of multiple markers to estimate the prevalence of source-specific fecal contamination. Finally,

we implement these models using both human markers to estimate intervention effects on the

prevalence of human fecal contamination in multiple exposure pathways.

Materials and Methods

Study setting and intervention

We characterized fecal contamination of households with children participating in the Maputo Sanitation (MapSan) study (clinicaltrials.gov NCT02362932), a prospective, controlled before and after health impact trial of an urban, onsite sanitation intervention.[40] The intervention was delivered to compounds (self-defined clusters of households sharing outdoor space) in low-income neighborhoods of Maputo, Mozambique, areas with high burdens of enteric disease and predominantly onsite sanitation infrastructure.[41,42] Similar compounds that did not receive the intervention were recruited to serve as control sites. At baseline, both intervention and control compounds shared sanitation facilities in poor condition.[26] The existing shared latrines in intervention compounds were replaced with pour-flush latrines that discharged aqueous effluent to infiltration pits and had sturdy, private superstructures. Intervention latrines were constructed between 2015 – 2016 by the nongovernmental organization (NGO) Water and Sanitation for the Urban Poor (WSUP), which selected intervention sites according to engineering and demand criteria (Table S1).[40]

Data collection

The intervention impact on fecal contamination was evaluated using a controlled before-and-after (CBA) study design.[5,43] Intervention compounds were enrolled immediately before the new latrine was opened for use, with concurrent enrollment of control compounds conducted at a similar frequency (Table S1). Follow-up visits to each compound were conducted approximately 12 months following baseline enrollment. We administered compound-, household-, and child-level surveys during both baseline and follow-up visits, as described elsewhere.[5,42] Concurrent with survey administration during May – August 2015, we opportunistically collected

environmental samples at a subset of MapSan study compounds from the shared outdoor space

and from each household with children participating in the health study (see Supporting

Information [SI]). During the 12-month follow-up phase in June – September 2016, we revisited

the original subset of compounds and collected environmental samples from additional study

compounds not sampled at baseline, as time permitted.

Detailed descriptions of environmental sample collection, processing, and analysis have

been published previously.[26] Briefly, we assessed fecal indicators in five environmental

compartments: compound source water, household stored water, latrine entrance soil, household

entrance soil, and household food preparation surfaces (see SI). Source water and latrine soil

were sampled once per compound on each visit, while stored water, food preparation surfaces,

and household soil were collected from each household with children enrolled in the health

impacts study. Samples were processed by membrane filtration, preceded by manual elution for

soil and swab samples, and the sample filters were analyzed for microbial indicators of fecal

contamination using both culture- and molecular-based detection.[25,26,44] We enumerated

culturable *E. coli* (cEC) from filters on modified mTEC broth (Hi-Media, Mumbai, India) and

immediately archived additional filters at -80°C for molecular analysis.[16,45] Archived filters were

analyzed by three locally validated real-time polymerase chain reaction (qPCR) assays targeting

fecal microbe genes. The EC23S857 (EC23S) assay targets *E. coli* and served as an indicator of

non-specific fecal contamination, while HF183/BacR287 (HF183) and Mnif both target microbes

specific to human feces and served as indicators of human-source fecal contamination.[46–48]

Limits of detection for each assay were previously determined using receiver operating

characteristic (ROC) analysis to identify optimal quantification cycle (Cq) cutoff values (see

SI).[26,49]

DNA was isolated from soil and surface sample filters using the DNeasy PowerSoil Kit

(Qiagen, Hilden, Germany) and from water sample filters with the DNA-EZ ST01 Kit

(GeneRite, North Brunswick, NJ, USA), with a positive control (PC) and negative extraction

control (NEC) included in each batch of up to 22 sample filters. PCs consisted of a clean filter

spiked with 2×10^8 copies of each composite DNA standard (Table S4).[26] Filters were treated

with 3 µg salmon testes DNA (MilliporeSigma, Burlington, MA, USA) immediately before

extraction as a specimen processing control (SPC) to assess PCR inhibition.[50,51] We tested each

extract with four qPCR assays using a CFX96 Touch thermocycler (Bio-Rad, Hercules, CA),

three targeting fecal microbes and Sketa22 targeting the salmon DNA SPC, with 10% of each

sample type analyzed in duplicate for all microbial targets.[52] Each reaction consisted of 12.5 µL

TaqMan Environmental Master Mix 2.0, 2.5 µL 10x primers/probe mix, 5 µL nuclease free

water (NFW), and 5 µL DNA template, for 25 µL total reaction volume. After an initial 10-

minute, 95°C incubation period, cycling conditions specified by the original developers were

followed for each assay (Table S3). Samples with Sketa22 quantification cycle (Cq) values > 3

above the mean Cq of extraction controls (NEC and PC) were considered inhibited and diluted

1:5 for further analysis. Each plate included three no-template controls (NTCs) and five-point,

ten-fold dilution series of three extracted PCs, corresponding to triplicate reactions with 10^5 –

10^1 or 10^6 – 10^2 target gene copies (gc). Target concentrations were estimated from calibration

curves fit to the standard dilution series using multilevel Bayesian regression with varying slopes

and intercepts by extraction batch and instrument run (see SI).[53] Fecal indicator concentrations

were $\log_{10}$ transformed and expressed as $\log_{10}$ colony forming units (cfu) or gc per 100 mL

water, 100 cm^2 surface, or 1 dry gram soil.

Estimating intervention effects

We used a difference-in-differences (DID) approach to estimate the effect of the

intervention on fecal indicator occurrence. DID enables unbiased estimation of the treatment

effect in the absence of randomization, including when different samples of each group are

observed pre- and post-treatment, under the "parallel trend" assumption that all unmeasured

time-varying covariates related to the outcome are constant across treatment groups and that

unmeasured covariates varying between treatment groups are constant through time.[43,54,55]

Although we estimated gene copy concentrations for all fecal indicators assessed by qPCR, we

treated the human markers as binary, diagnostic tests of the presence or absence of human fecal

contamination due to their relatively low baseline detection frequency (and limited availability of

concentration data as a result).[26] By contrast, *E. coli* was detected in the large majority of

baseline samples by both culture and qPCR approaches; treating such outcomes as

presence/absence would discard a great deal of information conveyed by the *E. coli*

concentration measurements, producing a binary outcome with very little variation. Direct DID

estimates for the mean concentration of non-specific indicators and the prevalence of human-

associated indicators were obtained using a bootstrap approach with 2000 samples. We

calculated the mean concentration or prevalence in each of the four design strata (pre-treatment

intervention compounds, post-treatment intervention compounds, pre-treatment control

compounds, and post-treatment controls) by sample type, from which the DID was calculated

directly (see SI). Bootstrap 95% compatibility intervals (CI) were obtained as the 2.5 and 97.5

percentile values of the bootstrap samples.[56]

We also conducted regression analyses incorporating potential confounding variables to

obtain conditional DID estimates. We used the product-term representation of the DID estimator,

in which binary indicators of treatment group, study phase, and their product (interaction) were

included as linear predictors. The coefficient on the product term provides the conditional DID

estimate.[54,57] Separate models were fit for each combination of fecal indicator and sample type

using Bayesian multilevel models with compound-varying intercepts. Censored linear regression

was used to estimate the intervention impact on the $\log_{10}$ concentration of non-specific indicators

and the effect of the intervention on human-associated indicator prevalence was estimated using

logistic regression and the prevalence odds ratio (POR) as the measure of effect.[58,59] Models

were fit with the package **brms** in **R** version 4.0.2 using 1500 warmup and 1000 sampling

iterations on four chains (see SI for prior distributions).[58,60] Estimates of the intervention effect

were summarized by the mean and central 95% CI of the resulting 4000 posterior draws.

Adjusted models included variables for selected compound, household, meteorological,

and sample characteristics. Compound population density, presence of domestic animals, and

asset-based household wealth scores were derived from household and compound surveys

administered during each study phase.[42,61] Previous day mean temperature and seven-day

antecedent rainfall were drawn from daily summary records for a local weather station. For

stored water samples, we considered whether the storage container was covered and if the mouth

was wide enough to admit hands. The surface material was considered for food surface swabs,

and for soil samples we accounted for sun exposure and visibly wet soil surfaces. Covariate data

sources and processing have been described previously.[26,42]

Conditional probability analysis

Both HF183 and Mnif were previously found to frequently misdiagnose human feces in

our study area.[26] An indicator's diagnostic accuracy is described by its sensitivity (Se), the

probability of detecting the indicator when contamination is present, and specificity (Sp), the

probability of not detecting the indicator when contamination is absent. The probability that a

positive sample is contaminated depends on the marker sensitivity and specificity and the

prevalence of human fecal contamination. This marginal probability of contamination can be

approximated as the frequency of indicator detection among all samples to explore indicator

reliability in a specific study.[31] We assessed the probability that human feces were present in an

environmental sample in which HF183 or Mnif was detected using Bayes' Theorem and the local

sensitivity and specificity of the two markers (see SI).[34–36] We calculated the conditional

probability of contamination for HF183 and Mnif separately and for each combination of the two

indicators by sample type. The marginal probability of contamination was approximated as the

detection frequency of HF183 among all samples of a given type.

Accounting for diagnostic accuracy

Fecal indicator measurements are used as proxies for unobserved fecal contamination to

estimate its prevalence and associations of interest, such as the effects of mitigation practices.

This approach is vulnerable to measurement error, illustrated by the limited diagnostic accuracy

of many host-associated fecal indicators.[16] Bias due to inaccurate diagnostic tests can be

mitigated by incorporating external information on the sensitivity and specificity of the test.[62]

The expected detection frequency, p, of a test with sensitivity Se and specificity Sp is given by

$$p = Se \times \pi + (1 - Sp)(1 - \pi) \tag{1}$$

for an underlying condition with prevalence π.[62,63] We adapted the approach of Gelman and

Carpenter to estimate the intervention effect on human fecal contamination prevalence from

observations of human-associated fecal indicators by incorporating external information on

indicator performance within a Bayesian hierarchical framework.[63] We included the product-

term representation of the DID estimator and other covariates as linear predictors of the

prevalence log-odds. Assuming indicator detection in the ith of n samples, y_i, was Bernoulli-

distributed with probability p_i, where p_i was related to the prevalence as shown in Equation (**1**),

the accuracy-adjusted prevalence model was

$$y_i \sim Bernoulli(p_i)$$

$$p_i = Se \times \pi_i + (1 - Sp)(1 - \pi_i) \tag{2}$$

$$logit(\pi_i) = \beta^0 + \beta^P P_i + \beta^T T_i + \beta^{DID} P_i \times T_i + \boldsymbol{X}_i \boldsymbol{\gamma}$$

where β^0 is the intercept; β^P, β^T, and β^{DID} are the parameters corresponding to indicators for

study phase (P), treatment group (T), and their product; and $\boldsymbol{\gamma}$ is a $p \times 1$ vector of regression

coefficients corresponding to the p additional covariates in the $n \times p$ matrix $\boldsymbol{X}$.

We fit three models that differed by definition of Se and Sp. In the simplest case (Model

1), we assumed a perfectly accurate test with $Se = Sp = 1$, thus $p = \pi$. The second model

(Model 2) incorporated observations from the local validation analysis by assuming

$$y^{Se} \sim binomial(n^{Se}, Se)$$

$$y^{Sp} \sim binomial(n^{Sp}, Sp) \tag{3}$$

241   for y^{Se} positive results in n^{Se} human fecal samples and y^{Sp} negative results in n^{Sp} non-human

242   fecal samples. Because our validation sample set was small and performance estimates vary

243   widely between studies, we fit a third model (Model 3) featuring a meta-analysis of indicator

244   sensitivity and specificity (see SI). We assumed the log-odds of the sensitivity in the kth study,

245    $Se_{[k]}$, were normally distributed with mean μ^{Se} and SD σ^{Se}, such that

$$y_{[k]}^{Se} \sim binomial\left(n_{[k]}^{Se}, Se_{[k]}\right)$$

$$logit\left(Se_{[k]}\right) \sim normal(\mu^{Se}, \sigma^{Se}) \tag{4}$$

with an equivalent structure for the specificity. We assigned $k = 1$ to our local validation study,

using $Se_{[1]}$ and $Sp_{[1]}$ as the values of Se and Sp in Equation (**2**).[26,63] This emphasized the local

performance data while allowing information from other settings to influence the estimates

through partial pooling, with the extent of pooling learned from the data (expressed through σ^{Se}

and σ^{Sp}).[59]

Modeling latent human fecal contamination

Fecal contamination can be understood as a latent environmental condition for which

fecal indicators serve as imperfect diagnostic tests.[64,65] Information from multiple fecal indicators

may be utilized by modeling each as arising from the same underlying contamination to

potentially improve inference. We extended the meta-analytic model (Model 3) to include

observations of both HF183 and Mnif in the same samples (Model 4), with separate detection

probabilities, p_i^{hf} and p_i^{mn}, obtained from indicator-specific sensitivity and specificity estimates

applied to the same underlying prevalence, π_i. As in previous models, the DID estimator and

other predictor variables were included in a linear model on the log-odds of π_i, assuming that

intervention effects and other covariates acted directly on the latent prevalence.

As environmental compartments from the same compound share sources of fecal

exposure, we extended the previous model to simultaneously consider observations of latrine

soil, household soil, and stored water in each compound (Model 5). Sample type-specific

prevalence variables, $\pi_i^{[type]}$, were modeled as linear deviations from a latent compound-level

prevalence π_j on the log-odds scale:

$$logit(\pi_i^{[type]}) = \alpha^{[type]} + X_i^{[type]}\gamma^{[type]} + logit(\pi_{[j]}^{comp})$$

$$logit\left(\pi_{[j]}^{comp}\right) = \alpha_{[j]}^{comp} + \beta^P P_{[j]} + \beta^T T_{[j]} + \beta^{DID} P_{[j]} \times T_{[j]} + X_{[j]}^{comp}\gamma^{comp}$$

$$\alpha_{[j]}^{comp} \sim normal(\mu^{comp}, \sigma^{comp})$$

$$\alpha^{[type]} \sim normal(0, \sigma^{type})$$

(5)

for sample i of a given $type$ (latrine soil, household soil, or stored water) in compound j, where

$\alpha_{[j]}^{comp}$ is a compound-varying intercept and $\alpha^{[type]}$ is a varying intercept by sample type.

Compound-level predictors, including the DID estimator terms, were placed on the compound-

prevalence log-odds.[63,66] Parameters for sample-level and meteorological predictors in $X_i^{[type]}$

were estimated separately for each sample type.

We coded each model in the probabilistic programming language **Stan** and fit the models

using the **RStan** interface with four chains of 1000 warmup and 1000 sampling iterations each,

for a total of 4000 posterior samples (see SI for Stan code and discussion of prior

distributions).[67,68] Models 1-3 were fit separately for HF183 and Mnif in each sample type

(latrine entrance soil, household entrance soil, and stored water), Model 4 was fit separately to

each sample type, and a single Model 5 fit was produced incorporating both indicators and all

sample types. In addition to the DID POR given by the product-term parameter, we used the

posterior predictive distribution to estimate the prevalence of human fecal contamination in each

stratum and to directly calculate DID on the probability scale.[59,69] Models were adjusted for the

same covariates as the DID regression models.

Ethical approval

This study was approved by the Institutional Review Board of the University of North

Carolina at Chapel Hill (IRB # 15-0963) and the associated health study was approved by the

Comité Nacional de Bioética para a Saúde (CNBS), Ministério da Saúde, Republic of

Mozambique (333/CNBS/14), the Ethics Committee of the London School of Hygiene and

Tropical Medicine (reference # 8345), and the Institutional Review Board of the Georgia

Institute of Technology (protocol # H15160). Environmental samples were only collected from

households with enrolled children for whom written, informed parental or guardian consent had

been given.

Results

Sample characteristics

We collected a total of 770 environmental samples from 507 unique locations at 139

households in 71 compounds. Samples were collected both pre- and post-intervention at 263

locations (52%), for a total of 526 paired samples and 244 unpaired samples (Table S2).

Characteristics expected to confound the relationship between sanitation and fecal contamination

were largely similar between treatment arms during each study phase (Table 1). Cumulative

precipitation was higher on average in intervention compounds at baseline and in control

compounds at follow-up. Water storage containers were also more frequently covered in

intervention (75%) than control households (57%) at baseline, though the majority of containers

were covered in all strata. Soil surfaces were more often visibly wet in control households (51%)

than intervention (33%) at follow-up, both of which were lower than at baseline (57% and 48%,

respectively). Most food preparation surfaces were plastic, though more often so in control

households during both study phases. A higher percentage of compounds from both treatment

arms reported owning domestic animals at follow-up (80–88%) than baseline (47–68%), which

may be related to differences in the questionnaire between survey phases. Median household

wealth was 40–45 on a 100-point index, with higher variance among controls at follow-up.

Median compound population density ranged from 5.5–8.1 residents/100 m^2.

Table 1. Characteristics of Maputo Sanitation study compounds and households selected for environmental sampling, samples
collected, and sampling dates, stratified by study phase and treatment arm

			before				after			
			control		intervention		control		intervention	
characteristic	level	metric	N	summary	N	summary	N	summary	N	summary
animals present	compound	n (%)	32	15 (47)	25	17 (68)	30	24 (80)	34	30 (88)
population density (persons/100 m^2)	compound	median (IQR^a)	29	5.5 (3.5)	23	8.1 (5.9)	28	5.9 (4.8)	33	6.7 (4.6)
wealth index (0 - 100)	household	median (IQR)	51	43 (12)	40	43 (12)	55	45 (19)	52	44 (14)
previous day mean temperature (°C)	date	median (IQR)	19	21 (2)	16	20 (2)	17	20 (1)	17	21 (3)
seven-day cumulative precipitation (mm)	date	median (IQR)	19	9 (3)	16	14 (3)	17	13 (39)	17	7 (0)
water container covered	sample	n (%)	44	25 (57)	28	21 (75)	38	21 (55)	47	30 (64)
narrow-mouth water container	sample	n (%)	44	13 (30)	28	10 (36)	38	13 (34)	47	14 (30)
plastic food surface material	sample	n (%)	34	30 (88)	23	18 (78)	29	26 (90)	36	29 (81)
shaded latrine soil	sample	n (%)	32	24 (75)	17	12 (71)	30	25 (83)	30	22 (73)
shaded household soil	sample	n (%)	42	31 (74)	28	24 (86)	35	32 (91)	39	31 (79)
wet latrine soil surface	sample	n (%)	32	20 (62)	17	13 (76)	30	18 (60)	30	21 (70)
wet household soil surface	sample	n (%)	42	24 (57)	27	13 (48)	35	18 (51)	39	13 (33)
latrine soil moisture (%)	sample	median (IQR)	33	9.8 (9.8)	23	8.4 (7.2)	30	10.0 (7.9)	30	8.7 (8.3)
household soil moisture (%)	sample	median (IQR)	49	9.9 (8.6)	35	6.9 (6.1)	47	7.8 (5.4)	43	5.4 (5.9)

310 [a] interquartile range
311

Fecal indicator occurrence

At least one fecal indicator was detected in 94% of samples (720/770) and *E. coli* was detected in 718 samples: by culture in 81% (611/755) and by qPCR in 86% (655/763). Mean cEC concentrations were lower at follow-up for all sample types in both treatment arms, a pattern not observed for EC23S concentrations (Figure 1). Of the 763 samples tested for human-associated indicators, 28% (217) were positive for at least one human marker. Human-associated indicators were common in soils (23–65% prevalence, across treatment groups and study phases) but only HF183 was regularly detected in stored water (10–22%) and both indicators were rare on food surfaces (0–9%). qPCR calibration curves (Table S5), detection limits (Table S6), and the results of laboratory quality controls are presented in the SI.

Bootstrap DID estimates suggest the intervention reduced EC23S concentrations on food preparation surfaces and HF183 prevalence in household soil but minimally impacted fecal indicator occurrence in other sample types (Table S7). Notably, HF183 prevalence in household soil was similar among intervention households in both study phases but increased among control compounds at follow-up. By contrast, model-based DID estimates, adjusted for potential confounding, were consistent with no intervention effect on food preparation surface EC23S concentration or household soil HF183 prevalence (Table S8). Adjusted models instead indicate the intervention reduced latrine soil concentrations of EC23S [mean difference: -1.2 (95% CI: -2.1, -0.30) $\log_{10}$ gc/dry g]. Although several sample characteristics were imbalanced between treatment arms and study phases (Table 1), estimates from models that adjusted for these variables were largely similar to the unadjusted models, with adjusted estimates marginally closer to the null in most cases (Table S8). EC23S concentrations in latrine soil were again the exception, with a substantially larger reduction obtained under the adjusted model than the

335   unadjusted estimate of -0.84 (95% CI: -1.6, -0.02) $\log_{10}$ gc/dry g. Due to low detection

336   frequency, models were not fit for either human marker on food surfaces or for Mnif in stored

337   water; source water samples were excluded from all analyses.[26]

338
339   **Figure 1. Bootstrap estimates of fecal indicator occurrence by study phase and treatment**
340   **arm. Points indicate mean $\log_{10}$ concentration for *E. coli* indicators and prevalence of**
341   **human-associated indicators, with bars presenting bootstrap 95% CIs.**
342
343

Conditional probability of human fecal contamination

The probability that a sample is contaminated with human feces given the detection of a

human indicator is a function of the indicator's sensitivity and specificity (Table S9) and the

prevalence of human contamination in the study environment. At 15% prevalence

(approximately the detection frequency of HF183 in stored water), the probability of human

contamination given a positive test was 26% for HF183 and 30% for Mnif. Only with prevalence

above 30–35% was detecting either indicator more likely than not to correctly diagnose human

fecal contamination. Combining test results from both indicators improved identification of

human contamination, increasing the probability of contamination to 45% when both markers

were positive and the prevalence was 15% (Figure 2). However, the two human markers

frequently disagreed when assessed in the same sample, conflicting in 44% of household soil,

43% of latrine soil, and 15% of stored water samples. Furthermore, at 44% prevalence (the

highest detection frequency for HF183, observed in latrine soils), there remained a >20% chance

that a sample positive for both indicators was not contaminated. Among lower-prevalence

sample types the conditional probability never reached 50%. Unless the background prevalence

in the study area was about 45% or greater, it is unlikely that the use of HF183 and Mnif reliably

identified human contamination in individual samples, particularly given the frequent

disagreement between the two markers.

362
Figure 2. Conditional probability of sample contamination with human feces given
detection status of both HF183 and Mnif for all values of human contamination prevalence.
Values of sensitivity and specificity were obtained using human and animal feces from the
study area, and are 64% and 67%, respectively, for HF183 and 71% and 70% for Mnif.
The dashed vertical lines indicate the HF183 detection frequency for each sample type to
illustrate relevant human contamination probabilities. FP: food preparation surfaces; SW:
stored water; HS: household entrance soil; LS: latrine entrance soil.
370
371

Prevalence of human fecal contamination

Posterior predictions from each of the five accuracy-adjusted models were used to estimate stratum-specific prevalence of human fecal contamination. To compare treatment assignments and study phases, we predicted prevalence for compounds with no animals or antecedent precipitation and the sample mean population density (7 persons/100 m^2), wealth score (46), and previous-day temperature (20.4 °C), in which soil surfaces were dry and shaded and water storage containers possessed wide, uncovered mouths. The prevalence estimates were notably imprecise; the 95% CI of the HF183 prevalence in post-treatment latrine soil ranged from 3% to 92% for Model 2 (Table 2). The 95% CI widths were similar for Model 1 and the bootstrap estimates but were substantially wider for the other four models, which accounted for FST marker sensitivity and specificity (see SI). The intervals narrowed somewhat when both indicators were considered (Model 4) and narrowed further when all sample types were incorporated (Model 5) but were still wider than the estimates that did not account for diagnostic accuracy.

Although we did not formally assess the pairwise differences between prevalence estimates, the wide and largely overlapping posterior predictive CIs indicate a limited ability to distinguish between prevalence estimates between different strata or models. The DID estimates on the probability scale were strongly consistent with no effect for all model specifications, which further suggests that the available data were insufficient to assess prevalence differences between strata. The corresponding prevalence odds ratio estimates obtained directly from the DID product term were likewise imprecise (Figure S1). Nonetheless, the model-based prevalence estimates were consistently more similar between study phase and treatment group than the corresponding bootstrap estimates. This trend was notable for Model 5, which assumed that time

and treatment effects acted directly on the compound-wide prevalence of human contamination,

thus affecting all three sample types equally. The compound-level prevalence estimates were

quite similar, particularly between study phases for the same treatment group: 27% (95% CI: 9-

52%) at baseline and 28% (9-53%) at follow-up for control compounds and 22% (6-50%) at

baseline and 22% (6-47%) at follow-up for intervention compounds. The corresponding

estimates for household soil were nearly identical to the compound-level estimates, with

somewhat higher estimates for latrine soil and lower for stored water. Although the physical

interpretation of this compound-level construct is uncertain, these estimates suggest that about a

quarter of compounds were measurably impacted by human fecal contamination, which was

unaffected by improvements to shared sanitation facilities.

Table 2. Bootstrap and adjusted model-based estimates human marker sensitivity and specificity, prevalence of human fecal contamination stratified by treatment arm and study phase, and effect of the sanitation intervention on human fecal contamination prevalence in soil and water from MapSan study compounds

	marker	sensitivity (95% CI)	specificity (95% CI)	N	prevalence estimate (95% CI)[a]				prevalence DID[b] (95% CI)
					control		intervention		
					before	after	before	after	
latrine soil									
bootstrap	HF183	1	1	116	0.33 (0.17, 0.50)	0.57 (0.39, 0.75)	0.43 (0.23, 0.64)	0.43 (0.26, 0.61)	-0.23 (-0.60, 0.14)
	Mnif	1	1	116	0.51 (0.35, 0.69)	0.50 (0.32, 0.68)	0.65 (0.45, 0.84)	0.36 (0.19, 0.54)	-0.27 (-0.63, 0.08)
model 1[c]	HF183	1	1	98	0.32 (0.17, 0.49)	0.42 (0.24, 0.60)	0.32 (0.15, 0.52)	0.37 (0.20, 0.57)	-0.04 (-0.22, 0.13)
	Mnif	1	1	98	0.44 (0.27, 0.63)	0.37 (0.20, 0.55)	0.43 (0.24, 0.65)	0.27 (0.13, 0.45)	-0.09 (-0.27, 0.07)
model 2[d]	HF183	0.60 (0.42, 0.79)	0.66 (0.53, 0.80)	98	0.38 (0.05, 0.88)	0.40 (0.05, 0.90)	0.38 (0.05, 0.89)	0.39 (0.03, 0.92)	-0.01 (-0.19, 0.18)
	Mnif	0.64 (0.47, 0.82)	0.66 (0.51, 0.81)	98	0.48 (0.09, 0.90)	0.44 (0.07, 0.90)	0.47 (0.07, 0.90)	0.39 (0.05, 0.92)	-0.04 (-0.25, 0.15)
model 3[e]	HF183	0.65 (0.45, 0.85)	0.68 (0.55, 0.82)	98	0.34 (0.05, 0.83)	0.37 (0.05, 0.85)	0.34 (0.04, 0.85)	0.36 (0.04, 0.88)	-0.01 (-0.19, 0.18)
	Mnif	0.70 (0.56, 0.83)	0.72 (0.58, 0.85)	98	0.49 (0.14, 0.84)	0.43 (0.11, 0.83)	0.47 (0.13, 0.84)	0.35 (0.07, 0.82)	-0.06 (-0.27, 0.13)
model 4[f]	HF183	0.64 (0.47, 0.82)	0.71 (0.57, 0.84)	98	0.39 (0.11, 0.73)	0.37 (0.10, 0.73)	0.37 (0.10, 0.74)	0.29 (0.07, 0.68)	-0.06 (-0.25, 0.11)
	Mnif	0.71 (0.58, 0.84)	0.71 (0.57, 0.84)						
model 5[g]	HF183	0.72 (0.57, 0.87)	0.85 (0.78, 0.91)	98	0.34 (0.12, 0.65)	0.35 (0.13, 0.65)	0.29 (0.08, 0.63)	0.28 (0.08, 0.60)	-0.02 (-0.17, 0.14)
	Mnif	0.71 (0.59, 0.83)	0.78 (0.68, 0.86)						
household soil									
bootstrap	HF183	1	1	176	0.17 (0.07, 0.28)	0.49 (0.35, 0.64)	0.36 (0.20, 0.52)	0.38 (0.24, 0.52)	-0.30 (-0.57, -0.01)
	Mnif	1	1	175	0.43 (0.30, 0.57)	0.25 (0.13, 0.39)	0.23 (0.09, 0.37)	0.24 (0.12, 0.38)	0.20 (-0.07, 0.46)
model 1	HF183	1	1	147	0.26 (0.15, 0.41)	0.43 (0.27, 0.58)	0.29 (0.15, 0.46)	0.41 (0.26, 0.58)	-0.04 (-0.21, 0.12)
	Mnif	1	1	146	0.37 (0.23, 0.52)	0.27 (0.15, 0.42)	0.27 (0.14, 0.43)	0.18 (0.09, 0.31)	0.01 (-0.13, 0.14)
model 2	HF183	0.60 (0.38, 0.80)	0.72 (0.61, 0.83)	147	0.28 (0.04, 0.73)	0.34 (0.03, 0.80)	0.27 (0.03, 0.74)	0.34 (0.02, 0.83)	0.00 (-0.18, 0.19)
	Mnif	0.57 (0.34, 0.80)	0.73 (0.63, 0.84)	146	0.30 (0.03, 0.78)	0.25 (0.02, 0.76)	0.25 (0.02, 0.77)	0.19 (0.01, 0.77)	-0.01 (-0.18, 0.14)
model 3	HF183	0.66 (0.43, 0.85)	0.74 (0.63, 0.85)	147	0.25 (0.04, 0.63)	0.33 (0.04, 0.74)	0.25 (0.03, 0.69)	0.33 (0.03, 0.80)	0.00 (-0.18, 0.20)
	Mnif	0.68 (0.50, 0.82)	0.76 (0.67, 0.86)	146	0.26 (0.03, 0.60)	0.20 (0.03, 0.52)	0.20 (0.02, 0.50)	0.13 (0.02, 0.40)	-0.01 (-0.16, 0.11)
model 4	HF183	0.69 (0.47, 0.87)	0.73 (0.63, 0.83)	146	0.20 (0.04, 0.44)	0.23 (0.03, 0.50)	0.15 (0.03, 0.37)	0.16 (0.02, 0.40)	-0.02 (-0.16, 0.11)
	Mnif	0.68 (0.51, 0.82)	0.75 (0.66, 0.84)						
model 5	HF183	0.72 (0.57, 0.87)	0.85 (0.78, 0.91)	146	0.26 (0.09, 0.49)	0.27 (0.10, 0.51)	0.22 (0.06, 0.47)	0.22 (0.06, 0.45)	-0.01 (-0.16, 0.12)
	Mnif	0.71 (0.59, 0.83)	0.78 (0.68, 0.86)						
stored water									
bootstrap	HF183	1	1	193	0.12 (0.04, 0.22)	0.10 (0.02, 0.20)	0.22 (0.10, 0.35)	0.19 (0.09, 0.30)	-0.01 (-0.21, 0.19)
model 1	HF183	1	1	170	0.23 (0.11, 0.38)	0.19 (0.09, 0.34)	0.28 (0.13, 0.48)	0.24 (0.11, 0.42)	0.00 (-0.14, 0.14)
model 2	HF183	0.60 (0.38, 0.81)	0.85 (0.78, 0.91)	170	0.15 (0.02, 0.40)	0.14 (0.02, 0.38)	0.17 (0.02, 0.47)	0.16 (0.01, 0.47)	0.00 (-0.13, 0.14)
model 3	HF183	0.67 (0.43, 0.85)	0.86 (0.79, 0.92)	170	0.15 (0.02, 0.38)	0.13 (0.02, 0.36)	0.17 (0.02, 0.45)	0.16 (0.02, 0.44)	0.00 (-0.13, 0.15)
model 5	HF183	0.72 (0.57, 0.87)	0.85 (0.78, 0.91)	169	0.19 (0.04, 0.43)	0.20 (0.03, 0.45)	0.16 (0.03, 0.40)	0.16 (0.02, 0.38)	-0.01 (-0.14, 0.11)
latent compound									

| model 5 | HF183 | 0.72 (0.57, 0.87) | 0.85 (0.78, 0.91) | 109 | 0.27 (0.09, 0.52) | 0.28 (0.09, 0.53) | 0.22 (0.06, 0.50) | 0.22 (0.06, 0.47) | -0.01 (-0.16, 0.13) |
| | Mnif | 0.71 (0.59, 0.83) | 0.78 (0.68, 0.86) | | | | | | |

[a] all models (excluding bootstrap estimates) were adjusted for population density, presence of animals, wealth score, temperature, antecedent precipitation, and sun exposure and surface wetness for soil samples and storage container mouth width and cover status for water samples

[b] difference-in-differences

[c] model 1: single sample type, single marker assuming perfect sensitivity and specificity

[d] model 2: single sample type, single marker with sensitivity and specificity from local validation study

[e] model 3: single sample type, single marker with meta-analytic sensitivity and specificity

[f] model 4: single sample type, two markers with meta-analytic sensitivity and specificity

[g] model 5: three sample types, two markers with meta-analytic sensitivity and specificity

Discussion

The provision of shared latrines reduced average soil concentrations of the molecular *E. coli* marker EC23S at latrine entrances by more than 1-$\log_{10}$ but did not have a comparable effect on culturable *E. coli*. EC23S latrine soil concentrations rose more in control compounds than they fell in intervention compounds, which under the parallel trends assumption is interpreted as a secular trend upwards that the intervention mitigated, for a much smaller absolute reduction than suggested by the DID estimate (Figure 1).[43] However, an opposite, downward trend was observed for all cEC concentrations. This discrepancy between two tests for the same organism complicates the interpretation of the relatively strong intervention effect estimated for EC23S. While the exact reasons for this discrepancy are yet to be determined, preliminary evidence from a related analysis suggests that the modified mTEC broth used for *E. coli* culture may have produced colonies of the same color and morphology for *Klebsiella* spp., which are commonly soil-derived and not specific to feces.[70] By contrast, the developers of EC23S reported 95% specificity to *E. coli* and cross reactions only with other *Escherichia* species, not *Klebsiella*.[46] Accordingly, EC23S potentially better reflected trends in fecal contamination, while cEC may have been confounded by soil microbes more susceptible to environmental conditions, such as the 2016 drought in southern Mozambique.[71]

A cluster-randomized trial in rural Bangladesh likewise found scant evidence of reductions in culturable *E. coli* concentrations from sanitation improvements.[72,73] Latrine provision also did not reduce the prevalence of pathogenic *E. coli* genes in soil, meaning neither culture- nor molecular-based measurements of soil *E. coli* were affected.[39] Other recent trials have not assessed intervention impacts on fecal contamination of soil, but several have evaluated contamination of drinking water, with some also testing child hands, food, or fomites.[15] As with

the present study, all found no effect of sanitation-only interventions on any environmental

compartment; combined water, sanitation, and hygiene interventions improved drinking water

quality in two studies.[13,14]

Measures of human-associated FST markers demonstrated that about a quarter of

compounds were impacted by human fecal contamination, with compound-level prevalence

estimates not statistically different at baseline and follow-up. Similarly, two cluster-randomized

trials, in India and Bangladesh, found no effect of rural sanitation interventions on the prevalence

of human-associated indicators in stored drinking water.[37,39] Both studies also assessed human

markers in mother and child hand rinse samples, which were not collected in this study. No

effect was observed for either hand type in India or on mother hands in Bangladesh, although the

human marker prevalence may have been reduced on child hands.[39]

Accounting for the diagnostic accuracy of FST markers revealed far greater uncertainty

about host-specific fecal contamination, both of individual samples and population averages,

than indicated by the raw indicator measurements. The relatively poor sensitivity and specificity

of both human markers in this setting severely limited their ability to identify specific samples

contaminated with human feces, but even moderate improvements in accuracy could

substantially increase FST marker utility. For example, a study in Singapore reported 75%

sensitivity and 89% specificity for HF183,[74] corresponding to a 55% chance a positive sample is

contaminated at 15% background prevalence and an 84% chance at 44% prevalence, compared

with 26% and 60%, respectively, for detection of HF183 in our study. Correcting for indicator

sensitivity and specificity to human-source contamination, coupled with the limited observations

of each sample type, yielded imprecise prevalence estimates that were consistent with both near

absence and almost omnipresence of contamination. While the reduced amplification efficiency

of HF183 (82%) may have contributed to its low sensitivity, it produced similar accuracy-

corrected estimates as Mnif, which was 95% efficient (Table S5). This imprecision inhibited

detecting intervention effects. The point estimates for the intervention effect were relatively

close to the null but the full posterior distributions were consistent with both large reductions and

substantial increases in prevalence attributable to the intervention. This analysis does not rule out

the possibility that sanitation improvements reduced the prevalence of human fecal

contamination. Rather, it strongly suggests that the tools used were inadequate, conveying too

little information to address the research question with an acceptable degree of confidence.

470   These limitations highlight the importance of conducting local validation studies for any

new FST application.[75] Accounting for diagnostic accuracy is unlikely to improve the strength or

precision of estimates, but may help mitigate overconfidence and overinterpretation by revealing

limitations of the available measurements. This practice could also be extended to account for

indicator sensitivity and specificity to strictly fecal targets, rather than environmental microbes

with non-fecal origins, although we lacked the appropriate data to implement such an analysis

for our two non-specific indicators, EC23S and cEC. As the diagnostic accuracy framework is

currently limited to binary outcomes, analysis of such high-prevalence indicators would benefit

from the development of analogous approaches for continuous outcomes. Given the

intermingling in low-income settings of humans and animals, and their gut microbiomes,

alternative FST targets such as mitochondrial DNA could prove more accurate.[76,77] Recent

technological advances also present opportunities for new approaches that might bypass the

limitations of the current FST paradigm, including portable, long-read sequencing platforms for

metagenomic-based source tracking and parallel PCR platforms that render simultaneous

analysis of multiple FST markers and comprehensive direct pathogen detection increasingly

feasible.[20,78–82] These technologies will also need to overcome the substantial variability, limited

analytical sensitivity, and matrix interference characteristic of environmental microbial

assessments.[16]

The low signal typical of environmental measurements suggests that study designs—

preferably longitudinal—that maximize observations on select pathways of greatest interest

should be prioritized to support more robust inference, regardless of analytical approach.[83] A

recent longitudinal analysis of *E. coli* concentrations in rural Bangladesh, collected at eight

timepoints over 2.5 years from 720 households, demonstrates the advantages of maximizing the

number of basic measurements across time. Although pooled estimates from certain sample

types achieved statistical significance, the sheer quantity of information available convincingly

demonstrated the lack of a physically meaningful sanitation intervention impacts on ambient

fecal contamination.[73]

Many have speculated that sanitation's apparent lack of effect may be due in part to

animal fecal contamination.[12,22] Animal feces often contain pathogens capable of infecting

humans and animal fecal biomass in domestic environments is estimated to far exceed that from

humans. [22,84–86] Inadequate management of child feces and fecal sludge, contamination of food

and water outside the home, and inadequate community-level drainage, solid waste, and

sanitation services all present potential pathways of continued contamination despite household

sanitation improvements.[24,87–92] Recognizing calls for "transformative" WASH to address these

multifarious hazards, sustained progress may require high standards of housing and public

services in addition to WASH improvements, necessitating multi-sectoral coordination and

financing.[12,93–95] Even small treatment effects may translate to positive economic benefits.[12]

Additionally, quality sanitation infrastructure can provide important benefits irrespective of

preventing pathogen exposure, particularly in crowded urban settlements.[96,97] For example,

previous research found users of MapSan intervention latrines and similar facilities in the same

neighborhoods reported reduced disgust and embarrassment about unhygienic conditions and

improved perceptions of security and privacy.[98] Based on the results of our study, we

recommend future research to understand the etiology and ecology of fecal pathogens in

domestic environments and beyond to help inform interventions needed to construct healthy

environments and to protect children's health.

Acknowledgment

This study was funded by the United States Agency for International Development under

Translating Research into Action, Cooperative Agreement No. GHS-A-00-09-00015-00, the Bill

and Melinda Gates Foundation grant OPP1137224, and the National Institute of Environmental

Health Sciences (T32ES007018). We gratefully acknowledge our implementing partner, Water

and Sanitation for the Urban Poor; the field research team at WE Consult; the laboratory staff at

the Mozambique National Institute of Health; and Olimpio Zavale, Olivia Ginn, and Anna

Stamatogiannakis for sampling support.

Supporting Information Available

Site selection criteria; samples collected; qPCR assay details; calibration curves;

detection limits; laboratory quality control; conditional probability; difference-in-differences

estimates; validation studies; diagnostic accuracy; accuracy-adjusted intervention effect

estimates; human fecal contamination prevalence estimates; prior distributions; model Stan code.

References

(1) Julian, T. R. Environmental Transmission of Diarrheal Pathogens in Low and Middle

Income Countries. *Environ. Sci. Process. Impacts* **2016**, *18* (8), 944–955.

https://doi.org/10.1039/C6EM00222F.

(2) Wagner, E. G.; Laniox, J. N. Excreta Disposal for Rural Areas and Small Communities.

Monogr. Ser. World Health Organ. **1958**, *39*, 1–182.

(3) Sclar, G. D.; Penakalapati, G.; Amato, H. K.; Garn, J. V.; Alexander, K.; Freeman, M. C.;

Boisson, S.; Medlicott, K. O.; Clasen, T. Assessing the Impact of Sanitation on Indicators

of Fecal Exposure along Principal Transmission Pathways: A Systematic Review. *Int. J.*

Hyg. Environ. Health **2016**, *219* (8), 709–723. https://doi.org/10.1016/j.ijheh.2016.09.021.

(4) Fuhrmeister, E. R.; Ercumen, A.; Pickering, A. J.; Jeanis, K. M.; Ahmed, M.; Brown, S.;

Arnold, B. F.; Hubbard, A. E.; Alam, M.; Sen, D.; Islam, S.; Kabir, M. H.; Kwong, L. H.;

Islam, M.; Unicomb, L.; Rahman, M.; Boehm, A. B.; Luby, S. P.; Colford, J. M.; Nelson,

K. L. Predictors of Enteric Pathogens in the Domestic Environment from Human and

Animal Sources in Rural Bangladesh. *Environ. Sci. Technol.* **2019**, *53* (17), 10023–10033.

https://doi.org/10.1021/acs.est.8b07192.

(5) Knee, J.; Sumner, T.; Adriano, Z.; Anderson, C.; Bush, F.; Capone, D.; Casmo, V.;

Holcomb, D. A.; Kolsky, P.; MacDougall, A.; Molotkova, E.; Braga, J. M.; Russo, C.;

Schmidt, W. P.; Stewart, J.; Zambrana, W.; Zuin, V.; Nalá, R.; Cumming, O.; Brown, J.

Effects of an Urban Sanitation Intervention on Childhood Enteric Infection and Diarrhea in

Maputo, Mozambique: A Controlled before-and-after Trial. *eLife* **2021**, *10*, e62278.

https://doi.org/10.7554/eLife.62278.

(6)   Null, C.; Stewart, C. P.; Pickering, A. J.; Dentz, H. N.; Arnold, B. F.; Arnold, C. D.;

Benjamin-Chung, J.; Clasen, T.; Dewey, K. G.; Fernald, L. C. H.; Hubbard, A. E.; Kariger,

P.; Lin, A.; Luby, S. P.; Mertens, A.; Njenga, S. M.; Nyambane, G.; Ram, P. K.; Colford, J.

M. Effects of Water Quality, Sanitation, Handwashing, and Nutritional Interventions on

Diarrhoea and Child Growth in Rural Kenya: A Cluster-Randomised Controlled Trial.

Lancet Glob. Health **2018**, *6* (3), e316–e329. https://doi.org/10.1016/S2214-

109X(18)30005-6.

(7)   Luby, S. P.; Rahman, M.; Arnold, B. F.; Unicomb, L.; Ashraf, S.; Winch, P. J.; Stewart, C.

P.; Begum, F.; Hussain, F.; Benjamin-Chung, J.; Leontsini, E.; Naser, A. M.; Parvez, S. M.;

Hubbard, A. E.; Lin, A.; Nizame, F. A.; Jannat, K.; Ercumen, A.; Ram, P. K.; Das, K. K.;

Abedin, J.; Clasen, T. F.; Dewey, K. G.; Fernald, L. C.; Null, C.; Ahmed, T.; Colford, J. M.

Effects of Water Quality, Sanitation, Handwashing, and Nutritional Interventions on

Diarrhoea and Child Growth in Rural Bangladesh: A Cluster Randomised Controlled Trial.

Lancet Glob. Health **2018**, *6* (3), e302–e315. https://doi.org/10.1016/S2214-

109X(17)30490-4.

(8)   Grembi, J. A.; Lin, A.; Karim, M. A.; Islam, M. O.; Miah, R.; Arnold, B. F.; McQuade, E.

T. R.; Ali, S.; Rahman, M. Z.; Hussain, Z.; Shoab, A. K.; Famida, S. L.; Hossen, M. S.;

Mutsuddi, P.; Rahman, M.; Unicomb, L.; Haque, R.; Taniuchi, M.; Liu, J.; Platts-Mills, J.

A.; Holmes, S. P.; Stewart, C. P.; Benjamin-Chung, J.; Colford, J. M.; Houpt, E. R.; Luby,

S. P. Effect of Water, Sanitation, Handwashing and Nutrition Interventions on

Enteropathogens in Children 14 Months Old: A Cluster-Randomized Controlled Trial in

Rural Bangladesh. *J. Infect. Dis.* **2020**, jiaa549. https://doi.org/10.1093/infdis/jiaa549.

(9)   Humphrey, J. H.; Mbuya, M. N. N.; Ntozini, R.; Moulton, L. H.; Stoltzfus, R. J.; Tavengwa,

N. V.; Mutasa, K.; Majo, F.; Mutasa, B.; Mangwadu, G.; Chasokela, C. M.; Chigumira, A.;

Chasekwa, B.; Smith, L. E.; Tielsch, J. M.; Jones, A. D.; Manges, A. R.; Maluccio, J. A.;

Prendergast, A. J. Independent and Combined Effects of Improved Water, Sanitation, and

Hygiene, and Improved Complementary Feeding, on Child Stunting and Anaemia in Rural

Zimbabwe: A Cluster-Randomised Trial. *Lancet Glob. Health* **2019**, *7* (1), e132–e147.

https://doi.org/10.1016/S2214-109X(18)30374-7.

(10) Pickering, A. J.; Crider, Y.; Sultana, S.; Swarthout, J.; Goddard, F. G.; Anjerul Islam, S.;

Sen, S.; Ayyagari, R.; Luby, S. P. Effect of In-Line Drinking Water Chlorination at the

Point of Collection on Child Diarrhoea in Urban Bangladesh: A Double-Blind, Cluster-

Randomised Controlled Trial. *Lancet Glob. Health* **2019**, *7* (9), e1247–e1256.

https://doi.org/10.1016/S2214-109X(19)30315-8.

(11) Reese, H.; Routray, P.; Torondel, B.; Sinharoy, S. S.; Mishra, S.; Freeman, M. C.; Chang,

H. H.; Clasen, T. Assessing Longer-Term Effectiveness of a Combined Household-Level

Piped Water and Sanitation Intervention on Child Diarrhoea, Acute Respiratory Infection,

Soil-Transmitted Helminth Infection and Nutritional Status: A Matched Cohort Study in

Rural Odisha,. *Int. J. Epidemiol.* **2019**, *48* (6), 1757–1767.

https://doi.org/10.1093/ije/dyz157.

(12) Whittington, D.; Radin, M.; Jeuland, M. Evidence-Based Policy Analysis? The Strange

Case of the Randomized Controlled Trials of Community-Led Total Sanitation. *Oxf. Rev.*

Econ. Policy **2020**, *36* (1), 191–221. https://doi.org/10.1093/oxrep/grz029.

(13) Ercumen, A.; Mertens, A.; Arnold, B. F.; Benjamin-Chung, J.; Hubbard, A. E.; Ahmed, M.

A.; Kabir, M. H.; Rahman Khalil, Md. M.; Kumar, A.; Rahman, Md. S.; Parvez, S. M.;

Unicomb, L.; Rahman, M.; Ram, P. K.; Clasen, T.; Luby, S. P.; Colford, J. M. Effects of

Single and Combined Water, Sanitation and Handwashing Interventions on Fecal

Contamination in the Domestic Environment: A Cluster-Randomized Controlled Trial in

Rural Bangladesh. *Environ. Sci. Technol.* **2018**, *52* (21), 12078–12088.

https://doi.org/10.1021/acs.est.8b05153.

(14) Pickering, A. J.; Swarthout, J.; Mureithi, M.; Mboya, J.; Arnold, B. F.; Wolfe, M.; Dentz,

H. N.; Lin, A.; Arnold, C. D.; Rao, G.; Stewart, C. P.; Ram, P. K.; Clasen, T.; Colford, J.

M.; Null, C. Can Individual and Integrated Water, Sanitation, and Handwashing

Interventions Reduce Fecal Contamination in the Household Environment? Evidence from

the WASH Benefits Cluster-Randomized Trial in Rural Kenya. *bioRxiv* **2019**.

https://doi.org/10.1101/731992.

(15) Goddard, F.; Ban, R.; Barr, D. B.; Brown, J.; Cannon, J.; Colford, J. M.; Eisenberg, J. N. S.;

Ercumen, A.; Petach, H.; Freeman, M. C.; Levy, K.; Luby, S. P.; moe, christine; Pickering,

A. J.; Sarnat, J. A.; Stewart, J. R.; Thomas, E. A.; Taniuchi, M.; Clasen, T. F. Measuring

Environmental Exposure to Enteric Pathogens in Low-Income Settings: Review and

Recommendations of an Interdisciplinary Working Group. *Environ. Sci. Technol.* **2020**,

acs.est.0c02421. https://doi.org/10.1021/acs.est.0c02421.

(16) Holcomb, D. A.; Stewart, J. R. Microbial Indicators of Fecal Pollution: Recent Progress and

Challenges in Assessing Water Quality. *Curr. Environ. Health Rep.* **2020**, *7*, 311–324.

https://doi.org/10.1007/s40572-020-00278-1.

(17) Harris, A. R.; Pickering, A. J.; Harris, M.; Doza, S.; Islam, Md. S.; Unicomb, L.; Luby, S.;

Davis, J.; Boehm, A. B. Ruminants Contribute Fecal Contamination to the Urban

Household Environment in Dhaka, Bangladesh. *Environ. Sci. Technol.* **2016**, *50* (9), 4642–

4649. https://doi.org/10.1021/acs.est.5b06282.

(18) Boehm, A. B.; Wang, D.; Ercumen, A.; Shea, M.; Harris, A. R.; Shanks, O. C.; Kelty, C.;

Ahmed, A.; Mahmud, Z. H.; Arnold, B. F.; Chase, C.; Kullmann, C.; Colford, J. M.; Luby,

S. P.; Pickering, A. J. Occurrence of Host-Associated Fecal Markers on Child Hands,

Household Soil, and Drinking Water in Rural Bangladeshi Households. *Environ. Sci.*

Technol. Lett. **2016**, acs.estlett.6b00382. https://doi.org/10.1021/acs.estlett.6b00382.

(19) Ercumen, A.; Pickering, A. J.; Kwong, L. H.; Arnold, B. F.; Parvez, S. M.; Alam, M.; Sen,

D.; Islam, S.; Kullmann, C.; Chase, C.; Ahmed, R.; Unicomb, L.; Luby, S. P.; Colford, J.

M. Animal Feces Contribute to Domestic Fecal Contamination: Evidence from E. Coli

Measured in Water, Hands, Food, Flies, and Soil in Bangladesh. *Environ. Sci. Technol.*

2017, *51* (15), 8725–8734. https://doi.org/10.1021/acs.est.7b01710.

(20) Baker, K. K.; Senesac, R.; Sewell, D.; Sen Gupta, A.; Cumming, O.; Mumma, J. Fecal

Fingerprints of Enteric Pathogen Contamination in Public Environments of Kisumu, Kenya,

Associated with Human Sanitation Conditions and Domestic Animals. *Environ. Sci.*

Technol. **2018**, *52* (18), 10263–10274. https://doi.org/10.1021/acs.est.8b01528.

(21) Wardrop, N. A.; Hill, A. G.; Dzodzomenyo, M.; Aryeetey, G.; Wright, J. A. Livestock

Ownership and Microbial Contamination of Drinking-Water: Evidence from Nationally

Representative Household Surveys in Ghana, Nepal and Bangladesh. *Int. J. Hyg. Environ.*

Health **2018**, *221* (1), 33–40. https://doi.org/10.1016/j.ijheh.2017.09.014.

(22) Prendergast, A. J.; Gharpure, R.; Mor, S.; Viney, M.; Dube, K.; Lello, J.; Berger, C.; Siwila,

J.; Joyeux, M.; Hodobo, T.; Hurt, L.; Brown, T.; Hoto, P.; Tavengwa, N.; Mutasa, K.;

Craddock, S.; Chasekwa, B.; Robertson, R. C.; Evans, C.; Chidhanguro, D.; Mutasa, B.;

Majo, F.; Smith, L. E.; Hirai, M.; Ntozini, R.; Humphrey, J. H.; Berendes, D. Putting the

"A" into WaSH: A Call for Integrated Management of Water, Animals, Sanitation, and

Hygiene. *Lancet Planet. Health* **2019**, *3* (8), e336–e337. https://doi.org/10.1016/S2542-

5196(19)30129-9.

(23) Navab-Daneshmand, T.; Friedrich, M. N. D.; Gächter, M.; Montealegre, M. C.; Mlambo, L.

S.; Nhiwatiwa, T.; Mosler, H.-J.; Julian, T. R. Escherichia Coli Contamination across

Multiple Environmental Compartments (Soil, Hands, Drinking Water, and Handwashing

Water) in Urban Harare: Correlations and Risk Factors. *Am. J. Trop. Med. Hyg.* **2018**, *98*

(3). https://doi.org/10.4269/ajtmh.17-0521.

(24) Robb, K.; Null, C.; Armah, G.; Teunis, P.; Moe, C. L.; Yakubu, H. Assessment of Fecal

Exposure Pathways in Low-Income Urban Neighborhoods in Accra, Ghana: Rationale,

Design, Methods, and Key Findings of the SaniPath Study. *Am. J. Trop. Med. Hyg.* **2017**,

1–26. https://doi.org/10.4269/ajtmh.16-0508.

(25) Pickering, A. J.; Julian, T. R.; Marks, S. J.; Mattioli, M. C.; Boehm, A. B.; Schwab, K. J.;

Davis, J. Fecal Contamination and Diarrheal Pathogens on Surfaces and in Soils among

Tanzanian Households with and without Improved Sanitation. *Environ. Sci. Technol.* **2012**,

46 (11), 5736–5743. https://doi.org/10.1021/es300022c.

(26) Holcomb, D. A.; Knee, J.; Sumner, T.; Adriano, Z.; de Bruijn, E.; Nalá, R.; Cumming, O.;

Brown, J.; Stewart, J. R. Human Fecal Contamination of Water, Soil, and Surfaces in

Households Sharing Poor-Quality Sanitation Facilities in Maputo, Mozambique. *Int. J.*

Hyg. Environ. Health **2020**, *226*, 113496. https://doi.org/10.1016/j.ijheh.2020.113496.

(27) Ishii, S.; Sadowsky, M. J. Escherichia Coli in the Environment: Implications for Water

Quality and Human Health. *Microbes Environ.* **2008**, *23* (2), 101–108.

https://doi.org/10.1264/jsme2.23.101.

(28) Byappanahalli, M. N.; Nevers, M. B.; Korajkic, A.; Staley, Z. R.; Harwood, V. J.

Enterococci in the Environment. *Microbiol. Mol. Biol. Rev.* **2012**, *76* (4), 685–706.

https://doi.org/10.1128/MMBR.00023-12.

(29) Price, H. D.; Adams, E. A.; Nkwanda, P. D.; Mkandawire, T. W.; Quilliam, R. S. Daily

Changes in Household Water Access and Quality in Urban Slums Undermine Global Safe

Water Monitoring Programmes. *Int. J. Hyg. Environ. Health* **2021**, *231*, 113632.

https://doi.org/10.1016/j.ijheh.2020.113632.

(30) Wyer, M. D.; Kay, D.; Morgan, H.; Naylor, S.; Clark, S.; Watkins, J.; Davies, C. M.;

Francis, C.; Osborn, H.; Bennett, S. Within-Day Variability in Microbial Concentrations at

a UK Designated Bathing Water: Implications for Regulatory Monitoring and the

Application of Predictive Modelling Based on Historical Compliance Data. *Water Res. X*

2018. https://doi.org/10.1016/j.wroa.2018.10.003.

(31) Jenkins, M. W.; Tiwari, S.; Lorente, M.; Gichaba, C. M.; Wuertz, S. Identifying Human and

Livestock Sources of Fecal Contamination in Kenya with Host-Specific Bacteroidales

Assays. *Water Res.* **2009**, *43* (19), 4956–4966.

https://doi.org/10.1016/j.watres.2009.07.028.

(32) Odagiri, M.; Schriewer, A.; Hanley, K.; Wuertz, S.; Misra, P. R.; Panigrahi, P.; Jenkins, M.

W. Validation of Bacteroidales Quantitative PCR Assays Targeting Human and Animal

Fecal Contamination in the Public and Domestic Domains in India. *Sci. Total Environ.*

2015, *502*, 462–470. https://doi.org/10.1016/j.scitotenv.2014.09.040.

(33) Feng, S.; Ahmed, W.; McLellan, S. L. Ecological and Technical Mechanisms for Cross-

Reaction of Human Fecal Indicators with Animal Hosts. *Appl. Environ. Microbiol.* **2019**,

86 (5). https://doi.org/10.1128/AEM.02319-19.

(34) Johnston, C.; Byappanahalli, M. N.; Gibson, J. M.; Ufnar, J. a.; Whitman, R. L.; Stewart, J.

R. Probabilistic Analysis Showing That a Combination of Bacteroides and

Methanobrevibacter Source Tracking Markers Is Effective for Identifying Waters

Contaminated by Human Fecal Pollution. *Environ. Sci. Technol.* **2013**, *47* (23), 13621–

13628. https://doi.org/10.1021/es403753k.

(35) Kildare, B. J.; Leutenegger, C. M.; McSwain, B. S.; Bambic, D. G.; Rajal, V. B.; Wuertz, S.

16S RRNA-Based Assays for Quantitative Detection of Universal, Human-, Cow-, and

Dog-Specific Fecal Bacteroidales: A Bayesian Approach. *Water Res.* **2007**, *41* (16), 3701–

3715. https://doi.org/10.1016/j.watres.2007.06.037.

(36) Curtis, K.; Gonzalez, R. A. Integrating Bayesian Analysis and Cumulative Probability

Generates High Confidence Using a Single Microbial Source Tracking Marker. *Environ.*

Sci. Technol. **2019**, acs.est.9b03843. https://doi.org/10.1021/acs.est.9b03843.

(37) Odagiri, M.; Schriewer, A.; Daniels, M. E.; Wuertz, S.; Smith, W. A.; Clasen, T.; Schmidt,

W.; Jin, Y.; Torondel, B.; Misra, P. R.; Panigrahi, P.; Jenkins, M. W. Human Fecal and

Pathogen Exposure Pathways in Rural Indian Villages and the Effect of Increased Latrine

Coverage. *Water Res.* **2016**, *100*, 232–244. https://doi.org/10.1016/j.watres.2016.05.015.

(38) Schriewer, A.; Odagiri, M.; Wuertz, S.; Misra, P. R.; Panigrahi, P.; Clasen, T.; Jenkins, M.

W. Human and Animal Fecal Contamination of Community Water Sources, Stored

Drinking Water and Hands in Rural India Measured with Validated Microbial Source

Tracking Assays. *Am. J. Trop. Med. Hyg.* **2015**, *93* (3), 509–516.

https://doi.org/10.4269/ajtmh.14-0824.

(39) Fuhrmeister, E. R.; Ercumen, A.; Pickering, A. J.; Jeanis, K. M.; Crider, Y.; Ahmed, M.;

Brown, S.; Alam, M.; Sen, D.; Islam, S.; Kabir, M. H.; Islam, M.; Rahman, M.; Kwong, L.

H.; Arnold, B. F.; Luby, S. P.; Colford, J. M.; Nelson, K. L. Effect of Sanitation

Improvements on Pathogens and Microbial Source Tracking Markers in the Rural

Bangladeshi Household Environment. *Environ. Sci. Technol.* **2020**, acs.est.9b04835.

https://doi.org/10.1021/acs.est.9b04835.

(40) Brown, J.; Cumming, O.; Bartram, J.; Cairncross, S.; Ensink, J.; Holcomb, D.; Knee, J.;

Kolsky, P.; Liang, K.; Liang, S.; Nala, R.; Norman, G.; Rheingans, R.; Stewart, J.; Zavale,

O.; Zuin, V.; Schmidt, W.-P. A Controlled, before-and-after Trial of an Urban Sanitation

Intervention to Reduce Enteric Infections in Children: Research Protocol for the Maputo

Sanitation (MapSan) Study, Mozambique. *BMJ Open* **2015**, *5* (6), e008215–e008215.

https://doi.org/10.1136/bmjopen-2015-008215.

(41) Devamani, C.; Norman, G.; Schmidt, W. A Simple Microbiological Tool to Evaluate the

Effect of Environmental Health Interventions on Hand Contamination. *Int. J. Environ. Res.*

Public. Health **2014**, *11* (11), 11846–11859. https://doi.org/10.3390/ijerph111111846.

(42) Knee, J.; Sumner, T.; Adriano, Z.; Berendes, D.; de Bruijn, E.; Schmidt, W.; Nalá, R.;

Cumming, O.; Brown, J. Risk Factors for Childhood Enteric Infection in Urban Maputo,

Mozambique: A Cross-Sectional Study. *PLoS Negl. Trop. Dis.* **2018**, *12* (11), e0006956.

https://doi.org/10.1371/journal.pntd.0006956.

(43) Schmidt, W. Randomised and Non-Randomised Studies to Estimate the Effect of

Community-Level Public Health Interventions: Definitions and Methodological

Considerations. *Emerg. Themes Epidemiol.* **2017**, *14* (1), 9. https://doi.org/10.1186/s12982-

017-0063-5.

(44) Capone, D.; Adriano, Z.; Berendes, D.; Cumming, O.; Dreibelbis, R.; Holcomb, D. A.;

Knee, J.; Ross, I.; Brown, J. A Localized Sanitation Status Index as a Proxy for Fecal

Contamination in Urban Maputo, Mozambique. *PLOS ONE* **2019**, *14* (10), e0224333.

https://doi.org/10.1371/journal.pone.0224333.

(45) USEPA. *Method 1603: Escherichia Coli (E.Coli) in Water by Membrane Filtration Using*

Modified Membrane-Thermotolerant Escherichia Coli Agar; 2009; p 13.

(46) Chern, E. C.; Siefring, S.; Paar, J.; Doolittle, M.; Haugland, R. A. Comparison of

Quantitative PCR Assays for Escherichia Coli Targeting Ribosomal RNA and Single Copy

Genes. *Lett. Appl. Microbiol.* **2011**, *52* (3), 298–306. https://doi.org/10.1111/j.1472-

765X.2010.03001.x.

(47) Green, H. C.; Haugland, R. A.; Varma, M.; Millen, H. T.; Borchardt, M. A.; Field, K. G.;

Walters, W. A.; Knight, R.; Sivaganesan, M.; Kelty, C. A.; Shanks, O. C. Improved HF183

Quantitative Real-Time PCR Assay for Characterization of Human Fecal Pollution in

Ambient Surface Water Samples. *Appl. Environ. Microbiol.* **2014**, *80* (10), 3086–3094.

https://doi.org/10.1128/AEM.04137-13.

(48) Johnston, C.; Ufnar, J. a; Griffith, J. F.; Gooch, J. a; Stewart, J. R. A Real-Time QPCR

Assay for the Detection of the NifH Gene of Methanobrevibacter Smithii, a Potential

Indicator of Sewage Pollution. *J. Appl. Microbiol.* **2010**, *109* (6), 1946–1956.

https://doi.org/10.1111/j.1365-2672.2010.04824.x.

(49) Nutz, S.; Döll, K.; Karlovsky, P. Determination of the LOQ in Real-Time PCR by Receiver

Operating Characteristic Curve Analysis: Application to QPCR Assays for Fusarium

Verticillioides and F. Proliferatum. *Anal. Bioanal. Chem.* **2011**, *401* (2), 717–726.

https://doi.org/10.1007/s00216-011-5089-x.

(50) Haugland, R. a; Siefring, S. C.; Wymer, L. J.; Brenner, K. P.; Dufour, A. P. Comparison of

Enterococcus Measurements in Freshwater at Two Recreational Beaches by Quantitative

Polymerase Chain Reaction and Membrane Filter Culture Analysis. *Water Res.* **2005**, *39*

(4), 559–568. https://doi.org/10.1016/j.watres.2004.11.011.

(51) Haugland, R. A.; Siefring, S.; Lavender, J.; Varma, M. Influences of Sample Interference

and Interference Controls on Quantification of Enterococci Fecal Indicator Bacteria in

Surface Water Samples by the QPCR Method. *Water Res.* **2012**, *46* (18), 5989–6001.

https://doi.org/10.1016/j.watres.2012.08.017.

(52) Haugland, R. a; Varma, M.; Sivaganesan, M.; Kelty, C.; Peed, L.; Shanks, O. C. Evaluation

of Genetic Markers from the 16S RRNA Gene V2 Region for Use in Quantitative Detection

of Selected Bacteroidales Species and Human Fecal Waste by QPCR. *Syst. Appl. Microbiol.*

2010, *33* (6), 348–357. https://doi.org/10.1016/j.syapm.2010.06.001.

(53) Sivaganesan, M.; Haugland, R. a.; Chern, E. C.; Shanks, O. C. Improved Strategies and

Optimization of Calibration Models for Real-Time PCR Absolute Quantification. *Water*

Res. **2010**, *44* (16), 4726–4735. https://doi.org/10.1016/j.watres.2010.07.066.

(54) Wing, C.; Simon, K.; Bello-Gomez, R. A. Designing Difference in Difference Studies: Best

Practices for Public Health Policy Research. *Annu. Rev. Public Health* **2018**, *39* (1), 453–

469. https://doi.org/10.1146/annurev-publhealth-040617-013507.

(55) Abadie, A. Semiparametric Difference-in-Differences Estimators. *Rev. Econ. Stud.* **2005**,

72 (1), 1–19. https://doi.org/10.1111/0034-6527.00321.

(56) Gelman, A.; Greenland, S. Are Confidence Intervals Better Termed "Uncertainty

Intervals"? *BMJ* **2019**, l5381. https://doi.org/10.1136/bmj.l5381.

(57) Gelman, A.; Hill, J. *Data Analysis Using Regression and Multilevel/Hierarchical Models*;

Cambridge University Press, 2007. https://doi.org/10.2277/0521867061.

(58) Bürkner, P.-C. Brms : An R Package for Bayesian Multilevel Models Using Stan. *J. Stat.*

Softw. **2017**, *80* (1). https://doi.org/10.18637/jss.v080.i01.

(59) McElreath, R. *Statistical Rethinking: A Bayesian Course with Examples in R and Stan*, 2nd

ed.; CRC texts in statistical science; Taylor and Francis, CRC Press: Boca Raton, 2020.

(60) R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for

Statistical Computing: Vienna, Austria, 2020.

(61) Schreiner, M.; Nsthandoca, H.; Lory, D.; Dd, D.; Yy, M. M. *A Simple Poverty Scorecard*

for Mozambique; Microfinance.com, 2013.

(62) Rogan, W. J.; Gladen, B. Estimating Prevalence from the Results of a Screening Test. *Am.*

J. Epidemiol. **1978**, *107* (1), 71–76. https://doi.org/10.1093/oxfordjournals.aje.a112510.

(63) Gelman, A.; Carpenter, B. Bayesian Analysis of Tests with Unknown Specificity and

Sensitivity. *J. R. Stat. Soc. Ser. C Appl. Stat.* **2020**, rssc.12435.

https://doi.org/10.1111/rssc.12435.

(64) Sima, L. C.; Ng, R.; Elimelech, M. Modeling Risk Categories to Predict the Longitudinal

Prevalence of Childhood Diarrhea in Indonesia. *Am. J. Trop. Med. Hyg.* **2013**, *89* (5), 884–

891. https://doi.org/10.4269/ajtmh.12-0540.

(65) Divelbiss, D. W.; Boccelli, D. L.; Succop, P. A.; Oerther, D. B. Environmental Health and

Household Demographics Impacting Biosand Filter Maintenance and Diarrhea in

Guatemala: An Application of Structural Equation Modeling. *Environ. Sci. Technol.* **2013**,

47 (3), 1638–1645. https://doi.org/10.1021/es303624a.

(66) Ghitza, Y.; Gelman, A. Deep Interactions with MRP: Election Turnout and Voting Patterns

Among Small Electoral Subgroups. *Am. J. Polit. Sci.* **2013**, *57* (3), 762–776.

https://doi.org/10.1111/ajps.12004.

(67) Carpenter, B.; Gelman, A.; Hoffman, M. D.; Lee, D.; Goodrich, B.; Betancourt, M.;

Brubaker, M.; Guo, J.; Li, P.; Riddell, A. Stan: A Probabilistic Programming Language. *J.*

Stat. Softw. **2017**, *76* (1). https://doi.org/10.18637/jss.v076.i01.

(68) Stan Development Team. *RStan: The R Interface to Stan*; 2020.

(69) Gelman, A.; Pardoe, I. 2. Average Predictive Comparisons for Models with Nonlinearity,

Interactions, and Variance Components. *Sociol. Methodol.* **2007**, *37* (1), 23–51.

https://doi.org/10.1111/j.1467-9531.2007.00181.x.

(70) Dengo-Baloi, L.; Berendes, D.; Holcomb, D.; Knee, J.; Langa, J.; Magaia, A.; Nalá, R.;

Brown, J. Increased Antibiotic Resistance in the Community in Maputo, Mozambique:

Preliminary Results of Microbial Analysis of a Pathogen Rich Environment, 2018.

(71) Araneda-Cabrera, R. J.; Bermúdez, M.; Puertas, J. Assessment of the Performance of

Drought Indices for Explaining Crop Yield Variability at the National Scale:

Methodological Framework and Application to Mozambique. *Agric. Water Manag.* **2021**,

246, 106692. https://doi.org/10.1016/j.agwat.2020.106692.

(72) Ercumen, A.; Pickering, A. J.; Kwong, L. H.; Mertens, A.; Arnold, B. F.; Benjamin-Chung,

J.; Hubbard, A. E.; Alam, M.; Sen, D.; Islam, S.; Rahman, Md. Z.; Kullmann, C.; Chase,

C.; Ahmed, R.; Parvez, S. M.; Unicomb, L.; Rahman, M.; Ram, P. K.; Clasen, T.; Luby, S.

P.; Colford, J. M. Do Sanitation Improvements Reduce Fecal Contamination of Water,

Hands, Food, Soil, and Flies? Evidence from a Cluster-Randomized Controlled Trial in

Rural Bangladesh. *Environ. Sci. Technol.* **2018**, *52* (21), 12089–12097.

https://doi.org/10.1021/acs.est.8b02988.

(73) Contreras, J. D.; Islam, M.; Mertens, A.; Pickering, A. J.; Kwong, L. H.; Arnold, B. F.;

Benjamin-Chung, J.; Hubbard, A. E.; Alam, M.; Sen, D.; Islam, S.; Rahman, M.; Unicomb,

L.; Luby, S. P.; Colford, J. M.; Ercumen, A. Longitudinal Effects of a Sanitation

Intervention on Environmental Fecal Contamination in a Cluster-Randomized Controlled

Trial in Rural Bangladesh. *Environ. Sci. Technol.* **2021**.

https://doi.org/10.1021/acs.est.1c01114.

(74) Nshimyimana, J. P.; Cruz, M. C.; Thompson, R. J.; Wuertz, S. Bacteroidales Markers for

Microbial Source Tracking in Southeast Asia. *Water Res.* **2017**, *118*, 239–248.

https://doi.org/10.1016/j.watres.2017.04.027.

(75) Stewart, J. R.; Boehm, A. B.; Dubinsky, E. A.; Fong, T.-T.; Goodwin, K. D.; Griffith, J. F.;

Noble, R. T.; Shanks, O. C.; Vijayavel, K.; Weisberg, S. B. Recommendations Following a

Multi-Laboratory Comparison of Microbial Source Tracking Methods. *Water Res.* **2013**, *47*

(18), 6829–6838. https://doi.org/10.1016/j.watres.2013.04.063.

(76) Zhu, K.; Suttner, B.; Pickering, A.; Konstantinidis, K. T.; Brown, J. A Novel Droplet

Digital PCR Human MtDNA Assay for Fecal Source Tracking. *Water Res.* **2020**, 116085.

https://doi.org/10.1016/j.watres.2020.116085.

(77) Pehrsson, E. C.; Tsukayama, P.; Patel, S.; Mejía-Bautista, M.; Sosa-Soto, G.; Navarrete, K.

M.; Calderon, M.; Cabrera, L.; Hoyos-Arango, W.; Bertoli, M. T.; Berg, D. E.; Gilman, R.

H.; Dantas, G. Interconnected Microbiomes and Resistomes in Low-Income Human

Habitats. *Nature* **2016**, *533* (7602), 212–216. https://doi.org/10.1038/nature17672.

(78) Bauza, V.; Madadi, V. O.; Ocharo, R. M.; Nguyen, T. H.; Guest, J. S. Microbial Source

Tracking Using 16S RRNA Amplicon Sequencing Identifies Evidence of Widespread

Contamination from Young Children's Feces in an Urban Slum of Nairobi, Kenya.

Environ. Sci. Technol. **2019**, *53* (14), 8271–8281. https://doi.org/10.1021/acs.est.8b06583.

(79) Hu, Y. O. O.; Ndegwa, N.; Alneberg, J.; Johansson, S.; Logue, J. B.; Huss, M.; Käller, M.;

Lundeberg, J.; Fagerberg, J.; Andersson, A. F. Stationary and Portable Sequencing-Based

Approaches for Tracing Wastewater Contamination in Urban Stormwater Systems. *Sci.*

Rep. **2018**, *8* (1), 1–13. https://doi.org/10.1038/s41598-018-29920-7.

(80) Acharya, K.; Khanal, S.; Pantha, K.; Amatya, N.; Davenport, R. J.; Werner, D. A

Comparative Assessment of Conventional and Molecular Methods, Including MinION

Nanopore Sequencing, for Surveying Water Quality. *Sci. Rep.* **2019**, *9* (1), 15726.

https://doi.org/10.1038/s41598-019-51997-x.

(81) Capone, D.; Berendes, D.; Cumming, O.; Holcomb, D. A.; Knee, J.; Konstantinidis, K. T.;

Levy, K.; Nala, R.; Risk, B. B.; Brown, J. Impact of an Urban Sanitation Intervention on

Enteric Pathogen Detection in Soils. *Environ. Sci. Technol.* **2021**, *in press*.

(82) Li, X.; Harwood, V. J.; Nayak, B.; Staley, C.; Sadowsky, M. J.; Weidhaas, J. A Novel

Microbial Source Tracking Microarray for Pathogen Detection and Fecal Source

Identification in Environmental Systems. *Environ. Sci. Technol.* **2015**, *49* (12), 7319–7329.

https://doi.org/10.1021/acs.est.5b00980.

(83) Ercumen, A.; Arnold, B. F.; Naser, A. Mohd.; Unicomb, L.; Colford, J. M.; Luby, S. P.

Potential Sources of Bias in the Use of Escherichia Coli to Measure Waterborne Diarrhoea

Risk in Low-Income Settings. *Trop. Med. Int. Health* **2017**, *22* (1), 2–11.

https://doi.org/10.1111/tmi.12803.

(84) Delahoy, M. J.; Wodnik, B.; McAliley, L.; Penakalapati, G.; Swarthout, J.; Freeman, M. C.;

Levy, K. Pathogens Transmitted in Animal Feces in Low- and Middle-Income Countries.

Int. J. Hyg. Environ. Health **2018**. https://doi.org/10.1016/j.ijheh.2018.03.005.

(85) Penakalapati, G.; Swarthout, J.; Delahoy, M. J.; McAliley, L.; Wodnik, B.; Levy, K.;

Freeman, M. C. Exposure to Animal Feces and Human Health: A Systematic Review and

Proposed Research Priorities. *Environ. Sci. Technol.* **2017**, *51* (20), 11537–11552.

https://doi.org/10.1021/acs.est.7b02811.

(86) Berendes, D. M.; Yang, P. J.; Lai, A.; Hu, D.; Brown, J. Estimation of Global Recoverable

Human and Animal Faecal Biomass. *Nat. Sustain.* **2018**, *1* (11), 679–685.

https://doi.org/10.1038/s41893-018-0167-0.

(87) Berendes, D. M.; Leon, J. S.; Kirby, A. E.; Clennon, J. A.; Raj, S. J.; Yakubu, H.; Robb, K.

A.; Kartikeyan, A.; Hemavathy, P.; Gunasekaran, A.; Roy, S.; Ghale, B. C.; Kumar, J. S.;

Mohan, V. R.; Kang, G.; Moe, C. L. Associations between Open Drain Flooding and

Pediatric Enteric Infections in the MAL-ED Cohort in a Low-Income, Urban Neighborhood

in Vellore, India. *BMC Public Health* **2019**, *19* (1), 926. https://doi.org/10.1186/s12889-

019-7268-1.

(88) Berendes, D.; Kirby, A.; Clennon, J. A.; Raj, S.; Yakubu, H.; Leon, J.; Robb, K.;

Kartikeyan, A.; Hemavathy, P.; Gunasekaran, A.; Ghale, B.; Kumar, J. S.; Mohan, V. R.;

Kang, G.; Moe, C. The Influence of Household- and Community-Level Sanitation and

Fecal Sludge Management on Urban Fecal Contamination in Households and Drains and

Enteric Infection in Children. *Am. J. Trop. Med. Hyg.* **2017**, *96* (6).

https://doi.org/10.4269/ajtmh.16-0170.

(89) Harris, M.; Alzua, M. L.; Osbert, N.; Pickering, A. Community-Level Sanitation Coverage

More Strongly Associated with Child Growth and Household Drinking Water Quality than

Access to a Private Toilet in Rural Mali. *Environ. Sci. Technol.* **2017**, *51* (12), 7219–7227.

https://doi.org/10.1021/acs.est.7b00178.

(90) Huda, T. Md. N.; Schmidt, W.-P.; Pickering, A. J.; Unicomb, L.; Mahmud, Z. H.; Luby, S.

P.; Biran, A. Effect of Neighborhood Sanitation Coverage on Fecal Contamination of the

Household Environment in Rural Bangladesh. *Am. J. Trop. Med. Hyg.* **2019**.

https://doi.org/10.4269/ajtmh.16-0996.

(91) Bauza, V.; Majorin, F.; Routray, P.; Sclar, G. D.; Caruso, B. A.; Clasen, T. Child Feces

Management Practices and Fecal Contamination: A Cross-Sectional Study in Rural Odisha,

India. *Sci. Total Environ.* **2019**, 136169. https://doi.org/10.1016/j.scitotenv.2019.136169.

(92) Waller, A.; Lakhanpaul, M.; Godfrey, S.; Parikh, P. Multiple and Complex Links between

BabyWASH and Stunting: An Evidence Synthesis. *J. Water Sanit. Hyg. Dev.* **2020**,

washdev2020265. https://doi.org/10.2166/washdev.2020.265.

(93) Cumming, O.; Arnold, B. F.; Ban, R.; Clasen, T.; Esteves Mills, J.; Freeman, M. C.;

Gordon, B.; Guiteras, R.; Howard, G.; Hunter, P. R.; Johnston, R. B.; Pickering, A. J.;

Prendergast, A. J.; Prüss-Ustün, A.; Rosenboom, J. W.; Spears, D.; Sundberg, S.; Wolf, J.;

Null, C.; Luby, S. P.; Humphrey, J. H.; Colford, J. M. The Implications of Three Major

New Trials for the Effect of Water, Sanitation and Hygiene on Childhood Diarrhea and

Stunting: A Consensus Statement. *BMC Med.* **2019**, *17* (1), 173.

https://doi.org/10.1186/s12916-019-1410-x.

(94) Pickering, A. J.; Null, C.; Winch, P. J.; Mangwadu, G.; Arnold, B. F.; Prendergast, A. J.;

Njenga, S. M.; Rahman, M.; Ntozini, R.; Benjamin-Chung, J.; Stewart, C. P.; Huda, T. M.

N.; Moulton, L. H.; Colford, J. M.; Luby, S. P.; Humphrey, J. H. The WASH Benefits and

SHINE Trials: Interpretation of WASH Intervention Effects on Linear Growth and

Diarrhoea. *Lancet Glob. Health* **2019**, *7* (8), e1139–e1146. https://doi.org/10.1016/S2214-

109X(19)30268-2.

(95) Husseini, M.; Darboe, M. K.; Moore, S. E.; Nabwera, H. M.; Prentice, A. M. Thresholds of

Socio-Economic and Environmental Conditions Necessary to Escape from Childhood

Malnutrition: A Natural Experiment in Rural Gambia. *BMC Med.* **2018**, *16* (1), 199.

https://doi.org/10.1186/s12916-018-1179-3.

(96) Tidwell, J.; Chipungu, J.; Ross, I.; Antwi-Agyei, P.; Alam, M.-U.; Tumwebaze, I. K.;

Norman, G.; Cumming, O.; Simiyu, S. Where Shared Sanitation Is the Only Immediate

Option: A Research Agenda for Shared Sanitation in Densely Populated Low-Income

Urban Settings. *Am. J. Trop. Med. Hyg.* **2020**. https://doi.org/10.4269/ajtmh.20-0985.

(97) Sclar, G. D.; Penakalapati, G.; Caruso, B. A.; Rehfuess, E. A.; Garn, J. V.; Alexander, K.

T.; Freeman, M. C.; Boisson, S.; Medlicott, K.; Clasen, T. Exploring the Relationship

between Sanitation and Mental and Social Well-Being: A Systematic Review and

Qualitative Synthesis. *Soc. Sci. Med.* **2018**.

https://doi.org/10.1016/j.socscimed.2018.09.016.

(98) Shiras, T.; Cumming, O.; Brown, J.; Muneme, B.; Nala, R.; Dreibelbis, R. Shared Latrines

in Maputo, Mozambique: Exploring Emotional Well-Being and Psychosocial Stress. *BMC*

Int. Health Hum. Rights **2018**, *18* (1), 30. https://doi.org/10.1186/s12914-018-0169-z.