1 Long Title: Dynamic evolution of MLKL in primate and poxvirus genomes indicates
2 necroptosis is a critical, not auxiliary, countermeasure during infection
3
4 Short Title: Dynamic evolution of host and viral necroptotic factors
5
6

7 Suzette Palmer[1], Sruthi Chappidi[1], Chelsea Pinkham[1] and Dustin C. Hancks[1*]
8
9 [1]Department of Immunology, University of Texas Southwestern Medical Center, Dallas,
10 Texas, United States of America
11
12 * Corresponding author
13 E-mail: dustin.hancks@utsouthwestern.edu (DH)

Abstract

Pathogen infection triggers host innate defenses which can lead to the activation of regulated cell death (RCD) pathways such as apoptosis. Given a key role in immunity, apoptotic effectors are often counteracted by pathogen-encoded antagonists. Mounting evidence indicates that programmed necrosis, which is mediated by the RIPK3/MLKL axis and termed necroptosis, evolved as a countermeasure to pathogen-mediated inhibition of apoptotic signaling. However, whether this emerging inflammatory RCD pathway functions primarily as a "back-up" or fundamental response remains inconclusive. We hypothesized that if necroptosis is an instrumental defense, then its effectors should display specific signatures associated with pathogen conflict that are rare in combination: rapid evolution, viral homolog hereafter virolog, and induction by cytokines (e.g. interferons). Our rapid evolution analysis across the necroptosis pathway revealed: 1) strong signatures of positive selection for RIPK3 and MLKL in primate genomes and to a lesser extent DAI/ZBP1, 2) elevated rates of amino acid substitution on multiple surfaces including the RIPK3/MLKL binding interface and 3) evidence supporting a means of activating RIPK3 independent of homotypic RHIM domain interactions. Interestingly, a poxvirus MLKL homolog has recently been identified that acts as a RIPK3 pseudosubstrate. Our findings indicate that poxvirus MLKLs are also subject to similar but distinct volatile patterns of evolution comparable to host necroptotic factors. Specifically, viral MLKLs have undergone numerous gains and losses in poxvirus evolution with some species harboring three distinct copies. Furthermore, we confirm that MLKL can be induced by cytokines like interferon gamma. In summary, MLKL displays all three hallmarks of pivotal immune factors of which only

38   OAS1, but not other factors like cGAS, APOBEC3G, or PKR, exhibits. These data

39   support the hypothesis that over evolutionary time, necroptosis has served as a key

40   battleground during infection and is therefore, not an auxiliary response.

Summary

Regulated cell death (RCD), such as apoptosis, is a common host defense against invading pathogens. Necroptosis, an inflammatory RCD pathway, is thought to have emerged as an auxiliary response when other cell death pathways are suppressed by pathogens during infection. In our analyses, we have identified genetic changes in host and viral factors associated with necroptosis that display signatures of adaptation and may have served as evolutionary countermeasures to shape infection outcomes. Consistent with repeated targeting by pathogen-encoded inhibitors, we found robust signatures of rapid evolution for the essential catalysts of necroptosis, RIPK3 and MLKL. Notably, an evolutionary signature specific to RIPK3 for a domain shared with other necroptotic factors suggests an undefined means to trigger this host defense pathway. In contrast, poxviruses appear to circumvent this pathway by constantly altering the number and nature of factors they deploy to suppress necroptosis including a mimic of MLKL, which was stolen from infected cells. Collectively, our findings provide new insights into host and viral genetics that may influence infection outcomes and the factors shaping the ability of pathogens to infect and spread to new species. Furthermore, these data support the notion that necroptosis is a fundamental, not auxiliary, host response during infection.

Introduction

Regulated cell death (RCD) programs are pivotal host defense responses against pathogens [1-3]. Consequently, pathogens have evolved a myriad of strategies to counteract these responses [4, 5], which includes the activation of apoptosis. Apoptosis is a well-characterized, non-inflammatory form of RCD that was first classified by its distinct morphology that includes membrane blebbing, apoptotic bodies, and nuclear and cytoplasmic condensation [6]. Both extrinsic as well as intrinsic cues can trigger a signaling cascade that involves the activation of effector caspases to execute apoptosis (Fig 1A) [7] which was characterized, in large part, using viral proteins [8].

Recently, a RCD pathway defined as programmed necrosis, hereafter necroptosis, was discovered. Necroptosis is triggered when specific apoptotic signaling and effector functions are suppressed during pathogen infection. Given the conditions under which it is activated, necroptosis has often been considered a necessary "back-up" host response to pathogen-mediated inhibition of apoptosis (Fig 1B) [9, 10]. In contrast to apoptotic cell death, necroptosis is inflammatory and characterized by cell and organelle swelling followed by plasma membrane rupture [7, 11, 12]. Apoptotic and necroptotic signaling also differ in their evolutionary dynamics over large timescales. Specifically, key effectors of cellular necroptotic signaling emerged later in evolution and display a "patchy" phylogenetic distribution [13, 14] which is in marked contrast to the more ancient and highly conserved components of apoptotic signaling [15-17]. Notably, the evolutionary dynamics of necroptotic signaling over more recent timescales is unknown but may inform determinants shaping contemporary infections.

Necroptosis can be initiated by at least three presumably independent receptors:

1) death receptor (DR) signaling, 2) the pathogen recognition receptor (PRR) Z-DNA

Binding Protein 1 (ZBP1) / DNA Activator of Interferon (DAI), and 3) Toll-like Receptor

3/4 (TLR3/TLR4) [7]. Activation of DR signaling triggers Receptor Interacting Protein

Kinase 1 (RIPK1) to self-oligomerize, which leads to the recruitment of Receptor

Interacting Protein Kinase 3 (RIPK3) via physical RIP Homotypic Interaction Motif

(RHIM) domain interactions. Activated RIPK3 subsequently binds and phosphorylates

Mixed Lineage Kinase-Like (MLKL) [18-20]. Phosphorylation of MLKL results in a

conformational change, enabling self-oligomerization of this factor to complete

necroptosis through MLKL-plasma membrane destabilization [21, 22].

Importantly, DR initiation of necroptosis appears to function when caspase-8

activity is inhibited during apoptosis [23, 24]. In contrast, DAI triggers necroptosis upon

binding foreign nucleic acids followed by activation of RIPK3 via RHIM-dependent

interactions [25, 26]. In addition, TLR3 and TLR4 signaling can activate RIPK3 via TIR-

domain-containing adaptor-inducing interferon-β (TRIF) also through physical RHIM

domain interactions [27, 28]. The diverse and distinct means of activating necroptosis

with convergence on the RIPK3/MLKL axis reflect the breadth of pathogens this

response may protect against.

Consistently, a growing list of viral- and bacterial-encoded inhibitors that target

discrete steps of necroptosis has emerged [29-36]. However, whether this pathogen-

mediated antagonism has shaped cellular necroptotic factors remains unknown. Strong

selective pressure may be imposed on host factors to escape direct binding by

pathogen-encoded antagonists. This pressure is often visible as elevated rates of amino

acid substitution, a signature of rapid evolution and a hallmark of positive selection.

Signatures of rapid evolution have been observed across orthologous sequences from

closely related species (e.g. primates) for several key immune factors: PKR [37], OAS1

[38, 39], MxA [40], TRIM5α [41], APOBEC3G [41], and cGAS [38, 39]. Studies of host-

pathogen co-evolution may reveal novel insights into cellular responses including

evidence for undefined host components, genetic determinants shaping infection

outcomes, and pathogen countermeasures of these defenses. We hypothesized that

analysis of signatures characteristic of host-pathogen conflict for both cellular and viral

components would illuminate whether necroptosis has served as a "back-up" or a core,

fundamental response over evolutionary time.

Consistent with pivotal roles in host defense across species, we report that the

necroptotic axis in primates - RIPK3 and MLKL - displays widespread and recurrent

signatures of rapid evolution. Interestingly, we found evidence for positive selection

within the RHIM domain of RIPK3, but not at homologous sites in RHIM domains of DAI,

TRIF, or RIPK1, that is proximal to a known target-site for a conserved bacterial

protease. Notably, these bacterial proteases have been shown to cleave all of the RHIM

domain containing necroptosis factors [30]. These data suggest an undefined means of

RHIM-independent RIPK3 activation. Evolutionary genetics analysis of a recently

described poxvirus MLKL homolog indicate that this antagonist has been subjected to

repeated gains and losses over viral evolution with some poxvirus genomes encoding

up to three mimics. In summary, our data demonstrate that MLKL belongs to a small

class of key immune factors, which includes OAS1 but not PKR or cGAS, that display all

three hallmarks of pivotal immune factors - rare in combination: 1) rapidly evolving, 2)

viral homolog, and 3) upregulation by cytokines such as interferons. Collectively, our

data are consistent with necroptosis being a fundamental host defense, not an auxiliary

one, shaping infection outcomes over evolution.

Results

Widespread signatures of rapid evolution in primate RIPK3 and MLKL

To determine if known components of the necroptosis pathway display

signatures of positive selection, we analyzed a matching set of twenty-one primate

sequences for *TRIF*, *DAI*, *RIPK1*, *RIPK3* and *MLKL* using a series of evolutionary

analyses. To test for recurrent positive selection, we used a combination of maximum-

likelihood based algorithms, which estimate rates of non-synonymous amino acid

replacements (dN) relative to synonymous (dS) amino acid substitutions. dN/dS values

> 1 are considered a hallmark of positive selection. Using codon-based models

implemented in Phylogenetic Analysis by Maximum Likelihood (PAML) [42], we detected

robust gene-wide rapid evolution signatures for *MLKL* [M7 vs. M8 (F3x4) $p < 5.400 \times 10^{-10}$],

RIPK3 [M7 vs. M8 (F3x4) $p < 3.056 \times 10^{-9}$] and to a lesser extent, *DAI* [M7 vs. M8

(F3x4) $p < 5.767 \times 10^{-3}$] but not *RIPK1* [M7 vs. M8 (F3x4) $p < 0.065$] or *TRIF* [M7 vs.

M8 (F3x4) $p < 0.066$] (Fig 2A, S1 File). The signature of selection for the main effectors

of the necroptosis pathway, RIPK3 and MLKL, is comparable to other key host defense

factors – PKR, cGAS, OAS1 – which are also upregulated by cytokines. Indeed, MLKL

is also upregulated by cytokines like interferons [43-45], which we confirm for IFNγ (Fig

2B).

To assay when and to what extent selection pressure has shaped necroptotic

signaling over primate evolution, we estimated dN/dS values across the primate

phylogeny (Fig 2C) for each factor. Consistent with our gene-wide tests of evolution, we

observed recurrent and widespread signatures of positive selection across primates for

RIPK3, *MLKL* and again to a lesser extent, *DAI*. Specifically, numerous lineages across

primate evolution for both *RIPK3* and *MLKL*, including branches in each major group,

show substitution patterns that are characteristic of genetic conflict. Strikingly, especially

strong signatures are evident for *RIPK3* in the lineage preceding the divergence of

Hominidae [10 nonsynonymous changes: 0 synonymous changes] and for *MLKL* in the

lineage prior to Hominidae divergence [dN/dS = 4.49] (Fig 2C). In contrast, only one

lineage with a rapid evolution signature was detected for both *TRIF* and *RIPK1*. Thus,

repeated innovation in the necroptosis pathway has been primarily focused on the

downstream axis of *RIPK3*/*MLKL* during primate evolution.

To uncover where strong selection pressure may have been imposed within the

effector proteins of necroptosis, dN/dS values were estimated for individual amino acid

positions using three distinct but commonly implemented methods: PAML [42], MEME

[46], and FUBAR [47] (Fig 3A, S1 File). The distribution and the number of rapidly

evolving sites are thought to reflect the number of protein surfaces in genetic conflict

with other factors, such as pathogen-encoded inhibitors [48]. In agreement with

necroptosis being triggered by diverse pathogens, predicted rapidly evolving sites are

distributed throughout the necroptosis proteins (*MLKL*, *RIPK3*, *DAI*, *RIPK1*, and *TRIF*).

In support of our preceding analysis, *MLKL* (20 total sites / 11 by multiple methods) and

RIPK3 (27 total sites / 7 by multiple methods) have the most sites identified (Fig 3A).

Interestingly, the majority (70%) of MLKL positively selected sites are localized to the

kinase-like domain, with 12/14 sites within this domain occurring between amino acid

positions 370-470 (Fig 3A). Positively selected sites mapped onto a previously

published co-crystal structure of mouse RIPK3 and MLKL [49] indicate that clusters of

sites represent distinct surfaces (Fig 3B). Analysis of space-filling models highlights that

the majority of the rapidly evolving sites are found on the protein surface for both RIPK3

and MLKL, which suggests they represent interfaces for protein-protein interactions (S1

Fig). Notably, there is a cluster of positively selected sites for RIPK3 that surrounds

MLKL and corresponds to the hydrophobic pocket where RIPK3 and MLKL interact via

hydrophobic interactions (Fig 3C). This signature may indicate the presence of MLKL

pseudosubstrates, as this pocket has been demonstrated previously to be necessary for

RIPK3:MLKL interaction and for necroptosis to occur [31, 49].

Rapid evolution exclusive to the RHIM domain of RIPK3 localizes to a known
bacterial protease cleavage site
190
191   RHIM-RHIM homotypic interactions between RIPK3 and established necroptosis

activators RIPK1, TRIF, and DAI are required for signal transduction. The critical nature

of this interaction is also supported by the identification of both viral- and bacterial-

encoded inhibitors that target the RHIM domains of these proteins. These include

cytomegalovirus M45 and herpes simplex virus ICP6 and ICP10, which all encode

RHIM domains and target host RHIM-dependent interactions [29, 33]. Another example

is EspL, a bacterial protease, which is encoded by diverse species, including

enteropathogenic *Escherichia coli* (EPEC), *Salmonella enterica,* and *Yersinia pestis*

[30]. Notably, EPEC EspL has been shown to cleave a specific motif (Q*G**N) in the

RHIM domains of human and mouse TRIF, DAI, RIPK3, and RIPK1, which results in the

suppression of necroptosis [30]. Interestingly, rapid evolution was evident in the RIPK3

RHIM domain but not detectable in homologous domains across primate *TRIF*, *DAI,* and

RIPK1 (Fig 3D). Indeed, *RIPK3* showed elevated rates of substitution by PAML analysis

at two sites immediately adjacent to the determined RHIM EspL cleavage motif –

G<u>DN</u>/NYL where "/" indicates cleavage; the sites 462D and 463N are rapidly evolving

(Fig 3D). The evolutionary signatures of necroptosis RHIM domains may indicate

modes of RIPK3 activation and/or functions independent of the RIPK1, TRIF, and DAI

RHIM domains.

Poxviral antagonism of the RIPK3:MLKL interface serves as a model for
antagonistic coevolution
211
The rapid evolution signature of RIPK3 near the hydrophobic pocket indicates

antagonism, likely by pseudosubstrates (Fig 3C). Indeed, a viral copy of MLKL (vMLKL),

which is encoded by numerous poxviruses has been recently identified [31]. Yet, the

origins and evolution of these vMLKLs remain unknown. We identified twenty-seven

distinct copies of vMLKL in the sequence database across the genomes of *Avipoxvirus*,

Clade II, along with two other poxvirus species that infect bats [eptesipoxvirus (EPTV)

and hypsugopoxvirus (HYPV)] (Fig 4D). vMLKL lacks the N-terminal bundle and brace

(NBB) domain but maintains the C-terminal kinase-like domain, to which RIPK3 binds to

phosphorylate the host cellular homolog [31] (Fig 4A). Ectopic expression of two

vMLKLs from poxvirus species that primarily infect mice were demonstrated to

competitively inhibit both mouse and human RIPK3 in binding and cell-culture

necroptosis assays [31]. Superposition of a homology model of myxoma virus (MYXV)

MLKL onto published crystal structures for mouse MLKL [49] and human MLKL [50]

indicated striking structural overlap with human and mouse MLKL (Fig 4B). This occurs

despite a broad-range of amino acid divergence between these host and viral MLKLs

(16-25% amino acid identity of the virus homologs to mouse and human MLKLs).

In agreement with the strong structural overlap, characterization of vMLKL

sequences indicate conservation of functional residues for human and mouse MLKL

including ATP binding, hydrogen-bonding, and protein stability (S2A Fig) [19-21, 49-51].

Several amino acid sites displaying positive selection for MLKL (Figs 2 and 3)

overlapped or were in close proximity to sites for hydrogen bonding and protein stability,

as demonstrated by the amino acid alignment (S2A Fig). Of note, there are several

intriguing examples of specific residue conservation between the primate positively

selected sites and the viral copies. One example is human MLKL F386 (F373 mouse),

which has been shown to be necessary for RIPK3/MLKL complex formation [31]. This

position exclusively has tyrosine, phenylalanine, or histidine amino acid variants across

the sampled primates, with this pattern of aromatic residues evident and conserved to

Aves MLKL virologs (S2B Fig). Consistently, poxvirus virologs have also maintained

only aromatic (phenylalanine or tyrosine) amino acids at the homologous position (S2A

Fig). These data suggest a key role for site 386 in host defense but also in

counteraction by viral mimics [52].

To further understand the evolutionary relationships between poxvirus vMLKLs,

we performed detailed sequence and phylogenetic analyses across corresponding viral

genomes. Full-length vMLKLs (excluding partial Vulture gryphus poxvirus and Hawaiian

goosepox virus sequences) display 18%-100% amino acid identity relative to each other

and cluster in a manner resembling known poxviral species relationships (Fig 4C).

Unexpectedly, this analysis identified two distinct vMLKL copies (e.g. paralogs), which

differ in amino acid identity, in the genomes of both eptesipoxvirus (EPTV vMLKL-A vs.

EPTV vMLKL-B : 45% amino acid identity) and hypsugopoxvirus (HYPV vMLKL-A vs.

HYPV vMLKL-B : 43% amino acid identity). Analysis of the location of the different

vMLKLs in each genome was conducted using CoGeBlast, which allows for individual

hit visualization. The results suggested distinct genomic locations for these genes (Fig

4E). Specifically, a majority of vMLKLs are located primarily on the right arm of the

poxvirus genome with EPTV having two copies that map to two distinct loci in this

region. In addition, both HYPV and EPTV have one additional copy that maps to the left

arm of the poxvirus genome. Interestingly, two of the vMLKLs in EPTV are present on

the inverted terminal repeat (ITR), potentially representing more recent acquisitions.

Likewise, fowlpox virus-282E4 vMLKL appears to be at a distinct inverted locus relative

to other *Avipoxvirus* species on the left arm of the genome (Fig 4E).

To confirm additional vMLKL copy numbers present in the eptesipoxvirus

genome, we next performed relative depth analysis, with approximately 1000 - fold

genome coverage (Fig 4F, S3 Fig), using the raw reads from the genome sequencing.

By mapping reads across the unique genome and one reference ITR, we found that

EPTV – A/C has a relative depth of two and EPTV – B has a relative depth of one,

which supports the presence of three copies in the eptesipoxvirus genome (Fig 4F). As

a control, we also conducted a similar analysis using reads from a recent myxoma virus

sequencing project [53] and found that the vMLKL relative depth remained near one,

indicating the presence of only one genomic copy (S3 Fig). These data demonstrate

that vMLKLs have different evolutionary histories and are likely not orthologous.

Phylogenetic distribution of vMLKLs can account for variable conservation of E3
across species
273
Poxviruses encode numerous immunomodulators to subdue and circumvent host

defenses including apoptosis. As a majority of poxviruses encode several characterized

anti-apoptotic proteins [54, 55] (Fig 4G, S4 Fig, S2 File) including the serpin class of

caspase pseudosubstrates [56, 57], these viruses would be expected to trigger

necroptosis. Indeed, the model poxvirus vaccinia has been shown recently to trigger

necroptosis through activation of DAI/ZBP1 in cultured cells and to counteract this

response using the Z - nucleic acid binding (Zα) domain of the E3 protein [32]. E3 is

considered a key host-range gene that antagonizes several host defenses. Interestingly,

some poxviruses that encode a repertoire of anti-apoptotic proteins lack E3 or possess

truncated E3 ORFs. For example, the *Avipoxvirus* family does not possess any

orthologs of E3 (Fig 4G, S2 File) and several Clade II poxviruses, including the well-

characterized myxoma virus, are also predicted to have N-terminally truncated E3

copies which lack the Zα domain that is important for antagonism of DAI [55] (Fig 4G,

S2 File). To gain insights into how a subset of poxviruses may circumvent necroptosis

given the presumed ability to trigger apoptosis, we compared the breadth of known anti-

apoptotic proteins, including serpins, across the poxvirus phylogeny relative to vMLKL

(Fig 4G). As vMLKLs are present in *Avipoxvirus*, hypsugopoxvirus, myxoma virus and

rabbit fibroma virus – the maintenance of this pseudosubstrate may account for the

tolerance of these species to a missing or truncated E3 protein.

Repeated gains and losses of vMLKL over poxvirus evolution

Our data illustrates that multiple vMLKL copies may exist in a given poxvirus

genome and that not all of the copies across the genomes may be orthologous. To

determine the relationship of poxvirus vMLKLs, we performed a synteny analysis using

five flanking upstream and downstream genes across the entire phylogeny (Fig 5, S3

File). These data indicate that while each vMLKL within *Avipoxvirus* and Clade II are

flanked largely by the same upstream and downstream genes, vMLKLs between the

species are flanked by different genes. The Clade II and *Avipoxvirus* species have

distinct copies as indicated by different upstream and downstream genes. Furthermore,

the lack of vMLKL in Yatapoxvirus species (Yaba monkey tumor virus and Yaba-like

disease virus), which belong to Clade II, revealed that this loss is likely due to a deletion

event consisting of multiple upstream and downstream genes in these lineages (Fig 5,

S3 File).

This analysis also provided insights regarding the multiple vMLKL copies in

eptesipoxvirus and hypsugopoxvirus. First, this analysis indicated that EPTV-B (EPTV-

WA-166) and HYPV-B (QDJ95132.1) are orthologous to each other and are likely

orthologous to the Clade II copies, as evidenced by some overlap in syntenic gene

neighbors (Fig 5). Second, EPTV-A (EPTV-WA-006) and HYPV-A (QDJ94987.1) share

gene neighbors indicating that these sequences are likely orthologs. EPTV-A shares

100% identity with EPTV-C (EPTV-WA-186) with both annotated as being located in the

inverted terminal repeats (ITRs). Consistently, EPTV-C has the same gene neighbors

as EPTV-A albeit inverted. The copies are labeled corresponding to their location

starting from the left arm of the poxvirus genome – e.g., A, B, C.

To further assess the evolutionary relatedness of vMLKLs, we performed whole

genome comparisons using CoreGenes (https://coregenes.ngrok.io/). This data was

visualized with RIdeogram [58] and focused on the copies encoded by *Avipoxvirus*,

Clade II and the two poxvirus species that primarily infect bats. Indeed, we found that

canarypox virus vMLKL and myxoma virus vMLKL localize to distinct chromosomal

locations with no shared surrounding regions of synteny (S5A Fig). In agreement with

our local synteny analysis above, the locus encoding Clade II vMLKL was likely lost in a

Yatapoxvirus ancestor following divergence as evidenced by global as well as local

synteny surrounding the vMLKL locus (S5B Fig). Lastly, we compared eptesipoxvirus

with myxoma virus (S5C Fig). The upstream regions of synteny of vMLKL indicate that

MYXV vMLKL is orthologous to the EPTV-B vMLKL copy, which is further supported by

MYXV vMLKL not being in the ITRs corresponding to EPTV-A/C. These data suggest

vMLKLs are at distinct loci in poxvirus genomes as a result of repeated gains and

losses during poxvirus genome evolution.

Phylogenetic analysis reveals origination of vMLKL copy from one horizontal
gene transfer event
332
The distinct copies of vMLKLs may have originated due to either 1) a single

transfer event from a host genome to an ancestral poxvirus followed by diversification or

2) multiple independent transfer events from presumably distinct host genomes to

different poxviruses. To address this, we performed a phylogenetic analysis using

amino acid sequences of vMLKLs with thirty-eight diverse host MLKLs. If distinct

vMLKLs were acquired via independent events, placement of viral sequences would

vary across the tree in a manner where they pair with different host sequences. Such

topology could reflect the source genome or that of a related, perhaps ancestral, animal

host. In contrast, if vMLKLs originated from a single transfer event, the tree topology

would group these sequences into one clade. Phylogenetic analysis using PhyML [59]

revealed that vMLKL virologs cluster as one group distinct from animal vMLKLs (Fig 6).

These data support a single transfer event from an ancestral host genome as the most

parsimonious origin for vMLKL (Fig 6, S4 File). The topology of the poxvirus branches

mirrors that of known species relationships. For example, all of the *Avipoxvirus* species

cluster together. Likewise, the Clade II species cluster together broadly, as well as at a

finer level such as in the *Leporipoxviruses* (e.g. MYXV and RFV). The tree topology also

supports the synteny analysis with HYPV-B and EPTV-B being most closely related

along with EPTV-A/C and HYPV-A vMLKL clustering distinctly.

Still, it remains unclear whether Clade II vMLKLs represent a distinct or

orthologous locus to Avipox vMLKL and which vMLKL locus represents the ancestral

locus. Indeed, our data for Clade II vMLKL could be explained, in part, by either 1) a

duplication of the Avipox vMLKL locus followed by a deletion event of the original locus

in the Clade II ancestor or 2) that Avipox and Clade II vMLKL are orthologous but

divergence no longer allows for syntenic comparison between these species. We

assessed the likelihood of these evolutionary histories by analyzing patterns of amino

acid variation shared between vMLKLs and mammalian MLKLs, which cluster closest to

vMLKLs (Fig 6). We evaluated innovations specific to vMLKLs but not host MLKLs

across our alignment. We identified seventeen amino acid residues largely conserved

between Avipox vMLKLs and mammalian MLKLs (Fig 7) but not Clade II, EPTV and

HYPV vMLKLs. Conversely, different variants at the same sites are shared across

Clade II, EPTV and HYPV vMLKLs but not Avipox or mammalian MLKLs. These data

support the hypothesis that Avipox vMLKL is the original locus and that Clade II vMLKL

is a distinct copy but derived from the Avipox vMLKL presumably through duplication.

Collectively, these data illustrate a volatile history of vMLKL since its acquisition by an

ancestral *Avipoxvirus*.

Discussion

Studies of host-pathogen co-evolution can reveal new insights that forge infection

outcomes. These include vulnerabilities in host defense and determinants that shape

pathogen host range as well as zoonoses. In addition, analysis of host and pathogen

effectors can illuminate new evolutionary paradigms such as the evolutionary

trajectories of viral mimics. Here, we focused on the emerging regulated cell death

program of necroptosis that is largely presumed to be a "back-up" response. Our data,

particularly the robust signatures of positive selection in RIPK3/MLKL, mirrors that of

key immune factors like cGAS, PKR, and OAS1 [37, 38]. These features support the

essentiality of RIPK3/MLKL in immunity during primate evolution. In addition, the

maintenance, as well as gains and losses of vMLKL throughout poxvirus evolution,

showcases the need to actively counteract the necroptosis program.

Proposed model for vMLKL acquisition
382
The findings from our evolutionary analysis for vMLKL are consistent with a

model consisting of a series of gains and losses of this virolog over time (Fig 8A). Our

data paired with those of Petrie *et al.*, which also suggested a host origin for vMLKL

[31], support the notion that this gene was initially acquired via one horizontal gene

transfer event (Fig 6). As vMLKL mirrors a truncated copy of cellular MLKL lacking

introns, we hypothesize that this transfer likely occurred via an RNA intermediate. The

N-terminal truncation in vMLKL could have transpired either during or after integration.

This copy at the original site of integration has been maintained through evolutionary

time in *Avipoxvirus* species and represents locus 1. Following the Avipox ancestral

divergence from a common ancestor shared with Clade II and HYPV/EPTV species,

locus 1 appears to have been duplicated to produce locus 2. We hypothesize that locus

1 was deleted while locus 2 was maintained in the shared ancestor Clade

II/EPTV/HYPV, and is present today in most contemporary genomes of Clade II

poxviruses and EPTV-B/HYPV-B. Yet, it remains possible - due to extensive

rearrangements resulting in the degradation of any detectable synteny by either method

we employed here - that locus 1 across Avipox represents the Clade II/EPTV/HYPV

vMLKL. In Yatapoxviruses, locus 2 vMLKL was lost in what appears to be a multi-gene

deletion event (Fig 5, S5 Fig). Following divergence of the EPTV/HYPV ancestor from

Clade II poxviruses, locus 2 was duplicated to a region in the ITR with a subsequent

duplication occurring in the other ITR. Both vMLKL copies in the eptesipox ITRs are

supported by our relative depth analysis (Fig 4F, S3 Fig). A similar analysis was

attempted but not possible with the hypsugopox genome data. Thus, vMLKL resides at

four distinct loci in poxvirus genomes across the phylogeny.

Positive selection signatures in the necroptotic signaling pathway are localized to

the RIPK3/MLKL axis

The signatures of rapid evolution in RIPK3 and MLKL resemble "molecular arms

races" dynamics with pathogen-encoded inhibitors [48] (Fig 2). The broad distribution of

rapidly evolving sites in both RIPK3 and MLKL is indicative of multiple, undefined

antagonists that have existed over evolutionary time (Fig 3). Interestingly, our analysis

across the core circuitry of necroptosis enabled two key observations. First, strong

selective pressure has been primarily exerted on the downstream components of the

pathway where upstream signaling converges (Figs 2 and 3). Second, our evolutionary

analysis across the pathway allowed the identification of rampant positive selection

exclusive to the RHIM-domain of RIPK3 but not the RHIM-domains of the upstream

activators DAI, TRIF, or RIPK1 (Fig 3). This is extremely striking given the elevated

dN/dS ratios for RIPK3, but not the other factors, at amino acid positions that define a

demonstrated cleavage site (Fig 3D) for the conserved bacterial protease EspL which is

capable of cleaving RHIM domains across this pathway. Notably, rapid evolution in the

adaptor protein MAVS has been shown to shape the ability of the hepatitis protease

NS3/4A to cleave this target [60]. Our data suggest that there may be RIPK3 activity that

is independent of activation by RHIM homotypic interactions (e.g., DAI/TRIF/RIPK1),

which is counteracted by EspL-like proteases. One possible explanation for this

signature is the presence of a yet undefined upstream activator of RIPK3, which lacks a

RHIM-domain, that is triggered by pathogenic bacteria. Given the data, we are unable to

infer whether this predicted activity of RIPK3 would also activate MLKL. Nevertheless, a

pathway perspective combined with evolutionary insights can drive new avenues of

investigation.

RIPK3/MLKL as a model for the maintenance of compatible, functional

interactions

Our study describes volatile evolution of vMLKL in poxvirus genomes. This work

extends on the recent discovery of this mimic, which showed two vMLKLs are capable

of counteracting necroptosis in cell culture [31]. Our integrated comparison of

established poxvirus immunomodulators with the phylogenetic distribution of vMLKL

appears to resolve the long-standing conundrum of how some poxvirus species persist

with N-terminally truncated or completely absent E3 ORFs (Fig 4F). In addition, the lack

of vMLKLs in orthopoxviruses can be explained, in part, by the recent identification of

vIRD encoded by this genus, which is an ankyrin-repeat domain containing protein that

targets RIPK3 for degradation [35]. The resolution of the evolutionary history of vMLKL

in our study may serve as a resource in guiding the functional analysis of host and viral

MLKL variants and their impact on RIPK3 binding and activity. It has not escaped our

attention that the RIPK3/MLKL axis, in part, resembles another well-defined host

defense axis - PKR/eIF2 α - which also consists of a kinase and its substrate

counteracted by a different poxvirus pseudosubstrate (K3L) [37]. However, a major

difference between the two systems is that RIPK3 and MLKL are rapidly evolving, in

contrast to PKR rapidly evolving and eIF2α being ultraconserved to yeast. This is

interesting given the PKR/eIF2α system has been useful in understanding how a rapidly

evolving factor maintains its interaction with a conserved factor while evolving to escape

antagonism [37]. Perhaps RIPK3/MLKL may be powerful in understanding how two key

host defense factors maintain their essential interaction to drive necroptosis while

escaping pathogen-mediated inhibition, particularly pseudosubstrates similar to vMLKL.

MLKL is a pivotal immune factor

Our study defines MLKL, which functions as the executioner of necroptosis, as a

factor that 1) is (up)regulated by immune signals (Fig 2B) [43-45], 2) has a viral

homolog/virolog (Figs 4,5,6 and 7)[31] and 3) is rapidly evolving (Figs 2 and 3). While

several key host defense factors typically display two of these signatures, to our

knowledge, only a select few are published which display all three of these hallmarks

(Fig 8B): OAS1 [38, 39, 61], and the recently described MISTRAV and MISTR1 [62].

OAS1 is one of the first interferon-stimulated genes identified and functions in host

defense against numerous, diverse viruses [63]. MISTR proteins appear to interface

with electron transport chain complexes to regulate stress responses and cellular

adaptation during infection [62]. We hypothesize this collection of three hallmarks may

serve as a guide to identify other pivotal host defense factors. Collectively, we interpret

this rare combination of signatures that are associated with OAS1 and now MLKL as

strong evidence that necroptosis is an indispensable, not auxiliary, host response that

has shaped infection outcomes over millions of years.

Methods

Protein modeling

Published crystal structures of m(ouse) RIPK3 (PDB: 4M66), mMLKL (PBD:4M68) and

co-crystal structure for mRIPK3:mMLKL (PDB: 4M69) were used for modeling following

pairwise alignment with h(uman)MLKL (PDB: 4BTF) or the predicted structure of

hRIPK3 [**20**, **49**]. Swiss-Model was used to predict the structure of hRIPK3 and myxoma

v(iral)MLKL [**64**]. UCSF Chimera (https://www.cgl.ucsf.edu/chimera/) was used to

perform analysis and visualize structures including rapidly evolving sites [**65**].

Positive selection analysis

Nucleotide sequences were obtained from NCBI and Ensembl databases (S1 File).

Multiple sequence alignments (MSA) were performed using MAFFT iterative refinement

method FFT-NS-i and visualized using Geneious Prime 2020.1.2

(https://www.geneious.com/) [66]. Indels were removed through manual trimming. dN/dS

lineage estimates were obtained using sample primate gene MSA and phylogenetic tree

[created using previously described primate lineage relationships [67]]. These served as

input for FreeRatio analysis implemented in PAML [42]. Nucleotide substitution (NS) site

analysis was performed with PAML using both F3x4 and F61 codon frequency models.

MEME and FUBAR analysis were performed using Datamonkey

(https://www.datamonkey.org/) to predict positively selected amino acid sites [46, 47,

68]. The S1 File contains additional findings from this analysis.

Identification of poxvirus vMLKL

489   Virologs to human MLKL were identified using tBLASTn and/or BLASTp [69].

490   Sequences were selected based on returned BLAST parameters (total score, E value

491   and percent identity) and the phylogenetic clustering of the sequences using PhyML

492   [59]. Scaled individual hit visualization was constructed using CoGeBlast [70].

Relative depth analysis

Eptesipox raw sequencing reads, published in Tu *et al.,* were obtained directly from Dr.

Chris Upton (University of Victoria, Victoria BC, Canada) and Dr. Yoshinori Nakazawa

(CDC, Atlanta, Georgia) [71] . NC_035460.1 was used as the eptesipoxvirus reference

sequence to map the reads. Myxoma virus raw sequencing reads were obtained from

European Nucleotide Archive (https://www.ebi.ac.uk/ena/browser/home), from Project

PRJNA513218 [53]. Accession IDs used for this analysis are SRR8402032,

SRR8402033, SRR8402034, and SRR8402035. As a control, AF170726.2 was used as

the myxoma virus reference sequence to map the reads. Raw sequencing reads were

processed using Trimmomatic-0.39 [72] to remove adaptors. Tanoti short read aligner

(https://bioinformatics.cvr.ac.uk/software/tanoti/) was then used to map reads to the

reference genome. SAMtools [73] was used to convert between file formats, merge files

and calculate coverage per basepair. Bins of 500 nucleotides were created for relative

depth analysis, which was calculated by dividing the bins by the mean of the non-ITR

genomic region. Data was visualized using gplots [74].

Phylogenetic analysis

Multiple Sequence Alignments were performed using MAFFT, with iterative refinement method L-INS-i, on amino acid sequences retrieved from NCBI and Ensembl databases (S4 File) [66]. Maximum Likelihood trees were inferred by PhyML using 100 bootstrap values [59].

Percent amino acid identity heatmap

Multiple Sequence Alignments were performed using MAFFT, with iterative refinement method L-INS-i, on amino acid sequences retrieved from NCBI database (S2 File) [66]. The R package gplots was used to create the heatmap found in Fig 4C [74, 75].

Poxvirus RCD-associated host range gene identification

Host range genes were identified using BLASTp and tBLASTn. Sequences were confirmed using reciprocal BLAST hits. Protein families were found using genomic database annotations and published results for included poxvirus genomes. Domain classification for homologs and protein families were confirmed using NCBI conserved genome database [76]. Results were compared to previous homolog studies performed by Bratke *et al.* 2013 and Haller *et al.* 2014, and can be found in S2 File [54, 55]. Proteins truncated at the N- and C-termini were not included in Fig 4G but can be found in the S2 File with appropriate label. An exception is necroptosis inhibitor E3, which is predicted to be N-terminally truncated in *Leporipoxviruses* [55]. Genes residing in the inverted terminal repeats (ITR) are counted as separate genes and thus two instances in order to stay consistent with the Fig 4C initial analysis, where the eptesipoxvirus ITR is included and analyzed. The p28/N1R and Ankyrin-repeat families were annotated using NCBI database annotations. The presence of these domains were confirmed

531   using NCBI conserved genome database or via BLASTp [76]. This analysis includes all

532   proteins possessing a p28/N1R-like or Ankyrin-repeat domain and does not predict if

533   these proteins are functional and/or likely truncated.

High resolution synteny analysis

Reciprocal BLAST hits (RBH) were performed for coding sequences upstream and

downstream of vMLKL genomic locations across poxvirus species included in this study.

Results can be found in the S2 File. Genomic database annotations and the NCBI

conserved genome database were used to confirm results [76]. Gene blocks

representing Ankyrin-repeat containing proteins without one distinct hit, but conserved

homology across poxvirus species are colored navy.

Whole genome comparison

543   Coregenes 5.0 (https://coregenes.ngrok.io/) was used to compare selected poxvirus

544   genomes [77]. Matches were identified using bidirectional best hit with an e-value =

545   1×10^{5}. RIdeogram [58] was used to visualize the output obtained from Coregenes 5.0.

Cell lines

547   HT29 and HeLa cell lines were obtained from ATCC. Both cell lines were cultured in

548   Corning DMEM with L-Glutamine, 4.5 g/L Glucose, and Sodium Pyruvate supplemented

549   with 10% FBS and 1X Gibco Antibiotic-Antimycotic solution. Cell lines were maintained

550   at 37°C in a humidified incubator at 5% CO_2.

Cell culture treatments

552   Cells were treated with either human IFNα [1000 U/mL (PBL Assay Science, USA) or

553   human IFNγ [1000 U/mL (ThermoFisher, USA)] diluted in DMEM and incubated for 24

554   hours. After 24 hours, cell lysates were harvested for western blots as described below.

Western blot analysis

Cells were collected using RIPA Lysis and Extraction Buffer (ThermoFisher, 89901) supplemented with Protease and Phosphatase Inhibitor Cocktail (Abcam, ab201119). Protein concentrations were measured using a Bradford assay. For all blots, 5 μg of protein lysates were loaded onto an SDS-PAGE gel and wet-transferred to 0.45 μM Immobilon-P PVDF membrane (Millipore, IPVH00010) at 200 mA for 90 minutes. Membranes were blocked with blocking buffer (5% milk in TBST) for 30 minutes at RT. Membranes were incubated in primary antibodies at 4°C overnight. The following primary antibodies used: MLKL (GeneTex, GTX107538), STAT1 D1K9Y Rabbit mAb (Cell Signaling Technology, 14994S), P-STAT1 Tyr701 58D6 Rabbit mAb (Cell Signaling Technology, 9167S) and β-Actin (Abcam, ab49900). Membranes were washed three times with TBST for 5 minutes and incubated with Goat Anti-Rabbit IgG (Bio-Rad, 170-6515) for 1 hour at RT. Membranes were washed three times with TBST and incubated with Clarity Max Western ECL Substrate (Bio-Rad, 1705062). Following incubation, blots were imaged using the Chemidoc MP Imager (Bio-Rad).

Acknowledgments

We express gratitude to other members of the Hancks Lab for feedback and discussion through the course of this project. We also thank Dr. Anant Gharpure and Dr. Sherry Haller for their comments on the manuscript. We also thank Dr. Chris Upton and Dr. Yoshinori Nakazawa as well as Dr. Chiara Chiapponi and Dr. Davide Lelli for their assistance with the eptesipoxvirus and hypsugopoxvirus data, respectively.

References

1.   Danthi P. Viruses and the Diversity of Cell Death. Annu Rev Virol. 2016;3(1):533-53.

2.   Jorgensen I, Rayamajhi M, Miao EA. Programmed cell death as a defence
against infection. Nat Rev Immunol. 2017;17(3):151-64.
3.   Lamkanfi M, Dixit VM. Manipulation of host cell death pathways during microbial
infections. Cell Host Microbe. 2010;8(1):44-54.
4.   Best SM. Viral subversion of apoptotic enzymes: escape from death row. Annu
Rev Microbiol. 2008;62:171-92.
5.   Barber GN. Host defense, viruses and apoptosis. Cell Death Differ.
2001;8(2):113-26.
6.   Kerr JF, Wyllie AH, Currie AR. Apoptosis: a basic biological phenomenon with
wide-ranging implications in tissue kinetics. Br J Cancer. 1972;26(4):239-57.
7.   Galluzzi L, Vitale I, Aaronson SA, Abrams JM, Adam D, Agostinis P, et al.
Molecular mechanisms of cell death: recommendations of the Nomenclature
Committee on Cell Death 2018. Cell Death Differ. 2018;25(3):486-541.
8.   Strasser A, O'Connor L, Dixit VM. Apoptosis signaling. Annu Rev Biochem.
2000;69:217-45.
9.   Pearson JS, Murphy JM. Down the rabbit hole: Is necroptosis truly an innate
response to infection? Cell Microbiol. 2017;19(8).
10.   Nailwal H, Chan FK. Necroptosis in anti-viral inflammation. Cell Death Differ.
2019;26(1):4-13.
11.   Green DR. The Coming Decade of Cell Death Research: Five Riddles. Cell.
2019;177(5):1094-107.
12.   Vanden Berghe T, Linkermann A, Jouan-Lanhouet S, Walczak H, Vandenabeele
P. Regulated necrosis: the expanding network of non-apoptotic cell death
pathways. Nat Rev Mol Cell Biol. 2014;15(2):135-47.
13.   Dondelinger Y, Hulpiau P, Saeys Y, Bertrand MJM, Vandenabeele P. An
evolutionary perspective on the necroptotic pathway. Trends Cell Biol.
2016;26(10):721-32.
14.   Brault M, Oberst A. Controlled detonation: evolution of necroptosis in pathogen
defense. Immunol Cell Biol. 2017;95(2):131-6.
15.   Green DR, Fitzgerald P. Just So Stories about the Evolution of Apoptosis. Curr
Biol. 2016;26(13):R620-R7.
16.   Koonin EV, Aravind L. Origin and evolution of eukaryotic apoptosis: the bacterial
connection. Cell Death Differ. 2002;9(4):394-404.
17.   Zmasek CM, Godzik A. Evolution of the animal apoptosis network. Cold Spring
Harb Perspect Biol. 2013;5(3):a008649.
18.   Wang H, Sun L, Su L, Rizo J, Liu L, Wang LF, et al. Mixed lineage kinase
domain-like protein MLKL causes necrotic membrane disruption upon
phosphorylation by RIP3. Mol Cell. 2014;54(1):133-46.
19.   Sun L, Wang H, Wang Z, He S, Chen S, Liao D, et al. Mixed lineage kinase
domain-like protein mediates necrosis signaling downstream of RIP3 kinase.
Cell. 2012;148(1-2):213-27.
20.   Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang JG, Alvarez-Diaz S, et
al. The pseudokinase MLKL mediates necroptosis via a molecular switch
mechanism. Immunity. 2013;39(3):443-53.

21.   Cai Z, Jitkaew S, Zhao J, Chiang HC, Choksi S, Liu J, et al. Plasma membrane
625   translocation of trimerized MLKL protein is required for TNF-induced necroptosis.
626   Nat Cell Biol. 2014;16(1):55-65.

627   22.   Chen X, Li W, Ren J, Huang D, He WT, Song Y, et al. Translocation of mixed
628   lineage kinase domain-like protein to plasma membrane leads to necrotic cell
629   death. Cell Res. 2014;24(1):105-21.

630   23.   Oberst A, Dillon CP, Weinlich R, McCormick LL, Fitzgerald P, Pop C, et al.
631   Catalytic activity of the caspase-8-FLIP(L) complex inhibits RIPK3-dependent
632   necrosis. Nature. 2011;471(7338):363-7.

633   24.   Kaiser WJ, Upton JW, Long AB, Livingston-Rosanoff D, Daley-Bauer LP, Hakem
634   R, et al. RIP3 mediates the embryonic lethality of caspase-8-deficient mice.
635   Nature. 2011;471(7338):368-72.

636   25.   Maelfait J, Liverpool L, Bridgeman A, Ragan KB, Upton JW, Rehwinkel J.
637   Sensing of viral and endogenous RNA by ZBP1/DAI induces necroptosis. EMBO
638   J. 2017;36(17):2529-43.

639   26.   Kaiser WJ, Upton JW, Mocarski ES. Receptor-interacting protein homotypic
640   interaction motif-dependent control of NF-kappa B activation via the DNA-
641   dependent activator of IFN regulatory factors. J Immunol. 2008;181(9):6427-34.

642   27.   Kaiser WJ, Sridharan H, Huang C, Mandal P, Upton JW, Gough PJ, et al. Toll-
643   like receptor 3-mediated necrosis via TRIF, RIP3, and MLKL. J Biol Chem.
644   2013;288(43):31268-79.

645   28.   He S, Liang Y, Shao F, Wang X. Toll-like receptors activate programmed
646   necrosis in macrophages through a receptor-interacting kinase-3-mediated
647   pathway. Proc Natl Acad Sci U S A. 2011;108(50):20054-9.

648   29.   Guo H, Omoto S, Harris PA, Finger JN, Bertin J, Gough PJ, et al. Herpes simplex
649   virus suppresses necroptosis in human cells. Cell Host Microbe. 2015;17(2):243-
650   51.

651   30.   Pearson JS, Giogha C, Muhlen S, Nachbur U, Pham CL, Zhang Y, et al. EspL is
652   a bacterial cysteine protease effector that cleaves RHIM proteins to block
653   necroptosis and inflammation. Nat Microbiol. 2017;2:16258.

654   31.   Petrie EJ, Sandow JJ, Lehmann WIL, Liang LY, Coursier D, Young SN, et al.
655   Viral MLKL Homologs Subvert Necroptotic Cell Death by Sequestering Cellular
656   RIPK3. Cell Rep. 2019;28(13):3309-19 e5.

657   32.   Koehler H, Cotsmire S, Langland J, Kibler KV, Kalman D, Upton JW, et al.
658   Inhibition of DAI-dependent necroptosis by the Z-DNA binding domain of the
659   vaccinia virus innate immune evasion protein, E3. Proc Natl Acad Sci U S A.
660   2017;114(43):11506-11.

661   33.   Upton JW, Kaiser WJ, Mocarski ES. Cytomegalovirus M45 cell death
662   suppression requires receptor-interacting protein (RIP) homotypic interaction
663   motif (RHIM)-dependent interaction with RIP1. J Biol Chem.
664   2008;283(25):16966-70.

665   34.   Mack C, Sickmann A, Lembo D, Brune W. Inhibition of proinflammatory and
666   innate immune signaling pathways by a cytomegalovirus RIP1-interacting
667   protein. Proc Natl Acad Sci U S A. 2008;105(8):3094-9.

35.   Liu Z, Nailwal H, Rector J, Rahman MM, Sam R, McFadden G, et al. A class of
viral inducer of degradation of the necroptosis adaptor RIPK3 regulates virus-
induced inflammation. Immunity. 2021.
36.   Fletcher-Etherington A, Nobre L, Nightingale K, Antrobus R, Nichols J, Davison
AJ, et al. Human cytomegalovirus protein pUL36: A dual cell death pathway
inhibitor. Proc Natl Acad Sci U S A. 2020;117(31):18771-9.
37.   Elde NC, Child SJ, Geballe AP, Malik HS. Protein kinase R reveals an
evolutionary model for defeating viral mimicry. Nature. 2009;457(7228):485-9.
38.   Hancks DC, Hartley MK, Hagan C, Clark NL, Elde NC. Overlapping Patterns of
Rapid Evolution in the Nucleic Acid Sensors cGAS and OAS1 Suggest a
Common Mechanism of Pathogen Antagonism and Escape. PLoS Genet.
2015;11(5):e1005203.
39.   Mozzi A, Pontremoli C, Forni D, Clerici M, Pozzoli U, Bresolin N, et al. OASes
and STING: adaptive evolution in concert. Genome Biol Evol. 2015;7(4):1016-32.
40.   Mitchell PS, Emerman M, Malik HS. An evolutionary perspective on the broad
antiviral specificity of MxA. Curr Opin Microbiol. 2013;16(4):493-9.
41.   Sawyer SL, Wu LI, Emerman M, Malik HS. Positive selection of primate
TRIM5alpha identifies a critical species-specific retroviral restriction domain. Proc
Natl Acad Sci U S A. 2005;102(8):2832-7.
42.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol.
2007;24(8):1586-91.
43.   Knuth AK, Rosler S, Schenk B, Kowald L, van Wijk SJL, Fulda S. Interferons
Transcriptionally Up-Regulate MLKL Expression in Cancer Cells. Neoplasia.
2019;21(1):74-81.
44.   Sarhan J, Liu BC, Muendlein HI, Weindel CG, Smirnova I, Tang AY, et al.
Constitutive interferon signaling maintains critical threshold of MLKL expression
to license necroptosis. Cell Death Differ. 2019;26(2):332-47.
45.   Rusinova I, Forster S, Yu S, Kannan A, Masse M, Cumming H, et al. Interferome
v2.0: an updated database of annotated interferon-regulated genes. Nucleic
Acids Res. 2013;41(Database issue):D1040-6.
46.   Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL.
Detecting individual sites subject to episodic diversifying selection. PLoS Genet.
2012;8(7):e1002764.
47.   Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, et
al. FUBAR: a fast, unconstrained bayesian approximation for inferring selection.
Mol Biol Evol. 2013;30(5):1196-205.
48.   Daugherty MD, Malik HS. Rules of engagement: molecular insights from host-
virus arms races. Annu Rev Genet. 2012;46:677-700.
49.   Xie T, Peng W, Yan C, Wu J, Gong X, Shi Y. Structural insights into RIP3-
mediated necroptotic signaling. Cell Rep. 2013;5(1):70-8.
50.   Murphy JM, Lucet IS, Hildebrand JM, Tanzer MC, Young SN, Sharma P, et al.
Insights into the evolution of divergent nucleotide-binding mechanisms among
pseudokinases revealed by crystal structures of human and mouse MLKL.
Biochem J. 2014;457(3):369-77.
51.   UniProt C. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res.
2019;47(D1):D506-D15.

52. Davies KA, Fitzgibbon C, Young SN, Garnish SE, Yeung W, Coursier D, et al.
Distinct pseudokinase domain conformations underlie divergent activation
mechanisms among vertebrate MLKL orthologues. Nat Commun.
2020;11(1):3060.
53. Kerr PJ, Eden JS, Di Giallonardo F, Peacock D, Liu J, Strive T, et al. Punctuated
Evolution of Myxoma Virus: Rapid and Disjunct Evolution of a Recent Viral
Lineage in Australia. J Virol. 2019;93(8).
54. Haller SL, Peng C, McFadden G, Rothenburg S. Poxviruses and the evolution of
host range and virulence. Infect Genet Evol. 2014;21:15-40.
55. Bratke KA, McLysaght A, Rothenburg S. A survey of host range genes in
poxvirus genomes. Infect Genet Evol. 2013;14:406-25.
56. Nichols DB, De Martini W, Cottrell J. Poxviruses Utilize Multiple Strategies to
Inhibit Apoptosis. Viruses. 2017;9(8).
57. Taylor JM, Barry M. Near death experiences: poxvirus regulation of apoptotic
death. Virology. 2006;344(1):139-50.
58. Zhaodong Hao DL, Ying Ge, Jisen Shi, Dolf Weijers, Guangchuang Yu, Jinhui
Chen. RIdeogram: drawing SVG graphics to visualize and map genome-wide
data on the idiograms. PeerJ Computer Science. 2020;6:e251.
59. Lefort V, Longueville JE, Gascuel O. SMS: Smart Model Selection in PhyML. Mol
Biol Evol. 2017;34(9):2422-4.
60. Patel MR, Loo YM, Horner SM, Gale M, Jr., Malik HS. Convergent evolution of
escape from hepaciviral antagonism in primates. PLoS Biol.
2012;10(3):e1001282.
61. Darby AC, McInnes CJ, Kjaer KH, Wood AR, Hughes M, Martensen PM, et al.
Novel host-related virulence factors are encoded by squirrelpox virus, the main
causative agent of epidemic disease in red squirrels in the UK. PLoS One.
2014;9(7):e96439.
62. Sorouri M, Chang T, Jesudhasan P, Pinkham C, Elde NC, Hancks DC.
Signatures of host-pathogen evolutionary conflict reveal MISTR-A conserved
MItochondrial STress Response network. PLoS Biol. 2020;18(12):e3001045.
63. Chakrabarti A, Jha BK, Silverman RH. New insights into the role of RNase L in
innate immunity. J Interferon Cytokine Res. 2011;31(1):49-57.
64. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al.
SWISS-MODEL: homology modelling of protein structures and complexes.
Nucleic Acids Res. 2018;46(W1):W296-W303.
65. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et
al. UCSF Chimera--a visualization system for exploratory research and analysis.
J Comput Chem. 2004;25(13):1605-12.
66. Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief Bioinform.
2019;20(4):1160-6.
67. Perelman P, Johnson WE, Roos C, Seuanez HN, Horvath JE, Moreira MA, et al.
A molecular phylogeny of living primates. PLoS Genet. 2011;7(3):e1001342.
68. Weaver S, Shank SD, Spielman SJ, Li M, Muse SV, Kosakovsky Pond SL.
Datamonkey 2.0: A Modern Web Application for Characterizing Selective and
Other Evolutionary Processes. Mol Biol Evol. 2018;35(3):773-7.

69.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment
761   search tool. J Mol Biol. 1990;215(3):403-10.
762   70.   Lyons E, Pedersen B, Kane J, Alam M, Ming R, Tang H, et al. Finding and
763   comparing syntenic regions among Arabidopsis and the outgroups papaya,
764   poplar, and grape: CoGe with rosids. Plant Physiol. 2008;148(4):1772-81.
765   71.   Tu SL, Nakazawa Y, Gao J, Wilkins K, Gallardo-Romero N, Li Y, et al.
766   Characterization of Eptesipoxvirus, a novel poxvirus from a microchiropteran bat.
767   Virus Genes. 2017;53(6):856-67.
768   72.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
769   sequence data. Bioinformatics. 2014;30(15):2114-20.
770   73.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The
771   Sequence Alignment/Map format and SAMtools. Bioinformatics.
772   2009;25(16):2078-9.
773   74.   Gregory R. Warnes BB, Lodewijk Bonebakker, Robert Gentleman, Wolfgang
774   Huber, Andy Liaw, Thomas Lumley, Martin Maechler, Arni Magnusson, Steffen
775   Moeller, Marc Schwartz and Bill Venables. gplots: Various R Programming Tools
776   for Plotting Data. R package version 3.0.3 ed2020.
777   75.   Team RC. R: A language and environment for statistical computing. R
778   Foundation for Statistical Computing. Vienna, Austria2013.
779   76.   Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al.
780   CDD/SPARCLE: the conserved domain database in 2020. Nucleic Acids Res.
781   2020;48(D1):D265-D8.
782   77.   Contreras-Moreira B, Vinuesa P. GET_HOMOLOGUES, a versatile software
783   package for scalable and robust microbial pangenome analysis. Appl Environ
784   Microbiol. 2013;79(24):7696-701.

Figure captions

Fig 1. Cell death pathways, a mechanism of host defense. (A) Apoptosis is a

regulated cell death pathway that acts as a key line of host defense against pathogens

and is executed by caspases via an extrinsic or intrinsic route [**7**]. **(B)** Necroptosis can

act as an alternative regulated cell death pathway in the event of inhibition of the

apoptosis pathway by pathogens [**10**] such as poxviruses. E3 [**32**] and vMLKL [**31**] are

recently described poxvirus factors that inhibit necroptotic signaling.

Fig 2. Rapid evolution of the necroptotic axis, RIPK3 and MLKL, across primate

lineages. A) Summary of rapid evolution results for M7 x M8 (F3x4) Likelihood ratio test

statistics (2δ) for necroptotic factors across a set of twenty-one matching primate

species. **B)** MLKL is upregulated by IFNγ treatment in HeLa and HT29 cell lines. **C)**

dN/dS values estimated using PAML across twenty-one primate lineages. Branches

colored in red display signatures of positive selection (dN/dS > 1, or greater than or

equal to 3 nonsynonymous amino acid changes relative to zero synonymous amino acid

changes).

Fig 3. Widespread signatures of positive selection across necroptosis protein

sequences. A) Rapidly evolving sites represented as triangles and shown for TRIF

(orange), DAI (pink), RIPK1 (red), RIPK3 (green) and MLKL (blue). Rapidly evolving

amino acid sites predicted using PAML (yellow triangle), FUBAR (blue triangle) and

MEME (green triangle). Sites with p-values < 0.05 or posterior probabilities (P) > 0.95

are listed above the protein cartoons and represented as triangles. Sites with p-values <

0.01 or posterior probabilities (P) > 0.99 are bolded and have an asterisk. Amino acid

alignment position refers to the human reference sequences. **B)** Location of rapidly

evolving sites (shown in red) on the h(uman)RIPK3 (green) homology model and the

hMLKL (light blue) crystal structure [20]. The homology model was predicted using

Swiss-Model [64] and aligned to crystal structures m(ouse)RIPK3 (silver), mMLKL

(silver), and mRIPK3:mMLKL (silver) [49]. **C)** Rapid evolution (red) at the mRIPK3

hydrophobic pocket. **D)** Rapid evolution exclusive to the RIPK3 RHIM-domain proximal

to the EspL cleavage motif (Q*G**N - highlighted in black). Sequences spanning the

EspL cleavage motif from 12 primates for RIPK3 (green), RIPK1 (red), TRIF (orange),

and DAI (pink) RHIM domains. The EspL cleavage site is underlined and displays a

black arrow above the amino acid. Sites with p-values < 0.05 or posterior probabilities

(P) > 0.95 are listed above the protein cartoons and represented as triangles. Sites with

p-values < 0.01 or posterior probabilities (P) > 0.99 are bolded and have an asterisk.

Amino acid alignment position refers to the human reference sequences. Note: PAML,

MEME, and FUBAR analysis were performed for RIPK1 without drill monkey, since this

region was trimmed in previous analyses.

Fig 4. vMLKL mimics are encoded in *Avipoxvirus*, Clade II, eptesipoxvirus, and

hypsugopoxvirus genomes. A) Protein structures of human MLKL (light blue) and

myxoma v(irus)MLKL (dark blue). **B)** Homology model for myxoma vMLKL (dark blue)

predicted using Swiss-Model and aligned onto previously published crystal structures

for hMLKL (light blue) and mMLKL (silver). **C)** Heatmap comparing percent identities of

vMLKL copies across poxvirus species which were included in this study. **D)** Cladogram

depicting distinct species containing a vMLKL copy with *Avipoxvirus* (green), Clade II

(blue) and other poxvirus species (yellow, pink and red) separated according to genus.

Note that eptesipoxvirus and hypsugopoxvirus contain multiple copies of vMLKL and

are labeled based on their position in the chromosome – A (yellow), B (pink), C (red).

Eptesipoxvirus (black dashed box) contains vMLKL copies located on the inverted

terminal repeats (ITR) represented by A (yellow) and C (red). **E)** Location of individual

and multiple copies of vMLKL across poxvirus genomes. Multiple copies of vMLKL are

highlighted (yellow, pink, and red) in eptesipoxvirus and hypsugopoxvirus**.**

Chromosomal cartoons created in CoGeBlast represent scaled locations of the vMLKL

copies for each of the poxvirus species. The genome accession numbers are located on

the top left corner of the chromosomes, and the nucleotide lengths are located on the

top right corner of the chromosomes. **F)** Multiple copies of vMLKL in the eptesipoxvirus

genome are supported by relative depth analysis. The distribution of reads that align

with the ITR and non-ITR regions is shown. Zoomed in images, represented by black

boxes, show the relative depth of the EPTV vMLKLs (black) and the two flanking

upstream and downstream genes (gray). The ITR relative depth mean is 1.852 and the

median is 1.899. The non-ITR relative depth mean is 1.000 and the median is 1.025. **G)**

Repertoire of poxvirus apoptosis and necroptosis antagonists explained, in part, by

evolutionary distribution of vMLKL. Only proteins predicted to be functional are shown.

Two exceptions, found in myxoma virus and rabbit fibroma virus, are represented by an

'*' and have N-terminally truncated E3 copies. In depth analysis can be found in S2 File.

Fig 5. High resolution synteny analysis suggests diverse evolutionary histories

for vMLKL. Analysis of upstream and downstream flanking genes indicate vMLKLs

(black) reside at different locations across poxvirus genomes. Protein homology was

determined using reciprocal BLAST hits for the 5 genes upstream (left of vMLKL) and 5

genes downstream (right of vMLKL) of vMLKL. While a gene window of 11 is shown

here, an expanded analysis is found in the S3 File. Genes oriented to the right

represent 5'-3' directionality while genes oriented to the left represent 3'-5' directionality.

The letter "U" represents a gene that is either unique to the individual species or unique

to the group of poxviruses that have maintained a vMLKL copy. Gene blocks that have

homologous genes (colored borders) that are outside the scope of this figure have

arrows pointing to the left (representing upstream) or to the right (representing

downstream). Two Yatapoxvirus species (Yaba monkey tumor virus and Yaba-like

disease virus) were included as they belong to clade II but lack vMLKL. ൠ represents a

deletion event. Species defined as "Other" (red) are either unclassified

(hypsugopoxvirus) or part of vespertilionpoxviruses (eptesipoxvirus).

Fig 6. Phylogenetic analysis suggests a singular origin for non-orthologous poxvirus MLKLs. A maximum-likelihood phylogenetic tree of diverse host and poxviral MLKL amino acid sequences produced using PhyML [59] is displayed. Consistent with a single horizontal gene transfer event, *Avipoxvirus* (green), Clade II (blue), and other poxvirus sequences (yellow, pink, and red) cluster together. Branch lengths represent amino acid changes per site. 100 bootstrap replicates were performed with branch support > 70% (*) or 90% indicated (**). The alignment of MLKL sequences was performed using MAFFT and can be found in the S4 File.

Fig 7. Amino acid variant analysis suggests Avipox vMLKL is the original locus, and Clade II vMLKL is an independent, but derived copy. A set of informative amino acid positions were identified in the alignment of mammalian and vMLKL sequences. Residues identical to consensus are shown as "." while "-" represents a gap in the sequence alignment. Amino acid site positions (shown above the alignment) are in relation to human MLKL. Alignment can be found in the S4 File.

Fig 8. Volatile evolution of MLKL and hallmarks associated with key host defense factors indicate that necroptosis is a pivotal, not auxiliary, immune defense. A) Model of poxvirus MLKL acquisition and evolution across lineages. Data indicate that vMLKL was likely acquired by a vertebrate host presumably through a single horizontal gene transfer event to a poxvirus ancestor (Fig.6). This copy at the original site of integration, located on the right arm, has been maintained in *Avipoxvirus* (green) genomes over time to present day (Fig 4). A duplication followed by a deletion event occurred in the common ancestor of Clade II (dark blue) and species classified as "Other" (red) following divergence from *Avipox* (Figs 5 and 7, S5 Fig). This second copy

also on the right arm has been maintained until the present in all Clade II species but

lost in Yatapoxvirus lineages (Fig.5). Subsequently, the second copy was duplicated to

the ITR in the ancestor of eptesipoxvirus and hypsugopoxvirus, which is maintained in

present-day lineages. While hypsugopoxvirus ITR sequence was not included in the

final genome assembly, our synteny analysis (Fig 5) suggests vMLKL is also located on

an ITR (dashed box) in this lineage. "ᘯ" represents predicted deletion events. Dashed

arrows represent the progression of evolutionary time (left to right). Scaled

chromosomal locations of the present day vMLKL copies are represented on the right.

B) MLKL displays all three key hallmarks often common to key host defense factors but

unique to OAS1 [38, 39, 61], MISTRAV, and MISTR1 [62]. These factors are rapidly

evolving, possess a viral homolog/virolog, and are regulated by immune signals

(includes upregulation and downregulation).

Supporting information

S1 Fig. MLKL and RIPK3 rapidly evolving sites localize to protein surfaces.

Positively selected sites (red) identified in Fig 3 mapped onto filled crystal structures of

RIPK3 and MLKL.

S2 Fig. Protein sequence alignments for human MLKL, mouse MLKL and

vMLKLs. A) Diagrams of hMLKL (light blue) and mMLKL (silver) with previously

established functional residues annotated [19-21, 49-51]. Amino acid alignment of

hMLKL, mMLKL, and poxvirus MLKLs was constructed using MAFFT, with functional

residues annotated on diagrams also highlighted on this alignment. Brackets

correspond to the starting point of vMLKL, which is truncated, relative to hMLKL

(alignment beginning at hMLKL N155). **B)** Alignment of human MLKL F386

(corresponding to mouse MLKL F373) with MLKLs from diverse host species. Aromatic

amino acid positions (red) are primarily conserved to Aves.

S3 Fig. Relative depth analysis of myxoma virus genome sequencing reads

indicates a single copy of vMLKL in this species. A) Table shows MYXV and EPTV

read depth (total number of reads in the BAM file) and effective depth (number of reads

in the BAM file that map to the reference genome) following merging of bam files.

Analysis performed using data from Kerr *et al.*, and Tu *et al.,* [53, 71]. Myxoma virus

analysis was performed as a control given complementary analysis (Fig 4F) and

predicted single-copy vMLKL in this and other clade II poxviruses. As the effective depth

for myxoma virus sequencing was lower than eptesipoxvirus data, read depth isolates

were pooled together to facilitate a more normalized approximation of the read counts

per nucleotide in the genome. **B)** Myxoma virus genome relative depth analysis - (reads

that align with the ITR and non-ITR regions), where "n =" represents the number of

myxoma virus isolates pooled together from Project PRJNA513218 [53]. Inset, relative

depth at MYVX vMLKL (black) locus including two flanking upstream and downstream

genes (gray). The non-ITR and ITR means and medians are present on each plot.

S4 Fig. Poxvirus Ankyrin and P28/N1R protein family ranges separated by genus.

Summary of ankyrin-repeat containing and P28/N1R proteins per genome analyzed in

Fig 4G. Functional protein predictions were not performed for these families.

S5 Fig. Whole genome synteny analysis illustrates the presence of distinct,

multiple copies of vMLKL across poxvirus genomes. The whole genome

comparisons are **A)** canarypox virus (*Avipoxvirus* - green) versus myxoma virus (Clade

II - blue), **B)** yaba-like disease virus (Clade II - blue) versus myxoma virus and **C)**

eptesipoxvirus (Other - red) versus myxoma virus. Sequence accession numbers are

centered at the bottom of each genome. Lines that extend between genomes represent

homologous regions, that were calculated using CoreGenes 5.0

(https://coregenes.ngrok.io/) [77]. Lines that are light gray represent conserved

homologous regions that are not part of the Fig 5 high resolution synteny analysis. Lines

in dark gray represent conserved homologous genes that are shown in Fig 5 and the S3

File. vMLKL is represented by a black line. "⚡" represents a deletion event.

S1 File. Supporting analysis for Figs 2 and 3 and S1 Fig.

S2 File. Supporting analysis for Fig 4 and S4 Fig.

S3 File. Supporting analysis for Fig 5.

S4 File. Supporting analysis for Figs 6 and 7.

Palmer et al. Fig 1

A) Apoptosis

Intrinsic Pathway
Mitochondria

Extrinsic Pathway
Death Receptor

Caspase 9

Caspase 8

Caspase 3/6/7

Apoptosis

Caspase Inhibition

B) Necroptosis

Death Receptor

Toll-like Receptor

RIPK1

TRIF

DAI

E3

RIPK3

vMLKL

MLKL

Necroptosis

none

Palmer et al. Fig 2

A)

	2δ	P-value
TRIF	5.481	0.065
DAI	10.311	**5.767 x 10⁻³**
RIPK1	5.434	0.066
RIPK3	39.212	**3.056 x 10⁻⁹**
MLKL	42.679	**5.400 x 10⁻¹⁰**

Palmer et al. Fig 3

A) Human MLKL

N — NBB | Kinase-like — C
1 — 471

Myxoma MLKL

N — Kinase-like — C
1 — 288

B)

Human MLKL
Myxoma MLKL
Mouse MLKL

C) Percent Identity
0 — 100

Avipox · Clade II · HYPV/EPTV

D)

Vultur Gryphus
Magpiepox
Canarypox
Shearwaterpox-2
Hawaiian Goosepox
Fowlpox-282E4
Shearwaterpox-1
Fowlpox-MN00.2
Flamingopox
Penguinpox
Pigeonpox
Turkeypox
Swinepox
Sheeppox
Goatpox
Lumpy Skin Disease
Deerpox-W-848-83
Deerpox-W-1170-84
Moosepox
Rabbit Fibroma
Myxoma
Cotia
BeAn
Eptesipox-B
Hypsugopox-B
Hypsugopox-A
Eptesipox-A
ITR
Eptesipox-C

E)

MK903864.1	293226
AY318871.1	359853
KX857215.1	351108
MG702259.1	308827
KX857216.1	326929
MH709124.1	298798
MF678796.1	293123
KJ859677.1	306862
KJ801920.1	282356
KP728110.2	188534
AF410153.1	146454
MN072628.1	149956
KC951854.1	150194
AF409137.1	150793
AY689436.1	166259
AY689437.1	170560
MG751778.1	164258
AF170722.1	159857
AF170726.2	161773
KM595078.1	185139
KY094066.1	163005
MK860688.1 (A ... B)	166600
NC_035460.1 (A ... B C)	176688

G)

	N1	F1/M11	T4	Serpins	C7 Family	E3
MPPV	0	0	0	4	0	0
CNPV	0	0	0	4	0	0
SWPV-2	0	0	0	4	0	0
FPV-282E4	0	0	0	3	0	0
SWPV-1	0	0	0	4	1	0
FPV-MN00.2	0	0	0	4	0	0
FGPV	0	0	0	5	1	0
PEPV	0	0	0	4	1	0
FEP	0	0	0	4	1	0
TKPV	0	0	0	2	1	0
SPV	0	1	0	1	1	1
SPPV	1	1	2	1	1	1
GTPV	1	1	2	1	1	1
LSDV	1	1	2	1	1	1
DPV-W-8448-83	1	1	2	4	1	1
DPV-W-1170-84	1	1	2	4	1	1
MOPV	1	1	2	4	1	1
RFV	1	1	2	1	3	1*
MYXV	1	1	2	3	3	1*
COTV	1	1	0	3	2	1
BAV	1	1	0	0	1	1
HYPV	1	1	0	1	1	0
EPTV	1	1	0	3	1	1

F)

EPTV- A/C

EPTV- B

Relative Depth

ITRs

Nucleotide Position
0 — 50,000 — 100,000 — 150,000

Palmer et al. Fig 5

Palmer et al. Fig 6

Palmer et al. Fig 7

Human MLKL

* site positions are in relation to human MLKL

Species	222Y	230K	257F	267G	290T	296D	313A	316G	320L	321H	328L	329H	389Y	394E	431E	441L	449R
Consensus	Y	K	F	G	T	D	A	G	L	H	L	H	Y	E	E	L	R
Human	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
African Green Monkey	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Bolivian Squirrel Monkey	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Bushbaby	.	E	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Mouse	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Rat	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Chinese Hamster	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Blind Mole Rat	.	.	L	-	-	-	-	-	.	.	.	.	.	.	.	.	.
Guinea Pig	.	.	.	.	S	.	.	.	.	.	.	.	.	.	.	.	.
Squirrel	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Pika	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Tree Shrew	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Westeen European Hedge Hog	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Pig	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Cow	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Sheep	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Alpaca	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Horse	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Megabat	.	.	.	.	.	.	.	.	.	M	.	C	.	.	.	.	.
Microbat	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Malayan Pangolin	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Florida Manatee	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Nine-banded Armadillo	.	.	.	.	.	.	.	.	.	.	.	.	.	.	Q	.	L
Magpiepox	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Canarypox	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Shearwaterpox-2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Fowlpox-282E4	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Shearwaterpox-1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Fowlpox-MN00.2	.	.	L	.	.	.	.	.	M	.	.	.	.	.	.	.	.
Flamingopox	.	.	L	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Penguinpox	.	.	L	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Pigeonpox	.	.	L	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Turkeypox	.	.	Y	.	S	K	.	.	.	.	.	.	.	.	.	.	.
Swinepox	F	.	I	A	Y	M	C	.	.	Y	-	Y	H	D	L	I	L
Sheeppox	F	R	I	A	Y	R	C	.	V	Y	-	Y	R	D	L	I	T
Goatpox	F	R	I	A	Y	R	C	.	V	Y	-	Y	R	D	L	I	T
Lumpy Skin Disease	F	R	I	A	Y	K	C	.	V	Y	-	Y	R	D	L	I	T
Deerpox-W-848-83	F	R	I	.	Y	N	C	S	.	Y	-	Y	L	D	F	I	T
Deerpox-W-1170-84	F	R	I	.	Y	N	C	S	.	Y	-	Y	L	D	F	I	T
Moosepox	F	R	I	.	Y	N	C	S	.	Y	-	Y	L	D	F	I	T
Rabbit Fibroma	F	R	V	.	Y	S	C	.	V	Y	-	Y	.	D	L	I	T
Myxoma	F	R	V	.	Y	Y	Y	C	.	V	C	-	Y	.	D	L	T
Cotia	.	I	K	A	P	L	S	S	F	Y	-	Y	Q	D	S	.	I
BeAn	.	I	K	A	P	L	.	S	.	Y	-	Y	Q	D	S	.	V
Hypsugopox-B	.	.	I	.	Y	I	.	A	Y	Y	V	Y	.	D	L	I	L
Eptesipox-B	F	.	I	.	Y	I	.	A	Y	Y	.	.	.	D	L	I	L
Hypsugopox-A	F	.	I	.	Y	.	.	A	Y	Y	V	Y	.	D	L	.	L
Eptesipox-A	F	.	I	.	Y	.	.	A	Y	Y	V	Y	F	D	L	.	L
Eptesipox-C	F	.	I	.	Y	.	.	A	Y	Y	V	Y	F	D	L	.	L

Mammalia (Human through Nine-banded Armadillo); *Poxviridae*: Avipox (Magpiepox through Turkeypox), Clade II (Swinepox through BeAn), Other (Hypsugopox-B through Eptesipox-C)

A)

B)

A)

B)

A)

Species	Number of Isolates	Read Depth	Effective Depth
Eptesipox	1	15,580,507	952,609
Myxoma	1	827,204	690,356
	2	1,821,118	1,659,533
	3	2,441,318	2,230,935
	4	3,424,938	3,102,827

B)

Palmer et al. S4 Fig

	Avipox	Clade II	HYPV/EPTV
Ankyrin Protein Family (proteins/genome)	16-62	4-16	4-8
P28/N1R Family (proteins/genome)	3-26	1-2	1

Palmer et al. S5 Fig