1 TITLE: Target enrichment and extensive population sampling help untangle the recent, rapid
2 radiation of *Oenothera* sect. *Calylophus*
3
4 RUNNING HEAD: Cooper et al., Phylogenomics of *Oenothera* sect. *Calylophus*
5
6 AUTHORS:
7

8 Benjamin J. Cooper[1,2,*], Michael J. Moore[3], Norman A. Douglas[4], Warren L. Wagner[5], Matthew
9 G. Johnson[1,6], Rick P. Overson[1,7], Angela J. McDonnell[1], Rachel A. Levin[8], Robert A. Raguso[9],
10 Hilda Flores Olvera[10], Helga Ochoterena[10], Jeremie B. Fant[1,2], Krissa A. Skogen[1,2], Norman J.
11 Wickett[1,2,*]
12
[1] The Negaunee Institute for Plant Conservation Science and Action, Chicago Botanic Garden,
1000 Lake Cook Rd., Glencoe, IL 60022, USA
[2] Northwestern University, Program in Plant Biology and Conservation, O.T. Hogan Hall, Room
6-140B, 2205 Tech Drive, Evanston, IL 60208, USA
[3] Oberlin College, Department of Biology, 119 Woodland St., Oberlin, OH 44074, USA
[4] Department of Biology, University of Florida, Gainesville, FL 32611, USA
[5] Department of Botany, MRC-166, Smithsonian Institution, PO Box 37012, Washington, DC
20013-7012, USA
[6] Department of Biological Sciences, Texas Tech University, Box 43131 Lubbock, TX 79409,
USA
[7] School of Sustainability, Arizona State University, PO Box 875502, Tempe, AZ 85287-5502,
USA
[8] Department of Biology, Amherst College, 25 East Drive, Amherst, MA, 01002, USA
[9] Department of Neurobiology & Behavior, Cornell University, 215 Tower Road, Ithaca, NY
14853, USA
[10] Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México,
Mexico City, Mexico
30
*Authors for correspondence: nwickett@chicagobotanic.org, benjamin.cooper06@gmail.com

ABSTRACT

Oenothera sect. *Calylophus* is a North American group of 13 recognized taxa in the

evening primrose family (Onagraceae) with an evolutionary history that may include

independent origins of bee pollination, edaphic endemism, and permanent translocation

heterozygosity. Like other groups that radiated relatively recently and rapidly, taxon boundaries

within *Oenothera* sect. *Calylophus* have remained challenging to circumscribe. In this study, we

used target enrichment, flanking non-coding regions, summary coalescent methods, tests for

gene flow modified for target-enrichment data, and morphometric analysis to reconstruct

phylogenetic hypotheses, evaluate current taxon circumscriptions, and examine character

evolution in *Oenothera* sect. *Calylophus*. Because sect. *Calylophus* comprises a clade with a

relatively restricted geographic range, we were able to extensively sample across the range of

geographic and morphological diversity in the group. We found that the combination of exons

and flanking non-coding regions led to improved support for species relationships. We

reconstructed potential hybrid origins of some accessions and note that if processes such as

hybridization are not taken into account, the number of inferred evolutionary transitions may be

artificially inflated. We recovered strong evidence for multiple origins of the evolution of bee

pollination from ancestral hawkmoth pollination, the evolution of edaphic specialization on

gypsum, and permanent translocation heterozygosity. This study applies newly emerging

techniques alongside dense infraspecific sampling and morphological analyses to effectively

address a relatively common but recalcitrant problem in evolutionary biology.

Keywords.– Gypsum Endemism, Onagraceae, *Oenothera* sect. *Calylophus*, Pollinator Shift,

Recent Radiation, Phylogenomics, Target Enrichment

INTRODUCTION

The challenges of reconstructing species histories for groups that arose through recent,

rapid radiations are well established. Phylogenetic signal can be obscured by processes such as

incomplete lineage sorting (ILS) and gene flow (Maddison and Knowles 2006; Knowles and

Chan 2008; Christie and Knowles 2015), resulting in short branch lengths and conflicting gene

tree topologies. Consequently, approaches that use few loci or concatenation may fail to resolve

the most accurate species tree (Eckert and Carstens 2008; Leaché et al. 2014; Xi et al. 2014;

Giarla and Esselstyn 2015). This may be particularly common in plants that are thought to have

experienced rapid or recent radiation with ongoing hybridization and high levels of ILS. The

application of target enrichment methods that efficiently sequence hundreds of nuclear loci,

coalescent-based phylogenetic methods that account for ILS and gene flow, and extensive

sampling of morphologically diverse populations across the geographic range should allow for

more accurate representations of phylogenetic relationships (Maddison and Knowles 2006;

Knowles and Chan 2008; Knowles 2009; Mamanova et al. 2010; Lemmon et al. 2012; Straub et

al. 2012; Bryson et al. 2014; Weitemier et al. 2014; Mandel et al. 2014; Stephens et al. 2015;

Johnson et al. 2016).

Oenothera sect. *Calylophus* currently comprises seven species (thirteen taxa) with a

center of diversity in western Texas, southern New Mexico, and north-central Mexico (Fig. 1;

Towner 1977; Turner and Moore 2014; Wagner 2021). Previous analyses suggest that *Oenothera*

sect. *Calylophus* forms a well-supported, morphologically coherent clade with a relatively

restricted geographic range (Towner 1977; Levin et al. 2004; Wagner et al. 2007; Turner and

Moore 2014; Wagner 2021). However, as with other groups that have experienced rapid

78 radiations, taxon boundaries within *Oenothera* sect. *Calylophus* have been challenging to define,

79 likely due to phenomena such as overlapping morphological boundaries, ongoing introgression,

80 and incomplete lineage sorting.

Figure 1

In the most comprehensive study of the group to date, Towner (1977) circumscribed taxa

using morphology, breeding system, geography, and ecology, but it was noted (and our field

observations confirm) that characters often overlap among taxa (Towner 1977). Taxa within

Oenothera sect. *Calylophus* are divided into two easily recognizable subsections: subsect.

Salpingia and subsect. *Calylophus* (Towner 1977; Wagner et al. 2007). Pollination varies

between the two subsections; flowers of subsect. *Salpingia* are adapted to vespertine pollination

by hawkmoths, except for *O. tubicula*, which opens in the morning and is primarily pollinated by

bees (Towner 1977), while taxa in subsect. *Calylophus* are predominantly bee-pollinated

(Towner 1977). Taxa in both subsections have geographic ranges that partially (or even largely)

overlap, resulting in occasional morphologically intermediate populations (Towner 1977).

Although confounding for morphological-based analyses, this observed pattern of reticulation is

consistent with a recent, rapid radiation occurring in parallel with climatic fluctuations and

increasing aridity in the region since the Pleistocene (Raven 1964; Towner 1977; Nason et al.

2002; Katinas et al. 2004). Hawkmoth pollination, which is ancestral in Onagraceae and common

in *Oenothera* sect. *Calylophus*, is known to result in long-distance pollen movement (Stockhouse

1973; Skogen et al. 2019). Therefore, gene flow may have been extensive over the evolutionary

history of hawkmoth-pollinated taxa, increasing the chances that processes such as historical

introgression may obscure phylogenetic signal in extant plants (Elrich and Raven 1969). With a

phylogenomic approach that samples hundreds of nuclear loci, we may better illuminate both the

history of these species and the key evolutionary processes related to speciation in this group.

Understanding speciation in angiosperms remains a fundamental question in evolutionary

biology (Barrett et al. 1996; Rajakaruna 2004; van der Niet et al. 2006; Wilson et al. 2007;

Crepet and Niklas 2009; Peakall et al. 2010; Xu et al. 2011; Van der Niet and Johnson 2012;

Boberg et al. 2014). *Oenothera* section *Calylophus* has an evolutionary history that likely

involves changes in reproductive system (pollinator functional group, breeding system) and

edaphic endemism. For example, there are thought to be two independent shifts between

pollinators from hawkmoth to bee pollination (Towner 1977; Fig. 1b), despite many studies in

other plant groups showing a directional bias in shifts from bee to hummingbird or hawkmoth

pollination (Barrett et al. 1996; Wilson et al. 2007; Thomson and Wilson 2008; Tripp and Manos

2008; Barrett 2013). However, pollinator shifts that do not follow this sequence may be more

likely when the extent of trait divergence and specialization does not completely inhibit

secondary pollinators such as bees, as has been suggested in *Oenothera* sect. *Calylophus*

(Stebbins 1970; Tripp and Manos 2008; Van Der Niet et al. 2014). Shifts to autogamy are also

frequent across angiosperms and in Onagraceae alone there are an estimated 353 shifts to modal

autogamy (i.e. mostly autogamous; Raven 1979). *Oenothera* sect. *Calylophus* also includes at

least one autogamous species, *O. serrulata*, which exhibits permanent translocation

heterozygosity, a phenomenon in which all chromosomes are translocationally heterozygous

(PTH; Towner 1977) (Fig. 1b). While the evolution of PTH has been assessed in molecular

phylogenetic analyses across Onagraceae (Johnson et al. 2009; Hollister et al. 2019), no study to

date has examined this transition in a well-sampled clade with extensive population sampling.

Lastly, abiotic ecological factors such as edaphic specialization are also known to drive

speciation in some groups (Rajakaruna 2004; van der Niet et al. 2006), including in *Oenothera*

sect. *Pachylophus* (Patsis et al. in press). For example, serpentine endemics represent ~10% of

the endemic flora in California even though serpentine soils account for about 1% of terrestrial

habitat in the state (Brady et al. 2005). Similarly, the Chihuahuan Desert is comprised of

numerous isolated islands of gypsum outcrops and current estimates suggest that at least 235 taxa

from 36 different plant families are gypsum endemics (Moore and Jansen 2007; Moore et al.

2014). It is suspected that gypsum endemism has also evolved independently in *Oenothera* sect.

Calylophus at least twice (Towner 1977; Turner and Moore 2014; Fig. 1b). Ultimately, to

understand the role that these transitions have played in shaping the diversity of *Oenothera* sect.

Calylophus, a robust phylogeny is required.

Here, we use target enrichment, summary coalescent methods, and morphometric

analyses to reconstruct a phylogenetic relationships, leveraging this information to re-examine

previous taxonomic concepts, test for instances of hybridization, and resolve the history of

pollinator shifts, PTH, and gypsum endemism in *Oenothera* sect. *Calylophus*. Target enrichment

is a cost-effective method for sequencing hundreds of loci across a high volume of samples,

producing highly informative datasets for phylogenetics (Lemmon et al. 2012; Straub et al. 2012;

Mandel et al. 2014; Weitemier et al. 2014; Heyduk et al. 2015; Stephens et al. 2015; Johnson et

al. 2016). While target enrichment is generally designed to capture coding regions, a significant

proportion of flanking non-coding regions can be recovered (the "splash-zone"; Weitemier et al.

2014). The inclusion of non-coding regions may be particularly informative for recent radiations,

since these regions are less constrained by selective pressures and may contain on average more

informative sites at shallower time scales (Folk et al. 2015). We included these flanking non-

coding regions in our sequence alignments to evaluate their impact on reconstructing lower-order

relationships. Importantly, we sampled extensively, including individuals from numerous

populations across the geographic and morphological ranges of all thirteen taxa in the section

(Fig. 1a). This study presents an example of how combining these molecular techniques with

dense sampling and morphological analysis can be used to effectively address a common

problem in evolutionary biology.

MATERIALS AND METHODS

A total of 194 individuals spanning the geographic, morphological, and ecological ranges

of all 13 recognized taxa in *Oenothera* sect. *Calylophus* [following Towner (1977) and Turner

and Moore (2014)] were included in this study (Fig. 1a, S1, S2) along with 8 outgroups

representing other major sections of *Oenothera* (*Eremia, Gaura, Kneiffia, Lavauxia, Oenothera,*

Pachylophus, and *Ravenia*) and other genera (*Chylismia* and *Eulobus*) in Onagraceae (S1, S2).

DNA was extracted from fresh, silica-dried leaf tissue (S3). PTH status was confirmed or

reassessed for individuals in subsect. *Calylophus* by assessing pollen fertility, when flowers were

present, using a modified Alexander stain (Alexander 1969, 1980; S3).

Target nuclear loci for enrichment were determined by clustering transcriptome

assemblies of *Oenothera serrulata* (1KP accession *SJAN*) and *Oenothera capillifolia* subsp.

berlandieri (1KP accession *EQYT*). Starting with the 956 phylogenetically informative

Arabidopsis loci identified by Duarte et al. (2010; S3), we identified 322 homologous, single-

copy loci in our clusters and used these in the probe design process. Libraries were enriched for

these loci using the MyBaits protocol (Arbor Biosciences, Ann Arbor, MI, USA) and sequenced

on an Illumina MiSeq (2 x 300 cycles, v3 chemistry; Illumina, Inc., San Diego, California,

USA). Raw reads have been deposited at the NCBI Sequence Read Archive (BioProject

PRJNA544074; See S3 for details). Reads were trimmed using Trimmomatic (Bolger et al. 2014;

S3) and trimmed, quality-filtered reads were assembled using HybPiper (Johnson et al. 2016).

From the assembled loci, we produced two datasets: **"exons",** constiting of exon-only

alignments, and **"supercontig",** consisting of alignments containing both the exon alignment and

flanking non-coding regions (the "splash-zone" per Weitemier 2014 and reconstructed using

supercontigs produced by HybPiper). We used these two datasets to test the most recent

taxonomic circumscription of the group with several methods: (1) phylogenetic inference of

concatenated alignments (two analyses: exons and supercontigs) using RAxML (Stamatakis

2014), (2) ASTRAL-II (Mirarab and Warnow 2015; Sayyari and Mirarab 2016) species tree

inference (two analyses: exons and supercontigs), (3) SVD Quartets (Chifman and Kubatko

2014, 2015) species tree inference (one analysis: supercontigs), (4) Phyparts (Smith et al. 2015;

one analysis: supercontigs), (5) IQtree (Minh et al. 2018) with both gene and site concordance

factors (one analysis: supercontigs).

We used HyDe (Blischak et al. 2018) to test for putative hybrid origins of selected taxa

and accessions by calculating D-Statistics (Green et al. 2010) for a set of hypotheses (S3). To

further characterize population-level processes or genetic structure within sect. *Calylophus*, we

extracted and filtered SNPs by mapping individual reads against reference supercontigs (see

https://github.com/lindsawi/HybSeq-SNP-Extraction) and used Discriminant Analysis of

Principal Components (Jombart et al. 2010) as implemented in the R package *adegenet* (Jombart

2008) and the snmf function in the LEA package (Frichot and François 2015) in R (R Core

Team, 2020; S3).

We evaluated current taxonomic concepts and patterns of morphological variation by

measuring character states for vegetative and floral structures that have been used historically to

discriminate taxa in *Oenothera* sect. *Calylophus* (Towner 1977): plant height, leaf length (distal),

leaf width (distal), leaf length/width ratio (distal), leaf length (basal), leaf width (basal), leaf

length/width (basal), sepal length, and sepal tip length (S3). Measurements were made for 125 of

the sequenced samples (S11); we were unable to measure all traits for 73 samples because we did

not have access to the herbarium vouchers, or the trait of interest was not captured on the

voucher, therefore some samples were dropped from the analysis due to missing values. Finally,

the number of transitions and inferred ancestral conditions of reproductive system were mapped

onto an ASTRAL species tree, with individuals grouped into species, using the stochastic

mapping function in the R package *phangorn* version 2.5.5 (Schliep 2011; S3).

RESULTS AND DISCUSSION

Sequencing and Phylogenetic Results

Sequencing resulted in a total of 80,273,296 pairs of 300-bp reads with an average of 625,323 reads per sample. Following quality filtering, assembly and alignment, we recovered 204 loci that were present in at least 70% of the samples. Across all datasets and analyses, *Oenothera* sect. *Calylophus* was monophyletic. At the subsection level there was strong agreement in topology between concatenation and coalescent-based trees. For example, subsect. *Calylophus* was recovered as sister to *Oenothera* subsect. *Salpingia* (minus *O. toumeyi*) with strong support across all analyses and *O. toumeyi* [considered by Towner (1977) to be in subsect. *Salpingia*; Fig. 2, S4-7] was recovered as sister to subsect. *Calylophus* across all trees, with strong bootstrap support. Within subsect. *Calylophus* there was poor resolution for currently recognized taxa in all analyses, whereas taxon relationships were better resolved in subsect. *Salpingia* (Fig. 2, S4-7). With coalescent-based tree reconstruction, most taxa *sensu* Towner (1977) were recovered as monophyletic with moderate to strong support (Fig. 2, S6-8). In contrast, both the exon and supercontig concatenation trees recovered most currently recognized taxa as non-monophyletic (S4, S5). Given that concatenation has been shown to produce incorrect topologies in the presence of high ILS (Roch and Steel 2015) and that *Oenothera* sect. *Calylophus* underwent recent radiation, we believe the paraphyly of taxa in both concatenation trees might be artifactual. We therefore interpret relationships based on our coalescent-based trees, which comprise the focus for the remainder of the paper (Fig. 2, S6-8).

Figure 2

To understand whether summary coalescent relationships display a consistent signal

across the genome, we quantified gene tree and site concordance using Phyparts and IQtree. We

found that gene tree concordance was highest at the deepest nodes at the species level where we

expected less ILS and more time between speciation events (Fig. 2, S8, S9). Correspondingly,

gene tree concordance was lowest at the subspecies level where increased sharing of ancestral

alleles and ongoing gene flow are more likely (Fig. 2, S8, S9). For example, within *O. hartwegii*

less than 1% of genes were concordant for bifurcations representing all currently recognized taxa

at the subspecies level (Fig. 2, S8, S9). For species-level nodes with high support and high gene

tree concordance, site concordance was also high; for example, *O. lavandulifolia* had high

bootstrap support in summary coalescent trees (BS = 100), high gene tree concordance (Phyparts

= 94% concordance, gCF = 74), and high site concordance (sCF = 83; Fig. 2, S9h). For

subspecies with high support, but low gene tree concordance, site concordance was moderate.

For example *O. hartwegii* subsp. *fendleri* had high bootstrap support (BS = 99), low gene tree

concordance (Phyparts = <1% concordance, gCF = 1), and moderate site concordance (sCF = 38;

Fig. 2, S9b). For subspecies that were monophyletic in our coalescent-based trees, but that had

low support and low gene tree concordance, site concordance was moderate with an average of

35% of sites in agreement for taxa at these nodes (S9[c,f,g,v]). For example *O. hartwegii* subsp.

filifolia had low bootstrap support, low gene tree concordance (Phyparts = <1% concordance,

gCF = 0), and moderate site concordance (sCF = 36; Fig. 2, S9c). This is an important finding

because while species tree relationships can be obscured by ILS, site concordance factors, which

may be less constrained and less subject to ILS at shallower evolutionary timescales, provide a

key alternative method of support (Minh et al. 2018).

In general, topologies of exon and supercontig datasets were similar, with no major

differences in clade membership, but the inclusion of the flanking non-coding regions increased

support at shallow nodes in our trees. However, this trend was not universal. For example, in

subsect. *Salpingia*, using the supercontig dataset decreased support slightly for one taxon (*O.*

hartwegii subsp. *filifolia*), and in subsect. *Calylophus* it led to paraphyly of another (*O.*

capillifolia subsp. *capillifolia*). For six other taxa, our results showed that using supercontigs

increased bootstrap support. Therefore, these results demonstrated a net benefit of including

flanking non-coding regions for resolving relationships among closely related taxa.

Hybridization and Geneflow

Using concatenated loci from the supercontig dataset, we used HyDe (Blischak et al.

2018) to test for signals of hybridization. We used 552,521 sites and tested 22 hypotheses for

either individuals or groups suspected to be of hybrid origin based on field observations of

morphological intermediacy, geographic location, and topological position in our coalescent-

based trees and found evidence of hybridization in three individuals representing two taxa, both

in subsect. *Salpingia*. The highest signal of hybridization, with a gamma value ($\hat{\gamma}$) of 0.947

suggesting more historic gene-flow, was observed in one of three sampled individuals of *O.*

tubicula subsp. *strigulosa* (*MJM1916.E*). This involved admixture between *O. tubicula* subsp.

tubicula and the clade consisting of *O. hartwegii* subsp. *hartwegii* and *O. hartwegii* subsp.

maccartii (Z-score = 5.585, p-value = 0.000, $\hat{\gamma}$ = 0.947; Fig. 3a, S10). We also detected

significant levels of hybridization, with $\hat{\gamma}$ ranging from 0.332 to 0.338 suggesting more

contemporary gene-flow, in two individuals in *O. hartwegii* subsp. *pubescens*, *BJC29* (Z-score =

2.378, p-value = 0.009, $\hat{\gamma}$ = 0.338) and *MJM594* (Z-score = 2.094, p-value = 0.018, $\hat{\gamma}$ = 0.332).

This more recent gene flow involved admixture between *O. hartwegii* subsp. *pubescens* and the

clade consisting of *O. hartwegii* subsp. *hartwegii* and *O. hartwegii* subsp. *maccartii* (Fig. 3a,

S10).

269   The finding that one individual of *O. tubicula* subsp. *strigulosa* may be of hybrid origin is

consistent with gene flow between *O. tubicula* subsp. *strigulosa* and its sister taxon *O. tubicula*

subsp. *tubicula* (sensu Towner 1977). In our coalescent-based analyses, the two subspecies of *O.*

tubicula were not recovered as sister taxa, and this relationship was strongly supported (S9[j-k]).

If the two *O. tubicula* taxa arose independently, this would support the hypothesis that bee

pollination arose in *Oenothera* sect. *Calylophus* independently three times. However, while the

summary coalescent analyses we utilized to estimate phylogenies accounted for ILS in tree

estimation, they did not account for gene flow (Meng and Kubatko 2009; Gerard et al. 2011;

Kubatko and Chifman 2019). Our HyDe results may support the hypothesis that *O. tubicula*

subsp. *strigulosa* has experienced gene flow from two closely related taxa, and may have hybrid

origins resulting from crossing between *O. tubicula* subsp. *tubicula* and *O. hartwegii* subsp.

hartwegii (Fig. 3a, S10). This is consistent with Towner's interpretation of this taxon, which he

hypothesized may represent a stabilized derivative of introgression between *O. tubicula* subsp.

tubicula and *O. hartwegii* subsp. *hartwegii* (Towner 1977). Therefore, the placement of *O.*

tubicula subsp. *strigulosa* as sister to the rest of the *O. hartwegii* species complex in our trees

may result from past gene flow and hence may not represent independent origins of bee

pollination in subsect. *Salpingia*. These results underscore the importance of explicitly including

tests for hybridization in phylogenetic studies. In the case of these data, estimating a species tree

given a set of gene trees within a coalescent framework without considering other non-ILS

sources of signal conflict could artificially inflate the number of inferred evolutionary transitions.

Figure 3

Our HyDe results also suggest that at least some of the morphological intermediacy and overlap among taxa in the group is due to continued, or at least recent, gene flow. For example, both *O. hartwegii* subsp. *pubescens* individuals that are inferred to have significant levels of admixture were collected from morphologically intermediate populations of *O. hartwegii* subsp. *pubescens* and *O. hartwegii* subsp. *hartwegii*. In addition, *O. hartwegii* subsp. *hartwegii* was a parent in all three hybridization events (S10). Thus, gene flow may explain this taxon's non-monophyly in our summary coalescent results. However, despite the often confounding patterns of overlapping morphological variation among closely related taxa in subsect. *Salpingia*, this pattern does not necessarily appear to be the result of admixture, as many of the tests for hybridization based on field observations were not significant (Fig 3a, S10). What is also clear from these results is that much like collecting hundreds of nuclear genes provides a more

nuanced picture of phylogenetic signal and taxon relationships, our results show that collecting

multiple individuals from across the geographic and morphological ranges is necessary for a

more complete picture of relationships among closely related taxa.

After filtering, we extracted a set of 9,728 single nucleotide polymorphisms (SNPs) from

both coding and non-coding regions. A Discriminant Analysis of Principal Components (DAPC;

Fig. 3b) using these data clearly distinguishes *Oenothera* subsect. *Salpingia* from *O.* subsect.

Calylophus, with *O. toumeyi* intermediate between the two, which is consistent with the

phylogenetic results presented here. Additionally, the DAPC identifies *O. lavandulifolia* as a

distinct genetic cluster from the remaining taxa in subsection *Salpingia*. The overlap between

taxa, for example between the remaining taxa in subsection *Salpingia*, is consistent with the high

levels of gene tree discordance identified by PhyParts (Fig. 3a). For this latter group of taxa, we

computed estimates of ancestry coefficients using snmf, which suggests a substantial amount of

shared ancestral polymorphisms while also showing some evidence of clear genetic structure

among taxa (Fig. 3c). Consistent with the phylogenetic analyses, there does not appear to be any

clear genetic distinction between *O. hartwegii* subsp. *hartwegii* and *O. hartwegii* subsp.

maccartii, whereas *O. hartwegii* subsp. *fendleri*, *O. tubicula* subsp. *tubicula,* and *O. hartwegii*

subsp. *filifolia* appear to be largely distinct.

Morphological Analysis

We conducted morphometric Principal Components Analysis (PCA) to determine if

morphological patterns were consistent with phylogentic results and to examine if specific

characters could be used to diagnose taxa as circumscribed by our phylogenetic analysis. Towner

(1977) observed overlapping and confounding patterns of morphological variation among taxa

within subsections, particularly within the *O. hartwegii* species complex. Despite this, because

some taxa (e.g., *O. hartwegii* subsp. *fendleri*) were strongly supported by our summary

coalescent trees we expected that they would be well distinguished in morphometric analysis.

Figure 4

The main traits that separated taxa in subsect. *Salpingia* were leaf traits and plant size, while

in subsect. *Calylophus* the main traits that separated taxa were sepal traits. In subsect. *Salpingia*,

PC1 accounted for 43.1% of variance in PCA, while PC2 accounted for 28.5% (Fig. 4b).

Morphological characters most associated with PC1 were leaf width (distal and basal), plant

height, and leaf length/width ratio (distal and basal). Those associated with PC2 were leaf length

(distal and basal), sepal length, and sepal tip length (Fig. 4b). In subsect. *Calylophus*, PC1

accounted for 44.4% of explained variance and PC2 accounted for 26.2% (Fig. 4c). The

characters most associated with PC1 in subsect. *Calylophus* include leaf length (distal and basal),

sepal length, and sepal tip length. Those most associated with PC2 were leaf length/width ratio

(distal and basal) and distal leaf width (Fig. 4c).

Our results support Towner's understanding of taxon boundaries by underscoring previous

difficulties in identifying individuals in this group based on morphology (Towner 1977). We

found substantial overlapping morphological variation among currently recognized taxa in both

subsections, though some taxa exhibited better grouping than others. The amount of overlap

between taxa was not a function of the strength of tree support for a given taxon in our summary

coalescent results. For example, *O. hartwegii* subsp. *fendleri*, a taxon that was well supported in

our summary coalescent trees, exhibited some of the highest degree of overlap with other taxa in

PCA space. Conversely, both *O. hartwegii* subsp. *filifolia* and *O. hartwegii* subsp. *pubescens*,

two taxa that formed poorly supported clades in our trees, formed clusters on the outer edges of

PCA space and had less overlap than other taxa (Fig. 4b). Interestingly, *O. hartwegii* subsp.

hartwegii, the taxon that was identified as a parent in all three instances of admixture in our

HyDe analysis, also overpas morphologically with most other taxa in subsect. *Salpingia* (Fig.

4b). This is not surprising given that it is widely distributed in northern Mexico and western

Texas and frequently comes into contact with related taxa resulting in sympatric populations and

occasional morphologically intermediate populations.

Implications for Reproductive Systems and Edaphic Endemism

Our results show that shifts from hawkmoth to bee pollination likely occurred twice in

sect. *Calylophus* (S3) and thus may be more common in *Oenothera* than previously thought. The

strongly supported sister relationship of *O. toumeyi* to remaining subsect. *Calylophus* in our

summary coalescent results is consistent with two independent shifts to bee pollination, once in

the ancestor of subsect. *Calylophus*, and another in subsect. *Salpingia* on the branch leading to

O. tubicula (Fig. 2). Independent shifts to bee pollination from hawkmoth pollination are perhaps

not surprising considering that within sect. *Calyophus*, hawkmoth-pollinated floral forms exhibit

variation between populations in hypanthium length and diameter and do not prevent occasional

pollination by bees (Lewis 2015; Towner 1977). Hawkmoth-pollinated taxa in sect *Calylophus*

exhibit vespertine anthesis, which separates them temporally from diurnal bees, but variation in

the timing of anthesis is also common between populations (Towner 1977), and hawkmoths are

documented to vary greatly in abundance spatiotemporally (Miller 1981; Campbell et al. 1997;

Artz et al. 2010). In cases of pollen limitation, night-blooming plants benefit from bimodal

pollination between moths and bees by acquiring pollinator assurance against yearly variation or

local extinction of specific pollinators, as shown for other *Oenothera* species (Barthell and

Knops 1997; Artz et al. 2010), *Lonicera japonica* (Miyake and Yahara 1998) and for night-

blooming *Ancistrophora* cacti (Schlumpberger et al. 2009).

Additionally, it has been shown that florivore-mediated selection drives floral trait shifts

in sect. *Calylophus* towards bee-pollinated floral forms (Jogesh et al. 2017; Bruzzese et al. 2019).

Variation in reproductive traits that allows some continued pollination by bees provides an

alternative mode of pollen transfer and may represent a mechanism for ensuring pollination.

While studies have shown that premating barriers contribute greatly to reproductive isolation

(Stanton et al. 2016), our results show that multiple, independent shifts from hawkmoth to bee

pollination and associated morphological changes, such as the shorter floral tube length of bee

pollinated flowers, may occur in sect. *Calylophus*, and hence may not be a particularly reliable

character for diagnosing taxa in this group.

Stochastic mapping (supplemental) suggests that there are multiple origins of permanent

translocation heterozygosity (PTH) in sect. *Calylophus* While ring chromosomes are common

and found in all taxa in sect. *Calylophus*, PTH is currently known from only one taxon, *O.*

serrulata. Because neighboring populations of *O. serrulata* and its putative progenitor O.

capillifolia subsp. *berlandieri* often resemble each other phenetically, Towner (1977)

hypothesized that *O. serrulata* may have originated multiple times through independent origins

of translocation heterozygosity in different geographic regions, and may be best recognized as "a

complex assemblage of populations having a common breeding system." However, this has

never before been explored in a phylogenetic context, nor has it been clearly demonstrated with

phylogenetic studies in Onagraceae. In our summary coalescent trees, all currently recognized

taxa in subsect. *Calylophus* were paraphyletic and *O. serrulata* was scattered throughout the

subsection (Fig. 2, S6, S7). Although support values are not always high for the positions of

various individuals of *O. serrulata*, there is at least one well defined, well supported split among

populations of *O. serrulata*. In our summary coalescent trees, the two *O. serrulata* accessions

from south Texas (*MJM970* & *MJM983*) grouped with other south Texas populations of *O.*

capillifolia subsp. *berlandieri* with generally strong support (Fig. 2, S6, S7). This relationship

was supported in PCA space as well, where *MJM983* was morphologically more similar to the

south Texas *O. capillifolia* subsp. *berlandieri* accessions than to other *O. serrulata* (Fig 4C). Our

results are therefore consistent with an independent origin of PTH in coastal Texas populations

of *O. serrulata*, demonstrating at minimum two origins of PTH (see *Taxonomic Implications*

below). However we cannot rule out other independent origins of PTH. Given the prevalence of

translocations among partial sets (i.e. not all seven) of homologous chromosomes in *O.* section

Calylophus (Towner 1977), and if this could be considered an intermediate step towards

"complete" PTH, perhaps it is not surprising to reconstruct multiple origins. Our results suggest

that a more complete assessment of the extent and distribution of this phenomenon in section

Calylophus is needed.

Independent origins of gypsum endemism in sect. *Calylophus* are also supported by our

analyses (supplemental). Edaphic specialization is a fundamental driver of speciation in plants

and contributes greatly to endemism and species diversity in areas with geologically distinct

substrates such as gypsum and serpentine outcrops (Kruckeberg 1984; Anacker et al. 2011;

Cacho and Strauss 2014; Moore et al. 2014). To date, two gypsum endemic taxa have been

described in sect. *Calylophus*, one in each subsection: *O. hartwegii* subsp. *filifolia*, which is

relatively widespread on gypsum in New Mexico and trans-Pecos Texas and only rarely

sympatric with other taxa, and the recently described *O. gayleana*, which is found in

southeastern New Mexico and adjacent western Texas, with disjunct populations in northern

Texas and western Oklahoma (Turner and Moore 2014). Despite low support and low gene tree

congruence in our analyses, the two gypsum endemic taxa had moderate sCF support (Fig. 2,

S9), much like other taxa with similarly low support and high levels of discordance. In

addition,while both gypsum endemics overlapped with other taxa in the morphometric analysis,

they occupied morphological extremes in PCA space (Fig. 4). Given that other well-supported

taxa also overlap morphologically, it is perhaps not surprising that the two gypsum endemic taxa

are not more differentiated from other taxa morphologically. Perhaps the strongest evidence in

our data for their recognition as distinct taxa is that we found no evidence of admixture between

either of these gypsum endemic taxa and other closely related taxa (Fig. 3a, S10).

Taxonomic Implications

The most consequential taxonomic result that arises from our analyses is the position of

O. toumeyi, a member of subsect. *Salpingia* as circumscribed by Towner (1977). In the present

study, this species is resolved as sister to subsect. *Calylophus* with strong support, rendering

subsect. *Salpingia* paraphyletic (Fig. 2, S9). Towner (1977) grouped *O. toumeyi* with *O.*

hartwegii due to similar floral and bud characters including large flowers and long floral tubes

suggestive of hawkmoth pollination, and rounded buds with long, free sepal-tips. Because the

breeding system is a defining difference in the current circumscription between the two

subsections in sect. *Calylophus*, this result supports abandoning subsections altogether in sect.

Calylophus.

434   Within subsect. *Salpingia* our results also suggest the need for revision. While our

phylogenetic analyses strongly support the current circumscription of *O. lavandulifolia* (sensu

Towner 1977) as a distinct species within subsect. *Salpingia* (Fig. 2, S9), the relationships of the

other two species *O. hartwegii* and *O. tubicula* are less clear. Towner (1977) differentiated these

two species by the breeding system and grouped the five subspecies of *O. hartwegii* together

based on a pattern of reticulate and intergrading variation in which taxa were distinguished from

one another by often slight differences in pubescence and leaf shape. Our morphometric analysis

confirmed this pattern; however, our phylogenetic results indicated that one taxon, *O. hartwegii*

subsp. *fendleri*, shares a closer relationship with the bee pollinated *O. tubicula* subsp. *tubicula*

than other taxa in the hawkmoth pollinated *O. hartwegii* species complex (Fig. 2, S6, S7). This

relationship was strongly supported and renders *O. hartwegii*, according to the current

circumscription, paraphyletic (Towner 1977). Based on strong phylogenetic support for this

clade, and its strong morphological distinctiveness as described by Towner (1977), we suggest

that *O. hartwegii* subsp. *fendleri* be elevated to the species rank along with both subspecies of *O.*

tubicula which were equally well supported in phylogenetic analysis and are geographically

isolated. Furthermore, our results support a the possible elevation to species rank for *O.*

hartwegii subsp. *filifolia*. While this taxon was poorly supported in our summary coalescent trees

(Fig. 2, S6, S7), we found no evidence of hybridization between this taxon and other closely

related taxa. In addition, *O. hartwegii* subsp. *fillifolia* is restricted to gypsum. Therefore, we

believe that the ecological distinctiveness and lack of gene flow of *O. hartwegii* subsp. *filifolia*

with other taxa in the *O. hartwegii* species complex warrants its elevation as a distinct species. In

light of these changes, and to maintain consistency in classification in the subsection, we feel

that despite the evidence of hybridization of *O. hartwegii* subsp. *pubescens* with *O. hartwegii*

subsp. *hartwegii*, it possesses a morphological distinctiveness that is supported by our

phylogenetic results. We therefore recommend *O. hartwegii* subsp. *pubescens* be elevated to the

species level, while *O. hartwegii* subsp. *hartwegii* and *O. hartwegii* subsp. *maccartii* be retained

as is, forming a polytypic species with two subspecies.

In contrast to the relatively clear divisions among taxa in subsect. *Salpingia,* none of the

four currently recognized taxa in the subsect. *Calylophus* were consistently recovered as

monophyletic. For example, *O. capillifolia* subsp. *capillifolia* was monophyletic in our exon-

only summary coalescent tree, but not in the "supercontig" tree, and *O. capillifolia* subsp.

berlandieri and *O. serrulata* were scattered throughout sect. *Calylophus* in both trees, perhaps

suggesting widespread gene flow and/or multiple origins of PTH (Fig. 2, S6). Importantly, our

results suggest that the circumscription of *O. gayleana* sensu Turner and Moore (2014) should be

amended. Specifically, we find that the populations of subsect. *Calylophus* from northern Texas

and western Oklahoma that were assigned to *O. gayleana* by Turner and Moore (2014; *MJM790-*

5, BJC71) may instead may belong to *O. serrulata* based on both their phylogenetic positions

(Fig. 2, S6, S7) and reduced pollen fertility (S12). These north Texas/western Oklahoma

populations seem to represent slightly narrower-leaved individuals of *O. serrulata*, which is a

common inhabitant of the extensive gypsum outcrops of this area (although it is not restricted to

gypsum there).

Finally, our results highlight an unrecognized cryptic taxon within *O. capillifolia* formed

by southern Texas coastal populations currently recognized as *O. capillifolia* subsp. *berlandieri*.

Towner (1977) described *O. capillifolia* as a polytypic species with two well-differentiated

morphological races. Though he noted the geographic and cytological distinction of the southern

Texas coastal populations of *O. capillifolia* subsp. *berlandieri*, these populations were included

in *O. capillifolia* subsp. *berlandieri* primarily because of completely overlapping morphological

variation. In our results, this cryptic clade of southern Texas coastal populations of *O. capillifolia*

subsp. *berlandieri* is the most phylogenetically well supported clade in subsect. *Calylophus* and

therefore may warrant taxonomic distinction based on our data (Fig. 2, S9r). Similarly, the

southern Texas coastal populations of *O. serrulata*, which is likely an independent origin of PTH

derived from this cryptic southern Texas coastal clade of *O. capillifolia* subsp. *berlandieri*, are

ecologically distinctive and geographically disjunct from other *O. serrulata* (occurring in coastal

dunes, unlike other populations in western Texas, Oklahoma, and northern Texas). In the past

they were considered distinctive enough to be described as a species, *Calylophus australis*

(Towner & Raven 1970). However, Towner (1977) later combined this species with *O. serrulata*

based on his decision to treat all PTH populations as *O. serrulata*. Combined with our results

here and the ecogeographic distinctiveness consistent with an independent origin of PTH in

coastal Texas, we believe this taxon also warrants recognition as a second PTH species in

Oenothera sect. *Calylophus*.

CONCLUSIONS

Here we describe a robust example of resolving a recent, rapid radiation using multiple

sources of evidence: (1) extensive sampling from populations throughout the geographic and

morphological range, (2) target enrichment for hundreds of nuclear genes, (3) the inclusion of

flanking non-coding regions, (4) gene tree-based hybridization inference, (5) SNPs extracted

from target enrichment data, and (6) morphometrics. Our results indicate that in recently radiated

species complexes with low sequence divergence and/or high levels of ILS that could be an

intractable problem with traditional loci, the use of targeted enrichment in addition to flanking

non-coding regions provides a net benefit and is essential to recover species-level resolution. Our

results also underscore the importance of summary coalescent methods and evaluating gene tree

discordance for resolving historical relationships in recalcitrant groups. By explicitly testing for

hybridization using gene tree approaches, we also demonstrate that the estimated number of

character state transitions may be artifactually inflated if hybridization is not taken into account.

This, in combination with morphometrics, provided key evolutionary insights where

relationships in summary coalescent methods may be obscured by gene flow. Importantly, our

study uncovers strong evidence for multiple origins of biologically important phenomena,

including the evolution of bee pollination, PTH, and edaphic specialization. Consequently,

Oenothera sect. *Calylophus* might represent a powerful system for understanding these

phenomena, especially with future genome sequence data.

DATA AVAILABILITY

Illumina reads generated for this study are available at the NCBI Sequence Read Archive under

BioProject PRJNA544074. Assembled exon and supercontig sequences, multiple sequence

alignments, and SNP files are available at http://dx.doi.org/10.5061/dryad.[NNNN].

ACKNOWLEDGEMENTS

Material for *O. serrulata* and *O. capillifolia* subsp. *berlandieri* that were used for designing

probes for target enrichment (1KP accession codes SJAN and EQTY, respectively) was

originally collected by R.A.R.; samples were grown, and RNA was extracted, in the lab of M.

Johnson and subsequently sequenced by the One Thousand Plant Transcriptomes (1KP)

initiative. We thank the following for access to field sites and permission to collect samples: U.S.

Bureau of Land Management (Colorado, New Mexico, Utah), U.S.D.A. Forest Service (Regions

2, 3, and 4), Carlsbad Caverns National Park, Guadalupe Mountains National Park, Big Bend

National Park, Palo Duro State Park, and White Sands Missile Range. We thank the Billie L.

Turner Plant Resources Center at the University of Texas at Austin for access to herbarium

collections, and we thank the following persons for help with collections: David Anderson,

George S. Hinton, Nidia Mendoza Díaz, Patrick Alexander, Christopher Martine, Rebecca

Drenovsky, Clare Muller, Jeffrey Sanders, Anna Brunner, Joseph Charboneau, Heather-Rose

Kates, and Sophia Weinmann. Finally, we thank Elliot Gardner for providing invaluable advice

during target enrichment and sequencing, and Daniel Bruzzese for help with collections and for

designing the flower silhouette used to signify PTH in figure 1 (available at www.phylopic.org).

FUNDING SOURCES

538   This work was supported by National Science Foundation grants DEB-1342873 to KAS, JBF,

539   and NJW, and DEB-1054539 to MJM. Additional support was provided by the National

540   Geographic Society, Oberlin College, the Nauganee Institute of the Chicago Botanic Garden,

541   The Shaw Fellowship, New Mexico Native Plant Society, American Society of Plant

542   Taxonomists, and the Society of Herbarium Curators.

543
544   REFERENCES
545

546   Alexander M.P. 1969. Differential Staining of Aborted and Nonaborted Pollen. Stain Technol.
547   44:117–122.
548   Alexander M.P. 1980. A Versatile Stain for Pollen Fungi, Yeast and Bacteria. Stain Technol.
549   55:13–18.
550   Anacker B.L., Whittall J.B., Goldberg E.E., Harrison S.P. 2011. Origins and consequences of
551   serpentine endemism in the California flora. Evolution (N. Y). 65:365–376.
552   Artz D.R., Villagra C. A., Raguso R. A. 2010. Spatiotemporal variation in the reproductive
553   ecology of two parapatric subspecies of *Oenothera cespitosa* (Onagraceae). Am. J. Bot.
554   97:1498–510.
555   Barrett S., Harder L., Worley A. 1996. The comparative biology of pollination and mating in
556   floweing plants. Philos. Trans. R. Soc. B Biol. Sci. 351:1271–1280.
557   Barrett S.C.H. 2013. The evolution of plant reproductive systems: how often are transitions
558   irreversible? Proc R Soc B. 280:20130913.
559   Barthell J. F., Knops J. M. 1997. Visitation of evening primrose by carpenter bees: evidence of
560   a" mixed" pollination syndrome. Southwest. Nat. 86-93.
561   Blischak P.D., Chifman J., Wolfe A.D., Kubatko L.S. 2018. HyDe: A python package for
562   genome-scale hybridization detection. Syst. Biol. 67:821–829.
563   Boberg E., Alexandersson R., Jonsson M., Maad J., Ågren J., Nilsson L.A. 2014. Pollinator
564   shifts and the evolution of spur length in the moth-pollinated orchid *Platanthera bifolia*.
565   Ann. Bot. 113:267–275.
566   Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina
567   sequence data. Bioinformatics. 30:2114–2120.
568   Brady K.U., Kruckeberg A.R., Bradshaw H.D. 2005. Evolutionary ecology of plant adaptation to
569   serpentine soils. Annu. Rev. Ecol. Evol. Syst. 36:243–266.
570   Bruzzese D.J., Wagner D.L., Harrison T., Jogesh T., Overson R.P., Wickett N.J., Raguso R.A.,
571   Skogen K.A. 2019. Phylogeny, host use, and diversification in the moth family Momphidae
572   (Lepidoptera: Gelechioidea). PLoS One. 14:e0207833.
573   Bryson R.W., Linkem C.W., Dorcas M.E., Lathrop A., Jones J.M., Alvarado-Díaz J., Grünwald
574   C.I., Murphy R.W. 2014. Multilocus species delimitation in the *Crotalus triseriatus* species
575   group (serpentes: Viperidae: Crotalinae), with the description of two new species. Zootaxa.
576   3826:475–496.
577   Cacho N.I., Strauss S.Y. 2014. Occupation of bare habitats, an evolutionary precursor to soil
578   specialization in plants. Proc. Natl. Acad. Sci. 111:15132–15137.
579   Campbell D.R.., Waser N.M.., Melendez-Ackerman E.J.. 1997. Analyzing pollinator-mediated
580   selection in a plant hybrid zone: Hummingbird visitation patterns on three spatial scales.
581   Am. Nat. 149:295–315.
582   Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model.
583   Bioinformatics. 30:3317–3324.
Chifman J., Kubatko L. 2015. Identifiability of the unrooted species tree topology under the

coalescent model with time-reversible substitution processes, site-specific rate variation,
and invariable sites. J. Theor. Biol. 374:35–47.
Christie M.R., Knowles L.L. 2015. Habitat corridors facilitate genetic resilience irrespective of
species dispersal abilities or population sizes. Evol. Appl. 8:454-463.
Crepet W.L., Niklas K.J. 2009. Darwin's second "abominable mystery": Why are there so many
angiosperm species? Am. J. Bot. 96:366–81.
Duarte J.M., Wall P.K., Edger P.P., Landherr L.L., Ma H., Pires J.C., Leebens-Mack J.,
dePamphilis C.W. 2010. Identification of shared single copy nuclear genes in *Arabidopsis,*
Populus, Vitis and *Oryza* and their phylogenetic utility across various taxonomic levels.
BMC Evol Biol. 10:61.
Eckert A.J., Carstens B.C. 2008. Does gene flow destroy phylogenetic signal? The performance
of three methods for estimating species phylogenies in the presence of gene flow. Mol.
Phylogenet. Evol. 49:832–842.
Ehrlich, P. R., & Raven, P. H. (1969). Differentiation of populations. Science. 165:1228–1232.
Folk R.A., Mandel J.R., Freudenstein J. V. 2015. A protocol for targeted enrichment of intron-
containing sequence markers for recent radiations: A phylogenomic example from
Heuchera (Saxifragaceae). Appl. Plant Sci. 3:1500039.
Frichot E, Mathieu F, Trouillon T, Bouchard G and François O (2014) Fast and efficient
estimation of individual ancestry coefficients. Genetics. 196:973-983.
Gerard D., Gibbs H.L., Kubatko L. 2011. Estimating hybridization in the presence of
coalescence using phylogenetic intraspecific sampling. BMC Evol. Biol. 11:291.
Giarla T.C., Esselstyn J.A. 2015. The challenges of resolving a rapid, recent radiation: Empirical
and simulated phylogenomic of Philippine shrews. Syst. Biol. 64(5):727-740.
Green R., Krause J., Briggs A., Rasilla Vives M., Fortea Pérez F. 2010. A draft sequence of the
neandertal genome. Science. 328:710–722.
Heyduk K., Trapnell D.W., Barrett C.F., Leebens-mack J.I.M. 2016. Phylogenomic analyses of
species relationships in the genus *Sabal* (Arecaceae) using targeted sequence capture. Biol.
J. Linn. Soc. 117(1):106-120.
Hollister J.D., Greiner S., Johnson M.T.J., Wright S.I. 2019. Hybridization and a loss of sex
shape genome-wide diversity and the origin of species in the evening primroses (Oenothera,
Onagraceae). New Phytol. 224:1372–1380.
Jogesh T., Overson R.P., Raguso R.A., Skogen K.A. 2017. Herbivory as an important selective
force in the evolution of floral traits and pollinator shifts. AoB Plants. 9(1):plw088.
Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett
N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics from high-
throughput sequencing reads using target enrichment. Appl. Plant Sci. 4:1600016.
Johnson M.T.J., Smith S.D., Rausher M.D. 2009. Plant sex and the evolution of plant defenses
against herbivores. Proc. Natl. Acad. Sci. 106:18079–18084.
Jombart, T. (2008) adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics 24:1403-1405.
Jombart T, Devillard S and Balloux, F (2010). Discriminant analysis of principal components: a
new method for the analysis of genetically structured populations. BMC Genetics. 11:94.
Katinas L., Crisci J., Wagner W., Hoch P. 2004. Geographical diversification of tribes
Epilobieae, Gongylocarpeae, and Onagreae (Onagraceae) in North America, based on
parsimony analysis of endemicity and track compatibility analysis. Ann. Missouri Bot.
91:159–185.

Knowles L.L. 2009. Estimating species trees: Methods of phylogenetic analysis when there is
incongruence across genes. Syst. Biol. 58:463–467.
Knowles L.L., Chan Y.-H. 2008. Resolving species phylogenies of recent evolutionary
radiations. Ann. Missouri Bot. Gard. 95:224–231.
Kruckeberg A. 1984. California Serpentines: Flora, Vegetation, Geology, Soils, and
Management Problems. Berkeley: Univ of California Press.
Kubatko L.S., Chifman J. 2019. An invariants-based method for efficient identification of hybrid
species from large-scale genomic data. BMC Evol. Biol. 19:1–13.
Leaché A.D., Harris R.B., Rannala B., Yang Z. 2014. The influence of gene flow on species tree
estimation: a simulation study. Syst. Biol. 63:17–30.
Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored hybrid enrichment for massively
high-throughput phylogenomics. Syst. Biol. 61:727–744.
Levin R., Wagner W., Hoch P. 2004. Paraphyly in Tribe Onagreae : Insights into Phylogenetic
Relationships of Onagraceae Based on Nuclear and Chloroplast Sequence Data. Syst. Bot.
29:147–164.
Lewis, Emily. 2015. Differences in Population Genetic Structure of Hawkmoth and Bee-
Pollinated Species of *Oenothera* (Onagraceae) Are More Pronounced at a Landscape
Scale. Northwestern University Libraries. Masters Thesis. Northwestern University.
Maddison W.P., Knowles L. 2006. Inferring Phylogeny Despite Incomplete Lineage Sorting.
Evol. Biol. 55:21–30.
Mamanova L., Coffey A.J., Scott C.E., Kozarewa I., Turner E.H., Kumar A., Howard E.,
Shendure J., Turner D.J. 2010. Target-enrichment strategies for next- generation
sequencing. Nat. Methods. 7:111–118.
Mandel J.R., Dikow R.B., Funk V. a, Masalia R.R., Staton S.E., Kozik A., Michelmore R.W.,
Rieseberg L.H., Burke J.M. 2014. A target enrichment method for gathering phylogenetic
information from hundreds of loci: An example from the Compositae. Appl. Plant Sci. 2:1–
6.
Meng C., Kubatko L.S. 2009. Detecting hybrid speciation in the presence of incomplete lineage
sorting using gene tree incongruence: A model. Theor. Popul. Biol. 75:35–45.
Miller R.B.. 1981. Hawkmoths and the Geographic Patterns of Floral Variation in *Aquilegia*
caerulea. Evolution (N. Y). 35:763–774.
Minh B.Q., Hahn M.W., Lanfear R. 2020. New methods to calculate concordance factors for
phylogenomic datasets. Mol. Biol. Evol. 37(9):2727-2733.
Mirarab S., Warnow T. 2015. ASTRAL-II: Coalescent-based species tree estimation with many
hundreds of taxa and thousands of genes. Bioinformatics. 31:i44–i52.
Miyake T., Yahara T. 1998. Why does the flower of *Lonicera japonica* open at dusk?. Can. J.
Bot. *76*(10):1806-1811.
Moore M., Jansen R. 2007. Origins and biogeography of gypsophily in the Chihuahuan Desert
plant group *Tiquilia* Subg. subg. *Eddya*. Syst. Bot. 32:392–414.
Moore M.J., Mota J.F., Douglas N.A., Flores Olvera H., Ochoterena H. 2014. Ecology assembly
evolution gypsophile floras. In: Rajakaruna N., Boyd R.S., Harris T.B., editors. Plant
Ecology and Evolution in Harsh Environments. New York: Nova Science Publishers, Inc. p.
97–128.
Nason J.D., Hamrick J.L., Fleming T.H. 2002. Historical vicariance and postglacial colonization
effects on the evolution of genetic structure in *Lophocereus*, a Sonoran Desert columnar
cactus. Evolution 56:2214–2226.

Van der Niet T., Johnson S.D. 2012. Phylogenetic evidence for pollinator-driven diversification
of angiosperms. Trends Ecol. Evol. 27:353–361.
van der Niet T., Johnson S.D., Linder H.P. 2006. Macroevolutionary data suggest a role for
reinforcement in pollination system shifts. Evolution 60:1596–1601.
Van Der Niet T., Peakall R., Johnson S.D. 2014. Pollinator-driven ecological speciation in
plants: New evidence and future perspectives. Ann. Bot. 113:199–211.
Peakall R., Ebert D., Poldy J., Barrow R.A., Francke W., Bower C.C., Schiestl F.P. 2010.
Pollinator specificity, floral odour chemistry and the phylogeny of Australian sexually
deceptive *Chiloglottis* orchids: implications for pollinator-driven speciation. New Phytol.
188:437–50.
Rajakaruna N. 2004. The Edaphic Factor in the Origin of Plant Species. Int. Geol. Rev. 46:471–
478.
Raven P.H. 1964. Catastrophic Selection and Edaphic Endemism. Evolution 18:336–338.
Raven P.H. 1979. A survey of reproductive biology in Onagraceae. New Zeal. J. Bot. 17:575–
593.
R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
Roch S., Steel M. 2015. Likelihood-based tree reconstruction on a concatenation of aligned
sequence data sets can be statistically inconsistent. Theor. Popul. Biol. 100:56–62.
Sayyari E., Mirarab S. 2016. Fast coalescent-based computation of local branch support from
quartet frequencies. Mol. Biol. Evol. 33(7):1654-68.
Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27(4):592-593.
Schlumpberger B. O., Cocucci A. A., Moré M., Sérsic A. N., Raguso R. A. 2009. Extreme
variation in floral characters and its consequences for pollinator attraction among
populations of an Andean cactus. Ann. Bot. *103*(9):1489-1500.
Skogen K.A., Overson R.P., Hilpman E.T., Fant J.B. 2019. Hawkmoth pollination facilitates
long-distance pollen dispersal and reduces isolation across a gradient of land-use change.
Ann. Missouri Bot. Gard. 104:495–511.
Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals
conflict, concordance, and gene duplications with examples from animals and plants. BMC
Evol. Biol. 15:150.
Stamatakis A. 2014. Stamatakis - 2014 - RAxML version 8 a tool for phylogenetic analysis and
post-analysis of large phylogenies. 2010–2011.
Stanton K., Valentin C.M., Wijnen M.E., Stutstman S., Palacios J.J., Cooley A.M. 2016.
Absence of postmating barriers between a selfing vs. Outcrossing chilean *Mimulus* species
pair. Am. J. Bot. 103:1030–1040.
Stebbins G.L. 1970. Adaptive Radiation of Reproductive Characteristics in Angiosperms, I:
Pollination Mechanisms. Annu. Rev. Ecol. Syst. 1:307–326.
Stephens J.D., Rogers W.L., Heyduk K., Cruse-Sanders J.M., Determann R.O., Glenn T.C.,
Malmberg R.L. 2015. Resolving phylogenetic relationships of the recently radiated
carnivorous plant genus *Sarracenia* using target enrichment. Mol. Phylogenet. Evol. 85:76–
87.
Stockhouse R.E.I. 1973. Biosystematic Studies of *Oenothera* L. Subgenus *Pachylophus*. Univ.
Microfilm. Ph.D. thesis. Colorado State University.
Straub S.C.K., Parks M., Weitemier K., Fishbein M., Cronn R.C., Liston A. 2012. Navigating the
tip of the genomic iceberg: Next-generation sequencing for plant systematics. Am. J. Bot.

99:349–64.
Swofford, D. L. 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods).
Version 4. Sinauer Associates, Sunderland, Massachusetts.
Thomson J.D., Wilson P. 2008. Explaining Evolutionary Shifts between Bee and Hummingbird
Pollination: Convergence, Divergence, and Directionality. Int. J. Plant Sci. 169:23–38.
Towner H.F., Raven P.H. 1970. A new species and some new combinations in *Calylophus*
(Onagraceae). Madrono. 20: 241-245.
Towner H.F. 1977. The Biosystematics of *Calylophus* (Onagraceae). Ann. Missouri Bot. Gard.
64:48–120.
Tripp E.A., Manos P.S. 2008. Is floral specialization an evolutionary dead-end? Pollination
system transitions in *Ruellia* (Acanthaceae). Evolution 62:1712–1737.
Turner B.L., Moore M.J. 2014. *Oenothera gayleana* (*Oenothera* sect. *Calylophus*, Onagraceae),
a new gypsophile from Texas, New Mexico, and Oklahoma. Phytologia.org. 96:200–206.
Wagner W., Hoch P., Raven P. 2007. Revised Classification of the Onagraceae. Syst. Bot.
Monogr. 83:1–240.
Wagner W.L. 2021. In press. In: Flora of North America Editorial Committee, eds. 1993+.
Flora of North America North of Mexico, 20+ vols. New York and Oxford: Oxford
University Press.
Weitemier K., Straub S.C.K., Cronn R.C., Fishbein M., Schmickl R., McDonnell A., Liston A.
2014. Hyb-Seq: Combining Target Enrichment and Genome Skimming for Plant
Phylogenomics. Appl. Plant Sci. 2:1400042.
Wilson P., Wolfe A.D., Armbruster W.S., Thomson J.D. 2007. Constrained lability in floral
evolution: Counting convergent origins of hummingbird pollination in Penstemon and
Keckiella. New Phytol. 176:883–890.
Xi Z., Liu L., Rest J.S., Davis C.C. 2014. Coalescent versus Concatenation Methods and the
Placement of *Amborella* as Sister to Water Lilies. Syst. Biol. 63:919–932.
Xu S., Schlüter P.M., Scopece G., Breitkopf H., Gross K., Cozzolino S., Schiestl F.P. 2011.
Floral isolation is the main reproductive barrier among closely related sexually deceptive
orchids. Evolution. 65:2606–20.

FIGURE LEGENDS
753
Figure 1. (a) Range map of all taxa in *Oenothera* sect. *Calylophus* (based on Towner 1977).
Sampling locations of leaf tissue samples (points; color corresponds to taxa in cladogram to the
right [Figure 1b]) and estimated taxon ranges (polygons; colors correspond to Figure 1b)
proposed by Towner (1977) and Turner and Moore (2014). (b) Estimated cladogram of
Oenothera sect. *Calylophus* sensu Towner (1977) and Turner and Moore (2014). Symbols to the
right of tip labels signify pollination system (bee, hawkmoth, or Permanent Translocation
Heterozygosity [PTH]) and the symbol color specifies whether a given taxon is a gypsum
endemic (purple = no, pink = yes). Photo panels: (c) *Oenothera hartwegii* subsp. *fendleri* (d)
Oenothera hartwegii subsp. *filifolia* (e) *Oenothera hartwegii* subsp. *hartwegii* (f) *Oenothera*
hartwegii subsp. *maccartii* (g) *Oenothera hartwegii* subsp. *pubescens* (h) *Oenothera tubicula*
subsp. *tubicula* (i) *Oenothera lavandulifolia* (j) *Oenothera tubicula* subsp. *strigulosa* (k)
Oenothera capillifolia subsp. *berlandieri* (l) *Oenothera capillifolia* subsp. *capillifolia* (m)
Oenothera gayleana (n) *Oenothera serrulata* (o) *Oenothera toumeyi*

767
Figure 2. ASTRAL-II summary coalescent tree constructed using the "Supercontig" dataset with
100 bootstraps. At relevent nodes, piecharts represent Phyparts analysis (blue = concordant,
green = most frequent conflict, yellow = all other conflict, black = uninformative gene trees), top
row of support values are bootstrap values from ASTRAL-II, and gCF and sCF from IQTree
(BS/gCF/sCF), bottom two support values are number of concordant gene trees for the node and
total number of gene trees minus the number of concordant gene trees at that node
(concord/discord). Colored tip points correspond to taxon designation.
775
776   **Figure 3.** (a) Summary of HyDe Analysis annotated on ASTRAL-III species tree constructed
777   using the "Supercontig" dataset; black arrows represent direction of admixture detected by HyDe
778   anaylsis. (b) Discriminant Analysis of Principal Components based on a filtered set of SNPs
779   extracted from the entire supercontig dataset , and (c) sNMF plot of inferred ancestry coefficients
780   using the same set of filtered SNPs but limited to subsection *Salpingia* (minus *O. lavandulifolia*)
781
782   **Figure 4.** Morphometric Principal Components Analysis (PCA) using 9 morphological
783   characters (plant height, leaf length [basal and distal], leaf width [basal and distal], leaf
784   length/width ratio (basal and distal), and sepal tip length) for (a) Section *Calylophus* (b)
785   Subsection *Salpingia* without *O. toumeyi*, and (c) Subsection *Calylohus* with *O. toumeyi*
786   included.

 SUPPLEMENTAL MATERIAL

788 S1.
789 See excel table "S1 Accessions and Seq Stats"
790

791 S2.
792 Number of leaf tissue accessions sequenced from each taxon

Taxon	No. of Accessions
O. lavandulifolia	16
O. tubicula subsp. *strigulosa*	3
O. tubicula subsp. *tubicula*	9
O. hartwegii subsp. *fendleri*	15
O. hartwegii subsp. *filifolia*	44
O. hartwegii subsp. *hartwegii*	12
O. hartwegii subsp. *maccartii*	9
O. hartwegii subsp. *pubescens*	26
O. hartwegii	1
O. toumeyi	5
O. capillifolia subsp. *berlandieri*	17
O. capillifolia subsp. *capillifolia*	11
O. gayleana	9
O. serrulata	17
Outgroups:	
O. pilosella	1
O. organensis	1
O. primiveris	1
O. tubifera	1
O. triloba	1
O. cespitosa subsp. *cespitosa*	1
O. suffrutescens	1
Chylismia scapoidea subsp. *scapoidea*	1
Eulobus californicus	1
Total	203

S3.
MATERIALS AND METHODS
Taxon Sampling, DNA Extraction, and Determination of PTH
A total of 194 individuals spanning the geographic, morphological, and ecological ranges
of all 13 recognized taxa in *Oenothera* sect. *Calylophus* [following Towner (1977) and Turner
and Moore (2014)] were included in this study (Fig. 1a, S1, S2) along with eight outgroups
representing other major sections of Oenothera (*Eremia*, *Gaura*, *Kneiffia*, *Lavauxia*, *Oenothera*,
Pachylophus, and *Ravenia*) and other genera (*Chylismia* and *Eulobus*) in Onagraceae (S1, S2).
All leaf tissue samples were collected from individuals in the field between 2007 and 2015 and
voucher specimens were deposited at the United States National Herbarium (US), with

duplicates in most cases at either the Nancy Rich Poole Herbarium (CHIC) or the George T.

Jones Herbarium at Oberlin College (OC; S1). DNA was extracted from fresh, silica-dried leaf

tissue using either (1) a modified CTAB protocol (Doyle 1987), (2) the Nucleon PhytoPure DNA

extraction kit (GE Healthcare Life Sciences, Pittsburgh, Pennsylvania, USA), or (3) a modified

CTAB and silicon dioxide purification protocol (Doyle 1987; Sharma and Purohit 2012; See S13

for details) followed by passing any extractions retaining a brown or yellow coloration through a

Qiagen Qiaquick PCR spin column for additional purification according to the manufacturer's

protocol (Qiagen, Venlo, Netherlands). The third DNA extraction method was used for difficult

to extract, polysaccharide-rich leaf tissue samples that yielded gooey, discolored DNA following

initial extraction. PTH status was determined for individuals in subsect. *Calylophus* using floral

morphology and/or when flowers were present by assessing pollen fertility using a modified

Alexander stain, as PTH taxa have a demonstrated 50% reduction in pollen fertility (Towner

1977). For accessions identified as either *O. capillifolia* subsp. *berlandieri* or *O. serrulata* that

had sufficient pollen available, pollen was removed from flowers and stained using a modified

Alexander stain (Alexander 1969, 1980). Accessions with less than 50% viable pollen were

assigned to *O. serrulata*, the only currently recognized PTH taxon in subsect. *Calylophus*. Pollen

count data are provided in Supplement 12 (See S12 for details).

Bait Design, Library Construction, Target Enrichment, and Sequencing

We targeted 322 orthologous, low-copy nuclear loci determined by clustering

transcriptomes of *Oenothera serrulata* (1KP accession *SJAN*) and *Oenothera berlandieri* (1KP

accession *EQYT*) to select a subset of the 956 phylogenetically informative *Arabidopsis* loci

identified by Duarte et al. (2010). Transcriptomes of two *Oenothera* species, *O. serrulata* and *O.*

berlandieri, were assembled and optimal isoforms were filtered for the longest reading frame

using custom Perl scripts. Assembled transcripts were aligned as amino acids to the 956 TAIR

loci of *Arabidopsis* in TranslatorX (Abascal et al. 2010). This alignment identified 956

orthologous sequences, from which 322 loci were randomly selected. BLAST searches of amino

acid sequences from these loci were carried out to ensure orthology between the transcript loci

and the *Arabidopsis* TAIR locus. The bait set was designed from these 322 loci, which were

selected from both *O. serrulata* and *O. berlanderi* sequences. A set of 19,994 120-bp baits tiled

across each locus with a 60 base overlap (2x tiling) was manufactured by Arbor Biosciences

(formerly MYcroarray, Ann Arbor, Michigan, USA). Sequencing libraries for 67 samples were

prepared with the Illumina TruSeq Nano HT DNA Library Preparation Kit (San Diego,

California, USA) following the manufacturer's protocol, except using half volumes beginning

with the second addition of Dynabeads MyOne Streptavidin C1 magnetic beads (Invitrogen,

Carlsbad, CA, USA). DNA samples were sheared using a Covaris M220 Focused-Ultrasonicator

(Covaris, Woburn, Maryland, USA) to a fragment length of ~550 bp (for an average insert size

of ~420 bp). The remaining 134 libraries were constructed by Rapid Genomics (Gainesville,

Florida, USA), with custom adapters. The Illumina i5 and i7 barcodes were used for all libraries.

Libraries were enriched for these loci using the MyBaits protocol (ArborBiosciences

2016) with combined pools of libraries totaling 1.2 μg of DNA (12 libraries/pool at 100

ng/library). Libraries with less than 100 ng of total recovered DNA were pooled together in

equimolar concentrations using available product, resulting in some pools with less than 1.2 μg

of DNA. The smallest successful pool contained four samples with 6 ng of library each.

Hybridization was performed at 65°C for approximately 18 hours. The enriched libraries were

reamplified with 14 to 18 PCR cycles and a final cleanup was performed using a Qiagen

QiaQuick PCR cleanup kit following the manufacturer's protocol to remove bead contamination
(Qiagen, Venlo, Netherlands). DNA concentrations were measured using a Qubit 2.0
Fluorometer (Life Technologies, Carlsbad, California, USA) and molarity was measured on an
Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, California, USA). A final
cleaning step using Dynabeads MyOne Streptavidin C1 magnetic beads was performed on pools
with adapter contamination as detected on the Bioanalyzer. Pools were sequenced in four runs at
equimolar ratios (4 nM), on an Illumina MiSeq (2 x 300 cycles, v3 chemistry; Illumina, Inc., San
Diego, California, USA) at the Pritzker DNA lab (Field Museum, Chicago, IL, USA). This
produced 80,273,296 pairs of 300-bp reads. Reads were demultiplexed and adapters trimmed
automatically by Illumina Basespace (Illumina 2016). Raw reads have been deposited at the
NCBI Sequence Read Archive (BioProject PRJNA544074).
886
Quality Filtering, Assembly and Alignment
A summary of read quality from each sample was produced using FastQC
(http://www.bioinformatics.babraham.ac.uk/people.html), which revealed read-through adapter
contamination in many of the poorer quality samples. To remove read-through contamination
and filter for quality, reads were trimmed for known Illumina adapters using Trimmomatic
(Bolger et al. 2014) with the following settings:
ILLUMINACLIP:<illumina_adapters.fasta>:2:30:10 LEADING:10 TRAILING:10
SLIDINGWINDOW:4:20 MINLEN:20. Trimmed, quality-filtered reads were assembled using
the HybPiper pipeline (Johnson et al. 2016) with default settings, followed by the intronerate.py
script, to produce both exons and the "splash zone" flanking non-coding region-containing
supercontigs. Only pairs with both mates surviving trimming and quality filtering were used for
HybPiper.
To compare the influence of "splash-zone" non-coding regions, two sets of alignments
were created: (1) exons alone, and (2) coding sequences plus the "splash-zone" (Hereby referred
to as supercontigs). For multiple sequence alignments of exons alone, protein and nucleotide
sequences assembled in HybPiper were gathered into fasta files by gene. For protein sequences
only, stop codons were changed to "X" using a sed command-line regular expression to facilitate
alignment, and sequences were aligned using MAFFT with settings: --auto --adjustdirection --
maxiterate 1000 (Katoh et al. 2002). Aligned protein sequences were then used to fit unaligned
nucleotide sequences into coding frame alignments using pal2nal with default settings (Suyama
et al. 2006). In-frame, aligned DNA sequences were trimmed to remove low-coverage positions
and sequences composed only of gaps using TrimAl with the automated setting, which is
optimized for maximum likelihood analyses (Gutíerrez et al. 2009). For supercontigs, nucleotide
sequences assembled using HybPiper were gathered into fasta files by gene, gene names were
removed from fasta headers using a command-line regular expression, and sequences were
aligned in MAFFT with settings: --auto --adjustdirection --maxiterate 1000 (Katoh et al. 2002).
Reverse compliment tags ("_R_") were removed from taxon names using a command-line
regular expression, and sequences were trimmed using TrimAl with previously listed settings
optimized for maximum likelihood analyses (Gutíerrez et al. 2009). To minimize the effects of
missing data on phylogenetic analyses, accessions with < 50% of loci passing quality filtering
were removed, and genes that were recovered across < 70% of the total remaining samples were
also removed. Following quality filtering, we recovered 204 high quality loci (present in at least
70% of samples) and an average of 625,323 reads per sample (S1). Final assemblies are available
at Dryad (http://dx.doi.org/10.5061/dryad.[NNNN]). Bioinformatics pipelines and analyses were

run at the high-performance computing cluster ("Treubia") at the Chicago Botanic Garden unless
otherwise specified.
923
Phylogenetic Reconstruction
We conducted phylogenetic analyses using two strategies for each set of alignments.
Alignments were concatenated and analyzed using maximum likelihood (ML) in RAxML
(Stamatakis 2014; hereafter referred to as "concatenation"), whereas coalescent-based analyses
were conducted using ML gene trees in ASTRAL-II (Mirarab and Warnow 2015; Sayyari and
Mirarab 2016) and using unlinked SNPs in SVDquartets (Chifman and Kubatko 2014, 2015)
implemented in PAUP* beta version 4.0a168 (Swofford 2003; S3). In concatenation analyses,
after aligning each gene separately in MAFFT, genes were concatenated, partitioned, and
maximum likelihood trees were reconstructed in RAxML Version 8 (Stamatakis 2014) using the
GRTCAT model with 100 "rapid-boostrapping" psuedoreplicates and *Chylismia scapoidea* as
the outgroup, on the CIPRES Science Gateway computing cluster (Miller et al. 2010). For
coalescent analyses, individual gene trees were first estimated using RAxML Version 8
(Stamatakis 2014), with 100 "rapid-bootstraping" psuedo-replicates and settings: -p 12345 -x
12345 -N 100 -c 25 -f a -m GTRCAT –s, and *Chylismia scapoidea* as the outgroup. Gene trees
based on supercontigs were not partitioned by codon position. Coalescent-based analyses of
accessions were conducted in ASTRAL-II (Mirarab and Warnow 2015; Sayyari and Mirarab
2016) with default settings using the best RAxML gene trees and their associated bootstrap files
as input, and in SVD Quartets (Chifman and Kubatko 2014, 2015) with default settings using an
inframe aligned supermatrix with all 204 loci and supercontigs. ASTRAL-II and SVD Quartets
analysis was performed with 100 multi-locus bootstraps.
To assess concordance among gene trees and provide additional support complementary
to bootstrap values, we conducted two additional analyses. First, we assessed raw gene tree
concordance using Phyparts (Smith et al. 2015). Prior to running Phyparts, nodes with < 33%
support in the supercontig RAxML gene trees were collapsed using the sumtrees command in
Dendropy (Sukumaran 2010). These gene trees were then re-rooted using *Chylismia scapoidea*
as the outgroup and ASTRAL-II was rerun using these collapsed, re-rooted gene trees as the
input files. Pie charts showing gene tree discordance were generated and overlaid on the
resulting ASTRAL-II tree using the PhypartsPiecharts script
(https://github.com/mossmatters/phyloscripts/tree/master/phypartspiecharts). Phyparts piecharts
and gene tree concordance values were also added to Figure 1 by importing the two data files
produced by the Phypartspiecharts.py into R and manually matching them to key nodes on our
ASTRAL-II supercontig tree using *ggtree* version 1.14.6 (G Yu, DK Smith, H Zhu, Y Guan
2017). We also generated gene and site concordance factors for our ASTRAL-II tree constructed
using supercontigs in IQTree v1.7-beta16 (Minh et al. 2018). IQtree calculates the gene
concordance factor (gCF) and accounts for incomplete taxon coverage among gene trees and
therefore may provide a more accurate representation of agreement among gene trees than other
methods. In addition to gCF, IQTree calculates the site concordance factor (sCF), which is
defined as the percentage of decisive nucleotide sites supporting a specific node (Minh et al.
2018). We used the RAxML gene trees produced for the ASTRAL-II supercontig analysis, and
supercontig alignments themselves, as the inputs for IQtree. For computing sCF, we randomly
sampled 100 quartets around each internal node. Finally, we mapped gCF and sCF values to the
ASTRAL-II supercontig tree produced in our previous summary coalescent analysis (Fig. 2). All

phylogenetic trees, with the exception of the full Phyparts picharts tree, were visualized using the
R package *ggtree* version 1.14.6 (G Yu, DK Smith, H Zhu, Y Guan 2017).
968
Ancestral State Reconstruction
To infer ancestral conditions and the number of transitions in reproductive system, we
used the *phangorn* (Schliep 2011) package in R. First, a Coalescent-based species tree with
accessions grouped into taxa using a mapping file was estimated in ASTRAL-III (Zhang et al.
2018) with default settings using the best RAxML gene trees and their associated bootstrap files,
from the supercontig alignments, as input. Next we time calibrated the ASTRAL-III species tree
to 1 million years based on estimates of other taxa in the genus (Evans et al. 2009) using the
makeChronosCalib function in the *ape* (Paradis et al. 2004) package in R, and estimated an
ultrametric tree using the chronos function in *ape* (Paradis et al. 2004) with settings: lambda = 1,
model = "relaxed". Finally, we performed marginal reconstruction of ancestral character states
using the maximum likelihood method using the optim.pml and ancestral.pml functions in the
phangorn (Schliep 2011) package in R.
981
Testing for Hybrid Orgins with HyDe
To test for putative hybrid origins of selected taxa, we used HyDe (Blischak et al. 2018)
to calculate D-Statistics (Green et al. 2010) for a set of hypotheses (S10). Briefly, HyDe
considers a four-taxon network of an outgroup and a triplet of ingroup populations to detect
hybridization from phylogenetic invariants that arise under the coalescent model with
hybridization. Introgression between P3 and either P1 or P2 influences the relative frequencies of
ABBA and BABA, and the D-statistic measures the imbalance between these frequencies. We
tested the triplets in (S10) and set *Chylismia scapoidea* as the outgroup. We considered
hypothesis tests significant at an overall $\alpha < 0.05$ level with estimates of γ between 0 and 1. Z-
scores greater than 3 are generally interpreted as strong evidence of introgression.
992
Population-level Analysis
To further characterize population-level processes or genetic structure within sect.
Calylophus, we extracted and filtered SNPs by mapping individual reads against reference
supercontigs (see https://github.com/lindsawi/HybSeq-SNP-Extraction). To account for
duplicates arising from PCR during HybSeq in SNP calling and filtering, first we selected the
sample with the highest target recovery rate and sequencing depth as a target reference sequence
(Oenothera_capillifolia_berlandieri_bjc19) and gathered supercontigs for this individual into a
single target FASTA file. We then ran BWA (Li and Durbin 2009) to align sequences, Samtools
'index' (Danecek et al. 2011) and GATK CreateSequenceDictionary (Poplin et al. 2017),
respectively, on the resulting target FASTA file. Next we ran a custom script
"variant_workflow.sh" using both read files from each *Calylophus* sample as input to create a
vcf file for each sample. SNP's were called for each individual using GATK (Poplin et al. 2017)
and the vcf file from each sample as input. The resulting vcf file created in the previous step was
filtered to remove indels using GATK and the original target FASTA file as input, and then
filtered again based on read mapping and quality with GATK VariantFiltration with settings: --
filterExpression "QD < 5.0 || FS > 60.0 || MQ < 40.0 || MQRankSum < -12.5 || ReadPosRankSum
< -8.0" (Poplin et al. 2017). Finally, we generated a reduced SNP file in FastStructure format
using PLINK (Purcell et al. 2007) to remove SNPs that did pass filter using the command: plink -
-vcf-filter --vcf Pachylophus.filtered.snps.vcf --const-fid --allow-extra-chr --geno --make-bed --

recode structure. Finally we used Discriminant Analysis of Principal Components (Jombart et al.
2010) as implemented in the R package *adegenet* (Jombart 2008) and the snmf function in the
LEA package (Frichot and François 2015) in R (R Core Team, 2020).
1015
Morphological Measurements and Analysis
To assess taxon boundaries and patterns of morphological variation, we measured
character states for the following key morphological structures that have been used historically to
discriminate taxa in sect. *Calylophus* (Towner 1977): plant height, leaf length (distal), leaf width
(distal), leaf length/width ratio (distal), leaf length (basal), leaf width (basal), leaf length/width
(basal), sepal length, and sepal tip length. Measurements were made with digital calipers when
possible, or with a standard metric ruler and dissecting scope, from voucher specimens of nearly
all sampled populations of sect. *Calylophus* included in our molecular phylogenetic analyses.
Measurements are provided in S11 and have been deposited at Dryad
(http://dx.doi.org/10.5061/dryad.[NNNN]). Morphological measurements were log transformed
using the R base function 'log' (R Core Team 2018) prior to Principal Components Analysis
(PCA), which was conducted in R using the *stats* package version 3.7 and the function
'prcomp'(R Core Team 2018). All 'NA' values were omitted from analysis. Plots of PCA results
were visualized using the *ggplot2* package in R (Wickham 2016).
1030

REFERENCES
Abascal F., Zardoya R., Telford M.J. 2010. TranslatorX: Multiple alignment of nucleotide
sequences guided by amino acid translations. Nucleic Acids Res. 38:7–13.
Alexander M.P. 1969. Differential Staining of Aborted and Nonaborted Pollen. Stain Technol.
44:117–122.
Alexander M.P. 1980. A Versatile Stain for Pollen Fungi, Yeast and Bacteria. Stain Technol.
55:13–18.
ArborBiosciences. 2016. In-Solution Sequence Capture for Targeted High-Throughput
Sequencing: Version 3.02. .
Blischak P.D., Chifman J., Wolfe A.D., Kubatko L.S. 2018. HyDe: A python package for
genome-scale hybridization detection. Syst. Biol. 67:821–829.
Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina
sequence data. Bioinformatics. 30:2114–2120.
Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model.
Bioinformatics. 30:3317–3324.
Chifman J., Kubatko L. 2015. Identifiability of the unrooted species tree topology under the
coalescent model with time-reversible substitution processes, site-specific rate variation,
and invariable sites. J. Theor. Biol. 374:35–47.
Danecek P., Auton A., Abecasis G., Albers C.A., Banks E., DePristo M.A., Handsaker R.E.,
Lunter G., Marth G.T., Sherry S.T., McVean G., Durbin R. 2011. The variant call format
and VCFtools. Bioinformatics. 27:2156–2158.
Doyle J.J. 1987. A Rapid DNA Isolation Procedure for Small Quantities of Fresh Leaf Tissue.
Phytochem Bull.:11–15.
Duarte J.M., Wall P.K., Edger P.P., Landherr L.L., Ma H., Pires J.C., Leebens-Mack J.,
dePamphilis C.W. 2010. Identification of shared single copy nuclear genes in Arabidopsis,
Populus, Vitis and Oryza and their phylogenetic utility across various taxonomic levels.
BMC Evol Biol. 10:61.

Evans M.E.K., Smith S. a, Flynn R.S., Donoghue M.J. 2009. Climate, niche evolution, and
diversification of the "bird-cage" evening primroses (Oenothera, sections Anogra and
Kleinia). Am. Nat. 173:225–40.
G Yu, DK Smith, H Zhu, Y Guan T.L. 2017. ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. Methods
Ecol. Evol.:28–36.
Green R., Krause J., Briggs A., Rasilla Vives M., Fortea Pérez F. 2010. A draft sequence of the
neandertal genome. Science (80-.). 328:710–722.
Gutíerrez S.C., Martínez J.M.S., Gabaldón T. 2009. TrimAl : a tool for automatic alignment
trimming. Bioinformatics.:1–2.
Illumina. 2016. BaseSpace Sequence Hub. Available from https://basespace.illumina.com/.
Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett
N.J. 2016. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics from High-
Throughput Sequencing Reads Using Target Enrichment. Appl. Plant Sci. 4:1600016.
Katoh K., Misawa K., Kuma K., Miyata T. 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30:3059–3066.
Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 25:1754–1760.
Miller M.A., Pfeiffer W., Schwartz T. 2010. Creating the CIPRES Science Gateway for
inference of large phylogenetic trees. 2010 Gatew. Comput. Environ. Work. GCE 2010.
Minh B.Q., Hahn M.W., Lanfear R. 2018. New methods to calculate concordance factors for
phylogenomic datasets. bioRxiv.:487801.
Mirarab S., Warnow T. 2015. ASTRAL-II: Coalescent-based species tree estimation with many
hundreds of taxa and thousands of genes. Bioinformatics. 31:i44–i52.
Paradis E., Claude J., Strimmer K. 2004. APE: Analyses of phylogenetics and evolution in R
language. Bioinformatics. 20:289–290.
Poplin R., Ruano-Rubio V., DePristo M.A., Fennell T.J., Carneiro M.O., Auwera G.A. Van der,
Kling D.E., Gauthier L.D., Levy-Moonshine A., Roazen D., Shakir K., Thibault J.,
Chandran S., Whelan C., Lek M., Gabriel S., Daly M.J., Neale B., MacArthur D.G., Banks
E. 2017. Scaling accurate genetic variant discovery to tens of thousands of samples.
bioRxiv.:201178.
Purcell S., Neale B., Todd-Brown K., Thomas L., Ferreira M.A.R., Bender D., Maller J., Sklar
P., De Bakker P.I.W., Daly M.J., Sham P.C. 2007. PLINK: A tool set for whole-genome
association and population-based linkage analyses. Am. J. Hum. Genet. 81:559–575.
R Core Team. 2018. R: A language and environment for statistical computing. .
Sayyari E., Mirarab S. 2016. Fast coalescent-based computation of local branch support from
quartet frequencies. Mol. Biol. Evol.:msw079.
Schliep K.P. 2011. phangorn: Phylogenetic analysis in R. Bioinformatics. 27:592–593.
Sharma P., Purohit S.D. 2012. An improved method of DNA isolation from polysaccharide rich
leaves of Boswellia serrata Roxb. Indian J. Biotechnol. 11:67–71.
Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals
conflict, concordance, and gene duplications with examples from animals and plants. BMC
Evol. Biol. 15:150.
Stamatakis A. 2014. Stamatakis - 2014 - RAxML version 8 a tool for phylogenetic analysis and
post-analysis of large phylogenies. :2010–2011.
Sukumaran J. and M.T.H. 2010. DendroPy: A Python library for phylogenetic computing.

Bioinformatics.:1569–1571.

Suyama M., Torrents D., Bork P. 2006. PAL2NAL: Robust conversion of protein sequence

alignments into the corresponding codon alignments. Nucleic Acids Res. 34:609–612.

Towner H.F.. 1977. The Biosystematics of Calylophus (Onagraceae). Ann. Missouri Bot. Gard.

64:48–120.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

1112

1113   S4.

1114
1115   Concatenation tree constructed using the exon-only dataset and 100 bootstraps. Bootstrap values
1116   indicated at relevant nodes.

1117   S5.

1118
1119   Concatenation tree constructed using the supercontig dataset and 100 bootstraps. Bootstrap
1120   values indicated at relevant nodes.

1121   S6.

1122
1123   ASTRAL-II summary coalescent tree constructed using the exon-only dataset and 100
1124   bootstraps. Bootstrap values indicated at relevant nodes.

1125   S7.

1126
1127   SVD Quartets summary coalescent tree constructed using the supercontig dataset with 100
1128   bootstraps. Bootstrap values indicated at relevant nodes.

1129

1130   S8.

1131
Phyparts piecharts ASTRAL-II tree constructed using the supercontig dataset. Piechart colors
correspond to: blue = concordant, green = top alternative bipartition, red = all other alternative
bipartitions, black = uninformative for that node.

S9.

Summary of support values for current and proposed taxonomic treatments, by analysis. 'e'

signifies trees based on exon-only data, "e+i" signifies trees based on supercontigs. '*p'* indicates

paraphyletic, unsupported taxon treatment according to tree topology.

Taxon	Concat. e	Concat. e+i	ASTRAL-II e	ASTRAL-II e+i	SVDQ e+i	Phyparts	gCF iQtree e+i	sCF iQtree e+i
a. *Oenothera hartwegii* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
b. *Oenothera hartwegii* subsp. *fendleri* (Towner 1977)	<50	*p*	81	99	*p*	1/ 188	1	38
c. *Oenothera hartwegii* subsp. *filifolia* (Towner 1977)	*p*	*p*	<50	<50	<50	1/ 190	0	36
d. *Oenothera hartwegii* subsp. *hartwegii* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
e. *Oenothera hartwegii* subsp. *macartii* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	
f. *Oenothera hartwegii* subsp. *hartwegii* + subsp. *macartii*	*p*	*p*	<50	<50	<50	0/ 200	0	33
g. *Oenothera hartwegii* subsp. *pubescens* (Towner 1977)	*p*	*p*	<50	<50	<50	0/ 200	0	36
h. *Oenothera lavandulifolia* (Towner 1977)	100	*p*	100	100	100	147/37	74	83
i. *Oenothera tubicula* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*1/ 188*	*p*	*p*
j. *Oenothera tubicula* subsp. *tubicula* (Towner 1977)	*p*	*p*	81	99	<50	2/ 171	1	38
k. *Oenothera tubicula* subsp. *strigulosa* (Towner 1977)	*p*	*p*	81	100	<50	*p*	0	37
l. *Oenothera fendleri* + *Oenothera tubicula* subsp. *tubicula*	*p*	*p*	100	100	<50	16/129	10	40
m. *Oenothera toumeyi* (Towner 1977)	100	100	100	100	100	125/65	62	56
n. *Oenothera toumeyi* + subsect. *Calylophus*	100	100	100	100	100	149/42	76	65
o. *Oenothera capillifolia* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
p. *Oenothera capillifolia* subsp. *capillifolia* (Towner 1977)	*p*	*p*	<50	*p*	<77	*p*	*p*	*p*
q. *Oenothera capillifolia* subsp. *berlandieri* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
r. *Oenothera capillifolia* subsp. *berlandieri* (New Taxon; South Texas coastal populations)	*p*	*p*	<50	100	100	*157/27*	79	86
s. *Oenothera serrulata* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
t. *Oenothera 'australis'* (New Taxon; South Texas *O. serrulata*)	*p*	100	*p*	*p*	*p*	10/175	*p*	*p*
u. *Oenothera gayleana* (Turner & Moore 2014)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
v. *Oenothera gayleana* (Revised Taxon)	*p*	*p*	*p*	<50	<50	0/ 193	0	35

1139

1140 S10.

1141 List of admixture hypotheses tested, Zscore, P-value and Gamma results from HyDe analysis.

1142 Each row represents a triplet set that was tested consisting of a putative hybrid individual and

1143 two parent groups; Parent 1 and Parent 2 (see "HyDe Group" Appendix I for group

1144 membership).

1145

Putative hybrid individuals	Parent 1 Group	Parent 2 Group	Zscore	P	Gamma
Ohar_X_fil_MJM629.6	CoreOharfil	CoreOharpub	-99999.9	1	-0.049
Ohar_X_fil_MJM628	CoreOharfil	CoreOharpub	-99999.9	1	-0.057
Ohar_X_fil_MJM626	CoreOharfil	CoreOharpub	-99999.9	1	-0.001
Ohar_X_fil_MJM793	CoreOharfil	CoreOharpub	-99999.9	1	-0.057
Ohar_X_fil_MJM794	CoreOharfil	CoreOharpub	-99999.9	1	-0.034
Ohar_X_fil_BJC81	CoreOharfil	CoreOharpub	-99999.9	1	-0.042
Oharhar_MJM1581	CoreOharpub	Oharhar/harmac	-99999.9	1	1.706
Ohar_X_pub_BJC29	CoreOharpub	Oharhar/harmac	2.378	0.009**	0.338
Oharpub_BJC30	CoreOharpub	Oharhar/harmac	-99999.9	1	-0.215
Oharpub_MJM2158.C	CoreOharpub	Oharhar/harmac	1.113	0.133	0.229
Ohar_X_pub_MJM1134.2	CoreOharpub	Oharhar/harmac	0.043	0.483	0.015
Ohar_X_pub_MJM2405.A	CoreOharpub	Oharhar/harmac	0.882	0.189	0.158
Ohar_X_pub_MJM594.4	CoreOharpub	Oharhar/harmac	2.094	0.018*	0.332
Ohar_X_pub_MJM615	CoreOharpub	Oharhar/harmac	0.413	0.340	0.096
Oharmac_MJM997.1	CoreOharpub	Oharhar/harmac	1.563	0.059	0.224
Ohar_X_har_MJM613	CoreOharpub	Oharhar/harmac	-99999.9	1	-0.495
Otub_X_str_MJM1916.E	Otubtub	Oharhar/harmac	5.585	0.000***	0.947
Oharfen_MJM2192.A	CoreOharfen	CoreOharfil	0.305	0.380	0.001
Oharfen_MJM781.3	CoreOharfen	CoreOharpub	0.025	0.490	9.69E-05
Ocapcap_MJM1015.2	CoreOcapcap	S. TX Ocapber	0.467	0.320	0.003
Ocap_X_ber_MJM767.3	central OK Ocapber	Pecos River/Southern Plains	-99999.9	1	-0.003

- Indicates value meets significance threshold of P < 0.05
** Indicates value meets significance threshold of P < 0.001
*** Indicates value meets significance threshold of P < 0.0001

1146

1147 S11.

1148 See excel table "S11 Morphometric Data".

1149 S12.

1150 See excel table "S12 Pollen Counts".

1151

1152 S13.

<div align="center">

Oenothera HybSeq CTAB Silica DNA extractions
b. cooper 11/10/2015

</div>

Adapted from JJSA Protocol, J. Fant's Protocol and Sharma and Puohit (2012) for Silica-dried Leaves of *Oenothera* sect. *Calylophus*

SAMPLE PREPARATION & GRINDING

1. Warm <u>CTAB buffer</u> to 65°C in water bath in the hood and pour liquid nitrogen into dipping canister (You will only need a few inches of liquid in the Dewar, make sure to pour it back into the large holding canister when you are done to prevent evaporation).
2. Put a small volume of 1:1 sterilized sand and 3 metal/ceramic beads into centrifuge tubes, then put tubes in freezer (-20) until envelopes are ready.
3. Keeping envelopes/leaf tissue on ice to prevent thawing; take envelopes and centrifuge tubes out of freezer and weigh/estimate small amount of dried tissue (e.g., 10-50 mg of leaves), put in labeled microfuge tubes. After each tube is filled, place tube in freezer.
   a. Make sure leaf tissue is broken up into small pieces when adding to ensure full grinding.
   b. Keeping tissue frozen prevents phenolic compounds from oxidizing, and enzymes from activating and breaking down DNA
   c. Follow DNA free lab protocols using bleach to prevent contamination/cross contamination of samples.
4. Mix 45 (= .3%) µl 2-mercaptoethanol to 15 ml <u>CTAB buffer</u> in small beaker (enough for ~15 samples...double for 30 samples, etc.)
   a. Mercaptoethanol must be measured and added in the fume hood.
   b. CTAB is a detergent that lyses (breaks down) the cell membranes. The PVP in the buffer helps bind the polysaccharides and might co-ppt
5. Dip tubes in Liquid Nitrogen (or store in -80 for ~30 minutes) and **immediately** Grind tissue **thoroughly** in Fast Prep machine (thorough grinding may require 2-4 cycles, re-dip tubes in liquid nitrogen after 2 cycles to ensure no thawing occurs).
6. **Immediately** after grinding add 600 µl of CTAB/mercaptoethanol buffer to each tube
   a. **BE CAREFUL.** It is important to prevent cross-contamination when opening tubes. **Powdered tissue can spread easily through the air or on your fingers.** Make sure open tubes are spaced apart from one another on the tray; open each tube away from other open tubes/caps; and place caps a safe distance from one another. Make sure leaf powder is not stuck to your gloves before touching consecutive tubes/caps.
   b. Grind on FastPrep for one cycle

EXTRACTION AND ISOLATION OF DNA (once leaf tissue is well-ground)

1. Incubate tubes for 60 minutes at 60°C on Thermoblock or water bath
   a. Shake tubes several times ~ every 10 minutes during incubation (for compacted samples, vortex on high setting to ensure mixing)
   b. When incubation is complete, turn Thermoblock down to 37°C
2. Add 600 µL chloroform-isoamyl (24:1) to tube and vortex on High setting to ensure thorough mixing.
   a. **BE CAREFUL.** It is important to prevent cross-contamination at this step when opening/handling tubes. Follow steps outlined previously.
   b. Chlorophyll and other pigments get transferred into the CI layer because pigments are non-polar and dissolve into the highly non-polar chloroform. The DNA remains in the aqueous layer.
3. Spin for 10 min at 9,000 RPM.
   a. After spinning, the upper layer of each tube should have a clear liquid: this is where your DNA is.
4. Transfer top (clear) liquid to a new (labeled) 1.5 ml eppindorf tube. **Do not be Greedy**, leaving some liquid in tube is fine – the important thing is not to pipette **ANY** interphase material.
   a. A yield of ~400-500 µL of supernatant is common.
   b. If you accidently disturb the interphase, or believe that you have pipetted interphase, return supernatant to microfuge tube and re-centrifuge for 10 min at 9000 rpm. Repeat step 4.

DNA PRECIPITATION

1. Add 1/10 volume 3M sodium acetate and ½ volume 5M NaCl (i.e. 500ul of solution add 250 ul of NaCl, 50 ul of Na-Acetate).
2. Add 2/3rd volume of cold isopropanol (to previous volumes = 500 ul isop)
   a. Isoproponal is heavy alcohol so allows DNA ppt with lower volume

3.   Mix & store in -20°C (freezer) for 1-2 hours
(NOTE: POTENTIAL STOPPING POINT FOR A DAY OR MORE…SAMPLES CAN STAY AT -20°C FOR DAYS)
4.   Spin for 3 min at **10K-12K RPM.**
5.   Decant the supernatant & drain; make sure pellet stays at bottom of tube.
a.   Pellets are usually white but sometimes they can even be brown, this is not necessarily a problem.
6.   Wash pellet with 500ul of <u>70</u>% EtOH & flick tubes to dislodge pellet from bottom, then vortex on high to
clean.
7.   Spin for 3 min at 10K-12K RPM.
8.   Decant the supernatant making sure pellet stays at bottom of tube
9.   Leave tubes open and place tubes on Speedvac (vacuum on -2.5, heat = low, spin) for ~20 minutes or until
all EtOH has evaporated.
a.   If pellets are not dry after 20 minutes, check frequently to prevent over drying which makes the pellet
1219   difficult to re-suspend.
b.   Alternatively, can air dry on thermoblock
(NOTE: POTENTIAL STOPPING POINT …SAMPLES CAN STAY AT 20°C FOR DAYS – place in cupboard but cover with kimwipe)
FINAL CLEANING & RESUSPENSION OF DNA
1.   Add 500 ul of 0.5 mL High Salt TE Buffer and close lids
2.   Incubate tubes on Thermoblock at 37-50°C (45) until pellet dissolves (15-30 minutes);
a.   Vortex frequently if this is taking a long time
(NOTE: POTENTIAL STOPPING POINT FOR A DAY OR MORE…SAMPLES CAN STAY AT -20°C FOR DAYS)
3.   Add 3 volumes of binding buffer (3 volumes for every 1 volume DNA pellet = ~100-150 ul BB), **let stand for**
20 minutes
4.   Centrifuge at 550 g for 10 min
5.   transfer supernatant to clean 2ml eppi tube leaving any colored or gelatinous precipitate behind.
6.   Add 300 µl of silica suspension and mix for 30 minutes by regular and frequent gentle inversion
7.   Centrifuge at 550 g for 10 minutes, discard supernatent
8.   Re-suspend silica pellet in 1.5 mL of wash buffer 1
9.   Centrifuge at 3000 g for 15 seconds, decant supernatent
10.   Re-suspend silica pellet in 1.5 mL of wash buffer 2
11.   Centrifuge at 3000 g for 15 seconds, decant supernatant, dry pellet completely on speed vac
12.   suspend silica pellet in 300 ul TE buffer, incubate at 50° mixing regularly by vortex
13.   centrifuge at 11600 g for 1 min. then transfer supernatant into clean 1.5 mL eppi tube.
14.   Precipitate DNA by adding 50 ul of 3M sodium Acetate and 500 uL 100% Etoh
15.   Store at -20° C for at least 2 hours
(NOTE: POTENTIAL STOPPING POINT FOR A DAY OR MORE…SAMPLES CAN STAY AT -20°C FOR DAYS)
16.   Centrifuge at 11600 g for 5 min, discard supernatant, dry pellet on speed vac
17.   Dissolve in 50 uL 1X TE buffer (it may not look like there is anything in the tube at this point, but there is
lots of pure DNA in there! Add the TE buffer)
18.   Store in -20°C (freezer)
19.   In rare circumstances that an extraction still has coloration at this stage (yellow or brown usually) you can
now put it through a Qiagen Qiaquick spin column or other proprietary cleanup column to further purify
the DNA.
1249
REAGENTS (recipes below) and SUPPLIES
•   CTAB buffer (in glass container in refrigerator)
•   Liquid Nitrogen
•   2-mercaptoethanol (in fume hood with gloves)
•   24:1Chloroform:isoamyl alcohol - glass container in fumehood use gloves and keep in hood -(Or 25:24:1
phenol-chloroform:isoamyl)
•   5M Nacl
•   3M M sodium acetate
•   Isopropanol and 70% and 100% EtOH
•   High Salt TE buffer
•   Binding Buffer

- Silica Matrix
- Wash Buffer 1
- Wash Buffer 2
- 1X TE buffer
- 1.5 ml eppi tubes
- 2 ml eppi tubes
1267

1268 **CTAB BUFFER** [100 mM Tris-HCl (pH=8.0), 1.4 M NaCl, 20 mM EDTA, 2% (or 4%) CTAB
1269 (hexadecyltrimethylammonium bromide), 2% PVP-40 (polyvinylpyrollidone, m.w. 40000), + 0.3% β-
1270 mercaptoethanol (add in fume hood in a beaker to a premeasured volume of CTAB buffer required for the number
1271 of extractions planned plus one).]
1272
For 200 mL CTAB buffer 2% (4%):
20 ml 1M Tris-Cl
1.4 M NaCl (16.36 g NaCl)
- The addition of NaCl at concentrations higher than 0.5 M, along with CTAB, is known to remove
polysaccharides during DNA extraction
8 ml 0.5 M EDTA
4 g CTAB (8 g for 4%)
4 g PVP-40
~120 ml dH2O, then fill to 200 mL
1282
1283 **TE buffer**: [10 mM Tris-HCl (pH=8.0), 1 mM EDTA.]

1284 10 mL 1 M Tris-HCl
1285 2 ml 0.5 M EDTA
1286 982 ml dH2O
1287
1288 **High Salt TE Buffer**: [10 mM Tris-HCl (pH=8.0), 1 mM EDTA, 1M NaCl.]
1289 10 ml 1 M Tris-HCl
1290 2 ml 0.5 M EDTA
1291 200 ml 5M NaCl
1292 788 ml dH2O
1293
Binding Buffer: [50mM Tris (pH7.5), 6M NaClO4, 1mM EDTA]
For 200 ml:
10mL Tris
168.552 g NaClO4
400 ul EDTA
top off to 200 mL with dH20
1300
1301 **Wash Buffer 1**: [3 volumes binding buffer, 1 volume water]
1302 For 60 samples:
1303 90 ml Binding Buffer
1304 30 ml dH20
1305
Wash Buffer 2: [1 volume 40 mM Tris (pH 8.0), 4 mM EDTA, 0.8 M NaCl, 1 Volume ethanol]
For 60 samples:
1.8 ml 1M tris
360 ul .5M EDTA
7.2 ml 5M NaCL

35.64 ml dh20
45 ml 100% etoh
1313
1314   **24:1 Chloroform: Isoamyl**
1315   240ml Chloroform
1316   10ml Isoamyl alcohol
1317
1318   25:24:1 Chloroform: Isoamyl Saturated with 10 mM Tris, pH 8.0, 1 mM EDTA. – Purchase to ensure right pH
1319   250ml Phenol
1320   240ml Chloroform
1321   10ml Isoamyl alcohol
1322
Silica suspension (use lab grade silicon dioxide 99%)
1. Suspend silicon dioxide powder in ~20 volumes of water (50 ml : 1L)
a. use magnetic stirring rod to bring silica into uniform suspension
2. Decant suspension into clean beaker leaving behind heavy sediment at bottom
(discard this)
3. Allow suspension to settle by gravity for 2-3 hours
4. Decant and discard supernatant with fine material keeping silicon dioxide that
has settled
5. Re-suspend silicon dioxide in dH20, repeating step until fine particles removed
(2-3X total is good)
a. use magnetic stirring rod to bring silica into uniform suspension
6. Re-suspend silicon dioxide in dH20 and transfer to 15 ml falcon tubes
7. Collect silicon dioxide by low speed centrifugation (up to 3200 rpm for 3 min)
8. Add ~15 ml of 10% bleach solution to each aliquot and mix frequently by
inversion/vortexing for 15-30 minutes
9. Re-suspend diatomite in dH20 by vortexing, then centrifuge at low speed for 3
minutes to pellet
10. Discard supernatent
11. Repeat step 9 and 10 two more times
12. Re-suspend in dH20 and autoclave
13. Re-suspend silica dioxide pellet in 1 volume PCR grade DNA/RNA free H2O
14. Aliquot into 1.5 ml eppi tubes and store at 4° C
15. Vortex thoroughly to bring silica into uniform suspension before using

1361   S14.

1362
1363   Ancestral State Reconstruction of reproductive system in sect. *Calylophus* using supercontigs
1364   and accessions grouped into taxa. Pie-charts on nodes represent likelihood of ancestral
1365   reproductive system at each node (teal = hawkmoth pollination, purple = bee pollination, yellow
1366   = PTH).
1367