1 TITLE: Target enrichment and extensive population sampling help untangle the recent, rapid
2 radiation of *Oenothera* sect. *Calylophus*
3
4 RUNNING HEAD: Cooper et al., Phylogenomics of *Oenothera* sect. *Calylophus*
5

6 AUTHORS
7 Benjamin J. Cooper[1,2,*], Michael J. Moore[3], Norman A. Douglas[4], Warren L. Wagner[5], Matthew
8 G. Johnson[1,6], Rick P. Overson[1,7], Angela J. McDonnell[1], Jeremie B. Fant[1,2], Krissa A. Skogen[1,2],
9 Norman J. Wickett[1,2,*]
10
[1]The Negaunee Institute for Plant Conservation Science and Action, Chicago Botanic Garden,
1000 Lake Cook Rd., Glencoe, IL 60022
[2]Northwestern University, Program in Plant Biology and Conservation, O.T. Hogan Hall, Room
6-140B, 2205 Tech Drive, Evanston, IL 60208
[3]Oberlin College, Department of Biology, 119 Woodland St., Oberlin, OH 44074
[4] Department of Biology, University of Florida, Gainesville, FL 32611
[5] Department of Botany, MRC-166, Smithsonian Institution, PO Box 37012, Washington, DC
20013-7012
[6] Department of Biological Sciences, Texas Tech University, Box 43131 Lubbock, TX 79409
[7] School of Sustainability, Arizona State University, PO Box 875502, Tempe, AZ 85287-5502
* Authors for correspondence: nwickett@chicagobotanic.org, benjamin.cooper06@gmail.com
22

23 ABSTRACT
24

Oenothera sect. *Calylophus* is a North American group of 13 recognized taxa in the
evening primrose family (Onagraceae) with an evolutionary history that may include
independent origins of bee pollination, edaphic endemism, and permanent translocation
heterozygosity. Like other groups that radiated relatively recently and rapidly, taxon boundaries
within *Oenothera* sect. *Calylophus* have remained challenging to circumscribe. In this study, we
used target enrichment, flanking non-coding regions, summary coalescent methods, tests for
gene flow modified for target-enrichment data, and morphometric analysis to reconstruct
phylogenetic hypotheses, evaluate current taxon circumscriptions, and examine character
evolution in *Oenothera* sect. *Calylophus*. Because sect. *Calylophus* comprises a clade with a
relatively restricted geographic range, we were able to comprehensively sample across the range
of geographic and morphological diversity in the group with extensive sampling. We found that
the combination of exons and flanking non-coding regions led to improved support for species
relationships. We reconstructed potential hybrid origins of some accessions and note that if
processes such as hybridization are not taken into account, the number of inferred evolutionary
transitions may be artificially inflated. We recovered strong evidence for multiple origins of the
evolution of bee pollination from the ancestral hawkmoth pollination, the evolution of edaphic
specialization on gypsum, and permanent translocation heterozygosity. This study applies newly
emerging techniques alongside dense infraspecific sampling and morphological analyses to
effectively address a relatively common but recalcitrant problem in evolutionary biology.
44
45 *Keywords.*– Gypsum Endemism, Onagraceae, *Oenothera* sect. *Calylophus*, Pollinator Shift,
46 Recent Radiation, Phylogenomics, Target Enrichment

INTRODUCTION

The challenges of reconstructing species histories for groups that arose through recent, rapid radiations are well established. Phylogenetic signal can be obscured by processes such as incomplete lineage sorting (ILS) and gene flow (Maddison and Knowles 2006; Knowles and Chan 2008; Christie and Knowles 2015), resulting in short branch lengths and conflicting gene tree topologies. Consequently, approaches that use few loci or concatenation may fail to resolve the most accurate species tree (Eckert and Carstens 2008; Leaché et al. 2014; Xi et al. 2014; Giarla and Esselstyn 2015). This may be particularly common in plants that are thought to have experienced rapid or recent radiation with ongoing hybridization and high levels of ILS , as is likely the case in the evening primrose genus *Oenothera* sect. *Calylophus* (Onagraceae). The application of target enrichment methods that efficiently sequence hundreds of nuclear loci, coalescent-based phylogenetic methods that account for ILS and gene flow, and extensive sampling of morphologically diverse populations across the geographic range may allow for more accurate representations of phylogenetic relationships in this group (Maddison and Knowles 2006; Knowles and Chan 2008; Knowles 2009; Mamanova et al. 2010; Lemmon et al. 2012; Straub et al. 2012; Bryson et al. 2014; Weitemier et al. 2014; Mandel et al. 2014; Stephens et al. 2015; Johnson et al. 2016).

Oenothera sect. *Calylophus* is currently considered to comprise seven species (thirteen taxa) with a center of diversity in western Texas, southern New Mexico, and north-central Mexico (Fig. 1; Towner 1977; Turner and Moore 2014; Wagner, in press). Previous analyses suggest that *Oenothera* sect. *Calylophus* forms a well-supported, morphologically coherent clade with a relatively restricted geographic range (Towner 1977; Levin et al. 2004; Wagner et al. 2007; Turner and Moore 2014; Wagner, in press). However, as with other groups that have experienced rapid radiations, taxon boundaries within *Oenothera* sect. *Calylophus* have been challenging to define, likely due to phenomena such as overlapping morphological boundaries, ongoing introgression, and incomplete lineage sorting.

In the most comprehensive study of the group to date, Towner (1977) circumscribed taxa using morphology, breeding system, geography, and ecology, but it was noted (and our field observations confirm) that characters often overlap among taxa (Towner 1977). Taxa within *Oenothera* sect. *Calylophus* are divided into two easily recognizable subsections: subsect. *Salpingia* and subsect. *Calylophus* (Towner 1977; Wagner et al. 2007). Pollination varies between the two subsections; flowers of subsect. *Salpingia* are adapted to vespertine pollination by hawkmoths, except for *O. tubicula*, which opens in the morning and is primarily pollinated by bees (Towner 1977). Taxa in subsect. *Calylophus* are predominantly bee-pollinated (Towner 1977) and have geographic ranges that partially (or even largely) overlap, resulting in occasional morphologically intermediate populations (Towner 1977). While confounding for morphological-based analyses, this observed pattern of reticulation is consistent with a recent, rapid radiation occurring in parallel with climatic fluctuations and increasing aridity in the region since the Pleistocene (Raven 1964; Towner 1977; Nason et al. 2002; Katinas et al. 2004). Hawkmoth pollination, which is ancestral in the family Onagraceae, and common in *Oenothera* sect. *Calylophus*, is known to result in long-distance pollen movement (Stockhouse 1973; Skogen et al. 2019); therefore, gene flow may have been extensive over the evolutionary history of hawkmoth-pollinated taxa, increasing the chances that processes such as historical introgression may obscure phylogenetic signal in extant plants (Elrich and Raven 1969). With a

phylogenomic approach that samples hundreds of nuclear loci, we may better illuminate both the
history of these species and the key evolutionary processes related to speciation in this group.
Understanding speciation in angiosperms remains a fundamental question in evolutionary
biology (Barrett et al. 1996; Rajakaruna 2004; van der Niet et al. 2006; Wilson et al. 2007;
Crepet and Niklas 2009; Peakall et al. 2010; Xu et al. 2011; Van der Niet and Johnson 2012;
Boberg et al. 2014). Section *Calylophus* has an evolutionary history that likely involves changes
in reproductive systems (pollinator functional group, breeding system, and autogamy) and
edaphic endemism. For example, there are thought to be two independent shifts between
pollinators in the section from hawkmoth to bee pollination (Towner 1977; Fig. 1b), despite
many studies in other plant groups showing a directional bias in shifts from bee to hummingbird
or hawkmoth pollination (Barrett et al. 1996; Wilson et al. 2007; Thomson and Wilson 2008;
Tripp and Manos 2008; Barrett 2013). However, pollinator shifts that do not follow this sequence
may be more likely when the extent of trait divergence and specialization does not completely
inhibit secondary pollinators such as bees, as has been suggested in *Oenothera* sect. *Calylophus*
(Stebbins 1970; Tripp and Manos 2008; Van Der Niet et al. 2014). Shifts to autogamy are also
frequent across angiosperms and in Onagraceae alone there are an estimated 353 shifts to modal
autogamy (Raven 1979). *Oenothera* sect. *Calylophus* also includes at least one autogamous
species, *O. serrulata*, which exhibits permanent translocation heterozygosity, a phenomenon in
which all chromosomes are translocationally heterozygous (PTH; Towner 1977) (Fig. 1b). While
the evolution of PTH has been assessed in molecular phylogenetic analyses across Onagraceae
(Johnson et al. 2009), no study to date has examined this transition in a well-sampled clade with
extensive population sampling. Lastly, abiotic ecological factors such as edaphic specialization
are also known to drive speciation in some groups (Rajakaruna 2004; van der Niet et al. 2006).
For example, serpentine endemics represent ~10% of the endemic flora in California even
though serpentine soils account for about 1% of terrestrial habitat in the state (Brady et al. 2005).
Similarly, the Chihuahuan Desert is comprised of numerous isolated islands of gypsum outcrops
and current estimates suggest that at least 235 taxa from 36 different plant families are gypsum
endemics (Moore and Jansen 2007; Moore et al. 2014). It is suspected that gypsum endemism
has also evolved independently in *Oenothera* sect. *Calylophus* at least twice (Towner 1977;
Turner and Moore 2014; Fig. 1b). Ultimately, to understand the role that these transitions have
played in shaping the diversity of *Oenothera* sect. *Calylophus*, a robust phylogeny is required.
Here, we use target enrichment, summary coalescent methods, and morphometric
analyses to reconstruct a phylogenetic hypothesis, examine previous taxonomic concepts, and
resolve the history of pollinator shifts, PTH, and gypsum endemism in *Oenothera* sect.
Calylophus. Target enrichment is a cost-effective method for sequencing hundreds of loci across
hundreds of samples, producing highly informative datasets for phylogenetics (Lemmon et al.
2012; Straub et al. 2012; Mandel et al. 2014; Weitemier et al. 2014; Heyduk et al. 2015;
Stephens et al. 2015; Johnson et al. 2016). While target enrichment is generally designed to
capture coding regions, a significant proportion of flanking non-coding regions can be recovered
(the "splash-zone"; Weitemier et al. 2014). The inclusion of non-coding regions may be
particularly informative for recent radiations, since these regions are less constrained by selective
pressures and may contain on average more informative sites at shallower time scales (Folk et al.
2015). We included these "splash-zone" regions in our sequence alignments to evaluate their
impact on reconstructing lower-order relationships. Importantly, we sampled extensively,
including individuals from numerous populations across the geographic and morphological
ranges of all thirteen taxa in the section (Fig. 1a). This study presents an example of how

combining these molecular techniques with dense sampling and morphological analysis can be
used to effectively address a common problem in evolutionary biology.
140
141 METHODS
142
A total of 194 individuals spanning the geographic, morphological, and ecological ranges
of all 13 recognized taxa in *Oenothera* sect. *Calylophus* [following Towner (1977) and Turner
and Moore (2014)] were included in this study (Fig. 1a, S1, S2) along with eight outgroups
representing other major sections of *Oenothera* (*Eremia, Gaura, Kneiffia, Lavauxia, Oenothera,*
Pachylophus, and *Ravenia*) and other genera (*Chylismia* and *Eulobus*) in Onagraceae (S1, S2).
DNA was extracted from fresh, silica-dried leaf tissue (S3). staiPTH status was determined for
individuals in subsect. *Calylophus* by assessing pollen fertility, when flowers were present, using
a modified Alexander stain (Alexander 1969, 1980; S3).
Target nuclear loci for enrichment were determined by clustering transcriptome
assemblies of *Oenothera serrulata* (1KP accession SJAN) and *Oenothera berlandieri* (1KP
accession EQYT). Starting with the 956 phylogenetically informative *Arabidopsis* loci identified
by Duarte et al. (2010; S3), we identified 322 homologous, single-copy loci in our clusters and
used these in the probe design process. Libraries were enriched for these loci using the MyBaits
protocol (Arbor Biosciences, Ann Arbor, MI, USA) and sequenced on an Illumina MiSeq (2 x
300 cycles, v3 chemistry; Illumina, Inc., San Diego, California, USA). Raw reads have been
deposited at the NCBI Sequence Read Archive (BioProject PRJNA544074; See S3 for details).
Reads were trimmed using Trimmomatic (Bolger et al. 2014; S3) and trimmed, quality-filtered
reads were assembled using HybPiper (Johnson et al. 2016). From the assembled loci, we
produced two datasets: **"exons"** – exon-only alignments, and **"supercontig"** – alignments
containing the exon alignment and flanking non-coding regions (the "splash-zone" per
Weitemier 2014 and reconstructed using supercontigs produced by HybPiper). We used these
two datasets to test the most recent taxonomic circumscription of the group with several
methods: (1) phylogenetic inference of concatenated alignments using RAxML (Stamatakis
2014; two analyses: exons and supercontigs), (2) ASTRAL-II (Mirarab and Warnow 2015;
Sayyari and Mirarab 2016) species tree inference (two analyses: exons and supercontigs), (3)
SVD Quartets (Chifman and Kubatko 2014, 2015) species tree inference (one analysis:
supercontigs), (4) Phyparts (Smith et al. 2015; one analysis: supercontigs), (5) IQtree (Minh et
al. 2018) with both gene and site concordance factors (one analysis: supercontigs).
We used HyDe (Blischak et al. 2018) to test for putative hybrid origins of selected taxa
and accessions by calculating D-Statistics (Green et al. 2010) for a set of hypotheses (S3). To
further characterize population-level processes or genetic structure within sect. *Calylophus*, we
extracted and filtered SNPs by mapping individual reads against reference supercontigs (see
https://github.com/lindsawi/HybSeq-SNP-Extraction) and used Discriminant Analysis of
Principal Components (Jombart et al. 2010) as implemented in the R package *adegenet* (Jombart
2008) and the snmf function in the LEA package (Frichot and François 2015) in R (R Core
Team, 2020; S3). We evaluated current taxonomic concepts and patterns of morphological
variation by measuring character states for morphological structures that have been used
historically to discriminate taxa in *Oenothera* sect. *Calylophus* (Towner 1977): Plant height, leaf
length (distal), leaf width (distal), leaf length/width ratio (distal), leaf length (basal), leaf width
(basal), leaf length/width (basal), sepal length, and sepal tip length (S3). Measurements were
made for 125 of the sequenced samples (S11); unfortunately, we were unable to measure all

traits for 73 samples because we did not have access to the herbarium vouchers, or the trait of
interest was not captured on the voucher, therefore some samples were dropped from the analysis
due to missing values. Finally, the number of transitions and inferred ancestral conditions of
reproductive system were mapped onto an ASTRAL species tree, with individuals grouped into
species, using the stochastic mapping function in the R package *phangorn* version 2.5.5 (Schliep
2011; S3).
190
191   RESULTS AND DISCUSSION
192
Sequencing and Phylogenetic Results
Sequencing resulted in a total of 80,273,296 pairs of 300-bp reads with an average of
625,323 reads per sample. Following quality filtering, assembly and alignment, we recovered
204 loci that were present in at least 70% of the samples. Across all datasets and analyses,
Oenothera sect. *Calylophus* was monophyletic. At the subsection level there was strong
agreement in topology between concatenation and coalescent-based trees. For example, subsect.
Calylophus was recovered as sister to subsect. *Salpingia* with strong support across all analyses
and *O. toumeyi* [considered by Towner (1977) to be in subsection *Salpingia*; Fig. 2, S4-7] was
recovered as sister to subsect. *Calylophus* across all trees, with strong bootstrap support. Within
subsect. *Calylophus* there was poor resolution for currently recognized taxa in all analyses,
whereas taxon relationships were better resolved in subsect. *Salpingia* (Fig. 2, S4-7). With
coalescent-based tree reconstruction, most taxa *sensu* Towner (1977) were recovered as
monophyletic with moderate to strong support (Fig. 2, S6-8). In contrast, both the exon and
supercontig concatenation trees recovered most currently recognized taxa as non-monophyletic
(S4, S5). Given that concatenation has been shown to produce incorrect topologies in the
presence of high ILS (Roch and Steel 2015) and that *Oenothera* sect. *Calylophus* underwent
recent radiation, we believe the paraphyly of taxa in both concatenation trees might be
artifactual. We, therefore, prefer to interpret relationships based on our coalescent-based trees,
which comprise the focus for the remainder of the paper (Fig. 2, S6-8).
To understand whether summary coalescent relationships display a consistent signal
across the genome, we quantified gene tree and site concordance using Phyparts and IQtree. We
found that gene tree concordance was highest at the deepest nodes at the species level where we
expected less ILS and more time between speciation events (Fig. 2, S8, S9). Correspondingly,
gene tree concordance was lowest at the subspecies level where increased sharing of ancestral
alleles and ongoing gene flow are more likely (Fig. 2, S8, S9). For example, within *O. hartwegii*
less than 1% of genes were concordant for bifurcations representing all currently recognized taxa
at the subspecies level (Fig. 2, S8, S9). For species-level nodes with high support and high gene
tree concordance, site concordance was also high; for example, *O. lavandulifolia* had high
bootstrap support in summary coalescent trees (BS = 100), high gene tree concordance (Phyparts
= 94% concordance, gCF = 74), and high site concordance (sCF = 83; Fig. 2, S9h). For
subspecies with high support, but low gene tree concordance, site concordance was moderate.
For example *O. hartwegii* subsp. *fendleri* had high bootstrap support (BS = 99), low gene tree
concordance (Phyparts = <1% concordance, gCF = 1), and moderate site concordance (sCF = 38;
Fig. 2, S9b). For subspecies that were monophyletic in our coalescent-based trees, but that had
low support and low gene tree concordance, site concordance was moderate with an average of
35% of sites in agreement for taxa at these nodes (S9[c,f,g,v]). For example *O. hartwegii* subsp.
filifolia had low bootstrap support, low gene tree concordance (Phyparts = <1% concordance,

gCF = 0), and moderate site concordance (sCF = 36; Fig. 2, S9c). This is an important finding
because while individual gene histories can be obscured by ILS, site concordance factors, which
may be less constrained and less subject to ILS at shallower evolutionary timescales, provide a
key alternative method of support.
In general, topologies of exon and supercontig datasets were similar, with no major
differences in clade membership, but the inclusion of the splash-zone increased support at
shallow nodes in our trees. However, this trend was not universal. For example, in subsect.
Salpingia, using the supercontig dataset decreased support slightly for one taxon (*O. hartwegii*
subsp. *filifolia*), and in subsect. *Calylophus* it led to paraphyly of another (*O. capillifolia* subsp.
capillifolia). For six other taxa, our results showed that using supercontigs increased bootstrap
support. Therefore, these results demonstrated a net benefit of including flanking non-coding
regions for resolving relationships among closely related taxa.
242
Hybridization and Geneflow
Using concatenated loci from the supercontig dataset, we used HyDe (Blischak et al.
2018) to test for signals of hybridization. We used 552,521 sites and tested 22 hypotheses for
either individuals or groups suspected to be of hybrid origin based on field observations of
morphological intermediacy, geographic location, and topological position in our coalescent-
based trees and found evidence of hybridization in three individuals representing two taxa, both
in subsect. *Salpingia*. The highest signal of hybridization, with a gamma value ($\hat{\gamma}$) of .947
suggesting more historic gene-flow, was observed in one individual of *O. tubicula* subsp.
strigulosa (MJM1916.E). This involved admixture between *O. tubicula* subsp. *tubicula* and the
clade consisting of *O. hartwegii* subsp. *hartwegii* and *O. hartwegii* subsp. *maccartii* (Z-score =
5.585, p-value = 0.000, $\hat{\gamma}$ = 0.947; Fig. 3a, S10). We also detected significant levels of
hybridization, with $\hat{\gamma}$ ranging from 0.332 to 0.338 suggesting more contemporary gene-flow, in
two individuals in *O. hartwegii* subsp. *pubescens*, BJC29 (Z-score = 2.378, p-value = 0.009, $\hat{\gamma}$ =
0.338) and MJM594 (Z-score = 2.094, p-value = 0.018, $\hat{\gamma}$ = 0.332). This more recent gene flow
involved admixture between *O. hartwegii* subsp. *pubescens* and the clade consisting of *O.*
hartwegii subsp. *hartwegii* and *O. hartwegii* subsp. *maccartii* (Fig. 3a, S10).
The finding that one individual of *O. tubicula* subsp. *strigulosa* may be of hybrid origin is
consistent with ongoing gene flow between *O. tubicula* subsp. *strigulosa* and its sister taxon *O.*
tubicula subsp. *tubicula*. In our coalescent-based analyses, the two subspecies of *O. tubicula*
were not recovered as sister taxa, and this relationship was strongly supported (S9[j-k]). If the
two *O. tubicula* taxa arose independently, this would support the hypothesis that bee pollination
arose in *Oenothera* sect. *Calylophus* independently three times. However, while the summary
coalescent analyses we utilized to estimate phylogenies accounted for ILS in tree estimation,
they did not account for ongoing gene flow (Meng and Kubatko 2009; Gerard et al. 2011;
Kubatko and Chifman 2019). Our HyDe results may support the hypothesis that *O. tubicula*
subsp. *strigulosa* has experienced gene flow from two closely related taxa, and may have hybrid
origins resulting from crossing between *O. tubicula* subsp. *tubicula* and *O. hartwegii* subsp.
hartwegii (Fig. 3a, S10). Therefore, the placement of *O. tubicula* subsp. *strigulosa* as sister to the
rest of the *O. hartwegii* species complex in our trees is likely the result of past gene flow and
does not represent independent origins of bee pollination in subsect. *Salpingia*. These results
underscore the importance of explicitly including tests for hybridization in phylogenetic studies. In
the case of these data estimating a species tree given a set of gene trees within a coalescent

framework without considering other, non-ILS sources of signal conflict could artificially inflate
the number of inferred evolutionary transitions.
277   Our HyDe results also suggest that at least some of the morphological intermediacy and
overlap among taxa in the group is due to continued, or at least recent, gene flow. For example,
both *O. hartwegii* subsp. *pubescens* individuals that are inferred to have significant levels of
admixture were collected from morphologically intermediate populations of *O. hartwegii* subsp.
pubescens and *O. hartwegii* subsp. *hartwegii*. In addition, *O. hartwegii* subsp. *hartwegii* was a
parent in all three hybridization events (S10). Thus, gene flow may explain this taxon's non-
monophyly in our summary coalescent results. However, despite the often confounding patterns
of overlapping morphological variation among closely related taxa in subsect. *Salpingia*, this
pattern does not necessarily appear to be the result of admixture, as many of the tests for
hybridization based on field observations were not significant (Fig 3a, S10). What is also clear
from these results is that much like collecting hundreds of nuclear genes provides a more
nuanced picture of phylogenetic signal and taxon relationships, our results show that collecting
multiple individuals from across the geographic and morphological ranges is necessary for a
more complete picture of relationships among closely related taxa.
291   After filtering, we extracted a set of 9728 single nucleotide polymorphisms (SNPs) from
both coding and non-coding regions. A Discriminant Analysis of Principal Components (DAPC;
Fig. 3b) clearly distinguishes subsection *Salpingia* from subsection *Calylophus*, with *O. toumeyi*
intermediate between the two, which is consistent with the phylogenetic results presented here.
Additionally, the DAPC identifies *O. lavandulifolia* as a distinct genetic cluster from the
remaining taxa in subsection *Salpingia*. The overlap between taxa, for example between the
remaining taxa in subsection *Salpingia*, is consistent with the high levels of gene tree
discordance identified by PhyParts (Fig. 3a). For this latter group of taxa, we computed estimates
of ancestry coefficients using sNMF, which suggests a substantial amount of shared ancestral
polymorphisms while also showing some evidence of clear genetic structure among taxa (Fig.
3c). Consistent with the phylogenetic analyses, there does not appear to be any clear genetic
distinction between *O. hartwegii* subsp. *hartwegii* and *O. hartwegii* subsp. *maccartii*, while *O.*
hartwegii subsp. *fendleri*, *O. tubicula* subsp. *tubicula,* and *O. hartwegii* subsp. *filifolia* appear to
be largely distinct.
305
Morphological Analysis
We conducted morphometric Principal Components Analysis (PCA) to determine if
morphological patterns were consistent with phylogentic results and to examine if specific
characters could be used to diagnose taxa as circumscribed by our phylogenetic analysis. Towner
(1977) observed overlapping and confounding patterns of morphological variation among taxa
within subsections, as for example in the *O. hartwegii* species complex. Despite this, because
some taxa (e.g., *O. hartwegii* subsp. *fendleri*) were strongly supported by our summary
coalescent trees we expected that they would be well distinguished in morphometric analysis.
In general, we found that the main traits that separated taxa in subsect. *Salpingia* were leaf
traits and plant size, while in subsect. *Calylophus* the main traits that separated taxa were sepal
traits. In subsect. *Salpingia*, PC1 accounted for 44.2% of variance in PCA, while PC2 accounted
for 28.1% (Fig. 4b). Morphological characters most associated with PC1 were leaf width (distal
and basal), plant height, and leaf length/width ratio (distal and basal). Those associated with PC2
were leaf length (distal and basal), sepal length, and sepal tip length (Fig. 4b). In subsect.
Calylophus, PC1 accounted for 43.8% of explained variance and PC2 accounted for 27.4% (Fig.

4c). The characters most associated with PC1 in subsect. *Calylophus* include leaf length (distal
and basal), sepal length, and sepal tip length. Those most associated with PC2 were leaf
length/width ratio (distal and basal) and distal leaf width (Fig. 4c).
Our results support Towner's understanding of taxon boundaries by underscoring previous
difficulties in identifying individuals in this group based on morphology. We found substantial
overlapping morphological variation among currently recognized taxa in both subsections,
though some taxa exhibited better grouping than others. The amount of overlap between taxa
was not a function of the strength of tree support for a given taxon in our summary coalescent
results. For example, *O. hartwegii* subsp. *fendleri*, a taxon that was well supported in our
summary coalescent trees, exhibited some of the highest degree of overlap with other taxa in
PCA space. Conversely, both *O. hartwegii* subsp. *filifolia* and *O. hartwegii* subsp. *pubescens*,
two taxa that formed poorly supported clades in our trees, formed clusters on the outer edges of
PCA space and had less overlap than other taxa (Fig. 4b). Interestingly, *O. hartwegii* subsp.
hartwegii, the taxon that was identified as a parent in all three instances of admixture in our
HyDe analysis, also overlaps in PCA with most other taxa in subsect. *Salpingia* (Fig. 4b). This is
not surprising given that it is widely distributed in northern Mexico and western Texas, and
frequently comes into contact with related taxa resulting in sympatric populations and occasional
morphologically intermediate populations.
339
Implications for Reproductive Systems and Edaphic Endemism
Our results show shifts from hawkmoth to bee pollination likely occurred twice in sect.
Calylophus (S3) and thus may be more common in Oenothera than previously thought. The
strongly supported sister relationship of *O. toumeyi* to remaining subsect. *Calylophus* in our
summary coalescent results is consistent with two independent shifts, once in the ancestor of
subsect. *Calylophus*, and another in subsect. *Salpingia* on the branch leading to *O. tubicula* (Fig.
2). Independent shifts to bee pollination from hawkmoth pollination are perhaps not surprising
considering that within sect. *Calyophus*, hawkmoth-pollinated floral forms exhibit plasticity in
hypanthium length and diameter and do not prevent occasional pollination by bees (Lewis et al.
in prep; Towner 1977). Hawkmoth-pollinated taxa in sect *Calylophus* exhibit vespertine anthesis,
which separates them temporally from diurnal bees, but plasticity in the timing of anthesis is also
common among populations (Towner 1977), and hawkmoths are documented to vary
spatiotemporally in abundance (Miller 1981; Campbell et al. 1997; Artz et al. 2010). Aditionally,
it has been shown that florivore-mediated selection drives floral trait shifts in sect. *Calylophus*
towards bee pollinated floral forms (Jogesh et al. 2017; Bruzzese et al. 2019). Plasticity in
reproductive traits that allow some continued pollination by bees provides an alternative mode of
pollen transfer and may represent a mechanism for ensuring pollination. While studies have
shown that premating barriers such as these contribute greatly to reproductive isolation (Stanton
et al. 2016), our results show that multiple, indepdendent shifts from hawkmoth to bee
pollination and associated morphological changes, such as the shorter corolla length of bee
pollinated flowers, may occur in sect. *Calylophus*, and hence may not be a particularly reliable
character for diagnosing taxa in this group.
Stochastic mapping (supplemental) suggests that there are multiple origins of permanent
translocation heterozygosity (PTH) in sect. *Calylophus* While ring chromosomes are common
and found in all taxa in sect. *Calylophus*, PTH is currently known from only one taxon, *O.*
serrulata. Because neighboring populations of *O. serrulata* and its putative progenitor O.
capillifolia subsp. *berlandieri* often resemble each other phenetically, Towner (1977)

hypothesized that *O. serrulata* may have originated multiple times through independent origins of translocation heterozygosity in different geographic regions, and may be best recognized as "a complex assemblage of populations having a common breeding system." However, this has never before been explored in a phylogenetic context, nor has it been clearly demonstrated with phylogenetic studies in Onagraceae. In our summary coalescent trees, all currently recognized taxa in subsect. *Calylophus* were paraphyletic and *O. serrulata* was scattered throughout the subsection (Fig. 2, S6, S7). Although support values are not always high for the positions of various populations of *O. serrulata*, there is at least one well defined, well supported split among populations of *O. serrulata*. In our summary coalescent trees, the two *O. serrulata* accessions from south Texas (MJM 970 & MJM 983) grouped with other south Texas populations of *O. capillifolia* subsp. *berlandieri* with generally strong support (Fig. 2, S6, S7). This relationship was supported in PCA space as well, where MJM 983 was morphologically more similar to the south Texas *O. capillifolia* subsp. *berlandieri* accessions than to other *O. serrulata* (Fig 4C). Our results are therefore consistent with an independent origin of PTH in coastal Texas populations of *O. serrulata*, demonstrating at minimum two origins of PTH (see *Taxonomic Implications* below). However we cannot rule out other independent origins of PTH, for example in the populations of *O. capillifolia* subsp. *berlandieri* occupying sand dunes in western Texas and southeastern New Mexico, or in the other taxa in subsect. *Calylophus*: *O. capillifolia* subsp. *berlandieri*, or the gypsum endemic *O. gayleana*.

Independent origins of gypsum endemism in sect. *Calylophus* are also supported by our analyses (supplemental). Edaphic specialization is a fundamental driver of speciation in plants and contributes greatly to endemism and species diversity in areas with geologically distinct substrates such as gypsum and serpentine outcrops (Kruckeberg 1984; Anacker et al. 2011; Cacho and Strauss 2014; Moore et al. 2014). To date, two gypsum endemic taxa have been described in sect. *Calylophus*, one in each subsection: *O. hartwegii* subsp. *filifolia*, which is relatively widespread on gypsum in New Mexico and trans-Pecos Texas and only rarely grows in sympatry with other taxa, and the recently described *O. gayleana*, which is found in southeastern New Mexico and adjacent western Texas, with disjunct populations in northern Texas and western Oklahoma (Turner and Moore 2014). Despite low support and low gene tree congruence in our analyses, the two gypsum endemic taxa had moderate sCF support (Fig. 2, S9), much like other taxa with similarly low support and high levels of discordance. In addition,while both gypsum endemics overlapped with other taxa in our morphometric analysis, they occupied morphological extremes in PCA space (Fig. 4). Given that other well-supported taxa are also not well differentiated in PCA space, it is not remarkable that the two gypsum endemic taxa were also not more differentiated from other taxa morphologically. Perhaps the strongest evidence in our data for their recognition is that we found no evidence of admixture between either of these gypsum endemic taxa and other closely related taxa (Fig. 3a, S10).

Taxonomic Implications

The most consequential taxonomic result that arises from our analysis is the position of *O. toumeyi*, a member of subsect. *Salpingia*, as sister to subsect. *Calylophus* with strong support, rendering subsect. *Salpingia* paraphyletic (Fig. 2, S9). The current taxonomy groups *O. toumeyi* with *O. hartwegii* due to similar floral and bud characters including large flowers and long floral tubes suggestive of hawkmoth pollination, and rounded buds with long, free sepal-tips (Towner 1977). Because the breeding system is a defining difference in the current circumscription

between the two subsections in sect. *Calylophus*, this result supports abandoning subsections
altogether in sect. *Calylophus*.
Within subsect. *Salpingia* our results also suggest the need for revision. While our
phylogenetic analyses strongly support the current circumscription of *O. lavandulifolia* (sensu
Towner 1977) as a distinct species within subsect. *Salpingia* (Fig. 2, S9), the relationships of the
other two species *O. hartwegii* and *O. tubicula* are less clear. Towner (1977) differentiated these
two species by the breeding system and grouped the five subspecies of *O. hartwegii* together
based on a pattern of reticulate and intergrading variation in which taxa were distinguished from
one another by often slight differences in pubescence and leaf shape. Our morphometric analysis
confirmed this pattern; however, our phylogenetic results indicated that one taxon, *O. hartwegii*
subsp. *fendleri*, shares a closer relationship with the bee pollinated *O. tubicula* subsp. *tubicula*
than other taxa in the hawkmoth pollinated *O. hartwegii* species complex (Fig. 2, S6, S7). This
relationship was strongly supported and renders *O. hartwegii*, according to the current
circumscription, paraphyletic (Towner 1977). Based on strong phylogenetic support for this
clade, and its strong morphological distinctiveness as described by Towner (1977), we suggest
that *O. hartwegii* subsp. *fendleri* be elevated to the species level along with both races of *O.*
tubicula which were equally well supported in phylogenetic analysis and are geographically
isolated. Furthermore, our results support a specific distinction for *O. hartwegii* subsp. *filifolia*.
While this taxon was poorly supported in our summary coalescent trees (Fig. 2, S6, S7), we
found no evidence of hybridization between this taxon and other closely related taxa. In addition,
O. hartwegii subsp. *fillifolia* is restricted to gypsum. Therefore, we believe that the ecological
distinctiveness and lack of gene flow of *O. hartwegii* subsp. *filifolia* with other taxa in the *O.*
hartwegii species complex warrants its elevation as a distinct species. In light of these changes,
and to maintain consistency in classification in the subsection, we feel that despite the evidence
of hybridization of *O. hartwegii* subsp. *pubescens* with *O. hartwegii* subsp. *hartwegii*, it
possesses a morphological distinctiveness that is supported by our phylogenetic results. We
therefore recommend this taxon be elevated to the species level, while *O. hartwegii* subsp.
hartwegii and *O. hartwegii* subsp. *maccartii* be retained as is, forming a polytypic species with
two subspecies.
In contrast to the relatively clear divisions among taxa in subsect. *Salpingia* in our
coalescent trees*,* none of the four currently recognized taxa in the subsect. *Calylophus* were
consistently recovered as monophyletic. For example, *O. capillifolia* subsp. *capillifolia* was
monophyletic in our exon-only summary coalescent tree, but not in the "supercontig" tree, and
O. capillifolia subsp. *berlandieri* and *O. serrulata* were scattered throughout sect. *Calylophus* in
both trees, perhaps suggesting widespread gene flow and/or multiple origins of PTH (Fig. 2, S6).
Importantly, our results suggest that the circumscription of *O. gayleana* sensu Turner and Moore
(2014) should be amended. Specifically, we find that the populations of subsect. *Calylophus*
from northern Texas and western Oklahoma that were assigned to *O. gayleana* by Turner and
Moore (2014; MJM790-5, BJC71) instead may belong to *O. serrulata* based on both their
phylogenetic positions (Fig. 2, S6, S7) and reduced pollen fertility (S12). These north
Texas/western Oklahoma populations seem to represent slightly narrower-leaved individuals of
O. serrulata, which is a common inhabitant of the extensive gypsum outcrops of this area
(although it is not restricted to gypsum there).
Finally, our results highlight an unrecognized cryptic taxon within *O. capillifolia* formed
by southern Texas coastal populations currently recognized as *O. capillifolia* subsp. *berlandieri*.
Towner (1977) described *O. capillifolia* as a polytypic species with two well-differentiated

morphological races. Though he noted the geographic and cytological distinction of the southern
Texas coastal populations of *O. capillifolia* subsp. *berlandieri*, these populations were included
in *O. capillifolia* subsp. *berlandieri* primarily because of, "completely overlapping
morphological variation." In our results, this cryptic clade of southern Texas coastal populations
of *O. capillifolia* subsp. *berlandieri* is the most phylogenetically well supported clade in subsect.
Calylophus and therefore may warrant taxonomic distinction based on our data (Fig. 2, S9r).
Similarly, the southern Texas coastal populations of *O. serrulata*, which is likely an independent
origin of PTH derived from this cryptic southern Texas coastal clade of *O. capillifolia* subsp.
berlandieri, are ecologically distinctive and geographically disjunct from other *O. serrulata*
(occurring in coastal dunes, unlike other populations in western Texas, Oklahoma, and northern
Texas). In the past they were considered distinctive enough to be described as a species,
Calylophus australis (Towner & Raven 1970). However, Towner (1977) later sunk this variation
into *O. serrulata* based on his decision to treat all PTH populations as *O. serrulata*. Combined
with our results here and the ecogeographic distinctiveness consistent with an independent origin
of PTH in coastal Texas, we believe this taxon also warrants recognition as a second PTH
species in *Oenothera* sect. *Calylophus*.
474
475 CONCLUSIONS
476
Here we describe a robust example of resolving a recent, rapid radiation using multiple
sources of evidence: (1) extensive sampling from populations throughout the geographic and
morphological range, (2) target enrichment for hundreds of nuclear genes, (3) the inclusion of
flanking non-coding regions, (4) gene tree-based hybridization inference, (5) SNPs extracted
from target enrichment data, and (6) morphometrics. Our results indicate that in recently radiated
species complexes with low sequence divergence and/or high levels of ILS that could be an
intractable problem with traditional loci, the use of targeted enrichment in addition to flanking
non-coding regions provides a net benefit and is essential to recover species-level resolution. Our
results also underscore the importance of summary coalescent methods and evaluating gene tree
discordance for resolving historical relationships in recalcitrant groups. By explicitly testing for
hybridization using gene tree approaches, we also demonstrate that the estimated number of
character state transitions may be artifactually inflated if hybridization is not taken into account.
This, in combination with morphometrics, provided key evolutionary insights where
relationships in summary coalescent methods may be obscured by gene flow. Importantly, our
study uncovers strong evidence for multiple origins of biologically important phenomena,
including PTH, the evolution of bee pollination, and the evolution of edaphic specialization.
Consequently, *Oenothera* sect. *Calylophus* might represent a powerful system for understanding
these phenomena, especially with future genome sequence data.
495
496 FUNDING SOURCES
497

498 This work was supported by National Science Foundation grants DEB-1342873 to KAS, JAF,
499 and NJW, and DEB-1054539 to MJM. Additional support was provided by the National
500 Geographic Society, Oberlin College;, The Chicago botanic Garden, The Shaw Fellowship, New
501 Mexico Native Plant Society, American Society of Plant Taxonomists, and The Society of
502 Herbarium Curators.

REFERENCES

Alexander M.P. 1969. Differential Staining of Aborted and Nonaborted Pollen. Stain Technol. 44:117–122.

Alexander M.P. 1980. A Versatile Stain for Pollen Fungi, Yeast and Bacteria. Stain Technol. 55:13–18.

Anacker B.L., Whittall J.B., Goldberg E.E., Harrison S.P. 2011. Origins and consequences of serpentine endemism in the California flora. Evolution (N. Y). 65:365–376.

Artz D.R., Villagra C. A., Raguso R. A. 2010. Spatiotemporal variation in the reproductive ecology of two parapatric subspecies of *Oenothera cespitosa* (Onagraceae). Am. J. Bot. 97:1498–510.

Barrett S., Harder L., Worley A. 1996. The comparative biology of pollination and mating in floweing plants. Philos. Trans. R. Soc. B Biol. Sci. 351:1271–1280.

Barrett S.C.H. 2013. The evolution of plant reproductive systems: how often are transitions irreversible? Proc R Soc B. 280:20130913.

Blischak P.D., Chifman J., Wolfe A.D., Kubatko L.S. 2018. HyDe: A python package for genome-scale hybridization detection. Syst. Biol. 67:821–829.

Boberg E., Alexandersson R., Jonsson M., Maad J., Ågren J., Nilsson L.A. 2014. Pollinator shifts and the evolution of spur length in the moth-pollinated orchid *Platanthera bifolia*. Ann. Bot. 113:267–275.

Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics. 30:2114–2120.

Brady K.U., Kruckeberg A.R., Bradshaw H.D. 2005. Evolutionary ecology of plant adaptation to serpentine soils. Annu. Rev. Ecol. Evol. Syst. 36:243–266.

Bruzzese D.J., Wagner D.L., Harrison T., Jogesh T., Overson R.P., Wickett N.J., Raguso R.A., Skogen K.A. 2019. Phylogeny, host use, and diversification in the moth family Momphidae (Lepidoptera: Gelechioidea). PLoS One. 14:e0207833.

Bryson R.W., Linkem C.W., Dorcas M.E., Lathrop A., Jones J.M., Alvarado-Díaz J., Grünwald C.I., Murphy R.W. 2014. Multilocus species delimitation in the *Crotalus triseriatus* species group (serpentes: Viperidae: Crotalinae), with the description of two new species. Zootaxa. 3826:475–496.

Cacho N.I., Strauss S.Y. 2014. Occupation of bare habitats, an evolutionary precursor to soil specialization in plants. Proc. Natl. Acad. Sci. 111:15132–15137.

Campbell D.R.., Waser N.M.., Melendez-Ackerman E.J.. 1997. Analyzing pollinator-mediated selection in a plant hybrid zone: Hummingbird visitation patterns on three spatial scales. Am. Nat. 149:295–315.

Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model. Bioinformatics. 30:3317–3324.

Chifman J., Kubatko L. 2015. Identifiability of the unrooted species tree topology under the coalescent model with time-reversible substitution processes, site-specific rate variation, and invariable sites. J. Theor. Biol. 374:35–47.

Christie M.R., Knowles L.L. 2015. Habitat corridors facilitate genetic resilience irrespective of species dispersal abilities or population sizes. Evol. Appl. 8:454-463.

Crepet W.L., Niklas K.J. 2009. Darwin's second "abominable mystery": Why are there so many angiosperm species? Am. J. Bot. 96:366–81.

Duarte J.M., Wall P.K., Edger P.P., Landherr L.L., Ma H., Pires J.C., Leebens-Mack J.,

dePamphilis C.W. 2010. Identification of shared single copy nuclear genes in *Arabidopsis,*
Populus, Vitis and *Oryza* and their phylogenetic utility across various taxonomic levels.
BMC Evol Biol. 10:61.

Eckert A.J., Carstens B.C. 2008. Does gene flow destroy phylogenetic signal? The performance
of three methods for estimating species phylogenies in the presence of gene flow. Mol.
Phylogenet. Evol. 49:832–842.

Ehrlich, P. R., & Raven, P. H. (1969). Differentiation of populations. Science. 165:1228–1232.

Folk R.A., Mandel J.R., Freudenstein J. V. 2015. A protocol for targeted enrichment of intron-
containing sequence markers for recent radiations: A phylogenomic example from
Heuchera (Saxifragaceae). Appl. Plant Sci. 3:1500039.

Frichot E, Mathieu F, Trouillon T, Bouchard G and Fran¸cois O (2014) Fast and efficient
estimation of individual ancestry coefficients. Genetics. 196:973-983.

Gerard D., Gibbs H.L., Kubatko L. 2011. Estimating hybridization in the presence of
coalescence using phylogenetic intraspecific sampling. BMC Evol. Biol. 11:291.

Giarla T.C., Esselstyn J.A. 2015. The challenges of resolving a rapid, recent radiation: Empirical
and simulated phylogenomic of Philippine shrews. Syst. Biol. 64(5):727-740.

Green R., Krause J., Briggs A., Rasilla Vives M., Fortea Pérez F. 2010. A draft sequence of the
neandertal genome. Science. 328:710–722.

Heyduk K., Trapnell D.W., Barrett C.F., Leebens-mack J.I.M. 2016. Phylogenomic analyses of
species relationships in the genus *Sabal* (Arecaceae) using targeted sequence capture. Biol.
J. Linn. Soc. 117(1):106-120.

Jogesh T., Overson R.P., Raguso R.A., Skogen K.A. 2017. Herbivory as an important selective
force in the evolution of floral traits and pollinator shifts. AoB Plants. 9(1):plw088.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett
N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics from high-
throughput sequencing reads using target enrichment. Appl. Plant Sci. 4:1600016.

Johnson M.T.J., Smith S.D., Rausher M.D. 2009. Plant sex and the evolution of plant defenses
against herbivores. Proc. Natl. Acad. Sci. 106:18079–18084.

Jombart, T. (2008) adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics 24:1403-1405.

Jombart T, Devillard S and Balloux, F (2010). Discriminant analysis of principal components: a
new method for the analysis of genetically structured populations. BMC Genetics. 11:94.

Katinas L., Crisci J., Wagner W., Hoch P. 2004. Geographical diversification of tribes
Epilobieae, Gongylocarpeae, and Onagreae (Onagraceae) in North America, based on
parsimony analysis of endemicity and track compatibility analysis. Ann. Missouri Bot.
91:159–185.

Knowles L.L. 2009. Estimating species trees: Methods of phylogenetic analysis when there is
incongruence across genes. Syst. Biol. 58:463–467.

Knowles L.L., Chan Y.-H. 2008. Resolving species phylogenies of recent evolutionary
radiations. Ann. Missouri Bot. Gard. 95:224–231.

Kruckeberg A. 1984. California Serpentines: Flora, Vegetation, Geology, Soils, and
Management Problems. Berkeley: Univ of California Press.

Kubatko L.S., Chifman J. 2019. An invariants-based method for efficient identification of hybrid
species from large-scale genomic data. BMC Evol. Biol. 19:1–13.

Leaché A.D., Harris R.B., Rannala B., Yang Z. 2014. The influence of gene flow on species tree
estimation: a simulation study. Syst. Biol. 63:17–30.

Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored hybrid enrichment for massively
high-throughput phylogenomics. Syst. Biol. 61:727–744.
Levin R., Wagner W., Hoch P. 2004. Paraphyly in Tribe Onagreae : Insights into Phylogenetic
Relationships of Onagraceae Based on Nuclear and Chloroplast Sequence Data. Syst. Bot.
29:147–164.
Maddison W.P., Knowles L. 2006. Inferring Phylogeny Despite Incomplete Lineage Sorting.
Evol. Biol. 55:21–30.
Mamanova L., Coffey A.J., Scott C.E., Kozarewa I., Turner E.H., Kumar A., Howard E.,
Shendure J., Turner D.J. 2010. Target-enrichment strategies for next- generation
sequencing. Nat. Methods. 7:111–118.
Mandel J.R., Dikow R.B., Funk V. a, Masalia R.R., Staton S.E., Kozik A., Michelmore R.W.,
Rieseberg L.H., Burke J.M. 2014. A target enrichment method for gathering phylogenetic
information from hundreds of loci: An example from the Compositae. Appl. Plant Sci. 2:1–
6.
Meng C., Kubatko L.S. 2009. Detecting hybrid speciation in the presence of incomplete lineage
sorting using gene tree incongruence: A model. Theor. Popul. Biol. 75:35–45.
Miller R.B.. 1981. Hawkmoths and the Geographic Patterns of Floral Variation in *Aquilegia*
caerulea. Evolution (N. Y). 35:763–774.
Minh B.Q., Hahn M.W., Lanfear R. 2020. New methods to calculate concordance factors for
phylogenomic datasets. Mol. Biol. Evol. 37(9):2727-2733.
Mirarab S., Warnow T. 2015. ASTRAL-II: Coalescent-based species tree estimation with many
hundreds of taxa and thousands of genes. Bioinformatics. 31:i44–i52.
Moore M., Jansen R. 2007. Origins and biogeography of gypsophily in the Chihuahuan Desert
plant group *Tiquilia* Subg. subg. *Eddya*. Syst. Bot. 32:392–414.
Moore M.J., Mota J.F., Douglas N.A., Flores Olvera H., Ochoterena H. 2014. Ecology assembly
evolution gypsophile floras. In: Rajakaruna N., Boyd R.S., Harris T.B., editors. Plant
Ecology and Evolution in Harsh Environments. New York: Nova Science Publishers, Inc. p.
97–128.
Nason J.D., Hamrick J.L., Fleming T.H. 2002. Historical vicariance and postglacial colonization
effects on the evolution of genetic structure in *Lophocereus*, a Sonoran Desert columnar
cactus. Evolution 56:2214–2226.
Van der Niet T., Johnson S.D. 2012. Phylogenetic evidence for pollinator-driven diversification
of angiosperms. Trends Ecol. Evol. 27:353–361.
van der Niet T., Johnson S.D., Linder H.P. 2006. Macroevolutionary data suggest a role for
reinforcement in pollination system shifts. Evolution 60:1596–1601.
Van Der Niet T., Peakall R., Johnson S.D. 2014. Pollinator-driven ecological speciation in
plants: New evidence and future perspectives. Ann. Bot. 113:199–211.
Peakall R., Ebert D., Poldy J., Barrow R.A., Francke W., Bower C.C., Schiestl F.P. 2010.
Pollinator specificity, floral odour chemistry and the phylogeny of Australian sexually
deceptive *Chiloglottis* orchids: implications for pollinator-driven speciation. New Phytol.
188:437–50.
Rajakaruna N. 2004. The Edaphic Factor in the Origin of Plant Species. Int. Geol. Rev. 46:471–
478.
Raven P.H. 1964. Catastrophic Selection and Edaphic Endemism. Evolution 18:336–338.
Raven P.H. 1979. A survey of reproductive biology in Onagraceae. New Zeal. J. Bot. 17:575–
593.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
    Statistical Computing, Vienna, Austria.

Roch S., Steel M. 2015. Likelihood-based tree reconstruction on a concatenation of aligned
    sequence data sets can be statistically inconsistent. Theor. Popul. Biol. 100:56–62.

Sayyari E., Mirarab S. 2016. Fast coalescent-based computation of local branch support from
    quartet frequencies. Mol. Biol. Evol. 33(7):1654-68.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27(4):592-593.

Skogen K.A., Overson R.P., Hilpman E.T., Fant J.B. 2019. Hawkmoth pollination facilitates
    long-distance pollen dispersal and reduces isolation across a gradient of land-use change.
    Ann. Missouri Bot. Gard. 104:495–511.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals
    conflict, concordance, and gene duplications with examples from animals and plants. BMC
    Evol. Biol. 15:150.

Stamatakis A. 2014. Stamatakis - 2014 - RAxML version 8 a tool for phylogenetic analysis and
    post-analysis of large phylogenies. 2010–2011.

Stanton K., Valentin C.M., Wijnen M.E., Stutstman S., Palacios J.J., Cooley A.M. 2016.
    Absence of postmating barriers between a selfing vs. Outcrossing chilean *Mimulus* species
    pair. Am. J. Bot. 103:1030–1040.

Stebbins G.L. 1970. Adaptive Radiation of Reproductive Characteristics in Angiosperms, I:
    Pollination Mechanisms. Annu. Rev. Ecol. Syst. 1:307–326.

Stephens J.D., Rogers W.L., Heyduk K., Cruse-Sanders J.M., Determann R.O., Glenn T.C.,
    Malmberg R.L. 2015. Resolving phylogenetic relationships of the recently radiated
    carnivorous plant genus *Sarracenia* using target enrichment. Mol. Phylogenet. Evol. 85:76–
    87.

Stockhouse R.E.I. 1973. Biosystematic Studies of *Oenothera* L. Subgenus *Pachylophus*. Univ.
    Microfilm. Ph.D. thesis. Colorado State University.

Straub S.C.K., Parks M., Weitemier K., Fishbein M., Cronn R.C., Liston A. 2012. Navigating the
    tip of the genomic iceberg: Next-generation sequencing for plant systematics. Am. J. Bot.
    99:349–64.

Swofford, D. L. 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods).
    Version 4. Sinauer Associates, Sunderland, Massachusetts.

Thomson J.D., Wilson P. 2008. Explaining Evolutionary Shifts between Bee and Hummingbird
    Pollination: Convergence, Divergence, and Directionality. Int. J. Plant Sci. 169:23–38.

Towner H.F., Raven P.H. 1970. A new species and some new combinations in *Calylophus*
    (Onagraceae). Madrono. 20: 241-245.

Towner H.F. 1977. The Biosystematics of *Calylophus* (Onagraceae). Ann. Missouri Bot. Gard.
    64:48–120.

Tripp E.A., Manos P.S. 2008. Is floral specialization an evolutionary dead-end? Pollination
    system transitions in *Ruellia* (Acanthaceae). Evolution 62:1712–1737.

Turner B.L., Moore M.J. 2014. *Oenothera gayleana* (*Oenothera* sect. *Calylophus*, Onagraceae),
    a new gypsophile from Texas, New Mexico, and Oklahoma. Phytologia.org. 96:200–206.

Wagner W., Hoch P., Raven P. 2007. Revised Classification of the Onagraceae. Syst. Bot.
    Monogr. 83:1–240.

Wagner W.L. In press. *Oenothera.* In: Flora of North America Editorial Committee, editor. Flora
    of North America North of Mexico. New York and Oxford. Vol. 10.

Weitemier K., Straub S.C.K., Cronn R.C., Fishbein M., Schmickl R., McDonnell A., Liston A.

2014. Hyb-Seq: Combining Target Enrichment and Genome Skimming for Plant
Phylogenomics. Appl. Plant Sci. 2:1400042.
Wilson P., Wolfe A.D., Armbruster W.S., Thomson J.D. 2007. Constrained lability in floral
evolution: Counting convergent origins of hummingbird pollination in Penstemon and
Keckiella. New Phytol. 176:883–890.
Xi Z., Liu L., Rest J.S., Davis C.C. 2014. Coalescent versus Concatenation Methods and the
Placement of *Amborella* as Sister to Water Lilies. Syst. Biol. 63:919–932.
Xu S., Schlüter P.M., Scopece G., Breitkopf H., Gross K., Cozzolino S., Schiestl F.P. 2011.
Floral isolation is the main reproductive barrier among closely related sexually deceptive
orchids. Evolution. 65:2606–20.

700
701
Figure 1. (a) Range map of all taxa in *Oenothera* sect. *Calylophus* (based on Towner 1977).
Sampling locations of leaf tissue samples (points; color corresponds to taxa in cladogram to the
right [Figure 1b]) and estimated taxon ranges (polygons; colors correspond to Figure 1b)
proposed by Towner (1977) and Turner and Moore (2014). (b) Estimated cladogram of
Oenothera sect. *Calylophus* sensu Towner (1977) and Turner and Moore (2014). Symbols to the
right of tip labels signify pollination system (bee, hawkmoth, or Permanent Translocation
Heterozygosity [PTH]) and the symbol color specifies whether a given taxon is a gypsum
endemic (purple = no, pink = yes). Photo panels: (c) *Oenothera hartwegii* subsp. *fendleri* (d)
Oenothera hartwegii subsp. *filifolia* (e) *Oenothera hartwegii* subsp. *hartwegii* (f) *Oenothera*
hartwegii subsp. *maccartii* (g) *Oenothera hartwegii* subsp. *pubescens* (h) *Oenothera tubicula*
subsp. *tubicula* (i) *Oenothera lavandulifolia* (j) *Oenothera tubicula* subsp. *strigulosa* (k)
Oenothera capillifolia subsp. *berlandieri* (l) *Oenothera capillifolia* subsp. *capillifolia* (m)
Oenothera gayleana (n) *Oenothera serrulata* (o) *Oenothera toumeyi*
715

Figure 2. ASTRAL-II summary coalescent tree constructing using the "Supercontig" dataset with 100 bootstraps. At relevent nodes, piecharts represent Phyparts analysis (blue = concordant, green = most conflict, red = all other conflict, black = uninformative gene trees), top row of support values are bootstrap values from ASTRAL-II, and gCF and sCF from IQTree (BS/gCF/sCF), bottom two support values are number of concordant gene trees for the node and total number of gene trees minus the number of concordant gene trees at that node (concord/discord). Colored tip points correspond to taxon designation.

Figure 3. (a) Summary of HyDe Analysis annotated on ASTRAL-III species tree constructed using the "Supercontig" dataset; black arrows represent direction of admixture detected by HyDe anaylsis. (b) Discriminant Analysis of Principal Components based on a filtered set of SNPs extracted from the entire supercontig dataset , and (c) sNMF plot of inferred ancestry coefficients using the same set of filtered SNPs but limited to subsection *Salpingia*.

732
733
734 Figure 4. Morphometric Principal Components Analysis (PCA) using 9 morphological characters
735 (plant height, leaf length [basal and distal], leaf width [basal and distal], leaf length/width ratio
736 (basal and distal), and sepal tip length) for (a) Section *Calylophus* (b) Subsection *Salpingia*
737 without *O. toumeyi*, and (c) Subsection *Calylohus* with *O. toumeyi* included.

SUPPLEMENTAL MATERIAL

S1
See excel table "S1 Accessions and Seq Stats"

S2
Number of leaf tissue accessions sequenced from each taxon

Taxon	No. of Accessions
O. lavandulifolia	16
O. tubicula subsp. *strigulosa*	3
O. tubicula subsp. *tubicula*	9
O. hartwegii subsp. *fendleri*	15
O. hartwegii subsp. *filifolia*	44
O. hartwegii subsp. *hartwegii*	12
O. hartwegii subsp. *maccartii*	9
O. hartwegii subsp. *pubescens*	26
O. hartwegii	1
O. toumeyi	5
O. capillifolia subsp. *berlandieri*	17
O. capillifolia subsp. *capillifolia*	11
O. gayleana	9
O. serrulata	17
Outgroups:	
O. pilosella	1
O. organensis	1
O. primiveris	1
O. tubifera	1
O. triloba	1
O. cespitosa subsp. *cespitosa*	1
O. suffrutescens	1
Chylismia scapoidea subsp. *scapoidea*	1
Eulobus californicus	1
Total	203

S3
SUPPLEMENTAL MATERIALS AND METHODS

Taxon Sampling, DNA Extraction, and Determination of PTH

A total of 194 individuals spanning the geographic, morphological, and ecological ranges of all 13 recognized taxa in *Oenothera* sect. *Calylophus* [following Towner (1977) and Turner and Moore (2014)] were included in this study (Fig. 1a, S1, S2) along with eight outgroups representing other major sections of Oenothera (*Eremia*, *Gaura*, *Kneiffia*, *Lavauxia*, *Oenothera*, *Pachylophus*, and *Ravenia*) and other genera (*Chylismia* and *Eulobus*) in Onagraceae (S1, S2). All leaf tissue samples were collected from individuals in the field between 2007 and 2015 and voucher specimens were deposited at the United States National Herbarium (US), with

duplicates in most cases at either the Nancy Rich Poole Herbarium (CHIC) or the George T.
Jones Herbarium at Oberlin College (OC; S1). DNA was extracted from fresh, silica-dried leaf
tissue using either (1) a modified CTAB protocol (Doyle 1987), (2) the Nucleon PhytoPure DNA
extraction kit (GE Healthcare Life Sciences, Pittsburgh, Pennsylvania, USA), or (3) a modified
CTAB and silicon dioxide purification protocol (Doyle 1987; Sharma and Purohit 2012; See S13
for detail) followed by passing any extractions retaining a brown or yellow coloration through a
Qiagen Qiaquick PCR spin column for additional purification according to the manufacturer's
protocol (Qiagen, Venlo, Netherlands). The third DNA extraction method was used for difficult
to extract, polysaccharide-rich leaf tissue samples that yielded gooey, discolored DNA following
initial extraction. PTH status was determined for individuals in subsect. *Calylophus* using floral
morphology and/or when flowers were present by assessing pollen fertility using a modified
Alexander stain, as PTH taxa have a demonstrated 50% reduction in pollen fertility (Towner
1977). For accessions identified as either *O. capillifolia* subsp. *berlandieri* or *O. serrulata* that
had sufficient pollen available, pollen was removed from flowers and stained using a modified
Alexander stain (Alexander 1969, 1980). Accessions with less than 50% viable pollen were
assigned to *O. serrulata*, the only currently recognized PTH taxon in subsect. *Calylophus*. Pollen
count data are provided in Supplement 12 (See S12 for details).
797
Bait Design, Library Construction, Target Enrichment, and Sequencing
We targeted 322 orthologous, low-copy nuclear loci determined by clustering
transcriptomes of *Oenothera serrulata* (1KP accession SJAN) and *Oenothera berlandieri* (1KP
accession EQYT) to select a subset of the 956 phylogenetically informative *Arabidopsis* loci
identified by Duarte et al. (2010). Transcriptomes of two *Oenothera* species, *O. serrulata* and *O.*
berlandieri, were assembled and optimal isoforms were filtered for the longest reading frame.
Assembled transcripts were aligned as amino acids to the 956 TAIR loci of *Arabidopsis* in
TranslatorX (Abascal et al. 2010). This alignment identified 956 orthologous sequences, from
which 322 loci were randomly selected. BLAST searches of amino acid sequences from these
loci were carried out to ensure orthology between the transcript loci and the *Arabidopsis* TAIR
locus. The bait set was designed from these 322 loci, which were selected from both *O. serrulata*
and *O. berlanderi* sequences. A set of 19,994 120-bp baits tiled across each locus with a 60 base
overlap (2x tiling) was manufactured by Arbor Biosciences (formerly MYcroarray, Ann Arbor,
Michigan, USA). Sequencing libraries for 67 samples were prepared with the Illumina TruSeq
Nano HT DNA Library Preparation Kit (San Diego, California, USA) following the
manufacturer's protocol, except using half volumes beginning with the second addition of
Dynabeads MyOne Streptavidin C1 magnetic beads (Invitrogen, Carlsbad, CA, USA). DNA
samples were sheared using a Covaris M220 Focused-Ultrasonicator (Covaris, Woburn,
Maryland, USA) to a fragment length of ~550 bp (for an average insert size of ~420 bp). The
remaining 134 libraries were constructed by Rapid Genomics (Gainesville, Florida, USA), with
custom adapters. The Illumina i5 and i7 barcodes were used for all libraries. Target sequences
can be accessed at https://github.com/wickettlab/HybSeqFiles.
Libraries were enriched for these loci using the MyBaits protocol (ArborBiosciences
2016) with combined pools of libraries totaling 1.2 μg of DNA (12 libraries/pool at 100
ng/library). Libraries with less than 100 ng of total recovered DNA were pooled together in
equimolar concentrations using available product, resulting in some pools with less than 1.2 μg
of DNA. The smallest successful pool contained four samples with 6 ng of library each.
Hybridization was performed at 65°C for approximately 18 hours. The enriched libraries were

reamplified with 14 to 18 PCR cycles and a final cleanup was performed using a Qiagen
QiaQuick PCR cleanup kit following the manufacturer's protocol to remove bead contamination
(Qiagen, Venlo, Netherlands). DNA concentrations were measured using a Qubit 2.0
Fluorometer (Life Technologies, Carlsbad, California, USA) and molarity was measured on an
Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, California, USA). A final
cleaning step using Dynabeads MyOne Streptavidin C1 magnetic beads was performed on pools
with adapter contamination as detected on the Bioanalyzer. Pools were sequenced in four runs at
equimolar ratios (4 nM), on an Illumina MiSeq (2 x 300 cycles, v3 chemistry; Illumina, Inc., San
Diego, California, USA) at the Pritzker DNA lab (Field Museum, Chicago, IL, USA). This
produced 80,273,296 pairs of 300-bp reads. Reads were demultiplexed and adapters trimmed
automatically by Illumina Basespace (Illumina 2016). Raw reads have been deposited at the
NCBI Sequence Read Archive (BioProject PRJNA544074).
838
Quality Filtering, Assembly and Alignment
A summary of read quality from each sample was produced using FastQC
(http://www.bioinformatics.babraham.ac.uk/people.html), which revealed read-through adapter
contamination in many of the poorer quality samples. To remove read-through contamination
and filter for quality, reads were trimmed for known Illumina adapters using Trimmomatic
(Bolger et al. 2014) with the following settings:
ILLUMINACLIP:<illumina_adapters.fasta>:2:30:10 LEADING:10 TRAILING:10
SLIDINGWINDOW:4:20 MINLEN:20. Trimmed, quality-filtered reads were assembled using
the HybPiper pipeline (Johnson et al. 2016) with default settings, followed by the intronerate.py
script, to produce both exons and the "splash zone" flanking non-coding region-containing
supercontigs. Only pairs with both mates surviving trimming and quality filtering were used for
HybPiper.
To compare the influence of "splash-zone" non-coding regions, two sets of alignments
were created: (1) exons alone, and (2) coding sequences plus the "splash-zone" (Hereby referred
to as supercontigs). For multiple sequence alignments of exons alone, protein and nucleotide
sequences assembled in HybPiper were gathered into fasta files by gene. For protein sequences
only, stop codons were changed to "X" using a sed command-line regular expression to facilitate
alignment, and sequences were aligned using MAFFT with settings: --auto --adjustdirection --
maxiterate 1000 (Katoh et al. 2002). Aligned protein sequences were then used to fit unaligned
nucleotide sequences into coding frame alignments using pal2nal with default settings (Suyama
et al. 2006). In-frame, aligned DNA sequences were trimmed to remove low-coverage positions
and sequences composed only of gaps using TrimAl with the automated setting, which is
optimized for maximum likelihood analyses (Gutíerrez et al. 2009). For supercontigs, nucleotide
sequences assembled using HybPiper were gathered into fasta files by gene, gene names were
removed from fasta headers using a command-line regular expression, and sequences were
aligned in MAFFT with settings: --auto --adjustdirection --maxiterate 1000 (Katoh et al. 2002).
Reverse compliment tags ("_R_") were removed from taxon names using a command-line
regular expression, and sequences were trimmed using TrimAl with previously listed settings
optimized for maximum likelihood analyses (Gutíerrez et al. 2009). To minimize the effects of
missing data on phylogenetic analyses, accessions with < 50% of loci passing quality filtering
were removed, and genes that were recovered across < 70% of the total remaining samples were
also removed. Following quality filtering, we recovered 204 high quality loci (present in at least
70% of samples) and an average of 625,323 reads per sample (S1). All pipelines and analyses

were run on the high-performance computing cluster at the Chicago Botanic Garden unless
otherwise specified.
874
Phylogenetic Reconstruction
We conducted phylogenetic analyses using two strategies for each set of alignments.
Alignments were concatenated and analyzed using maximum likelihood (ML) in RAxML
(Stamatakis 2014; hereafter referred to as "concatenation"), whereas coalescent-based analyses
were conducted using ML gene trees in ASTRAL-II (Mirarab and Warnow 2015; Sayyari and
Mirarab 2016) and using unlinked SNPs in SVDquartets (Chifman and Kubatko 2014, 2015)
implemented in PAUP* beta version 4.0a168 (Swofford 2003; S3). In concatenation analyses,
after aligning each gene separately in MAFFT, genes were concatenated, partitioned, and
maximum likelihood trees were reconstructed in RAxML Version 8 (Stamatakis 2014) using the
GRTCAT model with 100 "rapid-boostrapping" psuedoreplicates and *Chylismia scapoidea* as
the outgroup, on the CIPRES Science Gateway computing cluster (Miller et al. 2010). For
coalescent analyses, individual gene trees were first estimated using RAxML Version 8
(Stamatakis 2014), with 100 "rapid-bootstraping" psuedo-replicates and settings: -p 12345 -x
12345 -N 100 -c 25 -f a -m GTRCAT –s, and *Chylismia scapoidea* as the outgroup. Gene trees
based on supercontigs were not partitioned by codon position. Coalescent-based analyses of
accessions were conducted in ASTRAL-II (Mirarab and Warnow 2015; Sayyari and Mirarab
2016) with default settings using the best RAxML gene trees and their associated bootstrap files
as input, and in SVD Quartets (Chifman and Kubatko 2014, 2015) with default settings using an
inframe aligned supermatrix with all 204 loci and supercontigs. ASTRAL-II and SVD Quartets
analysis was performed with 100 multi-locus bootstraps.
To assess concordance among gene trees and provide additional support complementary
to bootstrap values, we conducted two additional analyses. First, we assessed raw gene tree
concordance using Phyparts (Smith et al. 2015). Prior to running Phyparts, nodes with < 33%
support in the supercontig RAxML gene trees were collapsed using the sumtrees command in
Dendropy (Sukumaran 2010). These gene trees were then re-rooted using *Chylismia scapoidea*
as the outgroup and ASTRAL-II was rerun using these collapsed, re-rooted gene trees as the
input files. Pie charts showing gene tree discordance were generated and overlaid on the
resulting ASTRAL-II tree using the PhypartsPiecharts script
(https://github.com/mossmatters/phyloscripts/tree/master/phypartspiecharts). Phyparts piecharts
and gene tree concordance values were also added to Figure 1 by importing the two data files
produced by the Phypartspiecharts.py into R and manually matching them to key nodes on our
ASTRAL-II supercontig tree using *ggtree* version 1.14.6 (G Yu, DK Smith, H Zhu, Y Guan
2017). We also generated gene and site concordance factors for our ASTRAL-II tree constructed
using supercontigs in IQTree v1.7-beta16 (Minh et al. 2018). IQtree calculates the gene
concordance factor (gCF) and accounts for incomplete taxon coverage among gene trees and
therefore may provide a more accurate representation of agreement among gene trees than other
methods. In addition to gCF, IQTree calculates the site concordance factor (sCF), which is
defined as the percentage of decisive nucleotide sites supporting a specific node (Minh et al.
2018). We used the RAxML gene trees produced for the ASTRAL-II supercontig analysis, and
supercontig alignments themselves, as the inputs for IQtree. For computing sCF, we randomly
sampled 100 quartets around each internal node. Finally, we mapped gCF and sCF values to the
ASTRAL-II supercontig tree produced in our previous summary coalescent analysis (Fig. 2). All

phylogenetic trees, with the exception of the full Phyparts picharts tree, were visualized using the
R package *ggtree* version 1.14.6 (G Yu, DK Smith, H Zhu, Y Guan 2017).

Ancestral State Reconstruction
To infer ancestral conditions and the number of transitions in reproductive system, we
used the *phangorn* (Schliep 2011) package in R. First, a Coalescent-based species tree with
accessions grouped into taxa using a mapping file was estimated in ASTRAL-III (Zhang et al.
2018) with default settings using the best RAxML gene trees and their associated bootstrap files,
from the supercontig alignments, as input. Next we time calibrated the ASTRAL-III species tree
to 1 million years based on estimates of other taxa in the genus (Evans et al. 2009) using the
makeChronosCalib function in the *ape* (Paradis et al. 2004) package in R, and estimated an
ultrametric tree using the chronos function in *ape* (Paradis et al. 2004) with settings: lambda = 1,
model = "relaxed". Finally, we performed marginal reconstruction of ancestral character states
using the maximum likelihood method using the optim.pml and ancestral.pml functions in the
phangorn (Schliep 2011) package in R.

Testing for Hybrid Orgins with HyDe
To test for putative hybrid origins of selected taxa, we used HyDe (Blischak et al. 2018)
to calculate D-Statistics (Green et al. 2010) for a set of hypotheses (S10). Briefly, HyDe
considers a four-taxon network of an outgroup and a triplet of ingroup populations to detect
hybridization from phylogenetic invariants that arise under the coalescent model with
hybridization. Introgression between P3 and either P1 or P2 influences the relative frequencies of
ABBA and BABA, and the D-statistic measures the imbalance between these frequencies. We
tested the triplets in (S10) and set *Chylismia scapoidea* as the outgroup. We considered
hypothesis tests significant at an overall $\alpha < 0.05$ level with estimates of γ between 0 and 1. Z-
scores greater than 3 are generally interpreted as strong evidence of introgression.

Population-level Analysis
To further characterize population-level processes or genetic structure within sect.
Calylophus, we extracted and filtered SNPs by mapping individual reads against reference
supercontigs (see https://github.com/lindsawi/HybSeq-SNP-Extraction). To account for
duplicates arising from PCR during HybSeq in SNP calling and filtering, first we selected the
sample with the highest target recovery rate and sequencing depth as a target reference sequence
(Oenothera_capillifolia_berlandieri_bjc19) and gathered supercontigs for this individual into a
single target FASTA file. We then ran BWA (Li and Durbin 2009) to align sequences, Samtools
'index' (Danecek et al. 2011) and GATK CreateSequenceDictionary (Poplin et al. 2017),
respectively, on the resulting target FASTA file. Next we ran a custom script
"variant_workflow.sh" using both read files from each *Calylophus* sample as input to create a
vcf file for each sample. SNP's were called for each individual using GATK (Poplin et al. 2017)
and the vcf file from each sample as input. The resulting vcf file created in the previous step was
filtered to remove indels using GATK and the original target FASTA file as input, and then
filtered again based on read mapping and quality with GATK VariantFiltration with settings: --
filterExpression "QD < 5.0 || FS > 60.0 || MQ < 40.0 || MQRankSum < -12.5 || ReadPosRankSum
< -8.0" (Poplin et al. 2017). We generated a reduced SNP file using PLINK (Purcell et al. 2007)
to remove SNPs that did pass filter using the command: plink --vcf-filter --vcf input.vcf --const-
fid --allow-extra-chr --geno --make-bed --recode structure. Finally we used Discriminant

Analysis of Principal Components (Jombart et al. 2010) as implemented in the R package
adegenet (Jombart 2008) and the snmf function in the LEA package (Frichot and François 2015)
in R (R Core Team, 2020) to identify genetic structure between and among individuals and
populations in our data.
967
968   *Morphological Measurements and Analysis*
969   To assess taxon boundaries and patterns of morphological variation, we measured
character states for the following key morphological structures that have been used historically to
discriminate taxa in sect. *Calylophus* (Towner 1977): plant height, leaf length (distal), leaf width
(distal), leaf length/width ratio (distal), leaf length (basal), leaf width (basal), leaf length/width
(basal), sepal length, and sepal tip length. Measurements were made with digital calipers when
possible, or with a standard metric ruler and dissecting scope, from voucher specimens of nearly
all sampled populations of sect. *Calylophus* included in our molecular phylogenetic analyses.
Measurements are provided in S11. Morphological measurements were log transformed using
the R base function 'log' (R Core Team 2018) prior to Principal Components Analysis (PCA),
which was conducted in R using the *stats* package version 3.7 and the function 'prcomp'(R Core
Team 2018). All 'NA' values were omitted from analysis. Plots of PCA results were visualized
using the *ggplot2* package in R (Wickham 2016).
981

REFERENCES
Abascal F., Zardoya R., Telford M.J. 2010. TranslatorX: Multiple alignment of nucleotide
sequences guided by amino acid translations. Nucleic Acids Res. 38:7–13.
Alexander M.P. 1969. Differential Staining of Aborted and Nonaborted Pollen. Stain Technol.
44:117–122.
Alexander M.P. 1980. A Versatile Stain for Pollen Fungi, Yeast and Bacteria. Stain Technol.
55:13–18.
ArborBiosciences. 2016. In-Solution Sequence Capture for Targeted High-Throughput
Sequencing: Version 3.02. .
Blischak P.D., Chifman J., Wolfe A.D., Kubatko L.S. 2018. HyDe: A python package for
genome-scale hybridization detection. Syst. Biol. 67:821–829.
Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina
sequence data. Bioinformatics. 30:2114–2120.
Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model.
Bioinformatics. 30:3317–3324.
Chifman J., Kubatko L. 2015. Identifiability of the unrooted species tree topology under the
coalescent model with time-reversible substitution processes, site-specific rate variation,
and invariable sites. J. Theor. Biol. 374:35–47.
Danecek P., Auton A., Abecasis G., Albers C.A., Banks E., DePristo M.A., Handsaker R.E.,
Lunter G., Marth G.T., Sherry S.T., McVean G., Durbin R. 2011. The variant call format
and VCFtools. Bioinformatics. 27:2156–2158.
Doyle J.J. 1987. A Rapid DNA Isolation Procedure for Small Quantities of Fresh Leaf Tissue.
Phytochem Bull.:11–15.
Duarte J.M., Wall P.K., Edger P.P., Landherr L.L., Ma H., Pires J.C., Leebens-Mack J.,
dePamphilis C.W. 2010. Identification of shared single copy nuclear genes in *Arabidopsis*,
Populus, Vitis and *Oryza* and their phylogenetic utility across various taxonomic levels.
BMC Evol Biol. 10:61.

Evans M.E.K., Smith S. a, Flynn R.S., Donoghue M.J. 2009. Climate, niche evolution, and
diversification of the "bird-cage" evening primroses (*Oenothera*, sections *Anogra* and
Kleinia). Am. Nat. 173:225–40.

G Yu, DK Smith, H Zhu, Y Guan T.L. 2017. ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. Methods
Ecol. Evol.:28–36.

Green R., Krause J., Briggs A., Rasilla Vives M., Fortea Pérez F. 2010. A draft sequence of the
neandertal genome. Science (80-.). 328:710–722.

Gutíerrez S.C., Martínez J.M.S., Gabaldón T. 2009. TrimAl : a tool for automatic alignment
trimming. Bioinformatics.:1–2.

Illumina. 2016. BaseSpace Sequence Hub. Available from https://basespace.illumina.com/.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett
N.J. 2016. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics from High-
Throughput Sequencing Reads Using Target Enrichment. Appl. Plant Sci. 4:1600016.

Katoh K., Misawa K., Kuma K., Miyata T. 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30:3059–3066.

Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 25:1754–1760.

Miller M.A., Pfeiffer W., Schwartz T. 2010. Creating the CIPRES Science Gateway for
inference of large phylogenetic trees. 2010 Gatew. Comput. Environ. Work. GCE 2010.

Minh B.Q., Hahn M.W., Lanfear R. 2018. New methods to calculate concordance factors for
phylogenomic datasets. bioRxiv.:487801.

Mirarab S., Warnow T. 2015. ASTRAL-II: Coalescent-based species tree estimation with many
hundreds of taxa and thousands of genes. Bioinformatics. 31:i44–i52.

Paradis E., Claude J., Strimmer K. 2004. APE: Analyses of phylogenetics and evolution in R
language. Bioinformatics. 20:289–290.

Poplin R., Ruano-Rubio V., DePristo M.A., Fennell T.J., Carneiro M.O., Auwera G.A. Van der,
Kling D.E., Gauthier L.D., Levy-Moonshine A., Roazen D., Shakir K., Thibault J.,
Chandran S., Whelan C., Lek M., Gabriel S., Daly M.J., Neale B., MacArthur D.G., Banks
E. 2017. Scaling accurate genetic variant discovery to tens of thousands of samples.
bioRxiv.:201178.

Purcell S., Neale B., Todd-Brown K., Thomas L., Ferreira M.A.R., Bender D., Maller J., Sklar
P., De Bakker P.I.W., Daly M.J., Sham P.C. 2007. PLINK: A tool set for whole-genome
association and population-based linkage analyses. Am. J. Hum. Genet. 81:559–575.

R Core Team. 2018. R: A language and environment for statistical computing. .

Sayyari E., Mirarab S. 2016. Fast coalescent-based computation of local branch support from
quartet frequencies. Mol. Biol. Evol.:msw079.

Schliep K.P. 2011. phangorn: Phylogenetic analysis in R. Bioinformatics. 27:592–593.

Sharma P., Purohit S.D. 2012. An improved method of DNA isolation from polysaccharide rich
leaves of Boswellia serrata Roxb. Indian J. Biotechnol. 11:67–71.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals
conflict, concordance, and gene duplications with examples from animals and plants. BMC
Evol. Biol. 15:150.

Stamatakis A. 2014. RAxML version 8 a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30(9):1312-1313.

Sukumaran J. and M.T.H. 2010. DendroPy: A Python library for phylogenetic computing.

Bioinformatics.:1569–1571.

Suyama M., Torrents D., Bork P. 2006. PAL2NAL: Robust conversion of protein sequence
alignments into the corresponding codon alignments. Nucleic Acids Res. 34:609–612.

Towner H.F.. 1977. The Biosystematics of *Calylophus* (Onagraceae). Ann. Missouri Bot. Gard.
64:48–120.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

S4

1064
1065   Concatenation tree constructing using the exon-only dataset and 100 bootstraps. Bootstrap values
1066   indicated at relevant nodes.

1067   **S5**

1068
1069   Concatenation tree constructing using the supercontig dataset and 100 bootstraps. Bootstrap
1070   values indicated at relevant nodes.

1072
1073 ASTRAL-II summary coalescent tree constructing using the exon-only dataset and 100
1074 bootstraps. Bootstrap values indicated at relevant nodes.

1075 **S7**

1076
1077 SVD Quartets summary coalescent tree constructing using the supercontig dataset with 100
1078 bootstraps. Bootstrap values indicated at relevant nodes.

1080
1081
Phyparts piecharts ASTRAL-II tree constructing using the supercontig dataset. Piechart colors
correspond to: blue = concordant, green = top alternative bipartition, red = all other alternative
bipartitions, black = uninformative for that node.

Summary of support values for current and proposed taxonomic treatments, by analysis. 'e'
signifies trees based on exon-only data, "e+i" signifies trees based on supercontigs. '*p'* indicates
paraphyletic, unsupported taxon treatment according to tree topology.

Taxon	Concat. e	Concat. e+i	ASTRAL-II e	ASTRAL-II e+i	SVDQ e+i	Phyparts	gCF iQtree e+i	sCF iQtree e+i
a. *Oenothera hartwegii* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
b. *Oenothera hartwegii* subsp. *fendleri* (Towner 1977)	<50	*p*	81	99	*p*	1/188	1	38
c. *Oenothera hartwegii* subsp. *filifolia* (Towner 1977)	*p*	*p*	<50	<50	<50	1/190	0	36
d. *Oenothera hartwegii* subsp. *hartwegii* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
e. *Oenothera hartwegii* subsp. *macartii* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	
f. *Oenothera hartwegii* subsp. *hartwegii* + subsp. *macartii*	*p*	*p*	<50	<50	<50	0/200	0	33
g. *Oenothera hartwegii* subsp. *pubescens* (Towner 1977)	*p*	*p*	<50	<50	<50	0/200	0	36
h. *Oenothera lavandulifolia* (Towner 1977)	100	*p*	100	100	100	147/37	74	83
i. *Oenothera tubicula* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*1/188*	*p*	*p*
j. *Oenothera tubicula* subsp. *tubicula* (Towner 1977)	*p*	*p*	81	99	<50	2/171	1	38
k. *Oenothera tubicula* subsp. *strigulosa* (Towner 1977)	*p*	*p*	81	100	<50	*p*	0	37
l. *Oenothera fendleri* + *Oenothera tubicula* subsp. *tubicula*	*p*	*p*	100	100	<50	16/129	10	40
m. *Oenothera toumeyi* (Towner 1977)	100	100	100	100	100	125/65	62	56
n. *Oenothera toumeyi* + subsect. *Calylophus*	100	100	100	100	100	149/42	76	65
o. *Oenothera capillifolia* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
p. *Oenothera capillifolia* subsp. *capillifolia* (Towner 1977)	*p*	*p*	<50	*p*	<77	*p*	*p*	*p*
q. *Oenothera capillifolia* subsp. *berlandieri* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
r. *Oenothera capillifolia* subsp. *berlandieri* (New Taxon; South Texas coastal populations)	*p*	*p*	<50	100	100	*157/27*	79	86
s. *Oenothera serrulata* (Towner 1977)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
t. *Oenothera 'australis'* (New Taxon; South Texas *O. serrulata*)	*p*	100	*p*	*p*	*p*	10/175	*p*	*p*
u. *Oenothera gayleana* (Turner & Moore 2014)	*p*	*p*	*p*	*p*	*p*	*p*	*p*	*p*
v. *Oenothera gayleana* (Revised Taxon)	*p*	*p*	*p*	<50	<50	0/193	0	35

1089 **S10**
1090 List of admixture hypotheses tested, Zscore, P-value and Gamma results from HyDe analysis.
1091 Each row represents a triplet set that was tested consisting of a putative hybrid individual and
1092 two parent groups; Parent 1 and Parent 2.
1093

Putative hybrid individuals	Parent 1 Group	Parent 2 Group	Zscore	P	Gamma
Ohar_X_fil_MJM629.6	CoreOharfil	CoreOharpub	-99999.9	1	-0.049
Ohar_X_fil_MJM628	CoreOharfil	CoreOharpub	-99999.9	1	-0.057
Ohar_X_fil_MJM626	CoreOharfil	CoreOharpub	-99999.9	1	-0.001
Ohar_X_fil_MJM793	CoreOharfil	CoreOharpub	-99999.9	1	-0.057
Ohar_X_fil_MJM794	CoreOharfil	CoreOharpub	-99999.9	1	-0.034
Ohar_X_fil_BJC81	CoreOharfil	CoreOharpub	-99999.9	1	-0.042
Oharhar_MJM1581	CoreOharpub	Oharhar/harmac	-99999.9	1	1.706
Ohar_X_pub_BJC29	CoreOharpub	Oharhar/harmac	2.378	0.009**	0.338
Oharpub_BJC30	CoreOharpub	Oharhar/harmac	-99999.9	1	-0.215
Oharpub_MJM2158.C	CoreOharpub	Oharhar/harmac	1.113	0.133	0.229
Ohar_X_pub_MJM1134.2	CoreOharpub	Oharhar/harmac	0.043	0.483	0.015
Ohar_X_pub_MJM2405.A	CoreOharpub	Oharhar/harmac	0.882	0.189	0.158
Ohar_X_pub_MJM594.4	CoreOharpub	Oharhar/harmac	2.094	0.018*	0.332
Ohar_X_pub_MJM615	CoreOharpub	Oharhar/harmac	0.413	0.340	0.096
Oharmac_MJM997.1	CoreOharpub	Oharhar/harmac	1.563	0.059	0.224
Ohar_X_har_MJM613	CoreOharpub	Oharhar/harmac	-99999.9	1	-0.495
Otub_X_str_MJM1916.E	Otubtub	Oharhar/harmac	5.585	0.000***	0.947
Oharfen_MJM2192.A	CoreOharfen	CoreOharfil	0.305	0.380	0.001
Oharfen_MJM781.3	CoreOharfen	CoreOharpub	0.025	0.490	9.69E-05
Ocapcap_MJM1015.2	CoreOcapcap	S. TX Ocapber	0.467	0.320	0.003
Ocap_X_ber_MJM767.3	central OK Ocapber	Pecos River/Southern Plains	-99999.9	1	-0.003

- Indicates value meets significance threshold of $P < 0.05$
** Indicates value meets significance threshold of $P < 0.001$
*** Indicates value meets significance threshold of $P < 0.0001$

1094

1095 **S11**
1096 See excel table "S11 Morphometric Data"
1097

1098 **S12**
1099 See excel table "S12 Pollen Counts"
1100

1101 **S13**
1102 DNA EXTRACTION PROTOCOL – see additional file

1104
1105 Ancestral State Reconstruction of reproductive system in sect. *Calylophus* using supercontigs
1106 and accessions grouped into taxa. Pie-charts on notes represent likelihood of ancestral
1107 reproductive system at each node (teal = hawkmoth pollination, purple = bee pollination, yellow
1108 = PTH).