**Comprehensive survey of microsporidia reveals extensive ecological and phenotypic diversity**

Brandon M. Murareanu[1*], Ronesh Sukhdeo[1*], Rui Qu[1*], Jason Jiang[1*], and Aaron W. Reinke[1#]

[1] Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada

*These authors contributed equally

#Corresponding author

aaron.reinke@utoronto.ca

## Abstract

Microsporidia are a large group of fungal-related obligate intracellular parasites. Though many microsporidia

species have been identified over the past 160 years, there is a lacking depiction of the full diversity of this

phylum. To systematically describe the characteristics of these parasites, we created a database of 1,428

species and their attributes, including the hosts they infect and spore characteristics. We find that microsporidia

have been reported to infect 15 metazoan and four protozoan phyla, with smaller phyla being underrepresented.

Most species are only reported to infect a single host, but those that are generalists are also more likely to infect

a broader set of host tissues. Strikingly, polar tubes are 3-fold longer in species that infect tissues besides the

intestine, suggesting that polar tube length is a determinant of tissue specificity. Phylogenetic analysis revealed

four clades which each contain microsporidia infecting hosts from all major habitats. Although related species

are more likely to infect similar hosts, we observe examples of changes in host specificity and convergent

evolution. Taken together, our results show that microsporidia display vast diversity in their morphology and the

hosts they infect, illustrating the flexibility of these parasites to evolve new traits.

## Introduction

24 Microsporidia are a large phylum of obligate intracellular parasites. Although they were once thought to be

25 protists, it is now clear they belong to a lineage of early diverging fungi[1]. Microsporidia was the first parasite to

26 have its genome sequenced, revealing the smallest eukaryotic genome with as few as ~2000 protein coding

genes[2,3]. Genome reduction results from the loss of many metabolic and regulatory genes, reduced non-coding DNA, and reduced gene sizes that encode for proteins smaller than their yeast orthlogs[4–7]. Microsporidia have also lost mitochondria and instead depend on reduced organelles known as mitosomes[8]. As a result, microsporidia rely heavily on their hosts for nutrients and have evolved many novel proteins for interacting with host cells[9,10]. Paradoxically, although microsporidia are amongst the simplest eukaryotes, there are at least 1500 known species that as a phylum can infect many different animals and most major tissues[11,12].

Microsporidia cause death and disease and are a threat to both animal and human health. They were first discovered in 1857 in silkworms and microsporidia infections have triggered historical collapses of the sericulture industry[13]. Microsporidia infections in animals are very common. For example, 50% of honey bee colonies and 60% of field-collected mosquitos have been reported to be infected with microsporidia[14,15]. These parasites are an emerging threat, with the most common species that infect honey bee colonies and shrimp farms only being discovered within the last two decades[16,17]. Microsporidia also cause problems in laboratory research, commonly infecting zebrafish facilities and causing fruit fly colonies to collapse[18,19]. Infections often result in delayed animal growth, reduction in offspring, and premature mortality[20–23]. Over a dozen species of microsporidia can infect humans and several of these species are also found in a large variety of different mammals and birds, leading to concern over zoonotic transfer[24–26]. In addition to the threats posed by microsporidia, they also have potential benefits, including being used as biocontrol agents against insect pests or preventing the spread of mosquito-vectored diseases[27,28].

All microsporidia produce environmentally resistant spores. These spores contain both the infectious sporoplasm and the unique polar tube infectious apparatus[29]. In the presence of a host, microsporidia fire their polar tube which is thought to either pierce the host cell or fuse with the host plasma membrane to form an invasion synapse[30]. Each species has spores of a characteristic size, but some species have multiple classes of spores[31,32]. While many infections are thought to result from the oral consumption of spores, other types of infections e.g. ocular, can also occur[26].

Although many microsporidia species have been characterized, their properties have not been examined systematically. Thus, the true diversity of these medically and agriculturally important parasites is unknown.

Here, we curate a database of microsporidia attributes for 1428 species. Using this data, we describe the history and geography of microsporidia discovery. We show that microsporidia can infect a diverse range of host species from all major habitats. We find that microsporidia that infect more hosts tend to infect more tissues. We observe large ranges for spore size, which correlates with the number of nuclei and polar tube length. We also find that the length of the polar tube is longer in species that infect other tissues besides the intestine. Phylogenetic analysis shows that microsporidia group into clades with predominant environmental habitats, but the ability to infect in other environments within a clade is also common. Related species mostly infect similar hosts, but dramatic changes in host specificity and convergent evolution also occur. Together, this work describes the diverse properties of microsporidia species and provides a resource for further understanding these enigmatic parasites.

**Materials and methods**

**Construction of microsporidia species properties database**

Each microsporidia species described from Victor Sprague's 1977 book was manually entered into an Excel table[33]. Each row represented a microsporidium, and columns were created for each species attribute recorded. Parenthesized text was used to denote which spore class or host an attribute corresponded to. The spore conditions (ex: fixed, fresh, etc.) under which morphological attributes were recorded were also denoted by parenthesized text. Species without a specific name (ex: Nosema sp.) were numbered to give each entry a unique name.

To identify additional species, we searched the journal databases Scopus, Web Of Science and JSTOR for papers published between 1977 and April 2020. The search terms ("microsporidia" OR "microsporidium" OR "microsporida") were used for all searches. Papers describing new microsporidia species or further describing existing species were used. Moreover, we looked at citing articles for papers describing new species, to find papers further describing those species. Non-English papers were translated into English using Google Translate. An additional 134 species were identified from a database of compiled by Kirk[34]. Papers describing these additional species were found by searching in the aforementioned databases, or with Google.

Microsporidia attributes for species described in journal articles were recorded as described above. For a subset

of papers where the full article could not be accessed, information was recorded from the abstract.

**Analysis of microsporidia discovery date and geographical location**

For each species, the earliest year from the "Date Identified" column of Table S1 was extracted. Species without

identification year data were excluded from this analysis. Data of the countries where each species had been

reported was extracted from the "Locality" column of the database. A species found in multiple countries was

counted for all of those countries. With the exception of oceans, species found in bodies of water were counted

as the country located closest to it. If a species was reported either in a country that no longer exists or a general

region, it was counted as the largest country that occupied its lands. For example, species found in the U.S.S.R.

were counted as species found in Russia. Species reported from each continent were determined according to

the categorization of worldpopulationreview.com.

**Determination of host taxonomy and environment**

The taxonomy for each host species was determined using Catalogue of Life

(http://www.catalogueoflife.org/annual-checklist/2019/), GBIF (https://www.gbif.org/), WoRMS

(http://www.marinespecies.org/), NCBI Taxonomy Browser

(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi) or Wikipedia (https://en.wikipedia.org/). Using

this approach, we annotated the taxonomy for 1422 out of 1449 host species. Host environment was determined

using Catalogue of Life, GBIF and WoRMS. Host environment data was also extracted from other online

databases, papers describing the host, or the original papers describing infection of the host by microsporidia

(Table S2). To classify moths, all non-Pyralidae moths were classified as "terrestrial". Pyralidae moths were

classified as freshwater and terrestrial, since they have aquatic larval stages[35]. If environment data was not

available at the species level, we used environment data from the closest taxonomic level, the highest being

order. Some species without environment data in the sources mentioned above, were assigned environments

by common knowledge, i.e., parakeets inhabiting a terrestrial environment. The environment of 1338 out of 1449

species was annotated. Experimental hosts for microsporidia were excluded from analysis, as they may not

reflect the natural host range of a microsporidia species[36].

**Categorizing and counting tissues infected**

All unique infected-tissue names were extracted from the "Site" column of Table S1. These names were sorted into 23 broader tissue categories (Table S3). Each unique tissue name was restricted to a single tissue category. Tissues that could not be clearly assigned to a category were classified as "Ambiguous". Infections of protist species were classified as "Protist". To count the number of tissues infected by each microsporidium, we counted how many tissues were listed for each microsporidium. Microsporidia species with systemic or ambiguous infections were excluded, as tissues in these categories could not be associated with a single tissue. The overall infection frequencies within tissue categories were determined using a custom R script. Each species was counted as infecting a tissue category one or zero times. Ambiguous and protist infections were not counted.

**Comparing infected hosts to infected tissues**

To compare the number of hosts infected to the number of tissues infected, microsporidia species were grouped based on the number of hosts they infect. The proportion of species infecting one tissue was calculated within each host bin. Species with systemic infections were counted as infecting multiple tissues. Experimental hosts were excluded from analysis.

**Determining microsporidia transmission mode**

Microsporidia's mode of transmission was either determined from literature transmission data or inferred from the tissue infected, as has been done previously[37]. ("Site" and "Transmission (described in literature)" columns in Table S1). Species without literature transmission data were classified as vertically transmitted if infecting germline, horizontally transmitted if infecting somatic tissue, or both. Transmission mode was not determined for species with systemic infections, as it is unclear what tissues they infect. When available, literature transmission data supplemented transmission mode determination. Species only infecting somatic tissues but with vertical transmission data from literature were classified as both. Species only infecting germline with horizontal transmission from literature were also classified as both. For species without tissue infection data, mode of transmission was determined solely through literature transmission data (horizontal, vertical, or both). When comparing rates of horizontal and vertical transmission across environments, transmission modes were

described for individual microsporidia host species. Only hosts in single environments were considered. Host

species that could not be matched to specific tissues or modes of transmission were excluded from analysis.

**Spore shape classification**

Spore shapes were extracted from the "Spore Shape" column in Table S1. These shapes were grouped and

placed into 20 shape categories (Table S4). Shapes that could not be categorized as a defined grouped shape

were labelled "ambiguous" and excluded from analysis. If a given spore was documented to be oval in addition

to another shape, the latter shape classification was kept and used for analysis. However, if a spore was

documented to be oval in addition to at least two other shapes, it was excluded from analysis.

**Analysis of spore dimensions**

Spore lengths and widths were extracted from "Spore Length Average" and "Spore Width Average" columns in

Table S1. If a given species had multiple measurements for a spore class, the "fresh" measurement was used

and the remaining "fixed" measurements were excluded. If a spore class only had "fixed" measurements

available, they were included in the analysis.

**Calculating spore volume**

Spore volume was estimated for all types of documented spores using the formula for an ovoid, as this shape

accounts for the majority of spores (Fig. 4A). The following formula was used: Volume= 4/3πabc, where a, b,

and c are the 3 semi-axes of the ovoid.  Width/2 was used for both a and b as width and depth of spores have

been shown to be similar[38]. Length/2 was used for c. When possible, priority was given to the "fresh"

measurements of documented spore for calculating volume.

**Calculating polar tube length**

Polar tube length was calculated by using the spore width and number of observed polar filament coils

documented. Fresh width values were prioritized for calculations. The following formula was used:

$Calculated\ polar\ tubule\ length = \pi \times \text{width of spore} \times \text{number of polar filament coils.}$

**Microsporidia tissue specificity**

To assess if microsporidia polar tube length affects which tissues are infected, hosts infected by multiple

microsporidia species were identified. Calculated polar tube lengths were organized into "Intestinal" infection if

the species only infected the gut/digestive system as determined in Table S3. If a species infecting the same

host infected non-intestinal tissues, it was classified as "Other Tissues". Every pair of microsporidia species

between these two classes was compared for every host (Table S5).

**Comparing polar tube length to volume**

To compare polar tube length to spore volume, only species with experimental polar tube values were

considered. Species with multiple spore classes were represented multiple times. Entries where spore class

could not be clearly matched to volume, polar tube length, or nucleus data were excluded. Least squares

regression was performed for all spores (Fig. S4), as well as uninucleate and binucleate spores individually to

control for the confounding effects of nuclei number. Correlation was calculated with Spearman $r_s$ to limit the

effect of outliers on correlation values.

**18S phylogenetic analysis of microsporidia species**

Using a modified version of the *read.Genbank* function from the *ape* package for R, microsporidia 18S sequences

and    associated    data    were    collected    from    the    NCBI    GenBank    nucleotide    database:

*https://www.ncbi.nlm.nih.gov/genbank/*. The search terms *("Microsporidia"[Organism] AND (rRNA[Title] OR*

*ribosomal[Title])) AND (small[Title] OR SSU[Title] OR 18s[Title] OR 16s[Title])* were used to collect 7830 unique

18S sequences representing 1107 unique microsporidia species identifiers. The longest sequence for each

unique species identifier was extracted from this collection for subsequent analysis. Only the sequences with

species identifiers corresponding to species in our database (Table S1) were included in subsequent steps. This

left us with 18S sequences for 273 unique microsporidia species. We also included a single 18S sequence for

*Rozella allomycis*, a commonly used outgroup in microsporidia phylogenies[39]. A list of accession numbers for all

274 18S sequences and corresponding species can be found in Table S6.

The 274 18S sequences were aligned with MAFFT, using the default parameters. The alignment was masked

using the *MaskAlignment* function from the *DECIPHER* package for R. Bayesian phylogenetic analysis of the

masked alignment was performed with MrBayes v 3.2.7, using two MCMC runs, each with one cold and three

heated chains, for 10 million generations, sampling every 1000 generations, with the first 2.5 million generations being discarded as burn-in. For this inference, we used the GTR model of substitution, with an *"invgamma"* model for rate variation. Maximum Likelihood phylogenetic analysis of the masked alignment was performed with RAxML-HPC2 using the default parameters and 1000 bootstrap iterations. 50% majority rule consensus trees were generated for both Bayesian and Maximum Likelihood analyses and were annotated using the *ggtree* and *treeio* packages for R. Clade groupings from both trees were compared by computing the fraction of species placed in the same clade in both trees. A complete list of parameters for the MrBayes and RAxML phylogenetic analyses can be found in Table S7. MAFFT, MrBayes and RAxML were all run on XSEDE accessed via the CIPRES Science Gateway: *https://www.phylo.org/*[40].

**Analysis of microsporidia evolutionary relationships**

All analysis of evolutionary relationships between the four major clades was done in R. The distribution of host environments was determined by counting the number of species in each clade that are found in each environment, expressed as a percentage of the total number of species in each clade. The distributions of spore volume and polar tube length were determined using the calculated spore volume and calculated polar tube length data (Table S1). In cases where species had multiple spore classes, all were included as separate data points in the analysis. To determine if any hosts were infected by multiple microsporidia species from different clades, the assortment of hosts for each clade were compared with each other to see if any hosts appeared in multiple clades.

Using the *dist.alignment* function from the *seqinr* package for R, a pairwise sequence similarity matrix for the 37,128 unique species pairs encompassing all 273 microsporidia species from the phylogenetic analysis was computed based on the unmasked MAFFT alignment. The fractions of species pairs within a given sequence similarity range that had at least one shared host taxonym, environment, and infected tissue were computed separately and displayed as distributions with 20 sequence similarity bins. The percentage similarities in spore volume and polar tube length in each species pair were computed separately according to the formula *(larger value / smaller value)\*100,* using the calculated spore volume and calculated polar tube length data (Table S1). In cases where species had multiple spore classes, values for each class were averaged prior to the percentage

similarity computation. The average percentage similarities in spore volume as well as polar tube length within

a given sequence similarity range were then computed and displayed as distributions with 20 sequence similarity

bins. For each distribution, species pairs that had at least one species with missing data for the given comparison

criterion were excluded.

**Data and code availability**

All microsporidia species and host data are available in Tables S1 and S2. All data analysis was performed using

Microsoft Excel and R version 3.6.1 or later accessed via RStudio version 1.2.5019 or later[41]. All code used for

data analysis can be found on the Microsporidia-Database GitHub repository

*https://github.com/bmurareanu/Microsporidia-Database*.

**Results**

**Generating a database of microsporidia species' characteristics**

To determine the reported diversity of microsporidia species, we comprehensively examined the literature. We

began with Victor Sprague's 1977 book, the most complete compendium of microsporidia published thus far,

which contains information on 692 species[33]. We then identified 684 journal articles containing reports of

identified microsporidia species. We recorded the described attributes of each species including the date and

location of discovery, host species, infected tissues, spore size, and polar tube length. Overall, we generated a

database containing information on 1,428 microsporidia species (Table S1).

We first examined the history of microsporidia species discovery. The original article describing the first species

of microsporidia was published in 1857[42]. Since 1881, on average ~100 novel species have been reported per

decade. The number of discovered species peaked in the 1970s with 246 species being reported. Although in

recent years fewer species have been reported than at this peak, during the last decade, on average 13 novel

species have been described every year (Fig. 1A).

We next determined the geographical locations from which microsporidia have been isolated. Although the

majority of described species have been found in Europe and North America, microsporidia have been found in

all continents except Antarctica (Fig. 1B). They have also been reported in 11 oceans or seas, including a species

found in the Weddell sea off the coast of Antarctica, and from methane seeps on the floor of the pacific ocean[43,44].

While the United States has reported the most species of microsporidia (283), 123 species have also been found

in the much smaller Czech Republic (Fig. 1C).

**Microsporidia infect a diversity of hosts and tissues**

Microsporidia have often been described as infecting nearly all animal taxa, but a full description of which host

phyla these parasites infect is lacking[26,45,46]. Using several databases describing animal and protozoan diversity,

we determined the taxonomical placement of 1449 microsporidia hosts (Table S2). We find that infections have

been reported in 15 metazoan and four protozoan phyla. The majority of the hosts in five of these phyla are

themselves parasites, demonstrating that hyperparasitism is a widespread microsporidian trait (Fig. 2A). There

are 17 metazoan phyla that have not been reported to be infected by microsporidia. These phyla have on average

32-fold fewer known species than phyla that have been reported to be infected with microsporidia. This suggests

that the absence of known microsporidia infections in these phyla may be due to a lack of sampling from these

hosts (Fig. 2B). We also observed a diversity of hosts within a phylum; the two phyla with the most hosts,

Chordates and Arthropods, contain hosts in 39 out of 155 orders and 42 out of 162 orders, respectively (Fig. 2C

and S1).

Because microsporidia are obligate intracellular parasites, the habitat of the host informs that of the microsporidia

species[12]. We determined the ecological environments of the different hosts in our database. Of the four major

habitat types, 55.1% of hosts are terrestrial, 34.0% of hosts are freshwater, 23.1% of hosts are marine, and

12.6% of hosts are brackish. Interestingly, 25.7% of hosts are classified as living in more than one environment.

This is exemplified by seven hosts, such as mosquitos (*Anopheles pseudopunctipennis)* and parasitic flatworms

(*Heterophyes heterophyes*), which inhabit all four environment types (Fig. 2D and Table S2).

Microsporidia are known to infect a range of different tissues within their hosts[12]. To investigate the extent of this

diversity, we classified infections into 23 tissue types (Table S3). The tissues infected by microsporidia

encompass all embryonic germ layers and most major organ systems[47]. The adipose tissue, intestine, muscles,

germline, and excretory system are the most commonly infected tissues. Infections have also been reported in

specialized tissues such as the gills of fish and oenocytes, which are lipid-processing cells in insects[48] (Fig. 3A).

Most microsporidia infections appear to be restricted to a single tissue, but ~11% of species can infect more than

four tissues (Fig. 3B).

Microsporidia can be specialists that infect only one or a few hosts, or generalists that can infect many hosts[5,45].

Over 80% of reported species (with host data) only infect a single host, but 3% can infect five or more hosts (Fig.

3C). To determine if there is a relationship between host range and tissue specificity, we examined if species

that infect more hosts also infect more tissues. Indeed, 34% of species that infect one host infect multiple tissues,

compared to 55% of species that infect four or more hosts infecting multiple tissues (Fig. 3D). This correlation

suggests that species with larger host ranges tend to infect a broader set of tissues.

**Diversity of microsporidia spore morphology**

All microsporidia have an extracellular spore form and the morphological properties of these spores can be used

to distinguish species[31]. To examine the diversity of microsporidia spore morphology, we first classified spores

into 20 unique shape descriptors (Table S4). Over 93% of classified spores belong to just six shape classes (Fig.

4A). Examination of spore size revealed a large diversity in both spore length (1 – 25 μm) and width (0.15 – 10

μm). The spores fall into three main groups with the average length/width ratio of spherical spores being 1, rod-

shaped spores being ~3.8 and together, oval, ovocylindrical, pyriform, and elliptical being ~1.9 (Fig. 4B and S2).

Using length and width measurements of spores, we estimated spore volumes, which range from

$3.5 \times 10^{-2} \ \mu m^3 - 1.3 \times 10^3 \ \mu m^3$.

Microsporidia spores contain polar tubes that fire in the presence of a host signal and are used to infect host

cells[30]. The length of the polar tube can be measured by artificially triggering spore firing by treating with different

chemicals or mechanical pressure[30,49]. The polar tube is coiled inside of the spore and the number of coils can

be determined using electron microscopy[29]. Using the spore diameter and the number of polar tube coils, we

estimated the length of the unfired polar tube. We observe a large range in both experimentally measured (2.5

μm – 420 μm) and estimated (1.5 μm – 752.1 μm) polar tube lengths. We compared estimated polar tube length

to calculated polar tube length and observed a significant correlation (Fig. 4C). In contrast to previous studies

which have reported that the length of fired tubes is longer than the estimated length of unfired tubes, we see

many examples of unfired polar tubes being both shorter and longer than those that are fired[38,50].

Microsporidia spores need to be able to accommodate both the polar tube and the sporoplasm. Sporoplasm size

measurements do not exist for most species, so we used the number of nuclei (either one or two) as a proxy for

the size of the sporoplasm. We examined the relationship between spore size and nuclei number and observed

that spores with two nuclei are on average ~22% larger than those with one nucleus (Fig. 4D). We next examined

the relationship between polar tube length and spore size. We observed that experimentally measured polar

tube lengths were positively and significantly corelated with spore volume (Fig. 4E and S3). Taken together,

these results suggest that although the number of nuclei and the length of the polar tube don't explain all of the

variation in spore volume, the size of intracellular contents is correlated with spore size.

One possible explanation for why polar tube length varies so widely, is that the polar tube specifies which tissues

are infected. This idea was proposed by Luallen and colleagues studying microsporidia infection in the nematode

*Caenorhabditis elegans*[6]. The authors observed that *Nematocida parisii*, which infects the intestine, has a polar

tube length of 4.03 μm, in contrast to that of *Nematocida displodere*, which infects the muscle and epidermis

from the intestinal lumen, and has a polar tube length of 12.55 μm. To determine if polar tube length correlates

with infected tissue type, we examined 11 hosts that were infected by at least one microsporidia species that

exclusively infects the digestive system and at least one species that infects other tissues. In 82% of

microsporidia species pairs, the polar tube was longer in the microsporidia species that infected other tissues

besides the digestive system. On average, species that infect other tissues were 3-fold longer than those that

were restricted to the digestive system. This suggests that one function of polar tube length is determining host-

tissue specificity (Fig. 4F and Table S5).

Microsporidia can either be transmitted horizontality, from host to host, or vertically, from parent to offspring[20,37].

We examined whether species were reported to infect the germline (vertical) or somatic tissue (horizontal) and

observed that 18% of species are potentially vertically transmitted (Fig. S4A). We also observed that

transmission mode is not dependent on host environment (Fig. S4B). Furthermore, spore volume and polar tube

length are not correlated with transmission mode (Fig. S4C-D).

**Evolutionary properties of microsporidia**

Microsporidia have previously been reported to group into five clades, classified based on host environmental

habitat[51]. Using our large collection of host and species information, we sought to revisit these groupings. We

performed Bayesian phylogenetic analysis on 273 18S rRNA sequences belonging to microsporidia species for

which we also had attribute data. The resulting tree recovers four clades (I,III,IV, and V) and is largely in

agreement with recently published microsporidia phylogenetic trees[52,53] (Fig. 5A and S5). We also assembled a

Maximum Likelihood phylogenetic tree using the same sequences, and the majority of species (~95%) were

placed in the same clade in both trees, further supporting our clade groupings (see methods). All subsequent

analysis was done using the Bayesian tree. We then determined the percentage of microsporidia species in each

clade that inhabit each of the 4 major environments. We mostly observed the previously determined

environmental groupings, with clade IV being terrestrial and clade III being marine. We also observed that clades

I and V have a higher percentage of microsporidia with freshwater hosts than the other clades, as has been

previously shown. However, clades I and V have a higher percentage of microsporidia with terrestrial than

freshwater hosts (Fig. 5B). Although there is a predominant habitat in each clade, between 26-44% of

microsporidia species in each clade do not infect a host in the predominant habitat. This suggests that evolution

of microsporidia within clades is not strictly restricted to hosts from a single environmental habitat.

We next examined the diversity of spore and polar tube length amongst clades (Fig. 5C and D). We observe that

the clades with the highest percentage of microsporidia infecting freshwater hosts (I and V) have on average

larger spores and more variation in spore size than the other two clades (Fig. 5C). This same trend exists in

polar tube length, but to a lesser extent (Fig. 5D).

Related species of microsporidia are known to infect very different hosts, but it is unclear how common of an

occurrence this is[7]. To determine the relationship between microsporidia species similarity and host similarity,

we first calculated the pairwise distances between 273 microsporidia 18S rRNA sequences (Fig. 6A). We then

compared how often related species infect the same host family, order, class, or phylum. We observed that while

42% of microsporidia species pairs with greater than 65% 18S similarity infect hosts from the same family, this was true for only 2% of pairs with less than 65% 18S similarity. These strong associations between microsporidia and host relatedness were also observed at the other taxonomic ranks (Fig. 6B). Although very closely related microsporidia almost always infect the same phylum, we observed several examples of species with greater than 90% similarity infecting different phyla, including between species that infect insects and the known human infecting species *Vittaforma corneae* and *Trachipleistophora hominis*[26]. To identify examples of convergent evolution, we determined how often a host species is infected by microsporidia from more than one clade and found 21 such cases (Fig. 5E). Taken together, our results indicate that microsporidia mostly evolve to infect similar hosts but can undergo dramatic hosts shifts.

Finally, we determined the relationship between microsporidia sequence relatedness and infected tissues, habitat, and spore similarity. We find that closely related species mostly infect the same tissues and share the same environmental habitats (Fig. 6C-D). Spore size and polar tube length are also more similar in closely related species, but to a lesser extent than that of host similarity (Fig. 6E-F). Even closely related species only share on average less than 75% similarity between the tissues they infect and their spore properties. These results suggest that although related species are likely to have shared characteristics, microsporidia are flexible in their ability to evolve new traits.

**Conclusions**

Here we report the largest systemic collection of microsporidia species characteristics assembled since Sprague in 1977[33]. Although the data we present is comprehensive, there are several limitations. First, we only curated named species that were reported with attribute data. There are at least 93 identified microsporidia species that we could not find information on and therefore did not include in our analysis[34]. Second, many microsporidia species names have changed throughout the years and this likely led to missing information[54]. Third, we do not include complete attribute data for every species as authors do not always report on the same properties. Additionally, some characteristics such as spore size were determined through multiple techniques with differing accuracy. Finally, for species that are the subject of many publications, we are unlikely to have captured all of the known data including the full breadth of hosts these species infect.

About 1500 microsporidia species have been identified, which is likely only a small fraction of the true number of species[55]. We find that identification of novel microsporidia species peaked in the 1970s, which is consistent with a previous report showing identification of microsporidia has lagged behind other parasites[56]. Sampling for microsporidia species has been biased with ~65% of the species identified so far being from Europe and North America. Africa, South America, and Oceania have been shown to have large levels of parasite-host diversity, suggesting that these parts of the world could contain many undiscovered microsporidia species[57].

Although the pace of microsporidia discovery has slowed, many new and diverse species are still being discovered. Since the 1990s, microsporidia were reported for the first time in six phyla (Cephalorhyncha, Porifera, Amoebozoa, Dicyemida, Gastrotricha, and Phoronida), suggesting that most types of animals may be infected by microsporidia. In support of this, although there are no named microsporidia species infecting tardigrades, animals in this phylum has been observed to be infected by microsporidia[58]. Environmental sampling has also revealed the existence of many species of unnamed microsporidia, especially a diversity of short-branched microsporidia[52,59,60]. Echinoderms comprise the largest phylum without any reported microsporidia infections and represent an interesting target for future efforts to uncover novel microsporidia diversity.

The molecular mechanisms that determine host and tissue specificity in microsporidia are mostly unknown. Here we show that microsporidia polar tube length is longer in species that infect tissues other than the intestine. In bilaterian animals, the intestine is in the middle of the animal and access to most other tissues requires going through the intestine[61]. This has been shown in *C. elegans*, where infection of the epidermis is dependent upon oral consumption of spores[6]. Microsporidia have been shown to be more selective for which types of cells they infect inside of animals than in cell culture, suggesting that access to tissues could determine specificity[62,63]. We also show that there is a tendency for microsporidia species that infect more hosts to also infect more tissues. This is likely even more common than our data indicates, as it only takes a single infected animal to determine tissue distribution, but many identifications to determine host range[31]. Our data suggest that the evolutionary forces that drive species to be generalists also tends to generate species with broad tissue specificity.

In 2005 Vossbrinck and Debrunner-Vossbrinck proposed that microsporidia compose five major clades that group according to the environmental habitat of each species host[51]. Although these environmental groupings

provide an attractive explanation of microsporidia evolution, none of the clades only contained hosts from a

single environment[64]. Clade II was not well supported in this original phylogeny and has not been consistently

recovered in more recent phylogenetic trees[20,31,52,59]. The position of *Nematocida* in these phylogenies has not

been constant and often differs from the basal position recovered from trees based on whole genome

sequencing[53,65]. Environmental sampling from diverse environments recovered microsporidia from many

different clades, demonstrating that there is not a strict correspondence between clades and host environment.

For example, many samples from marine environments were found in clade IV (Terresporidia). Intriguingly,

several OTUs (operational taxonomy unit) were found in multiple environments, providing support for some

species having multiple hosts from different environments or the same host being infected in multiple

environments[52]. Our analysis suggests that although clades do belong predominantly to a single environment,

microsporidia can evolve to infect hosts in other habitats. A host often exists in more than one habitat and some

microsporidia have multiple hosts that exist in distinct habitats. Because of this, many species of microsporidia

are found in multiple environments, likely aiding their ability to evolve to infect new hosts.

In conclusion, we have assembled a collection of microsporidia attributes for 1,428 species. This data will be a

useful comparative resource which will support further species identification. In addition to the questions that we

have examined, this database will aid in addressing other systematic inquires about microsporidia diversity.

**Acknowledgements**

We thank Hala Tamim El Jarkass, Alexandra R. Willis, JiHae Jeon, and Robert J. Luallen for providing helpful

comments on the manuscript. This work was supported by the Natural Sciences and Engineering Research

Council of Canada (Grant #522691522691 and Undergraduate Student Research Awards to B.M., R.S., R.Q.,

and J.J) and an Alfred P. Sloan Research Fellowship FG2019-12040 (to A.W.R.).

**Competing Interests:** The authors declare that they have no competing interests.

**References**

1.    Keeling, P. Five Questions about Microsporidia. *PLOS Pathogens* **5**, e1000489 (2009).

2.    Katinka, M. D. *et al.* Genome sequence and gene compaction of the eukaryote parasite Encephalitozoon cuniculi.

*Nature* **414**, 450–453 (2001).

3.    Corradi, N., Pombert, J.-F., Farinelli, L., Didier, E. S. & Keeling, P. J. The complete sequence of the smallest known

nuclear genome from the microsporidian Encephalitozoon intestinalis. *Nat Commun* **1**, 77 (2010).

4.    Nakjang, S. *et al.* Reduction and expansion in microsporidian genome evolution: new insights from comparative

genomics. *Genome Biol Evol* **5**, 2285–2303 (2013).

5.    Desjardins, C. A. *et al.* Contrasting host–pathogen interactions and genome evolution in two generalist and specialist

microsporidian pathogens of mosquitoes. *Nature Communications* **6**, 1–12 (2015).

6.    Luallen, R. J. *et al.* Discovery of a Natural Microsporidian Pathogen with a Broad Tissue Tropism in Caenorhabditis

elegans. *PLoS Pathog* **12**, e1005724 (2016).

7.    Wadi, L. & Reinke, A. W. Evolution of microsporidia: An extremely successful group of eukaryotic intracellular

parasites. *PLoS Pathog* **16**, e1008276 (2020).

8.    Williams, B. A. P., Hirt, R. P., Lucocq, J. M. & Embley, T. M. A mitochondrial remnant in the microsporidian

Trachipleistophora hominis. *Nature* **418**, 865–869 (2002).

9.    Reinke, A. W., Balla, K. M., Bennett, E. J. & Troemel, E. R. Identification of microsporidia host-exposed proteins

reveals a repertoire of rapidly evolving proteins. *Nat Commun* **8**, 14023 (2017).

10.  Franchet, A., Niehus, S., Caravello, G. & Ferrandon, D. Phosphatidic acid as a limiting host metabolite for the

proliferation of the microsporidium Tubulinosema ratisbonensis in Drosophila flies. *Nature Microbiology* **4**, 645–655

(2019).

11.  Vávra, J. & Lukeš, J. Chapter Four - Microsporidia and 'The Art of Living Together'. in *Advances in Parasitology* (ed.

Rollinson, D.) vol. 82 253–319 (Academic Press, 2013).

12.  Stentiford, G. D., Feist, S. W., Stone, D. M., Bateman, K. S. & Dunn, A. M. Microsporidia: diverse, dynamic, and

emergent pathogens in aquatic systems. *Trends in Parasitology* **29**, 567–578 (2013).

13.  Buhroo, Z. Microsporidiosis in silkworms with particular reference to mulberry silkworm (Bombyx Mori L.). (2018).

14. Trzebny, A., Slodkowicz-Kowalska, A., Becnel, J. J., Sanscrainte, N. & Dabert, M. A new method of metabarcoding

Microsporidia and their hosts reveals high levels of microsporidian infections in mosquitoes (Culicidae). *Molecular*

*Ecology Resources* **20**, 1486–1504 (2020).

15. Emsen, B. *et al.* Higher prevalence and levels of Nosema ceranae than Nosema apis infections in Canadian honey

bee colonies. *Parasitology research* **115**, (2015).

16. Martín-Hernández, R. *et al.* Nosema ceranae in Apis mellifera: a 12 years postdetection perspective. *Environmental*

*Microbiology* **20**, 1302–1329 (2018).

17. Chayaburakul, K., Nash, G., Pratanpipat, P., Sriurairatana, S. & Withyachumnarnkul, B. Multiple pathogens found in

growth-retarded black tiger shrimp Penaeus monodon cultivated in Thailand. *Diseases of Aquatic Organisms* **60**, 89–

96 (2004).

18. Niehus, S., Giammarinaro, P., Liégeois, S., Quintin, J. & Ferrandon, D. Fly culture collapse disorder: detection,

prophylaxis and eradication of the microsporidian parasite Tubulinosema ratisbonensis infecting Drosophila

melanogaster. *Fly* **6**, 193–204 (2012).

19. Sanders, J. L., Watral, V. & Kent, M. L. Microsporidiosis in Zebrafish Research Facilities. *ILAR J* **53**, 106–113 (2012).

20. Troemel, E. R., Félix, M.-A., Whiteman, N. K., Barrière, A. & Ausubel, F. M. Microsporidia Are Natural Intracellular

Parasites of the Nematode Caenorhabditis elegans. *PLoS Biol* **6**, e309 (2008).

21. Balla, K. M., Luallen, R. J., Bakowski, M. A. & Troemel, E. R. Cell-to-cell spread of microsporidia causes

Caenorhabditis elegans organs to form syncytia. *Nat Microbiol* **1**, 16144 (2016).

22. Botías, C., Martín-Hernández, R., Barrios, L., Meana, A. & Higes, M. Nosema spp. infection and its negative effects on

honey bees (Apis mellifera iberiensis) at the colony level. *Veterinary research* **44**, 25 (2013).

23. Gupta, S. K., Hossain, Z., Nanu, M. M. & Mondal, K. Impact of microsporidian infection on growth and development

of silkworm Bombyx mori L. (Lepidoptera: Bombycidae). *Agriculture and Natural Resources* **50**, 388–395 (2016).

24. Li, W., Feng, Y. & Santin, M. Host Specificity of Enterocytozoon bieneusi and Public Health Implications. *Trends in*

*Parasitology* **35**, 436–451 (2019).

25. Hinney, B., Sak, B., Joachim, A. & Kváč, M. More than a rabbit's tale – Encephalitozoon spp. in wild mammals and

birds. *International Journal for Parasitology: Parasites and Wildlife* **5**, 76–87 (2016).

26. Han, B. & Weiss, L. M. Microsporidia: Obligate Intracellular Pathogens within the Fungal Kingdom. *Microbiol Spectr*

**5**, (2017).

27. Herren, J. K. *et al.* A microsporidian impairs Plasmodium falciparum transmission in Anopheles arabiensis

mosquitoes. *Nature Communications* **11**, 1–10 (2020).

28. Bjørnson, S. & Oi, D. Microsporidia Biological Control Agents and Pathogens of Beneficial Insects. in *Microsporidia*

635–670 (John Wiley & Sons, Ltd, 2014). doi:10.1002/9781118395264.ch25.

29. Vávra, J. & Larsson, J. I. R. Structure of Microsporidia. in *Microsporidia* 1–70 (John Wiley & Sons, Ltd, 2014).

doi:10.1002/9781118395264.ch1.

30. Han, B., Takvorian, P. M. & Weiss, L. M. Invasion of Host Cells by Microsporidia. *Front. Microbiol.* **11**, (2020).

31. Zhang, G. *et al.* A Large Collection of Novel Nematode-Infecting Microsporidia and Their Diverse Interactions with

Caenorhabditis elegans and Other Related Nematodes. *PLoS Pathog* **12**, e1006093 (2016).

32. Vizoso, D. B. & Ebert, D. Within-host dynamics of a microsporidium with horizontal and vertical transmission:

Octosporea bayeri in Daphnia magna. *Parasitology* **128**, 31–38 (2004).

33. Sprague, V. Annotated List of Species of Microsporidia. in *Comparative Pathobiology: Volume 2 Systematics of the*

*Microsporidia* (eds. Bulla, L. A. & Cheng, T. C.) 31–334 (Springer US, 1977). doi:10.1007/978-1-4613-4205-2_2.

34. Kirk, P. M. Microsporidia: Unicellular spore-forming protozoan parasites (version Nov 2015). *In: Species 2000 & ITIS*

*Catalogue of Life, 2020-09-01 Beta*.

35. Pabis, K. What is a moth doing under water? Ecology of aquatic and semi-aquatic Lepidoptera. *Knowl. Manag.*

*Aquat. Ecosyst.* 42 (2018) doi:10.1051/kmae/2018030.

36. Solter, L. F. & Maddox, J. V. Physiological Host Specificity of Microsporidia as an Indicator of Ecological Host

Specificity. *Journal of Invertebrate Pathology* **71**, 207–216 (1998).

37. Terry, R. S. *et al.* Widespread vertical transmission and associated host sex–ratio distortion within the eukaryotic

phylum Microspora. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**, 1783–1789 (2004).

38. Jaroenlak, P. *et al.* 3-Dimensional organization and dynamics of the microsporidian polar tube invasion machinery.

*PLOS Pathogens* **16**, e1008738 (2020).

39. James, T. Y. *et al.* Shared Signatures of Parasitism and Phylogenomics Unite Cryptomycota and Microsporidia.

*Current Biology* **23**, 1548–1553 (2013).

40. Miller, M. A., Pfeiffer, W. & Schwartz, T. Creating the CIPRES Science Gateway for inference of large phylogenetic

trees. in *2010 Gateway Computing Environments Workshop (GCE)* 1–8 (2010). doi:10.1109/GCE.2010.5676129.

41. R Development Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for Statistical

Computing, 2019).

42. Franzen, C. Microsporidia: A Review of 150 Years of Research. *TOPARAJ* **2**, 1–34 (2008).

43. Sapir, A. *et al.* Microsporidia-nematode associations in methane seeps reveal basal fungal parasitism in the deep

sea. *Front. Microbiol.* **5**, (2014).

44. Czaker, R. Wittmannia antarctica N. G., N. Sp. (Nosematidae), a New Hyperparasite in the Antarctic Dicyemid

Mesozoan Kantharella antarctica. *Journal of Eukaryotic Microbiology* **44**, 438–446 (1997).

45. Stentiford, G. D. *et al.* Microsporidia – Emergent Pathogens in the Global Food Chain. *Trends Parasitol* **32**, 336–348

(2016).

46. Jarkass, H. T. E. & Reinke, A. W. The ins and outs of host-microsporidia interactions during invasion, proliferation

and exit. *Cellular Microbiology* **22**, e13247 (2020).

47. Technau, U. & Scholz, C. B. Origin and evolution of endoderm and mesoderm. *Int. J. Dev. Biol.* **47**, 531–539 (2003).

48. Oenocytes in insects | Invertebrate Survival Journal. http://www.isj.unimo.it/index.php/ISJ/article/view/268.

49. Kudo, R. Notes on Nosema apis Zander. *The Journal of Parasitology* **7**, 85–90 (1920).

50. Weidner, E. Ultrastructural study of microsporidian invasion into cells. *Z. Parasitenk.* **40**, 227–242 (1972).

51. Vossbrinck, C. R. & Debrunner-Vossbrinck, B. A. Molecular phylogeny of the Microsporidia: ecological,

ultrastructural and taxonomic considerations. *Folia Parasitologica* **52**, 131–142 (2013).

52. Williams, B. A. P., Hamilton, K. M., Jones, M. D. & Bass, D. Group-specific environmental sequencing reveals high

levels of ecological heterogeneity across the microsporidian radiation. *Environmental Microbiology Reports* **10**, 328–

336 (2018).

53. Cormier, A. *et al.* Draft Genome Sequences of Thelohania contejeani and Cucumispora dikerogammari, Pathogenic

Microsporidia of Freshwater Crustaceans. *Microbiol Resour Announc* **10**, (2021).

54. Tokarev, Y. S. *et al.* A formal redefinition of the genera Nosema and Vairimorpha (Microsporidia: Nosematidae) and

reassignment of species based on molecular phylogenetics. *Journal of Invertebrate Pathology* **169**, 107279 (2020).

55. Larsen, B. B., Miller, E. C., Rhodes, M. K. & Wiens, J. J. Inordinate Fondness Multiplied and Redistributed: the

Number of Species on Earth and the New Pie of Life. *The Quarterly Review of Biology* **92**, 229–265 (2017).

56. Costello, M. J. Parasite Rates of Discovery, Global Species Richness and Host Specificity. *Integr Comp Biol* **56**, 588–

599 (2016).

57. Jorge, F. & Poulin, R. Poor geographical match between the distributions of host diversity and parasite discovery

effort. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20180072 (2018).

58. Rost-Roszkowska, M. M., Poprawa, I. & Kaczmarek, Ł. Autophagy as the cell survival in response to a microsporidian

infection of the midgut epithelium of Isohypsibius granulifer granulifer (Eutardigrada: Hypsibiidae). *Acta Zoologica*

**94**, 273–279 (2013).

59. Ardila-Garcia, A. M. & Fast, N. M. Microsporidian Infection in a Free-Living Marine Nematode. *Eukaryotic Cell* **11**,

1544–1551 (2012).

60. Bass, D. *et al.* Clarifying the Relationships between Microsporidia and Cryptomycota. *Journal of Eukaryotic*

*Microbiology* **65**, 773–782 (2018).

61. Hejnol, A. & Martín-Durán, J. M. Getting to the bottom of anal evolution. *Zoologischer Anzeiger - A Journal of*

*Comparative Zoology* **256**, 61–74 (2015).

62. Franzen, C., Hösl, M., Salzberger, B. & Hartmann, P. UPTAKE OF ENCEPHALITOZOON SPP. AND VITTAFORMA

CORNEAE (MICROSPORIDIA) BY DIFFERENT CELLS. *para* **91**, 745–749 (2005).

63. Monaghan, S. R. *et al.* Animal cell cultures in microsporidial research: their general roles and their specific use for

fish microsporidia. *In Vitro Cell.Dev.Biol.-Animal* **45**, 135 (2009).

64. Larsson, J. I. Molecular versus morphological approach to microsporidian classification. *Folia Parasitologica* **52**, 143–

144 (2013).

65. Cuomo, C. A. *et al.* Microsporidian genome analysis reveals evolutionary strategies for obligate intracellular growth.

*Genome Res.* **22**, 2478–2488 (2012).

 **Figures**

542 **Fig. 1. History and geography of microsporidia species discovery. (A)** Number of new microsporidia species

543 described per decade (n = 1415 microsporidia species). (**B-C**) Geographical location of microsporidia, with each

544 species depicted once for each country or ocean where they have been reported (n = 1301 microsporidia

545 species). **(B)** Number of microsporidia species found on each continent. **(C)** Location of microsporidia species

546 identified, by country and ocean. The number of species found in each country is depicted as a heatmap,

     according to the scale at the bottom. Countries in grey indicate that no species have been reported. Numbers in

     blue circles describe species found in respective oceans.

**Fig 2. Microsporidia infect a diverse range of hosts from all major environments.** Host taxonomy and

environment was determined for each reported microsporidia species (see methods). **(A)** Distribution of

microsporidia hosts by metazoan (left) and protist (right) phyla. The number of host species in each phylum

(blue) and the total number of extant species in each phylum (red) are shown. Phyla where the majority of

microsporidia hosts are parasites are indicated (**)(n = 1422 host species). **(B)** Number of described species

from phyla reported to be infected by microsporidia (red) (n = 15 phyla) and those that have not been reported

to be infected by microsporidia (blue) (n = 17 phyla). The Mann-Whitney U test was used to compare mean ranks

between the groups. **(C)** Frequency of Chordate orders reported to be infected by microsporidia (n = 39 orders).

**(D)** Venn diagram of the major environments that microsporidia hosts dwell in (n = 1339 host species).

**Fig. 3. Microsporidia infect a diversity of tissues and species that infect more hosts tend to infect a broader set of tissues. (A-B)** The tissues described to be infected by microsporidia were placed into 23 tissue-types (see methods). **(A)** The number of microsporidia species reported to infect each tissue type (n = 1111 species). **(B)** Distribution of the number of tissue types infected by each microsporidia species (n = 1009 species). **(C)** Distribution of the number of hosts infected by each microsporidia species (n = 1422 species). **(D)** Proportion of species that infect more than one tissue compared to how many hosts are infected by that species (n = 1066). Species that are described to have general or systemic infections are considered to infect more than one tissue. Significant difference between proportions were determined with pairwise chi-square tests, with Bonferroni p-value correction ($\alpha = 0.05/6$ comparisons $= 8.3 \times 10^{-3}$).

**Fig. 4. Microsporidia spores display a diversity of shapes, sizes, and polar tube lengths. (A-B)** Spore shapes were placed into 20 morphology classes (see methods and Table S4). **(A)** The frequency of each type of spore shape type (n = 1103 spores). **(B)** Spore length and width dimensions plotted for the six most common spore shapes (n = 807 spores). **(C)** Comparison of experimentally determined and calculated polar tubule lengths (n = 34 spores). **(D)** Median spore volume for uninucleate (n = 467) and binucleate (n = 263) spores. The p-value was determined by the Mann–Whitney U test. **(E)** Comparison between experimental polar filament lengths and spore volume (n = 37 spores for uninucleate and 26 species for binucleate). Correlation was calculated with spearman $r_s$. **(F)** Relationship between polar tubule length and tissues infected. Significance was calculated using a paired T-test between "Intestinal" species and "Other Tissues" species infecting the same host.

**Fig. 5. Bayesian 18S Phylogeny of Microsporidia Species Reveals Four Phenotypically Diverse Clades.**

**(A)** 18S phylogeny of 273 microsporidia species, inferred under a Bayesian framework (see methods). Clades

are colored and number of hosts indicated according to the legends at the right. Species not belonging to any of

the four main clades are colored black. **(B-D)** Analysis of microsporidia phenotypes in each of the four major

clades. **(B)** Frequency of host environments by clade. **(C)** Spore volume distribution by clade. **(D)** Calculated

polar tube length distribution by clade. Significance evaluated using pairwise Mann-Whitney U-tests, with

Bonferroni p-value adjustments applied to correct for multiple testing ($\alpha = 0.05/6 = 8.3 \times 10^{-3}$) **(E)** Number of

hosts infected by microsporidia species from one, two or three different clades (n = 408 microsporidia hosts).

**Fig. 6. Evolutionary relatedness of microsporidia phenotypes. (A)** Distribution of species pairs according to

sequence similarity (n = 37,128 comparisons). **(B-F)** Distribution of sequence pair similarity and frequency of

shared **(B)** host taxonomy (n = 35,778 comparisons), **(C)** environments (n = 34,453 comparisons), **(D)** infected

tissues (n = 27,730 comparisons), **(E)** difference in spore volume (n = 32,131 comparisons), **(F)** difference in

calculated polar tube length (n = 17,955 comparisons).

**Supplemental material**

**Fig. S1. Orders of Arthropods infected by microsporidia.**

**Fig. S2. Relationship between microsporidia spore dimensions and shape classification.**

**Fig. S3. Relationship between polar tube length and spore volume.**

**Fig. S4. Microsporidia host environment and spore morphology are independent of infection transmission mode.**

**Fig. S5. Bayesian 18S phylogeny of microsporidia species.**

**Table S1. Database of microsporidia species attributes.**

**Table S2. Taxonomical descriptions of microsporidia hosts.**

**Table S3. Microsporidia host-tissues classifications.**

**Table S4. Microsporidia spore-shape classifications.**

**Table S5. Polar tube lengths of species pairs that infect only the intestine and also other tissues in a common host.**

**Table S6. Accession numbers for microsporidia DNA sequences.**

**Table S7. Parameters for MrBayes phylogenetic analysis.**

620 **Fig. S1. Frequency of Arthropod order infections.**

621 Frequency of Arthropod orders reported to be infected by microsporidia (n = 46 orders).

623 **Fig. S2. Relationship between microsporidia spore dimensions and shape classification.**

624 Spore length and width dimensions for the six most common shape groups (Table S4) with the 16 remaining

625 shapes classified as other (n=1023 spores).

A

B

627 **Fig. S3. Relationship between polar tube length, nuclei number and spore volume.**

628 **A.** Comparison between experimental polar filament lengths and spore volume for spores of any nuclei number

629 (n = 147). Correlation was calculated with spearman $r_s$. **B.** Median spore experimental tube length for uninucleate

630 (n = 39) and binucleate (n = 29) spores. The p-value was determined by the Mann–Whitney U test.

**Fig. S4. Microsporidia host environment and spore morphology are independent of infection transmission mode.** The transmission mode of each species of microsporidia was inferred from tissues infected, with those that only infect somatic tissues being classified as horizontally transmitted, those that only infect germline tissues being classified as vertically transmitted, and those that infect both types of tissues being classified as both (see methods). **(A)** Venn diagram of transmission mode of microsporidia species (n = 1075 microsporidia species). **(B)** Comparison of infection transmission mode to host environment (n = 553 hosts). Only hosts classified as belonging to a single environment were analyzed. **(C)** Comparison of infection transmission mode to experimentally measured polar tube values (n = 141 spores) and **(D)** calculated polar tube values (n = 478 spores). Species with multiple polar tube values were represented multiple times. **(E)**

642  Comparison of infection transmission mode to spore volume (n = 1135 spores). Species with multiple spore sizes

643  were represented multiple times.

Loma_wallae
Loma_pacificodae
•• Loma_morhua
Loma_lingcodae
Loma_embiotocia
Loma_richardi
Loma_kenti
•• Loma_salmonae
Loma_sp
Loma_branchialis
Pseudoloma_neurophilia
Microsporidium_luciopercae
•• Ichthyosporidium_giganteum
Glugea_jazanensis
Glugea_arabica
Glugea_sardinellensis
Pleistophora_finisterrensis
Glugea_anomala
Glugea_plecoglossi
•• Glugea_stephani
• Glugea_hertwigi
Glugea_pagri
Glugea_atherinae
Glugea_gasterostei
Loma_psittaca
Loma_acerinae
Microsporidium_cypselurus
Pleistophora_beebei
Ovipleistophora_arlo
Ovipleistophora_diplostomuri
•• Pleistophora_hyphessobryconis
Ovipleistophora_mirandellae
•• Vavraia_culicis
Trachipleistophora_hominis
Heterosporis_saurida
Heterosporis_lessepsianus
Heterosporis_anguillarum
Pleistophora_pagri
Pleistophora_aegyptiaca
Pleistophora_sp_1
Pleistophora_hippoglossoideos
Pleistophora_typicalis
• Pleistophora_ehrenbaumi
Pleistophora_mulleri
Pleistophora_sp_2
Pleistophora_sp_3
•• Pleistophora_ladogensis
Myosporidium_merluccius
• Myosporidium_spraguei
Potaspora_morhaphis
Potaspora_aequidens
Potaspora_macrobrachium
Microsporidium_sp_SH
Kabatana_takedai
Kabatana_rondoni
Spraguea_lophii
Glugea_americanus
Microgemma_tincae
•• Tetramicra_brevifilum
Microgemma_caulleryi
Microgemma_carolinus
Hyperspora_aquatica
Cucumispora_ornata
•• Cucumispora_dikerogammari
Thelohania_butleri
Myospora_metanephrops
• Perezia_nelsoni
•• Facilispora_margolisi
•• Dictyocoela_muelleri
Dictyocoela_gammarellum
• Dictyocoela_roeselum
Dictyocoela_duebenum
Unikaryon_legeri
Ameson_sp
Ameson_herrnkindi
Ameson_michaelis
Nadelspora_canceri
Ameson_pulvis
Ameson_metacarcini
Thelohania_montirivulorum
• Thelohania_contejeani
Thelohania_parastaci

**Fig. S5. Bayesian 18S phylogeny of microsporidia species.** Phylogenetic tree of 273 microsporidia species

and *Rozella allomycis*. The tree was generated and labelled for clades and environments the same as in Fig 5.

The species name of each branch is shown on the right, and the Bayesian posterior probability support value is

shown at each node.