Tetrameric UvrD helicase is located at the *E. coli* replisome due to frequent replication blocks

Adam J. M Wollman[1,2,3,], Aisha H. Syeda[1,2], Jamieson A. L. Howard[1,2], Alex Payne-Dwyer[1,2], Andrew Leech[4], Dominika Warecka[2], Colin Guy[5, 6], Peter McGlynn[2,6], Michelle Hawkins[2] and Mark C. Leake[1,2]

The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First Authors

[1] School of Physics, Engineering and Technology, University of York, York YO10 5DD, United Kingdom.

[2] Department of Biology, University of York, York YO10 5DD, United Kingdom.

[3] Current address: Biosciences Institute, Newcastle University, NE1 7RU, United Kingdom.

[4] Bioscience Technology Facility, Department of Biology, University of York, York YO10 5DD, United Kingdom

[5] Current address: Covance Laboratories Ltd., Otley Road, Harrogate, HG3 1PY, United Kingdom

[6] Previous address: School of Medical Sciences, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, United Kingdom

* To whom correspondence should be addressed. To whom correspondence should be addressed.

Tel: +44 (0)1904322697. Email: mark.leake@york.ac.uk

Present Address: School of Physics, Engineering and Technology, and Department of Biology, University of York, York YO10 5DD, United Kingdom

SUMMARY

DNA replication in all organisms must overcome nucleoprotein blocks to complete genome duplication. Accessory replicative helicases in *Escherichia coli*, Rep and UvrD, help remove these blocks and aid the re-initiation of replication. Mechanistic details of Rep function have emerged from recent live cell studies; however, the division of UvrD functions between its activities in DNA repair and role as an accessory helicase remain unclear in live cells. By integrating super-resolved single-molecule fluorescence microscopy with biochemical analysis, we find that UvrD self-associates into tetrameric assemblies and, unlike Rep, is not recruited to a specific replisome protein despite being found at approximately 80% of replication forks. Instead, its colocation with forks is likely due to the very high frequency of replication blocks composed of DNA-bound proteins, including RNA polymerase and factors involved in repairing DNA damage. Deleting *rep* and DNA repair factor genes *mutS* and *uvrA*, and inhibiting transcription through RNA polymerase mutation and antibiotic inhibition,

37 indicates that the level of UvrD at the fork is dependent on UvrD's function. Our
38 findings show that UvrD is recruited to sites of nucleoprotein blocks via different
39 mechanisms to Rep and plays a multi-faceted role in ensuring successful DNA
40 replication.
41
INTRODUCTION
Replication and transcription of DNA occur simultaneously in *Escherichia coli,* making
conflicts between the bacterial replisome, the molecular replication machine
comprising in excess of 10 different proteins, and RNA polymerase (RNAP) inevitable
(1–4). Collisions between replication and transcription machineries hinder replication
fork progression and cause genome instability (5–14). RNAP can pause, stall and
backtrack while actively transcribing, and any immobile RNAP presents long-lived
barriers to replisomes (15, 16); RNAP is the most common nucleoprotein obstacle for
translocating replisomes. DNA is also frequently damaged in normal growth conditions
due to exposure to endogenous and exogenous DNA damaging agents, which also
presents significant obstacles for replication (17–19).
Several mechanisms exist to reduce conflict between the replisome and nucleoprotein
blocks. For example, transcription elongation and termination factors reduce the
number of immobile RNAPs on DNA (1, 18, 20–24). Head-on collisions were originally
thought to be more harmful (25–27) and therefore more important to resolve; a view
supported by the occurrence of highly transcribed genes encoded on the leading
strand, presumably to ensure co-directional transcription and replication (26, 28, 29).
However, co-directional collisions of the replisome and RNAP also impact fork
progression (3, 18, 30).
Accessory replicative helicases, identified in prokaryotes and eukaryotes, promote
fork movement through resolving nucleoprotein blocks (1, 12, 13, 31, 32). The Rep
and UvrD accessory helicases in *E. coli*, as well as the Pif1 helicase in *Saccharomyces*
cerevisiae, perform important roles in helping to clear nucleoprotein complexes (33).
More recently, Pif1 has been proposed to be a general displacement helicase for
replication bypass of protein blocks, a function similar to the Rep and UvrD helicases
(34). These helicases can reduce replisome pausing at several types of native
transcription complexes in living cells (13, 22, 35), and of engineered nucleoprotein
blocks in biochemical experiments *in vitro* (1, 31).

Rep is the primary accessory helicase in *E. coli*, however, UvrD, a superfamily 1

helicase that translocates 3'-5' along ssDNA, partially compensates for its absence *in*

vivo, attributed to the high degree of homology between these two helicases and the

abundance of UvrD inside cells (12, 21, 36–38). Rep also promotes replication through

stalled transcription elongation complexes (1, 12, 13, 39, 40). UvrD's role in nucleotide

excision repair along with UvrABC proteins, as well as in methyl-directed mismatch

repair along with the MutHLS proteins, is well characterised (41–46). UvrD interacts

directly with RNAP but the functional importance of this interaction has been unclear

until recently (47–51) . UvrD has been suggested to promote backtracking of stalled

RNAP as a first step in transcription-coupled repair (50, 52); however, other studies

argue against UvrD playing any role in coupling nucleotide excision repair to stalled

transcription complexes (48, 49, 53–57).

Our aim in this study was to establish the role of UvrD during DNA replication. The

primary objective was to test whether UvrD is present at active replication forks using

dual-color single-molecule imaging of fluorescently labelled UvrD and replication forks

in living *E. coli* cells. We conclude that UvrD is present at replication forks but is not

recruited via specific protein interactions with the replisome, which means that it

operates in a fundamentally different way to that of Rep. By using super-resolved

millisecond single-molecule fluorescence microscopy of mGFP-UvrD in live cells co-

expressing a fork marker, DnaQ-mCherry, we show that 80% of DnaQ foci colocalise

with UvrD. Stepwise photobleaching intensity analysis reveals tetrameric UvrD foci *in*

vivo, to be compared with earlier observations performed *in vitro* using analytical

ultracentrifugation that found a mixture of UvrD dimers and tetramers (58). Surface

plasmon resonance (SPR) of UvrD in combination with each replisome component

identified no specific replisome protein interaction that would explain replisome-UvrD

colocalisation. Deletion of Rep, mismatch repair protein MutS and nucleotide excision

repair protein UvrA, and disruption of transcription using either mutated RNAP or

antibiotic inhibition, significantly reduces the number of UvrD or level of observed

colocalisation of UvrD with the fork. Coupled with *in vitro* block-dependent differences

in UvrD replication promotion efficiency, we conclude that the colocation of UvrD with

the replisome stems primarily from UvrD resolving frequent nucleoprotein blocks to

the DNA replication machinery, as opposed to it binding specifically to a replisome

component, contrasting distinctly with the replisome-binding mode of operation of Rep.

RESULTS

UvrD is present at the majority of replication forks

Figure 1: Super-resolved single-molecule light microscopy indicates fork-UvrD colocation. (A). Slimfield microscope schematic and micrographs of mGFP-UvrD, DnaQ-mCherry. Cell membrane and DnaQ/UvrD foci indicated as white dashed and non-dashed lines respectively. (B) and (C) Histogram showing the number of DnaQ and UvrD foci detected per cell respectively, SD errors. N=45 cells.

We investigated UvrD's role in DNA replication using super-resolved single-molecule fluorescence microscopy to track dynamic patterns of fluorescently labelled UvrD localization relative to the replication fork in live cells (Figure 1A). We performed dual-color Slimfield (59, 60) to assess the degree of colocalisation between UvrD and the replisome, using the same imaging protocol used previously to confirm Rep colocalisation with the replisome (13). We employed DnaQ-mCherry as a fork marker - DnaQ encodes a core epsilon subunit of DNAP (13, 61–64) - along with genomically integrated mGFP-UvrD to report on the localization of UvrD. These strains phenocopied wild type for growth and in plasmid loss assays for mGFP-UvrD function retention (Supplementary Figures S1, S2, Tables S1-4). *In vitro* unwinding assays indicate that the nucleoprotein block removal activity of mGFP-UvrD was comparable to wild type (Supplementary Figure S3; note, a summary of all activity comparisons between mGFP-UvrD and wild type UvrD is shown in Table S5). Using Slimfield, we observed 1-2 DnaQ foci per cell, as well as a background 'pool' of unbound DnaQ molecules in the cytoplasm (61), as in the previous study for Rep, corresponding to the two moving forks which appear as a single focus near the start of replication when they are separated by less than the 200-300 nm diffraction limit of our microscope (Figure 1A,B Supplementary Figure S4). We also observed 1-2 UvrD foci per cell (and associated pool), of which $67 \pm 7\%$ ($\pm$SEM) were colocalised with DnaQ foci and $84 \pm 5\%$ of DnaQ foci colocalised with UvrD (Figure 1B and C). We calculated the

probability of random overlap between foci by modelling nearest-neighbor separation
as a Poisson distribution (65), indicating 20%. To confirm that colocalisation between
DnaQ and UvrD was not just a function of the nucleoid association of UvrD, we
performed Slimfield on Heat-stable nucleoid-structuring protein (H-NS) tagged with
GFP - H-NS associates with the nucleoid but not specifically with the fork - and co-
labelled DnaQ-mCherry. We found similar numbers of H-NS foci to UvrD
(Supplementary Figure S5) but only 38 ± 4% DnaQ foci were colocalised with H-NS
foci implying that, similar to Rep (13), UvrD is present at the majority of forks.
141
142   **UvrD foci are tetrameric assemblies in live bacteria**
143

145   **Figure 2 UvrD stoichiometry shows tetrameric periodicity.** The distribution of UvrD foci
146   stoichiometry rendered as a kernel density estimate for foci colocalised (A) and not colocalised
147   (B) with DnaQ foci. Distributions were fitted with multiple Gaussian peaks (black dotted lines)
148   with the peak value ± error determined by the half width at half maximum indicated above
149   each peak
150

Slimfield also enabled the stoichiometry of these foci (i.e., the number of fluorescently-
labelled molecules within each focus) to be determined by utilizing a method involving
stepwise photobleaching analysis (61). Here, we determined the brightness of a
single-molecule of mGFP or mCherry fluorescent protein using Slimfield
(Supplementary Figure S7), and used these values to normalize the initial brightness
of tracked foci to estimate the stoichiometry. We found that DnaQ had the same range
of 2-6 molecules per focus (Supplementary Figure S8) as found previously,
corresponding to 2-3 polymerases per fork (13, 61). The distribution of the
stoichiometry of UvrD foci which are colocalised with DnaQ had a distinct lowest-order

peak corresponding to approximately four molecules, with subsequent peaks at
multiples of four (Figure 2A). UvrD foci that were not colocalised with DnaQ also
contained a distinct lowest-order peak at four, but subsequent peaks were less clearly
tetrameric.
Although we utilised a definitive monomeric variant of GFP which contained an A206K
mutation to inhibit dimerization (66), we did not want to dismiss possible non-
physiological influences of fluorescent protein labelling of UvrD in its self-assembly
process. Using Size Exclusion Chromatography Multi-Angle Laser Light Scattering
(SEC-MALLS) we found qualitatively that the majority of UvrD runs off in a monomeric
fraction for both purified wild type UvrD and for mGFP-UvrD (Supplementary Figure
S9). However, SEC-MALLS has known limitations in differentiating between
monomeric and oligomeric fractions due to non-specific binding to the SEC resin
(depending on buffering conditions), resulting in proteins eluting later than expected
and co-elution of monomeric species and higher mass oligomers. In addition, there
are sensitivity difficulties for detecting components present in relatively small
quantities – we estimate the error on the cumulative weight fraction to be ~20% on our
system, consistent with the reported sensitivity error reported by others (67, 68).
To overcome these limitations, we performed high-precision high-sensitivity Slimfield
of purified mGFP-UvrD conjugated to a glass coverslip via anti-GFP antibody (69)
enabling detection of fluorescent foci bound to the surface with single-molecule
sensitivity. Slimfield indicated between ~10-20 surface-bound foci per field of view
(Supplementary Figure S9A), whose 200-300 nm width (half width at half maximum,
HWHM) was comparable to that determined for single purified GFP molecules. We
measured the fluorescence brightness of each focus using the same automated
single-particle tracking software as for Slimfield *in vivo* (Supplementary Figure S9B),
whose tracks photobleached in a stepwise manner consistent with the presence of
just a single GFP molecule. The amplitude of the probability distribution function of
foci brightness values (Supplementary Figure S9B) indicated that the proportion of
dimers (corresponding to a brightness of approximately twice the modal value) is ~3%
that of the corresponding value for monomers. Although it is not possible to entirely
exclude non-physiological effects of any fluorescent protein on the cellular activities of
a tagged protein, the finding from this single-molecule assay shows that the proportion
of monomeric mGFP-UvrD is substantially higher than dimeric mGFP-UvrD and so, if

there is an influence of mGFP on the self-assembly process of UvrD, it is unlikely to be a dominant factor.

At higher concentrations, UvrD has been observed to exist in a monomer-dimer-tetramer equilibrium *in vitro* (58). DNA unwinding experiments *in vitro* suggest UvrD must function as at least a dimer (70) or a trimer (71), however the oligomeric state of functional UvrD is disputed (reviewed in (72)). Our results are consistent with UvrD existing predominantly as a tetramer, or a dimer of dimers, with multiple tetramers or dimer-dimers sometimes present at forks. It should be noted that we often observe only a single DnaQ focus corresponding to two forks which are separated by less than the diffraction limit of our microscope (Supplementary Figure S8) such that some multiple tetrameric UvrD foci correspond to two replisomes. The less distinct peaks in the UvrD foci that are not colocalised with DnaQ may indicate that other UvrD species (monomers, dimers or trimers) are also present.

We determined the total cell copy number of UvrD by estimating the total contribution of mGFP-UvrD fluorescence to the observed intensity in whole cells (73), indicating approximately 800 molecules of UvrD per cell (Supplementary Figure S10), similar to that reported previously for Rep (13), and in line with other proteins that have a small active population detected as foci and a larger population of diffusive molecules in a protein pool (61, 73–76). This is a lower level of UvrD than was reported using transcription rate measurements (77) or high-throughput proteomics (78) but is comparable to recent estimates using ribosome profiling (79).

There is no obvious replisome interaction partner for UvrD at the fork

Our earlier findings concerning Rep indicated that colocalisation with the fork and recruitment to the replisome were mediated by protein-specific interaction with the replicative helicase DnaB (80), consistent with Rep and DnaB both exhibiting hexameric stoichiometry (13, 61). We sought to determine whether UvrD is also recruited by interacting with a specific replisome component, expanding on a previous study which used SPR to test whether purified UvrD interacted with each replisome component(12). In this study, UvrD, under conditions consistent with predominantly monomeric UvrD (Supplementary Figure S9), exhibited no direct interaction with primase, SSB, β sliding clamp, the DNA polymerase III $\alpha\epsilon$, $\chi\psi$ or γ complexes; or θ, δ, δ', χ, and γ subunits (12). However, it did identify a putative weak UvrD-Tau interaction which we interrogated further with more sensitive SPR experiments. We

were able to replicate the weak UvrD-Tau interaction; however, as a control we tested

Rep and found this also interacted with Tau to a similar extent (Supplementary Figure

S11). Tau interactions with either helicase were ~100x lower affinity than UvrD with

known interaction partner UvrB (Supplementary Figure S11). We therefore conclude

that the Tau interaction with UvrD is likely to be non-specific and that, unlike Rep, UvrD

does not have an obvious, direct interaction partner at the replisome, at least from the

suite of replisome proteins tested. Since our findings suggest that UvrD may exist as

as a tetramer at the replisome, it is also unlikely to be recruited by Rep, which has

shown putative interaction (81).

The presence of UvrD at the fork is mediated by specific replication blocks

Figure 3: Perturbing UvrD function using mutations and rifampicin antibiotic intervention influences fork-UvrD colocation. A. and B. The mean proportion of colocalised DnaQ and UvrD or Rep foci per cell respectively. Error bars represent standard error in the mean and statistical significance by Brunner-Munzel and Student's t-test $p<0.05$ indicated by *. Gray bar indicates levels of predicted random colocalisation. C. and D. Jitter plots of colocalised and not colocalised UvrD stoichiometry respectively. Bars show mean UvrD stoichiometry with error bars showing the standard error in the mean and statistical significance by Brunner-Munzel and Student's t-tests $p<0.05$ indicated by *. N=30 cells.

Alternatively, we hypothesized that the association of UvrD with the replication fork might be dependent on its activity in resolving nucleoprotein blocks. To test, we systematically impaired key DNA replication repair processes in which UvrD is implicated, as well as perturbing RNAP replicative blocks by disruption of transcription through both genetic mutation and antibiotic inhibition (Figure 3). We investigated how these perturbations altered fork association through UvrD colocalisation with DnaQ. As a control, we compared these to Rep colocalisation with DnaQ which can be abrogated down to random levels of colocalisation, indicating no interaction with the fork, by mutations which stop Rep interacting with DnaB and PriC (Figure 3A). We perturbed UvrD's activity in resolving DNA damage at replication blocks by deleting *mutS* or *uvrA*, rendering the strains defective in UvrD-associated mismatch repair or nucleotide excision repair respectively (43, 82). We reduced the need for UvrD to remove transcriptional blocks to replication (49) by introducing the *rpoB*35* mutation, which destabilizes open complexes formed during initiation of transcription (83, 84); and by using the antibiotic rifampicin, which blocks elongation of RNA transcripts (85). All of these strains were healthy under normal growth conditions (Supplementary Figure S1, Table S4) and displayed the same phenotype for DnaQ foci number and stoichiometry within measurement error (Supplementary Figure S8).

For the transcriptional block perturbations, the *rpoB*35* mutant dual labelled strain exhibited no difference in the proportion of colocalised DnaQ or UvrD foci (Figure 3A and B). This mutant and all others tested exhibited the same predominant tetrameric stoichiometry trends as wild type (Supplementary Figure S12), with clear peaks at tetramer intervals in the colocalised stoichiometry distribution and less clear but still largely tetrameric intervals in the stoichiometry distribution for UvrD foci that are not colocalised with DnaQ. This provides further evidence for tetrameric UvrD assemblies *in vivo*. We compared the mean stoichiometry for each mutant to that of wild type (Figure 3C and D). Unexpectedly, the *rpoB*35* mutant produced a statistically

significant increase in mean colocalised stoichiometry, corresponding to approximately one extra UvrD tetramer at the replication fork (Figure 3C). This finding may indicate the level of instability conferred by this mutation has multiple downstream effects that alter UvrD localisation significantly at the fork. Alternatively, it could mean that *rpoB*35* does not affect the formation of the types of transcriptional block that are acted upon by UvrD. Rifampicin treatment produced a clearer response, reducing the number of DnaQ foci colocalised with UvrD by ~20% (Figure 3A) and the mean stoichiometry of colocalised and not colocalised UvrD foci by around one UvrD tetramer (Figure 3C and D). Single-molecule observations of RNAP have shown it to be significantly more mobile under rifampicin treatment (86), implying fewer RNAPs bound to DNA and fewer blocks to replication. Our rifampicin treatment results are consistent with fewer transcriptional blocks to replication reducing the amount of UvrD recruited to the replisome and support the hypothesis that UvrD is recruited by these replicative blocks.

By deleting *mutS* or *uvrA* we removed UvrD's role in mismatch repair and nucleotide excision repair respectively. In the Δ*mutS* mutant we observed no change in UvrD colocalisation with DnaQ, but did measure an increase in the mean UvrD stoichiometry colocalised with DnaQ by around one tetramer (Figure 3A). In the Δ*uvrA* mutant we observed a decrease of ~25% in UvrD colocalisation with DnaQ (Figure 3B) with no change in the number of UvrD foci (Supplementary Figure S8). This agrees with a recent model that UvrA and UvrD are continuously associated with a subpopulation of RNAP to form a pre-transcription-coupled-repair complex (pre-TCRC); UvrD is then recruited at DNA lesions to facilitate DNA repair (50).

UvrD has been shown to compensate for Rep in *rep* null mutants (12). We also tested what effect this mutation would have in Slimfield *in vivo*. Surprisingly, *rep* deletion resulted in a steep drop in UvrD copy number (Supplementary Figure S10). The degree of colocalisation between UvrD and DnaQ remained unchanged compared to wild type (Figure 3A and B) but foci stoichiometry was reduced to a single UvrD tetramer (Figure 3C and D and Supplementary Figure 11), consistent with the observed lower copy number and availability of UvrD. The drop in UvrD copy number may reflect complex regulatory shifts in response to losing a key protein in Rep, although it is interesting that even at low copy numbers, a UvrD tetramer remains

308   recruited to the fork. Double knockouts, deficient in both Rep and UvrD, are lethal (12)

309   so this minimal helicase presence must be the limit required for survival.

Dynamic association of UvrD with the fork depends on DNA repair processing

Figure 4: UvrD dynamics are heterogeneous. A. Distribution of time over which UvrD and DnaQ foci were colocalised (blue) fitted with an exponential (red) to yield a characteristic dwell time. B. Distribution of microscopic diffusion coefficients of wild type UvrD foci (gray) fitted with a three-state gamma distribution model containing a relatively immobile population with $D = 0.1$ μm^2/s (red), a transiently immobile population, $D = 0.5$ μm^2/s (blue) and a mobile population, D=1.2 μm^2/s (green). C. The dwell times for each UvrD strain. Error bars represent the 95% confidence intervals on the fit. D. The proportion of UvrD foci in the immobile

320 population for each UvrD strain colocalised or not colocalised with DnaQ foci location.
321 Statistically significant differences p<0.05 from wild type indicated by *. E. PALM dwell time
322 and colocalisation proportion of detected tracks. N=45 cells for wild type and 30 per mutant,
323 with ~300 trajectories.
324

By tracking UvrD foci, we were also able to characterize their dynamics in wild type

and repair impaired cells. We measured the exposure time per frame multiplied by the

number of image frames in which UvrD foci were colocalised with DnaQ foci and fitted

the distribution of these dwell times with a single exponential function. We also

calculated the apparent microscopic diffusion coefficient of individual foci (Figure 4

and Supplementary Figure S13). The mean UvrD dwell time at the fork (corresponding

to the exponential fit time constant) for the wild type strain was measured to be

approximately 4 ms, indicating a relatively dynamic association of UvrD at the

replication fork. The diffusion coefficients (D) were best fit with a three parameter

Gamma model, containing immobile (D = 0.1 μm^2/s), transiently immobile population

(D = 0.5 μm^2/s) and mobile (D = 1.2 μm^2/s) populations similar to Rep. In wild type,

we found that approximately 10% of UvrD foci were immobile, whether colocalised

with the fork or not, with the remaining foci split between the other two populations.

We applied the same approach to cells which had been perturbed using either a range

of mutations which were known to affect DNA replication repair and block resolution,

or antibiotic treatment known to affect expression of RNAP. All perturbations resulted

in a marginal though statistically insignificant increase in mean dwell times to between

6-9 ms. The Δ*rep* and *rpoB*35* mutations resulted in no change in mobility from wild

type. Rifampicin treatment however, resulted in an increase in the proportion of

immobile foci both at and away from the fork. This finding appeared at first

counterintuitive since rifampicin treatment results in more mobile RNAP (86), however

it is unclear what effect rifampicin has on the population of RNAP molecules that

remain bound to DNA. These results may suggest that such blocks provide an

increased barrier to replication, decreasing UvrD mobility as it attempts to resolve

blocks. In Δ*mutS* and Δ*uvrA* strains, the proportion of immobile foci decreased at the

fork, again consistent with the hypothesis that UvrD association with the fork is

dependent on DNA damage resolution activity. One alternative explanation for these

results is that UvrD is less mobile when resolving DNA damage blocks to replication

and more mobile when resolving transcriptional blocks, as reducing or eliminating

UvrD's role in these processes decreased or increased the immobile fraction in DNA
repair and transcription impaired cells respectively (Figure 4).
356
***In vitro* biochemical experiments support the block-dependent UvrD function**
model
To investigate the role of different nucleoprotein blocks in UvrD recruitment to the
replisome, we reconstituted replisomes on plasmid templates containing *oriC* and
block sites. This *in vitro* approach allowed us to specify the type of nucleoprotein block
and test whether the ability of UvrD to promote replication is affected by whether UvrD
recruitment occurs before or after the replisome encounters the block. In these
experiments, UvrD or a Rep control were included either from the start of replication,
or were added after the replisome encountered the nucleoprotein block. Blocked fork
resolution was probed by examining replication products using denaturing gel
electrophoresis.
We first generated stalled transcription elongation complexes using the $P_{lacUV5\ 52C}$
promotor which allows RNAP to be stalled via nucleotide deprivation. The $P_{lacUV5\ 52C}$
promotor contains no C residues between +1 and +52 on the non-transcribed strand,
enabling transcription to be stalled by the omission of CTP. Stalled RNAP acts as a
barrier to the replisome in both head-on and co-directional orientations and prevents
the formation of full-length replication products (1). Arresting the replisome at this
RNAP block resulted in replication products consisting of four truncated leading
strands. The strand sizes matched those expected for replisomes moving clockwise
or counter-clockwise from *oriC* and encountering $P_{lacUV5\ 52C}$ or the promoters within the
ColE1 plasmid origin of replication (Figure 5A, lane 2 and Figure 5C) (1). As previously
found, inclusion of Rep or UvrD promotes movement of replisomes through stalled
RNAP (1). This is shown by a decrease in all four truncated leading strand products
and a concomitant increase in production of full-length leading strands (Figure 5A,
lanes 3 and 5 and 5E) (1). Addition of UvrD once the replisome had already
encountered the block gave variable results that ranged from an inhibitory to a stronger
effect, although differences were not statistically significant at $p < 0.05$ (Figure 5A,
compare lane 5 with 6, Figure 5E). Similar results were also obtained for Rep (Figure
5A, compare lane 3 with 4, Figure 5E). Neither Rep nor UvrD are capable of pushing
stalled RNAP from the $P_{lacUV5\ 52C}$ stall site in the absence of replication, so these
results are dependent on active replisomes encountering blocks (1).

Figure 5: Helicase pre-association affects the promotion of replication fork movement through different nucleoprotein blocks A. Denaturing agarose gel of replication products formed from pPM872 (1) in the absence and presence of RNAP block, Rep and UvrD added pre- and post-collision (*) as indicated. B. Denaturing agarose gel of replication products formed from pPM594 (12) in the absence and presence of EcoRI E111G block, Rep and UvrD added pre- and post-collision (*) as indicated. Truncated leading strand products formed by

replisomes originating from *oriC* and colliding with RNAP (C) and EcoRI E111G (D).
Quantification of lanes in (A) and (B) from three technical replicates shown in (E) and (F),
with standard error in the mean indicated and each replicate result shown as a cross. Note, we
were also able to confirm that the mGFP tag used in the UvrD microscopy experiments does
not prevent UvrD from dislodging nucleoprotein blocks in this context (see Supplementary
Figure S14).
401

Since UvrD has been shown to interact with RNAP (47–49), we reasoned that it could

be quickly recruited to an RNAP block no matter the time of addition. We used a

different nucleoprotein block that does not bind UvrD to test whether UvrD-block pre-

association is a requirement for its promotion of replication. Modified restriction

enzymes are a useful model for nucleoprotein blocks because their block strength can

be modified by varying the number of recognition sites on DNA templates. We chose

a mutant restriction enzyme (EcoRI E111G) that binds to its recognition sequence but

has reduced cleavage activity (87). EcoRI E111G blocks to replication can be removed

by Rep or UvrD helicase activity (12). We tested whether UvrD or a Rep control could

promote replication through an array of eight EcoRI sites when added before or after

initiation of replication (Figure 5B and D). Rep promotion of replication through this

barrier was unaffected by time of addition (Figure 5B, lane 3 and 4, 5F). Presumably

this is because the Rep-DnaB interaction ensures it will always be present at

replication fork blocks as it travels with the replisome. Surprisingly, adding UvrD post-

collision did not impede its promotion of replication; in fact, the resolution of these

collisions appeared to be more efficient because the proportion of full-length

replication products increased (Figure 5B, compare lane 5 with 6, 5F). Quantification

of full-length replication products as a proportion of reaction products showed that late

UvrD addition increased full-length replication products to 84% (n=6) compared to

41% (n=3) when UvrD was present before fork collision. This contrasts with the RNAP

block results where the UvrD-RNAP interaction (47–50) might be responsible for

localising UvrD to the block and facilitating resolution, resulting in no difference when

present pre- or post-collision. Similarly, Rep associates directly with the replication

fork, resulting in no difference when present pre- or post-collision, for both blocks.

These results support the absence of a UvrD-replisome interaction and suggest that

UvrD might be recruited to stalled replication forks at nucleoprotein blocks using a

different mechanism than Rep. We have no direct evidence to support a hypothesis

that UvrD recognises a specific DNA structure at a blocked fork, and note previous

studies which indicate no binding specificity to DNA beyond requiring 3´ssDNA (88,

89). Combining these data with reductions in colocalisation between UvrD and DnaQ
fork marker when replication blocks are perturbed *in vivo*, leads us to propose that
UvrD is present at the replication fork due to the high frequency of different types of
blocks to replication.
435
DISCUSSION
Here, we show that most replisomes are associated with UvrD. However, unlike the
established Rep association, UvrD has no replisome interaction partner. UvrD instead
associates with the replisome through the frequent blocks to replication encountered
by the fork. Similarly to Rep, *in vitro* evidence suggests that UvrD functions poorly as
a helicase when it is monomeric (90). We showed previously that functional Rep foci
are hexameric *in vivo* but that this oligomerisation state is due to interactions with the
hexameric DnaB replicative helicase (13). Again, UvrD functions differently and
appears to self-associate into tetramers, as evidenced in our findings by tetrameric
periodicity both at and away from the fork (Figure 2 and Supplementary Figure 11),
consistent with previous ultracentrifugation experiments *in vitro* (58). It is possible that
UvrD may not have evolved to fulfil exactly the same functions as Rep; in many
enzyme-catalysed reactions, if one enzyme is depleted then another will be available
to take its place, but with a reduced catalytic efficiency (91). On initial inspection, this
might appear to be the case for Rep and UvrD, in light of the lower efficiency of block
resolution for UvrD compared to Rep, noting that Rep deletion has a profound impact
on replication whereas UvrD deletion does not (92). However, we find that Rep and
UvrD do have different modes of colocation at a replication block. These differing
mechanisms of replication promotion suggest a model in which different accessory
helicases have evolved to fulfil the same overall function of resolving a block to DNA
replication but in alternative ways. Even with different block removal efficiency, such
redundancy may confer increased robustness, helping to ensure that DNA replication
is carried out correctly.
We used single-molecule Slimfield to show that a high proportion of forks are
colocated with UvrD *in vivo*. Although the raw imaging data is still subject to diffraction-
limited resolution, localization tracking algorithms pinpoint fluorescently labelled UvrD
to 40 nm lateral precision, smaller than the 50 nm diameter of the replisome machinery
itself (61) and equivalent to ~100 bp DNA. Thus, our measurements for UvrD and
DnaQ are good indicators of genuine colocation of UvrD and the fork. Importantly, the

degree of colocation we observed is much higher than expectations of random overlap
of the optical images of DnaQ-mCherry and mGFP-UvrD.

Using SPR analysis we failed to find evidence for direct interaction partners at the
replisome apart from the Tau protein. However, as a control, we tested Tau's
interaction with Rep and also found a positive interaction. Thus, we conclude that the
interaction of UvrD with Tau is most likely non-specific and that UvrD recruitment to
the replisome is mediated by blocks, rather than by a specific replisome interaction
partner. A practical limitation of SPR measurement is that the immobilisation
procedure does not normally preserve oligomer subunit interactions if they are not
very tight, so one would expect to observe UvrD monomers on the surface. Where
detection of an oligomer is specifically required, optimisation must be performed to
stabilise it (e.g. cross-linking) for immobilisation. It is possible at very high
immobilisation levels to have some "apparent" oligomer generated simply by the
proximity of subunits but this is not a practically useful strategy. UvrD's tetrameric form
may also affect its interactions with replisome components. These limitations mean
that SPR alone is semi-quantitative in not having the capability to report on the precise
oligomeric state of UvrD on the SPR chip.

We probed UvrD's function at the replication fork by perturbing its known DNA repair
processing functions, knocking out *mutS* and *uvrA* to block mismatch and nucleotide
excision repair respectively. We also introduced a mutant RNAP with *rpoB*35* and
treated the wild type with rifampicin to reduce the occurrence of transcriptional blocks
to translocating replication machinery. Deleting *uvrA* and rifampicin treatment
produced clear results, reduced fork colocalisation, but deleting *mutS* and introducing
the *rpoB*35* mutation did not. However, these latter perturbations did result in an
identical phenotype of one extra UvrD tetramer per fork. Indeed, a recent study has
confirmed the continuous physical association of UvrA and UvrD with RNAP. Our
results are in agreement with their model proposed that UvrA and UvrD form a pre-
TCRC that subsequently facilitates recruitment of UvrD at sites of DNA lesions.(50)
The *rpoB*35* mutation improves cell viability to increased DNA lesions (83). Taken
together, our results indicate that transcription-mediated lesions are a frequent
impediment to active replisomes resulting in transcription-replication conflicts, that
UvrD helps to clear via its association with RNAP. This intriguing link between

mutations requires further study, possibly of these dual labelled strains in response to

increased DNA lesions.

The only function of UvrD we did not probe directly was its role in RecA-mediated

recruitment. We believe the results from a *recA* knockdown would be difficult to

interpret due to its pleiotropic effects. RecA is the principal recombinase and induces

the complex SOS response resulting in the induction of UvrD expression (93, 94).

Furthermore, UvrD removes RecA from RecA-DNA complexes, thus functioning as an

antirecombinase (95–97). Dissecting these UvrD functions from other effects that a

recA mutation may have on the cell would be difficult. Applying the perturbations in

combination, i.e. double knockdowns or knockdowns plus rifampicin, would likely lead

to confusing results since too many deleterious perturbations may be lethal or change

normal cell function too radically to be biologically relevant in the intended way.

For the suite of mutant strains investigated we found small but significant

colocalisation changes in UvrD, implying that UvrD-fork colocation is due to the

frequent blocks to replication it encounters. Recent biochemical findings *in vitro* show

that blocks to replication occur very frequently (1, 19, 98) potentially accounting for our

observations here.

We show that forks stalled at transcriptional blocks can be rescued by addition of

UvrD, prior to or post-stalling. If UvrD is to help perform this function then it must be

present during the resolution of collision between the replication and transcription

machinery, however, our estimate for dwell time at the fork of just a few milliseconds

- compared to seconds for DnaQ (62) - suggests a non-negligible dissociation

constant; UvrD is not processive *per se* but undergoes rapid turnover at the fork such

that there is a high likelihood for UvrD being present at any given time, but not

necessarily the same set of molecules. Interestingly, we observed a similar dwell time

for Rep (13), despite it having a clear association partner in DnaB at the fork. However,

using a defective restriction enzyme EcoRI E111G block produced a different effect,

with late UvrD addition improving block resolution. This restriction enzyme block is

artificial and less biologically relevant but may indicate that the known specific UvrD

interaction with RNAP is important. Improved replication resolution with late addition

of UvrD is intriguing, implying increased affinity for, or activity at, pre-stalled replication

forks compared to UvrD being present as the fork stalls. Complete understanding

requires further study of replication block resolution in different conditions.

Our findings suggest that UvrD diffuses and transiently interacts with DNA either at
random, through interactions with RNAP, or interactions with a specific conformation
of the blocked replication fork. At the DNA, UvrD is predominantly tetrameric, either
pre-assembling in solution or on the DNA itself. The frequent sampling leads to UvrD
often being in the vicinity of the replisome, allowing UvrD to resolve frequent blocks to
replication. It is also possible that certain barriers to replication may be harder for Rep
to resolve, and such barriers can be more efficiently removed by UvrD, however, it is
still likely that they do not change the cell's ability to replicate the genome in light of
uvrD deletion still resulting in viable cells. Indeed, UvrD shares a high degree of
structural similarity with Rep, although the two proteins differ significantly in their amino
acid sequences (31, 99). While Rep is recruited to the fork through specific interactions
with DnaB, no such interactions were found for UvrD. Stochastic impairment of the
ability of Rep to interact with DnaB may necessitate block processing at the fork by
UvrD. In summary, our results highlight the contribution of UvrD action to facilitating
accurate genome duplication at active replisomes.

Acknowledgments

We thank Maria Chechik for help with mGFP-UvrD purification. We thank Prof. Nigel
Savery (University of Bristol) for the gift of the pETDuet-UvrD overexpression plasmid,
and Dr Chris Hill (University of York) for the donation of ^{32}P. We thank the Molecular
Interactions Laboratory in the Bioscience Technology Facility at the University of York
for technical assistance and support. Supported by BBSRC via grants BB/N006453/1
(M.L., P.M.), BB/R001235/1 (M.L.,P.M.), BB/N014863/1 (M.H.).

Author Contributions

Conceptualization: AJMW, AS, PMG, MH, MCL
Methodology: AJMW, AS, JALH, AL, CG, PMG, MH, MCL
Software: AJMW
Investigation: AJMW, AS, JALH, APD, AL, DW, CG, MH
Visualization: AJMW, AS, MH
Supervision: AJMW, AS, MH, PMG, MCL
Writing—original draft: AJMW, AS, MH, MCL

Writing—review & editing: AJMW, AS, JALH, AL, DW, PMG, MH, MCL

Declaration of interests

The authors declare no competing interests.

MATERIALS AND METHODS

Strain construction

All strains used in this study (listed in full in SI Table S1, with associated plasmids

and primers in SI Tables S2 and S3) are derivatives of the laboratory wild type strain

TB28. Tagging of *dnaQ-mCherry-<kan>* (C-terminally labelled) is described in (13).

mGFP-uvrD-<kan> (N-terminally labelled) was amplified from plasmid pAS79 (*uvrD*$^+$)

with primers oAS145 and oJGB383 having 50 bp homology to either end of the

native *uvrD* locus. The resulting PCR product had homology either side such that

recombination with the chromosome would result in integration of the PCR product at

the native locus under the control of the native promoter. Prior to integration, all PCR

products were treated with DpnI, gel purified, and introduced by electroporation into

cells expressing the lambda Red genes from the plasmid pKD46 (55). The

recombinants were selected for kanamycin resistance and screened for ampicillin

sensitivity. The colonies obtained were verified for integration by PCR and

sequencing. The *uvrD* recombinants were verified by PCR amplification using

primers oPM319 and oPM320, and sequencing using the primers oJGB417,

oJGB418, oPM407, oPM409, oPM411, and oMKG71. Where required, the

kanamycin resistance gene was removed by expressing Flp recombinase from the

plasmid pCP20 (100) to generate kanamycin sensitive strains carrying the

fluorescent protein (FP) fusions. Dual labelled strains were created by introducing

the kanamycin tagged FP alleles by standard P1 mediated transduction into single

labelled strains carrying the required FP allele after removing the linked kanamycin

marker.

Determination of cell doubling time

E. coli strains were grown overnight in LB medium at 37°C at 200 rpm shaking. The

saturated overnight cultures were diluted 100-fold into fresh LB or washed once with

1X 56 salts and diluted 100-fold in fresh 1X 56 salts with 0.2% glucose as the carbon

source. Aliquots of 100 µl each of the diluted cultures in fresh media were pipetted

into individual wells of 96 well clear flat bottom sterile microplates (Corning). The

microplates containing the diluted cultures were incubated in a BMG LABTECH

SPECTROstar Nano microplate reader at 37°C and the optical density (OD_{600})

values were recorded at defined time intervals. The time taken for the optical density

values to increase 2-fold during the exponential growth phase of the culture was

taken as the cell doubling time (SI Table S4, Supplementary Figure S1).

Protein Overexpression and Purification

mGFP-uvrD was excised from pAS79 using XhoI and SalI and ligated into pET14b

cut with XhoI, creating pAS152 encoding N-terminal histidine-tagged mGFP-UvrD.

pAS152 was used to overexpress the His-mGFP-UvrD fusion in BL21(DE3)/pLysS.

A single colony was used to inoculate 10 mL of LB containing carbenicillin and

chloramphenicol at 50 µg/mL and was incubated overnight at 37°C and 180 rpm.

The overnight culture was clarified by centrifugation at 3,220 xg and 4°C, before the

pellet was resuspended in 1 mL of LB and used to inoculate 1 L of LB containing

carbenicillin and chloramphenicol at 50 µg/mL. The culture was then incubated at

37°C and 180 rpm until an OD_{600} of 0.6, at which point it was cooled to 20°C and

IPTG was added to a 1 mM final concentration and incubation was continued at

20°C and 180 rpm overnight. Cells were pelleted by centrifugation at 3,500 xg and

4°C for 20 minutes, the pellet was resuspended in 20 mL of 50 mm Tris-HCl, pH 7.5,

at 4 °C, and 10% sucrose (w/v) before flash freezing with liquid nitrogen and storing

at -80°C.

Cells were thawed on ice and Tris pH 8.3, EGTA, EDTA, DTT, NaCl and lysozyme

were added to final concentrations of 50 mM, 0.5 mM, 5 mM, 1 mM, 200 mM and 0.2

mg/mL respectively. The suspension was incubated on ice for 30 minutes before the

addition of sodium deoxycholate to a final concentration of 0.05% (w/v) and allowed

to stand on ice for a further 30 minutes. NaCl was then added to a final concentration

of 500 mM before lysis by sonication on ice. The sample was clarified by

centrifugation at 30,600 xg and 4°C for 20 minutes and the supernatant collected.

The supernatant was subjected to polymin P precipitation, polymin P was added

dropwise to a final concentration of 0.075% (w/v) under stirring at 4°C, stirring was

continued for 10 minutes before centrifugation at 30,600g and 4°C for 20 minutes. The supernatant was collected and ammonium sulfate was added gradually to it whilst stirring at 4°C to a 50% saturation, the pellet was collected by centrifugation as previously and resuspended in binding buffer (20 mM Tris pH 8.3, 500 mM NaCl) + 5 mM imidazole. The sample was loaded onto a 5 mL HisTrap FF column that had been pre-equilibrated with binding buffer + 5 mM imidazole. The column was then washed with 15 column volumes of binding buffer + 5 mM imidazole before it was developed with a 20-column volume gradient of binding buffer + 5 to 300 mM imidazole. Peak fractions were monitored by their green colour as well as by SDS-PAGE, the peak fractions containing his-mGFP-UvrD were pooled and diluted in buffer A (20 mM Tris pH8.3, 1mM EDTA, 0.5 mM EGTA, 1mM DTT) to a conductivity equivalent to that of buffer A + 100 mM NaCl. The peak fractions were then loaded onto a 5 mL HiTrap Q column equilibrated with buffer A + 100 mM NaCl, the column was washed with 10 column volumes of buffer A + 100 mM NaCl and a gradient from 10 mM to 650 mM NaCl was developed over 20 column volumes. Peak fractions were assessed by SDS-PAGE and those containing his-mGFP-UvrD were loaded onto a HiLoad 26/60 Superdex 200 gel filtration column equilibrated with buffer B (20 mM Tris pH8.3, 300 mM NaCl), the column was developed with buffer B and peak fractions containing pure his-mGFP-UvrD as determined by SDS-PAGE were pooled and loaded onto a 5 mL HisTrap FF column that had been pre-equilibrated with binding buffer + 5 mM imidazole. The column was then washed with 3-column volumes of binding buffer + 5 mM imidazole before it was developed with a 6-column volume step of binding buffer + 200 mM imidazole. Peak fractions were then dialysed against storage buffer (20 mM Tris pH8.4, 300 mM NaCl, 10% glycerol) aliquoted and flash frozen in liquid nitrogen before storing at -80°C. Protein concentration was determined by Bradford's assay.

UvrD was purified as previously described (56). Briefly UvrD was overexpressed in BL21-AI from a pETDuet vector, via induction by IPTG and arabinose. UvrD was purified from the soluble cell extract after lysis through the use of affinity (heparin agarose), anion exchange (Q-sepharose), and gel filtration chromatography. Pure UvrD was then stored at -80ºC after dialysis into storage buffer or used immediately in Size Exclusion Chromatography - Multi-Angle Laser Light Scattering (SEC-

MALLS) analysis or subsequently diluted to 1 nM for single-molecule *in vitro*

experimentation.

Biotinylated UvrD and Rep were purified as previously described (12). Briefly, the

proteins were overexpressed in a BL21 (DE3) containing pBirAcm, encoding a biotin

ligase via induction by IPTG and the addition of biotin to the growth media. The

biotinylated proteins were then purified from the soluble cell extracts by ammonium

sulphate precipitation, and affinity chromatography (softlink avidin and heparin

agarose). Pure proteins were then dialysed into storage buffer overnight and stored

at -80°C

Helicase assay

Assays for unwinding of streptavidin-bound forks were performed using a substrate

made by annealing oligonucleotides oPM187B20 (5' end labelled with ^{32}P) and

oPM188B34. Reactions were performed in 10 μL volumes containing 40 mM HEPES

(pH 8); 10 mM DTT; 10 mM magnesium acetate; 2 mM ATP; 0.1 mg ml^{-1} BSA and

1 nM forked DNA substrate. Reactions were carried out as described earlier (13,

101). Briefly, the reaction mixture was pre-incubated at 37°C for five minutes with or

without 1 μM streptavidin (Sigma-Aldrich), to which helicase (as indicated) and biotin

(Sigma-Aldrich) were added to 100 μM (acting as a trap for free streptavidin) and

incubated at 37°C for another 10 minutes. Reactions were stopped by addition of 2.5

μl of 2.5% SDS, 200 mM EDTA and 10 mg ml^{-1} of proteinase K. Reactions were then

analysed by non-denaturing gel electrophoresis on 10% polyacrylamide TBE gels.

The quantification of the unwinding and displacement of streptavidin from the fork

was carried out as described in (102).

Preparation of cells for microscopy

Cells were grown overnight in LB to saturation. The overnight cultures were

inoculated the next day at 1:1000 dilution in 1X 56 salts supplemented with 0.2%

glucose as the carbon source. The dilutions were then grown at 30°C to mid-log

phase.

Rifampicin treatment

Cells prepared for microscopy were treated with rifampicin at a final concentration of
50 µg/mL for 30 minutes at 30°C.

Microscopy and image analysis

Imaging was performed on a bespoke dual-colour single-molecule microscope (13).
Excitation from Obis 488 nm and 561 nm wavelength 50mW lasers (run at 20 mW)
was reduced to 10 µm at full width half maximum field in the sample plane producing
Slimfield (59) producing a mean excitation intensity of ~0.25 mW/µm^2. Lasers were
digitally modulated to 5 ms period to produce alternating laser excitation using
National Instruments dynamic I/O module NI 9402. Excitation was coupled into a
Zeiss microscope body with the sample mounted on a Mad City Labs nanostage.
Images were magnified to 80nm/pixel and imaged using an Andor Ixon 128 emCCD
camera. Color channels were split red/green using a bespoke color splitter consisting
of a dual-pass green/red dichroic mirror centered at long-pass wavelength 560 nm
and emission filters with 25 nm bandwidths centered at 542 nm and 594 nm.

Cells were imaged on agarose pads suffused with minimal media as described
previously (103). Foci were detected and tracked using previously described
bespoke MATLAB software (104). In brief, bright candidate foci were detected in
each frame by image transformation and thresholding based on Otsu's method. A
17x17 pixel region of interest (ROI) is drawn around each candidate to define the
local background, while a radius 5 circle is drawn in the center (the foreground) and
subjected to iterative Gaussian masking (69). Foci were accepted if their signal to
noise ratio was above 0.4, defined as the mean intensity inside the foreground
divided by the standard deviation of the local background. Foci in successive frames
were linked together into trajectories based on nearest distance, provided it was < 5
pixels. Linked foci were accepted as "tracks" nominally if they persist for at least 4
consecutive image frames.

Characteristic intensity of mGFP or mCherry was determined from the distribution of
foci intensity values towards the end of the photobleach confirmed by overtracking
foci beyond their bleaching and applying an edge-preserving filter (105, 106) to the
raw intensity data to generate individual steps of the characteristic intensity, due to
photobleaching (Supplementary Figure S7). This intensity was used to determine the
stoichiometry of foci by fitting the first 3 intensity points with a straight line and

dividing the intercept by this characteristic intensity. The number of peaks in the Gaussian fits to UvrD was set by running a peak fitting algorithm over the wild type distribution. This number of Gaussians was then used for mutant distributions unless two or more of the Gaussians converged on the same/similar peak value, in which case they were removed. For DnaQ, two peaks were fit as used previously (61). The width of the obtained Gaussian fits was in line with previous measurements (61) and a function of several sources of noise in the challenging live-cell environment, including: fluorophores broad intensity distribution (73) (Supplementary Figure S6); camera noise; cell autofluorescence; and mobility of the protein complexes themselves.

GFP and mCherry images were aligned based on the peak of the 2D cross correlation between their respective brightfield images. Colocalisation between foci and the probability of random colocalisation was determined as described previously (65, 107).

The microscopic apparent diffusion coefficients (D) were determined by fitting a straight line to the first three mean squared displacements (MSD) values constrained through the equivalent localization precision MSD as determined from the intensity (73, 108). D distributions were fit by three Gamma distributions as described previously (109–111). Dwell time was calculated as the number of frames that each trajectory was colocalised with the fork position, as determined by the DnaQ foci detected at time zero.

Single-molecule *in vitro* imaging

10 µg/mL mouse anti-GFP (Roche. Cat no. 11814460001) in PBS was incubated on a tunnel slide for 5 min at room temperature, before washing with 100 µL of imaging buffer (20 mM Tris pH 7.8, 5 mM $MgCl_2$, 1 mM DTT). 1 nM of purified mGFP-UvrD in imaging buffer was then flowed onto the slide and incubated at room temperature for 5 min before washing with 100 µL of imaging buffer. The ends of the channel were then sealed for imaging. We used a similar single-molecule fluorescence microscope design (107) which enabled Slimfield as for the live cell imaging in order to image the antibody immobilised mGFP-UvrD.

SEC-MALLS

Size Exclusion Chromatography – Multi-Angle Laser Light Scattering (SEC-MALLS)

experiments used Wyatt HELEOS-II multi-angle light scattering and rEX refractive

index detectors linked to a Shimadzu HPLC system (SPD-20A UV detector, LC20-

AD isocratic pump system, DGU-20A3 degasser and SIL-20A autosampler) with a

G.E. Superdex S200 10/300 column at a flow rate of 0.5 mL/min in buffer of 20 mM

Tris pH 8.3 (at 4degC) 1 mM EDTA, 1 mm EGTA, 1 mM DTT, 200 mM NaCl, 5%

glycerol buffer. Data were analysed using Wyatt Astra 7 software. MWs were

estimated using the Zimm fit method with degree 1. A value of 0.182 mL/g was used

for protein refractive index increment (dn/dc).

Surface plasmon resonance (SPR)

SPR was performed at 25°C on a BIAcore T200 instrument as in (12). Immobilisation

of *E. coli* UvrD and Rep was performed onto streptavidin-coated SA sensor chips

whilst the indicated concentrations of Tau and UvrB were passed over in 10 mM

HEPES pH 7.4, 3 mM EDTA, 150 mM NaCl, 10 mM $MgCl_2$ and 0.05% Tween 20 at

20 µl min^{-1}. This buffer differed from that used in the DNA replication assays to

minimise non-specific interactions with the surface-immobilised streptavidin. We

performed experiments comparable to our previous study (12) in the range ca. 200-

4000 nM.

Replication assay

Replication assays were carried out using pPM872 as the template for RNAP blocks

(1) or pPM594 (12) for EcoRI E111G blocks. Plasmid pPM872 contains $P_{lacUV5\ 52C}$, a

strong promoter in which the first 52 nucleotides of the transcript lack cytosine

residues and are then followed by four consecutive cytosines. This enables

transcription of $P_{lacUV5\ 52C}$ to be stalled by the omission of CTP. Assays were

performed in 40 mM HEPES (pH 8); 10 mM DTT; 10 mM magnesium acetate; 2 mM

ATP; 0.2 mM GTP, 0.2 mM UTP; 0.04 mM dNTPs; and 0.1 mg/ml BSA. Reactions

(15 µl) contained 2 nM plasmid template, 50 nM DNA polymerase III αεθ complex,

25 nM τ clamp loader complex, 160 nM DnaB and DnaC monomers, 1 µM SSB, 80

nM β clamp, 30 nM HU, 200 nM DnaG, 300 nM DnaA. Helicases were added as

indicated. Rep and UvrD used at 200 nM. *E. coli* RNAP holoenzyme from NEB (1

U/µl) was used at 1/3 dilution. Dilution was determined empirically to match RNAP

replication inhibition levels from (1). EcoRI E111G was purified as in (112) and used

at 200 nM. Reactions were assembled on ice and initiated by addition of DnaA and

incubation for 4 min at 37°C, followed by addition of 60 units SmaI (Promega) plus

0.4 MBq [α^{32}P] dCTP (222 TBq/mmol). Standard reactions with the helicase present

in the initial protein mix were carried out at 37°C for 1 minute and then terminated by

addition of 1 µl of 0.5 M EDTA. Delayed helicase addition was carried out by adding

Rep/UvrD after 1 minute and incubation for a further minute at 37°C before

termination with EDTA. Ethanol precipitated replication products were analysed by

denaturing agarose gel electrophoresis (0.7% agarose in 2 mM EDTA, 30 mM NaOH

for 400 volt hours, standard run was 16 hours at 25 V), phosphorimaging and

autoradiography. 5'-labelled HindIII- digested lambda DNA was used as a marker.

Gels were quantified using Quantity One® (Bio-Rad) software. Full-length plasmid

replication products were quantified as a proportion of summed blocked products

and normalised for length (for EcoRI E111G block), or total lane signal (RNAP

block).

795
796

798
799 **REFERENCES**
800
1.   M. Hawkins, *et al.*, Direct removal of RNA polymerase barriers to replication by
accessory replicative helicases. *Nucleic Acids Res.* **47** (2019).
2.   R. T. Pomerantz, M. O'Donnell, The replisome uses mRNA as a primer after
colliding with RNA polymerase. *Nature* **456**, 762–767 (2008).
3.   H. Merrikh, C. Machón, W. H. Grainger, A. D. Grossman, P. Soultanas, Co-
directional replication-transcription conflicts lead to replication restart. *Nature*
470, 554–7 (2011).
4.   A. H. Syeda, J. Atkinson, R. G. Lloyd, P. McGlynn, The balance between
recombination enzymes and accessory replicative helicases in facilitating
genome duplication. *Genes (Basel).* **7** (2016).
5.   A. Helmrich, M. Ballarino, L. Tora, Collisions between replication and
transcription complexes cause common fragile site instability at the longest
human genes. *Mol. Cell* **44**, 966–977 (2011).
6.   Y. Takeuchi, T. Horiuchi, T. Kobayashi, Transcription-dependent
recombination and the role of fork collision in yeast rDNA. *Genes Dev.* **17**,
1497–1506 (2003).
7.   A. L. de Septenville, S. Duigou, H. Boubakri, B. Michel, Replication fork
reversal after replication-transcription collision. *PLoS Genet.* **8** (2012).

8.   C. J. Rudolph, A. L. Upton, A. Stockum, C. A. Nieduszynski, R. G. Lloyd,
Avoiding chromosome pathology when replication forks collide. *Nature* **500**,
608–611 (2013).
9.   B. Michel, G. Grompone, M. J. Florès, V. Bidnenko, Multiple pathways process
stalled replication forks. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 12783–12788
(2004).
10.   E. V. Mirkin, S. M. Mirkin, Mechanisms of transcription-replication collisions in
bacteria. *Mol. Cell. Biol.* **25**, 888–895 (2005).
11.   P. McGlynn, N. J. Savery, M. S. Dillingham, The conflict between DNA
replication and transcription. *Mol. Microbiol.* **85**, 12–20 (2012).
12.   C. P. Guy, *et al.*, Rep provides a second motor at the replisome to promote
duplication of protein-bound DNA. *Mol. Cell* **36**, 654–66 (2009).
13.   A. H. Syeda, *et al.*, Single-molecule live cell imaging of Rep reveals the
dynamic interplay between an accessory replicative helicase and the
replisome. *Nucleic Acids Res.* **47**, 6287–6298 (2019).
14.   P. McGlynn, Linking transcription with DNA repair, damage tolerance, and
genome duplication. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 15314–15315 (2010).
15.   D. Dutta, K. Shatalin, V. Epshtein, M. E. Gottesman, E. Nudler, Linking RNA
polymerase backtracking to genome instability in E. coli. *Cell* **146**, 533–543
(2011).
16.   D. Duchi, *et al.*, RNA Polymerase Pausing during Initial Transcription. *Mol. Cell*
63, 939–950 (2016).
17.   G. Ghosal, J. Chen, DNA damage tolerance: a double-edged sword guarding
the genome. *Transl. Cancer Res.* **2**, 107–129 (2013).
18.   S. Hamperl, M. J. Bocek, J. C. Saldivar, T. Swigut, K. A. Cimprich,
Transcription-Replication Conflict Orientation Modulates R-Loop Levels and
Activates Distinct DNA Damage Responses. *Cell* **170**, 774-786.e19 (2017).
19.   K. Hizume, H. Araki, Replication fork pausing at protein barriers on
chromosomes. *FEBS Lett.* **593**, 1449–1458 (2019).
20.   K. K. Myka, M. E. Gottesman, DksA and DNA double-strand break repair.
Curr. Genet. **65**, 1297–1300 (2019).
21.   K. K. Myka, *et al.*, Inhibiting translation elongation can aid genome duplication
in Escherichia coli. *Nucleic Acids Res.* **45**, 2571–2584 (2017).
22.   S. M. Mangiameli, C. N. Merrikh, P. A. Wiggins, H. Merrikh, Transcription leads
to pervasive replisome instability in bacteria. *Elife* **6** (2017).
23.   V. Kamarthapu, *et al.*, ppGpp couples transcription to DNA repair in E. coli.
Science **352**, 993–996 (2016).
24.   B. J. Brewer, When polymerases collide: replication and the transcriptional
organization of the E. coli chromosome. *Cell* **53**, 679–686 (1988).
25.   A. Srivatsan, A. Tehranchi, D. M. MacAlpine, J. D. Wang, Co-orientation of
replication and transcription preserves genome integrity. *PLoS Genet.* **6**
(2010).
26.   B. Liu, B. M. Alberts, Head-on collision between a DNA replication apparatus
and RNA polymerase transcription complex. *Science* **267**, 1131–1137 (1995).
27.   J. D. Wang, M. B. Berkmen, A. D. Grossman, Genome-wide coorientation of
replication and transcription reduces adverse effects on replication in Bacillus
subtilis. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 5608–5613 (2007).
28.   T. S. Sankar, B. D. Wastuwidyaningtyas, Y. Dong, S. A. Lewis, J. D. Wang,
The nature of mutations induced by replication–transcription collisions. *Nature*
535, 178–181 (2016).

29. D. Ivanova, *et al.*, Shaping the landscape of the Escherichia coli chromosome:
replication-transcription encounters in cells with an ectopic replication origin.
Nucleic Acids Res. **43**, 7865–7877 (2015).
30. A. Helmrich, M. Ballarino, E. Nudler, L. Tora, Transcription-replication
encounters, consequences and genomic instability. *Nat. Struct. Mol. Biol.* **20**,
412–418 (2013).
31. J. G. Brüning, J. L. Howard, P. McGlynn, Accessory replicative helicases and
the replication of protein-bound DNA. *J. Mol. Biol.* **426**, 3917–3928 (2014).
32. K. R. McDonald, *et al.*, Pfh1 Is an Accessory Replicative Helicase that
Interacts with the Replisome to Facilitate Fork Progression and Preserve
Genome Integrity. *PLoS Genet.* **12** (2016).
33. J. E. Sokoloski, A. G. Kozlov, R. Galletto, T. M. Lohman, Chemo-mechanical
pushing of proteins along single-stranded DNA. *Proc. Natl. Acad. Sci. U. S. A.*
113, 6194–6199 (2016).
34. G. D. Schauer, *et al.*, Replisome bypass of a protein-based R-loop block by
Pif1. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 30354–30361 (2020).
35. M. K. Gupta, *et al.*, Protein-DNA complexes are the primary sources of
replication fork pausing in Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **110**,
7252–7257 (2013).
36. O. Adebali, Y. Y. Chiou, J. Hu, A. Sancar, C. P. Selby, Genome-wide
transcription-coupled repair in Escherichia coli is mediated by the Mfd
translocase. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E2116–E2125 (2017).
37. N. Kang, E. Choi, S. G. Kim, J. Hwang, Inhibitory effect of UvrD and DinG on
the replication of ColE1-derived plasmids in Escherichia coli. *Plasmid* **81**, 32–
41 (2015).
38. V. Epshtein, UvrD helicase: an old dog with a new trick: how one step
backward leads to many steps forward. *Bioessays* **37**, 12–19 (2015).
39. H. Boubakri, A. L. De Septenville, E. Viguera, B. Michel, The helicases DinG,
Rep and UvrD cooperate to promote replication across transcription units in
vivo. *EMBO J.* **29**, 145–157 (2010).
40. X. Liu, J. X. Seet, Y. Shi, P. R. Bianco, Rep and UvrD Antagonize One Another
at Stalled Replication Forks and This Is Exacerbated by SSB. *ACS Omega* **4**,
5180–5196 (2019).
41. P. R. Caron, S. R. Kushner, L. Grossman, Involvement of helicase II (uvrD
gene product) and DNA polymerase I in excision mediated by the uvrABC
protein complex. *Proc. Natl. Acad. Sci. U. S. A.* **82**, 4925–4929 (1985).
42. B. Van Houten, Nucleotide excision repair in Escherichia coli. *Microbiol. Rev.*
54, 18–51 (1990).
43. A. L. Lu, S. Clark, P. Modrich, Methyl-directed repair of DNA base-pair
mismatches in vitro. *Proc. Natl. Acad. Sci. U. S. A.* **80**, 4639–4643 (1983).
44. S. W. Matson, A. B. Robertson, The UvrD helicase and its modulation by the
mismatch repair protein MutL. *Nucleic Acids Res.* **34**, 4089–4097 (2006).
45. L. Manelyte, *et al.*, The unstructured C-terminal extension of UvrD interacts
with UvrB, but is dispensable for nucleotide excision repair. *DNA Repair*
(Amst). **8**, 1300–1310 (2009).
46. A. B. Robertson, S. R. Pattishall, E. A. Gibbons, S. W. Matson, MutL-catalyzed
ATP hydrolysis is required at a post-UvrD loading step in methyl-directed
mismatch repair. *J. Biol. Chem.* **281**, 19949–19959 (2006).
47. E. J. Gwynn, *et al.*, The conserved C-terminus of the PcrA/UvrD helicase
interacts directly with RNA polymerase. *PLoS One* **8** (2013).

48. K. Sanders, *et al.*, The structure and function of an RNA polymerase
interaction domain in the PcrA/UvrD helicase. *Nucleic Acids Res.* **45**, 3875–
3887 (2017).
49. V. Epshtein, *et al.*, UvrD facilitates DNA repair by pulling RNA polymerase
backwards. *Nature* **505**, 372–377 (2014).
50. B. K. Bharati, *et al.*, Crucial role and mechanism of transcription-coupled DNA
repair in bacteria. *Nature* **604**, 152–159 (2022).
51. I. Urrutia-Irazabal, J. R. Ault, F. Sobott, N. J. Savery, M. S. Dillingham,
Analysis of the pcra-rna polymerase complex reveals a helicase interaction
motif and a role for pcra/uvrd helicase in the suppression of r-loops. *Elife* **10**
(2021).
52. B. Martinez, B. K. Bharati, V. Epshtein, E. Nudler, Pervasive Transcription-
coupled DNA repair in E. coli. *Nat. Commun.* **13** (2022).
53. N. M. Haines, Y. I. T. Kim, A. J. Smith, N. J. Savery, Stalled transcription
complexes promote DNA repair at a distance. *Proc. Natl. Acad. Sci. U. S. A.*
111, 4037–4042 (2014).
54. A. J. Smith, N. J. Savery, Effects of the bacterial transcription-repair coupling
factor during transcription of DNA containing non-bulky lesions. *DNA Repair*
(Amst). **7**, 1670–1679 (2008).
55. A. J. Smith, N. J. Savery, RNA polymerase mutants defective in the initiation of
transcription-coupled DNA repair. *Nucleic Acids Res.* **33**, 755–764 (2005).
56. J. Atkinson, *et al.*, Stimulation of UvrD helicase by UvrAB. *J. Biol. Chem.* **284**,
9612–9623 (2009).
57. R. P. Wong, N. García-Rodríguez, N. Zilio, M. Hanulová, H. D. Ulrich,
Processing of DNA Polymerase-Blocking Lesions during Genome Replication
Is Spatially and Temporally Segregated from Replication Forks. *Mol. Cell* **77**,
3-16.e4 (2020).
58. N. K. Maluf, T. M. Lohman, Self-association equilibria of Escherichia coli UvrD
helicase studied by analytical ultracentrifugation. *J. Mol. Biol.* **325**, 889–912
(2003).
59. M. Plank, G. H. Wadhams, M. C. Leake, Millisecond timescale slimfield
imaging and automated quantification of single fluorescent protein molecules
for use in probing complex biological processes. *Integr. Biol.* **1**, 602–612
(2009).
60. A. Badrinarayanan, R. Reyes-Lamothe, S. Uphoff, M. C. Leake, D. J. Sherratt,
In vivo architecture and action of bacterial structural maintenance of
chromosome proteins. *Science* **338**, 528–31 (2012).
61. R. Reyes-Lamothe, D. J. Sherratt, M. C. Leake, Stoichiometry and architecture
of active DNA replication machinery in Escherichia coli. *Science* **328**, 498–501
(2010).
62. T. R. Beattie, *et al.*, Frequent exchange of the DNA polymerase during
bacterial chromosome replication. *Elife* **6**, e21763 (2017).
63. R. Reyes-Lamothe, C. Possoz, O. Danilova, D. J. Sherratt, Independent
positioning and action of Escherichia coli replisomes in live cells. *Cell* **133**, 90–
102 (2008).
64. A. L. Payne-Dwyer, A. H. Syeda, J. W. Shepherd, L. Frame, M. C. Leake,
RecA and RecB: probing complexes of DNA repair proteins with mitomycin C
in live Escherichia coli with single-molecule sensitivity. *bioRxiv*,
2021.10.04.463068 (2021).
65. I. Llorente-Garcia, *et al.*, Single-molecule in vivo imaging of bacterial

respiratory complexes indicates delocalized oxidative phosphorylation.
Biochim. Biophys. Acta **1837**, 811–24 (2014).

66. D. A. Zacharias, J. D. Violin, A. C. Newton, R. Y. Tsien, Partitioning of lipid-
modified monomeric GFPs into membrane microdomains of live cells. *Science*
296, 913–916 (2002).

67. A. Oliva, M. Llabrés, J. B. Fariña, Comparative study of protein molecular
weights by size-exclusion chromatography and laser-light scattering. *J. Pharm.*
Biomed. Anal. **25** (2001).

68. E. Folta-Stogniew, K. R. Williams, Determination of molecular masses of
proteins in solution: Implementation of an HPLC size exclusion
chromatography and laser light scattering service in a core laboratory. *J.*
Biomol. Tech. **10** (1999).

69. M. C. Leake, *et al.*, Stoichiometry and turnover in single, functioning
membrane protein complexes. *Nature* **443**, 355–358 (2006).

70. N. K. Maluf, C. J. Fischer, T. M. Lohman, A Dimer of Escherichia coli UvrD is
the active form of the helicase in vitro. *J. Mol. Biol.* **325**, 913–935 (2003).

71. H. Yokota, Y. A. Chujo, Y. Harada, Single-molecule imaging of the oligomer
formation of the nonhexameric Escherichia coli UvrD helicase. *Biophys. J.* **104**,
924–933 (2013).

72. H. Yokota, Quantitative and kinetic single-molecule analysis of DNA unwinding
by Escherichia coli UvrD helicase. *Biophys. physicobiology* **19**, n/a (2022).

73. A. J. M. Wollman, M. C. Leake, Millisecond single-molecule localization
microscopy combined with convolution analysis and automated image
segmentation to determine protein concentrations in complexly structured,
functional cells, one cell at a time. *Faraday Discuss.* **184**, 401–24 (2015).

74. A. H. Syeda, *et al.*, Single-molecule live cell imaging of Rep reveals the
dynamic interplay between an accessory replicative helicase and the
replisome. *Nucleic Acids Res.* **47**, 6287–6298 (2019).

75. A. J. Wollman, *et al.*, Transcription factor clusters regulate genes in eukaryotic
cells. *Elife* **6** (2017).

76. S. Shashkova, A. J. M. Wollman, M. C. Leake, S. Hohmann, The yeast Mig1
transcriptional repressor is dephosphorylated by glucose-dependent and -
independent mechanisms. *FEMS Microbiol. Lett.* **364**, 245 (2017).

77. H. M. Arthur, P. B. Eastlake, Transcriptional control of the uvrD gene of
Escherichia coli. *Gene* **25**, 309–316 (1983).

78. Y. Taniguchi, *et al.*, Quantifying E. coli proteome and transcriptome with
single-molecule sensitivity in single cells. *Science* **329**, 533–8 (2010).

79. G. W. Li, D. Burkhardt, C. Gross, J. S. Weissman, Quantifying absolute protein
synthesis rates reveals principles underlying allocation of cellular resources.
Cell **157**, 624–635 (2014).

80. J. Atkinson, M. K. Gupta, P. McGlynn, Interaction of Rep and DnaB on DNA.
Nucleic Acids Res. **39**, 1351–1359 (2011).

81. S. V. Rajagopala, *et al.*, The binary protein-protein interaction landscape of
Escherichia coli. *Nat. Biotechnol. 2014 323* **32**, 285–290 (2014).

82. D. K. Orren, C. P. Selby, J. E. Hearst, A. Sancar, Post-incision steps of
nucleotide excision repair in Escherichia coli. Disassembly of the UvrBC-DNA
complex by helicase II and DNA polymerase I. *J. Biol. Chem.* **267**, 780–788
(1992).

83. P. McGlynn, R. G. Lloyd, Modulation of RNA polymerase by (p)ppGpp reveals
a RecG-dependent mechanism for replication fork progression. *Cell* **101**, 35–

45 (2000).

84.   Y. N. Zhou, D. J. Jin, The rpoB mutants destabilizing initiation complexes at
1021   stringently controlled promoters behave like "stringent" RNA polymerases in
1022   Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 2908–2913 (1998).

1023   85.   E. A. Campbell, *et al.*, Structural mechanism for rifampicin inhibition of
1024   bacterial rna polymerase. *Cell* **104**, 901–912 (2001).

1025   86.   M. Stracy, *et al.*, Live-cell superresolution microscopy reveals the organization
1026   of RNA polymerase in the bacterial nucleoid. *Proc. Natl. Acad. Sci. U. S. A.*
1027   **112**, E4390-9 (2015).

1028   87.   K. King, S. J. Benkovic, P. Modrich, Glu-111 is required for activation of the
1029   DNA cleavage center of EcoRI endonuclease. *J. Biol. Chem.* **264**, 11807–
1030   11815 (1989).

1031   88.   J. A. Ali, N. K. Maluf, T. M. Lohman, An oligomeric form of E. coli UvrD is
1032   required for optimal helicase activity. *J. Mol. Biol.* **293** (1999).

1033   89.   E. Curti, S. J. Smerdon, E. O. Davis, Characterization of the helicase activity
1034   and substrate specificity of Mycobacterium tuberculosis UvrD. *J. Bacteriol.* **189**
1035   (2007).

1036   90.   B. Nguyen, Y. Ordabayev, J. E. Sokoloski, E. Weiland, T. M. Lohman, Large
1037   domain movements upon UvrD dimerization and helicase activation. *Proc.*
1038   *Natl. Acad. Sci.* **114**, 12178–12183 (2017).

1039   91.   A. Wagner, Robustness against mutations in genetic networks of yeast. *Nat.*
1040   *Genet. 2000 244* **24**, 355–361 (2000).

1041   92.   J. Atkinson, *et al.*, Localization of an accessory helicase at the replisome is
1042   critical in sustaining efficient genome duplication. *Nucleic Acids Res.* **39**, 949–
1043   957 (2011).

1044   93.   S. C. Massoni, S. J. Sandler, Specificity in suppression of SOS expression by
1045   recA4162 and uvrD303. *DNA Repair (Amst).* **12**, 1072–1080 (2013).

1046   94.   K. H. Maslowska, K. Makiela-Dzbenska, I. J. Fijalkowska, The SOS system: A
1047   complex and tightly regulated response to DNA damage. *Environ. Mol.*
1048   *Mutagen.* **60**, 368–384 (2019).

1049   95.   R. C. Centore, S. J. Sandler, UvrD limits the number and intensities of RecA-
1050   green fluorescent protein structures in Escherichia coli K-12. *J. Bacteriol.* **189**,
1051   2915–2920 (2007).

1052   96.   V. Petrova, *et al.*, Active displacement of RecA filaments by UvrD translocase
1053   activity. *Nucleic Acids Res.* **43**, 4133–4149 (2015).

1054   97.   M. M. Cox, Regulation of bacterial RecA protein function. *Crit. Rev. Biochem.*
1055   *Mol. Biol.* **42**, 41–63 (2007).

1056   98.   N. Raghunathan, S. Goswami, J. K. Leela, A. Pandiyan, J. Gowrishankar, A
1057   new role for Escherichia coli Dam DNA methylase in prevention of aberrant
1058   chromosomal replication. *Nucleic Acids Res.* **47**, 5698–5711 (2019).

1059   99.   H. Yokota, Roles of the C-Terminal Amino Acids of Non-Hexameric Helicases:
1060   Insights from Escherichia coli UvrD. *Int. J. Mol. Sci.* **22**, 1–19 (2021).

1061   100.   K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in
1062   Escherichia coli K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97**,
1063   6640–6645 (2000).

1064   101.   J. G. Brüning, J. A. L. Howard, K. K. Myka, M. S. Dillingham, P. McGlynn, The
1065   2B subdomain of Rep helicase links translocation along DNA with protein
1066   displacement. *Nucleic Acids Res.* **46**, 8917–8925 (2018).

1067   102.   J. G. Brüning, J. A. L. Howard, P. McGlynn, Use of streptavidin bound to
1068   biotinylated DNA structures as model substrates for analysis of nucleoprotein

complex disruption by helicases. *Methods* **108**, 48–55 (2016).

103. A. J. M. Wollman, A. H. Syeda, P. McGlynn, M. C. Leake, Single-Molecule Observation of DNA Replication Repair Pathways in E. coli. *Adv. Exp. Med. Biol.* **915**, 5–16 (2016).

104. H. Miller, *et al.*, Superresolution imaging of single DNA molecules using stochastic photoblinking of minor groove and intercalating dyes. *Methods* **88**, 81–8 (2015).

105. M. C. Leake, D. Wilson, B. Bullard, R. M. Simmons, The elasticity of single kettin molecules using a two-bead laser-tweezers assay. *FEBS Lett.* **535**, 55–60 (2003).

106. M. C. Leake, D. Wilson, M. Gautel, R. M. Simmons, The elasticity of single titin molecules using a two-bead optical tweezers assay. *Biophys. J.* **87**, 1112–35 (2004).

107. A. J. M. Wollman, *et al.*, Critical roles for EGFR and EGFR-HER2 clusters in EGF binding of SW620 human carcinoma cells. *J. R. Soc. Interface* **19** (2022).

108. M. C. Leake, *et al.*, Variable stoichiometry of the TatA component of the twin-arginine protein transport system observed by in vivo single-molecule imaging. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 15376–81 (2008).

109. H. Miller, *et al.*, High-Speed Single-Molecule Tracking of CXCL13 in the B-Follicle. *Front. Immunol.* **9**, 1073 (2018).

110. J. Cosgrove, *et al.*, B cell zone reticular cell microenvironments shape CXCL13 gradient formation. *Nat. Commun.* **11** (2020).

111. X. Jin, *et al.*, Membraneless organelles formed by liquid-liquid phase separation increase bacterial fitness. *Sci. Adv.* **7** (2021).

112. S. C. Cheng, R. Kim, K. King, S. H. Kim, P. Modrich, Isolation of gram quantities of EcoRI restriction and modification enzymes from an overproducing strain. *J. Biol. Chem.* **259**, 11571–11575 (1984).

113. S. H. Chung, R. A. Kennedy, Forward-backward non-linear filtering technique for extracting small biological signals from noise. *J. Neurosci. Methods* **40**, 71–86 (1991).

Supplementary Information

Tetrameric UvrD helicase is located at the _E. coli_ replisome due to frequent replication blocks

Adam J. M Wollman[1,2,3,], Aisha H. Syeda[1,2], Jamieson A. L. Howard[1,2], Alex Payne-Dwyer[1,2], Andrew Leech[4] , Dominika Warecka[2], Colin Guy[5, 6], Peter McGlynn[2,6], Michelle Hawkins[2] and Mark C. Leake[1,2]

The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First Authors

[1] School of Physics, Engineering and Technology, University of York, York YO10 5DD, United Kingdom.

[2] Department of Biology, University of York, York YO10 5DD, United Kingdom.

[3] Current address: Biosciences Institute, Newcastle University, NE1 7RU, United Kingdom.

[4] Bioscience Technology Facility, Department of Biology, University of York, York YO10 5DD, United Kingdom

[5] Current address: Covance Laboratories Ltd., Otley Road, Harrogate, HG3 1PY, United Kingdom

[6] Previous address: School of Medical Sciences, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, United Kingdom

* To whom correspondence should be addressed. To whom correspondence should be addressed. Tel: +44 (0)1904322697. Email: mark.leake@york.ac.uk

Present Address: School of Physics, Engineering and Technology, and Department of Biology, University of York, York YO10 5DD, United Kingdom

1126 **Supplementary Tables**

1127 **Table S1 Strains used in this study**

Strain	Relevant genotype	Source or derivation
BW25113 derivatives:		
BW25113	*rrnB ΔlacZ4787 hsdR514 Δ(araBAD)567 Δ(rhaBAD)568 rph-1*	(1)
JW2703	*ΔmutS::<kan>*	(1)
JW3786	*ΔuvrD::<kan>*	(1)
JW4019	*ΔuvrA::<kan>*	(1)
MG1655 derivatives:		
TB12	*ΔlacIZYA::<kan>*	
TB28	*ΔlacIZYA*	(2, 3)
TB28 derivatives		
N6577	*ΔlacIZYA Δrep::cat*	(4)
N6661	*ΔlacIZYA::<> Δrep::dhfr* / pAM407 (*uvrD+ lacZ+*, a pRC7 derivative)	(4)
N6661	*ΔlacIZYA::<> Δrep::dhfr* / pAM407 (*uvrD+ lacZ+*, a pRC7 derivative)	(4)
N6644	*ΔlacIZYA::<> uvrD::dhfr Δrep::cat* / pAM407 (*uvrD+ lacZ+*, a pRC7 derivative)	(4)
JGB166	*ΔuvrD* / pKD46	JW3786 transformed with pKD46
AS97	TB28/pKD46	(5)
AS446	*dnaQ-mCherry-<kan>*	
AS448	*dnaQ-mCherry-<>*	(5)
AS512	*mGFP-uvrD-<kan>*	*mGFP-uvrD-<kan>* recombineered into JGB166
AS517	*ΔlacIZYA::<> Δrep::cat mGFP-uvrD-<kan>* / pAM407 (*uvrD+ lacZ+*, a pRC7 derivative)	N6661 X P1(AS512)
AS564	*dnaQ-mCherry-<kan> rpoB*35*	PM486 X P1(AS446)
AS571	*dnaQ-mCherry-<> rpoB*35*	AS564 to km^S with pCP20
AS580	*dnaQ-mCherry-<> mGFP-uvrD-<kan>*	AS448 X P1(AS512)
AS582	*dnaQ-mCherry-<> mGFP-uvrD-<kan> rpoB*35*	AS571 X P1(AS512)
AS590	*dnaQ-mCherry-<> mGFP-uvrD-<kan> Δrep::cat*	AS580 X P1(N6577)
AS650	*dnaQ-mCherry-<> mGFP-uvrD-<>*	AS580 to km^S with pCP20
AS656	*dnaQ-mCherry-<> mGFP-uvrD-<> ΔuvrA::<kan>*	AS650 X P1(JW4019)
AS658	*dnaQ-mCherry-<> mGFP-uvrD-<> ΔmutS::<kan>*	AS650 X P1(JW2703)

1128

1129

1130 **Table S2 Plasmids used in this study**

Plasmid	Description	Antibiotic	Reference
pCP20	Yeast Flp recominase expression plasmid	Amp	(6)
pKD46	λ Red recombinase expression plasmid	Amp	(6)
pAS79	pUC18 *mGFP-uvrD-<kan>*	Amp, kan	This study
pAS152	pET14b-mGFP uvrD	Amp	This study

1131
1132

1133 **Table S3 Oligonucleotides used in this study**

Primer	Sequence (5' - 3')
oJGB383	TGAATGATTTTTTAGGCCAAATAAGGTGCGCAGCACCGCATCCGGCAACGCATA TGAATATCCTCCTTAG
oJGB417	TACAAGACACGTGCTGAAGTC
oJGB418	TGCTAGTTGAACGCTTCCATC
oMKG71	CGGTGCCCTGAATGAACTGC
oPM319	CTTGTTGGATCAGACCGGAAAATG
oPM320	TGGCAACGCTATCCTTTTGTCA
oPM407	GCGGCAGGCAATGTGGTA
oPM409	CCTTTGAGCGTGTGGTGA
oPM411	CAGCGGCTACAAGCTCGG
oPM187B20	GTCGGATCCTCTAGACAGC(BiodT)CCATGATCACTGGCACTGGTAGAATTCGGC
oPM188B34	AACGTCATAGACGATTACATTGCTACATGGAGC(BiodT)GTCTAGAGGATCCGAC
oAS84	CCCGTCTCGATCTGGTGCAG
oAS85	TTGCTGCAAAAATCGCCCAAG
oAS132	CCCGTCTCGATCTGGTGCAGAAGAAAGGCGGAAGTTGCCTCTGGCGAGCAGGC TGGCTCCGCTGCTGG
oAS133	TTGCTGCAAAAATCGCCCAAGTCGCTATTTTTAGCGCCTTTCACAGGTATCATAT GAATATCCTCCTTAG
oAS145	AGGGCGTTGCGCTTCTCCGCCCAACCTATTTTTACGCGGCGGTGCCAATGAGTA AAGGAGAAGAACTT

1134

1135

1136 **Table S4 Cell doubling times of labelled strains.** Values expressed are the means of three
1137 independent replicates, with standard deviation and standard errors indicated.

Strain (Relevant genotype)	LB medium			Minimal medium		
	Mean doubling time (min)	SD (min)	SEM (min)	Mean doubling time (min)	SD (min)	SEM (min)
WT	23.72	1.11	0.64	62.80	17.06	9.85
dnaQ-mCherry	26.85	0.91	0.52	62.22	18.63	10.76
mGFP-uvrD	24.79	0.37	0.22	59.45	1.80	1.04
dnaQ-mCherry mGFP-uvrD	25.27	1.51	0.87	71.03	7.59	4.38
ΔmutS	24.59	0.55	0.32	107.91	13.96	8.06
ΔuvrA	27.99	0.23	0.13	92.25	5.86	3.38
ΔrpoB*35	33.98	0.17	0.10	83.11	30.83	17.80
Δrep	24.30	2.03	1.17	62.74	1.20	0.70
dnaQ-mCherry mGFP-uvrD ΔmutS	26.85	0.68	0.39	110.84	34.91	20.16
dnaQ-mCherry mGFP-uvrD ΔuvrA	24.13	0.56	0.33	91.80	5.18	2.99
dnaQ-mCherry mGFP-uvrD ΔrpoB*35	32.81	0.70	0.40	99.11	78.68	45.42
dnaQ-mCherry mGFP-uvrD Δrep	26.54	1.90	1.10	62.17	2.98	1.72

1138
1139

1140 **Table S5 Comparison of wild type and labelled UvrD**
1141

Method	Figure	Result
Growth rates	S1	No significant difference
Plasmid loss assays	S2	mGFP-UvrD can rescue cells where UvrD deletion is lethal
In vitro unwinding assay	S3	Nucleoprotein block removal and helicase activity of mGFP-UvrD and biotinylated UvrD was comparable to the wild type
SEC-MALLS	S9	No significant detectable difference in oligomeric state
In vitro replication assay	Fig. 5 and S14	Wild type UvrD and mGFP-UvrD are able to overcome blocks to replication

1149 **Supplementary figures**

1150

1151

1152

Supplementary Figure S1. Growth curves in LB medium. Cultures were grown in LB as

described in the text. The OD_{600} values are plotted on a log scale on the vertical axis with a

linear scale of time in hours on the horizontal axis. The genotypes of the strains are indicated

in boxes within the figures. SD error bars (shown just on every 5th consecutive point for clarity

here), taken from N=3 replicate cultures. Plots in panel A are growth curves of single and dual

labelled wild-type strains, while those in panel B are dual labelled and unlabelled mutant

strains. The unlabelled wild-type strain is included in all panels as a reference.

1160

| uvrD⁺/ Δrep uvrD⁺ (N6577) | uvrD⁺/ Δrep ΔuvrD (N6644) | uvrD⁺/ Δrep mGFP- uvrD-<kan>/ (AS517) |

0.91 (1120/1228)   0.004 (17/4737)   0.66 (730/1099)

1161

Supplementary Figure S2: Testing of mGFP-UvrD fusions for retention of function. We

transformed the strain carrying a chromosomal *mGFP-uvrD* allele with a derivative of pRC7

carrying a wild-type *uvrD* allele (pPM407), which is an unstable low copy plasmid that also

carries the *lacZYA* genes (3). Presence of this plasmid confers blue colour to the colonies on

plates containing Xgal when strains are chromosomally deleted for the *lac* operon. In rapidly

growing cells, cells require either *rep* or *uvrD* for viability, and the loss of both is lethal (4, 7).

Thus, *Δrep* cells with a functional *uvrD* allele are viable and can lose pRC7*uvrD*, evidenced

by appearance of white colonies on LB Xgal IPTG plates, but cells lacking *uvrD* function

cannot survive upon loss of the plasmid resulting in recovery of only blue colonies. The *Δrep*

mGFP-uvrD cells produced white colonies on Xgal media, indicating that the *mGFP-uvrD*

fusion is functional.

1173

1174

Supplementary Figure S3: mGFP-UvrD fusion is functional *in vitro*. (A)(i) Representative native polyacrylamide TBE gel showing UvrD, mGFP-UvrD and biotinylated-UvrD unwinding of forked DNA containing biotin on both strands within the duplex region (8, 9). Lanes 1-2 contain markers indicating the position of single stranded or forked DNA as indicated. Lanes 3-14 contain the products of unwinding the forked DNA by the indicated helicase at 1, 10, 50 and 100 nM. (ii) Quantification of the unwinding of the forked substrate by the indicated helicases. Data points shown are averages of band intensities (taken from two gels) with standard errors indicated by bars on the points. (B)(i) Representative native polyacrylamide TBE gels showing UvrD, mGFP-UvrD and biotinylated-UvrD unwinding of forked DNA containing biotin on both strands within the duplex region (8, 9). Lanes 1-4 contain markers indicating the position of single stranded or forked DNA +/- streptavidin as indicated. Lanes 5-16 contain the products of unwinding the forked DNA +/- streptavidin by the indicated helicase at 1, 10, 50 and 100 nM. (ii) Quantification of the unwinding of the forked, streptavidin bound substrate by the indicated helicases. Data points shown are averages of band intensities (taken from two gels) with standard errors indicated by bars on the points.

1190

Supplementary Figure S4: Cells typically appear to have two DnaQ foci, but more rarely show just a single DnaQ focus corresponding to two forks separated by less than the optical resolution limit. A. and B. Dual color Slimfield of mGFP-UvrD:DnaQ-mCherry, scale bar 1 micron. Segmented outlines of cell bodies (white dash) and detected foci (white circles) indicated. C. Mean corrected intensity line profile through a colocalized foci in B.

1197

1198

1199 **Supplementary Figure S5: H-NS associates with the nucleoid but not specifically**

1200 **with the fork.** A. Dual color Slimfield of H-NS-mGFP-:DnaQ-mCherry, scale bar 1 micron.

1201 Segmented outlines of cell bodies indicated (white dash). B. Histogram showing the number

1202 of H-NS foci detected per cell, SD errors.

1203

1204

1205

Supplementary Figure S6: Brightness of single mGFP and mCherry molecules.

Characteristic intensity distributions rendered as kernel density estimates of single A. mGFP-

UvrD and B. DnaQ-mCherry obtained using Slimfield. Peak±full width at half maximum

indicated. Distributions calculated from the tracked foci intensity distributions from the end of

the photobleach process such that only single fluorophore molecules are detected. Number

of molecules per foci before bleaching is determined by dividing the initial foci intensity by

these values for the equivalent fluorophore.

1213

1215 UvrD-GFP full dynamic range   UvrD-GFP foci
1216
Supplementary Figure S7: Single-molecule image analysis. A. Foci intensity as a
function of time for detected UvrD tracks (blue) overlaid by Chung-Kennedy (CK) filtered
intensity (squares). CK filtering is an edge preserving filter allowing noise to be filtered out
while retaining steps.(113) Examples are shown of intensity traces for single UvrD tracks
showing tetramers exhibiting multiple photobleach steps and monomers detected towards
the end of photobleaching. Foci were 'overtracked' beyond the last detected bright frame to
illustrate steps. **B**. UvrD-GFP micrograph showing the full dynamic range (left) and adjusted
to illustrate a foci (right)
1225

1229

Supplementary Figure S8: DnaQ tracking characterization. Top row: Distribution of
DnaQ foci stoichiometry as a kernel density estimate for all strains and conditions (red) with
two Gaussian fit overlaid as black dotted lines. Peak ± half width at half maximum shown
above. Middle row: Corresponding histogram of number of foci detected per cell. Bottom
row: histogram of the total amount of DnaQ in foci per cell. N=30-50 cells per strain or
condition.

1237

Supplementary Figure S9: SEC-MALLS of UvrD/mGFP UvrD   Size Exclusion
Chromatography - Multi-Angle Laser Light Scattering (SEC-MALLS) of 4 μM UvrD (red) and
mGFP-UvrD (blue) cumulative weight fraction as a function of molar mass in number of
UvrD/mGFP-UvrD, shading indicates 16% indicative error bounds consistent with sensitivity
error measurements reported from polydispersed heterogenous samples using SEC-MALLS
(68), and consistent with our own estimation of sensitivity using our SEC-MALLS system.
Inset shows the normalised UV absorbance as a function of elution time through the column
with the monomer peaks indicated.

1246

Slimfield of surface-immobilized mGFP-UvrD

A

B

Supplementary Figure S9: Single-molecule microscopy of purified mGFP-UvrD

indicates most foci contain monomeric mGFP-UvrD. A. Representative Slimfield of

surface-immobilized mGFP-UvrD. B. Automated single-particle tracking output for mGFP-

UvrD showing probability distribution for mGFP-UvrD foci brightness normalized by the

single-molecule dye brightness, peak (±HWHM) corresponding to 1 molecule indicated

(arrow) for n=7,384 foci. *Inset*: example overlaid stepwise photobleach traces for four

different mGFP-UvrD foci indicating a consistent single-molecule step to zero brightness

whose size is comparable to the measured modal brightness across all foci (dots: raw data,

lines: Chung-Kennedy edge-preserving filter output)**.**

1261
1262

1263
Supplementary Figure S10: A. Boxplot of the total number of mGFP-UvrD molecules per
cell, estimated by numerical integration of the whole cell fluorescence. Median is shown as
red line, bottom and top of the blue box mark the 25th and 75th percentiles and whiskers
extend to the most extreme points not considered outliers (2.7 standard deviations covering
99.3% of normally distributed data) with outliers beyond this shown as red +. N=30 cells per
strain. B Number of trajectories per cell used in the stoichiometry determination
1270

1271

1272

Supplementary Figure S11: Surface plasmon resonance measurements of helicase
interactions with Tau A.   Titration response sensorgrams for UvrD:UvrB, UvrD:Tau and
Rep:Tau with indicated analyte concentrations. B.   Binding affinity plotted as association
constant (K_a, uM-1) for the UvrD:UvrB, UvrD:Tau and Rep:Tau interactions assuming a single
sites binding model.   The respective dissociation constants K_d are 231 nM, 1.4 µM and
2.3 µM.SD error bars.

1279

1280   **Supplementary Figure S12:** Perturbed UvrD stoichiometry. The distribution of UvrD foci

1281   stoichiometry rendered as a kernel density estimate for foci colocalised (Left) and non-

1282   colocalised (Right) with DnaQ foci. Distributions were fitted with multiple Gaussian fits (black

1283   dotted lines) with the peak value $\pm$ half width at half maximum indicated above each peak.

1284

1285

1286

1287

Supplementary Figure S13: UvrD perturbed dynamics. A-E Left, distribution of time UvrD

and DnaQ foci were colocalised (blue) fit with an exponential (red) to yield a characteristic

dwell time. Right, distribution of microscopic diffusion coefficients of UvrD foci (grey) fitted with

a three-state gamma distribution model containing a relatively immobile population with

$D = 0.1$ μm²/s (red), a transiently immobile population, $D = 0.5$ μm²/s (blue) and a mobile

population, $D = 1.2$ μm²/s (green). Errors represent 95% confidence intervals. N~300

foci/condition.

1295
1296 **Supplementary Figure S14:** In vitro replication assay using mGFP-UvrD. Denaturing

1297 agarose gel of replication products formed from pPM872 (1) in the absence and presence of

1298 EcoRI E111G block, mGFP-UvrD and UvrD added pre- and post-collision (*) as indicated.

1299

Supplementary references

1. Baba,T., Ara,T., Hasegawa,M., Takai,Y., Okumura,Y., Baba,M., Datsenko,K.A., Tomita,M., Wanner,B.L. and Mori,H. (2006) Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.*, **2**, 2006.0008.

2. Bachmann,B.J. (1996) Derivations and genotypes of some mutant derivatives of Escherichia coli K-12. . In Neidhardt,F.C., Curtiss III,R., Ingraham,J.L., Lin,E.C.C., Low,K.B., Magasanik,B., Reznikoff,W.S., Riley,M., Schaechter,M., Umbarger,H.E. (eds), *Escherichia coli and Salmonella cellular and molecular biology*. ASM Press, Washington, DC, pp. 2460–2488.

3. Bernhardt,T.G. and de Boer,P.A.J. (2004) Screening for synthetic lethal mutants in Escherichia coli and identification of EnvC (YibP) as a periplasmic septal ring factor with murein hydrolase activity. *Mol. Microbiol.*, **52**, 1255–1269.

4. Guy,C.P., Atkinson,J., Gupta,M.K., Mahdi,A.A., Gwynn,E.J., Rudolph,C.J., Moon,P.B., van Knippenberg,I.C., Cadman,C.J., Dillingham,M.S., *et al.* (2009) Rep provides a second motor at the replisome to promote duplication of protein-bound DNA. *Mol. Cell*, **36**, 654–666.

5. Syeda,A.H., Wollman,A.J.M., Hargreaves,A.L., Howard,J.A.L., Brüning,J.-G., McGlynn,P. and Leake,M.C. (2019) Single-molecule live cell imaging of Rep reveals the dynamic interplay between an accessory replicative helicase and the replisome. *Nucleic Acids Res.*, **47**, 6287–6298.

6. Datsenko,K.A. and Wanner,B.L. (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA*, **97**, 6640–6645.

7. Boubakri,H., de Septenville,A.L., Viguera,E. and Michel,B. (2010) The helicases DinG, Rep and UvrD cooperate to promote replication across transcription units in vivo. *EMBO J.*, **29**, 145–157.

8. Brüning,J.-G., Howard,J.A.L. and McGlynn,P. (2016) Use of streptavidin bound to biotinylated DNA structures as model substrates for analysis of nucleoprotein complex disruption by helicases. *Methods*, **108**, 48–55.

9. Brüning,J.-G., Howard,J.A.L., Myka,K.K., Dillingham,M.S. and McGlynn,P. (2018) The 2B subdomain of Rep helicase links translocation along DNA with protein displacement. *Nucleic Acids Res.*, **46**, 8917–8925.