1
2 Variation in self-compatibility among genotypes and across ontogeny in a self-fertilizing
3 vertebrate
4
5

6 Authors: Jennifer D. Gresham (Emory University), Anna Clark (University of Alabama), Chloe
7 M.T. Keck (University of Alabama), Alexis E. Longmire (University of North Carolina, Chapel
8 Hill), Abye E. Nelson (University of Alabama), Haylee M. Quertermous (University of
9 Alabama), Ashley B. White (University of Alabama), and Ryan L. Earley (University of
10 Alabama)
11
12 Corresponding author: Jennifer D. Gresham, jdgresham20@gmail.com and 256-998-1747

13
14

15   Abstract

Mixed mating strategies offer the benefits of both self-fertilizing one's own eggs (selfing) and

outcrossing, while limiting the costs of both methods. The economics of mixed mating is further

determined by individual self-compatibility. In gynodioecious (hermaphrodites, females) and

androdioecious (hermaphrodites, males) species, the level of self-compatibility of the

hermaphrodites also acts as a selection pressure on the fitness of the other sex. Mangrove rivulus

fish populations are comprised of selfing hermaphrodites and males that result from

hermaphrodites changing sex. Although hermaphrodites overwhelmingly reproduce through

internal selfing, they occasionally oviposit unfertilized eggs. Males can externally fertilize these

eggs. Here, we reveal that fecundity and self-compatibility varies within individuals across

ontogeny until about 365 days post hatch, and among individuals derived from lineages that vary

in their propensity to change sex. Hermaphrodites from high sex changing lineages were

significantly less fecund and self-compatible than hermaphrodites from low sex changing

lineages. These differences in self-compatibility and fecundity have the potential to drive

evolutionary changes on mating strategy and the fitness of males in populations of the mangrove

rivulus. This study also illustrates the importance of including lineage variation when estimating

the costs and benefits of mixed mating strategies.

Introduction

33              Introduction

34  As a reproductive strategy, self-fertilization has many benefits. It overcomes the "cost of

35 males," wherein only one half of the population can produce offspring compared to the entire

36 population in asexual or self-fertilizing species (Maynard Smith 1971, 1978). Self-fertilization

37 also assures reproduction when pollinators and/or mates are scarce (Darwin 1876; Baker 1955;

38 Stebbins 1957; Pannell et al. 2015), eliminates physical damage associated with mating and/or

39 sexually transmitted parasites or pathogens (Fowler and Partridge 1989; Clutton-Brock and

40 Parker 1995), and maintains well-adapted genotypes in stable environments (Allard et al. 1972;

41 Dolgin et al. 2007). However, selection for selfing is often counteracted by selection for

42 outcrossing. Benefits of outcrossing include combatting inbreeding depression in two ways: i)

43 preventing accumulation of deleterious alleles in single genotypes, and ii) maintaining

44 heterozygotes that may be more fit than homozygous individuals (Muller 1932). Ultimately,

45 outcrossing represents a bet hedging strategy because it increases the variability of offspring

46 fitness against a changing abiotic environment, or against coevolving parasites, predators and

47 competitors (Muller 1932; Bell 1988; Lively 2010; Morran et al. 2011; Hartfield and Keightley

48 2012).

49  Many plant and invertebrate species have evolved a mixed mating strategy of selfing and

50 outcrossing. Mixed mating allows individuals and populations to reap the rewards of both

51 strategies as biotic and abiotic factors change the fitness landscape across space and time.

52 Because the fitness landscape is variable, it would be expected that self-compatibility also would

53 vary both within and among populations due to changes in the magnitude and/or type of

54 selection acting for and against selfing and outcrossing (Escobar et al. 2011). Plant populations

55 that are often mate- and/or pollinator-limited at the range edge or in ephemeral habitats have

increased self-compatibility compared to other populations, as reported for the small angiosperm

Leavenworthia alabamica (Busch 2005). Populations found near the center of the range and in

more permanent habitats are much more likely to be self-incompatible (Busch 2005). When

among-individual variation in self-compatibility is heritable, it provides the genetic and

phenotypic background on which selection can drive evolutionary change in mating strategies.

There are multiple examples of heritable variation in the self-compatibility of plants, including in

the creeping bellflower, *Campanula rapunculoide* (Stephenson et al. 2000), and the Carolina

horsenettle *Solanum carolinense* (Travers et al. 2004). Genetic and phenotypic variation in self-

compatibility, along with environmental variability, can thus drive variation in mating strategy,

as they alter the economics of selfing and outcrossing.

In mixed mating populations, self-compatibility can also vary across ontogeny, perhaps

as a plastic response to environmental conditions. Individuals in populations that have higher

rates of outcrossing often become more self-compatible with increasing age. This is often the

result of selection favoring a "waiting" strategy for outcrossing opportunities when individuals

are young, presumably to derive the benefits of variable and more fit progeny but also to assure

some reproduction even if mates and/or pollinators are limited. This has been documented in

plants (*Campanula rapunculoide*; Stephenson, Good, and Vogler 2000, *Solanum carolinense;*

Travers, Mena-Ali, and Stephenson 2004), and snails (*Physa acute*; Tsitrone, Jarne, and David

2003), but few other self-fertilizing species. Populations can also evolve different mating

strategies based on age structure. If younger individuals are more populous, and mating

opportunities are plentiful enough to make selfing unnecessary, age-dependent changes in self-

compatibility may be lost so that little or no selfing occurs. Alternatively, if populations consist

predominantly of older individuals, selfing may continue to be an evolutionary stable component

of a mixed mating strategy.

In androdioecious species, in which populations consist of hermaphrodites and males,

variation in self-compatibility should also result in variation in the proportion of males within a

population. Populations consisting of individuals with lower self-compatibility should maintain a

higher proportion of males, as the "cost of males" is decreased by more outcrossing

opportunities. Hypotheses concerning the evolution and variation of self-compatibility and

mixed mating have been examined in many plant and invertebrate systems (Busch 2005; Winn et

al. 2011; Wright et al. 2013), but not in a vertebrate species. Here we present evidence for

individual variation in self-compatibility in one of only two vertebrates known to utilize a mixed

mating strategy, the mangrove rivulus fish (*Kryptolebias marmoratus*).

The mangrove rivulus (hereafter "rivulus"), is a small euryhaline killifish that inhabits

high elevation mangrove forests of Florida, the Bahamas, and Central America. Populations

consist primarily of self-fertilizing hermaphrodites with varying proportions of males (no

females). Rates of outcrossing and selfing also vary among populations and correlate with the

proportion of males in the population; outcrossing rates increase as the proportion of males

increase (Mackiewicz et al. 2006*b*; Tatarenkov et al. 2009). While self-fertilization happens

internally, males can only achieve reproductive success if/when hermaphrodites lay unfertilized

eggs, and there is no evidence that hermaphrodites outcross with each other (Furness et al. 2015).

Rivulus males result from sex change, after which selfing hermaphrodites that make the sexual

transition become obligate outcrossers. There also appears to be genetic variation for the

propensity to change sex (Gresham et al. 2020, Turner 2006). Hermaphrodites fertilize the vast

majority of their own eggs (> 94%, Harrington 1971, personal observations), giving males very

limited opportunities for reproductive success. Previously, it was reported that while individuals

that change sex to male are significantly more likely to survive environmental challenges than

those that remain hermaphrodite, they are much less likely to lay any eggs before sex change

occurs (only 2 out of 180, Gresham et al. 2020). This presents the problem of how individuals

that change sex might successfully pass their sex-changing alleles on to the next generation.

We hypothesized that the proportion of fertilized eggs (a measure of self-compatibility)

would change with age and/or the propensity of a given lineage to change sex. First, we

predicted that self-compatibility would increase with age. Cole and Noakes (1997) reported

histological evidence that ovarian tissue matures before spermatogenic tissue. We predicted that

there would be a period immediately after sexual maturity where oviposition of unfertilized eggs

would be more likely. Second, we predicted that overall fecundity and/or self-compatibility

would be greater in lineages that are more likely to change sex. This prediction is based on two

competing ideas about how sex-changing alleles are successfully maintained in rivulus

populations: i) selection should favor individuals in high sex changing lineages to be very fecund

and self-compatible prior to changing sex, so as to compensate for any lost reproductive success

after the transition, and/or ii) selection should favor individuals derived from lineages that

change sex frequently to be highly fecund and self-compatible so that the siblings of sex

changers recoup the lost reproductive success of their "brothers."

120   Methods

We designed and executed an experiment to measure the fecundity and self-compatibility

of 213 individual fish representing 100 genetically distinct lineages (determined via

microsatellite genotyping using 32 loci; (Mackiewicz et al. 2006*a*; Tatarenkov et al. 2010). We

calculated the propensity to change sex for each of the 100 lineages as the proportion of

individuals in each lineage that transition from hermaphrodite to male under common garden

conditions (Gresham et al. 2020), which ranged from 0 - 68.6%. All experimental fish were

raised from hatching in individual plastic containers (Rubbermaid® Take-a-long Deep Squares)

filled with approximately 200 mL 25‰ synthetic saltwater (prepared with Instant Ocean® sea

salt and aged tap water) until 30 days post hatching (dph), at which time water volume was

increased to 600 mL (25‰). At 60 dph, we suspended a large marble-sized ball of egg-laying

substrate (Poly-fil®) from one top corner of the container. Beginning at 67 dph, each container

was checked weekly for eggs in the substrate, around the top edge of the container, and on the

bottom of the container. Fish were checked weekly for eggs until they changed sex to male or

were 365-371 dph, ensuring that each remaining hermaphrodite was at least 365 dph and had

been checked weekly 43 times.

Eggs were collected into 59 mL plastic cups (Fabri-Kal® Greenware GXL250PC) filled

with approximately 6 mL 25‰ synthetic saltwater. Eggs were immediately counted and viewed

under a stereomicroscope. Eggs were photographed and scored as either fertilized, unfertilized,

or inviable. Fertilized and unfertilized eggs were identified by the presence or absence of a

perivitelline space, respectively. Inviable eggs were identified by their marked discoloration and

opaque appearance (Figure 1). Unfertilized and inviable eggs were removed from the cup and

discarded. Fertilized eggs were left in the cups, and water was replaced every 7 days. Cups were

checked every day for hatchlings. Hatchlings from the experimental animals were then scored as

alive or dead, with live hatchlings being euthanized in 4ºC water. Experimental fish were also

observed for signs of sex change each week when the tubs were checked for eggs, as described in

(Scarsella et al. 2018); orange freckles or orange skin is the most reliable external character of

sex change. Individuals that changed sex (35 of the 213) were separated from the others and the

date and age recorded. All experimental animals were isolated from the general colony for the

duration of the experiment, kept on a 12h light: 12 h dark photoperiod, and fed a 4 mL

suspension of brine shrimp (*Artemia*) nauplii (~2000 shrimp) six days per week. Room

temperature was also maintained at 26.45 ± 1.6 °C (mean ± SEM). The University of

Alabama Institutional Animal Care and Use Committee approved all procedures (IACUC

Protocol #18-10-1644).

Figure 1. Scoring of eggs. Eggs were scored as A) fertilized, B) unfertilized, or C) inviable. Fertilized eggs were identified by the presence of the perivitelline space (ps). Unfertilized eggs lack the perivitelline space. Inviable eggs were identified by their opaque or discolored appearance.

Fecundity and Self-compatibility Statistical Analysis

We examined the relationship between fecundity (number of eggs laid) and both the

propensity for a given lineage to change sex, and the individual's age. To determine whether

fecundity varied as a function of the lineages' propensities to change sex, we first ran a nominal

logistic model with 'laid eggs' as a categorical dependent variable (yes = 1, no = 0) and lineage

sex change propensity as the independent variable. We used a general linear model to examine

the relationship between total number of eggs laid (continuous, dependent) and lineage sex

change propensity (independent). Residuals of the total number of eggs were left skewed, so we

transformed the number of eggs with square root (x+1). To determine whether fecundity (total

number of eggs laid) varied as a function of age, we used a generalized regression model with a

zero-inflated Poisson distribution and parent ID as a random effect to account for the fact that

each parent was checked for eggs multiple times. An initial plot of the raw data suggested a

possible non-linear relationship between age and fecundity so, we constructed linear, quadratic

and cubic models and compared model fits using Akaike's Information Criterion (AIC). The

linear, quadratic and cubic age terms were scaled by mean and standard deviation (i.e., z-score)

prior to running the models. The cubic model fit best (Table 3.1), so we report the results of this

model.

We modeled self-compatibility in two ways: by the proportion of eggs that were fertilized

and by hatch success of fertilized eggs. To determine whether the proportion of fertilized eggs

varied as a function of the lineages' propensity to change sex, we ran a general linear model. To

determine whether the proportion of fertilized eggs varied as a function of the parents' age when

the eggs were collected, we ran a general linear model with parent ID as a random effect. To

determine whether the proportion of fertilized eggs varied as a function of the total number of

eggs laid by each parent, we ran a general linear model. We calculated hatch success as the

proportion of fertilized eggs that resulted in a live hatchling. To determine whether hatch success

varied as a function of the lineages' propensity to change sex, we ran a general linear model.

Models were run in JMP Pro version 15.0.0 (JMP®, Version 15 Pro 2019) and in RStudio using

the glmmTMB package (RStudioTeam 2016; Brooks et al. 2017; R Core Team 2018). Data files

will be deposited on GitHub upon publication acceptance.

The experiment included 213 individual fish from 100 different lineages. Thirty-five of

those fish changed sex to male (16.4%) before the experiment ended at 365 - 371 dph. Of those

35, three individuals laid eggs first (9% of males, 1% of the total cohort). We collected 14,310

eggs, had 10,267 hatchlings (71.7% of eggs hatched), and 10,059 live hatchlings (98% of

hatchlings were alive). The three individuals that changed sex to male laid a total of 69 eggs by

197 dph (the oldest age at which a fish was identified as male), which equated to, on average, 23

$\pm$ 13 (SEM) eggs per individual among the three, but 1.97 $\pm$ 1.4 eggs per capita for all

individuals that changed sex to male. The remaining 178 hermaphrodites laid a total of 7,341

eggs before they collectively aged to 197 dph (41.2 $\pm$ 2.2 eggs per individual). We started

checking for eggs from each fish at 67 dph and collected a single egg from one individual on the

first egg check; the egg was fertilized. Overall, the proportion of eggs that were fertilized was

93.4%. The proportion of eggs that were unfertilized was 2.5%, and the proportion identified as

"inviable" was 4.1%.

Our results indicate that fecundity initially increases with age until about 175 dph, then

decreases until it plateaus at about 350 dph (Table 1 Figure 2A). The proportion of fertilized

eggs also increased significantly with age, but not the total fecundity of the parent (Table 1,

Figure 2B). Because we did not record the hatch success of each individual collection (i.e.,

clutch) of eggs, we were not able to analyze whether hatch success changed with age. The

lineages' propensity to change sex was significantly related to fecundity and self-compatibility.

When individuals that changed sex to male were included in the model, the odds of laying any

eggs decreased significantly as the propensity to change sex increased (Table 2). However, this

relationship became non-significant when individuals that changed sex to male were excluded

(Table 2). Conversely, the number of eggs laid decreased significantly as the propensity to

change sex increased, whether or not males were included in the model (Table 2 Figure 3A).

Both the proportion of fertilized eggs and the hatch success of those eggs also decreased

significantly with increasing propensity to change sex (Table 2, Figure 3B, C).

Table 1. Summary of fecundity and self-compatibility models with age of parent fish as the dependent variable. AIC = Akaike's Information Criterion; ZI = zero inflation; F = F-ratio; P = P-value; Z = Z-value; df = degrees of freedom.

Fecundity						
Models	AICc	ΔAICc	ZI parameter estimate (p-value)	Estimate	Z	P
# eggs ~_age + age^2 + age^3	33140.0	0.0	-0.98 (<0.0001)	age: 3.06 age^2: -5.76 age^3: 2.52	10.73 9.85 7.94	<0.0001 <0.0001 <0.0001
# eggs ~_age + age^2	33198.5	58.5	-0.92 (<0.0001)	age: 0.85 age^2: -1.14	13.01 -17.67	<0.0001 <0.0001
# eggs ~_age	33527.0	387.0	-0.83 (<0.0001)	age: -0.29	-29.58	<0.0001
Proportion of fertilized eggs						
			Estimate	F	df	P
Parent Age			0.000069	4.08	1	0.04
Number of Eggs Laid			0.000093	0.63	1	0.43

Table 2. Summary of fecundity, self-compatibility, and hatch success models with lineage sex change propensity as the dependent variable. Fecundity was square root (x+1) transformed to ensure normality. The models for proportion of fertilized eggs and hatch success already excluded the males that did not lay any eggs, thus separate models were not needed. Significant effects are shown in bold. F = F-ratio, P = P-value, df = degrees of freedom.

Probability of Laying Eggs	Odds Ratio	χ2	df	P
Lineage sex change propensity, **including** males	0.003	23.85	1	**<0.0001**
Lineage sex change propensity, **excluding** males	0.006	2.62	1	0.11
Fecundity (# of eggs laid)	Estimate	F	df	P
Lineage sex change propensity, **including** males	-4.17	36.29	1	**<0.0001**
Lineage sex change propensity, **excluding** males	-4.25	9.09	1	**0.003**
Proportion of fertilized eggs	Estimate	F	df	P
Lineage sex change propensity, **including** males	-0.10	4.86	1	**0.029**
Hatch Success	Estimate	F	df	P
Lineage sex change propensity, **including** males	-0.36	33.85	1	**<0.0001**

Figure 2. Age effects on the number of eggs collected and the proportion of those eggs that were fertilized. A) Age affected fecundity cubically; the number of eggs oviposited increased initially, then decreased between 170 – 180 dph, then plateaued around 350 dph. B) The proportion of eggs that were fertilized increased significantly with fish age. The additional straight line at the top of the graph is a string of points representing all of the egg collections where 100% of the eggs were fertilized.

Figure 3. Lineage sex change propensity effects on A) fecundity, B) the proportion of eggs that were fertilized, and C) hatch success of fertilized eggs. Lineage sex change propensity is measured as the proportion of each lineage that change sex in isolation under common garden conditions in our colony (lineage % male).

204   Discussion

Mixed mating strategies offer the fitness rewards of both outcrossing with conspecifics

and fertilizing one's own eggs. The cost-to-benefit ratio of each strategy is expected to vary

across space and time as the external and internal environments also change (Maynard Smith

1971; Lively and Lloyd 1990; Lehtonen et al. 2012; Layman et al. 2017; Lynch et al. 2018).

There is also variation among individuals in characters that affect mating strategy: fecundity and

self-compatibility (Stephenson et al. 2000; Travers et al. 2004; Busch 2005). Variation in self-

compatibility, combined with external factors such as pollinator and/or mate availability or

parasite pressure, has the potential to drive evolutionary changes in mating strategy. For instance,

if self-compatible individuals have greater reproductive fitness (produce more offspring and/or

offspring have greater survival rates) and survive just as well as self-incompatible individuals,

then the population may evolve to complete or near complete selfing. Comparatively, if there are

plenty of mates/pollinators and/or inbreeding depression that increases the benefits to cost ratio

of outcrossing, populations will evolve to complete or near complete outcrossing. We tested, in

the only self-fertilizing hermaphroditic vertebrate, the hypothesis that self-compatibility and

fecundity would vary with age and as a function of the lineages' propensities to change sex.

The most important finding was that both fecundity and self-compatibility decreased

as lineages became more likely to change sex, which is the exact opposite of what we predicted.

We predicted that fecundity and/or self-compatibility would increase with the lineages'

propensity to change sex for one of two reasons: either individuals that eventually change sex

would produce more eggs with greater self-compatibility, or the remaining hermaphrodites from

high sex changing lineages would produce at least as many eggs with equal self-compatibility as

those from lineages less likely to change sex. Given that neither of these predictions were

supported, it remains unclear how sex change from a self-fertilizing hermaphrodite to an obligate

outcrossing male is maintained in rivulus populations. Theory suggests that males incur a cost

because, when compared to self-fertilizing hermaphrodites, they cannot produce offspring and

also reduce the parent's genetic contribution to offspring by one half (Maynard Smith 1971,

1978). Males must overcome this cost by producing many extremely fit offspring.

Prior experiments have provided evidence of outbreeding depression on fecundity

(Gresham et al. 2020) in several ecologically relevant conditions, increasing the cost of males

and outcrossing. However, Ellison et al. (2011) provide evidence of potential inbreeding

depression under parasitic stress, therefore decreasing the cost of males. Male size also increases

with heterozygosity, possibly allowing more heterozygous males to secure more resources and

have greater access to unfertilized eggs (Molloy et al. 2011). The results from this experiment

suggest that the cost of being male is highly variable among lineages. Only three individuals laid

eggs before changing sex to male, indicating that they were not as fecund as self-fertilizing

hermaphrodites. The odds of laying any eggs decreased as sex change propensity increased when

all individuals were included in the model. However, when we excluded individuals that changed

sex, there was not an effect of lineage sex change propensity, demonstrating that the

hermaphrodites from high sex changing lineages are just as likely to lay *an* egg as those from

low sex changing lineages. In contrast, hermaphrodites from high sex changing lineages laid

significantly fewer eggs, even if the males were excluded from the model. Self-compatibility and

hatch success also decreased as propensity to change sex increased. These results indicate that

individuals from high sex changing lineages that remain hermaphrodite are not very fecund or

self-compatible. This exemplifies that the cost of changing sex to male might vary among

genetically distinct lineages. In particular, individuals from low sex changing, highly fecund and

self-compatible lineages are likely to experience the highest cost of sex change.

We found that fecundity initially increases with age, then decreases, and finally plateaus

as individuals age from about 60 dph to about 370 dph. The only other report of potential age-

related changes in fecundity in rivulus comes from Harrington (1971). From 37 fish, he reported

significant seasonality in "ovarian function" (i.e., fecundity) over three years when exposed to

natural day light cycles, but not a significant change in fecundity with age. The seasonal decline

reported by Harrington (1971), which occurred from September through December, aligns with

the end of our experiment. The fish in our experiment hatched and then were removed from the

experiment over a four-month period between 30 August 2018 and 22 December 2019. It is

possible that the association between age and fecundity that we found, particularly the shallow

decline in egg production from 180-300 dph represents the same seasonal decline described by

Harrington (1971). He allowed his fish to experience natural daylight cycles. Our fish were kept

on a constant photoperiod and a relatively constant temperature, and so should have been

removed from any external signals of season. To truly understand how fecundity is affected by

age and seasonality, we need additional years of records from multiple lineages.

Overall, 93.4% of the eggs we collected were fertilized. This is very close to the

proportion of eggs fertilized reported by Harrington (1971). From six fish of one lineage and

1,056 eggs, he identified 94.3% as fertilized. We recorded 2.5% unfertilized and 4.1% inviable,

compared to Harrington (1971), who reported 4% unfertilized and 1.7% inviable. It is likely that

most of the inviable eggs we recorded were unfertilized when oviposited, but then became

inviable and discolored by the time we collected them. Harrington (1971) collected eggs every

day and was more likely to record unfertilized eggs before they became inviable. Our first

fertilized egg was collected at 67 dph, indicating the egg was oviposited sometime between 60

dph and 67 dph. This is the earliest reported oviposition in rivulus.

Regardless of the cost of changing sex to male, there is still the issue of passing sex

changing alleles to the next generation. How do lineages that are less fecund, less self-

compatible, and that frequently change sex to male not disappear from populations? It is possible

that many of these lineages do go extinct but, some Belizean populations maintain relatively high

male proportions (> 20%) and non-trivial rates of outcrossing (Davis 1990; Turner et al. 1992;

Tatarenkov et al. 2009, 2015), which suggests that males and outcrossing are under positive

selection in some places. Males must be able to successfully find unfertilized eggs, but we are

not sure how that happens. There have been multiple attempts to mate males and hermaphrodites

in the lab. Mackiewicz et al. (2006*a*) paired multiple nearly-senescent hermaphrodites and males

in individual tubs and confirmed two out of 32 offspring were outcrossed progeny. Nakamura et

al. (2008) surgically removed eggs from two hermaphrodites for *in vitro* outcrossing, but only

one out of thirteen hatchlings were outcrossed instead of self-fertilized. Our laboratory has tried

pairing hermaphrodites and males in different ratios with no outcrossing success. In behavior

trials, reported interactions between pairs of hermaphrodites and males have ranged from no

courtship behaviors (Martin 2007) to significantly more courtship behaviors than same sex pairs

(Luke and Bechler 2010) or courtship behaviors by the males with reciprocal attacks by the

hermaphrodites (Turner et al. 1992). It is clear that we have not yet reproducibly identified

conditions that favor outcrossing.

There often are opposing selection pressures for and against biparental sex (outcrossing).

Mixed mating individuals can benefit from outcrossing when the costs of biparental sex are low,

but self when the costs of biparental sex are high or when mates and/or pollinators are sparse.

The relative benefit of selfing or outcrossing among individuals is also dependent on the

individual level of self-compatibility, which can have a genetic basis and/or respond to the

external environment in a plastic or flexible manner. Individuals and lineages that are not very

self-compatible cannot exploit the benefits of selfing as much as individuals that are more self-

compatible, even when the costs of outcrossing are high. In the mangrove rivulus fish, we have

found significant variation in self-compatibility that correlates with age and with the lineages'

propensities to change sex. Individuals from lineages with a higher sex change propensity laid

significantly fewer eggs and a significantly lower proportion of the eggs were self-fertilized (i.e.,

they were less self-compatible). Also, the proportion of fertilized eggs increased with age.

In wild populations, interactions between individuals' level of self-compatibility and

external conditions have the potential to drive evolutionary changes. The internal and external

environments can alter the age structure of populations and/or the economics of outcrossing and

selfing. In populations that have few or no lineages that frequently change sex (such as those

recently founded by highly self-compatible individuals), lineages with higher propensities to

change sex would likely struggle to become established, if offspring of selfed and outcrossed

matings are equally fit. In these populations, individuals that change sex frequently will find an

environment with very few unfertilized eggs, and very few opportunities to pass their sex

changing alleles on to the next generation. However, once high sex change propensity lineages

become established, possibly due to additional migration or outcrossed progeny having a fitness

advantage over selfed progeny, these lineages can maintain sex change and males. Indeed,

among rivulus populations across the geographic range we find a significant increase in the rate

of outcrossing that correlates with an increase in the proportion of sex changing lineages

(Mackiewicz et al. 2006*b*; Tatarenkov et al. 2015). We also find variation among lineages within

in a population in the propensity to change sex, but the magnitude of variation is population

specific (Gresham et al. 2020).

Males, and lineages that frequently produce males, may also have better success in

populations with many young hermaphrodites. While we did not find evidence that younger

hermaphrodites reproduce as "pure females" and only lay unfertilized eggs, younger

hermaphrodites did lay more unfertilized eggs. If the fitness of progeny derived from selfing and

outcrossing is relatively equal, population age structure can act as a selection pressure for or

against lineages that frequently change sex. For instance, populations that lose a greater

proportion of juveniles and young adults to predation will have proportionally fewer unfertilized

eggs available for males than those populations in which younger individuals thrive; selecting

against high sex changing lineages. These results open up fascinating opportunities for continued

laboratory studies on lineage variation in traits such as size, fecundity, and sex change across

different environments, and continued field studies to track demographic and population genetic

characteristics such as selfing and outcrossing rates, the proportion of males, and population size

and density.

References

335   Allard, R. W., G. R. Babbel, M. T. Clegg, and a L. Kahler. 1972. Evidence for coadaptation in
336   *Avena barbata*. Proceedings of the National Academy of Sciences of the United States of
337   America 69:3043–8.
338

339   Baker, H. 1955. Self-compatibility and establishment after "long-distance" dispersal. Evolution
340   9:347–349.
341

342   Bell, G. 1988. Recombination and the immortality of the germ line. Journal of Evolutionary
343   Biology 1:67–82.
344

345   Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.
346   Skaug, et al. 2017. glmmTMB balances speed and flexibility among packages for zero-
347   inflated generalized linear mixed modeling. The R Journal 9:378–400.
348

349   Busch, J. W. 2005. The evolution of self-compatibility in geographically peripheral populations
350   of *Leavenworthia alabamica* (Brassicaceae). American Journal of Botany 92:1503–1512.
351

352   Clutton-Brock, T. H., and G. A. Parker. 1995. Sexual coercion in animal societies. Animal
353   Behaviour 49:1345–1365.
354

355   Cole, K. S., and D. L. G. Noakes. 1997. Gonadal development and sexual allocation in mangrove
356   killifish, *Rivulus marmoratus* (Pisces: Atherinomorpha). Copeia 1997:596–600.
357

358   Darwin, C. 1876. The effects of cross and self fertilisation in the vegetable kingdom. D.
359   Appleton.
360

361   Davis, W. P. 1990. Field observations of the ecology and habits of mangrove rivulus.
362   Ichthyological Exploration of Freshwaters 1:123–134.
363

364   Dolgin, E. S., B. Charlesworth, S. E. Baird, and A. D. Cutter. 2007. Inbreeding and outbreeding
365   depression in *Caenorhabditis* nematodes. Evolution 61:1339–1352.
366

367   Ellison, A., J. Cable, and S. Consuegra. 2011. Best of both worlds? association between
368   outcrossing and parasite loads in a selfing fish. Evolution 65:3021–3026.
369

370   Escobar, J. S., J. R. Auld, A. C. Correa, J. M. Alonso, Y. K. Bony, M. A. Coutellec, J. M. Koene,
371   et al. 2011. Patterns of mating-system evolution in hermaphroditic animals: Correlations
372   among selfing rate, inbreeding depression, and the timing of reproduction. Evolution
373   65:1233–1253.
374

375   Fowler, K., and L. Partridge. 1989. A cost of mating in female fruitflies. Nature 338:760–761.
376

377   Furness, A. I., A. Tatarenkov, and J. C. Avise. 2015. A genetic test for whether pairs of
378   hermaphrodites can cross-fertilize in a selfing killifish. Journal of Heredity 106:749–752.

379 Gresham, J. D., K. M. Marson, A. Tatarenkov, and R. L. Earley. 2020. Sex change as a survival
380   strategy. Evolutionary Ecology 34:27–40.
381
382 Harrington Jr, R. W. 1971. How ecological and genetic factors interact to determine when self-
383   fertlizing hermaphrodites of *Rivulus marmoratus* change into functional secondary males,
384   with a reappraisal of the modes of intersexuality among fishes. Copeia 1971:389–432.
385
386 Hartfield, M., and P. D. Keightley. 2012. Current hypotheses for the evolution of sex and
387   recombination. Integrative Zoology 7:192–209.
388
389 JMP®, Version 15 Pro. 2019. SAS Institute Inc., Cary, NC.
390
391 Layman, N. C., M. T. R. Fernando, C. R. Herlihy, and J. W. Busch. 2017. Costs of selfing
392   prevent the spread of a self-compatibility mutation that causes reproductive assurance.
393   Evolution 71:884–897.
394
395 Lehtonen, J., M. D. Jennions, and H. Kokko. 2012. The many costs of sex. Trends in Ecology &
396   Evolution 27:172–178.
397
398 Lively, C. M. 2010. A review of Red Queen models for the persistence of obligate sexual
399   reproduction. Journal of Heredity 101:S13–S20.
400
401 Lively, C. M., and D. G. Lloyd. 1990. The cost of biparental sex under individual selection. The
402   American Naturalist 135:489–500.
403
404 Luke, K. N., and D. L. Bechler. 2010. The role of dyadic interactions in the mixed-mating
405   strategies of the mangrove rivulus *Kryptolebias marmoratus*. Current Zoology 56:6–17.
406
407 Lynch, Z. R., M. J. Penley, and L. T. Morran. 2018. Turnover in local parasite populations
408   temporarily favors host outcrossing over self-fertilization during experimental evolution.
409   Ecology and Evolution.
410
411 Mackiewicz, M., A. Tatarenkov, A. Perry, J. R. Martin, J. F. Elder, D. L. Bechler, and J. C.
412   Avise. 2006*a*. Microsatellite documentation of male-mediated outcrossing between
413   inbred laboratory strains of the self-fertilizing mangrove killifish (*Kryptolebias
414   marmoratus*). The Journal of Heredity 97:508–13.
415
416 Mackiewicz, M., A. Tatarenkov, D. S. Taylor, B. J. Turner, and J. C. Avise. 2006*b*. Extensive
417   outcrossing and androdioecy in a vertebrate species that otherwise reproduces as a self-
418   fertilizing hermaphrodite. Proceedings of the National Academy of Sciences of the
419   United States of America 103:9924–8.
420
421 Martin, S. B. 2007. Association behaviour of the self-fertilizing *Kryptolebias marmoratus*
422   (Poey): The influence of microhabitat use on the potential for a complex mating system.
423   Journal of Fish Biology 71:1383–1392.
424

425   Maynard Smith, J. 1971. What use is sex? Journal of Theoretical Biology 30:319–335.
426
427   Maynard Smith, J. 1978. The evolution of sex. Cambridge Univ Press, Cambridge.
428
429   Molloy, P. P., E. A. Nyboer, and I. M. Côté. 2011. Male-Male Competition in a Mixed-Mating
430   Fish. Ethology 117:586–596.
431
432   Morran, L. T., O. G. Schmidt, I. a Gelarden, R. C. Parrish, and C. M. Lively. 2011. Running with
433   the Red Queen: host-parasite coevolution selects for biparental sex. Science (New York,
434   N.Y.) 333:216–8.
435
436   Muller, H. J. 1932. Some genetic aspects of sex. American Society of Naturalists 703:118–138.
437
438   Nakamura, Y., K. Suga, Y. Sakakura, T. Sakamoto, and A. Hagiwara. 2008. Genetic and growth
439   differences in the outcrossings between two clonal strains of the self-fertilizing mangrove
440   killifish. Canadian Journal of Zoology 86:976–982.
441
442   Pannell, J. R., J. R. Auld, Y. Brandvain, M. Burd, J. W. Busch, P. Cheptou, J. K. Conner, et al.
443   2015. The scope of Baker's law. New Phytologist 208:656–667.
444
445   R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for
446   Statistical Computing, Vienna, Austria.
447
448   RStudioTeam. 2016. RStudio: Integrated Development for R. RStudio, Inc., Boston.
449
450   Scarsella, G. E., J. D. Gresham, and R. L. Earley. 2018. Relationships between external sexually
451   dimorphic characteristics and internal gonadal morphology in a sex-changing fish.
452   Journal of Zoology 305:133–140.
453
454   Stebbins, G. L. 1957. Self fertilization and population variability in the higher plants. The
455   American Naturalist 91:337.
456
457   Stephenson, A. G., S. V Good, and D. W. Vogler. 2000. Interrelationships among inbreeding de-
458   pression, plasticity in the self-incompatibility system, and the breeding system of
459   *Campanula rapunculoides* L. (Campanulaceae). Annals of Botany 85:211–219.
460
461   Tatarenkov, A., R. L. Earley, B. M. Perlman, D. S. Taylor, B. J. Turner, and J. C. Avise. 2015.
462   Genetic subdivision and variation in selfing rates among Central American populations of
463   the mangrove rivulus, *Kryptolebias marmoratus*. Journal of Heredity 106:276–284.
464
465   Tatarenkov, A., S. M. Q. Lima, D. S. Taylor, and J. C. Avise. 2009. Long-term retention of self-
466   fertilization in a fish clade. Proceedings of the National Academy of Sciences of the
467   United States of America 106:14456–14459.
468
469
470

471   Tatarenkov, A., B. C. Ring, J. F. Elder, D. L. Bechler, and J. C. Avise. 2010. Genetic
472   composition of laboratory stocks of the self-fertilizing fish *Kryptolebias marmoratus*: a
473   valuable resource for experimental research. PloS one 5:e12863.
474

475   Travers, S. E., J. Mena-Ali, and A. G. Stephenson. 2004. Plasticity in the self-incompatibility
476   system of *Solanum carolinense*. Plant Species Biology 19:127–135.
477

478   Tsitrone, A., P. Jarne, and P. David. 2003. Delayed Selfing and Resource Reallocations in
479   Relation to Mate Availability in the Freshwater Snail *Physa acuta*. American Naturalist
480   162:474–488.
481

482   Turner, B. J., W. P. Davis, and D. S. Taylor. 1992. Abundant males in populations of a selfing
483   hermaphrodite fish, *Rivulus marmoratus*, from some Belize cays. Journal of Fish Biology
484   40:307–310.
485

486   Winn, A. A., E. Elle, S. Kalisz, P. Cheptou, C. G. Eckert, C. Goodwillie, M. O. Johnston, et al.
487   2011. Analysis of inbreeding depression in mixed-mating plants provides evidence for
488   selective interference and stable mixed mating. Evolution 65:3339–3359.
489

490   Wright, S. I., S. Kalisz, and T. Slotte. 2013. Evolutionary consequences of self-fertilization in
491   plants. Proceedings. Biological sciences / The Royal Society 280:20130133.
492