Surfactin stimulated by pectin molecular patterns and root exudates acts as a key driver of *Bacillus*-plant mutualistic interaction

Grégory Hoff[1#], Anthony Arguelles-Arias[1], Farah Boubsi[1], Jelena Prsic[1], Thibault Meyer[1*], Heba M. M. Ibrahim[2], Sebastien Steels[1], Patricio Luzuriaga[1], Aurélien Legras[1], Laurent Franzil[1], Michelle Lequart[3], Catherine Rayon[3], Victoria Osorio[4], Edwin de Pauw[4], Yannick Lara[5], Estelle Deboever[6], Barbara de Coninck[2], Philippe Jacques[1], Magali Deleu[6], Emmanuel Petit[3], Olivier Van Wuytswinkel[3], Marc Ongena[1]

[1] Microbial Processes and Interactions, TERRA Teaching and Research Center, BioEcoAgro, Joint Research Unit/UMR transfrontalière 1158, University of Liège - Gembloux Agro-Bio Tech, Gembloux, Belgium

[2] Division of Plant Biotechnics, Department of Biosystems, Faculty of Bioscience Engineering, KU Leuven, Leuven, Belgium

[3] Unité Biologie des Plantes et Innovation, BioEcoAgro, Joint Research Unit/UMR transfrontalière 1158, Université de Picardie Jules Verne, UFR des Sciences, Amiens, France

[4] Mass Spectrometry Laboratory, MolSys Research Unit, Department of Chemistry, University of Liège, Liège, Belgium

[5] Astrobiology, UR-ASTROBIOLOGY, Geology Department, University of Liège, Liège, Belgium

[6] Molecular Biophysics at Interfaces Laboratory, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium

Corresponding authors: Grégory Hoff g.hoff@uu.nl
                           Marc Ongena marc.ongena@uliege.be

Current address: Ecology and Biodiversity, Department of Biology, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands

* Current address: UMR Ecologie Microbienne, F-69622, University of Lyon, Université Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup, Villeurbanne, France

Abstract

Bacillus velezensis is considered as model species belonging to the so-called *B. subtilis*
complex that typically evolved to dwell in the soil rhizosphere niche and establish intimate
association with plant roots. This bacterium provides protection to its natural host against
diseases and represents one of the most promising biocontrol agents. However, the
molecular basis of the cross-talk that this bacterium establishes with its natural host has
been poorly investigated. We show here that these plant-associated bacteria have evolved
some polymer-sensing system to perceive their host and that in response, they increase the
production of the surfactin-type lipopeptide. Furthermore, we demonstrate that surfactin
synthesis is favoured upon growth on root exudates and that this lipopeptide is a key
component used by the bacterium to optimize biofilm formation, motility and early root
colonization. In this specific nutritional context, the bacterium also modulates qualitatively
the pattern of surfactin homologues co-produced *in planta* and mainly forms variants that
are the most active at triggering plant immunity. Surfactin represents a shared good as it
reinforces the defensive capacity of the host.

Introduction

Soil is among the richest ecosystems in terms of microbial diversity, but only a subset of these microbes has evolved to efficiently establish in the competitive and nutrient-enriched rhizosphere layer surrounding plant roots [1]. The rhizosphere includes plant beneficial bacteria dwelling on the rhizoplane as multicellular biofilm communities, feeding on exuded carbohydrates [2,3], and, in turn, contributing to host fitness via growth stimulation and protection against phytopathogens [4,5]. This biocontrol activity is mediated via competition for nutrients and space, direct growth inhibition of the pathogenic (micro)organisms and more indirectly, by stimulating the host defensive capacity in an immunization-like process which leads to induced systemic resistance (ISR, [6,7]). This ISR mechanism results in enhanced defense lines and reduced disease symptoms upon perception of plant beneficial microbes [6,8].

From an ecological viewpoint, rhizosphere establishment and persistence of these beneficial bacteria rely on various traits but efficient root colonization and high competitiveness toward the surrounding microbiological network are pivotal. It is hypothesized that the potential to produce a wide range of chemically diverse and bioactive secondary metabolites (BSMs) acting as signals and/or antimicrobials is a common key feature of these beneficial bacteria [5,9,10]. Members of the *Bacillus velezensis* species are considered as archetypes of plant-associated beneficial bacilli and are among the most prolific BSMs producers with more than 12% of their genome devoted to the synthesis of compounds contributing to both ecological competence and biocontrol activity [11–15]. Among their BSM arsenal, the cyclic lipopeptide surfactin, is synthesized non-ribosomally by a multi-modular mega-enzyme machinery (encoded by the *srfA* operon) and is formed as a mix of naturally co-produced homologues varying in the length of the fatty acid chain. This multifunctional compound is of particular interest because it retains important roles in key developmental processes such as bacterial motility, biofilm formation and root colonization [16–18], but also because it represents the best described *Bacillus* triggers for plant immunity [6,8]. The potential of surfactin to stimulate ISR has been demonstrated on various plants including Solanaceae like tobacco and tomato on which it acts as main if not sole elicitor formed by *B. subtilis* and *B. velezensis* species [10,19]. In support to its key role in interaction with the host plant, we also previously reported that surfactin is promptly formed in the

course of early colonization and that its production is stimulated upon sensing root tissues

[20].

However, in contrast to the well-studied interactions between plants and microbial

pathogens or nitrogen-fixing bacteria [21], relatively little is known on the molecular basis of

cooperative interactions between plants and beneficial bacteria such as *B. velezensis*

[11,20,22]. More specifically, how and to what extent the expression of key bacterial BSMs may

be modulated by plant factors is poorly understood. A better knowledge is not only critical

for providing new insights in rhizosphere chemical ecology but also for optimizing the use of

these species as biocontrol agents, which still suffer from insufficient efficacy in practice [23].

Here, we investigated the molecular interaction driving the early steps of partnership

establishment between plant roots and *B. velezensis*. We show that cell wall pectin acts in

synergy with soluble root exudates as plant host cues perceived by *B. velezensis*. In

response, the bacterium stimulates the production of specific surfactin variants as key

components of its secretome to further improve the fitness of both partners *i.e.* early root

colonization and thus rhizosphere competence of the bacterium and priming of immunity in

the host plant.

Results

Pectin fragments of high polymerization degree act as host cues triggering surfactin production

We previously described that early production of surfactin (as a mix of naturally co-produced homologues varying in the length of the fatty acid chain, Fig. 1) is stimulated in contact with root tissues and several plant cell wall-associated polymers (PCWP) [20]. In this work, we further investigated this phenomenon focusing on the impact of pectin, as it represents complex sugar polymers typically found in the plant primary cell wall and particularly abundant in the middle lamella layer [24]. We first tested the effect of crude pectin extracted from tobacco root PCWP (referred as cPec, Supplementary Fig. 1 for composition and related structure). An 8-fold increase of surfactin production was detected at the early exponential growth phase (OD_{600}=0.2-0.25) in *B. velezensis* GA1 liquid cultures supplemented with cPec compared to an un-supplemented culture (Fig. 1ab). Surfactin production was also 10 times enhanced upon addition at the same concentration of pure commercially available homogalacturonan (HG) with high degree of polymerization (DP) (Supplementary Fig. 2) but low level of methyl-esterification (HGLM) according to the manufacturer (Fig. 1b). HG was tested as the most abundant pectic polysaccharide constituent, which represents 65% of crude primary cell wall pectin [24]. Production of this lipopeptide was also enhanced to a similar level upon addition of highly methylated HG (HGHM), showing that the degree of methyl-esterification of the polymer is not a major trait influencing perception by the bacterium (Supplementary Fig. 3). Altogether, this supports a key role of the pectin backbone as plant molecular pattern that is sensed by the bacterium to stimulate surfactin synthesis.

Interestingly, by screening the CAZy database [25] for genes encoding carbohydrate-active enzymes potentially involved in PCWP degradation by *B. velezensis*, two putative pectate/pectin lyases encoding genes were detected. These two genes, referred as *pelA* and *pelB* (accessions *GL331_08735* and *GL331_04125 in B. velezensis* GA1, respectively), are highly conserved among all sequenced *Bacillus* genomes that belong to the "Operational Group *B. amyloliquefaciens*" (Supplementary table 1) [26]. *pelA* and *pelB* are readily expressed in GA1 *in vitro* and the corresponding enzymes efficiently convert HG into

unsaturated oligogalacturonides with consistent activity occurring at the beginning of
stationary phase (Supplementary Fig. 4ab). However, the bacterial perception of oligomers
with lower polymerization degree compared to HG is not obvious since oligogalacturonides
(OG) did not stimulate surfactin biosynthesis (Fig. 1b, Supplementary Fig. 5 for OG
characterization). Supplementation with galacturonic acid (GA) led to a reduction of
surfactin production at mid exponential phase (OD_{600}=0.35, Supplementary Fig. 6 for growth
curves, Fig. 1c). Surfactin production is thus specifically boosted upon sensing long degree of
polymerization (DP) polymers, but is somehow inhibited in presence of GA constituting the
pectin backbone. Such HG-driven surfactin stimulation also occurs in other *B. velezensis*
isolates tested (FZB42, QST713 and S499) and to a lower extent *B. pumilus* QST 2808. It does
not occur in the non-rhizosphere dwelling isolates *B. amyloliquefaciens* DSM7 or *B. subtilis*
ATCC 21332 (Fig. 1c) suggesting that this trait may be specific to bacilli with a plant-
associated lifestyle.

133 The root nutritional context favors early surfactin production

Bacillus velezensis quickly colonize tomato plantlets in a gnotobiotic system and
forms visible biofilm-like structures covering the main root and embedding lateral roots after
24-48h post inoculation (Fig 2a). This is correlated with consistent *srfAA* gene expression and
surfactin production rate in the cell population at these early times but it was maintained
albeit to a lower level, over the investigated timeframe of seven days (Fig. 2ab). Since
surfactin enhancement linked to the perception of the pectin backbone is only transient (Fig
1b), we hypothesized that root exudates, constantly secreted by the plant, may also
positively impact the synthesis of the lipopeptide. Surfactin production rate was thus
compared upon growth in a classical laboratory medium (LB) and in a root exudate-
mimicking medium (REM) reflecting the content of carbohydrates typically released by
tomato or tobacco roots [27]. It revealed an earlier and higher production by cells growing in
REM (Fig 2c). Surfactin production in REM medium is initiated earlier and is more efficient in
B. velezensis compared to other closely related but non plant-associated species such as *B.*
amyloliquefaciens or *B. subtilis* (Fig. 2d).

Addition of HG in REM medium compared to LB revealed a cumulative effect of this
PCWP and root exudates on surfactin production (Fig 3a). This could be of clear ecological

benefit for the bacterium since surfactin is known to favor motility of multicellular communities and biofilm formation [16,28,29]. However, a recent study questioned the real role of surfactin in these key functions, since it production appears as non-essential for pellicle biofilm formation in *B. subtilis* NCIB 3610, suggesting a strain dependant role [30]. We previously reported that motility and biofilm formation are boosted upon growth on root exudates [27]. Here we show that HG supplementation also favors *B. velezensis* GA1 spreading on low-agar medium (Fig 3b) and early biofilm formation based on pellicle development at the air-liquid interface [31] (Fig 3c). The role of surfactin in swarming, pellicle formation and early root colonization was further confirmed for *B. velezensis* GA1. Indeed, swarming motility on low agar plates was almost reduced to zero in a surfactin deficient mutant, and the same mutant was more than 3 times less efficient to produce pellicles at the air liquid interface and to promptly colonize tomato roots after 1 day post inoculation when compared to the WT (Fig 3def). Collectively, these data allow correlating the positive impact of PCWP on bacterial motility, biofilm formation and early root colonization through an anticipated surfactin production in *B. velezensis*.

Surfactin induction by PCWP is not linked to major transcriptional changes.

Both HG and root exudates stimulate surfactin production in GA1. However, while no activation of the *srfA* biosynthetic gene cluster was observed upon HG addition (Supplementary Fig. 7), an early and high surfactin gene expression was measured in *srfAp::gfp* cells growing in REM compared to LB medium (Fig. 4a). To unravel transcriptome wide changes in GA1 associated with the perception of HG, RNA-sequencing was performed on cells grown in REM with our without addition of HG and collected at various time points (lag, early exponential and a mid-exponential phases). The data confirmed that HG perception is not linked to an increased expression of the *srfA* operon but also revealed a quite limited and transient transcriptional reprogramming with only 58 genes differentially expressed over this timeframe (Supplementary table 2). Remarkably, more than 30% of these genes are involved in stress response or cell wall modifications and are down regulated in the presence of HG (Fig. 4b). We thus hypothesize that a long-term co-evolution process may have facilitated *Bacillus* establishment on the roots by the inhibition of a costly stress response after perception of HG. Addition of HG also leads to a 4.2-fold reduced expression of *flgM* encoding an inhibitor of SigD, the σ factor involved in the activation of

motility related genes [32]. This may contribute to enhanced spreading of multicellular
communities in addition to the positive effect of surfactin mentioned above.

Root exudates drive the bacterium to form surfactin homologues with long fatty acid chain
(LFAC) and variants enriched in valine

The NRPS machinery works as an assembly line in which each module is responsible
for recruiting and binding a specific amino acid to the nascent peptide after a first lipo-
initiation step for binding the fatty acid (FA) taken up from the cellular pool (Supplementary
Fig. 8) [33,34]. In that way, surfactin is typically composed of saturated C_{12} to C_{19} FA of the
linear, iso or anteiso type of branching [35]. Beside an increased production of surfactin, we
also observed an effect on the pattern of surfactin variants synthesized by *B. velezensis* in
the presence of artificial plant exudates, as well as in naturally produced exudates and in
planta upon root colonization (Supplementary Fig. 9). Indeed, UPLC-MS profiling revealed
that the surfactin pattern produced by GA1 in REM medium is enriched in surfactin *iso*-C_{14}
(*i*C_{14}) and other variants compared to LB medium (Fig. 5a). They correspond to variants of
the canonical structure with substitution of Leu by Val for the last residue of the cyclic
peptide moiety (Val$_7$) and, to a much lower extent, to the same substitution in position 2
(Val$_2$, Fig. 5a, see Supplementary Fig. 10). Valine is used both as precursor for the synthesis
of branched fatty acids with an even number of carbons, and as a building block by the NRPS
to form the peptide moiety. Supplementation of the medium with deuterated L-Val-d^8
resulted in an additional increase in the proportions of surfactin *iso*-C14 and Val$_7$ isoforms
labeled at the expected positions in the peptide and in the fatty acid tail (Supplementary Fig.
11). Based on these data, the higher relative proportions of *i*C_{14}Val^7 formed in REM, but also
in planta (Fig. 5c), most probably result from some enrichment of the intracellular pool in
valine upon growth in the presence of root exudates (Supplementary Discussion). Given the
reduced specificity of NRPS domains involved in selection and activation of leucine at
positions 2 and 7, the megaenzyme would preferably bind valine as it is more available in the
pool.

As already described in *B. subtilis* [36,37], the pleiotropic regulator CodY acts as
repressor of surfactin synthesis in *B. velezensis* GA1 as illustrated by the 1.9-fold increase in
production by the Δ*codY* mutant of strain GA1. Interestingly, CodY activity/*codY* expression is

also itself negatively impacted by high cellular concentrations in branched chain amino acids
[38]. Both quantitative and qualitative changes in surfactin production upon growth in
exudates could therefore be, at least partly, due to a lower CodY activity (Supplementary
Discussion). In support to the role played by this regulator, a similar impact on surfactin
pattern were observed by deleting *codY* in GA1 or by supplementing the culture medium of
the wild-type with valine (Fig. 5d).

217 Long fatty acid chain surfactins act as key triggers of receptor-independent plant immunity

Based on the potential of surfactin as host immunity elicitor [9,39], we next wanted to
evaluate the possible relevance of quantitative and qualitative modulation of the surfactin
pattern driven by the plant for its own benefit.

Upon application as root treatment, pure surfactin used as mixture of isoforms
formed in REM medium, induced systemic resistance in hydroponically-grown tobacco plants
providing approximatively 45-50% significant disease reduction on leaves subsequently
infected with the pathogen *Botrytis cinerea* (Fig. 6a). The various isoforms were then HPLC-
purified and tested individually revealing that only long fatty acid homologues (C_{14}/C_{15})
provided systemic protection to a similar level whereas short fatty acid homologues (C_{12}/C_{13})
were inactive (Fig. 6b). Moreover, plant immunization by surfactin is dose-dependent and
concentrations up to 5 µM are sufficient to significantly stimulate ISR (Fig. 6c). Interestingly,
such low µM concentration are actually in the range of those that could accumulate in the
root vicinity within a few days upon colonization by GA1 (Supplementary Fig. 12).

We next wanted to correlate this systemic protection induced by the lipopeptide
with its potential to trigger locally early immune-related events such as the extracellular
burst in reactive oxygen species (ROS) involved in defense and signaling in pathogen-
triggered immunity (PTI) [40,41]. By contrast with flagellin (epitope Flg22), one of the best
characterized Microbe-Associated Molecular Patterns (MAMPs) isolated from bacterial
pathogens, treatment with surfactin did not induced burst in apoplastic ROS in root tissues
(Fig. 6d). However, surfactin-mediated ROS signaling still occurs since a clear cytoplasmic
ROS accumulation was observed (Fig 6e). Little information is available about the spatio-
temporal dynamics of such ROS burst but it may originate from different organelles and has
been occasionally described in response to perception of biotic and abiotic stresses [42,43].

Using cytoplasmic ROS as marker, the same trend as for ISR tests could be observed

regarding the influence of the structure on the activity of surfactin since long fatty acid

homologues but not short ones efficiently stimulated early immune reaction (Fig 6f). This

means that a single additional methylene group in the fatty acid tail of the molecule (C_{14}

versus C_{13}) likely determines its immunization potential (Fig 6b,f). By contrast, substitution of

Leu^7 by a Val in the C_{14} homologue does not impact activity suggesting that the peptide

moiety is not essential for perception by plant cells. In addition, the µM concentrations

required for optimal eliciting activity of surfactin are very high compared with PAMPs active

in the nM range [44]. Our previous data showed that surfactin elicitation is still active after

treating cells with proteases or after a first application indicating that there is no saturation

of high-affinity but low abundance binding sites on receptors [45,46]. All this indicates that

surfactin is perceived by plant cells via a mechanism independent of high-affinity pattern-

recognition receptors (PRRs) involved in MAMP perception [40,41,44,47,48]. We therefore

postulated that surfactin perception relies on some interaction with the lipid phase of the

plant plasma membrane. Binding experiments via isothermal titration calorimetry and

leakage assays based on the release of fluorescent probe were performed using liposomes

prepared with lipids specific to plant plasma membrane (PLPC/sitosterol/glucosylCeramide).

It revealed that long fatty acid homologues have a higher affinity for these vesicles than the

short fatty acid forms and display a higher destabilizing effect on the lipid bilayer when

added at concentrations of 5 µM or higher (Fig. 6gh). These biophysical data thus correlated

well with the contrasting biological activities of longer C_{14}/C_{15} and shorter C_{12}/C_{13} surfactin

homologues.

A similar cytosolic and dose-dependent ROS signature was observed in tomato roots

upon surfactin treatment (Supplementary Fig. 13). According to the priming concept [49], we

previously showed that ISR triggered by the lipopeptide in that plant as well as in tobacco

and *Arabidopsis*, is not associated with a fast and strong expression of defensive

mechanisms before pathogen infection [20,39]. In order to verify that surfactin elicitation does

not cause a massive release of antimicrobials from plant tissues, tomato roots were pre-

treated with the lipopeptide before inoculation with *B. velezensis*. As expected, it did not

impacted subsequent colonization in terms of rate and dynamics compared to untreated

plants indicating the absence of potential adverse effects on the bacterial partner (Fig. 6i).

Concluding remarks

A large part of the interactions between bacteria and plants is known to be mediated

by small-size secreted products [50]. However, a better understanding of the chemical cross-

talk at the plant-bacteria interface and its impact on bacterial ecology, plant fitness and

immune responses remains challenging. In epiphytic soil bacilli, root exudates induce

expression of an array of genes involved in various functions such as chemotaxis and

nutrient acquisition [51–53]. Our data further illustrate that utilization of this cocktail of

molecules released by roots but also the perception of some cell wall polymers may also

drive these bacteria to efficiently produce key components of the secondary metabolome

and more specifically the multifunctional surfactin lipopeptide [20]. As an amphiphilic

molecule and powerful biosurfactant, surfactin is presumably viewed as membrane active

compound with potent antimicrobial activity. However, this lipopeptide is poorly

antibacterial and antifungal [54]. In *B. velezensis*, more obvious ecological functions of this

CLP are to contribute to motility, biofilm formation and roots colonization. An enhanced

production upon host perception thus constitutes a major force driving successful

rhizosphere establishment.

Homogalacturonan acts as a cue to enhance surfactin secretion by bacterial cells but

no transcriptional induction of the corresponding biosynthesis operon was observed.

Surfactin synthesis is integrated in a complex network involving several pleiotropic

regulators acting directly or indirectly on the expression of the *srfA* operon [55–58]. However,

we hypothesize that surfactin induction by HG may rather rely on post-transcriptional

changes as reported for the effect of the DegU and YczE regulators on production of another

CLP, bacillomycin D [59]. Despite the relatively close genetic proximity of the two species, our

data suggest that regulation of surfactin could be slightly different in *B. velezensis* and *B.*

subtilis. As it represents a key infochemical devoted to cross-talk with the host plant,

surfactin regulation may have been fine-tuned in rhizosphere species to better fit with the

nutritional or more broadly ecological context.

Deciphering the mechanism by which *B. velezensis* recognizes pectin and enhances

surfactin production would help to identify candidate genes and pathways that are

responsible for plant sensing, ensuring persistence on roots which globally remains very

poorly known for beneficial rhizobacteria. We are currently investigating whether some cell

surface proteins may act as receptors for homogalacturonan perception and binding as recently described for *Sphingomonas* sp. [60], another beneficial species living in association with plants [61]. Some insights could be obtained by scrutinizing the few genes conserved in *B. velezensis* but missing in non-plant-associated *B. amyloliquefaciens* strains that are not responsive to pectin [62]. Interestingly, shorter fragments of HG and galacturonic acid do not stimulate surfactin secretion. It is therefore tempting to hypothesize that sensing unaltered polymer could indicate a healthy host suitable for bacterial colonization while the perception of monomers or low DP oligomers may reflect a dead or infected plant that is unable to adequately provide resources.

Our data illustrate for the first time that *B. velezensis* can also modulate qualitatively its surfactin pattern by growing in its natural nutritional context, *i.e.* on root exudates. Substitution of leucine by valine in the peptide part is not expected to impact the contribution of the lipopeptide to colonization by the producing strain itself considering the minor effect of these structural changes on motility and biofilm formation potential [18]. Small modifications in the peptide sequence may nevertheless avoid surfactin hijacking for use as signal prompting heterologous biofilm formation by closely related competitor species [18]. Based on our observations, the most obvious benefit of an increased proportion of long fatty acid chain homologues is for the host plant since they represent the most active forms for priming immunity with no impact on host fitness [20,39], by contrast with PTI [63,64]. As the bacterial partner does not have to face strong defensive responses from this reaction, it ensures positive mutualistic co-habitation allowing establishment of populations on roots. Persistence of threshold populations is necessary for consistent production of other specialized secondary metabolites more directly involved in warding off both microbial competitors and plant soilborne pathogens in the context of biocontrol.

Surfactin stimulation upon sensing host molecular patterns may thus reflect an aspect of plant-*Bacillus* coevolution as it makes a shared good out of this multifunctional lipopeptide. To some extent, it might represent a facet of the plant-driven selection process resulting in active recruitment of this bacterium as species that provides beneficial functions. Other bacterial genera such as *Pseudomonas* also prevailing in the rhizosphere microbiome actively produce CLPs with similar roles as surfactin. Evaluating whether their synthesis is also modulated by plant cues would conceptually allow broadening the significance of these

335   lipopeptide-mediated inter-kingdom interactions for bacterial ecology, plant health and

336   biocontrol.

Methods

Bacterial media and growth conditions. Cultures were performed at 26°C in root exudates mimicking medium (EM)[27] or in LB medium. To test the effect of plant cell wall polymers, each specific plant polysaccharide was added at a final concentration of 0.1% in the culture medium. Low (HGLM, <5%) and high (HGHM, >95%) methylated homogalacturonan were provided from Elicityl Oligotech whereas oligogalacturonides and D-galacturonic acid were provided from Sigma.

Strains construction. All the bacteria strains used in this study are listed in Supplementary table 3. All the primers used in this study are available upon request. To follow the expression level of the *srf* operon in GA1, we constructed a *gfp* transcriptional fusion under the control of the *srf* promoter and integrated it into the *amyE* locus. First, a GA1 *amyE* amplicon containing a native *KasI* restriction site was integrated in the PGEMT easy. In parallel, a *cat-gfp* cassette containing respectively (i) a chloramphenicol resistance gene (cat) and (ii) a promoterless *gfpmut3.1* gene was amplified with primers containing *KasI* sites at their 5' extremities using the pGFP star as a matrix [65]. The pGEMT *amyE* plasmid and the *cat-gfp* amplicon were both digested by *KasI* (NEB) and the two linear fragments with compatible 5' overhangs were ligated together to obtain the PGEMT *amyEup-cat-gfp-amyEdw* plasmid. To construct the final mutation cassette, an overlap extension PCR was assessed by following the method developed by Bryksin and Matsumura [66]. One first fragment containing the upper *amyE* homologous region and the *cat* gene, and one second fragment englobing the *gfpmut3.1* gene and the lower *amyE* homologous region were both amplified using the PGEMT *amyEup-cat-gfp-amyEdw* plasmid as a matrix. A third fragment was amplified using GA1 genome as matrix with chimeric primers designed to obtain a *srf* promoter amplicon flanked by 20 bp connectors in 5' and 3' containing respectively homologies to the upper and lower *amyE* fragments. All three fragments were joined together with a second PCR race to obtain the final cassette. *B. velezensis* GA1 transformation was performed after modification from the protocol developped by Jarmer *et al.* [67].Briefly, one colony was inoculated into LB liquid medium at 37°C (160 rpm) during 6h and cells were washed two times with peptone water. Until 1µg of the recombinant cassette was added to the GA1 cells suspension adjusted to an OD_{600nm} of 0.01 into MMG liquid medium (19 g l-1 K_2HPO_4 anhydrous; 6 g l-1 KH_2PO_4; 1 g l-1 Na_3 citrate anhydrous; 0.2

g l-1 MgSO$_4$ 7H$_2$O; 2 g l-1 Na$_2$SO4; 50 µM FeCl$_3$ (sterilized by filtration at 0.22 µm); 2µM

MnSO$_4$; 8 g l-1 Glucose; 2 g l-1 L-glutamic acid; pH 7.0). Cells were incubated at 37°C with

shaking, and colonies who integrated the cassette by a double crossing over event were

selected on LB plate supplemented with chloramphenicol. Proper integration of the *cat-gfp*

locus was verified by PCR. Knock-out mutant strains were constructed by gene replacement

by homologous recombination. A cassette containing a chloramphenicol resistance gene

flanked respectively by 1 kb of the upstream region and 1 kb of the downstream region of

the targeted gene was constructed by a three partners overlap PCR. This recombination

cassette was also introduced in *B. velezensis* GA1 by inducing natural competence as

described above [67]. Double homologous recombination event was selected by

chloramphenicol resistance. Deletion was confirmed by PCR analysis with the corresponding

upstream and downstream primers.

Fluorescence measurement. Fluorescence accumulation was evaluated thanks to the

channel FL1 of a BD accuri C6 flow cytometer (Biosciences) with the following parameters:

20000 events, medium flow rate (35 µl.min^{-1}), FSC threshold of 20000.

Genome sequencing. GA1 genome sequence was reconstructed using a combined approach

of two sequencing technologies which generated short paired end reads and long reads. The

resulted sequences were then used for hybrid assembly. More precisely, genomic DNA was

extracted and purified from *B. velezensis* GA1 using the GeneJET Genomic DNA purification

(ThermoFisher scientific). First half of extracted DNA was sent to the GIGA sequencing

facility (Liège, Belgium), and use as DNA template for illumina MiSeq sequencing after being

prepared using nextera library kit illumina. Sequencing run generated 150 bp paired-end

read, which were trimmed and corrected using an in-house python script and SPAdes 3.14

[68] before assembly. The second half of extracted DNA was used to generate long reads with

a MinION Oxford nanopore plateform. DNA library was constructed using the Rapid

Sequencing kit (SQK-RAD0004, Oxford nanopore). Adapters were trimmed from generated

reads with Porechop software (https://github.com/rrwick/Porechop). Trimmed reads were

then filtered by size (>500) and Q-score (>10) using NanoFilt implemented in NanoPack [69].

Finally, hybrid assembly was performed using hybridSPAdes algorithm implemented in

SPAdes 3.14 [70].

Transcriptome library preparation and sequencing. RNA extraction was performed for each

sample using the NucleoSpin RNA kit (Macherey-Nagel). Total RNAs were quantified using

Nanodrop (Thermofisher). For sequencing, all samples were sent to the GIGA-genomics

platform in Liège. Quality was assessed using the RNA 6000 Nano Chip on a 2100 Bioanalyzer

(Agilent). cDNA libraries were prepared employing Universal Prokaryotic RNA-Seq,

Prokaryotic AnyDeplete kit (Nugen) according to the manufacturer's instructions, with .

cDNA libraries were quantified and normalized by using the KAPA SYBR Fast Mastermix

(Sigma-Aldrich) with P5-P7 Illumina primers according to the manufacturer's instructions.

Prepared libraries were sequenced on a NextSeq 550 device (Illumina) by using the following

parameters : paired end, 80 cycles read 1, 8 cycles index, 80 cycles read 2.

RNA-seq data analysis. The raw RNA-seq reads were trimmed using Trimmomatic v0.39 [71].

We performed quality control on the trimmed reads using FastQC v0.11.8 (Babraham

Bioinformatics). Trimmed reads were mapped to the GA1 reference genome (see section

"genome sequencing" for accession numbers) using BWA-mem v0.7.17 [72] with the

following settings: mem -k 50 -B 40 -v 1. At least 95.4% of reads uniquely mapped to the

annotated reference genome (Supplementary table 4). SAMtools v1.9 [73] was used to

generate the BAM files and their indices. To calculate the read counts, the python-based tool

HTSeq v0.9 [74] was employed with the following parameters: htseq-count -q -s no -f. The

Cufflinks function cuffnorm [75] was used to generate the FPKM (fragments per kilobase of

transcript per million mapped reads) tables using the following settings: --compatible-hits-

norm --library-norm-method classic-fpkm. Genes with low reads counts (<25) were removed

before further analysis. Differential expression analysis was conducted according to the

DESeq2 pipeline (10.1186/s13059-014-0550-8) with cut-off parameters of $p < 0.05$ and log2-

fold-change > 1.5.

Motility and biofilm assays. Swarming motility assays were performed according to

Molinatto *et al.* 2017 [76]. Diameter of the bacterial swarming pattern was measured 48h

after inoculation on REM soft agar plates (0.8% agar) supplemented or not with 0.1% HGLM.

Quantification of total biofilm was performed by crystal violet staining. Strain of interest was

inoculated at a final OD_{600} of 0.1 in a 96 wells microplate containing 200 µl of REM medium

supplemented or not with 0.1 % HGLM. The plate was incubated at 30°C during 24h without

shaking. Medium and planctonic cells were discarded and wells were washed with PBS.

Biofilm pellicle was stained with 0.1% crystal violet during 10 min and washed with PBS. The

stained biofilm was dissolved with 30% acetic acid. Absorbance was measured at 595 nm.

Plant growth conditions and roots colonization assays. For sterilization, tomato seeds were

first immersed in a 70% ethanol solution during 2 minutes, transferred in a 20% bleach

solution under shaking for 20 minutes and rinsed three times with sterile water. Sterilized

tomato seeds were pre-germinated on solid Hoagland medium at 22°C under a 16h/8h

night/day cycle. After 4 days, 5μL of cultures containing the strain of interest and calibrated

at $OD_{600}=1$ were deposited on the root top. After 1 and 3 days of colonization, roots were

harvested, deposited separately in a peptone water solution supplemented with 0.1% of

Tween, and vortexed vigorously to tear off the bacterial cells from the roots. Several

dilutions were plated on LB media to evaluate the level of colonization. Measurements of

surfactin production by GA1 cells colonizing roots were performed on 1x1x0.7 cm pieces of

gelified medium containing roots based on the assumption that the produced lipopeptide

diffused to a maximal distance of 5 mm from each part of the root and is uniformly

distributed over the surface as we previously observed via imaging-MS [77]. A 10-fold

concentration factor was applied to estimate concentrations around the root surface in

order to take into account diffusion constraints in a solid matrix. Surfactin was quantified by

UPLC-MS as described below.

Plant cell wall extraction. Tobacco seeds were sterilized as described above for tomato

seeds and deposited on Hoagland plates at 22°C during one week for a successful

germination process. Each plantlet was then transferred in a seedholder filled with soft agar

and put in Araponics boxes containing the nutritive solution described above. Cell wall

extraction was performed on 6 weeks old plants grown at 22°C with a 16h/8h day-night

alternance. Roots were harvested, lyophilized and reduced to powder using a Retsch MM400

grinder. 500 mg of powder was resuspended in 40 ml of ethanol 80% at 90°C for 20 min. The

insoluble cell wall fraction was recovered by centrifugation and the pellet obtained was

washed once with water to obtain the Alcoholic Insoluble Residue (AIR) used for

fractionation. AIR was freeze-dried before use in fractionation protocol. Sequential

extraction of root cell walls was performed using a protocol derived from Carpita [78] and

Silva *et al.* [79]. Dry AIR was resuspended in 40 ml water and incubated at 100°C for 20 min.

Supernatant was recovered after centrifugation as a soluble pectic fraction (cPEC).

LC-MS analyses. Detection of metabolites and quantification was performed by LC-MS. 10 µL of samples were analyzed using UPLC–MS with UPLC (Acquity H-class, Waters) coupled to a single quadrupole mass spectrometer (SQD mass analyzer, Waters) using an C18 column (Acquity UPLC BEH C18 2.1 mm × 50 mm, 1.7 µm). Elution was performed at 40°C with a constant flow rate of 0.6 mL/min using a gradient of Acetonitrile (solvent B) and water (solvent A) both acidified with 0.1% formic acid as follows: starting at 15% B during 2 min, solvent B was then raised from 15% to 95% in 5 min and maintained at 95% up to 9.5 min before going back to initial conditions at 9.8 min during 3 minutes before next injection if needed. Compounds were detected in electrospray positive ion mode by setting SQD parameters as follows: source temperature 130°C; desolvation temperature 400°C, and nitrogen flow: 1000 L.h-1 with mass range from m/z 800 to 1550. Surfactins were quantified based on their retention times and masses compared to commercial standards (98% purity, Lipofabrik).

Induction of systemic resistance and ROS measurements. ISR assays were performed as previously described [39] on 4 weeks-old tobacco plants cultivated in hydroponic conditions using Hoagland solution as nutrient base. Plants were treated with pure surfactin at the root level and infected on leaves by applying a spore suspension of the phytopathogen *Botrytis cinerea* prepared as detailed previously [39]. Spreading lesions occurred starting from 48h post-infection and the diameter size was measured two days later. Five plants were used per treatment and experiments were repeated independently at least twice. For determination of cytoplasmic ROS stimulation, fluorescent probe (DCFH-DA) was used. Plants used in this experiment were grown on Hoagland medium for two weeks as described above. Experiments were performed on nine samples per treatment each containing three root segments (approx 100 mg FW) collected from different plants (n=9). Roots were treated with 50 µM DCFH-DA for 10 minutes, rinsed with PBS upon removing the probe, and finally treated. All the operations were conducted in a 96-well black microplate. Fluorescence measurements were performed on a Spark (Tecan) microplate reader (exc 485 nm; em = 535 nm) with readings every 10 minutes. Stimulation of apoplastic hydrogen peroxide production in root cells was measured via chemiluminescence (ferricyanide-catalysed oxidation of luminol). Means and standard deviations were calculated from measurements performed on three samples per treatment, each containing three root segments

(approximatively 100 mg FW) collected from different plants. Extracellular ROS in tomato

roots was conducted according to Bisceglia *et al.* [80] with minor changes. Namely, instead of

leaf discs, tomato roots, three segments (approximatively 100 mg FW from the same plant)

per sample, were used. Plants were grown for two weeks on Hoagland medium, and

chemiluminescence was measured in Tecan Spark plate reader.

ITC analysis. ITC analyses were performed with a VP-ITC Microcalorimeter (Microcal). The

calorimeter cell (volume of 1.4565 mL) was filled with a 10 µM (below the CMC

concentration) surfactin solution in buffer (Tris 10mM, NaCl 150mM, 1mM EDTA at pH 8.5).

The syringe was filled with a suspension of LUV at a lipid concentration of 5 mM. A series of

10µl injections was performed at constant time intervals (6 min) at 25°C. The solution in the

titration cell was stirred at 305 RPM. Prior to each analysis, all solutions were degassed using

a sonicator bath. The heats of dilution of vesicles were determined by injecting vesicles in

buffer and subtracted from the heats determined in the experiments. Data were processed

by software Origin 7 (Originlab) using the cumulative model described by Heerklotz and

Seelig [81]. All measurements were repeated at least three times with two different vesicle

preparations.

Leakage assays. Membrane permeabilization was followed as described by Van Bambeke *et*

al. [82]. Release of 8-hydroxypyrene-1,3,6 trisulfonic acid (HTPS) coentrapped with and

quenched by p-xylene-bis-pyridinium bromide (DPX) from liposomes can be monitored by

the fluorescence increase upon dilution following their leakage from the vesicles. Surfactin

C12 or Surfactin C14 was added from a stock solution in DMSO and fluorescence intensities

were immediately recorded. The percentage of HPTS released was defined as [(Ft-

Fcontr)/(Ftot- Fcontr)] /100, where Ft is the fluorescence signal measured after 15 min in the

presence of Surfactin C12 or Surfactin C14, Fcontr is the fluorescence signal measured at the

same time for control liposomes, and Ftot is the total fluorescence signal obtained after

complete disruption of the liposomes by 0.05% Triton X-100. All fluorescence determinations

were performed at room temperature on a Perkin Elmer LS-50B Fluorescence

Spectrophotometer (Perkin-Elmer Ltd.) using λexc of 450 nm and a λem of 512 nm.

Statistical analyses. All statistical analyses were performed on GraphPad prism. Before each

statistical analysis, variance homoscedasticity was verified by using a Brown-Forsythe test.

ANOVA analysis was used for multiple comparison and significant differences were indicate

by different letters. Statistical differences between means were evaluated by two-tailed

Student's t-test. Number or biological replicates used for each experiment are indicated in

the corresponding figure legend. P-Values are indicated in the figure legends.

Data availability

The RNA-seq datasets produced for this study are deposited at https://www.ebi.ac.uk/ena/

under the project reference PRJEB39762. All other datasets analyzed for this study are

included in the supplementary files. The Genome Resulting assembly of the GA1 strain was

deposited in the GenBank database under the accession numbers CP046386 and CP046387.

Acknowledgment

This work was supported by the EU Interreg V France-Wallonie-Vlaanderen portfolio SmartBiocontrol (Bioprotect and Bioscreen projects, avec le soutien du Fonds européen de développement régional - Met steun van het Europees Fonds voor Regionale Ontwikkeling), by the PDR research project ID 26084552 from the F.R.S.-FNRS (National fund for Scientific Research in Belgium) and by the EOS project ID 30650620 from the FWO/F.R.S.-FNRS. FB is recipient of a F.R.I.A. fellowship (Formation à la Recherche dans l'Industrie et l'Agriculture) and MO is senior research associate at the F.R.S.-F.N.R.S. We are grateful to the KU Leuven HPC infrastructure and the Flemish Supercomputer Center (VSC) for providing the computational resources and services to perform the RNA-seq analysis. We gratefully acknowledge Claire Bertrand and Loïc Ongena for critically reading the manuscript.

Bibliography

1. Andrews, J. H. & Harris, R. F. The Ecology and Biogeography of Microorganisms on Plant Surfaces. *Annu. Rev. Phytopathol.* **38**, 145–80 (2000).

2. Zhalnina, K. *et al.* Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. *Nat. Microbiol.* **3**, 470–480 (2018).

3. Vieira, S. *et al.* Drivers of the composition of active rhizosphere bacterial communities in temperate grasslands. *ISME J.* **14**, 463–475 (2020).

4. Vacheron, J. *et al.* Plant growth-promoting rhizobacteria and root system functioning. *Front. Plant Sci.* **4**, 1–19 (2013).

5. Backer, R. *et al.* Plant growth-promoting rhizobacteria: Context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Frontiers in Plant Science* vol. 871 (2018).

6. Pieterse, C. M. J. *et al.* Induced Systemic Resistance by Beneficial Microbes. *Annu. Rev. Phytopathol.* **52**, 347–375 (2014).

7. Köhl, J., Kolnaar, R. & Ravensberg, W. J. Mode of action of microbial biological control agents against plant diseases: Relevance beyond efficacy. *Front. Plant Sci.* **10**, 1–19 (2019).

8. van Loon, L. C., Bakker, P. A. H. M. & Pieterse, C. M. J. Systemic Resistance Induced By Rhizosphere Bacteria. *Annu. Rev. Phytopathol.* **36**, 453–483 (1998).

9. Ongena, M. *et al.* Surfactin and fengycin lipopeptides of *Bacillus subtilis* as elicitors of induced systemic resistance in plants. *Environ. Microbiol.* **9**, 1084–1090 (2007).

10. Ongena, M. & Jacques, P. *Bacillus* lipopeptides: versatile weapons for plant disease biocontrol. *Trends Microbiol.* **16**, 115–125 (2008).

11. Wu, K. *et al.* Pectin enhances bio-control efficacy by inducing colonization and secretion of secondary metabolites by *Bacillus amyloliquefaciens* SQY 162 in the rhizosphere of tobacco. *PLoS One* **10**, 1–17 (2015).

12. Saxena, A. K., Kumar, M., Chakdar, H., Anuroopa, N. & Bagyaraj, D. J. *Bacillus* species in soil as a natural resource for plant health and nutrition. *J. Appl. Microbiol.* **128**, 1583–1594 (2020).

13. Chen, X. H. *et al.* Comparative analysis of the complete genome sequence of the plant growth-promoting bacterium *Bacillus amyloliquefaciens* FZB42. *Nat. Biotechnol.* **25**, 1007–1014 (2007).

14.   Chen, X. H. *et al.* Genome analysis of *Bacillus amyloliquefaciens* FZB42 reveals its potential for
biocontrol of plant pathogens. *J. Biotechnol.* **140**, 27–37 (2009).

15.   Molinatto, G. *et al.* Complete genome sequence of *Bacillus amyloliquefaciens subsp.*
plantarum S499, a rhizobacterium that triggers plant defences and inhibits fungal
phytopathogens. *J. Biotechnol.* **238**, 56–59 (2016).

16.   van Gestel, J., Vlamakis, H. & Kolter, R. From Cell Differentiation to Cell Collectives: *Bacillus*
subtilis Uses Division of Labor to Migrate. *PLoS Biol.* **13**, 1–29 (2015).

17.   López, D., Vlamakis, H., Losick, R. & Kolter, R. Paracrine signaling in a bacterium. *Genes Dev.*
23, 1631–1638 (2009).

18.   Aleti, G. *et al.* Surfactin variants mediate species-specific biofilm formation and root
colonization in *Bacillus*. *Environ. Microbiol.* **18**, 2634–2645 (2016).

19.   Pršić, J. & Ongena, M. Elicitors of Plant Immunity Triggered by Beneficial Bacteria. *Front. Plant*
Sci. **11**, 1–12 (2020).

20.   Debois, D. *et al.* Plant polysaccharides initiate underground crosstalk with bacilli by inducing
synthesis of the immunogenic lipopeptide surfactin. *Environ. Microbiol. Rep.* **7**, 570–582
(2015).

21.   Gage, D. J. Infection and Invasion of Roots by Symbiotic, Nitrogen-Fixing Rhizobia during
Nodulation of Temperate Legumes. *Microbiol. Mol. Biol. Rev.* **68**, 280–300 (2004).

22.   Beauregard, P. B., Chai, Y., Vlamakis, H., Losick, R. & Kolter, R. *Bacillus subtilis* biofilm
induction by plant polysaccharides. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E1621-30 (2013).

23.   Fan, B. *et al.* *Bacillus velezensis* FZB42 in 2018: The gram-positive model strain for plant
growth promotion and biocontrol. *Front. Microbiol.* **9**, 1–14 (2018).

24.   Mohnen, D. Pectin structure and biosynthesis. *Curr. Opin. Plant Biol.* **11**, 266–277 (2008).

25.   Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P. M. & Henrissat, B. The
carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* **42**, 490–495 (2014).

26.   Fan, B., Blom, J., Klenk, H. P. & Borriss, R. *Bacillus amyloliquefaciens*, *Bacillus velezensis*, and
Bacillus siamensis Form an 'Operational Group *B. amyloliquefaciens*' within the *B. subtilis*
species complex. *Front. Microbiol.* **8**, 22 (2017).

27.   Nihorimbere, V. *et al.* Impact of rhizosphere factors on cyclic lipopeptide signature from the
plant beneficial strain *Bacillus amyloliquefaciens* S499. *FEMS Microbiol. Ecol.* **79**, 176–191
(2012).

28. Chen, Y. *et al.* Biocontrol of tomato wilt disease by *Bacillus subtilis* isolates from natural
environments depends on conserved genes mediating biofilm formation. *Environ. Microbiol.*
15, 848–864 (2013).

29. Grau, R. R. *et al.* A duo of potassium-responsive histidine kinases govern the multicellular
destiny of *Bacillus subtilis*. *MBio* **6**, 1–16 (2015).

30. Thérien, M. *et al.* Surfactin production is not essential for pellicle and root-associated biofilm
development of *Bacillus subtilis*. *Biofilm* **2**, 100021 (2020).

31. Vlamakis, H., Chai, Y., Beauregard, P., Losick, R. & Kolter, R. Sticking together: Building a
biofilm the *Bacillus subtilis* way. *Nat. Rev. Microbiol.* **11**, 157–168 (2013).

32. Caramori, T., Barillà, D., Nessi, C., Sacchi, L. & Galizzi, A. Role of FlgM in σD-dependent gene
expression in *Bacillus subtilis*. *J. Bacteriol.* **178**, 3113–3118 (1996).

33. Strieker, M., Tanović, A. & Marahiel, M. A. Nonribosomal peptide synthetases: Structures and
dynamics. *Curr. Opin. Struct. Biol.* **20**, 234–240 (2010).

34. Süssmuth, R. D. & Mainz, A. Nonribosomal Peptide Synthesis—Principles and Prospects.
Angew. Chemie - Int. Ed. **56**, 3770–3821 (2017).

35. Diomandé, S. E., Nguyen-The, C., Guinebretière, M. H., Broussolle, V. & Brillard, J. Role of fatty
acids in *Bacillus* environmental adaptation. *Front. Microbiol.* **6**, 1–20 (2015).

36. Serror, P. & Sonenshein, A. L. CodY is required for nutritional repression of Bacillus subtilis
genetic competence. *J. Bacteriol.* **178**, 5910–5915 (1996).

37. Dhali, D. *et al.* Genetic engineering of the branched fatty acid metabolic pathway of *Bacillus*
subtilis for the overproduction of surfactin C14 isoform. *Biotechnol. J.* **12**, 1–10 (2017).

38. Brinsmade, S. R., Kleijn, R. J., Sauer, U. & Sonenshein, A. L. Regulation of CodY activity through
modulation of intracellular branched-chain amino acid pools. *J. Bacteriol.* **192**, 6357–6368
(2010).

39. Cawoy, H. *et al.* Plant Defense Stimulation by Natural Isolates of *Bacillus* Depends on Efficient
Surfactin Production. *Mol. Plant-Microbe Interact.* **27**, 87–100 (2014).

40. Saijo, Y., Loo, E. P. iian & Yasuda, S. Pattern recognition receptors and signaling in plant–
microbe interactions. *Plant J.* **93**, 592–613 (2018).

41. Waszczak, C., Carmody, M. & Kangasjärvi, J. Reactive Oxygen Species in Plant Signaling. *Annu.*
Rev. Plant Biol. **69**, 209–236 (2018).

42.   Mignolet-Spruyt, L. *et al.* Spreading the news: Subcellular and organellar reactive oxygen
638   species production and signalling. *J. Exp. Bot.* **67**, 3831–3844 (2016).

639   43.   Ashtamker, C., Kiss, V., Sagi, M., Davydov, O. & Fluhr, R. Diverse subcellular locations of
640   cryptogein-induced reactive oxygen species production in tobacco bright yellow-2 cells. *Plant*
641   *Physiol.* **143**, 1817–1826 (2007).

642   44.   Bigeard, J., Colcombet, J. & Hirt, H. Signaling mechanisms in pattern-triggered immunity (PTI).
643   *Mol. Plant* **8**, 521–539 (2015).

644   45.   Jourdan, E. *et al.* Insights into the defense-related events occurring in plant cells following
645   perception of surfactin-type lipopeptide from Bacillus subtilis. *Mol. Plant-Microbe Interact.* **22**,
646   456–468 (2009).

647   46.   Henry, G., Deleu, M., Jourdan, E., Thonart, P. & Ongena, M. The bacterial lipopeptide surfactin
648   targets the lipid fraction of the plant plasma membrane to trigger immune-related defence
649   responses. *Cell. Microbiol.* **13**, 1824–1837 (2011).

650   47.   Zipfel, C. & Oldroyd, G. E. D. Plant signalling in symbiosis and immunity. *Nature* **543**, 328–336
651   (2017).

652   48.   Schellenberger, R. *et al.* Apoplastic invasion patterns triggering plant immunity: plasma
653   membrane sensing at the frontline. *Mol. Plant Pathol.* **20**, 1602–1616 (2019).

654   49.   Martinez-Medina, A. *et al.* Recognizing Plant Defense Priming. *Trends Plant Sci.* **21**, 818–822
655   (2016).

656   50.   Levy, A. *et al.* Genomic features of bacterial adaptation to plants. *Nat. Genet.* **50**, 138–150
657   (2018).

658   51.   Fan, B. *et al.* Transcriptomic profiling of *Bacillus amyloliquefaciens* FZB42 in response to maize
659   root exudates. *BMC Microbiol.* **12**, (2012).

660   52.   Feng, H. *et al.* Identification of chemotaxis compounds in root exudates and their sensing
661   chemoreceptors in plant-growth-promoting rhizobacteria *Bacillus amyloliquefaciens* SQR9.
662   *Mol. Plant-Microbe Interact.* **31**, 995–1005 (2018).

663   53.   Zhang, N. *et al.* Whole transcriptomic analysis of the plant-beneficial rhizobacterium *Bacillus*
664   *amyloliquefaciens* SQR9 during enhanced biofilm formation regulated by maize root exudates.
665   *BMC Genomics* **16**, 1–21 (2015).

666   54.   Raaijmakers, J. M., de Bruijn, I., Nybroe, O. & Ongena, M. Natural functions of lipopeptides
667   from *Bacillus* and *Pseudomonas*: More than surfactants and antibiotics. *FEMS Microbiol. Rev.*

34, 1037–1062 (2010).

55.   Hayashi, K. *et al.* The H_2O_2 Stress-Responsive Regulator PerR Positively Regulates *srfA*
Expression in *Bacillus subtilis*. **187**, 6659–6667 (2014).

56.   Wolf, D. *et al.* The quorum-sensing regulator ComA from *Bacillus subtilis* activates
transcription using topologically distinct DNA motifs. *Nucleic Acids Res.* **44**, 2160–2172 (2015).

57.   Mariappan, A., Makarewicz, O., Chen, X. H. & Borriss, R. Two-component response regulator
DegU controls the expression of bacilysin in plant-growth-promoting bacterium *Bacillus*
amyloliquefaciens FZB42. *J. Mol. Microbiol. Biotechnol.* **22**, 114–125 (2012).

58.   Zhi, Y., Wu, Q. & Xu, Y. Genome and transcriptome analysis of surfactin biosynthesis in
Bacillus amyloliquefaciens MT45. *Sci. Rep.* **7**, 1–13 (2017).

59.   Koumoutsi, A., Chen, X. H., Vater, J. & Borriss, R. DegU and YczE positively regulate the
synthesis of bacillomycin D by *Bacillus amyloliquefaciens* strain FZB42. *Appl. Environ.*
Microbiol. **73**, 6953–6964 (2007).

60.   Konishi, H., Hio, M., Kobayashi, M., Takase, R. & Hashimoto, W. Bacterial chemotaxis towards
polysaccharide pectin by pectin-binding protein. *Sci. Rep.* **10**, 1–12 (2020).

61.   Muller, D. B., Schubert, O. T., Rost, H., Aebersold, R. & Vorholt, J. A. Systems-level proteomics
of two ubiquitous leaf commensals reveals complementary adaptive traits for phyllosphere
colonization. *Mol. Cell. Proteomics* **15**, 3256–3269 (2016).

62.   Hossain, M. J. *et al.* Deciphering the conserved genetic loci implicated in plant disease control
through comparative genomics of *Bacillus amyloliquefaciens subsp. plantarum*. *Front. Plant*
Sci. **6**, 1–14 (2015).

63.   Birkenbihl, R. P., Liu, S. & Somssich, I. E. Transcriptional events defining plant immune
responses. *Curr. Opin. Plant Biol.* **38**, 1–9 (2017).

64.   Huot, B., Yao, J., Montgomery, B. L. & He, S. Y. Growth-defense tradeoffs in plants: A
balancing act to optimize fitness. *Molecular Plant* (2014) doi:10.1093/mp/ssu049.

65.   Trauth, S. & Bischofs, I. B. Ectopic integration vectors for generating fluorescent promoter
fusions in *Bacillus subtilis* with minimal dark noise. *PLoS One* **9**, (2014).

66.   Bryksin, A. V. & Matsumura, I. Overlap extension PCR cloning: a simple and reliable way to
create recombinant plasmids. *Biotechniques* **48**, 463–465 (2010).

67.   Jarmer, H., Berka, R., Knudsen, S. & Saxild, H. H. Transcriptome analysis documents induced
competence of *Bacillus subtilis* during nitrogen limiting conditions. *FEMS Microbiol. Lett.* **206**,

197–200 (2002).

68. Bankevich, A. *et al.* SPAdes: A new genome assembly algorithm and its applications to single-
cell sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

69. De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C. NanoPack:
Visualizing and processing long-read sequencing data. *Bioinformatics* **34**, 2666–2669 (2018).

70. Antipov, D., Korobeynikov, A., McLean, J. S. & Pevzner, P. A. HybridSPAdes: An algorithm for
hybrid assembly of short and long reads. *Bioinformatics* **32**, 1009–1015 (2016).

71. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014).

72. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**, 1754–1760 (2009).

73. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).

74. Anders, S., Pyl, P. T. & Huber, W. HTSeq-A Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166–169 (2015).

75. Trapnell, C. *et al.* Transcript assembly and abundance estimation from RNA-Seq reveals
thousands of new transcripts and switching among isoforms. *Nat. Biotechnol.* **28**, 511–515
(2011).

76. Molinatto, G. *et al.* Key impact of an uncommon plasmid on *Bacillus amyloliquefaciens subsp.*
plantarum S499 developmental traits and lipopeptide production. *Front. Microbiol.* **8**, 1–18
(2017).

77. Debois, D., Ongena, M., Cawoy, H. & De Pauw, E. MALDI-FTICR MS imaging as a powerful tool
to identify *Paenibacillus* antibiotics involved in the inhibition of plant pathogens. *J. Am. Soc.*
Mass Spectrom. **24**, 1202–1213 (2013).

78. Carpita, N. C. Cell wall development in maize coleoptiles. *Plant Physiol.* **76**, 205–212 (1984).

79. Silva, G. B. *et al.* Cell wall polysaccharides from fern leaves: Evidence for a mannan-rich Type
III cell wall in *Adiantum raddianum*. *Phytochemistry* **72**, 2352–2360 (2011).

80. Bisceglia, N., Gravino, M. & Savatin, D. Luminol-based Assay for Detection of Immunity
Elicitor-induced Hydrogen Peroxide Production in *Arabidopsis thaliana* Leaves. *Bio-Protocol* **5**,
(2015).

81.   Heerklotz, H. & Seelig, J. Titration calorimetry of surfactant-membrane partitioning and
membrane solubilization. *Biochim. Biophys. Acta - Biomembr.* **1508**, 69–85 (2000).

82.   Van Bambeke, F. *et al.* Biophysical studies and intracellular destabilization of pH-sensitive
liposomes. *Lipids* **35**, 213–223 (2000).

Figure 1: Impact of pectin on early surfactin production. a Surfactin (cyclic structure represented up) production in a root exudates mimicking (REM) medium at early growth phase (OD_{600}=0.2) with (red chromatogram) or without (blue chromatogram) addition of crude pectin extract added to the GA1 cultures. The main peak represents C15 surfactin whereas the minor left and right peaks represents C14 and C16 surfactins, respectively. **b** Surfactin accumulation in the early (left panel, OD_{600}=0.2) and mid (right panel, OD_{600}=0.35) exponential growth phase of GA1 cultures in REM medium supplemented with different sized pectin fragments : homogalacturonan low methylated (HGLM), DP>150; oligogalacturonides (OG), DP=15; galacturonic acid (GA), DP=1. Means ± std err. from three biological replicates of one representative experiment are shown. Significate difference between each condition is indicated by different letters, p-value < 0.01. **c** Comparison of surfactin induction level by HGLM in the early exponential growth phase for different *Bacillus* species : *Bacillus velezensis* (*B; v*), *Bacillus amyloliquefaciens* (*B. a*), *Bacillus subtilis* (*B. s*) and *Bacillus pumilus* (*B. p*). For each strain tested, surfactin accumulation was normalized with the control condition without HGLM represented by the black dotted line. Means ± std err. from three biological replicates are shown.

Figure 2: Impact of the specific rhizosphere nutritional context on early surfactin production. a Evaluation of bacterial population (black line, left axis) and relative *srfAA* expression on roots (grey bars, right axis) in a time frame of seven days post inoculation (dpi). *Bacillus* progression on roots characterized by a biofilm formation was assessed by microscopy at each time point (upper part). **b** Surfactin production rate on roots. Means ± std err. from three biological replicates of one representative experiment are shown **c** Surfactin accumulation measured by UPLC-MS in a 8h time course experiment in REM medium (grey bars) compared to LB medium (red bars). Means ± std err. from three biological replicates of one representative experiment are shown *** P-value <0.001, ** P-value <0.01, * P-value <0.05 **d** Comparison of early surfactin accumulation (µM of surfactin on y axis linked to OD_{600} on x axis) in different *Bacillus* species, including *B. velezensis* (GA1 and S499 in green), *B. pumilus* (QST 2808 in orange), *B. amyloliquefaciens* (DSM 7 in red) and *B. subtilis* (ATCC 21332 in blue). Circle symbols are representing plant associated bacteria whereas triangle symbols are representing non-plant associated bacteria.

Figure 3: Ecological importance of an early surfactin accumulation. a Evaluation of HGLM and root exudates synergistic effect on early surfactin production. Time course experiment for surfactin quantification was performed in REM (grey curves) and LB (red curves) medium with (circle symbols) or without (square symbols) addition of HGLM. Means ± std err. from three biological replicates of one representative experiment are shown **b** Swarming potential of *B. velezensis* GA1 on soft agar plates after addition of HGLM or not. Means ± std err. from seven biological replicates of one representative experiment are shown. **c** Evaluation of *B. velezensis* ability to form pellicles on microwells plates after addition of HGLM or not. Means ± std err. from eight biological replicates of one representative experiment are shown. Pellicle formation is illustrated on the right **d** Comparison of *B. velezensis* GA1 WT and (red) and a Δ*srfAA* mutant (grey) for their swarming potential in a time course study. Means ± std err. from three biological replicates of one representative experiment are shown. Time course study is illustrated right. **e** Comparison of pellicle formation between GA1 WT strain (red) and a Δ*srfAA* mutant (grey). Means ± std err. from eight biological replicates of one representative experiment are shown **** P-value <0.0001. **f** *In vitro* comparison of root colonization ability of GA1 (red boxes) and GA1 Δ*srfAA* (grey boxes) on tomato plantlets. Means ± std err. from eight biological replicates of one representative experiment are shown *** P-value <0.001, ns non significant.

Figure 4: Impact of plant triggers perception on *Bacillus* transcriptome. a Surfactin expression measured by fluorescence in the GA1 P*srfAp::gfp* reporter strain in a 24h time course study in EM medium (grey bars) compared to LB medium (red bars). Means ± std err. from three biological replicates of one representative experiment are shown *** P-value <0.001. **b** Classification of the different genes carrying a significant fold change (1.5 log2) 5 and 8 hours after addition of HG when compared to the control condition. The outer circle represents the proportion of up (dark blue) and down (red) regulated genes. The inner circle represents the proportion of genes belonging to the different functional family described in the legend.

Figure 5: Qualitative impact of root exudates on surfactin production. a Comparison of surfactin pattern in REM and LB medium. Based on MS-MS analyses, nine different surfactin forms were identified (a: C_{12}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; b: C_{13}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; c : *iso*-C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; c': *n*-C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; d : C_{15}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; e: C_{13}-Glu-Leu-Leu-Val-Asp-Leu-Val ; f: C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Val ; g: C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Val and h: C_{14}-Glu-Val-Leu-Val-Asp-Leu-Val.) **b** Relative proportions of surfactin variants in LB, REM, REM supplemented with valine, and *in planta*. **c** Qualitative and quantitative role of CodY on surfactin production. In a WT strain, 95% of the surfactin molecules are carrying a Leu in position 7 (grey bars) and only 5% are carrying a Val (red bars) whereas in Δ*codY* mutant almost 25% of the surfactin molecules are carrying a Val in position 7 and 75% a Leu. In addition, amount of total surfactin production rate of 150 % can be observed in Δ*codY* mutant compared to WT strain. Proportion of iso-C14 is also affected by CodY, 36 % of total C_{14} are iso-fatty acid (grey bars) and 64% are linear (red bars) in WT strain whereas in Δ*codY* mutant 55% of C_{14} are *iso*-C_{14} and 45 % are linear. Again, total amount of C_{14} is higher in Δ*codY* mutant (increase of 190 %).

Figure 6: Impact of surfactin homologues on Solanaceae plant immunity. abc Systemic resistance induced in hydroponically-grown tobacco by surfactin and expressed as reduction of *B. cinerea* infection (illustration of the reduction in the diameter of spreading lesions on infected leaves) in plants treated at the root level prior to pathogen inoculation on leaves compared to control plants. Data represent results grouped from two independent experiments with similar results and each involving 5 plants with 4 lesions on the second leave (n=40). **a** Effect of surfactin homologues (SF mix) as naturally co-produced by the bacterium (C12/C13/C14/C15 in relative proportions 8/17/33/42%) **** P-value <0.0001 **b** Effect of HPLC-purified surfactin homologues applied at 10 µM with fatty acid chains from C12 to C15. Significate difference between each condition is indicated by different letters, p-value < 0.05 **c** Effect of the most active C14 homologue tested at various concentrations. Significate difference between each condition is indicated by different letters, p-value < 0.05 **de** Stimulation of oxidative burst in root tissues upon treatment with a SF mix and to the response observed by treating roots with flagellin (flg22, 1 µM) used as positive control. **d** Stimulation of apoplastic ROS accumulation (DCFH-DA fluorescent probe) in root tissues upon treatment with a surfactin mix applied at 15 µM. Means and standard deviations are shown for one representative experiment performed on nine samples per treatment each containing three root segments (approx 100 mg FW) collected from different plants (n=9). Similar trend was obtained in an independent assay. **e** Stimulation of cytoplasmic hydrogen peroxide production in root cells. Means and s.d. were calculated from measurements performed on three samples per treatment each containing three root segments (approx 100 mg FW) collected from different plants. Data represent values obtained from two independent experiments (n=6 per treatment). **f** Stimulation of cytoplasmic hydrogen peroxide production in root cells after treatment with C_{12} and C_{14} surfactin homologues as representative of short and long fatty acid chains respectively. Flg22 as used as control. Significate difference between each condition is indicated by different letters, p-value < 0.0001. **g** Binding coefficient (K) of Surfactin homologues (C_{12} to C_{15}) to large unilamellar vesicles (LUV) composed by PLPC/Sitosterol/Glucosylceramide (60:20:20 molar ratio). Means ± std err. from three to five biological replicates of one representative experiment are shown Significate difference between each condition is indicated by different letters, p-value < 0.05 **h** Release of 8-hydroxypyrene-1,3,6 trisulfonic acid (HPTS) from PLPC/Sitosterol/Glucosylceramide (60:20:20 molar ratio) LUV, upon addition of surfactin C_{12} or C_{14} at different concentrations. The ordinate shows the amount of HPTS released after 15 min in the presence of the C_{12} or C_{14} as a percentage of the total amount released by Triton X-100. **i** Influence of roots pretreatment with 10µM of surfactin (blue boxes) compared to non-treated roots (red boxes) on *B. velezensis* GA1 root colonization. Means ± std err. from five biological replicates of one representative experiment are shown. Ns= non significate.