Surfactin stimulated by pectin molecular patterns and root exudates acts as a key driver of *Bacillus*-plant mutualistic interaction

Grégory Hoff[1,2*], Anthony Arguelles-Arias[1], Farah Boubsi[1], Jelena Prsic[1], Thibault Meyer[1,3], Heba M. M. Ibrahim[4], Sebastien Steels[1], Patricio Luzuriaga[1], Aurélien Legras[1], Laurent Franzil[1], Michelle Lequart[5], Catherine Rayon[5], Victoria Osorio[6], Edwin de Pauw[6], Yannick Lara[7], Estelle Deboever[8], Barbara de Coninck[4], Philippe Jacques[1], Magali Deleu[8], Emmanuel Petit[5], Olivier Van Wuytswinkel[5], Marc Ongena[1*]

[1] Microbial Processes and Interactions, TERRA Teaching and Research Center, BioEcoAgro, Joint Research Unit/UMR transfrontalière 1158, University of Liège - Gembloux Agro-Bio Tech, Gembloux, Belgium

[2] Ecology and Biodiversity, Department of Biology, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands

[3] UMR Ecologie Microbienne, F-69622, University of Lyon, Université Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup, Villeurbanne, France

[4] Division of Plant Biotechnics, Department of Biosystems, Faculty of Bioscience Engineering, KU Leuven, Leuven, Belgium

[5] Unité Biologie des Plantes et Innovation, BioEcoAgro, Joint Research Unit/UMR transfrontalière 1158, Université de Picardie Jules Verne, UFR des Sciences, Amiens, France

[6] Mass Spectrometry Laboratory, MolSys Research Unit, Department of Chemistry, University of Liège, Liège, Belgium

[7] Astrobiology, UR-ASTROBIOLOGY, Geology Department, University of Liège, Liège, Belgium

[8] Molecular Biophysics at Interfaces Laboratory, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium

*   Grégory Hoff g.hoff@uu.nl
    Marc Ongena marc.ongena@uliege.be

Abstract

Bacillus velezensis is considered as model species belonging to the so-called *B. subtilis* complex that typically evolved to dwell in the soil rhizosphere niche and establish intimate association with plant roots. This bacterium provides protection to its natural host against diseases and represents one of the most promising biocontrol agents. However, the molecular basis of the cross-talk that this bacterium establishes with its natural host has been poorly investigated. We show here that these plant-associated bacteria have evolved some polymer-sensing system to perceive their host and that in response, they increase the production of the surfactin-type lipopeptide. Furthermore, we demonstrate that surfactin synthesis is favoured upon growth on root exudates and that this lipopeptide is a key component used by the bacterium to optimize biofilm formation, motility and early root colonization. In this specific nutritional context, the bacterium also modulates qualitatively the pattern of surfactin homologues co-produced *in planta* and mainly forms variants that are the most active at triggering plant immunity. Surfactin represents a shared good as it reinforces the defensive capacity of the host.

Importance

Within the plant-associated microbiome, some bacterial species are of particular interest due to the disease protective effect they provide via direct pathogen suppression and/or stimulation of host immunity. While these biocontrol mechanisms are quite well characterized, we still poorly understand the molecular basis of the cross talk these beneficial bacteria initiate with their host. Here we show that the model species *Bacillus velezensis* stimulates production of the surfactin lipopeptide upon sensing pectin as cell surface molecular pattern and upon feeding on root exudates. Surfactin favors bacterial rhizosphere fitness on one hand and primes the plant immune system on the other hand. Our data therefore illustrate how both partners use this multifunctional compound as unique shared good to sustain mutualistic interaction.

Keywords:

Lipopeptides, plant-associated bacteria*,* cell wall polymers, plant immunity, molecular crosstalk

Introduction

Soil is among the richest ecosystems in terms of microbial diversity, but only a subset of these microbes has evolved to efficiently establish in the competitive and nutrient-enriched rhizosphere layer surrounding plant roots (1). The rhizosphere includes plant beneficial bacteria dwelling on the rhizoplane as multicellular biofilm communities, feeding on exuded carbohydrates (2, 3), and, in turn, contributing to host fitness via growth stimulation and protection against phytopathogens (4, 5). This biocontrol activity is mediated via competition for nutrients and space, direct growth inhibition of the pathogenic (micro)organisms and more indirectly, by stimulating the host defensive capacity in an immunization-like process which leads to induced systemic resistance (ISR, (6, 7)). This ISR mechanism results in enhanced defense lines and reduced disease symptoms upon perception of plant beneficial microbes (6, 8).

From an ecological viewpoint, rhizosphere establishment and persistence of these beneficial bacteria rely on various traits but efficient root colonization and high competitiveness toward the surrounding microbiological network are pivotal. It is hypothesized that the potential to produce a wide range of chemically diverse and bioactive secondary metabolites (BSMs) acting as signals and/or antimicrobials is a common key feature of these beneficial bacteria (5, 9, 10). Members of the *Bacillus velezensis* species are considered as archetypes of plant-associated beneficial bacilli and are among the most prolific BSMs producers with more than 12% of their genome devoted to the synthesis of compounds contributing to both ecological competence and biocontrol activity (11–15). Among their BSM arsenal, the cyclic lipopeptide surfactin, is synthesized non-ribosomally by a multi-modular mega-enzyme machinery (encoded by the *srfA* operon) and is formed as a mix of naturally co-produced homologues varying in the length of the fatty acid chain. This multifunctional compound is of particular interest because it retains important roles in key developmental processes such as bacterial motility, biofilm formation and root colonization (16–18), but also because it represents the best described *Bacillus* triggers for plant immunity (6, 8). The potential of surfactin to stimulate ISR has been demonstrated on various plants including Solanaceae like tobacco and tomato on which it acts as main if not sole elicitor formed by *B. subtilis* and *B. velezensis* species (10, 19). In support to its key role in interaction with the host plant, we also previously reported that surfactin is promptly formed in the course of early colonization and that its production is stimulated upon sensing root tissues (20).

However, in contrast to the well-studied interactions between plants and microbial
pathogens or nitrogen-fixing bacteria (21), relatively little is known on the molecular basis
of cooperative interactions between plants and beneficial bacteria such as *B. velezensis*
(11, 20, 22). More specifically, how and to what extent the expression of key bacterial
BSMs may be modulated by plant factors is poorly understood. A better knowledge is not
only critical for providing new insights in rhizosphere chemical ecology but also for
optimizing the use of these species as biocontrol agents, which still suffer from insufficient
efficacy in practice (23). Here, we investigated the molecular interaction driving the early
steps of partnership establishment between plant roots and *B. velezensis*. We show that
cell wall pectin acts in synergy with soluble root exudates as plant host cues perceived by
B. velezensis. In response, the bacterium stimulates the production of specific surfactin
variants as key components of its secretome to further improve the fitness of both partners
i.e. early root colonization and thus rhizosphere competence of the bacterium and priming
of immunity in the host plant.
106
107

Results

Pectin fragments of high polymerization degree act as host cues triggering surfactin production

We previously described that early production of surfactin, as a mix of naturally co-produced homologues varying in the length of the fatty acid chain, is stimulated in contact with root tissues and several plant cell wall-associated polymers (PCWP) (20). In this work, we further investigated this phenomenon focusing on the impact of pectin, as it represents complex sugar polymers typically found in the plant primary cell wall and particularly abundant in the middle lamella layer (24). We first tested the effect of crude pectin extracted from tobacco root PCWP (referred as cPec, Fig. 1ab for composition and related structure). An 8-fold increase of surfactin production was detected at the early exponential growth phase (OD_{600}=0.2-0.25) in *B. velezensis* GA1 liquid cultures supplemented with cPec compared to an un-supplemented culture (Fig. 1cd). Surfactin production was also 10 times enhanced upon addition at the same concentration of pure commercially available homogalacturonan (HG) with high degree of polymerization (DP) (Fig. S1ab) but low level of methyl-esterification (HGLM) according to the manufacturer (Fig. 1d). HG was tested as the most abundant pectic polysaccharide constituent, which represents 65% of crude primary cell wall pectin (24). Production of this lipopeptide was also enhanced to a similar level upon addition of highly methylated HG (HGHM), showing that the degree of methyl-esterification of the polymer is not a major trait influencing perception by the bacterium (Fig. S2). Altogether, this supports a key role of the pectin backbone as plant molecular pattern that is sensed by the bacterium to stimulate surfactin synthesis.

Interestingly, by screening the CAZy database (25) for genes encoding carbohydrate-active enzymes potentially involved in PCWP degradation by *B. velezensis*, two putative pectate/pectin lyases encoding genes were detected. These two genes, referred as *pelA* and *pelB* (accessions *GL331_08735* and *GL331_04125 in B. velezensis* GA1*, respectively)*, are highly conserved among all sequenced *Bacillus* genomes that belong to the "Operational Group *B. amyloliquefaciens*" (Table S1) (26). *pelA* and *pelB* are readily expressed in GA1 *in vitro* and the corresponding enzymes efficiently convert HG into unsaturated oligogalacturonides with consistent activity occurring at the beginning of stationary phase (Fig. S2). However, the bacterial perception of oligomers with lower

polymerization degree compared to HG is not obvious since oligogalacturonides (OG) did
not stimulate surfactin biosynthesis (Fig. 1d, Fig. S1c for OG characterization).
Supplementation with galacturonic acid (GA) led to a reduction of surfactin production at
mid exponential phase (OD_{600}=0.35, Fig. 1d). Surfactin production is thus specifically
boosted upon sensing long degree of polymerization (DP) polymers, but is somehow
inhibited in presence of GA constituting the pectin backbone. Such HG-driven surfactin
stimulation also occurs in other *B. velezensis* isolates tested (FZB42, QST713 and S499)
and to a lower extent *B. pumilus* QST 2808. It does not occur in the non-rhizosphere
dwelling isolates *B. amyloliquefaciens* DSM7 or *B. subtilis* ATCC 21332 (Fig. 1e)
suggesting that this trait may be specific to bacilli with a plant-associated lifestyle.

The root nutritional context favors early surfactin production

Bacillus velezensis quickly colonize tomato plantlets in a gnotobiotic system and
forms visible biofilm-like structures covering the main root and embedding lateral roots
after 24-48h post inoculation (Fig. 2a). This is correlated with consistent *srfAA* gene
expression and surfactin production rate in the cell population at these early times but it
was maintained albeit to a lower level, over the investigated timeframe of seven days (Fig.
2ab). Since surfactin enhancement linked to the perception of the pectin backbone is only
transient (Fig. 1d), we hypothesized that root exudates, constantly secreted by the plant,
may also positively impact the synthesis of the lipopeptide. Surfactin production rate was
thus compared upon growth in a classical laboratory medium (LB) and in a root exudate-
mimicking medium (REM) reflecting the content of carbohydrates typically released by
tomato or tobacco roots (27). It revealed an earlier and higher production by cells growing
in REM (Fig. 2c). Surfactin production in REM medium is initiated earlier and is more
efficient in *B. velezensis* compared to other closely related but non plant-associated
species such as *B. amyloliquefaciens* or *B. subtilis* (Fig. 2d).

Addition of HG in REM medium compared to LB revealed a cumulative effect of
this PCWP and root exudates on surfactin production (Fig 3a). This could be of clear
ecological benefit for the bacterium since surfactin is known to favor motility of multicellular
communities and biofilm formation (16, 28, 29). However, a recent study questioned the
real role of surfactin in these key functions, since it production appears as non-essential
for pellicle biofilm formation in *B. subtilis* NCIB 3610, suggesting a strain dependant role
(30). We previously reported that motility and biofilm formation are boosted upon growth
on root exudates (27). Here we show that HG supplementation also favors *B. velezensis*

GA1 spreading on low-agar medium (Fig 3b) and early biofilm formation based on pellicle

development at the air-liquid interface (31) (Fig 3c). The role of surfactin in swarming,

pellicle formation and early root colonization was further confirmed for *B. velezensis* GA1.

Indeed, swarming motility on low agar plates was almost reduced to zero in a surfactin

deficient mutant, and the same mutant was more than 3 times less efficient to produce

pellicles at the air liquid interface and to promptly colonize tomato roots after 1 day post

inoculation when compared to the WT (Fig 3def). Collectively, these data allow correlating

the positive impact of PCWP on bacterial motility, biofilm formation and early root

colonization through an anticipated surfactin production in *B. velezensis*.

183   Surfactin induction by PCWP is not linked to major transcriptional changes.

Both HG and root exudates stimulate surfactin production in GA1. However, while

no activation of the *srfA* biosynthetic gene cluster was observed upon HG addition (Fig.

4a), an early and high surfactin gene expression was measured in *srfAp*::*gfp* cells growing

in REM compared to LB medium (Fig. 4b). To unravel transcriptome wide changes in GA1

associated with the perception of HG, RNA-sequencing was performed on cells grown in

REM with our without addition of HG and collected at various time points (lag, early

exponential and a mid-exponential phases). The data confirmed that HG perception is not

linked to an increased expression of the *srfA* operon but also revealed a quite limited and

transient transcriptional reprogramming with only 58 genes differentially expressed over

this timeframe (Table 1). Remarkably, more than 30% of these genes are involved in

stress response or cell wall modifications and are down regulated in the presence of HG

(Fig. 4c). We thus hypothesize that a long-term co-evolution process may have facilitated

Bacillus establishment on the roots by the inhibition of a costly stress response after

perception of HG. Addition of HG also leads to a 4.2-fold reduced expression of *flgM*

encoding an inhibitor of SigD, the σ factor involved in the activation of motility related

genes (32). This may contribute to enhanced spreading of multicellular communities in

addition to the positive effect of surfactin mentioned above.

201   Root exudates drive the bacterium to form surfactin homologues with long fatty

202   acid chain (LFAC) and variants enriched in valine

The NRPS machinery works as an assembly line in which each module is

responsible for recruiting and binding a specific amino acid to the nascent peptide after a

first lipo-initiation step for binding the fatty acid (FA) taken up from the cellular pool (Fig.

1A) (33, 34). In that way, surfactin is typically composed of saturated C_{12} to C_{19}-FA of the

linear, iso or anteiso type of branching (35). Beside an increased production of surfactin,

we also observed an effect on the pattern of surfactin variants synthesized by *B.*

velezensis in the presence of artificial plant exudates, as well as in naturally produced

exudates and in planta upon root colonization (Fig. S4). Indeed, UPLC-MS profiling

revealed that the surfactin pattern produced by GA1 in REM medium is enriched in

surfactin *iso*-C_{14} (iC_{14}) and other variants compared to LB medium (Fig. 5b). They

correspond to variants of the canonical structure with substitution of Leu by Val for the last

residue of the cyclic peptide moiety (Val_7) and, to a much lower extent, to the same

substitution in position 2 (Val_2, Fig. 5c, see Fig. S5). Valine is used both as precursor for

the synthesis of branched fatty acids with an even number of carbons, and as a building

block by the NRPS to form the peptide moiety. Supplementation of the medium with

deuterated L-Val-d^8 resulted in an additional increase in the proportions of surfactin *iso*-

C14 and Val_7 isoforms labeled at the expected positions in the peptide and in the fatty acid

tail (Fig. S6). Based on these data, the higher relative proportions of $iC_{14}Val^7$ formed in

REM, but also *in planta* (Fig. 5c), most probably result from some enrichment of the

intracellular pool in valine upon growth in the presence of root exudates (Supplementary

Discussion). Given the reduced specificity of NRPS domains involved in selection and

activation of leucine at positions 2 and 7, the megaenzyme would preferably bind valine

as it is more available in the pool.

As already described in *B. subtilis* (36, 37), the pleiotropic regulator CodY acts as

repressor of surfactin synthesis in *B. velezensis* GA1 as illustrated by the 1.9-fold increase

in production by the Δ*codY* mutant of strain GA1. Interestingly, CodY activity/*codY*

expression is also itself negatively impacted by high cellular concentrations in branched

chain amino acids (38). Both quantitative and qualitative changes in surfactin production

upon growth in exudates could therefore be, at least partly, due to a lower CodY activity

(Supplementary Discussion). In support to the role played by this regulator, a similar

impact on surfactin pattern were observed by deleting *codY* in GA1 or by supplementing

the culture medium of the wild-type with valine (Fig. 5d).

Long fatty acid chain surfactins act as key triggers of receptor-independent plant
immunity

Based on the potential of surfactin as host immunity elicitor (9, 39), we next wanted

to evaluate the possible relevance of quantitative and qualitative modulation of the

surfactin pattern driven by the plant for its own benefit.

Upon application as root treatment, pure surfactin used as mixture of isoforms

formed in REM medium, induced systemic resistance in hydroponically-grown tobacco

plants providing approximatively 45-50% significant disease reduction on leaves

subsequently infected with the pathogen *Botrytis cinerea* (Fig. 6a). The various isoforms

were then HPLC-purified and tested individually revealing that only long fatty acid

homologues (C_{14}/C_{15}) provided systemic protection to a similar level whereas short fatty

acid homologues (C_{12}/C_{13}) were inactive (Fig. 6b). Moreover, plant immunization by

surfactin is dose-dependent and concentrations up to 5 μM are sufficient to significantly

stimulate ISR (Fig. 6c). Interestingly, such low μM concentration are actually in the range

of those that could accumulate in the root vicinity within a few days upon colonization by

GA1 (Fig. S7).

We next wanted to correlate this systemic protection induced by the lipopeptide

with its potential to trigger locally early immune-related events such as the extracellular

burst in reactive oxygen species (ROS) involved in defense and signaling in pathogen-

triggered immunity (PTI) (40, 41). By contrast with flagellin (epitope Flg22), one of the best

characterized Microbe-Associated Molecular Patterns (MAMPs) isolated from bacterial

pathogens, treatment with surfactin did not induced burst in apoplastic ROS in root tissues

(Fig. 6d). However, surfactin-mediated ROS signaling still occurs since a clear cytoplasmic

ROS accumulation was observed (Fig 6e). Little information is available about the spatio-

temporal dynamics of such ROS burst but it may originate from different organelles and

has been occasionally described in response to perception of biotic and abiotic stresses

(42, 43). Using cytoplasmic ROS as marker, the same trend as for ISR tests could be

observed regarding the influence of the structure on the activity of surfactin since long fatty

acid homologues but not short ones efficiently stimulated early immune reaction (Fig 6f).

This means that a single additional methylene group in the fatty acid tail of the molecule

(C_{14} versus C_{13}) likely determines its immunization potential (Fig 6b,f). By contrast,

substitution of Leu[7] by a Val in the C_{14} homologue does not impact activity suggesting that

the peptide moiety is not essential for perception by plant cells. In addition, the μM

concentrations required for optimal eliciting activity of surfactin are very high compared

with PAMPs active in the nM range (44). Our previous data showed that surfactin elicitation

is still active after treating cells with proteases or after a first application indicating that

there is no saturation of high-affinity but low abundance binding sites on receptors (45,

46). All this indicates that surfactin is perceived by plant cells via a mechanism

independent of high-affinity pattern-recognition receptors (PRRs) involved in MAMP

perception (40, 41, 44, 47, 48). We therefore postulated that surfactin perception relies on

some interaction with the lipid phase of the plant plasma membrane. Binding experiments

via isothermal titration calorimetry and leakage assays based on the release of fluorescent

probe were performed using liposomes prepared with lipids specific to plant plasma

membrane (PLPC/sitosterol/glucosylCeramide). It revealed that long fatty acid

homologues have a higher affinity for these vesicles than the short fatty acid forms and

display a higher destabilizing effect on the lipid bilayer when added at concentrations of 5

µM or higher (Fig. 6gh). These biophysical data thus correlated well with the contrasting

biological activities of longer C_{14}/C_{15} and shorter C_{12}/C_{13} surfactin homologues.

According to the priming concept (49), we previously showed that ISR triggered by

the lipopeptide in that plant as well as in tobacco and *Arabidopsis*, is not associated with

a fast and strong expression of defensive mechanisms before pathogen infection (20, 39).

In order to verify that surfactin elicitation does not cause a massive release of

antimicrobials from plant tissues, tomato roots were pre-treated with the lipopeptide before

inoculation with *B. velezensis*. As expected, it did not impacted subsequent colonization

in terms of rate and dynamics compared to untreated plants indicating the absence of

potential adverse effects on the bacterial partner (Fig. 6i).

291
292

Discussion

A large part of the interactions between bacteria and plants is known to be mediated by small-size secreted products (50). However, a better understanding of the chemical cross-talk at the plant-bacteria interface and its impact on bacterial ecology, plant fitness and immune responses remains challenging. In epiphytic soil bacilli, root exudates induce expression of an array of genes involved in various functions such as chemotaxis and nutrient acquisition (51–53). Our data further illustrate that utilization of this cocktail of molecules released by roots but also the perception of some cell wall polymers may also drive these bacteria to efficiently produce key components of the secondary metabolome and more specifically the multifunctional surfactin lipopeptide (20). As an amphiphilic molecule and powerful biosurfactant, surfactin is presumably viewed as membrane active compound with potent antimicrobial activity. However, this lipopeptide is poorly antibacterial and antifungal (54). In *B. velezensis*, more obvious ecological functions of this CLP are to contribute to motility, biofilm formation and roots colonization. An enhanced production upon host perception thus constitutes a major force driving successful rhizosphere establishment.

Homogalacturonan acts as a cue to enhance surfactin secretion by bacterial cells but no transcriptional induction of the corresponding biosynthesis operon was observed. Surfactin synthesis is integrated in a complex network involving several pleiotropic regulators acting directly or indirectly on the expression of the *srfA* operon (55–58). However, we hypothesize that surfactin induction by HG may rather rely on post-transcriptional changes as reported for the effect of the DegU and YczE regulators on production of another CLP, bacillomycin D (59). Despite the relatively close genetic proximity of the two species, our data suggest that regulation of surfactin could be slightly different in *B. velezensis* and *B. subtilis*. As it represents a key infochemical devoted to cross-talk with the host plant, surfactin regulation may have been fine-tuned in rhizosphere species to better fit with the nutritional or more broadly ecological context.

Deciphering the mechanism by which *B. velezensis* recognizes pectin and enhances surfactin production would help to identify candidate genes and pathways that are responsible for plant sensing, ensuring persistence on roots which globally remains very poorly known for beneficial rhizobacteria. We are currently investigating whether some cell surface proteins may act as receptors for homogalacturonan perception and

binding as recently described for *Sphingomonas* sp. (60), another beneficial species living
in association with plants (61). Some insights could be obtained by scrutinizing the few
genes conserved in *B. velezensis* but missing in non-plant-associated *B.*
amyloliquefaciens strains that are not responsive to pectin (62). Interestingly, shorter
fragments of HG and galacturonic acid do not stimulate surfactin secretion. It is therefore
tempting to hypothesize that sensing unaltered polymer could indicate a healthy host
suitable for bacterial colonization while the perception of monomers or low DP oligomers
may reflect a dead or infected plant that is unable to adequately provide resources.

Our data illustrate for the first time that *B. velezensis* can also modulate
qualitatively its surfactin pattern by growing in its natural nutritional context, *i.e.* on root
exudates. Substitution of leucine by valine in the peptide part is not expected to impact
the contribution of the lipopeptide to colonization by the producing strain itself considering
the minor effect of these structural changes on motility and biofilm formation potential (18).
Small modifications in the peptide sequence may nevertheless avoid surfactin hijacking
for use as signal prompting heterologous biofilm formation by closely related competitor
species (18). Based on our observations, the most obvious benefit of an increased
proportion of long fatty acid chain homologues is for the host plant since they represent
the most active forms for priming immunity with no impact on host fitness (20, 39), by
contrast with PTI (63, 64). As the bacterial partner does not have to face strong defensive
responses from this reaction, it ensures positive mutualistic co-habitation allowing
establishment of populations on roots. Persistence of threshold populations is necessary
for consistent production of other specialized secondary metabolites more directly
involved in warding off both microbial competitors and plant soilborne pathogens in the
context of biocontrol.

Surfactin stimulation upon sensing host molecular patterns may thus reflect an
aspect of plant-*Bacillus* coevolution as it makes a shared good out of this multifunctional
lipopeptide. To some extent, it might represent a facet of the plant-driven selection process
resulting in active recruitment of this bacterium as species that provides beneficial
functions. Other bacterial genera such as *Pseudomonas* also prevailing in the rhizosphere
microbiome actively produce CLPs with similar roles as surfactin. Evaluating whether their
synthesis is also modulated by plant cues would conceptually allow broadening the
significance of these lipopeptide-mediated inter-kingdom interactions for bacterial
ecology, plant health and biocontrol.

Materials and Methods

Bacterial media and growth conditions. Cultures were performed at 26°C in root exudates mimicking medium (EM) (27) or in LB medium. To test the effect of plant cell wall polymers, each specific plant polysaccharide was added at a final concentration of 0.1% in the culture medium. Low (HGLM, <5%) and high (HGHM, >95%) methylated homogalacturonan were provided from Elicityl Oligotech whereas oligogalacturonides and D-galacturonic acid were provided from Sigma.

Strains construction. All the bacteria strains used in this study are listed in table 2. All the primers used in this study are available upon request. To follow the expression level of the *srf* operon in GA1, we constructed a *gfp* transcriptional fusion under the control of the *srf* promoter and integrated it into the *amyE* locus. First, a GA1 *amyE* amplicon containing a native *KasI* restriction site was integrated in the PGEMT easy. In parallel, a *cat-gfp* cassette containing respectively (i) a chloramphenicol resistance gene (cat) and (ii) a promoterless *gfpmut3.1* gene was amplified with primers containing *KasI* sites at their 5' extremities using the pGFP star as a matrix (65). The pGEMT *amyE* plasmid and the *cat-gfp* amplicon were both digested by *KasI* (NEB) and the two linear fragments with compatible 5' overhangs were ligated together to obtain the PGEMT *amyEup-cat-gfp-amyEdw* plasmid. To construct the final mutation cassette, an overlap extension PCR was assessed by following the method developed by Bryksin and Matsumura (66). One first fragment containing the upper *amyE* homologous region and the *cat* gene, and one second fragment englobing the *gfpmut3.1* gene and the lower *amyE* homologous region were both amplified using the PGEMT *amyEup-cat-gfp-amyEdw* plasmid as a matrix. A third fragment was amplified using GA1 genome as matrix with chimeric primers designed to obtain a *srf* promoter amplicon flanked by 20 bp connectors in 5' and 3' containing respectively homologies to the upper and lower *amyE* fragments. All three fragments were joined together with a second PCR race to obtain the final cassette. *B. velezensis* GA1 transformation was performed after modification from the protocol developed by Jarmer *et al.* (67). Briefly, one colony was inoculated into LB liquid medium at 37°C (160 rpm) during 6h and cells were washed two times with peptone water. Until 1µg of the recombinant cassette was added to the GA1 cells suspension adjusted to an OD_{600nm} of 0.01 into MMG liquid medium (19 g l-1 K_2HPO_4 anhydrous; 6 g l-1 KH_2PO_4; 1 g l-1 Na_3 citrate anhydrous; 0.2 g l-1 $MgSO_4$ $7H_2O$; 2 g l-1 Na_2SO4; 50 µM $FeCl_3$ (sterilized by filtration at 0.22 µm); 2µM $MnSO_4$; 8 g l-1 Glucose; 2 g l-1 L-glutamic acid; pH 7.0). Cells were incubated at 37°C with shaking, and colonies who integrated the cassette by a double crossing over event were selected on LB plate supplemented with chloramphenicol. Proper integration of the *cat-gfp* locus was verified by PCR. Knock-out mutant strains were constructed by gene replacement by homologous recombination. A cassette containing a chloramphenicol resistance gene flanked respectively by 1 kb of the upstream region and 1 kb of the downstream region of the targeted gene was constructed by a three partners overlap PCR. This recombination cassette was also introduced in *B. velezensis* GA1 by inducing

natural competence as described above (67). Double homologous recombination event was
selected by chloramphenicol resistance. Deletion was confirmed by PCR analysis with the
corresponding upstream and downstream primers.

Fluorescence measurement. Fluorescence accumulation was evaluated thanks to the channel
FL1 of a BD accuri C6 flow cytometer (Biosciences) with the following parameters: 20000 events,
medium flow rate (35 µl.min^{-1}), FSC threshold of 20000.

Genome sequencing. GA1 genome sequence was reconstructed using a combined approach of
two sequencing technologies which generated short paired end reads and long reads. The resulted
sequences were then used for hybrid assembly. More precisely, genomic DNA was extracted and
purified from *B. velezensis* GA1 using the GeneJET Genomic DNA purification (ThermoFisher
scientific). First half of extracted DNA was sent to the GIGA sequencing facility (Liège, Belgium),
and use as DNA template for illumina MiSeq sequencing after being prepared using nextera library
kit illumina. Sequencing run generated 150 bp paired-end read, which were trimmed and corrected
using an in-house python script and SPAdes 3.14 (68) before assembly. The second half of
extracted DNA was used to generate long reads with a MinION Oxford nanopore plateform. DNA
library was constructed using the Rapid Sequencing kit (SQK-RAD0004, Oxford nanopore).
Adapters were trimmed from generated reads with Porechop software
(https://github.com/rrwick/Porechop). Trimmed reads were then filtered by size (>500) and Q-score
(>10) using NanoFilt implemented in NanoPack (69). Finally, hybrid assembly was performed using
hybridSPAdes algorithm implemented in SPAdes 3.14 (70).

Transcriptome library preparation and sequencing. RNA extraction was performed for each
sample using the NucleoSpin RNA kit (Macherey-Nagel). Total RNAs were quantified using
Nanodrop (Thermofisher). For sequencing, all samples were sent to the GIGA-genomics platform
in Liège. Quality was assessed using the RNA 6000 Nano Chip on a 2100 Bioanalyzer (Agilent).
cDNA libraries were prepared employing Universal Prokaryotic RNA-Seq, Prokaryotic AnyDeplete
kit (Nugen) according to the manufacturer's instructions, with . cDNA libraries were quantified and
normalized by using the KAPA SYBR Fast Mastermix (Sigma-Aldrich) with P5-P7 Illumina primers
according to the manufacturer's instructions. Prepared libraries were sequenced on a NextSeq 550
device (Illumina) by using the following parameters : paired end, 80 cycles read 1, 8 cycles index,
80 cycles read 2.

RNA-seq data analysis. The raw RNA-seq reads were trimmed using Trimmomatic v0.39 (71).
We performed quality control on the trimmed reads using FastQC v0.11.8 (Babraham
Bioinformatics). Trimmed reads were mapped to the GA1 reference genome (see section "genome
sequencing" for accession numbers) using BWA-mem v0.7.17 (72) with the following settings: mem
-k 50 -B 40 -v 1. At least 95.4% of reads uniquely mapped to the annotated reference genome
(Table S2). SAMtools v1.9 (73) was used to generate the BAM files and their indices. To calculate

the read counts, the python-based tool HTSeq v0.9 (74) was employed with the following
parameters: htseq-count -q -s no -f. The Cufflinks function cuffnorm (75) was used to generate the
FPKM (fragments per kilobase of transcript per million mapped reads) tables using the following
settings: --compatible-hits-norm --library-norm-method classic-fpkm. Genes with low reads counts
(<25) were removed before further analysis. Differential expression analysis was conducted
according to the DESeq2 pipeline (10.1186/s13059-014-0550-8) with cut-off parameters of $p<0.05$
and log2-fold-change > 1.5.

Motility and biofilm assays. Swarming motility assays were performed according to Molinatto *et*
al. 2017 (76). Diameter of the bacterial swarming pattern was measured 48h after inoculation on
REM soft agar plates (0.8% agar) supplemented or not with 0.1% HGLM. Quantification of total
biofilm was performed by crystal violet staining. Strain of interest was inoculated at a final OD_{600} of
0.1 in a 96 wells microplate containing 200 µl of REM medium supplemented or not with 0.1 %
HGLM. The plate was incubated at 30°C during 24h without shaking. Medium and planctonic cells
were discarded and wells were washed with PBS. Biofilm pellicle was stained with 0.1% crystal
violet during 10 min and washed with PBS. The stained biofilm was dissolved with 30% acetic acid.
Absorbance was measured at 595 nm.

Plant growth conditions and roots colonization assays. For sterilization, tomato seeds were
first immersed in a 70% ethanol solution during 2 minutes, transferred in a 20% bleach solution
under shaking for 20 minutes and rinsed three times with sterile water. Sterilized tomato seeds
were pre-germinated on solid Hoagland medium at 22°C under a 16h/8h night/day cycle. After 4
days, 5µL of cultures containing the strain of interest and calibrated at $OD_{600}=1$ were deposited on
the root top. After 1 and 3 days of colonization, roots were harvested, deposited separately in a
peptone water solution supplemented with 0.1% of Tween, and vortexed vigorously to tear off the
bacterial cells from the roots. Several dilutions were plated on LB media to evaluate the level of
colonization. Measurements of surfactin production by GA1 cells colonizing roots were performed
on 1x1x0.7 cm pieces of gelified medium containing roots based on the assumption that the
produced lipopeptide diffused to a maximal distance of 5 mm from each part of the root and is
uniformly distributed over the surface as we previously observed via imaging-MS (77). A 10-fold
concentration factor was applied to estimate concentrations around the root surface in order to take
into account diffusion constraints in a solid matrix. Surfactin was quantified by UPLC-MS as
described below.

Plant cell wall extraction. Tobacco seeds were sterilized as described above for tomato seeds
and deposited on Hoagland plates at 22°C during one week for a successful germination process.
Each plantlet was then transferred in a seedholder filled with soft agar and put in Araponics boxes
containing the nutritive solution described above. Cell wall extraction was performed on 6 weeks
old plants grown at 22°C with a 16h/8h day-night alternance. Roots were harvested, lyophilized

and reduced to powder using a Retsch MM400 grinder. 500 mg of powder was resuspended in 40

ml of ethanol 80% at 90°C for 20 min. The insoluble cell wall fraction was recovered by

centrifugation and the pellet obtained was washed once with water to obtain the Alcoholic Insoluble

Residue (AIR) used for fractionation. AIR was freeze-dried before use in fractionation protocol.

Sequential extraction of root cell walls was performed using a protocol derived from Carpita (78)

and Silva *et al.* (79). Dry AIR was resuspended in 40 ml water and incubated at 100°C for 20 min.

Supernatant was recovered after centrifugation as a soluble pectic fraction (cPEC).

Monosaccharide composition analysis using HPAEC-PAD. Before monosaccharide

composition analysis, cPec fraction was dialyzed during 24h against a large volume of water and

freeze-dried. 2 mg of dried fraction material was hydrolyzed in 1 ml of 2M Trifluoroacetic acid (TFA)

at 121°C for 90 min. TFA was evaporated under nitrogen gas flux and the hydrolysed dried residue

was resuspended in 1 ml water, filtered on 0,2 μm cartridge and stored in vials at 20° before

HPAEC-PAD. High Performance Anion Exchange Chromatography with Pulsed Amperometric

detection (HPAEC-PAD) was used for neutral and acidic monosaccharide composition analysis

using a Dionex DX-500 system (Dionex Corporation) equipped with a Carbopac PA-1 analytical

column (4 mm x 250 mm). The elution was performed with a flow rate of 1 mL.min^{-1} in a gradient

mode. The gradient for neutral sugars (eluent A: deionized water, eluent B: 160 mM NaOH and

eluent C: 200 mM NaOH) was 10% B for 25 min, 100% B for 10 min and finally an equilibration

step with 10% B (15 min). The gradient for uronic acid (eluent A: 160 mM NaOH and eluent B: 160

mM NaOH + 600 mM AcONa) was 0% B for 5 minutes, 30 minutes of linear gradient from 0 to

100% B, 100% B for 5 minutes and finally an equilibration step with 0% B (10 minutes). Detection

was performed with a pulsed amperometric ED50 detector (Dionex Corporation). 20 mL of sample

was injected with an autosampler. Each carbohydrate concentration was determined after

integration of the respective areas (Chromeleon management system, Dionex) and comparison

with standard curves.

LC-MS analyses. Detection of metabolites and quantification was performed by LC-MS. 10 µL of

samples were analyzed using UPLC–MS with UPLC (Acquity H-class, Waters) coupled to a single

quadrupole mass spectrometer (SQD mass analyzer, Waters) using an C18 column (Acquity UPLC

BEH C18 2.1 mm × 50 mm, 1.7 µm). Elution was performed at 40°C with a constant flow rate of

0.6 mL/min using a gradient of Acetonitrile (solvent B) and water (solvent A) both acidified with

0.1% formic acid as follows: starting at 15% B during 2 min, solvent B was then raised from 15%

to 95% in 5 min and maintained at 95% up to 9.5 min before going back to initial conditions at 9.8

min during 3 minutes before next injection if needed. Compounds were detected in electrospray

positive ion mode by setting SQD parameters as follows: source temperature 130°C; desolvation

temperature 400°C, and nitrogen flow: 1000 L.h-1 with mass range from m/z 800 to 1550.

Surfactins were quantified based on their retention times and masses compared to commercial

standards (98% purity, Lipofabrik).

Induction of systemic resistance and ROS measurements. ISR assays were performed as previously described (39) on 4 weeks-old tobacco plants cultivated in hydroponic conditions using Hoagland solution as nutrient base. Plants were treated with pure surfactin at the root level and infected on leaves by applying a spore suspension of the phytopathogen *Botrytis cinerea* prepared as detailed previously (39). Spreading lesions occurred starting from 48h post-infection and the diameter size was measured two days later. Five plants were used per treatment and experiments were repeated independently at least twice. For determination of cytoplasmic ROS stimulation, fluorescent probe (DCFH-DA) was used. Plants used in this experiment were grown on Hoagland medium for two weeks as described above. Experiments were performed on nine samples per treatment each containing three root segments (approx 100 mg FW) collected from different plants (n=9). Roots were treated with 50 µM DCFH-DA for 10 minutes, rinsed with PBS upon removing the probe, and finally treated. All the operations were conducted in a 96-well black microplate. Fluorescence measurements were performed on a Spark (Tecan) microplate reader (exc 485 nm; em = 535 nm) with readings every 10 minutes. Stimulation of apoplastic hydrogen peroxide production in root cells was measured via chemiluminescence (ferricyanide-catalysed oxidation of luminol). Means and standard deviations were calculated from measurements performed on three samples per treatment, each containing three root segments (approximatively 100 mg FW) collected from different plants. Extracellular ROS in tomato roots was conducted according to Bisceglia *et al.* (80) with minor changes. Namely, instead of leaf discs, tomato roots, three segments (approximatively 100 mg FW from the same plant) per sample, were used. Plants were grown for two weeks on Hoagland medium, and chemiluminescence was measured in Tecan Spark plate reader.

ITC analysis. ITC analyses were performed with a VP-ITC Microcalorimeter (Microcal). The calorimeter cell (volume of 1.4565 mL) was filled with a 10 µM (below the CMC concentration) surfactin solution in buffer (Tris 10mM, NaCl 150mM, 1mM EDTA at pH 8.5). The syringe was filled with a suspension of LUV at a lipid concentration of 5 mM. A series of 10µl injections was performed at constant time intervals (6 min) at 25°C. The solution in the titration cell was stirred at 305 RPM. Prior to each analysis, all solutions were degassed using a sonicator bath. The heats of dilution of vesicles were determined by injecting vesicles in buffer and subtracted from the heats determined in the experiments. Data were processed by software Origin 7 (Originlab) using the cumulative model described by Heerklotz and Seelig (81). All measurements were repeated at least three times with two different vesicle preparations.

Leakage assays. Membrane permeabilization was followed as described by Van Bambeke *et al.* (82). Release of 8-hydroxypyrene-1,3,6 trisulfonic acid (HTPS) coentrapped with and quenched by p-xylene-bis-pyridinium bromide (DPX) from liposomes can be monitored by the fluorescence increase upon dilution following their leakage from the vesicles. Surfactin C12 or Surfactin C14 was added from a stock solution in DMSO and fluorescence intensities were immediately recorded. The

percentage of HPTS released was defined as [(Ft- Fcontr)/(Ftot- Fcontr)] /100, where Ft is the fluorescence signal measured after 15 min in the presence of Surfactin C12 or Surfactin C14, Fcontr is the fluorescence signal measured at the same time for control liposomes, and Ftot is the total fluorescence signal obtained after complete disruption of the liposomes by 0.05% Triton X-100. All fluorescence determinations were performed at room temperature on a Perkin Elmer LS-50B Fluorescence Spectrophotometer (Perkin-Elmer Ltd.) using λexc of 450 nm and a λem of 512 nm.

Statistical analyses. All statistical analyses were performed on GraphPad prism. Before each statistical analysis, variance homoscedasticity was verified by using a Brown-Forsythe test. ANOVA analysis was used for multiple comparison and significant differences were indicate by different letters. Statistical differences between means were evaluated by two-tailed Student's t-test. Number or biological replicates used for each experiment are indicated in the corresponding figure legend. P-Values are indicated in the figure legends.

Data availability The RNA-seq datasets produced for this study are deposited at https://www.ebi.ac.uk/ena/ under the project reference PRJEB39762. All other datasets analyzed for this study are included in the supplementary files. The Genome Resulting assembly of the GA1 strain was deposited in the GenBank database under the accession numbers CP046386 and CP046387.

562
Acknowledgments

564
This work was supported by the EU Interreg V France-Wallonie-Vlaanderen portfolio
SmartBiocontrol (Bioprotect and Bioscreen projects, avec le soutien du Fonds européen
de développement régional - Met steun van het Europees Fonds voor Regionale
Ontwikkeling), by the PDR research project ID 26084552 from the F.R.S.-FNRS (National
fund for Scientific Research in Belgium) and by the EOS project ID 30650620 from the
FWO/F.R.S.-FNRS. FB is recipient of a F.R.I.A. fellowship (Formation à la Recherche
dans l'Industrie et l'Agriculture) and MO is senior research associate at the F.R.S.-
F.N.R.S. We are grateful to the KU Leuven HPC infrastructure and the Flemish
Supercomputer Center (VSC) for providing the computational resources and services to
perform the RNA-seq analysis. We gratefully acknowledge Claire Bertrand and Loïc
Ongena for critically reading the manuscript.

References

1.   Andrews JH, Harris RF. 2000. The Ecology and Biogeography of Micoroorganisms on Plant Surfaces. Annu Rev Phytopathol 38:145–80.

2.   Zhalnina K, Louie KB, Hao Z, Mansoori N, Da Rocha UN, Shi S, Cho H, Karaoz U, Loqué D, Bowen BP, Firestone MK, Northen TR, Brodie EL. 2018. Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. Nat Microbiol 3:470–480.

3.   Vieira S, Sikorski J, Dietz S, Herz K, Schrumpf M, Bruelheide H, Scheel D, Friedrich MW, Overmann J. 2020. Drivers of the composition of active rhizosphere bacterial communities in temperate grasslands. ISME J 14:463–475.

4.   Vacheron J, Desbrosses G, Bouffaud M-L, Touraine B, Moënne-Loccoz Y, Muller D, Legendre L, Wisniewski-Dyé F, Prigent-Combaret C. 2013. Plant growth-promoting rhizobacteria and root system functioning. Front Plant Sci 4:1–19.

5.   Backer R, Rokem JS, Ilangumaran G, Lamont J, Praslickova D, Ricci E, Subramanian S, Smith DL. 2018. Plant growth-promoting rhizobacteria: Context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. Front Plant Sci. Frontiers Media S.A.

6.   Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, Van Wees SCM, Bakker PAHM. 2014. Induced Systemic Resistance by Beneficial Microbes. Annu Rev Phytopathol 52:347–375.

7.   Köhl J, Kolnaar R, Ravensberg WJ. 2019. Mode of action of microbial biological control agents against plant diseases: Relevance beyond efficacy. Front Plant Sci 10:1–19.

8.   van Loon LC, Bakker PAHM, Pieterse CMJ. 1998. Systemic Resistance Induced By Rhizosphere Bacteria. Annu Rev Phytopathol 36:453–483.

9.   Ongena M, Jourdan E, Adam A, Paquot M, Brans A, Joris B, Arpigny JL, Thonart P. 2007. Surfactin and fengycin lipopeptides of Bacillus subtilis as elicitors of induced systemic resistance in plants. Environ Microbiol 9:1084–1090.

10.   Ongena M, Jacques P. 2008. Bacillus lipopeptides: versatile weapons for plant disease biocontrol. Trends Microbiol 16:115–125.

11.   Wu K, Fang Z, Guo R, Pan B, Shi W, Yuan S, Guan H, Gong M, Shen B, Shen Q. 2015. Pectin enhances bio-control efficacy by inducing colonization and secretion of secondary metabolites by Bacillus amyloliquefaciens SQY 162 in the rhizosphere of tobacco. PLoS One 10:1–17.

12.   Saxena AK, Kumar M, Chakdar H, Anuroopa N, Bagyaraj DJ. 2020. Bacillus species in soil as a natural resource for plant health and nutrition. J Appl Microbiol 128:1583–1594.

13.   Chen XH, Koumoutsi A, Scholz R, Eisenreich A, Schneider K, Heinemeyer I, Morgenstern B, Voss B, Hess WR, Reva O, Junge H, Voigt B, Jungblut PR, Vater

J, Süssmuth R, Liesegang H, Strittmatter A, Gottschalk G, Borriss R. 2007. Comparative analysis of the complete genome sequence of the plant growth-promoting bacterium Bacillus amyloliquefaciens FZB42. Nat Biotechnol 25:1007–1014.

14. Chen XH, Koumoutsi A, Scholz R, Schneider K, Vater J, Süssmuth R, Piel J, Borriss R. 2009. Genome analysis of Bacillus amyloliquefaciens FZB42 reveals its potential for biocontrol of plant pathogens. J Biotechnol 140:27–37.

15. Molinatto G, Puopolo G, Sonego P, Moretto M, Engelen K, Viti C, Ongena M, Pertot I. 2016. Complete genome sequence of Bacillus amyloliquefaciens subsp. plantarum S499, a rhizobacterium that triggers plant defences and inhibits fungal phytopathogens. J Biotechnol 238:56–59.

16. van Gestel J, Vlamakis H, Kolter R. 2015. From Cell Differentiation to Cell Collectives: Bacillus subtilis Uses Division of Labor to Migrate. PLoS Biol 13:1–29.

17. López D, Vlamakis H, Losick R, Kolter R. 2009. Paracrine signaling in a bacterium. Genes Dev 23:1631–1638.

18. Aleti G, Lehner S, Bacher M, Compant S, Nikolic B, Plesko M, Schuhmacher R, Sessitsch A, Brader G. 2016. Surfactin variants mediate species-specific biofilm formation and root colonization in Bacillus. Environ Microbiol 18:2634–2645.

19. Pršić J, Ongena M. 2020. Elicitors of Plant Immunity Triggered by Beneficial Bacteria. Front Plant Sci 11:1–12.

20. Debois D, Fernandez O, Franzil L, Jourdan E, de Brogniez A, Willems L, Clément C, Dorey S, De Pauw E, Ongena M. 2015. Plant polysaccharides initiate underground crosstalk with bacilli by inducing synthesis of the immunogenic lipopeptide surfactin. Environ Microbiol Rep 7:570–582.

21. Gage DJ. 2004. Infection and Invasion of Roots by Symbiotic, Nitrogen-Fixing Rhizobia during Nodulation of Temperate Legumes. Microbiol Mol Biol Rev 68:280–300.

22. Beauregard PB, Chai Y, Vlamakis H, Losick R, Kolter R. 2013. Bacillus subtilis biofilm induction by plant polysaccharides. Proc Natl Acad Sci U S A 110:E1621-30.

23. Fan B, Wang C, Song X, Ding X, Wu L, Wu H, Gao X, Borriss R. 2018. Bacillus velezensis FZB42 in 2018: The gram-positive model strain for plant growth promotion and biocontrol. Front Microbiol 9:1–14.

24. Mohnen D. 2008. Pectin structure and biosynthesis. Curr Opin Plant Biol 11:266–277.

25. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. 2014. The carbohydrate-active enzymes database (CAZy) in 2013. Nucleic Acids Res 42:490–495.

26. Fan B, Blom J, Klenk HP, Borriss R. 2017. Bacillus amyloliquefaciens, Bacillus velezensis, and Bacillus siamensis Form an "Operational Group B.

amyloliquefaciens" within the B. subtilis species complex. Front Microbiol 8:22.

27.   Nihorimbere V, Cawoy H, Seyer A, Brunelle A, Thonart P, Ongena M. 2012. Impact
of rhizosphere factors on cyclic lipopeptide signature from the plant beneficial strain
Bacillus amyloliquefaciens S499. FEMS Microbiol Ecol 79:176–191.

28.   Chen Y, Yan F, Chai Y, Liu H, Kolter R, Losick R, Guo JH. 2013. Biocontrol of
tomato wilt disease by Bacillus subtilis isolates from natural environments depends
on conserved genes mediating biofilm formation. Environ Microbiol 15:848–864.

29.   Grau RR, De Oña P, Kunert M, Leñini C, Gallegos-Monterrosa R, Mhatre E, Vileta
D, Donato V, Hölscher T, Boland W, Kuipers OP, Kovács ÁT. 2015. A duo of
potassium-responsive histidine kinases govern the multicellular destiny of Bacillus
subtilis. MBio 6:1–16.

30.   Thérien M, Kiesewalter HT, Auria E, Charron-Lamoureux V, Wibowo M, Maróti G,
Kovács ÁT, Beauregard PB. 2020. Surfactin production is not essential for pellicle
and root-associated biofilm development of Bacillus subtilis. Biofilm 2:100021.

31.   Vlamakis H, Chai Y, Beauregard P, Losick R, Kolter R. 2013. Sticking together:
Building a biofilm the Bacillus subtilis way. Nat Rev Microbiol 11:157–168.

32.   Caramori T, Barillà D, Nessi C, Sacchi L, Galizzi A. 1996. Role of FlgM in σD-
dependent gene expression in Bacillus subtilis. J Bacteriol 178:3113–3118.

33.   Strieker M, Tanović A, Marahiel MA. 2010. Nonribosomal peptide synthetases:
Structures and dynamics. Curr Opin Struct Biol 20:234–240.

34.   Süssmuth RD, Mainz A. 2017. Nonribosomal Peptide Synthesis—Principles and
Prospects. Angew Chemie - Int Ed 56:3770–3821.

35.   Diomandé SE, Nguyen-The C, Guinebretière MH, Broussolle V, Brillard J. 2015.
Role of fatty acids in Bacillus environmental adaptation. Front Microbiol 6:1–20.

36.   Serror P, Sonenshein AL. 1996. CodY is required for nutritional repression of
Bacillus subtilis genetic competence. J Bacteriol 178:5910–5915.

37.   Dhali D, Coutte F, Arias AA, Auger S, Bidnenko V, Chataigné G, Lalk M, Niehren
J, de Sousa J, Versari C, Jacques P. 2017. Genetic engineering of the branched
fatty acid metabolic pathway of Bacillus subtilis for the overproduction of surfactin
C14 isoform. Biotechnol J 12:1–10.

38.   Brinsmade SR, Kleijn RJ, Sauer U, Sonenshein AL. 2010. Regulation of CodY
activity through modulation of intracellular branched-chain amino acid pools. J
Bacteriol 192:6357–6368.

39.   Cawoy H, Mariutto M, Henry G, Fisher C, Vasilyeva N, Thonart P, Dommes J,
Ongena M. 2014. Plant Defense Stimulation by Natural Isolates of Bacillus
Depends on Efficient Surfactin Production. Mol Plant-Microbe Interact 27:87–100.

40.   Saijo Y, Loo EP iian, Yasuda S. 2018. Pattern recognition receptors and signaling
in plant–microbe interactions. Plant J 93:592–613.

41.   Waszczak C, Carmody M, Kangasjärvi J. 2018. Reactive Oxygen Species in Plant

Signaling. Annu Rev Plant Biol 69:209–236.

42.   Mignolet-Spruyt L, Xu E, Idänheimo N, Hoeberichts FA, Mühlenbock P, Brosche M,
Van Breusegem F, Kangasjärvi J. 2016. Spreading the news: Subcellular and
organellar reactive oxygen species production and signalling. J Exp Bot 67:3831–
3844.

43.   Ashtamker C, Kiss V, Sagi M, Davydov O, Fluhr R. 2007. Diverse subcellular
locations of cryptogein-induced reactive oxygen species production in tobacco
bright yellow-2 cells. Plant Physiol 143:1817–1826.

44.   Bigeard J, Colcombet J, Hirt H. 2015. Signaling mechanisms in pattern-triggered
immunity (PTI). Mol Plant 8:521–539.

45.   Jourdan E, Henry G, Duby F, Dommes J, Barthélemy JP, Thonart P, Ongena M.
2009. Insights into the defense-related events occurring in plant cells following
perception of surfactin-type lipopeptide from Bacillus subtilis. Mol Plant-Microbe
Interact 22:456–468.

46.   Henry G, Deleu M, Jourdan E, Thonart P, Ongena M. 2011. The bacterial
lipopeptide surfactin targets the lipid fraction of the plant plasma membrane to
trigger immune-related defence responses. Cell Microbiol 13:1824–1837.

47.   Zipfel C, Oldroyd GED. 2017. Plant signalling in symbiosis and immunity. Nature
543:328–336.

48.   Schellenberger R, Touchard M, Clément C, Baillieul F, Cordelier S, Crouzet J,
Dorey S. 2019. Apoplastic invasion patterns triggering plant immunity: plasma
membrane sensing at the frontline. Mol Plant Pathol 20:1602–1616.

49.   Martinez-Medina A, Flors V, Heil M, Mauch-Mani B, Pieterse CMJ, Pozo MJ, Ton
J, van Dam NM, Conrath U. 2016. Recognizing Plant Defense Priming. Trends
Plant Sci 21:818–822.

50.   Levy A, Salas Gonzalez I, Mittelviefhaus M, Clingenpeel S, Herrera Paredes S,
Miao J, Wang K, Devescovi G, Stillman K, Monteiro F, Rangel Alvarez B, Lundberg
DS, Lu TY, Lebeis S, Jin Z, McDonald M, Klein AP, Feltcher ME, Rio TG, Grant SR,
Doty SL, Ley RE, Zhao B, Venturi V, Pelletier DA, Vorholt JA, Tringe SG, Woyke T,
Dangl JL. 2018. Genomic features of bacterial adaptation to plants. Nat Genet
50:138–150.

51.   Fan B, Carvalhais LC, Becker A, Fedoseyenko D, Von Wirén N, Borriss R. 2012.
Transcriptomic profiling of Bacillus amyloliquefaciens FZB42 in response to maize
root exudates. BMC Microbiol 12.

52.   Feng H, Zhang N, Du W, Zhang H, Liu Y, Fu R, Shao J, Zhang G, Shen Q, Zhang
R. 2018. Identification of chemotaxis compounds in root exudates and their sensing
chemoreceptors in plant-growth-promoting rhizobacteria bacillus amyloliquefaciens
SQR9. Mol Plant-Microbe Interact 31:995–1005.

53.   Zhang N, Yang D, Wang D, Miao Y, Shao J, Zhou X, Xu Z, Li Q, Feng H, Li S, Shen
Q, Zhang R. 2015. Whole transcriptomic analysis of the plant-beneficial
rhizobacterium Bacillus amyloliquefaciens SQR9 during enhanced biofilm

formation regulated by maize root exudates. BMC Genomics 16:1–21.

54. Raaijmakers JM, de Bruijn I, Nybroe O, Ongena M. 2010. Natural functions of
lipopeptides from Bacillus and Pseudomonas: More than surfactants and
antibiotics. FEMS Microbiol Rev 34:1037–1062.

55. HayashiTaku K. 2014. The H 2 O 2 Stress-Responsive Regulator PerR Positively
Regulates srfA Expression in Bacillus subtilis The H 2 O 2 Stress-Responsive
Regulator PerR Positively Regulates srfA Expression in Bacillus subtilis 187:6659–
6667.

56. Wolf D, Rippa V, Mobarec JC, Sauer P, Adlung L, Kolb P, Bischofs IB. 2015. The
quorum-sensing regulator ComA from Bacillus subtilis activates transcription using
topologically distinct DNA motifs. Nucleic Acids Res 44:2160–2172.

57. Mariappan A, Makarewicz O, Chen XH, Borriss R. 2012. Two-component response
regulator DegU controls the expression of bacilysin in plant-growth-promoting
bacterium bacillus amyloliquefaciens FZB42. J Mol Microbiol Biotechnol 22:114–
125.

58. Zhi Y, Wu Q, Xu Y. 2017. Genome and transcriptome analysis of surfactin
biosynthesis in Bacillus amyloliquefaciens MT45. Sci Rep 7:1–13.

59. Koumoutsi A, Chen XH, Vater J, Borriss R. 2007. DegU and YczE positively
regulate the synthesis of bacillomycin D by Bacillus amyloliquefaciens strain
FZB42. Appl Environ Microbiol 73:6953–6964.

60. Konishi H, Hio M, Kobayashi M, Takase R, Hashimoto W. 2020. Bacterial
chemotaxis towards polysaccharide pectin by pectin-binding protein. Sci Rep 10:1–
12.

61. Muller DB, Schubert OT, Rost H, Aebersold R, Vorholt JA. 2016. Systems-level
proteomics of two ubiquitous leaf commensals reveals complementary adaptive
traits for phyllosphere colonization. Mol Cell Proteomics 15:3256–3269.

62. Hossain MJ, Ran C, Liu K, Ryu C-MM, Rasmussen-Ivey CR, Williams MA, Hassan
MK, Choi S-KK, Jeong H, Newman M, Kloepper JW, Liles MR. 2015. Deciphering
the conserved genetic loci implicated in plant disease control through comparative
genomics of Bacillus amyloliquefaciens subsp. plantarum. Front Plant Sci 6:1–14.

63. Birkenbihl RP, Liu S, Somssich IE. 2017. Transcriptional events defining plant
immune responses. Curr Opin Plant Biol 38:1–9.

64. Huot B, Yao J, Montgomery BL, He SY. 2014. Growth-defense tradeoffs in plants:
A balancing act to optimize fitness. Mol Plant.

65. Trauth S, Bischofs IB. 2014. Ectopic integration vectors for generating fluorescent
promoter fusions in Bacillus subtilis with minimal dark noise. PLoS One 9.

66. Bryksin A V., Matsumura I. 2010. Overlap extension PCR cloning: a simple and
reliable way to create recombinant plasmids. Biotechniques 48:463–465.

67. Jarmer H, Berka R, Knudsen S, Saxild HH. 2002. Transcriptome analysis
documents induced competence of Bacillus subtilis during nitrogen limiting

conditions. FEMS Microbiol Lett 206:197–200.

68. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
Nikolenko SI, Pham S, Prjibelski AD, Pyshkin A V., Sirotkin A V., Vyahhi N, Tesler
G, Alekseyev MA, Pevzner PA. 2012. SPAdes: A new genome assembly algorithm
and its applications to single-cell sequencing. J Comput Biol 19:455–477.

69. De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. 2018. NanoPack:
Visualizing and processing long-read sequencing data. Bioinformatics 34:2666–
2669.

70. Antipov D, Korobeynikov A, McLean JS, Pevzner PA. 2016. HybridSPAdes: An
algorithm for hybrid assembly of short and long reads. Bioinformatics 32:1009–
1015.

71. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina
sequence data. Bioinformatics 30:2114–2120.

72. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25:1754–1760.

73. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R. 2009. The Sequence Alignment/Map format and SAMtools.
Bioinformatics 25:2078–2079.

74. Anders S, Pyl PT, Huber W. 2015. HTSeq-A Python framework to work with high-
throughput sequencing data. Bioinformatics 31:166–169.

75. Trapnell C, Williams B a, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg
SL, Wold BJ, Pachter L. 2011. Transcript assembly and abundance estimation from
RNA-Seq reveals thousands of new transcripts and switching among isoforms. Nat
Biotechnol 28:511–515.

76. Molinatto G, Franzil L, Steels S, Puopolo G, Pertot I, Ongena M. 2017. Key impact
of an uncommon plasmid on bacillus amyloliquefaciens subsp. plantarum S499
developmental traits and lipopeptide production. Front Microbiol 8:1–18.

77. Debois D, Ongena M, Cawoy H, De Pauw E. 2013. MALDI-FTICR MS imaging as
a powerful tool to identify paenibacillus antibiotics involved in the inhibition of plant
pathogens. J Am Soc Mass Spectrom 24:1202–1213.

78. Carpita NC. 1984. Cell wall development in maize coleoptiles. Plant Physiol
76:205–212.

79. Silva GB, Ionashiro M, Carrara TB, Crivellari AC, Tiné MAS, Prado J, Carpita NC,
Buckeridge MS. 2011. Cell wall polysaccharides from fern leaves: Evidence for a
mannan-rich Type III cell wall in Adiantum raddianum. Phytochemistry 72:2352–
2360.

80. Bisceglia N, Gravino M, Savatin D. 2015. Luminol-based Assay for Detection of
Immunity Elicitor-induced Hydrogen Peroxide Production in Arabidopsis thaliana
Leaves. Bio-Protocol 5.

81. Heerklotz H, Seelig J. 2000. Titration calorimetry of surfactant-membrane

partitioning and membrane solubilization. Biochim Biophys Acta - Biomembr
1508:69–85.

82. Van Bambeke F, Kerkhofs A, Schanck A, Remacle C, Sonveaux E, Tulkens PM,
Mingeot-Leclercq M-P. 2000. Biophysical studies and intracellular destabilization of
pH-sensitive liposomes. Lipids 35:213–223.

83. Arguelles-Arias A, Ongena M, Halimi B, Lara Y, Brans A, Joris B, Fickers P. 2009.
Bacillus amyloliquefaciens GA1 as a source of potent antibiotics and other
secondary metabolites for biocontrol of plant pathogens. Microb Cell Fact 8:1–12.

84. Pandin C, Le Coq D, Deschamps J, Védie R, Rousseau T, Aymerich S, Briandet R.
2018. Complete genome sequence of *Bacillus velezensis* QST713: A biocontrol
agent that protects Agaricus bisporus crops against the green mould disease. J
Biotechnol 278:10–19.

85. Serrano L, Manker D, Brandi F, Cali T. 2013. The use of *Bacillus subtilis* QST 713
and *Bacillus pumilus* qst 2808 as protectant fungicides in conventional application
programs for black leaf streak control. Acta Hortic 986:149–156.

832
833

Figures and Tables

Figure 1: Impact of pectin on early surfactin production. a Sugar composition analysis of crude pectin (cPec) extracted from tobacco roots. Composition is expressed as Molar Ratio percentage (Molar %) for each fraction. Galacturonic acid (orange) constituting the pectin backbone (**b** for schematization) is the main sugar of the cPec fraction. Other minor sugars (rhamnose, galactose, arabinose…) are typically found in the pectin side chains (Mohnen et al. 2008, **b**). **b** Schematization of pectin structure. Homogalacturonan (HG) contains an assembly of at least 100 galacturonic acid (GalA) residues that can be acetyl or methyl esterified. Rhamnogalacturonan I (RGI) is constituted by a succession of GalA-Rha dimers, each one containing an alternance of rhamnosyl and galacturonic acid units. The Rha unit can be branched with variable neutral sugar side chains including essentially galactosyl and/or arabinosyl units. Rhamnogalacturonan II (RGII) structure is well conserved within the HG polymer. RGII englobes 9 GalA units substituted by four side chains with complex sugars, including apiose, DHA, aceric acid and KDO, neutral sugars like, rhamnose, galactose, arabinose, xylose, and fucose or also organic acids such as galacturonic and glucuronic acid. RGII can also complex with Bore allowing a crosslink between two HG molecules. **c** Surfactin (cyclic structure represented up) production in a root exudates mimicking (REM) medium at early growth phase (OD_{600}=0.2) with (red chromatogram) or without (blue chromatogram) addition of crude pectin extract added to the GA1 cultures. The main peak represents C15 surfactin whereas the minor left and right peaks represents C14 and C16 surfactins, respectively. **d** Surfactin accumulation in the early (left panel, OD_{600}=0.2) and mid (right panel, OD_{600}=0.35) exponential growth phase of GA1 cultures in REM medium supplemented with different sized pectin fragments : homogalacturonan low methylated (HGLM), DP>150; oligogalacturonides (OG), DP=15; galacturonic acid (GA), DP=1. Means ± std err. from three biological replicates of one experiment are shown. Significate difference between each condition is indicated by different letters, p-value < 0.01. **e** Comparison of surfactin induction level by HGLM in the early exponential growth phase for different *Bacillus* species : *Bacillus velezensis* (*B; v*), *Bacillus amyloliquefaciens* (*B. a*), *Bacillus subtilis* (*B. s*) and *Bacillus pumilus* (*B. p*). For each strain tested, surfactin accumulation was normalized with the control condition without HGLM represented by the black dotted line. Means ± std err. from three biological replicates are shown.

Figure 2: Impact of the specific rhizosphere nutritional context on early surfactin production. a Evaluation of bacterial population (black line, left axis) and relative *srfAA* expression on roots (grey bars, right axis) in a time frame of seven days post inoculation (dpi). *Bacillus* progression on roots characterized by a biofilm formation was assessed by microscopy at each time point (upper part). **b** Surfactin production rate on roots. Means ± std err. from three biological replicates of one experiment are shown **c** Surfactin accumulation measured by UPLC-MS in a 8h time course experiment in REM medium (grey bars) compared to LB medium (red bars). Means ± std err. from three biological replicates of one experiment are shown *** P-value <0.001, ** P-value <0.01, * P-value <0.05 **d** Comparison of early surfactin accumulation (µM of surfactin on y axis linked to OD_{600} on x axis) in different *Bacillus* species, including *B. velezensis* (GA1 and S499 in green), *B. pumilus* (QST 2808 in orange), *B. amyloliquefaciens* (DSM 7 in red) and *B. subtilis* (ATCC 21332 in blue). Circle symbols are representing plant associated bacteria whereas triangle symbols are representing non-plant associated bacteria.

Figure 3: Ecological importance of an early surfactin accumulation. a Evaluation of HGLM and root exudates synergistic effect on early surfactin production. Time course experiment for surfactin quantification was performed in REM (grey curves) and LB (red curves) medium with (circle symbols) or without (square symbols) addition of HGLM. Means ± std err. from three biological replicates of one experiment are shown **b** Swarming potential of *B. velezensis* GA1 on soft agar plates after addition of HGLM or not. The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=7 biological replicates of one experiment). **c** Evaluation of *B. velezensis* ability to form pellicles on microwells plates after addition of HGLM or not. The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=8 biological replicates of one experiment). Pellicle formation is illustrated on the right **d** Comparison of *B. velezensis* GA1 WT and (red) and a ∆*srfAA* mutant (grey) for their swarming potential in a time course study. Means ± std err. from three biological replicates of one experiment are shown. Time course study is illustrated right. **e** Comparison of pellicle formation between GA1 WT strain (red) and a ∆*srfAA* mutant (grey)**.** The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=8 biological replicates of one experiment) **** P-value <0.0001. **f** *In vitro* comparison of root colonization ability of GA1 (red boxes) and GA1 ∆*srfAA* (grey boxes) on tomato plantlets. The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=7 biological replicates of one experiment) *** P-value <0.001, ns non significant.

Figure 4: Impact of plant triggers perception on *Bacillus* transcriptome. a Surfactin expression measured by fluorescence in the GA1 *PsrfAp::gfp* reporter strain at early exponential phase in REM medium (grey bars) compared to REM medium supplemented with HGLM (red bars). Means ± std err. from three biological replicates of one representative experiment are shown ns = non significate **b** Surfactin expression measured by fluorescence in the GA1 P*srfAp::gfp* reporter strain in a 24h time course study in EM medium (grey bars) compared to LB medium (red bars). Means ± std err. from three biological replicates of one representative experiment are shown *** P-value <0.001. **c** Classification of the different genes carrying a significant fold change (1.5 log2) 5 and 8 hours after addition of HG when compared to the control condition. The outer circle represents the proportion of up (dark blue) and down (red) regulated genes. The inner circle represents the proportion of genes belonging to the different functional family described in the legend.

Figure 5: Qualitative impact of root exudates on surfactin production. **a** Representation of the NRPS machinery leading to the assembly of the surfactin molecule. This mega-enzyme is organized in 7 functional units called modules which are each responsible for the incorporation of one amino acid building block into the growing peptide chain. Each module is subdivided into different domains including an adenylation (A, violet circle) and a peptidyl carrier protein (PCP, red circle) catalyzing the peptide initiation, and one condensation domain (C, brown circle) responsible for peptide elongation. The termination of the peptide synthesis is performed by a thioesterase domain (TE, blue circle) in the last module. Modules 3 and 6 also possess an epimerization domain (E, green circle). Surfactin molecule contains a 7 amino acids chain structured as follow: L-Glu – L-Leu – D-Leu – L-Val – L-Asp – D-Leu – L-Leu. In some specific variants, Leu in position 2 and/or 7can be substituted by a Val and more rarely by an Ile, and inversely, Val in position 4 can be substituted by a Leu and also more rarely by a Ile. In addition to

the amino acid chain variability, multiple homologs with the same peptidic core but
differences in terms of fatty acid chain length (C_{12} to C_{17}) or isomerisation of this latter (iso,
anteiso or linear configuration) can also be produced. **b** Comparison of surfactin pattern
in REM and LB medium. Based on MS-MS analyses, nine different surfactin forms were
identified (a: C_{12}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; b: C_{13}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; c
: *iso*-C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; c': *n*-C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; d : C_{15}-
Glu-Leu-Leu-Val-Asp-Leu-Leu ; e: C_{13}-Glu-Leu-Leu-Val-Asp-Leu-Val ; f: C_{14}-Glu-Leu-
Leu-Val-Asp-Leu-Val ; g: C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Val and h: C_{14}-Glu-Val-Leu-Val-
Asp-Leu-Val.) **c** Relative proportions of surfactin variants in LB, REM, REM supplemented
with valine, and *in planta.* **d** Qualitative and quantitative role of CodY on surfactin
production. In a WT strain, 95% of the surfactin molecules are carrying a Leu in position
7 (grey bars) and only 5% are carrying a Val (red bars) whereas in Δ*codY* mutant almost
25% of the surfactin molecules are carrying a Val in position 7 and 75% a Leu. In addition,
amount of total surfactin production rate of 150 % can be observed in Δ*codY* mutant
compared to WT strain. Proportion of iso-C14 is also affected by CodY, 36 % of total C_{14}
are iso-fatty acid (grey bars) and 64% are linear (red bars) in WT strain whereas in Δ*codY*
mutant 55% of C_{14} are *iso*-C_{14} and 45 % are linear. Again, total amount of C_{14} is higher in
Δ*codY* mutant (increase of 190 %).

Figure 6: Impact of surfactin homologues on Solanaceae plant immunity. abc
Systemic resistance induced in hydroponically-grown tobacco by surfactin and expressed
as reduction of *B. cinerea* infection (illustration of the reduction in the diameter of
spreading lesions on infected leaves) in plants treated at the root level prior to pathogen
inoculation on leaves compared to control plants. Data represent results grouped from two
independent experiments with similar results and each involving 5 plants with 4 lesions on
the second leave (n=40). The box plots encompass the 1st and 3rd quartile, the whiskers
extend to the minimum and maximum points, and the midline indicates the median (n=7
biological replicates of one experiment). **a** Effect of surfactin homologues (SF mix) as
naturally co-produced by the bacterium (C12/C13/C14/C15 in relative proportions
8/17/33/42%) **** P-value <0.0001   **b** Effect of HPLC-purified surfactin homologues
applied at 10 µM with fatty acid chains from C12 to C15. Significate difference between
each condition is indicated by different letters, p-value < 0.05 **c** Effect of the most active
C14 homologue tested at various concentrations. Significate difference between each
condition is indicated by different letters, p-value < 0.05 **de** Stimulation of oxidative burst
in root tissues upon treatment with a SF mix and to the response observed by treating
roots with flagellin (flg22, 1 µM) used as positive control. **d** Stimulation of apoplastic ROS
accumulation (DCFH-DA fluorescent probe) in root tissues upon treatment with a surfactin
mix applied at 15 µM. Means and standard deviations are shown for one representative
experiment performed on nine samples per treatment each containing three root segments
(approx 100 mg FW) collected from different plants (n=9). Similar trend was obtained in
an independent assay. **e** Stimulation of cytoplasmic hydrogen peroxide production in root
cells. Means and s.d. were calculated from measurements performed on three samples
per treatment each containing three root segments (approx 100 mg FW) collected from
different plants. Data represent values obtained from two independent experiments (n=6
per treatment). **f** Stimulation of cytoplasmic hydrogen peroxide production in root cells
after treatment with C_{12} and C_{14} surfactin homologues as representative of short and long
fatty acid chains respectively. Flg22 was used as control. The box plots encompass the
1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the
midline indicates the median (n=6 biological replicates of one experiment).Significate
difference between each condition is indicated by different letters, p-value < 0.0001. **g**
Binding coefficient (K) of Surfactin homologues (C_{12} to C_{15}) to large unilamellar vesicles

(LUV) composed by PLPC/Sitosterol/Glucosylceramide (60:20:20 molar ratio).
Means ± std err. from three to five biological replicates of one representative experiment
are shown Significate difference between each condition is indicated by different letters,
p-value < 0.05 **h** Release of 8-hydroxypyrene-1,3,6 trisulfonic acid (HPTS) from
PLPC/Sitosterol/Glucosylceramide (60:20:20 molar ratio) LUV, upon addition of surfactin
C_{12} or C_{14} at different concentrations. The ordinate shows the amount of HPTS released
after 15 min in the presence of the C_{12} or C_{14} as a percentage of the total amount released
by Triton X-100. **i** Influence of roots pretreatment with 10µM of surfactin (blue boxes)
compared to non-treated roots (red boxes) on *B. velezensis* GA1 root colonization. The
box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and
maximum points, and the midline indicates the median (n=5 biological replicates of one
experiment). Ns= non significate.

Table 1: Differentially expressed genes in *B. velezensis* GA1 after HGLM perception
995

Locus_tag	Name	Fold change		Conservation				Category	Informations
		5h	8h	*B. amylo.* DSM7		*B. subtilis* 168			
				QC (%)	ID (%)	QC (%)	ID (%)		
GL331_00010	*floT (yuaG)*		-3.048	100	93.2	97	81.46	Stress	Inner membrane protein, flotillin-like protein
GL331_00015	*yuaF*	-3.016		99	93.4	94	68.97	Stress	Involved in the control of membrane fluidity
GL331_00985	*yusU*	-3.072		100	95.12	99	76.37	Unknown	Unknown
GL331_01085	*liaH*		-4.878	100	93.95	100	77.21	Stress	*lia* operon expression modulator, similar to phage shock protein, resistance against oxidative stress and cell wall antibiotics
GL331_01090	*liaI*		-3.846	100	93.96	100	69.71		Resistance against oxidative stress and cell wall antibiotics
GL331_01180	*fhuD*	-3.176		100	95.1	99	77.31	Transport	Hydroxamate siderophore ABC transporter
GL331_01520	*opuCA*		3.066	100	95.61	99	82.35	Transport	Glycine betaine/carnitine/choline/arsenobetaine/arsenocholine ABC transporter
GL331_02210	*hpf*	-4.636		100	97	100	83.71	Stress	General stress protein, required for ribosome dimerization in the stationary phase, protects essential ribosomal proteins
GL331_02265	*flgN*	-3.608		100	96.48	76	76.55	Motility	Flagellar filament assembly protein
GL331_02270	*flgM*	-4.334		100	98.5	100	77.15		Negative regulator of flagellin synthesis, anti-*sigD*

Locus	Gene	FC1	FC2					Category	Description
GL331_02275	yvyF	-4.562		100	95.71	100	78.38		Unknown
GL331_02335	tuaF		-5.05	100	92.51	100	70.91	Cell wall	Teichuronic acid biosynthesis protein
GL331_02355	tuaB		-3.262	100	94.41	100	73.25	Cell wall	Polymer export
GL331_03100	maeA		-3.69	100	97.29	100	82.88	Metabolism	Malate utilization
GL331_03645	ywcE		-3.516	100	96.93	89	85.04	Sporulation	Holin, spore morphogenesis and germination protein
GL331_03950	cydB	-5.742	-3.506	100	95.38	100	79.45	Metabolism	Cytochrome d ubiquinol oxidase subunit 2
GL331_03955	cydA	-4.792		100	96.52	100	82.3		Cytochrome d ubiquinol oxidase subunit 1
GL331_04110			-3.374	/	/	/	/	Unknown	Hypothetical protein
GL331_04120	wapA		-4.134	/	/	92	77.81	Cell wall	Cell wall-associated protein precursor
GL331_04135	yxiE	-3.784		100	95.53	98	78.25	Stress	Universal stress protein
GL331_04235		-3.672		100	94.18	89	75.79	Transport	Hydroxamate siderophore ABC transporter
GL331_04550			3.198	99	96.39	/	/	Transport	Transport protein (putative quinolone resistance)
GL331_04560	thiF		3.138	/	/	/	/	Metabolism	Thiazole biosynthesis adenylyltransferase
GL331_04675			-3.768	100	92.69	/	/	Unknown	PIG-L family deacetylase
GL331_05550	ctsR	-3.034		100	96.77	100	86.45	Regulation	Transcriptional repressor, protein synthesis, modification and degradation
GL331_06040	btr	-3.352		100	91.23	72	68.27	Regulation	Transcriptional activator, regulation of iron acquisition
GL331_06530	yceF		-3.108	100	96.77	100	82.43	Stress	General stress protein, manganese resistance protein
GL331_06540	yceH		-3.104	100	95.22	98	83.68		Similar to toxic anion resistance protein
GL331_06585		-3.342		100	94.65	97	81.42	Metabolism	L-lactate dehydrogenase
GL331_07095	yczJ		-4.126	100	95.64	99	74.38	Unknown	Unknown
GL331_07100			-3.124						
GL331_07310	gsiB	-3.068		100	96.6	93	89.97	Stress	General stress protein, response to glucose starvation or to water deficits

Locus	Gene							Category	Description
GL331_08210	pspA		-3.664	100	93.74	96	74.52	Stress	Phage shock protein A homolog, paralogous to liaH
GL331_08215	ydjG		-3.246	99	94.43	99	75.05	Stress	Unknown
GL331_08225	ydjI		-3.014	99	97.01	98	77.93		Unknown
GL331_09875	nhaX	-3.04		99	92.59	99	74.9	Stress	Universal stress protein
GL331_10225	yhzC	-3.306		100	94.87	100	85.04	Unknown	Unknown
GL331_10530	argJ		3.852	99	94.82	99	77.23		
GL331_10535	argB		5.39	100	91.51	100	71.5	Metabolism	Biosynthesis of arginine
GL331_10540	argD		4.64	100	93.96	96	74.62		
GL331_10545	carA		5.726	100	93.48	100	73.46		
GL331_10715	cwlQ	-3.67		100	92.43	75	77.02	Cell wall	Bifunctional cell wall hydrolase
GL331_14205	iseA	-5.102		100	93.19	100	72	Cell wall	Cell wall endopeptidases and cell separation inhibitor
GL331_15495	sigX		-3.024	100	97.44	99	87.18	Sigma factor	RNA polymerase sigma factor
GL331_15575	ribH		3.352	100	95.91	100	81.29		
GL331_15580	ribA		3.946	100	94.57	100	78.11	Metabolism	Riboflavin biosynthesis
GL331_15585	ribE		3.896	100	94.29	100	74.88		
GL331_15590	ribD		4.102	100	94.44	99	76.73		
GL331_15930	loaP	-3.848		/	/	/	/	Regulation	Antiterminator involved in regulation of polyketide synthesis
GL331_16515	pstBB		-5.384	99	92.7	93	73.36		
GL331_16520	pstBA		-4.924	100	92.32	89	77	Metabolism, Transport	High-affinity phosphate uptake, phosphate ABC transporter
GL331_16525	pstA		-4.038	100	94.8	99	79.07		
GL331_17335			-5.61	100	89.17	48	75.86	Unknown	Unknown
GL331_17345	safA		-4.26	99	89.73	42	76.04	Sporulation	Major organizer of the inner spore coat
GL331_17895	pftB	-3.294		100	95.03	100	81.73		Pyruvate transporter

GL331_17900	*pftA*	-3.958		100	97.28	83	79.89	Metabolism, Transport	
GL331_18085	*ytzJ*	-3.29		100	98.96	100	84.38	Unknown	Unknown
GL331_18115	*argH*		5.002	100	95.06	99	83.98	Metabolism	Biosynthesis of arginine
GL331_18120	*argG*		5.144	100	95.3	100	81.77		

Table 2 : Strains used in this study

Strain	Characteristics	Sources
Bacillus velezensis		
B. velezensis GA1	Wild-type strain	(83)
B. velezensis GA1 P*srf_gfp*	*amyE*::P*srf_gfp*+*chl* ; Chl+	This study
*B. velezensis*GA1 Δ*srfAA*	Δ*srfAA*::*chl* ; Chl+	This study
*B. velezensis*GA1 Δ*codY*	Δ*codY*::*chl* ; Chl+	This study
B. velezensis S499	Wild-type strain	(15)
B. velezensis FZB42	Wild-type strain	(13)
B. velezensis QST713	Wild-type strain	(84)
Bacillus amyloliquefaciens		
B. amyloliquefaciens DSM7	Wild-type strain	ATCC
Bacillus subtilis		
B. subtilis ATCC 21332	Wild-type strain	ATCC
Bacillus pumilus		
B. pumilus QST 2808	Wild-type strain	(85)
Escherichia coli		
E. coli dh5α	Wild-type strain	CGSC
E. coli dh5α pGEM-T Easy *amyE*	pGEM-T Easy *amyE ;* Amp+	This study
E. coli dh5α pGEM-T Easy *amyE*up-*cat-gfp-amyE*dw	pGEMT-T Easy *amyE*up-*cat-gfp-amyE*dw ; Amp+ Chl+	This study
E. coli dh5α pGFP_Star	pGFP-Star ; Chl+	This study

Figure 1: Impact of pectin on early surfactin production. a Sugar composition analysis of crude pectin (cPec) extracted from tobacco roots. Composition is expressed as Molar Ratio percentage (Molar %) for each fraction. Galacturonic acid (orange) constituting the pectin backbone (**b** for schematization) is the main sugar of the cPec fraction. Other minor sugars (rhamnose, galactose, arabinose…) are typically found in the pectin side chains (Mohnen et al. 2008, **b**). **b** Schematization of pectin structure. Homogalacturonan (HG) contains an assembly of at least 100 galacturonic acid (GalA) residues that can be acetyl or methyl esterified. Rhamnogalacturonan I (RGI) is constituted by a succession of GalA-Rha dimers, each one containing an alternance of rhamnosyl and galacturonic acid units. The Rha unit can be branched with variable neutral sugar side chains including essentially galactosyl and/or arabinosyl units. Rhamnogalacturonan II (RGII) structure is well conserved within the HG polymer. RGII englobes 9 GalA units substituted by four side chains with complex sugars, including apiose, DHA, aceric acid and KDO, neutral sugars like, rhamnose, galactose, arabinose, xylose, and fucose or also organic acids such as galacturonic and glucuronic acid. RGII can also complex with Bore allowing a crosslink between two HG molecules. **c** Surfactin (cyclic structure represented up) production in a root exudates mimicking (REM) medium at early growth phase (OD_{600}=0.2) with (red chromatogram) or without (blue chromatogram) addition of crude pectin extract added to the GA1 cultures. The main peak represents C15 surfactin whereas the minor left and right peaks represents C14 and C16 surfactins, respectively. **d** Surfactin accumulation in the early (left panel, OD_{600}=0.2) and mid (right panel, OD_{600}=0.35) exponential growth phase of GA1 cultures in REM medium supplemented with different sized pectin fragments : homogalacturonan low methylated (HGLM), DP>150; oligogalacturonides (OG), DP=15; galacturonic acid (GA), DP=1. Means ± std err. from three biological replicates of one experiment are shown. Significate difference between each condition is indicated by different letters, p-value < 0.01. **e** Comparison of surfactin induction level by HGLM in the early exponential growth phase for different *Bacillus* species : *Bacillus velezensis* (*B; v*), *Bacillus amyloliquefaciens* (*B. a*), *Bacillus subtilis* (*B. s*) and *Bacillus pumilus* (*B. p*). For each strain tested, surfactin accumulation was normalized with the control condition without HGLM represented by the black dotted line. Means ± std err. from three biological replicates are shown.

Figure 2: Impact of the specific rhizosphere nutritional context on early surfactin production. a Evaluation of bacterial population (black line, left axis) and relative *srfAA* expression on roots (grey bars, right axis) in a time frame of seven days post inoculation (dpi). *Bacillus* progression on roots characterized by a biofilm formation was assessed by microscopy at each time point (upper part). **b** Surfactin production rate on roots. Means ± std err. from three biological replicates of one experiment are shown **c** Surfactin accumulation measured by UPLC-MS in a 8h time course experiment in REM medium (grey bars) compared to LB medium (red bars). Means ± std err. from three biological replicates of one experiment are shown *** P-value <0.001, ** P-value <0.01, * P-value <0.05 **d** Comparison of early surfactin accumulation (µM of surfactin on y axis linked to OD_{600} on x axis) in different *Bacillus* species, including *B. velezensis* (GA1 and S499 in green), *B. pumilus* (QST 2808 in orange), *B. amyloliquefaciens* (DSM 7 in red) and *B. subtilis* (ATCC 21332 in blue). Circle symbols are representing plant associated bacteria whereas triangle symbols are representing non-plant associated bacteria.

Figure 3: Ecological importance of an early surfactin accumulation. a Evaluation of HGLM and root exudates synergistic effect on early surfactin production. Time course experiment for surfactin quantification was performed in REM (grey curves) and LB (red curves) medium with (circle symbols) or without (square symbols) addition of HGLM. Means ± std err. from three biological replicates of one experiment are shown **b** Swarming potential of *B. velezensis* GA1 on soft agar plates after addition of HGLM or not. The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=7 biological replicates of one experiment). **c** Evaluation of *B. velezensis* ability to form pellicles on microwells plates after addition of HGLM or not. The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=8 biological replicates of one experiment). Pellicle formation is illustrated on the right **d** Comparison of *B. velezensis* GA1 WT and (red) and a Δ*srfAA* mutant (grey) for their swarming potential in a time course study. Means ± std err. from three biological replicates of one experiment are shown. Time course study is illustrated right. **e** Comparison of pellicle formation between GA1 WT strain (red) and a Δ*srfAA* mutant (grey)**.** The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=8 biological replicates of one experiment) **** P-value <0.0001. **f** *In vitro* comparison of root colonization ability of GA1 (red boxes) and GA1 Δ*srfAA* (grey boxes) on tomato plantlets. The box plots encompass the 1st and 3rd quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=7 biological replicates of one experiment) *** P-value <0.001, ns non significant.

Figure 4: Impact of plant triggers perception on *Bacillus* transcriptome. a Surfactin expression measured by fluorescence in the GA1 *PsrfAp::gfp* reporter strain at early exponential phase in REM medium (grey bars) compared to REM medium supplemented with HGLM (red bars). Means ± std err. from three biological replicates of one representative experiment are shown ns = non significate **b** Surfactin expression measured by fluorescence in the GA1 P*srfAp::gfp* reporter strain in a 24h time course study in EM medium (grey bars) compared to LB medium (red bars). Means ± std err. from three biological replicates of one representative experiment are shown *** P-value <0.001. **c** Classification of the different genes carrying a significant fold change (1.5 log2) 5 and 8 hours after addition of HG when compared to the control condition. The outer circle represents the proportion of up (dark blue) and down (red) regulated genes. The inner circle represents the proportion of genes belonging to the different functional family described in the legend.

Figure 5: Qualitative impact of root exudates on surfactin production. **a** Representation of the NRPS machinery leading to the assembly of the surfactin molecule. This mega-enzyme is organized in 7 functional units called modules which are each responsible for the incorporation of one amino acid building block into the growing peptide chain. Each module is subdivided into different domains including an adenylation (A, violet circle) and a peptidyl carrier protein (PCP, red circle) catalyzing the peptide initiation, and one condensation domain (C, brown circle) responsible for peptide elongation. The termination of the peptide synthesis is performed by a thioesterase domain (TE, blue circle) in the last module. Modules 3 and 6 also possess an epimerization domain (E, green circle). Surfactin molecule contains a 7 amino acids chain structured as follow: L-Glu – L-Leu – D-Leu – L-Val – L-Asp – D-Leu – L-Leu. In some specific variants, Leu in position 2 and/or 7can be substituted by a Val and more rarely by an Ile, and inversely, Val in position 4 can be substituted by a Leu and also more rarely by a Ile. In addition to the amino acid chain variability, multiple homologs with the same peptidic core but differences in terms of fatty acid chain length (C_{12} to C_{17}) or isomerisation of this latter (iso, anteiso or linear configuration) can also be produced. **b** Comparison of surfactin pattern in REM and LB medium. Based on MS-MS analyses, nine different surfactin forms were identified (a: C_{12}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; b: C_{13}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; c : *iso*-C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; c': *n*-C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; d : C_{15}-Glu-Leu-Leu-Val-Asp-Leu-Leu ; e: C_{13}-Glu-Leu-Leu-Val-Asp-Leu-Val ; f: C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Val ; g: C_{14}-Glu-Leu-Leu-Val-Asp-Leu-Val and h: C_{14}-Glu-Val-Leu-Val-Asp-Leu-Val.) **c** Relative proportions of surfactin variants in LB, REM, REM supplemented with valine, and *in planta.* **d** Qualitative and quantitative role of CodY on surfactin production. In a WT strain, 95% of the surfactin molecules are carrying a Leu in position 7 (grey bars) and only 5% are carrying a Val (red bars) whereas in $\Delta codY$ mutant almost 25% of the surfactin molecules are carrying a Val in position 7 and 75% a Leu. In addition, amount of total surfactin production rate of 150 % can be observed in $\Delta codY$ mutant compared to WT strain. Proportion of iso-C14 is also affected by CodY, 36 % of total C_{14} are iso-fatty acid (grey bars) and 64% are linear (red bars) in WT strain whereas in $\Delta codY$ mutant 55% of C_{14} are *iso*-C_{14} and 45 % are linear. Again, total amount of C_{14} is higher in $\Delta codY$ mutant (increase of 190 %).

Figure 6: Impact of surfactin homologues on Solanaceae plant immunity. abc Systemic resistance induced in hydroponically-grown tobacco by surfactin and expressed as reduction of *B. cinerea* infection (illustration of the reduction in the diameter of spreading lesions on infected leaves) in plants treated at the root level prior to pathogen inoculation on leaves compared to control plants. Data represent results grouped from two independent experiments with similar results and each involving 5 plants with 4 lesions on the second leave (n=40). The box plots encompass the 1^{st} and 3^{rd} quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=7 biological replicates of one experiment). **a** Effect of surfactin homologues (SF mix) as naturally co-produced by the bacterium (C12/C13/C14/C15 in relative proportions 8/17/33/42%) ******** P-value <0.0001 **b** Effect of HPLC-purified surfactin homologues applied at 10 µM with fatty acid chains from C12 to C15. Significate difference between each condition is indicated by different letters, p-value < 0.05 **c** Effect of the most active C14 homologue tested at various concentrations. Significate difference between each condition is indicated by different letters, p-value < 0.05 **de** Stimulation of oxidative burst in root tissues upon treatment with a SF mix and to the response observed by treating roots with flagellin (flg22, 1 µM) used as positive control. **d** Stimulation of apoplastic ROS accumulation (DCFH-DA fluorescent probe) in root tissues upon treatment with a surfactin mix applied at 15 µM. Means and standard deviations are shown for one representative experiment performed on nine samples per treatment each containing three root segments (approx 100 mg FW) collected from different plants (n=9). Similar trend was obtained in an independent assay. **e** Stimulation of cytoplasmic hydrogen peroxide production in root cells. Means and s.d. were calculated from measurements performed on three samples per treatment each containing three root segments (approx 100 mg FW) collected from different plants. Data represent values obtained from two independent experiments (n=6 per treatment). **f** Stimulation of cytoplasmic hydrogen peroxide production in root cells after treatment with C_{12} and C_{14} surfactin homologues as representative of short and long fatty acid chains respectively. Flg22 was used as control. The box plots encompass the 1^{st} and 3^{rd} quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=6 biological replicates of one experiment).Significate difference between each condition is indicated by different letters, p-value < 0.0001. **g** Binding coefficient (K) of Surfactin homologues (C_{12} to C_{15}) to large unilamellar vesicles (LUV) composed by PLPC/Sitosterol/Glucosylceramide (60:20:20 molar ratio). Means ± std err. from three to five biological replicates of one representative experiment are shown Significate difference between each condition is indicated by different letters, p-value < 0.05 **h** Release of 8-hydroxypyrene-1,3,6 trisulfonic acid (HPTS) from PLPC/Sitosterol/Glucosylceramide (60:20:20 molar ratio) LUV, upon addition of surfactin C_{12} or C_{14} at different concentrations. The ordinate shows the amount of HPTS released after 15 min in the presence of the C_{12} or C_{14} as a percentage of the total amount released by Triton X-100. **i** Influence of roots pretreatment with 10µM of surfactin (blue boxes) compared to non-treated roots (red boxes) on *B. velezensis* GA1 root colonization. The box plots encompass the 1^{st} and 3^{rd} quartile, the whiskers extend to the minimum and maximum points, and the midline indicates the median (n=5 biological replicates of one experiment). Ns= non significate.