Frequency of quorum sensing mutations in *Pseudomonas aeruginosa* strains isolated from different environments

Kathleen O'Connor, Conan Y. Zhao & Stephen P. Diggle*

Center for Microbial Dynamics & Infection, School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30332.

*Correspondence: stephen.diggle@biosci.gatech.edu

Keywords: *Pseudomonas aeruginosa*, quorum sensing, LasR, cystic fibrosis

Abstract

Pseudomonas aeruginosa uses quorum sensing (QS) to coordinate the expression of multiple genes necessary for establishing and maintaining infection. *lasR* QS mutations have been shown to frequently arise in cystic fibrosis (CF) lung infections, however, there has been far less emphasis on determining whether QS system mutations arise across other environments. To test this, we utilized 852 publicly available sequenced *P. aeruginosa* genomes from the *Pseudomonas* International Consortium Database (IPCD) to study *P. aeruginosa* QS mutational signatures. We found that across all isolates, LasR is the most variable protein sequence compared to other QS proteins. In order to study isolates by source, we focused on a subset of 654 isolates collected from CF, wounds, and non-infection environmental isolates, where we could clearly identify their source. Using this sub-set analysis, we found that LasR mutations are not specific to CF lungs, but are common across all environments. We then used amino acid length as a proxy for observing loss of function in LasR proteins among the strains. We found that truncated LasR proteins are more abundant in *P. aeruginosa* strains isolated from human infection than the environment. Overall, our findings suggest that the evolution of *lasR* QS mutations in *P. aeruginosa* are common and not limited to infection environments.

Introduction

Pseudomonas aeruginosa is a Gram-negative opportunistic pathogen equipped with a large genome, which enables it to be metabolically versatile and capable of occupying a range of different habitats, especially human and animal impacted environments [1, 2]. It is intrinsically resistant to many classes of antibiotic, and it produces a range of tissue damaging extracellular products such as exoenzymes and phenazine pigments in order to aid its dissemination and spread within a host [2]. *P. aeruginosa* is one of the most prominent bacterial pathogens that colonizes cystic fibrosis (CF) lungs and it is often the dominant CF lung pathogen, particularly as the

41 infection becomes more chronic over time [3]. In addition, *P. aeruginosa* frequently infects
42 chronic wounds, often in conjunction with other microbial species [4].
43
One of the major adaptations of *P. aeruginosa* during chronic infection is the loss of quorum
sensing (QS). QS in *P. aeruginosa* regulates the expression of hundreds of genes, including those
that encode for secreted products and virulence factors [5, 6]. In *P. aeruginosa,* QS is regulated by
a complex hierarchical network of genes, composed of two complete *N*-acyl homoserine lactone
(AHL) circuits, LasR-LasI and RhlR-RhlI, as well as an orphan regulator termed QscR [5, 7]. The
Las and Rhl systems are composed of LuxR-LuxI pairs, homologous to other bacterial QS systems.
The LuxR-type receptors (LasR, RhlR) act as transcriptional regulators, and the LuxI-type proteins
(LasI, RhlI) are signal synthases. LasI produces 3-oxo-dodecanoyl-L-homoserine lactone (3OC12-
HSL), and RhlI produces butanoyl-L-homoserine lactone (C4-HSL) [5]. Both signals can function
in a combinatorial manner to regulate certain genes [8, 9]. Working in conjunction with the two
AHL systems is an alkyl-quinolone (AQ) system, comprising the *pqsABCDE* operon and *pqsH,*
pqsL and *pqsR* (*mvfR*) genes. These genes drive the synthesis and response of 2-heptyl-3-hydroxy-
4-quinolone (the Pseudomonas quinolone signal PQS), which is used as a QS signal and which
also has iron chelating properties [10, 11].
58
LasR was first identified in 1991 as a regulator of the *lasB* (elastase) gene [12]. It has since been
described as a key QS regulator in the well-studied laboratory strains PAO1 and PA14 [2]*,* where
it has been shown to sit at the top of the QS hierarchy, regulating both the *rhl* and *pqs* systems [5,
6]. *lasR* mutants have frequently been isolated from CF lungs [13-16] and more recently, some CF
strains use RhlR to regulate the *rhl* and *pqs* systems in the absence of a functional LasR [6, 16-
19]. The decoupling of the AHL QS hierarchy requires the inactivation of MexT, a regulator of
the multi-drug efflux pump operon MexEF-OprN [18, 20]. PqsE and RhlR have also been shown
to function as a ligand:receptor pair [21].
67
The ecological and evolutionary implications of QS re-wiring remain to be explored, and the
drivers for *lasR* mutation before and during infection are unknown. To date, little is known about
QS mutations outside of an infection environment, and so in this study, we explored the diversity
and frequency of QS mutations across a range of ecologically distinct environments in order to
determine (i) which QS genes are frequently mutated; (ii) mutational signatures, or patterns in QS
gene mutation specific to isolate source.
74
75 **Results**
76
We utilized the published sequences of 852 *P. aeruginosa* isolates from the International
Pseudomonas Consortium Database (IPCD); a database representing a range of *P. aeruginosa*
strains from different sources including rivers, infections and plants [22]. We queried a number of
key QS genes from the *las*, *rhl and pqs* systems against gene sequences from PAO1 using BLASTn

for all 852 isolates. We determined the putative amino acid sequence for each gene and calculated
dissimilarity scores using BLOSUM80, an empirical amino acid substitution matrix [13]. All
analyses were conducted in R version 4.3.
84
We first looked at the number of sequences we had for each QS protein, and the diversity of the
protein sequences. When we queried each QS gene nucleotide sequence against the 852 isolates,
the query returned less than 852 sequences for each gene. This disparity is likely due to gaps in
sequences, gene deletions, and extensive mutations, preventing BLASTn from returning a query.
Fig. 1A shows that a LasR query returned the fewest number of sequences, suggesting that there
are many strains that contain large deletions in LasR, truncations, or are missing the LasR gene
entirely. After translating the sequences, we found that LasR also had the most unique protein
sequences across 852 isolates (Fig. 1B). Given this finding, we analyzed whether there was a
mutational signature for the *las* system in order to determine whether certain kinds of mutation or
divergence were specific to *las* genes, and if these mutations were specific to isolate source. We
created PCA plots of LasR (Fig. 1C) and LasI (Fig. 1D) proteins from all returned isolates and
found that the LasR protein was distributed across the PCA plot, and the most divergent strains for
LasR were truncated. Compared to LasR, the other key QS proteins were more conserved across
isolates.
99
After analyzing all QS proteins, we then specifically focused on LasR. To determine if there were
LasR mutations specific to each environment, we categorized the strains by source. Using data
from the IPCD, we selected a subset of 654 strains labeled as "environmental", "cystic fibrosis" or
"CF", and "wound" or "ulcer" or "burn" and reclassified them as environmental (209 strains), CF
(396 strains), or wound (wound, ulcer and burn) (49 strains); 654 total. The remaining 198 strains
from the original set of 852 strains were of uncertain origin and therefore not used in this particular
analysis. To establish a threshold by which a protein could be deemed functional or not, we looked
at truncated LasR proteins within each environment. We compared the amino acid length of LasR
in the IPCD strains to the PAO1 LasR protein - which is equal in length to many commonly
researched strains including PA14, PAK and an epidemic CF strain, LESB58. Our assumption was
that a truncated protein due to shortened DNA sequence or an early stop site, would lead to a
nonfunctional protein. We used a stringent 100% length as a cut-off, and any protein shorter than
full-length was considered truncated. Fig. 2A shows the proportion of each group that had
truncated LasR proteins with CF, environmental, and wound isolates having 20%, 11% and 30%
truncations respectively. We also used a PCA plot to visualize LasR amino acid variation by
environment using BLOSUM80 generated dissimilarity scores (Fig. 2B). Overall, we found that
lasR mutations are ubiquitous across all environments, but there is a larger percentage of strains
with truncated LasR proteins found in infection environments.
118
119
120

Discussion

In *P. aeruginosa*, *lasR* QS mutants are frequently isolated from human chronic infection, but it has remained unclear whether such mutants specifically evolve in infection environments or are common across multiple environments. Using a publicly available database of 852 fully sequenced isolates from CF, wounds and non-infection based (environmental) isolates, we determined the frequency and pattern of *lasR* and other QS mutations in *P. aeruginosa*. We found that (i) LasR is the most variable protein of all the major QS proteins; (ii) *lasR* mutations are found in isolates across all environments, suggesting that any environment can drive the evolution of these mutations.

But what does drive the evolution of *lasR* mutations and what fitness benefits do *lasR* mutations provide to *P. aeruginosa* isolates or populations? First, *lasR* mutants could arise in populations through social cheating, where mutants exploit the social interactions and exoproducts produced by *lasR* intact cells [23, 24]. Controlled experiments have shown that *lasR* mutants can socially exploit wild type cells *in vitro* [24] and *in vivo* [25], although it is unclear whether the spatial structuring found within infections will allow the close proximity of different isolates to allow for regular cheating. Importantly, QS genes have recently been shown to be down-regulated during infection compared to *in vitro* conditions, questioning the long-held belief that a functional QS system is essential for *P. aeruginosa* to establish and persist in human infections [6, 26]. This would likely reduce any fitness benefits of being a *lasR* mutant persisting via social cheating. Second, *lasR* mutants may have increased fitness in particular environments due to certain phenotypes driven by the mutation being beneficial. For example, *lasR* mutations have previously been shown to confer a growth advantage with particular carbon and nitrogen sources, including amino acids [27]. Third, *lasR* mutants may be more competitive than *lasR* positive cells, which provides fitness benefits against other *P. aeruginosa* strains or other species.

There are, however, likely evolutionary benefits for both the maintenance and loss of LasR so that both *lasR* positive and negative strains can stably coexist in heterogenous populations and contribute to an overall community function. In recent support of this idea, it has been shown that (i) *lasR*- strains overproduce Rhl-associated factors and cross-feed wild type cells in low iron environments, which will likely impact infection dynamics of mixed populations [28]; (ii) mixed *lasR* +/- populations display decreased virulence in mouse models of infection [25]; (iii) mixed populations exhibit enhanced tolerance to beta-lactam antibiotics [29]. Taken together, this suggests there are likely to be considerable fitness advantages to cells growing in heterogeneous QS populations, perhaps as a bet-hedging mechanism for future disturbance events.

Overall, our work highlights that *lasR* mutations are the most commonly found QS mutation across different environments, although we do not know whether mutations in the *lasR* gene always results in a loss of QS function. Indeed, recent studies on QS in *P. aeruginosa* has revealed that

the complex and intertwined *las*, *rhl*, *and pqs* systems can be rewired in the event that *lasR*
becomes mutated [16, 18][9, 19]. It is not always clear whether these strains are entirely QS-null
or if they have re-wired their QS systems to circumvent the loss of *lasR*. Further work is needed
to determine why strains lose functional LasR proteins, and what fitness benefits the strains or
community gains. Future work should more strongly focus on the ecology of mixed QS-
phenotypes to better understand QS-involvement in infection and other environments. With
ongoing work identifying QS-inhibitors targeting the *las* QS system, the frequency of *lasR* mutated
strains found in our study suggests that this particular pursuit is likely to fail.
169
170 **Materials and Methods**
171
Querying QS genes from the International Pseudomonas Consortium Database. Using
nucleotide sequences from PAO1, we queried QS genes (see Fig. 1) using BLASTn for isolates
from the IPCD [12]. We chose this strain because it is a fully sequenced, frequently used lab strain.
We then translated these sequences into protein sequences calculating putative amino acid
sequence similarities using BLOSUM80 [13]. First, we compared genes found in each isolate
against our reference strain, PAO1, normalized against the similarity of the reference against itself.
We then calculated the mean dissimilarity score of all isolates compared to PAO1. Some isolates
were missing genes due to sequencing errors or true truncations, the number of isolates with a
given gene present was under 852 for all genes. All analyses, including translation steps were
conducted in R version 4.3. All code and files are available on Github
(https://github.gatech.edu/login?return_to=https%3A%2F%2Fgithub.gatech.edu%2Fkoconnor36
%2FFrequency_of_quorum_sensing_mutations_in_Pa2021).
184
185 **Creating an IPCD database using BLASTn.** We pulled IPCD data from GenBank. We used the
186 makeblastdb/ command to generate a database of all isolate contigs.
187
188 **Using BLASTn to find QS genes for each isolate.** Using our generated database, we queried the
189 PAO1 sequence from each gene, found from Genbank, against the database. We generated csv
190 files for each gene which included the gene sequences for each isolate.
191
192 **Translating nucleotide to amino acid sequence.** We translated genes to proteins using a custom
193 R script. We first queried only for sequences starting with a canonical ATG start codon. We
194 exclude sequences with fully unresolvable nucleotides (coded as "-"), but allowed fuzzy codons
195 so long as they resolved to unambiguous amino acids. We translated the sequences meeting these
196 criteria using the translate function from the BioStrings R package (v.2.58.0).
197
198 **Calculating dissimilarity scores for isolates' QS proteins.** All sequence analyses were
199 performed in R (v.4.0.2) using the Biostrings package v.2.58.0. We compared isolate protein
200 sequences to PAO1 protein sequences using BLOSUM80, a matrix designed to compared protein

201 sequences within species. We found that close to 50% of all isolate proteins were identical to
202 PAO1.
203
Determining truncation rates for LasR and categorizing isolates by location. We determined
the length of the reference LasR protein, from PAO1, compared to each isolate protein. If the
isolate protein was 100% or less of the length of the PAO1 protein, we categorized it as truncated.
Sequences were categorized as CF-originated (CF), environmental (ENV), or wound (WND). If
the sequence was entered into IPCD as environmental, we adopted that label. Additionally, we
included sequences labeled from animal hosts as environmental. For CF, we only included
sequences with sources explicitly labeled as CF. For wound, we included sequences labeled as
wound, ulcer, and burn.
212

213 **Author contributions.** SPD and KOC designed the study. KOC and CYZ performed the *in silico*
214 analysis of the data. All authors contributed to the writing of the manuscript.
215
216 **Competing interests.** The authors declare no competing interests.
217
218 **Funding and acknowledgements.** We wish to thank The Cystic Fibrosis Foundation
219 (DIGGLE18I0 and DIGGLE20G0) to SPD for funding. We also thank members of the Diggle Lab
220 and Jon Gerhart for helpful discussion.

221
222 **Figure legends**
223
Figure 1. Determining variability in QS proteins between *P. aeruginosa* isolates from the
IPCD. (A) We created a database of 852 isolates and used PAO1 to search for the QS proteins of
each isolate. Due to the variation in each isolate's genome and due to gaps in sequencing, each
protein queried returned fewer than 852 sequences (shown in gray). We also determined the
number of unique sequences for each protein and found that LasR had the highest number of
unique sequences (in color). (B) Using a custom dissimilarity metric (BLOSUM80), we calculated
mean dissimilarity scores. We found that LasR had the highest mean dissimilarity score compared
to all QS proteins, and the largest variation. We conducted principal component analyses (PCA)
and plotted the similarity scores of all isolates for the LasR protein (C) and the LasI protein (D).
233
Figure 2. LasR truncations are found in isolates from all sources. To observe the fraction of
truncated proteins across all environments, we categorized the isolates into 3 groups: cystic fibrosis
(CF), environmental (ENV) or wound (WND). (A) We show the number of truncated proteins out
of the total number of isolates in each group. The PCA plot (B) depicts the similarity scores for all
groups compared, and we see the arm of truncated proteins consisted primarily of CF and WND
isolates.
240
241

References

1.   **Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG et al.** Common virulence factors for bacterial pathogenicity in plants and animals. *Science* 1995;268(5219):1899-1902.

2.   **Diggle SP, Whiteley M**. Microbe Profile: *Pseudomonas aeruginosa*: opportunistic pathogen and lab rat. *Microbiology* 2020;166(1):30-33.

3.   **Elborn JS**. Cystic fibrosis. *The Lancet* 2016;388(10059):2519-2531.

4.   **Ibberson CB, Whiteley M**. The social life of microbes in chronic infection. *Curr Opin Microbiol* 2020;53:44-50.

5.   **Whiteley M, Diggle SP, Greenberg EP**. Progress in and promise of bacterial quorum sensing research. *Nature* 2017;551(7680):313-320.

6.   **Azimi S, Klementiev AD, Whiteley M, Diggle SP.** Bacterial Quorum Sensing During Infection. *Annu Rev Microbiol* 2020;74:201-219.

7.   **Schuster M, Sexton DJ, Diggle SP, Greenberg EP**. Acyl-homoserine lactone quorum sensing: from evolution to application. *Annu Rev Microbiol*, Review 2013;67(1):43-63.

8.   **Cornforth DM, Popat R, McNally L, Gurney J, Scott-Phillips TC et al.** Combinatorial quorum sensing allows bacteria to resolve their social and physical environment. *Proc Natl Acad Sci U S A* 2014;111(11):4280-4284.

9.   **Gurney J, Azimi S, Brown SP, Diggle SP**. Combinatorial quorum sensing in Pseudomonas aeruginosa allows for novel cheating strategies. *Microbiology (Reading)* 2020;166(8):777-784.

10.   **Dubern JF, Diggle SP**. Quorum sensing by 2-alkyl-4-quinolones in *Pseudomonas aeruginosa* and other bacterial species. *Mol Biosyst*, Review 2008;4(9):882-888.

11.   **Diggle SP, Matthijs S, Wright VJ, Fletcher MP, Chhabra SR et al.** The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS play multifunctional roles in quorum sensing and iron entrapment. *Chem Biol* 2007;14(1):87-96.

12.   **Gambello MJ, Iglewski BH**. Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. *J Bacteriol* 1991;173(9):3000-3009.

13.   **Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR et al.** Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. *Proc Natl Acad Sci U S A* 2006;103(22):8487-8492.

14.   **Ciofu O, Mandsberg LF, Bjarnsholt T, Wassermann T, Hoiby N**. Genetic adaptation of *Pseudomonas aeruginosa* during chronic lung infection of patients with cystic fibrosis: strong and weak mutators with heterogeneous genetic backgrounds emerge in *mucA* and/or *lasR* mutants. *Microbiology* 2010;156(Pt 4):1108-1119.

15.   **Jiricny N, Molin S, Foster K, Diggle SP, Scanlan PD et al.** Loss of social behaviours in populations of *Pseudomonas aeruginosa* infecting lungs of patients with cystic fibrosis. *PLoS One* 2014;9(1):e83124.

16.   **Feltner JB, Wolter DJ, Pope CE, Groleau MC, Smalley NE et al.** LasR Variant Cystic Fibrosis Isolates Reveal an Adaptable Quorum-Sensing Hierarchy in *Pseudomonas aeruginosa*. *mBio* 2016;7(5):e01513-01516-01519.

17.   **Diggle SP, Winzer K, Chhabra SR, Worrall KE, Camara M et al.** The *Pseudomonas aeruginosa* quinolone signal molecule overcomes the cell density-dependency of the quorum sensing hierarchy, regulates *rhl*-dependent genes at the onset of stationary phase and can be produced in the absence of LasR. *Mol Microbiol* 2003;50(1):29-43.

18. **Kostylev M, Kim DY, Smalley NE, Salukhe I, Greenberg EP et al.** Evolution of the
Pseudomonas aeruginosa quorum-sensing hierarchy. *Proc Natl Acad Sci U S A*
2019;116(14):7027-7032.
19. **Cruz RL, Asfahl KL, Van den Bossche S, Coenye T, Crabbe A et al.** RhlR-Regulated
Acyl-Homoserine Lactone Quorum Sensing in a Cystic Fibrosis Isolate of *Pseudomonas*
aeruginosa. mBio 2020;11(2).
20. **Oshri RD, Zrihen KS, Shner I, Omer Bendori S, Eldar A**. Selection for increased
quorum-sensing cooperation in *Pseudomonas aeruginosa* through the shut-down of a drug
resistance pump. *ISME J* 2018;12(10):2458-2469.
21. **Mukherjee S, Moustafa DA, Stergioula V, Smith CD, Goldberg JB et al.** The PqsE and
RhlR proteins are an autoinducer synthase-receptor pair that control virulence and biofilm
development in *Pseudomonas aeruginosa. Proc Natl Acad Sci U S A* 2018;115(40):E9411-E9418.
22. **Freschi L, Vincent AT, Jeukens J, Emond-Rheault JG, Kukavica-Ibrulj I et al.** The
Pseudomonas aeruginosa Pan-Genome Provides New Insights on Its Population Structure,
Horizontal Gene Transfer, and Pathogenicity. *Genome Biol Evol* 2019;11(1):109-120.
23. **West SA, Griffin AS, Gardner A, Diggle SP**. Social evolution theory for microorganisms.
Nat Rev Microbiol, Review 2006;4(8):597-607.
24. **Diggle SP, Griffin AS, Campbell GS, West SA**. Cooperation and conflict in quorum-
sensing bacterial populations. *Nature* 2007;450(7168):411-414.
25. **Rumbaugh KP, Diggle SP, Watters CM, Ross-Gillespie A, Griffin AS et al.** Quorum
sensing and the social evolution of bacterial virulence. *Curr Biol* 2009;19(4):341-345.
26. **Cornforth DM, Dees JL, Ibberson CB, Huse HK, Mathiesen IH et al.** *Pseudomonas*
aeruginosa transcriptome during human infection. *Proc Natl Acad Sci U S A* 2018;115(22):E5125-
E5134.
27. **D'Argenio DA, Wu M, Hoffman LR, Kulasekara HD, Deziel E et al.** Growth
phenotypes of *Pseudomonas aeruginosa lasR* mutants adapted to the airways of cystic fibrosis
patients. *Mol Microbiol* 2007;64(2):512-533.
28. **Mould DL, Botelho NJ, Hogan DA**. Intraspecies Signaling between Common Variants
of *Pseudomonas aeruginosa* Increases Production of Quorum-Sensing-Controlled Virulence
Factors. *mBio* 2020;11(4).
29. **Azimi S, Roberts AEL, Peng S, Weitz JS, McNally A et al.** Allelic polymorphism shapes
community function in evolving *Pseudomonas aeruginosa* populations. *ISME J* 2020;14(8):1929-
1942.

Figure 1

Figure 2