Frequency of quorum sensing mutations in *Pseudomonas aeruginosa* strains isolated from different environments

Kathleen O'Connor, Conan Y. Zhao, Madeline Mei & Stephen P. Diggle*

Center for Microbial Dynamics & Infection, School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30332.

*Correspondence: sdiggle3@gatech.edu

Keywords: *Pseudomonas aeruginosa*, quorum sensing, LasR, cystic fibrosis

Abstract

Pseudomonas aeruginosa uses quorum sensing (QS) to coordinate the expression of multiple genes necessary for establishing and maintaining infection. It has previously been shown that *lasR* QS mutations frequently arise in cystic fibrosis (CF) lung infections, however, there has been far less emphasis on determining whether other QS system mutations arise during infection or in other environments. To test this, we utilized 852 publicly available sequenced *P. aeruginosa* genomes from the *Pseudomonas* International Consortium Database (IPCD) to study *P. aeruginosa* QS mutational signatures. To study isolates by source, we focused on a subset of 654 isolates collected from CF, wounds, and non-infection environmental isolates, where we could clearly identify their source. We also worked a small collection of isolates *in vitro* to determine the impact of *lasR* and *pqs* mutations on isolate phenotypes. We found that *lasR* mutations are common across all environments and are not specific to infection nor a particular infection type. We also found that the *pqs* system proteins PqsA, PqsH, PqsL and MexT, a protein of increasing importance to the QS field, are highly variable. Conversely, RsaL, a negative transcriptional regulator of the *las* system, was found to be highly conserved, suggesting selective pressure to repress *las* system activity. Overall, our findings suggest that QS mutations in *P. aeruginosa* are common and not limited to the *las* system; however, LasR is unique in the frequency of putative loss-of-function mutations.

Introduction

Pseudomonas aeruginosa is a Gram-negative opportunistic pathogen that can be problematic in a number of infection types and environments [1]. One of the major adaptations of *P. aeruginosa* during chronic infection is the loss of quorum sensing (QS) [2–8]. In *P. aeruginosa,* QS regulates the expression of hundreds of genes, including those that encode for secreted products and virulence factors [9, 10]. It is regulated via a complex hierarchical network, composed of two *N*-

acyl homoserine lactone (AHL) circuits known as LasR-LasI and RhlR-RhlI, two orphan regulators termed QscR and VqsR, and a negative transcriptional regulator of the *las* system, RsaL [9–19]. The Las and Rhl systems are composed of LuxR-LuxI pairs, which are homologous to other Gram-negative bacterial QS systems. The LuxR-type receptors (LasR, RhlR) act as transcriptional regulators, and the LuxI-type proteins (LasI, RhlI) are signal synthases. LasI produces 3-oxo-dodecanoyl-L-homoserine lactone (3OC12-HSL), and RhlI produces N-butanoyl-L-homoserine lactone (C4-HSL). Both signals can function in a combinatorial manner to synergistically regulate genes [20, 21]. Working in conjunction with the two AHL systems, is an alkyl-quinolone (AQ) system, comprising the *pqsABCDE* operon and *pqsH, pqsL* and *pqsR* (*mvfR*) genes. These genes drive the synthesis and response of 2-heptyl-3-hydroxy-4-quinolone (the *Pseudomonas* quinolone signal; PQS) [22, 23].

LasR was first identified in 1991 as a regulator of the *lasB* (elastase) gene [24]. It has since been described as a key QS regulator in the well-studied laboratory strains PAO1 and PA14, where it has been shown to sit at the top of the QS hierarchy, regulating both the *rhl* and *pqs* systems [9, 11–15]. *lasR* mutants have frequently been isolated from cystic fibrosis (CF) lungs and more recently, it has been shown that some CF strains use RhlR to regulate the *rhl* and *pqs* systems in the absence of functional LasR [25–30]. The decoupling of the AHL QS hierarchy reportedly requires the inactivation of MexT [25, 27, 31], a regulator of the multi-drug efflux pump operon MexEF-OprN [32, 33]. It has also been shown that *mexT* mutation in PAO1 can decouple public *rhlR*-regulated traits from private metabolic *lasR*-regulated traits [20]. PqsE and RhlR have also been suggested to function as a ligand:receptor pair in some QS 're-wired' strains [34–36].

Previous work has shown that *lasR* mutations are also found outside the CF lung, but the extent and variation of these mutations is still being elucidated [26, 30]. The degree of mutation in other QS genes in infection and environmental strains of *P. aeruginosa* remains unknown. In this study, we explored the diversity and frequency of QS mutations across a range of ecologically distinct environments to determine (i) which QS genes are frequently mutated; (ii) mutational signatures, or patterns in other genes associated with *lasR* mutations; (iii) *lasR* gene mutation frequency specific to isolate source; and (iv) the phenotypic outcome of QS mutations.

Results

We utilized the published sequences of 852 *P. aeruginosa* isolates from the International *Pseudomonas* Consortium Database (IPCD); a database representing a range of *P. aeruginosa* strains from different sources including rivers, human infection and plants [37]. We queried key QS genes from the *las*, *rhl and pqs* systems, as well as the QS regulators *qscR, rsaL* and *vqsR* against gene sequences from PAO1 (GCF_000006765.1) for all 852 isolates. For some analyses, we additionally looked at *mexT* and *psdR,* two genes that have been associated with *lasR* mutations *in vitro* [25, 31, 38, 39]. We used two analysis pipelines to look at the variation in QS genes in *P.*

aeruginosa. One pipeline used NCBI BLASTn [40] and BLOSSUM80 [41], and the second analysis utilized GAMMA [42]. All analyses were conducted in R version 4.3 [43].

Quorum sensing genes are highly variable at the nucleotide level. We first analyzed the diversity of nucleotide sequence for each QS gene for all 852 isolates and looked at which isolates had mutations in multiple QS-related genes. We found a large diversity in sequences and that few isolates had QS genes identical to PAO1 (GCF_000006765.1). The *rhl* system was especially variable, with 849 and 847 isolates differing from PAO1 for *rhlR* and *rhlI* respectively (Fig. 1A). The *las* system was more conserved, with 553 and 290 isolates varying from PAO1 for *lasR* and *lasI* respectively (Fig. 1A). Genes involved in 2-alkyl-4-quinolone (AQ) biosynthesis and response showed a large variation in mutations between genes, with *pqsE* as the least variable (479 mutated isolates), and *pqsH* and *pqsL* as the most variable (845, 847 respectively) (Fig. 1C). The orphan QS transcriptional regulators *qscR* and *vqsR* were highly variable (772 and 559 respectively). However, the negative *las* system transcriptional regulator *rsaL* was relatively conserved (218 mutated isolates) (Fig. 1D).

Negative and positive regulators of the *las* system are antipodal in sequence variation. We counted the number of unique protein sequences found in our database for each QS gene. We added *mucA* to our analysis as a reference gene that is frequently mutated in CF lung infections [44, 45], and *rpsL,* which encodes the 30S ribosomal protein S12; a conserved housekeeping gene [46, 47]. When we queried each QS gene nucleotide sequence against the 852 isolates using BLASTn, the query returned less than 852 sequences for each gene. This disparity is likely due to gaps in sequences, gene deletions, and extensive mutations, preventing BLASTn from returning a query. Additionally, for this analysis, we removed all truncated nucleotide sequences from the analysis, to avoid including sequencing near the end of a contig. Fig. 2A shows the number of protein sequences analyzed for each gene. After MucA, a LasR query returned the fewest number of sequences – 756 out of 852 isolates, suggesting that there are many strains that contain large deletions in LasR, truncations, or are lacking the LasR gene entirely. After translating the sequences, we found that LasR had the most unique protein sequences (259) across 852 isolates (Table 1, Fig. 2A) compared to other QS genes and MucA. The next most variable QS proteins, PqsH, PqsA, and PqsL contained 189, 170, and 169 unique protein sequences, respectively (Table 1; Fig. 2A). We found that AHL signal synthases were highly conserved; LasI and RhlI had only 61 and 87 unique protein sequences respectively. RsaL was the most conserved of all studied proteins, with only 18 unique sequences. Compared to LasR, the other key QS proteins were more conserved across isolates. Our protein sequence findings appear to contrast with our nucleotide sequence analysis, but this is likely due to primarily silent mutations in the *rhl* system. To control for possible bias due to differences in protein length, we normalized gene sizes to PAO1 by dividing the lengths of each gene by the PAO1 reference gene length. We found that the shortest proteins, RspL and RsaL were the most conserved, while the longest genes, PqsH, PqsA, and PqsL were all highly variable. The most variable protein, LasR is of intermediate length, compared to RsaL and Pqs proteins (Table 1).

We next plotted the BLOSUM80 dissimilarity scores for all isolates for each gene and found that LasR had a large number of highly dissimilar sequences (Fig. 2B). The highly dissimilar scores are caused by truncation mutations due to an early stop site leading to a shortened protein. Transcriptional regulators PqsR and RhlR also have a number of truncation mutations, however, there are fewer than LasR. While PqsA, PqsL and PqsH all encode many unique protein sequences, most isolates cluster towards the bottom, indicating that they have dissimilarity scores close to 0, indicating that the proteins have a near identical function.

MexT, PqsL and LasR have the highest mutation rates. We used the protein sequence data from GAMMA to look at the mutation rate of each QS protein (Table 2). For this analysis, we were able to include truncated nucleotide sequences, and only excluded sequences where it was indicated that the gene was located at the contig edge. We included MexT and PsdR proteins in this analysis. For the majority of genes, we compared isolates to the PAO1 (GCF_000006765.1) gene sequence, with the exception of PqsD and RhlI which we compared to the PAK strain of *P. aeruginosa*. The MexT sequence for PAO1 has an 8 bp insertion sequence, which was not commonly found in the IPCD database of isolates, as well as many other coding mutations. We instead used the sequence for isolate U0330A as it coded for a protein shared by 128 other isolates, the highest number of strains sharing one MexT sequence. The MexT gene is highly variable, and is mutated in our Nottingham PAO1 lab strain (NPAO1) and the PAO1 reference used (GCF_000006765.1). Among the QS genes, we found that PqsL had the highest mutation rate, followed by LasR. We defined mutation rate in this analysis as the percentage of non-wildtype protein sequences out of queried isolates. RsaL and LasI were highly conserved, with low mutation rates.

Using GAMMA generated protein mutation data, we next calculated the average codon similarity of all isolates to a reference (Table 2). High codon similarity suggests similar protein function, while low codon similarity indicates a change or loss of protein function. For the majority of genes we used PAO1 (GCF_000006765.1) as a reference, however the codon similarity average for several proteins were low. To address this, for RhlI and PqsD we used PAK as the reference strain, and for MexT we used U0330A as these strains were likely to be better models based on higher similarity averages for each protein. Similar to our analysis using BLOSUM80, we found that LasR had the lowest average protein similarity of all QS genes, followed by MexT, which we attributed to the high rate of truncation mutations. All other QS proteins had a codon similarity average of greater than 0.99, indicating that while there is variation in the protein sequence, many of these mutations likely lead to a similarly functioning protein.

LasR protein mutations are common across all environments but are more divergent in human wounds. In agreement with previous studies, LasR was the most variable gene in our protein analysis (Table 1; Fig. 2), and we found that the highly dissimilar LasR sequences were due to protein truncations [2, 30, 48]. To determine if LasR mutations vary by environment, we categorized the strains by source. Using data from the IPCD, we selected a subset of 654 strains labeled as "environmental", "cystic fibrosis" or "CF", and "wound" or "ulcer" or "burn" and reclassified them as environmental (209 strains), CF (396 strains), or wound (wound, ulcer, and burn) (49 strains); 654 total. The remaining 198 strains from the original set of 852 strains were of uncertain origin and therefore excluded from this analysis. To establish a threshold by which a

protein could be deemed functional or not, we looked at truncated LasR proteins within each environment. We compared the amino acid length of LasR in the IPCD strains to the PAO1 LasR protein – which is equal in length to many commonly researched strains including PA14, PAK and the Liverpool epidemic CF strain LESB58. Our assumption was that a truncated protein due to an early stop site, would lead to a nonfunctional protein. We used a stringent 100% length as a cut-off, and any protein shorter than full-length was considered truncated. Fig. 3A shows the proportion of each group that had truncated LasR proteins with CF, environmental, and wound isolates having 20%, 11% and 30% truncations respectively. We then plotted the protein sequence dissimilarities categorized by isolate source, and found highly dissimilar isolates, primarily in CF and wounds (Fig. 3B). Overall, we found that *lasR* mutations are ubiquitous across all environments, but there is a larger percentage of strains with truncated LasR proteins found in infection environments. We additionally calculated the mean and median codon similarity scores for each environment, with 1 being a protein identical to PAO1. We found that the wound group was the most dissimilar compared to PAO1, with the lowest mean similarity score (0.893), and CF and environmental were similar (CF: 0.953, Env: 0.948). The wound group also had the lowest median similarity score (0.995), followed by CF (0.996) and environmental had the highest median similarity score (1.0).

***lasR* mutation is not a predictor of exoprotease production or colony autolysis in selected environmental and infection isolates**. To accompany our genomic analysis, we selected 12 isolates from the IPCD in order to perform phenotypic assays to assess QS function. We included our lab strain NPAO1 as well as the clean deletion mutants PAO1Δ*lasR*, PAO1Δ*rhlR*, PAO1Δ*lasR*Δ*rhlR*, PAO1Δ*lasI*. Our PAO1Δ*lasR* strain made high levels of protease that were not significantly lower than NPAO1 (89 % of NPAO1 levels, $p = 0.338$). A *lasR* gentamicin insertion mutant made 75 % of the protease levels of NPAO1 ($p = 0.113$) (Fig. 4). We found that unlike mutations in LasR, mutations in RhlR had a significant effect on the protease levels produced compared to NPAO1 (PAO1Δ*rhlR*: 45 %, $p = 0.013$; PAO1Δ*lasR*Δ*rhlR* double mutant: 20 %, $p = 0.011$). Compared with both Δ*lasR* mutants, NPAO1Δ*lasI* made lower amounts of protease (60 %, $p = 0.025$) (Fig. 4). Isolates with mutated LasR proteins that produce significantly less protease than wildtype included A17, CND03, CPHL2000, Jp115 (p values respectively: 0.01, 0.01, 0.02, 0.01). PT31M, a German environmental isolate contains a truncation mutation in LasR, yet it produced an intermediate amount of protease compared to NPAO1 with an intact LasR (69 %, $p = 0.062$). The Belgian river environmental strain W15Dec14 also produced an intermediate amount of protease (67 %, $p = 0.065$) and is wildtype for LasR, but does contain coding mutations in RhlI. All other isolates with wildtype LasR proteins produced protease levels that were not significantly different from NPAO1 (Fig. 4).

Lysis in QS mutants has long been observed in the lab, specifically in *lasR*, *lasI* and/or *pqs* mutants [48–51]. We used QS deletion mutant strains to determine how QS mediates colony autolysis. PAO1Δ*lasR* and PAO1Δ*pqsL* showed similar lysis phenotypes and a metallic sheen, while PAO1Δ*pqsH* caused colony plaquing (Fig. 5). PAO1Δ*pqsA*Δ*lasR,* PAO1Δ*pqsA*Δ*pqsH* and PAO1Δ*pqsA*Δ*pqsL* mutations led to a loss of colony autolysis (Fig. 5). We found that all LasR wildtype strains we tested from the IPCD showing lysis, contained multiple mutations in the *pqs* quinolone system, as follows: SWPA15J=NSWPA15a (PqsC,D,E), 5BR2 (PqsD,L), Co398373

(PqsD,L), Jp238 (PqsB,D,L), W15D14 (PqsE,D,L), Aa249 (PqsB,D,L), JD303 (PqsA,B,D). Strains with wildtype LasR showing lysis were from mixed environmental, wound and CF sources. There were two *lasR* mutant isolates that showed no lysis - A17 and CPHL2000, both sourced from human wounds (Fig. 5). Isolates A17 and CPHL2000 have 1 and 2 SNPs in LasR respectively, both creating a change in amino acid sequence (A17: L110P, CPHL2000: E196G). As expected, the 3 isolates with truncated LasR (CND03, Jp1155, and PT31M) demonstrated lysis.

Discussion

In *P. aeruginosa*, *lasR* QS mutants are frequently isolated from human chronic infection and environmental sources, but less is known about other QS genes [2, 48, 52]. Using a publicly available database of 852 fully sequenced isolates from CF, wounds and non-human associated (environmental) strains [37], we determined the mutation frequency and variation of *lasR* and other QS genes in *P. aeruginosa*. To determine how QS genotype impacts phenotypic function, we used 12 strains that were sequenced as part of the IPCD study [37]. We found that (i) by multiple metrics, LasR is the most variable QS protein; (ii) LasR mutations are found in isolates across all environments, suggesting that any environment can drive the evolution of these mutations; (iii) the negative *las* system regulator RsaL is well-conserved; (iv) signal synthases LasI and RhlI are conserved compared to their transcriptional regulator pairs; (v) coding mutations in *pqs* system genes are common and may impact colony autolysis, however they are less divergent than LasR mutations; (vi) MexT is highly variable and mutations are common.

Our work supports the long-held belief that LasR is a commonly mutated QS protein as we showed that LasR has the highest number of unique protein sequences compared to other QS genes. A high number of isolates carried mutations in LasR compared to the PAO1 reference protein sequence. The IPCD strains also had the lowest average codon similarity for the LasR gene when compared to other QS regulon genes, which we attributed to nonsense mutations leading to protein truncation, and other large changes in amino acid sequence. Our results also highlight that LasR mutations are found in environmental isolates and are not specific to human infections. Truncation mutations are more commonly found in human wound infections, suggesting that there might be some selective pressure to evolve loss of function mutations. Mean and median similarity scores suggest there is not a large difference between CF and environmental isolates, and instead we found that wound isolates are enriched for loss of function mutations. The wound group was the smallest group of isolates we queried thus a larger exploration of wound strains could provide a deeper understanding of this phenomenon.

Interestingly, the most highly conserved QS protein was RsaL, which negatively regulates the *las* QS system [16, 53, 54]. RsaL and LasR are antipodal in variation across isolates and perform opposite jobs in regulating the *las* system. There may be a benefit to tightly conserving the "off" switch of the *las* system while frequently mutating and losing the function of the "on" switch, suggesting that a key element of the QS regulatory cascade is down-regulating *las*-controlled QS in *P. aeruginosa*. Given this complexity, it remains a challenge to understand why the *las* system evolved and what fitness benefits it provides *P. aeruginosa* in different environments. There may be evolutionary benefits for both the maintenance and loss of LasR so that both *lasR* positive and negative strains can stably coexist in heterogenous populations and contribute to an overall community function. In support of this idea, it has been shown that (i) *lasR*- strains overproduce

Rhl-associated factors and cross-feed wild type cells in low iron environments, which will likely impact infection dynamics of mixed populations [55]; (ii) mixed *lasR +/-* populations display decreased virulence in mouse models of infection [56]; and (iii) mixed populations exhibit enhanced tolerance to beta-lactam antibiotics [57]. These data suggest there are likely considerable fitness advantages to cells growing in heterogeneous QS populations, perhaps as a bet-hedging mechanism for future disturbance events.

The signal synthases of the two AHL QS systems in *P. aeruginosa* are conserved compared to their transcriptional regulator pairs. It has been hypothesized that there could be evolutionary mechanisms to conserve signal synthases while mutating transcriptional regulators [58]. We saw a significant difference in protease production by the clean deletion of the regulator mutant LasR and the signal synthase mutant LasI, where deletion of LasI resulted in a larger decrease in protease production. The reason for this could be further explored and may be impacted by the MexT mutation in NPAO1, common to other PAO1 lab strains, or the fact that the 3' end of RsaL is also deleted in a clean LasR deletion mutation [59]. There could also be an evolutionary benefit to maintaining some protease function in the absence of fully functioning QS, supporting the maintenance of signal synthases while mutating positive transcriptional regulators.

Recent studies on QS in *P. aeruginosa* has revealed that the complex and intertwined *las*, *rhl* and *pqs* systems can be re-wired if *lasR* becomes mutated [25, 27, 31, 34–36]. It is not always clear whether these strains are entirely QS-null or if they have re-wired their QS systems to circumvent the loss of *lasR*. In our *in vitro* work, we looked at lysis and protease secretion as indicators of QS function, which have previously been used to screen for LasR function [48, 60]. We found an environmental isolate, PT31M, with a truncated *lasR* that produces high levels of protease. PT31M may be an example of a re-wired isolate showing QS independent of LasR, suggesting this adaptation is not specific to CF lung infections. Colony lysis has been used as a predictor for *las* system function, however this study has shown little correlation between lysis and *lasR* genotype, potentially due to *pqs* mutation mediated lysis seen in NPAO1Δ*pqsH* and NPAO1Δ*pqsL*.

Overall, our work shows that LasR is uniquely mutated compared to other QS genes, future work should more strongly focus on the ecology of mixed QS-phenotypes to better understand QS-involvement in infection and other environments. With ongoing work identifying QS-inhibitors targeting the *las* QS system, the frequency of *lasR* mutated strains found in our study suggests that this particular pursuit might be improved by targeting a less variable QS gene, such as *rhl* system genes or RsaL.

Materials and Methods

Querying QS genes from the International *Pseudomonas* Consortium Database. Using nucleotide sequences from PAO1 (GCF_000006765.1), we queried QS genes using BLASTn for isolates from the IPCD [37]. We chose this strain because it is a fully sequenced, frequently used lab strain. We first compared strains with nucleotide mutations relative to PAO1 in each of the QS genes of interest, to determine how frequently strains exhibit nucleotide polymorphisms across multiple QS genes. We then translated these sequences into protein sequences calculating putative amino acid sequence similarities using BLOSUM80 [41]. First, we compared genes found in each

isolate against our reference strain, PAO1, normalized against the similarity of the reference against itself. We then calculated the mean dissimilarity score of all isolates compared to PAO1. Some isolates were missing genes due to sequencing errors or true truncations, the number of isolates with a given gene present was under 852 for all genes. All analyses, including translation steps were conducted in R version 4.3. All code and files are available on a publicly available database:

https://github.gatech.edu/koconnor36/Frequency_of_quorum_sensing_mutations_in_Pa2021

Creating an IPCD database using BLASTn. We pulled IPCD data from GenBank from the PRJNA325248 BioProject (https://www.ncbi.nlm.nih.gov/bioproject/325248), and downloaded contigs as a multifasta file. We used the makeblastdb/ command to generate a database of all isolate contigs.

Using BLASTn to find QS genes for each isolate. Using our generated database, we queried the PAO1 sequence for each gene found on Pseudomonas.com [61], against the database. We generated csv files for each gene which included the gene sequences for each isolate.

Comparing strains with nucleotide polymorphisms in multiple QS genes. We used the csv files for each gene generated from the BLASTn analysis to isolate all accession information of strains with <100% query coverage and <100% identity to the PAO1 sequence of each QS gene. For each QS gene, we used this list of NCBI accession IDs, corresponding to unique strains, to visualize the number of strains with nucleotide mutations in one or more genes with InteractiVenn [62]. We then calculated the number of strains and the percentage of strains in the IPCD database with at least one mutation in each QS gene. We generated accession lists of mutated strains for each QS gene using a custom R script (v.4.0.2) and generated Fig. 1 with InteractiVenn. All analyses were conducted in R version 4.3.

Translating nucleotide to amino acid sequence. We translated genes to proteins using a custom R script. We first queried only for sequences starting with a canonical ATG start codon. VqsR is an exception as it begins with the alternative start codon "GTG". We translated the sequences meeting these criteria using the translate function from the BioStrings R package (v.2.58.0) [63].

Calculating dissimilarity scores for isolates' QS proteins. All sequence analyses were performed in R (v.4.0.2) using the Biostrings package v.2.58.0. We compared isolate protein sequences to PAO1 protein sequences using BLOSUM80, a matrix designed to compare protein sequences within species.

Determining truncation rates for LasR and categorizing isolates by location. We determined the length of the reference LasR protein, from PAO1, compared to each isolate protein. If the isolate protein was 100% or less of the length of the PAO1 protein, we categorized it as truncated. Sequences were categorized as CF-originated (CF), environmental (ENV), or wound (WND). If the sequence was entered into IPCD as environmental, we adopted that label. Additionally, we included sequences labeled from animal hosts as environmental. For CF, we only included sequences with sources explicitly labeled as CF or cystic fibrosis. For wound, we included sequences labeled as wound, ulcer, and burn.

Protein mutation analysis using GAMMA: We used GAMMA [42] to find the mutations in QS genes for all isolates. We created a multifasta of genes: *lasR,I, rhlR,I, pqsA,B,C,D,E,H,L,R, qscR, rsaL, vqsR, mucA, rpsL, mexT, psdR* to query against the IPCD PRJNA325248 multifasta. We ran commands as described in the GAMMA GitHub [42] https://github.com/rastanton/GAMMA, and generated a GAMMA file with mutations reported for all genes in all IPCD isolates. All other sequences are available on our project Github https://github.gatech.edu/koconnor36/Frequency_of_quorum_sensing_mutations_in_Pa2021.

Phenotypic assays. Isolates sequenced as part of the IPCD were donated by the Brown lab at Georgia Tech. We cultured isolates on LB agar plates, and confirmed their identity by sequencing the *lasR* gene, using colony *OneTaq* PCR (New England Biolabs). For protease activity, we inoculated colonies in 5 mL of LB broth, and grew strains in biological triplicates for 16-18 hours at 37 °C, 200 rpm. We pelleted cells and used the supernatant to run the Pierce Fluorescent Protease Assay Kit (Thermo). Readings were quantified using the TPCK Trypsin standard in triplicate, and normalized by strain optical density (OD at 600_{nm}). All readings were compared to NPAO1. For colony autolysis experiments, we inoculated colonies in 5 mL of LB broth, and grew strains for 16-18 hours at 37 °C, 200 rpm. We tested colony autolysis by spotting 5 μL of overnight cultures on LB agar plates, and growing colonies for 24 hours at 37 °C.

Author contributions. SPD and KO designed the study. KO, CYZ, and MM performed the *in silico* analysis of the data. KO performed the *in vitro* analysis. All authors contributed to the writing of the manuscript.

Competing interests. The authors declare no competing interests.

Funding and acknowledgements. We wish to thank The Cystic Fibrosis Foundation (DIGGLE20G0) and The National Institute for Health (R01AI153116) to SPD for funding. We also thank members of the Diggle Lab and Jon Gerhart for helpful discussion. We thank the Brown Lab at Georgia Tech for donating strains from the IPCD collection.

Figure legends

Table 1. Variation in protein sequence encoded by *P. aeruginosa* isolates. Using the sequences returned by BLASTn, we translated genes with full length nucleotide sequences to determine the number of unique protein sequences encoded by the isolates in the database. We found that LasR had the most unique sequences across queried isolates, and RsaL encoded the fewest unique sequences.

Table 2. Mutation rate and mean codon similarity of QS proteins in *P. aeruginosa*. We used GAMMA to query QS genes against 852 isolates. We used PAK as a reference for RhlI and PqsD, U0330A as a reference for MexT and PAO1 as a reference for all other genes. GAMMA assigns isolates with a codon similarity measure which reflects the type of amino acid change or truncation. A low score reflects a likely change in function of the protein, and a score of 1 reflects the isolate is identical to the reference. We found that LasR had the lowest mean codon similarity and RsaL the highest.

Figure 1. Comparison of nucleotide sequence variation among QS genes of *P. aeruginosa* isolates of the IPCD. There is a high degree of variation in nucleotide sequences in many of the QS genes, obscuring patterns in mutation overlap between genes within an isolate. (A) A Venn diagram of *las* and *rhl* system genes are plotted together, showing how many isolates share mutations in multiple genes. 246 isolates are mutated for all 4 genes, and few isolates have mutations in only a single gene. (B) A Venn diagram for *pqsABCD* shows a similar pattern and 683 isolates are mutated in all 4 genes, while few isolates are mutated in only one *pqs* gene. (C) Few isolates have only one *pqs* gene mutation, and 430 isolates have mutations in all *pqsEHLR* genes. (D) A Venn diagram of the negative QS regulators *vqsR, rsaL, and qscR* show that 132 isolates share mutations in all 3 genes.

Figure 2. Determining variability in QS proteins between *P. aeruginosa* isolates from the IPCD. (A) We created a database of 852 isolates and used PAO1 to search for the QS proteins of each isolate. Due to the variation in each isolate's genome and due to gaps in sequencing, each protein queried returned fewer than 852 sequences (shown in gray). We also determined the number of unique sequences for each protein and found that LasR had the highest number of unique sequences (in color). The next highest were PqsH, PqsA, and PqsL. RsaL had the fewest number of unique sequences. (B) Using a custom dissimilarity metric (BLOSUM80), we calculated mean dissimilarity scores. We found that LasR had the highest mean dissimilarity score compared to all QS proteins, and the largest variation.

Figure 3. LasR truncations are found in isolates from all sources. To observe the fraction of truncated proteins across all environments, we categorized the isolates into 3 groups: cystic fibrosis (CF), environmental (ENV) or wound (WND). (A) We show the number of truncated proteins out of the total number of isolates in each group. The protein variation plot (B) depicts the similarity scores for all groups compared, and we see the highly dissimilar truncated proteins consisted primarily of CF and WND isolates.

Figure 4. Exoprotease secretion by environmental and clinical isolates. We measured the supernatant exoprotease levels produced by stationary phase isolates, compared to PAO1, PAO1$\Delta lasR$ and PAO1$\Delta lasR\Delta rhlR$. We normalized the readings by optical density (OD 600_{nm}) and compared all isolates and clean mutants to PAO1.

Figure 5. Colony autolysis phenotypes of isolates compared to lab strain QS mutants. We spotted overnight cultures on LBA plates to observe the colony autolysis phenotypes exhibited by clean QS deletion mutants alongside environmental and clinical isolates.

References

1.   **Diggle SP, Whiteley M**. Microbe profile: *Pseudomonas aeruginosa*: Opportunistic pathogen and lab rat. *Microbiology (United Kingdom)* 2020;166:30–33.
2.   **Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, *et al.*** Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. *Proc Natl Acad Sci U S A* 2006;103:8487–8492.

3.   **LaFayette SL, Houle D, Beaudoin T, Wojewodka G, Radzioch D, *et al.*** Cystic fibrosis–adapted *Pseudomonas aeruginosa* quorum sensing *lasR* mutants cause hyperinflammatory responses. *Science Advances* 2015;1:1–15.

4.   **Hansen SK, Rau MH, Johansen HK, Ciofu O, Jelsbak L, *et al.*** Evolution and diversification of *Pseudomonas aeruginosa* in the paranasal sinuses of cystic fibrosis children have implications for chronic lung infection. *ISME Journal* 2012;6:31–45.

5.   **Ciofu O, Mandsberg LF, Bjarnsholt T, Wassermann T, Høiby N**. Genetic adaptation of *Pseudomonas aeruginosa* during chronic lung infection of patients with cystic fibrosis: Strong and weak mutators with heterogeneous genetic backgrounds emerge in *mucA* and/or *lasR* mutants. *Microbiology (N Y)* 2010;156:1108–1119.

6.   **Williams P, Cámara M**. Quorum sensing and environmental adaptation in *Pseudomonas aeruginosa*: a tale of regulatory networks and multifunctional signal molecules. *Current Opinion in Microbiology* 2009;12:182–191.

7.   **Galán-Vásquez E, Luna B, Martínez-Antonio A**. The Regulatory Network of *Pseudomonas aeruginosa*. *Microbial Informatics and Experimentation* 2011;1:3.

8.   **Lee J, Zhang L**. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. *Protein and Cell* 2014;6:26–41.

9.   **Whiteley M, Diggle SP, Greenberg EP, Wilson EO**. Bacterial quorum sensing: the progress and promise of an emerging research area. *Nature* 2017;551:313–320.

10.   **Azimi S, Klementiev AD, Whiteley M, Diggle SP**. Bacterial Quorum Sensing during Infection. *Annual Review of Microbiology* 2020;74:201–219.

11.   **Wagner VE, Bushnell D, Passador L, Brooks AI, Iglewski BH**. Microarray analysis of *Pseudomonas aeruginosa* quorum-sensing regulons: Effects of growth phase and environment. *Journal of Bacteriology* 2003;185:2080–2095.

12.   **Schuster M, Lostroh CP, Ogi T, Greenberg EP**. Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: A transcriptome analysis. *Journal of Bacteriology* 2003;185:2066–2079.

13.   **Fuqua WC, Winans SC, Greenberg EP**. Quorum sensing in bacteria: The LuxR-LuxI family of cell density- responsive transcriptional regulators. *Journal of Bacteriology* 1994;176:269–275.

14.   **Ding F, Oinuma KI, Smalley NE, Schaefer AL, Hamwy O, *et al.*** The *Pseudomonas aeruginosa* Orphan Quorum Sensing Signal Receptor QscR Regulates Global Quorum Sensing Gene Expression by Activating a Single Linked Operon. *mBio*;9. Epub ahead of print 2018. DOI: https://doi .org/10.1128/mBio.01274-18.

15.   **Schuster M, Greenberg EP**. A network of networks: Quorum-sensing gene regulation in *Pseudomonas aeruginosa*. *International Journal of Medical Microbiology* 2006;296:73–81.

16.   **Rampioni G, Bertani I, Zennaro E, Polticelli F, Venturi V, *et al.*** The quorum-sensing negative regulator RsaL of *Pseudomonas aeruginosa* binds to the *lasI* promoter. *Journal of Bacteriology* 2006;188:815–819.

17.   **Rampioni G, Schuster M, Greenberg EP, Zennaro E, Leoni L**. Contribution of the RsaL global regulator to *Pseudomonas aeruginosa* virulence and biofilm formation. *FEMS Microbiology Letters* 2009;301:210–217.

18.   **Liang H, Deng X, Ji Q, Sun F, Shen T, *et al.*** The *Pseudomonas aeruginosa* global regulator VqsR directly inhibits QscR To control quorum-sensing and virulence gene expression. *Journal of Bacteriology* 2012;194:3098–3108.

19. **Juhas M, Wiehlmann L, Huber B, Jordan D, Lauber J,** *et al.* Global regulation of quorum sensing and virulence by VqsR in *Pseudomonas aeruginosa*. *Microbiology (N Y)* 2004;150:831–841.

20. **Gurney J, Azimi S, Brown SP, Diggle SP**. Combinatorial quorum sensing in *Pseudomonas aeruginosa* allows for novel cheating strategies. *Microbiology (United Kingdom)* 2020;166:777–784.

21. **Cornforth DM, Popat R, McNally L, Gurney J, Scott-Phillips TC,** *et al.* Combinatorial quorum sensing allows bacteria to resolve their social and physical environment. *Proc Natl Acad Sci U S A* 2014;111:4280–4284.

22. **Dubern JF, Diggle SP**. Quorum sensing by 2-alkyl-4-quinolones in *Pseudomonas aeruginosa* and other bacterial species. *Molecular BioSystems* 2008;4:882–888.

23. **Heeb S, Fletcher MP, Chhabra SR, Diggle SP, Williams P,** *et al.* Quinolones: From antibiotics to autoinducers. *FEMS Microbiology Reviews* 2011;35:247–274.

24. **Gambello MJ, Iglewski BH**. Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. *Journal of Bacteriology* 1991;173:3000–3009.

25. **Kostylev M, Kim DY, Smalley NE, Salukhe I, Peter Greenberg E,** *et al.* Evolution of the *Pseudomonas aeruginosa* quorum-sensing hierarchy. *Proc Natl Acad Sci U S A* 2019;116:7027–7032.

26. **Feltner JB, Wolter DJ, Pope CE, Groleau M, Smalley NE,** *et al.* LasR Variant Cystic Fibrosis Isolates Reveal an Adaptable Quorum-. *mBio* 2016;7:e01513-16.

27. **Cruz RL, Asfahl KL, Bossche S van den, Coenye T, Crabbé A,** *et al.* RhlR-Regulated Acyl-Homoserine Lactone Quorum Sensing in a Cystic Fibrosis Isolate of *Pseudomonas aeruginosa*. *mBio*;11. https://doi.org/10.1128/mBio.00532-20 (2020).

28. **Thomason MK, Voichek M, Dar D, Addis V, Fitzgerald D,** *et al.* A rhli 5= UTR-derived sRNA regulates RhlR-dependent quorum sensing in *Pseudomonas aeruginosa*. *mBio*;10. Epub ahead of print 2019. DOI: 10.1128/mBio.02253-19.

29. **Chen R, Déziel E, Groleau MC, Schaefer AL, Greenberg EP**. Social cheating in a *Pseudomonas aeruginosa* quorum-sensing variant. *Proc Natl Acad Sci U S A* 2019;116:7021–7026.

30. **Groleau M, Taillefer H, Vincent AT, Constant P, Déziel E**. *Pseudomonas aeruginosa* isolates defective in function of the LasR quorum sensing regulator are frequent in diverse environmental niches . *Environmental Microbiology* 2021;00:1–14.

31. **Oshri RD, Zrihen KS, Shner I, Omer Bendori S, Eldar A**. Selection for increased quorum-sensing cooperation in *Pseudomonas aeruginosa* through the shut-down of a drug resistance pump. *ISME Journal* 2018;12:2458–2469.

32. **Schweizer HP**. Efflux as a mechanism of resistance to antimicrobials in *Pseudomonas aeruginosa* and related bacteria: Unanswered questions. *Genetics and Molecular Research* 2003;2:48–62.

33. **Köhler T, Epp SF, Curty LK, Pechère JC**. Characterization of MexT, the regulator of the MexE-MexF-OprN multidrug efflux system of *Pseudomonas aeruginosa*. *Journal of Bacteriology* 1999;181:6300–6305.

34. **Mukherjee S, Moustafa DA, Stergioula V, Smith CD, Goldberg JB,** *et al.* The PqsE and RhlR proteins are an autoinducer synthase–receptor pair that control virulence and biofilm development in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 2018;115:E9411–E9418.

35. **Farrow JM, Sund ZM, Ellison ML, Wade DS, Coleman JP, *et al.*** PqsE functions independently of PqsR-Pseudomonas quinolone signal and enhances the *rhl* quorum-sensing system. *Journal of Bacteriology* 2008;190:7043–7051.

36. **García-Reyes S, Cocotl-Yañez M, Soto-Aceves MP, González-Valdez A, Servín-González L, *et al.*** PqsR-independent quorum-sensing response of *Pseudomonas aeruginosa* ATCC 9027 outlier-strain reveals new insights on the PqsE effect on RhlR activity. *Molecular Microbiology* 2021;1113–1123.

37. **Freschi L, Vincent AT, Jeukens J, Emond-Rheault JG, Kukavica-Ibrulj I, *et al.*** The *Pseudomonas aeruginosa* Pan-Genome Provides New Insights on Its Population Structure, Horizontal Gene Transfer, and Pathogenicity. *Genome Biology and Evolution* 2019;11:109–120.

38. **Asfahl KL, Walsh J, Gilbert K, Schuster M.** Non-social adaptation defers a tragedy of the commons in *Pseudomonas aeruginosa* quorum sensing. *ISME Journal* 2015;9:1734–1746.

39. **Smalley NE, Schaefer AL, Asfahl KL, Perez C, Greenberg EP, *et al.*** Evolution of the Quorum Sensing Regulon in Cooperating Populations of *Pseudomonas aeruginosa*. *mBio* 2022;13:1–19.

40. **Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ.** Basic local alignment search tool. *Journal of Molecular Biology* 1990;215:403–410.

41. **Henikoff S, Henikoff JG.** Amino acid substitution matrices from protein blocks. *Proc Natl Acad Sci U S A* 1992;89:10915–10919.

42. **Stanton RA, Vlachos N, Halpin AL.** GAMMA: a tool for the rapid identification, classification and annotation of translated gene matches from sequencing data. *Bioinformatics* 2021;1–3.

43. **Team Rs.** RStudio: Integrated Development Environment for R.

44. **Martin DW, Schurr MJ, Mudd MH, Govan JRW, Holloway BW, *et al.*** Mechanism of conversion to mucoidy in *Pseudomonas aeruginosa* infecting cystic fibrosis patients. *Proc Natl Acad Sci U S A* 1993;90:8377–8381.

45. **Deretic V, Martin DW, Schurr MJ, Mudd MH, Hibler NS, *et al.*** Conversion to mucoidy in *Pseudomonas aeruginosa*. *Bio/Technology* 1993;11:1133–1136.

46. **Weir TL, Stull VJ, Badri D, Trunck LA, Schweizer HP, *et al.*** Global gene expression profiles suggest an important role for nutrient acquisition in early pathogenesis in a plant model of *Pseudomonas aeruginosa* infection. *Applied and Environmental Microbiology* 2008;74:5784–5791.

47. **Dumas JL, van Delden C, Perron K, Köhler T.** Analysis of antibiotic resistance gene expression in *Pseudomonas aeruginosa* by quantitative real-time-PCR. *FEMS Microbiology Letters* 2006;254:217–225.

48. **D'Argenio DA, Wu M, Hoffman LR, Kulasekara HD, Déziel E, *et al.*** Growth phenotypes of *Pseudomonas aeruginosa lasR* mutants adapted to the airways of cystic fibrosis patients. *Molecular Microbiology* 2007;64:512–533.

49. **D'Argenio DA, Calfee MW, Rainey PB, Pesci EC.** Autolysis and autoaggregation in *Pseudomonas aeruginosa* colony morphology mutants. *Journal of Bacteriology* 2002;184:6481–6489.

50. **Déziel E, Lépine F, Milot S, He J, Mindrinos MN, *et al.*** Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. *Proc Natl Acad Sci U S A* 2004;101:1339–1344.

51. **Hazan R, Que YA, Maura D, Strobel B, Majcherczyk PA, *et al.*** Auto Poisoning of the Respiratory Chain by a Quorum-Sensing-Regulated Molecule Favors Biofilm Formation and Antibiotic Tolerance. *Current Biology* 2016;26:195–206.

52. **Marie-Christine Groleaua, Hélène Taillefera, Antony T. Vincentb PC and ED**. *Pseudomonas aeruginosa* isolates defective in function of the LasR quorum sensing regulator are frequent in diverse environmental niches. *bioRxiv*. Epub ahead of print 2021. DOI: https://doi.org/10.1101/2021.03.25.437011.

53. **de Kievit T, Seed PC, Nezezon J, Passador L, Iglewski BH**. RsaL, a novel repressor of virulence gene expression in *Pseudomonas aeruginosa*. *Journal of Bacteriology* 1999;181:2175–2184.

54. **Rampioni G, Schuster M, Greenberg EP, Bertani I, Grasso M, *et al.*** RsaL provides quorum sensing homeostasis and functions as a global regulator of gene expression in *Pseudomonas aeruginosa*. *Molecular Microbiology* 2007;66:1557–1565.

55. **Mould DL, Botelho NJ, Hogan DA**. Intraspecies signaling between common variants of *Pseudomonas aeruginosa* increases production of quorum-sensing-controlled virulence factors. *mBio* 2020;11:1–16.

56. **Rumbaugh KP, Diggle SP, Watters CM, Ross-Gillespie A, Griffin AS, *et al.*** Quorum Sensing and the Social Evolution of Bacterial Virulence. *Current Biology* 2009;19:341–345.

57. **Azimi S, Roberts AEL, Peng S, Weitz JS, McNally A, *et al.*** Allelic polymorphism shapes community function in evolving *Pseudomonas aeruginosa* populations. *ISME Journal* 2020;14:1929–1942.

58. **Eldar A**. Social conflict drives the evolutionary divergence of quorum sensing. *Proc Natl Acad Sci U S A* 2011;108:13635–13640.

59. **LoVullo ED, Schweizer HP**. *Pseudomonas aeruginosa mexT* is an indicator of PAO1 strain integrity. *Journal of Medical Microbiology* 2020;69:139–145.

60. **Sandoz KM, Mitzimberg SM, Schuster M**. Social cheating in *Pseudomonas aeruginosa* quorum sensing. *Proc Natl Acad Sci U S A* 2007;104:15876–15881.

61. **Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, *et al.*** Enhanced annotations and features for comparing thousands of Pseudomonas genomes in the Pseudomonas genome database. *Nucleic Acids Research* 2016;44:D646–D653.

62. **Heberle H, Meirelles VG, da Silva FR, Telles GP, Minghim R**. InteractiVenn: A web-based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics* 2015;16:1–7.

63. **Pagès H, Aboyoun P, Gentleman R DS**. Biostrings: Efficient manipulation of biological strings.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Gene	Gene Length	Genes Returned	Unique Protein Sequences	Unique Sequences/ Gene Length
rsaL	243	806	18	0.07
rpsL	372	829	27	0.07
lasI	606	809	61	0.10
rhlI	606	827	87	0.14
vqsR	807	816	89	0.11
qscR	714	823	98	0.14
pqsB	852	822	103	0.12
pqsE	906	818	117	0.13
mucA	585	713	120	0.21
pqsR	999	818	122	0.12
pqsD	1014	820	125	0.12
pqsC	1047	823	128	0.12
rhlR	726	819	133	0.18
pqsL	1197	827	169	0.14
pqsA	1554	790	170	0.11
pqsH	1149	829	189	0.16
lasR	720	756	259	0.36

Table 1. Protein Data

Table 2. Mutation Rate

Gene	Total	Mutant	Percent Mutant	Codon Similarity
LasR	826	425	51.45%	0.9343
MucA	850	257	30.24%	0.9450
MexT	847	719	84.89%	0.9687
RhlR	850	104	12.24%	0.9925
PqsA	826	302	36.56%	0.9945
LasI	832	33	3.97%	0.9949
PqsE	844	101	11.97%	0.9951
PqsD	847	240	28.34%	0.9953
PqsL	850	447	52.59%	0.9953
VqsR	849	256	30.15%	0.9955
QscR	848	101	11.91%	0.9967
PqsR	852	200	23.47%	0.9970
RhlI	850	198	23.29%	0.9981
PqsB	847	260	30.70%	0.9984
PqsH	848	336	39.62%	0.9986
PsdR	844	38	4.50%	0.9990
PqsC	847	164	19.36%	0.9993
RpsL	852	21	2.46%	0.9998
RsaL	830	6	0.72%	0.9999

Table S1. Nucleotide sequence mutation frequency		
Gene	Mutants	Mutation Frequency (%)
lasR	553	64.91
lasI	290	34.04
rhlR	849	99.65
rhlI	847	99.41
pqsA	843	98.94
pqsB	685	80.40
pqsC	842	98.83
pqsD	843	98.94
pqsE	479	56.22
pqsH	845	99.18
pqsL	847	99.41
pqsR	789	92.61
qscR	772	90.61
rsaL	218	25.59
vqsR	559	65.61

S1. Nucleotide sequence mutation frequency		
Gene	Mutants	Mutation Frequency (%)
lasR	553	64.91
lasI	290	34.04
rhlR	849	99.65
rhlI	847	99.41
pqsA	843	98.94
pqsB	685	80.40
pqsC	842	98.83
pqsD	843	98.94
pqsE	479	56.22
pqsH	845	99.18
pqsL	847	99.41
pqsR	789	92.61
qscR	772	90.61
rsaL	218	25.59
vqsR	559	65.61

S2: Mutation profile of QS genes for 12 isolates

Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
A17	PAO1_lasR	139500	140220	Mutant	L110P,	1	1	0	0.9958	0.9986	1	720	720	+
	PAO1_lasI	140583	141189	Native	No coding mutations	0	0	0	1	1	1	606	606	+
	PAO1_rhlR	111501	112227	Native	No coding mutations	0	3	0	1	0.9959	1	726	726	-
	PAO1_rhlI	110716	111322	Mutant	S62G,D83E,	2	9	3	0.9901	0.9851	1	606	606	-
	PAO1_pqsA	266692	268246	Mutant	N180T,	1	3	2	0.9981	0.9981	1	1554	1554	-
	PAO1_pqsB	265847	266699	Mutant	P200L,	1	3	0	0.9965	0.9965	1	852	852	-
	PAO1_pqsC	264808	265855	Native	No coding mutations	0	7	1	1	0.9933	1	1047	1047	-
	PAO1_pqsD	263752	264766	Mutant	E176D,	1	6	2	0.997	0.9941	1	1014	1014	-
	PAO1_pqsE	262853	263759	Native	No coding mutations	0	0	0	1	1	1	906	906	-
	PAO1_pqsH	0	1054	Contig Edge (truncation)	223 SNPs in 1-1054,96 coding mutations,truncation at codon 268 (of 351 codons)	128	318	148	0.6658	0.7232	0.9173	1054	1149	
	PAO1_pqsL	246548	247745	Mutant	E330G,	1	4	0	0.9975	0.9967	1	1197	1197	+
	PAO1_pqsR	259613	260612	Mutant	R289L,	1	2	1	0.997	0.998	1	999	999	+
	PAO1_qscR	1815	2529	Native	No coding mutations	0	2	0	1	0.9972	1	714	714	-
	PAO1_rsaL	140209	140452	Native	No coding mutations	0	0	0	1	1	1	243	243	-
	PAO1_vqsR	6714	7521	Native	No coding mutations	0	1	0	1	0.9988	1	807	807	-
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
SWPA15J = NSWPA15a	PAO1_lasR	82843	83563	Native	No coding mutations	0	0	0	1	1	1	720	720	-
	PAO1_lasI	81874	82480	Native	No coding mutations	0	0	0	1	1	1	606	606	-
	PAO1_rhlR	593582	594308	Native	No coding mutations	0	5	1	1	0.9931	1	726	726	+
	PAO1_rhlI	594487	595093	Mutant	S62G,D83E,	2	8	3	0.9901	0.9868	1	606	606	+
	PAO1_pqsA	330200	331754	Native	No coding mutations	0	4	1	1	0.9974	1	1554	1554	+
	PAO1_pqsB	331747	332599	Native	No coding mutations	0	2	0	1	0.9977	1	852	852	+
	PAO1_pqsC	332591	333638	Mutant	L58P,	1	4	0	0.9971	0.9962	1	1047	1047	+
	PAO1_pqsD	333680	334694	Mutant	E176D,	1	6	2	0.997	0.9941	1	1014	1014	+
	PAO1_pqsE	334687	335593	Mutant	S257P,	1	5	1	0.9967	0.9945	1	906	906	+
	PAO1_pqsH	196503	197652	Native	No coding mutations	0	5	0	1	0.9956	1	1149	1149	+
	PAO1_pqsL	50584	51781	Native	No coding mutations	0	3	0	1	0.9975	1	1197	1197	-
	PAO1_pqsR	337835	338834	Native	No coding mutations	0	1	0	1	0.999	1	999	999	-
	PAO1_qscR	2085	2799	Native	No coding mutations	0	3	0	1	0.9958	1	714	714	-
	PAO1_rsaL	82611	82854	Native	No coding mutations	0	0	0	1	1	1	243	243	+
	PAO1_vqsR	190036	190843	Native	No coding mutations	0	1	0	1	0.9988	1	807	807	+
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
CND03	PAO1_lasR	50721	51430	Indel Truncation	11 bp Deletion at 440,truncation at codon 229 (of 240 codons),87 coding mutations	87	24	10	0.6375	0.9667	0.9847	709	720	+
	PAO1_lasI	51793	52399	Native	No coding mutations	0	1	0	1	0.9983	1	606	606	+
	PAO1_rhlR	109740	110466	Nonstop	15 coding mutations	15	36	21	0.938	0.9504	1	726	726	-
	PAO1_rhlI	109019	109625	Mutant	S62G,	1	8	1	0.995	0.9868	1	606	606	-
	PAO1_pqsA	171136	172690	Mutant	P263A,R347Q,	2	16	6	0.9961	0.9897	1	1554	1554	-
	PAO1_pqsB	170291	171143	Mutant	V87I,A208D,A281V,	3	10	1	0.9894	0.9883	1	852	852	-
	PAO1_pqsC	169252	170299	Mutant	T311A,	1	19	3	0.9971	0.9819	1	1047	1047	-
	PAO1_pqsD	168196	169210	Mutant	A243V,	1	16	1	0.997	0.9842	1	1014	1014	-
	PAO1_pqsE	167297	168203	Native	No coding mutations	0	3	0	1	0.9967	1	906	906	-
	PAO1_pqsH	173924	175073	Native	No coding mutations	0	7	1	1	0.9939	1	1149	1149	+
	PAO1_pqsL	65246	66443	Nonstop	D175E,R179H,A232S,R240C,A395S,P396L,L397E,G398A,*399P ,	9	31	12	0.9774	0.9741	1	1197	1197	+
	PAO1_pqsR	164057	165056	Mutant	A314V,	1	2	1	0.997	0.998	1	999	999	+
	PAO1_qscR	1814	2528	Native	No coding mutations	0	5	0	1	0.993	1	714	714	-
	PAO1_rsaL	51419	51662	Native	No coding mutations	0	0	0	1	1	1	243	243	-
	PAO1_vqsR	167457	168264	Mutant	D41H,R63I,V66I,T116A,	4	11	2	0.9851	0.9864	1	807	807	+
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand

Strain	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
5BR2	PAO1_lasR	21434	22154	Native	No coding mutations	0	0	0	1	1	1	720	720	+
	PAO1_lasI	22517	23123	Native	No coding mutations	0	0	0	1	1	1	606	606	+
	PAO1_rhlR	586980	587706	Native	No coding mutations	0	2	0	1	0.9972	1	726	726	+
	PAO1_rhlI	587885	588491	Mutant	S62G,D83E,	2	9	3	0.9901	0.9851	1	606	606	+
	PAO1_pqsA	242065	243619	Native	No coding mutations	0	2	1	1	0.9987	1	1554	1554	-
	PAO1_pqsB	241220	242072	Native	No coding mutations	0	0	0	1	1	1	852	852	-
	PAO1_pqsC	240181	241228	Native	No coding mutations	0	7	1	1	0.9933	1	1047	1047	-
	PAO1_pqsD	239125	240139	Mutant	E176D,	1	6	2	0.997	0.9941	1	1014	1014	-
	PAO1_pqsE	238226	239132	Native	No coding mutations	0	0	0	1	1	1	906	906	-
	PAO1_pqsH	17520	18669	Native	No coding mutations	0	6	1	1	0.9948	1	1149	1149	-
	PAO1_pqsL	14990	16187	Mutant	D175E,E330G,	2	5	1	0.995	0.9958	1	1197	1197	+
	PAO1_pqsR	234986	235985	Native	No coding mutations	0	1	0	1	0.999	1	999	999	+
	PAO1_qscR	5192	5906	Mutant	G197X,	1	3	0	0.9958	0.9958	1	714	714	+
	PAO1_rsaL	22143	22386	Native	No coding mutations	0	0	0	1	1	1	243	243	-
	PAO1_vqsR	24329	25136	Native	No coding mutations	0	1	0	1	0.9988	1	807	807	-
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
Co398373	PAO1_lasR	82710	83430	Native	No coding mutations	0	0	0	1	1	1	720	720	-
	PAO1_lasI	81741	82347	Native	No coding mutations	0	0	0	1	1	1	606	606	-
	PAO1_rhlR	46340	47066	Native	No coding mutations	0	6	1	1	0.9917	1	726	726	-
	PAO1_rhlI	45555	46161	Mutant	S62G,D83E,	2	9	3	0.9901	0.9851	1	606	606	-
	PAO1_pqsA	163393	164947	Native	No coding mutations	0	2	1	1	0.9987	1	1554	1554	-
	PAO1_pqsB	162548	163400	Native	No coding mutations	0	0	0	1	1	1	852	852	-
	PAO1_pqsC	161509	162556	Native	No coding mutations	0	3	1	1	0.9971	1	1047	1047	-
	PAO1_pqsD	160453	161467	Mutant	E176D,	1	5	2	0.997	0.9951	1	1014	1014	-
	PAO1_pqsE	159554	160460	Native	No coding mutations	0	1	1	1	0.9989	1	906	906	-
	PAO1_pqsH	24290	25439	Native	No coding mutations	0	5	0	1	0.9956	1	1149	1149	+
	PAO1_pqsL	30827	32024	Mutant	R30H,	1	2	0	0.9975	0.9983	1	1197	1197	-
	PAO1_pqsR	156313	157312	Native	No coding mutations	0	2	1	1	0.998	1	999	999	+
	PAO1_qscR	1815	2529	Native	No coding mutations	0	3	0	1	0.9958	1	714	714	-
	PAO1_rsaL	82478	82721	Native	No coding mutations	0	0	0	1	1	1	243	243	+
	PAO1_vqsR	17823	18630	Native	No coding mutations	0	1	0	1	0.9988	1	807	807	+
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
CPHL2000	PAO1_lasR	798489	799209	Mutant	E196G,	1	2	1	0.9958	0.9972	1	720	720	+
	PAO1_lasI	799572	800178	Native	No coding mutations	0	2	0	1	0.9967	1	606	606	+
	PAO1_rhlR	658430	659156	Native	No coding mutations	0	4	0	1	0.9945	1	726	726	-
	PAO1_rhlI	657645	658251	Mutant	S62G,D83E,	2	9	3	0.9901	0.9851	1	606	606	-
	PAO1_pqsA	330679	332233	Mutant	P263A,	1	3	2	0.9981	0.9981	1	1554	1554	+
	PAO1_pqsB	332226	333078	Native	No coding mutations	0	1	0	1	0.9988	1	852	852	+
	PAO1_pqsC	333070	334117	Native	No coding mutations	0	3	1	1	0.9971	1	1047	1047	+
	PAO1_pqsD	334159	335173	Mutant	H103P,E176D,	2	7	3	0.9941	0.9931	1	1014	1014	+
	PAO1_pqsE	335166	336072	Native	No coding mutations	0	1	1	1	0.9989	1	906	906	+
	PAO1_pqsH	161904	163053	Native	No coding mutations	0	1	0	1	0.9991	1	1149	1149	+
	PAO1_pqsL	25594	26791	Native	No coding mutations	0	2	0	1	0.9983	1	1197	1197	-
	PAO1_pqsR	338314	339313	Truncation	truncation at codon 14 (of 333 codons),2 coding mutations	2	3	1	0.994	0.997	1	999	999	-
	PAO1_qscR	1119	1833	Native	No coding mutations	0	2	0	1	0.9972	1	714	714	-
	PAO1_rsaL	799198	799441	Native	No coding mutations	0	0	0	1	1	1	243	243	-
	PAO1_vqsR	155437	156244	Mutant	R23Q,	1	2	0	0.9963	0.9975	1	807	807	-
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
Jp238	PAO1_lasR	82822	83542	Native	No coding mutations	0	0	0	1	1	1	720	720	-
	PAO1_lasI	81853	82459	Native	No coding mutations	0	0	0	1	1	1	606	606	-
	PAO1_rhlR	791947	792673	Native	No coding mutations	0	6	1	1	0.9917	1	726	726	-
	PAO1_rhlI	791162	791768	Mutant	S62G,D83E,	2	8	3	0.9901	0.9868	1	606	606	-
	PAO1_pqsA	542456	544010	Native	No coding mutations	0	1	0	1	0.9994	1	1554	1554	+
	PAO1_pqsB	541611	542463	Mutant	A50V,	1	1	0	0.9965	0.9988	1	852	852	-
	PAO1_pqsC	540572	541619	Native	No coding mutations	0	3	1	1	0.9971	1	1047	1047	-
	PAO1_pqsD	539516	540530	Mutant	L120F,E176D,	2	7	3	0.9941	0.9931	1	1014	1014	-
	PAO1_pqsE	538617	539523	Native	No coding mutations	0	0	0	1	1	1	906	906	-
	PAO1_pqsH	108635	109784	Mutant	N91S,	1	6	1	0.9974	0.9948	1	1149	1149	+

Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
	PAO1_pqsL	184199	185396	Native	No coding mutations	0	2	0	1	0.9983	1	1197	1197	+
	PAO1_pqsR	535376	536375	Native	No coding mutations	0	2	1	1	0.998	1	999	999	+
	PAO1_qscR	1815	2529	Native	No coding mutations	0	2	0	1	0.9972	1	714	714	-
	PAO1_rsaL	82590	82833	Native	No coding mutations	0	0	0	1	1	1	243	243	+
	PAO1_vqsR	102168	102975	Native	No coding mutations	0	0	0	1	1	1	807	807	+
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
Jp1155	PAO1_lasR	365051	365760	Indel Truncation	11 bp Deletion at 197,truncation at codon 229 (of 240 codons),161 coding mutations	161	20	6	0.3292	0.9722	0.9847	709	720	-
	PAO1_lasI	364082	364688	Native	No coding mutations	0	1	0	1	0.9983	1	606	606	-
	PAO1_rhlR	10191	10917	Native	No coding mutations	0	3	0	1	0.9959	1	726	726	+
	PAO1_rhlI	11096	11702	Mutant	R22L,S62G,	2	7	2	0.9901	0.9884	1	606	606	-
	PAO1_pqsA	836864	838418	Mutant	P263A,R347Q,	2	13	4	0.9961	0.9916	1	1554	1554	-
	PAO1_pqsB	836019	836871	Mutant	G279E,	1	5	1	0.9965	0.9941	1	852	852	-
	PAO1_pqsC	834980	836027	Native	No coding mutations	0	17	3	1	0.9838	1	1047	1047	-
	PAO1_pqsD	833924	834938	Mutant	A243V,	1	14	1	0.997	0.9862	1	1014	1014	-
	PAO1_pqsE	833025	833931	Native	No coding mutations	0	6	0	1	0.9934	1	906	906	-
	PAO1_pqsH	58848	59997	Native	No coding mutations	0	9	2	1	0.9922	1	1149	1149	-
	PAO1_pqsL	22214	23411	Nonstop	D175E,S206A,A232S,T378A,A395S,P396L,L397E,G398A,*399P,	9	31	13	0.9774	0.9741	1	1197	1197	-
	PAO1_pqsR	829785	830784	Native	No coding mutations	0	0	0	1	1	1	999	999	+
	PAO1_qscR	1983	2697	Mutant	S64N,	1	5	0	0.9958	0.993	1	714	714	-
	PAO1_rsaL	364819	365062	Native	No coding mutations	0	0	0	1	1	1	243	243	+
	PAO1_vqsR	65657	66464	Mutant	R63I,V66I,T116A,	3	10	1	0.9888	0.9876	1	807	807	-
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
W15Dec14	PAO1_lasR	83344	84064	Native	No coding mutations	0	0	0	1	1	1	720	720	-
	PAO1_lasI	82375	82981	Native	No coding mutations	0	0	0	1	1	1	606	606	-
	PAO1_rhlR	728506	729232	Native	No coding mutations	0	2	0	1	0.9972	1	726	726	+
	PAO1_rhlI	729412	730018	Mutant	S62G,D83E,	2	8	3	0.9901	0.9868	1	606	606	+
	PAO1_pqsA	561139	562693	Native	No coding mutations	0	3	0	1	0.9981	1	1554	1554	-
	PAO1_pqsB	560294	561146	Native	No coding mutations	0	2	0	1	0.9977	1	852	852	-
	PAO1_pqsC	559255	560302	Native	No coding mutations	0	5	1	1	0.9952	1	1047	1047	-
	PAO1_pqsD	558199	559213	Mutant	E176D,	1	7	2	0.997	0.9931	1	1014	1014	-
	PAO1_pqsE	557300	558206	Mutant	H71X,E103G,	2	4	1	0.9934	0.9956	1	906	906	-
	PAO1_pqsH	196286	197435	Native	No coding mutations	0	3	1	1	0.9974	1	1149	1149	+
	PAO1_pqsL	50331	51528	Mutant	H75D,T378A,	2	6	1	0.995	0.995	1	1197	1197	+
	PAO1_pqsR	554059	555058	Native	No coding mutations	0	2	1	1	0.998	1	999	999	+
	PAO1_qscR	421613	422327	Native	No coding mutations	0	2	0	1	0.9972	1	714	714	+
	PAO1_rsaL	83112	83355	Native	No coding mutations	0	0	0	1	1	1	243	243	+
	PAO1_vqsR	189819	190626	Native	No coding mutations	0	1	0	1	0.9988	1	807	807	+
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
Aa249	PAO1_lasR	85774	86494	Native	No coding mutations	0	0	0	1	1	1	720	720	-
	PAO1_lasI	84805	85411	Native	No coding mutations	0	0	0	1	1	1	606	606	-
	PAO1_rhlR	111501	112227	Native	No coding mutations	0	3	0	1	0.9959	1	726	726	-
	PAO1_rhlI	110716	111322	Mutant	S62G,D83E,	2	9	3	0.9901	0.9851	1	606	606	+
	PAO1_pqsA	563765	565319	Native	No coding mutations	0	2	1	1	0.9987	1	1554	1554	-
	PAO1_pqsB	562920	563772	Mutant	P200L,	1	3	0	0.9965	0.9965	1	852	852	-
	PAO1_pqsC	561881	562928	Native	No coding mutations	0	7	1	1	0.9933	1	1047	1047	-
	PAO1_pqsD	560825	561839	Mutant	E176D,	1	6	2	0.997	0.9941	1	1014	1014	-
	PAO1_pqsE	559926	560832	Native	No coding mutations	0	0	0	1	1	1	906	906	-
	PAO1_pqsH	91040	92189	Native	No coding mutations	0	5	0	1	0.9956	1	1149	1149	-
	PAO1_pqsL	225165	226362	Mutant	E330G,	1	4	0	0.9975	0.9967	1	1197	1197	+
	PAO1_pqsR	489416	490415	Native	No coding mutations	0	1	0	1	0.999	1	999	999	-
	PAO1_qscR	1815	2529	Native	No coding mutations	0	2	0	1	0.9972	1	714	714	-
	PAO1_rsaL	85542	85785	Native	No coding mutations	0	0	0	1	1	1	243	243	+
	PAO1_vqsR	97849	98656	Native	No coding mutations	0	1	0	1	0.9988	1	807	807	-
Strain:	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand

Strain	Gene	Start	Stop	Match_Type	Description	Codon_Changes	BP_Changes	Transversions	Codon_Percent	BP_Percent	Percent_Length	Match_Length	Target_Length	Strand
PT31M														
	PAO1_lasR	46184	46891	Indel Truncation	13 bp Deletion at 380,truncation at codon 173 (of 240 codons),107 coding mutations	107	22	6	0.5542	0.9694	0.9819	707	720	+
	PAO1_lasI	47254	47860	Native	No coding mutations	0	2	1	1	0.9967	1	606	606	+
	PAO1_rhlR	219207	219933	Native	No coding mutations	0	5	1	1	0.9931	1	726	726	-
	PAO1_rhlI	218422	219028	Mutant	S62G,D83E,	2	9	3	0.9901	0.9851	1	606	606	-
	PAO1_pqsA	286400	287954	Native	No coding mutations	0	2	2	1	0.9987	1	1554	1554	-
	PAO1_pqsB	285555	286407	Mutant	D47N,	1	2	0	0.9965	0.9977	1	852	852	-
	PAO1_pqsC	284516	285563	Native	No coding mutations	0	3	0	1	0.9971	1	1047	1047	-
	PAO1_pqsD	283460	284474	Mutant	E176D,	1	6	2	0.997	0.9941	1	1014	1014	-
	PAO1_pqsE	282561	283467	Native	No coding mutations	0	0	0	1	1	1	906	906	-
	PAO1_pqsH	151924	153073	Mutant	N91S,	1	7	2	0.9974	0.9939	1	1149	1149	-
	PAO1_pqsL	25330	26527	Native	No coding mutations	0	4	0	1	0.9967	1	1197	1197	-
	PAO1_pqsR	279320	280319	Native	No coding mutations	0	1	0	1	0.999	1	999	999	+
	PAO1_qscR	424624	425338	Native	No coding mutations	0	0	0	1	1	1	714	714	+
	PAO1_rsaL	46880	47123	Native	No coding mutations	0	1	0	1	0.9959	1	243	243	-
	PAO1_vqsR	158733	159540	Native		0	7	1	1	0.9913	1	807	807	-
JD303	PAO1_lasR	318802	319522	Native	No coding mutations	0	0	0	1	1	1	720	720	+
	PAO1_lasI	319885	320491	Native	No coding mutations	0	0	0	1	1	1	606	606	+
	PAO1_rhlR	799484	800210	Native	No coding mutations	0	6	1	1	0.9917	1	726	726	-
	PAO1_rhlI	798699	799305	Mutant	S62G,D83E,	2	10	2	0.9901	0.9835	1	606	606	-
	PAO1_pqsA	190222	191776	Mutant	P356L,	1	2	0	0.9981	0.9987	1	1554	1554	+
	PAO1_pqsB	191769	192621	Mutant	A50V,	1	1	0	0.9965	0.9988	1	852	852	+
	PAO1_pqsC	192613	193660	Native	No coding mutations	0	6	1	1	0.9943	1	1047	1047	+
	PAO1_pqsD	193702	194716	Mutant	E176D,	1	7	2	0.997	0.9931	1	1014	1014	+
	PAO1_pqsE	194709	195615	Native	No coding mutations	0	0	0	1	1	1	906	906	+
	PAO1_pqsH	150411	151560	Native	No coding mutations	0	3	1	1	0.9974	1	1149	1149	+
	PAO1_pqsL	187133	188330	Native	No coding mutations	0	6	0	1	0.995	1	1197	1197	+
	PAO1_pqsR	197857	198856	Native	No coding mutations	0	0	0	1	1	1	999	999	-
	PAO1_qscR	829886	830600	Native	No coding mutations	0	1	0	1	0.9986	1	714	714	+
	PAO1_rsaL	319511	319754	Native	No coding mutations	0	0	0	1	1	1	243	243	-
	PAO1_vqsR	157220	158027	Native	No coding mutations	0	0	0	1	1	1	807	807	-