Chelator sensing and lipopeptide interplay mediates molecular

interspecies interactions between soil bacilli and pseudomonads

Sofija Andric[1*], Thibault Meyer[1,Φ *], Augustin Rigolet[1], Anthony Argüelles Arias[1], Sébastien Steels[1],

Grégory Hoff[1,#], Monica Höfte[2], René De Mot[3], Andrea McCann[4], Edwin De Pauw[4] and Marc

Ongena[1]

[1]Microbial Processes and Interactions Laboratory, Terra Teaching and Research Center, Gembloux Agro-Bio

Tech, University of Liège, Gembloux, Belgium

[2]Laboratory of Phytopathology, Department of Plants and Crops, Faculty of Bioscience engineering, Ghent

University, Gent, Belgium

[3] Centre of Microbial and Plant Genetics, Faculty of Bioscience Engineering, University of Leuven, Heverlee,

Belgium

[4] Mass Spectrometry Laboratory, , MolSys Research Unit, Department of Chemistry, University of Liège,

Belgium

[Φ] Current address: UMR Ecologie Microbienne, F-69622, University of Lyon, Université Claude Bernard Lyon

1, CNRS, INRAE, VetAgro Sup, Villeurbanne, France

[#] Current address: Ecology and Biodiversity, Department of Biology, Utrecht University, Padualaan 8, 3584

CH, Utrecht, The Netherlands

* Equal contribution

Abstract

Some bacterial species are important members of the rhizosphere microbiome and confer protection to the host plant against pathogens. However, our knowledge of the multitrophic interactions determining the ecological fitness of these biocontrol bacteria in their highly competitive natural niche is still limited. In this work, we investigated the molecular mechanisms underlying interactions between *B. velezensis,* considered as model plant-associated and beneficial species in the *Bacillus* genus, and *Pseudomonas* as a rhizosphere-dwelling competitor. Our data show that *B. velezensis* boosts its arsenal of specialized antibacterials upon the perception of the secondary siderophore enantio-pyochelin produced by phylogenetically distinct pseudomonads and some other genera. We postulate that *B. velezensis* has developed some chelator sensing systems to learn about the identity of its surrounding competitors. Illustrating the multifaceted molecular response of *Bacillus*, surfactin is another crucial component of the secondary metabolome mobilized in interbacteria competition. Its accumulation not only enhances motility but, unexpectedly, the lipopeptide also acts as a chemical trap that reduces the toxicity of other lipopeptides released by *Pseudomonas* challengers. This in turn favors the persistence of *Bacillus* populations upon competitive root colonization. Our work thus highlights new ecological roles for bacterial secondary metabolites acting as key drivers of social interactions.

Soil is one of the richest ecosystems in terms of microbial diversity and abundance[1]. However, the scarcity of resources makes it one of the most privileged environments for competitive interspecies interactions[2,3]. A subset of the bulk soil microbes has evolved to dwell in the rhizosphere compartment surrounding roots due to continued nutrient-enriched exudation from the plant. Compared to bulk soil, microbial warfare in the rhizosphere is presumably even more intense as the habitat is spatially restricted and more densely populated[3]. Besides rivalry for nutrients (exploitative), interference competition is considered a key factor driving microbial interactions and community assembly. This competition can involve signal interference or toxins deployed by contact-dependent delivery systems[4,5] but is mainly mediated at distance through the emission of various molecular weapons. The molecular basis of interference interactions and their phenotypic outcomes between diverse soil bacterial species have been amply investigated in the last decade[6,7].

Bacilli belonging to the *B. subtilis* complex are ubiquitous members of the rhizosphere microbiome[2,8,9]. Among these species, *B. velezensis* has emerged as plant-associated model bacilli, displaying strong potential as biocontrol agent reducing diseases caused by phytopathogens[10]. *B. velezensis* distinguishes itself from other species of the *B. subtilis* group by its richness in biosynthetic gene clusters (BGCs, representing up to 13% of the whole genome) responsible for the synthesis of bioactive secondary metabolites (BSMs)[11,12]. This chemically-diverse secondary metabolome includes volatiles, terpenes, non-ribosomal (NR) dipeptides, cyclic lipopeptides (CLPs) and polyketides (PKs), but also ribosomally synthesized lantibiotics and larger bacteriocins (RiPP)[13,14]. BSMs are involved in biocontrol activity via direct inhibition of pathogenic microbes and/or via stimulation of the plant immune system[15,16]. From an ecological viewpoint, BSMs also contribute to competitiveness in the rhizosphere niche thanks to multiple and

complementary functions as drivers of developmental traits, as antimicrobials, or as signals

initiating cross-talk with the host plant[17–19].

Mostly guided by practical concerns for use as biocontrol agents, research on BSMs has

mainly focused on the characterization of their biological activities. However, the impact of

environmental factors that may modulate their expression under natural conditions still remains

poorly understood. It includes interactions with other organisms sharing the niche. Some recent

reports illustrate how soil bacilli may adapt their behavior upon sensing bacterial competitors but

almost exclusively focusing on developmental traits (sporulation, biofilm formation, or motility)[7].

Unlike other genera such as *Streptomyces*, it remains largely unknown to what extent bacilli in

general and *B. velezensis* in particular, may modulate the expression of their secondary

metabolome upon interaction with other bacteria[7,20]. In this work, we investigated the molecular

outcomes of interspecies interactions in which *B. velezensis* may engage. We selected

Pseudomonas as challenger considering that species of this genus are also highly competitive and

commonly encountered in rhizosphere microbiomes[8]. We performed experiments under nutritional

conditions mimicking the oligotrophic rhizosphere environment and used contact-independent

settings for pairwise interaction which probably best reflect the real situation in soil. Our data

revealed that the two bacteria initiate multifaceted interactions mostly mediated by the non-

ribosomally synthesized components of their secondary metabolome. We pointed out unsuspected

roles for some of these BSMs in the interaction context. Beyond its role as a metal chelator, the

Pseudomonas secondary siderophore enantio-pyochelin (E-PCH) acts as a signal triggering dual

production of PKs and RiPP in *Bacillus,* while specific lipopeptides modulate the inhibitory

interaction between the two species. This results in marked phenotypic changes in *B. velezensis*

such as higher antibacterial potential, enhanced motility and protective effect via chemical trapping.

We also illustrate the relevance of these outcomes in the context of competitive root colonization.

Results

B. velezensis modulates its secondary metabolome and boosts antibacterial activity upon sensing *Pseudomonas* metabolites

We used *B. velezensis* strain GA1 as a BSM-rich and genetically amenable isolate representative of the species. Genome mining with AntiSMASH 5.0[21] confirmed the presence of all gene clusters necessary for the biosynthesis of known BSMs typically formed by this bacterium (Supplementary Table 1). Based on the exact mass and absence of the corresponding peaks in deletion mutants, most of the predicted non-ribosomal (NR) secondary metabolites were identified in cell-free crude supernatants via optimized UPLC-MS (Supplementary Fig. 1). It includes the whole set of cyclic lipopeptides (CLPs of the surfactin, fengycin and iturin families) and polyketides (PKs difficidin, macrolactin and bacillaene) with their multiple co-produced structural variants, as well as the siderophore bacillibactin. We verified that all these compounds are readily formed in the so-called exudate-mimicking (EM) medium reflecting the specific content in major carbon sources typically released by roots of *Solanaceae* (such as tomato) plants[22]. In addition to these NR products, genes encoding RiPPs such as amylocyclicin and amylolysin are also present in GA1, but these compounds could not be reliably detected in culture broths. The NR dipeptide bacilysin was also predicted but not detected. We selected as the main interaction partner the plant-associated *Pseudomonas* sp. strain CMR12a based on its biocontrol potential and its production of multiple secondary metabolites[23–27]. Genome mining confirmed the potential of CMR12a to synthesize a range of BSMs, including antimicrobial phenazines, the siderophores pyoverdine (PVD) (structure confirmation in Supplementary Fig. 2) and E-PCH as well as two structurally distinct CLPs, sessilins and orfamides (Supplementary Table 1). In contrast to *Bacillus*, the capacity to co-produce two different CLPs is a quite rare trait for non-phytopathogenic pseudomonads and represented an additional criterion for selecting strain CMR12a for this study[23,27–29]. In the case of CMR12a, according to the exact mass and absence of the corresponding peaks in deletion mutants, all these

compounds were detected in EM culture broth but most of them are more efficiently produced upon

growth in casamino acids medium (CAA) commonly used for *Pseudomonas* cultivation

(Supplementary Fig. 1).

Our prime objective was to evaluate the intrinsic potential of *B. velezensis* to react to the

perception of *Pseudomonas* metabolites in an experimental setting avoiding interferences due to

diffusion constraints in a semi-solid matrix or due to the formation of impermeable biofilm

structures. The first assays were performed by growing GA1 in agitated liquid EM medium

supplemented or not with (sterile) BSM-containing spent medium of CAA-grown CMR12a (CFS,

cell-free supernatant). At a low dose (2% (v/v)), the addition of this CFS extract led to a marked

increase in the production of some GA1 NR metabolites. Significantly higher amounts were

measured for surfactins, bacillaene or its dehydrated variant dihydrobacillaene (2H-bae), difficidin

or its oxidized form, and bacillibactin (Fig. 1a, b) but not for other compounds such as fengycins,

iturins and macrolactins (Fig. 1a).

The boost in BSMs synthesis triggered by *Pseudomonas* CFS was associated with an

increase in the antibacterial activity of the corresponding extracts when tested for growth inhibition

of *Xanthomonas campestris* and *Clavibacter michiganensis* used respectively as representative of

Gram-negative and Gram-positive plant pathogenic bacteria of agronomical importance[30] (Fig. 1c).

Since most of the BSMs are not commercially available and our attempt to purify PKs failed due to

chemical instability, we could not use individual compounds for their specific involvement in

bacterial inhibition. As an alternative, we generated and tested a range of GA1 knockout mutants

including the Δ*sfp* derivative specifically repressed in 4'-phosphopantetheinyl transferase essential

for the proper functioning of the PK and NRP biosynthesis machinery. Full loss of anti-

Xanthomonas activity in Δ*sfp* extracts indicated a key role for NR BSMs and ruled out the possible

involvement of other chemicals known for their antibacterial activity such as bacilysin or RiPPs (Fig.

1d). Loss of function of mutants specifically repressed in the synthesis of individual compounds

pointed out the key role of (oxy)difficidin and to a lower extent of 2H-bae in *Xanthomonas* inhibition

(Fig. 1d). These two PKs are also responsible for GA1 inhibitory activity toward other important

bacterial phytopathogens such as *Pectobacterium carotovorum*, *Agrobacterium tumefaciens* and

Rhodococcus fasciens but are not involved in the inhibition of plant pathogenic *Pseudomonas*

species for which bacilysin may be the active metabolite (Supplementary Fig. 3). However, as

illustrated below, *B. velezensis* does not display significant toxicity against CMR12a and other non-

pathogenic soil *Pseudomonas* isolates tested here. Stimulation of PKs synthesis upon sensing

CMR12a is not specific to GA1 and was also observed in other *B. velezensis* strains with well-

known biocontrol potential such as S499, FZB42 and QST713[31–33] (Supplementary Fig. 4).

In contrast to *Xanthomonas*, enhanced antibiotic activity against *Clavibacter* is not mediated

by NR products as shown by the fully conserved activity in the Δ*sfp* mutant (Fig. 1d). Therefore, we

suspected from genomic data and literature[34] that RiPPs such as amylocyclicin could be involved in

inhibition. This hypothesis was supported by the 80% reduction in antibiotic potential observed for

the Δ*acnA* mutant knocked out for the corresponding biosynthesis gene (Fig. 1d). Besides, RT-

qPCR data revealed a highly induced expression of *acnA* gene in GA1 cells upon supplementation

with CMR12a CFS (Fig. 1e). However, we were not able to provide evidence for higher

accumulation of the mature peptide in the medium. Enhanced expression of the *acnA* gene in

presence of *Pseudomonas* products was also observed for strain S499 (Supplementary Fig. 5).

E-PCH acts as a signal sensed by *Bacillus* to stimulate polyketide production

We next wanted to identify the signaling molecules secreted by *Pseudomonas* that are sensed by

Bacillus cells and lead to improved BSMs production. For that purpose, we used 2H-bae as an

indicator of the *Bacillus* response because it represents the most consistent and highly boosted

polyketide. We first compared the triggering potential of CFS obtained from knockout mutants of

CMR12a specifically lacking the different identified metabolites (Supplementary Fig. 1). Only

extracts from mutants impaired in the production of siderophores and more specifically E-PCH were

significantly affected in PKs-inducing potential (Fig. 2a). Possible involvement of this compound

was supported by the drastic reduction in the activity of CFS prepared from CMR12a culture in CAA

medium supplemented with Fe^{3+} where siderophore expression is repressed (Fig. 2b,

Supplementary Fig. 6). We also performed bioactivity-guided fractionation and data showed that

only extracts containing PVD and/or E-PCH displayed consistent PKs-triggering activity

(Supplementary Fig. 7). HPLC-purified compounds were also tested independently at a

concentration similar to the one measured in CFS CAA extract revealing a much higher PK-

triggering activity for E-PCH compared to the main PVD isoform (Fig. 2b). Dose-dependent assays

further indicated that supplementation with PVD, as strong chelator[35], caused iron limitation in the

medium which is sensed by GA1. It is supported by the marked increase in production of the

siderophore bacillibactin in GA1 wild-type (Fig. 2c) and by the reduced growth of the ΔdhbC mutant,

repressed in bacillibactin synthesis, upon PVD addition (Fig. 2d, Supplementary Fig. 8). This last

result indicates that PVD in its ferric form cannot be taken up by GA1 despite the presence of

several transporters for exogenous siderophores in *B. velezensis* similar to those identified in *B.*

subtilis[36,37] based on genome comparison (Supplementary Table 2). Therefore, we assumed that

iron stress mediated by PVD only induces a rather limited boost in PKs production. We validated

that such response is not due to iron starvation by supplementing GA1 culture with increasing

doses of the 2,2'-dipyridyl (DIP) chemical chelator that cannot be taken up by *Bacillus* cells (Fig.

2b). By contrast, the addition of E-PCH with a much lower affinity for iron does not activate

bacillibactin synthesis (Fig. 2c) and does not affect ΔdhbC growth at the concentrations used (Fig.

2d, Supplementary Fig. 8). We conclude that the activity of this compound referred to as secondary

siderophore is not related to iron-stress. If internalized, E-PCH can cause oxidative stress and

damage in other bacteria as reported for *E. coli*[38,39]. However, the absence of toxicity toward GA1

indicates that E-PCH is not taken up by *Bacillus* cells and thus clearly acts as a signal molecule

perceived at the cell surface. PKs boost also occurred upon addition of CFS obtained from other

Pseudomonas isolates producing pyochelin-type siderophores, such as *P. protegens* Pf-5[40].

However, PKs stimulation was similarly observed in response to *P. tolaasii* CH36[41] which does not

form pyochelin, indicating that other unidentified BSMs may act as triggers in other strains

(Supplementary Fig. 9).

Enhanced motility as distance-dependent and surfactin-mediated response of *Bacillus*

Surfactin production is stimulated by CMR12a CFS and by pure E-PCH (Fig. 1a and

Supplementary Fig. 10). Based on mutant loss-of-function analysis, this multifunctional CLP does

not contribute to the antibacterial potential of *B. velezensis* (Fig. 1d and Supplementary Fig. 3) but

is known to be notably involved in developmental processes of multicellular communities such as

biofilm formation and motility[42]. Therefore, we wanted to test a possible impact of *Pseudomonas* on

the motile phenotype of *B. velezensis* upon co-cultivation on plates. We observed distance-

dependent enhanced motility on medium containing high agar concentrations (1.5% m/v) which

phenotypically resembles the sliding-type of motility illustrated by typical "van Gogh bundles"[42] (Fig.

3a). This migration pattern is flagellum independent but depends on multiple factors including the

synthesis of surfactin which reduces friction at the cell-substrate interface[42]. We thus suspected

such enhanced motility to derive from an increased formation of the lipopeptide. This was

supported by the almost full loss of migration of the Δ*srfaA* mutant in these interaction conditions

(Fig. 3b). Moreover, spatial mapping via MALDI-FT-ICR MS imaging confirmed a higher

accumulation of surfactin ions in the interaction zone and around the *Bacillus* colony when growing

at a short or intermediate distance from the *Pseudomonas* challenger, compared to the largest

distance where the motile phenotype is much less visible (Fig. 3c). These data indicate that *Bacillus*

cells in the microcolony perceive a soluble signal diffusing from the *Pseudomonas* colony over a

limited distance.

Interplay between CLPs drives antagonistic interactions

Besides modulating secondary metabolite synthesis, we further observed that confrontation with *Pseudomonas* may also lead to some antagonistic outcomes. GA1 growth as planktonic cells is slightly inhibited upon supplementation of the medium with 2% v/v CMR12a CFS but this inhibition is much more marked at a higher dose (Supplementary Fig. 11). To identify the *Pseudomonas* compound retaining such antibiotic activity, we tested the effect of CFS from various CMR12a mutants impaired in the synthesis of lipopeptides and/or phenazines. Even if some contribution of other compounds cannot be ruled out, it revealed that the CLP sessilin is mainly responsible for toxic activity toward GA1 grown in liquid cultures but also when the two bacteria are grown at close proximity on gelified EM medium (Fig. 4a). Nevertheless, we observed that the sessilin-mediated inhibitory effect is markedly reduced by delaying CFS supplementation until 6 h of *Bacillus* culture instead of adding it at the beginning of incubation (Fig. 4b). This suggested that early secreted *Bacillus* compounds may counteract the toxic effect of sessilin. We hypothesized that surfactin can play this role as it is the first detectable BSM to accumulate in significant amounts in the medium early in the growth phase. We tested the surfactin-deficient mutant in the same conditions and observed that its growth is still strongly affected indicating that no other GA1 compound may be involved in toxicity alleviation. Chemical complementation with purified surfactins restored growth to a large extent, providing further evidence for a protective role of the surfactin lipopeptide (Fig. 4b). Such sessilin-dependent inhibition also occurred when bacteria were confronted on solid CAA medium (Fig. 4c-I) favoring *Pseudomonas* BSM production. In these conditions, the formation of a white precipitate in the interaction zone was observed with CMR12a wild-type but not when GA1 was confronted with the ΔsesA mutant (Fig. 4c). UPLC-MS analysis of ethanol extracts from this white-line area confirmed the presence of sessilin ions but also revealed an accumulation of surfactin from GA1 in the confrontation zone (Fig. 4d). The involvement of surfactins in precipitate formation was confirmed by the absence of this white-line upon testing the ΔsrfaA mutant of GA1

(Fig. 4c-I, II). The loss of surfactin production and white-line formation was associated with a higher

sensitivity of the *Bacillus* colony to the sessilin toxin secreted by *Pseudomonas*. Altogether, these

data indicate that surfactin acts as a chemical trap and inactivates sessilin via co-aggregation into

insoluble complexes.

A similar CLP-dependent antagonistic interaction and white-line formation were observed

upon co-cultivation of GA1 with *P. tolaasii* strain CH36 producing tolaasin (Fig. 4c-II), a CLP

structurally very similar to sessilin (only differing by two amino acid residues, Supplementary Fig.

12). However, this chemical aggregation is quite specific regarding the type of CLP involved, since

it was not visible upon the interaction of GA1 with other *Pseudomonas* strains forming different CLP

structural groups that are not toxic for *Bacillus* (Fig. 4c-III, see Supplementary Fig. 12 and 13 for

identification and structures). Sessilin/tolaasin-dependent toxicity and white-line formation were

also observed when other surfactin-producing *B. velezensis* isolates were confronted with CMR12a

and CH36 (Supplementary Fig. 14 and 15, respectively). Although the chemical basis and the

stoichiometry of such molecular interaction remain to be determined, it probably follows the same

rules as observed for the association between sessilins/tolaasins and other endogenous

Pseudomonas CLPs such as WLIP or orfamides[28] or between CLPs and other unknown

metabolites[43,44].

BSMs-mediated interactions drive competitive root colonization

Our *in vitro* data point out how *B. velezensis* may modulate its secondary metabolome when

confronted with *Pseudomonas*. To appreciate the relevance of our findings in a more realistic

context, we next evaluated whether such BSMs interplay may also occur upon root co-colonization

of tomato plantlets and possibly impact *Bacillus* fitness. When inoculated independently, CMR12a

colonized roots more efficiently than GA1 within the first 3 days, most probably due to a higher

intrinsic growth rate[45]. Upon co-inoculation, the CMR12a colonization rate was not affected but GA1

populations were reduced compared to mono-inoculated plantlets (Fig. 5a). UPLC-MS analysis of

methanolic extracts prepared from co-bacterized roots (and surrounding medium) revealed

substantial amounts of E-PCH (Supplementary Fig. 16) indicating that the molecule is readily

formed under these conditions and could therefore also act as a signal *in planta.* Probably due to

the low populations of GA1, we could not detect *Bacillus* PKs and RiPPs in these extracts.

However, a significantly enhanced expression of gene clusters responsible for the synthesis of

bacillaene, difficidin and amylocyclicin was observed in GA1 cells co-inoculated with *Pseudomonas*

compared to single inoculation (Fig. 5b). It indicated that the metabolite response observed in GA1

in vitro cultures in EM medium may also occur upon competitive colonization where the bacteria

feed exclusively on root exudates.

Lipopeptides involved in interference interaction are also readily formed upon single and

dual root colonization (Fig. 5d). We hypothesized that the inhibitory effect of sessilin may impact the

colonization potential of GA1 in presence of CMR12a which was confirmed by the increase in GA1

populations co-inoculated with the ΔsesA mutant (Fig. 5c). Moreover, colonization by the ΔsrfaA

mutant is more impacted compared to WT when co-cultivated with CMR12a and a significant gain

in root establishment is recovered upon co-colonization with the ΔsesA mutant (Fig. 5d). The

sessilin-surfactin interplay thus also occurs *in planta.* Sessilin would confer a competitive

advantage to CMR12a during colonization by inhibiting GA1 development but efficient surfactin

production on roots may provide some protection to the *Bacillus* cells.

Discussion

It has been recently reported that *Pseudomonas* toxin delivery via Type VI secretion system

and antibiotic (2,4-diacetylphloroglucinol) production may impact biofilm formation and sporulation

in *B. subtilis*[46,47]. However, our current understanding of the molecular basis of interactions between

soil bacilli and pseudomonads is still rather limited. Here we show that the model species *B.*

velezensis can mobilize a substantial part of its secondary metabolome in response to

Pseudomonas competitors. To our knowledge, it is the first evidence for enhanced synthesis of

both broad-spectrum polyketides and RiPP in *Bacillus* upon a perception of other bacteria, in

contact-independent *in vitro* settings and upon competitive root colonization. This correlates with an

enhanced antibacterial potential which is of interest for biocontrol but which can also be considered

as a defensive strategy to persist in its natural competitive niche. Upon sensing *Pseudomonas*, *B.*

velezensis calls on its antibiotic arsenal but also recruits its surfactin lipopeptide to improve

multicellular mobility. This may be viewed as an escape mechanism enabling *Bacillus* cells to

relocate after detecting harmful challengers. Improved motility of *B. subtilis* has been already

described upon sensing competitors such as *Streptomyces venezuelae*[7,48] but no relationship was

established with enhanced production of BSMs potentially involved in the process. We also

highlight a new role for surfactin acting as a chemical shield to counteract the toxicity of exogenous

CLPs. Intraspecies CLP co-precipitation has been reported[23] but our results make sense of this

phenomenon in the context of interference interaction between two different genera. *In planta*, this

new function of surfactin contributes to *Bacillus* competitiveness for root invasion. This has to be

added to other previously reported implications of surfactin in *B. subtilis* interspecies interactions,

such as interfering with the growth of closely related species in synergy with cannibalism toxins[49],

inhibiting the development of *Streptomyces* aerial hyphae[50], or participating in the expansion and

motility of the interacting species[47]. We postulate that such *Bacillus* metabolite response largely

contributes to mount a multi-faceted defensive strategy in order to gain fitness and persistence in

its natural competitive niche.

Furthermore, we exemplify that PKs stimulation in *B. velezensis* is mainly mediated by the

Pseudomonas secondary siderophore pyochelin, although it cannot be excluded that other

secreted products may also play a role. *Bacillus* perceives pyochelin in a way independent of iron

stress and piracy, indicating that beyond its iron-scavenging function, this siderophore may also act

as infochemical in interspecies cross-talk. In the pairwise system used here, E-PCH signaling

superimposes the possible effect of iron limitation in the external medium which may also result in

enhanced production of antibacterial metabolites by *Bacillus,* as occasionally reported[51]. That said,

due to the limitation in bioavailable iron, almost all known rhizobacterial species have adapted to

produce their iron-scavenging molecules to compete for this essential element[52–54]. Siderophore

production is thus widely conserved among soil-borne bacteria[55]. It means that upon recognition of

exogenous siderophores, any isolate may somehow identify surrounding competitors. However,

some of these siderophores are structurally very variable and almost strain-specific (such as PVDs

from fluorescent pseudomonads) while some others are much more widely distributed across

species and even genera (enterobactin-like, citrate)[52]. In both cases, their recognition would not

provide proper information about the producer because they are too specific or too general,

respectively. Interestingly, the synthesis of E-PCH and its structurally very close enantio form is

conserved in several but not all *Pseudomonas* sp.[56–58] as well as in a limited number of species

belonging to other genera such as *Burkholderia*[59] and *Streptomyces*[60,61]. We therefore hypothesize

that *Bacillus* may have evolved some chelator-sensing systems targeting siderophores that are

conserved enough to be detected but restricted to specific microbial phylogenetic groups. With this

mechanism, soil bacilli would rely on siderophores as public goods to accurately identify

competitors and respond in an appropriate way like remodeling its BSM secretome. This novel

concept of chelator sensing represents a new facet of siderophore-mediated social interactions.

Whether it is used for other secondary siderophores than E-PCH and if so, whether this adaptative

trait can be generalized to other soil-dwelling species deserves to be further investigated given its

possible impact on soil bacterial ecology. Beyond the notion of specialized metabolites, we point

out unsuspected functions for some bacterial small molecules in the context of interactions between

clades that are important members of the plant-associated microbiome.

Methods

Bacterial strains and growth conditions

Strains and plasmids used in this study are listed in Supplementary Table 3. *B. velezensis* strains were grown at 30 °C on, half diluted, recomposed exudate solid medium (EM)[22] or in liquid EM with shaking (160 rpm). Deletion mutants of *B. velezensis* were selected on appropriate antibiotics (chloramphenicol at 5 µg/ml, phleomycin at 4 µg/ml, kanamycin at 25 µg/ml) on Lysogeny broth (LB) (10 g l^{-1} NaCl, 5 g l^{-1} yeast extract and 10 g l^{-1} tryptone). *Pseudomonas* sp. strains were grown on King B (20 g l^{-1} of bacteriological peptone, 10 g l^{-1} of glycerol and 1.5 g l^{-1} of K_2HPO_4, 1.5 g l^{-1} of $MgSO_4.7H_2O$, pH = 7) and casamino acid (CAA) solid and liquid medium (10 g l^{-1} casamino acid, 0.3 g l^{-1} K_2HPO_4, 0.5 g l^{-1} $MgSO_4$ and pH = 7) with shaking (120 rpm), at 30 °C. The phytopathogenic bacterial strains were grown on LB and EM solid and liquid media and with shaking (150 rpm), at 30 °C.

Construction of deletion mutants of *B. velezensis* GA1

All deletion mutants were constructed by marker replacement. Briefly, 1 kb of the upstream region of the targeted gene, an antibiotic marker (chloramphenicol, phleomycin or kanamycin cassette) and downstream region of the targeted gene were PCR amplified with specific primers (Supplementary Table 3). The three DNA fragments were linked by overlap PCR to obtain a DNA fragment containing the antibiotic marker flanked by the two homologous recombination regions. This latter fragment was introduced into *B. velezensis* GA1 by natural competence induced by nitrogen limitation[62]. Homologous recombination event was selected by chloramphenicol resistance (phleomycin resistance for double mutants or kanamycin resistance for triple mutants) on LB medium. All gene deletions were confirmed by PCR analysis with the corresponding UpF and DwR specific primers and by the loss of the corresponding BSMs production.

Transformation of the *B. velezensis* GA1 strain was performed following the protocol

previously described[62] with some modifications. One fresh GA1 colony was inoculated into LB liquid

medium at 37 °C (160 rpm) until reaching an OD_{600nm} of 1.0. Afterwards, cells were washed one

time with peptone water and one time with a modified Spizizen minimal salt liquid medium (MMG)

(19 g l^{-1} K_2HPO_4 anhydrous; 6 g l^{-1} KH_2PO_4; 1 g l^{-1} Na_3 citrate anhydrous; 0.2 g l^{-1} $MgSO_4$ $7H_2O$; 2 g

l^{-1} Na_2SO_4; 50 μM $FeCl_3$ (sterilized by filtration at 0.22 μm); 2 μM $MnSO_4$; 8 g l^{-1} glucose; 2 g l^{-1} L-

glutamic acid; pH 7.0), 1 μg of DNA recombinant fragment was added to the GA1 cells suspension

adjusted to an OD_{600nm} of 0.01 into MMG liquid medium. One day after incubation at 37 °C with

shaking at 165 rpm, bacteria were spread on LB plates supplemented with the appropriate

antibiotic to select positive colonies.

Construction of deletion mutants of *Pseudomonas* sp. CMR12a

E-PCH and PVD mutants of *Pseudomonas* sp. CMR12a were constructed using the I-SceI system

and the pEMG suicid vector[63,64]. Briefly, the upstream and downstream region flanking the *pchA*

(*C4K39_5481*) or the *pvdI* (*C4K39_6027*) genes were PCR amplified (primers listed in the

Supplementary Table 3), linked via overlap PCR and inserted into the pEMG vector. The resulting

plasmid (Supplementary Table 3) was integrated by conjugation into the *Pseudomonas* sp.

CMR12a chromosome via homologous recombination. Kanamycin (25μg/mL) resistant cells were

selected on King B agar plates and transformed by electroporation with the pSW-2 plasmid

(harboring I-SceI system). Gentamycin (20μg/ml) resistant colonies on agar plates were transferred

to King B medium with and without kanamycin to verify the loss of the antibiotic (kanamycin)

resistance. *Pseudomonas* mutants were identified by PCR with the corresponding UpF and DwR

specific primers and via the loss of E-PCH or/and PVD production (see section Secondary

metabolites analysis).

***Pseudomonas* sp. cell-free supernatant**

Pseudomonas sp. strains were grown overnight on LB solid medium, at 30 °C. The cell suspension was adjusted to OD_{600nm} 0.05 by resuspension in 100 ml of CAA and when appropriate supplemented with 20 µg/l of $FeCl_3.6H_2O$ (iron supplementation). Cultures were shaken at 120 rpm at 30 °C for 48 h and then centrifuged at 5000 rpm at room temperature (22 °C) for 20 min. The supernatant was filter-sterilized (0.22 µm pore size filters) and stored at -20 °C until use.

Dual interactions

B. velezensis strains were grown overnight on LB solid medium, at 30 °C. Cells were resuspended in 2 ml of EM liquid medium to a final OD_{600nm} of 0.1 in which 1, 2, or 4% v/v (depending on the experiment and indicated in the figures legends) of *Pseudomonas* CFS were added while the control remained un-supplemented. *B. velezensis* liquid cultures were shaken in an incubator at 300 rpm at 30 °C for 24 h. Additionally, 2 ml of the (co-)culture supernatants were sampled at 8 h and 24 h, centrifugated at 5000 rpm at room temperature (approx. 22 °C) for 10 min to extract supernatants and collect the cells. Further, cell-free (co-)culture supernatants were filter-sterilized (0.22µm) and used for analytical analysis of secondary metabolites and antibacterial assays. For some experiments using 2H-bae as a marker, the CFS obtained from the double mutant sessilins and orfamides (Δ*sesA-ofaBC*) was used instead of CFS from CMR12a wild-type because it yielded a higher response and lower inhibition interferences by CLPs. The remaining cells, after supernatant collection, were stored at -80 °C to avoid RNA degradation, until performing RT-qPCR analysis.

Antimicrobial activity assays

Antibacterial activity of the *B. velezensis* supernatant generated after dual interaction with *Pseudomonas* CFS was tested against *X. campestris* pv. *campestris* and *C. michiganensis* subsp. *michiganensis.* The activity of co-culture supernatants was quantified in microtiter plates (96-well)

filled with 250 µl of LB liquid medium, inoculated at OD_{600nm} = 0.1 with *X. campestris* pv. *campestris*

and *C. michiganensis* subsp. *michiganensis* and supplemented with 2% or 6% v/v of the

supernatants, respectively. The activity of (co-)culture supernatants was estimated by measuring

the pathogen OD_{600nm} every 30min during 24 h with a Spectramax® (Molecular Devices,

Wokingham, UK), continuously shaken, at 30 °C. For estimating the activity of co-culture

supernatants on a solid medium, 5 µl supernatant was applied to a sterile paper disk (5 mm

diameter). After drying, disks were placed on LBA square plates previously inoculated with a

confluent layer of *X. campestris* pv. *campestris*, *C. michiganensis* subsp. *michiganensis, P.*

carotovorum, P. fuscovaginae, P. cichorii, A. tumefaciens or R. fascians. LB liquid medium was

used as a negative control. Plates were incubated at 25 °C for 48 h. Three repetitions were done

and the inhibition zones from the edge of the paper discs to the edge of the zone were measured.

Antibacterial activity of the different *Pseudomonas* strains on the *B. velezensis* strains growth were

tested by adding different % (v/v) of the corresponding *Pseudomonas* CFS in microtiter plates (96-

well) filled with 250 µl of EM liquid medium. *B. velezensis* OD_{600nm} after 7 h was measured with a

Spectramax® (Molecular Devices, Wokingham, UK).

RNA isolation and RT-qPCR

RNA extraction and DNAse treatment were carried out using the NucleoSpin RNA Kit (Macherey

Nagel, Germany), following the Gram + manufacturer's protocol. RNA quality and quantity were

performed with Thermo scientific NanoDrop 2000 UV-vis Spectrophotometer. Primer 3 program

available online was used for primer design and primers were synthesized by Eurogentec. The

primer efficiency was evaluated and primer pairs showing an efficiency between 90 and 110% in

the qPCR analysis were selected. Reverse transcriptase and RT-qPCR reactions were conducted

using the Luna® Universal One-Step RT-qPCR Kit (New England Biolabs, Ipswich, MA, United

States). The reaction was performed with 50 ng of total RNA in a total volume of 20 µL: 10 µL of

luna universal reaction mix, 0.8 μL of each primer (10 μM), 5 μL of cDNA (50ng), 1 μL of RT

Enzyme MIX, 2.4 µl of Nuclease-free water. The thermal cycling program applied on the ABI

StepOne was: 55 °C for 10 min, 95 °C for 1 min, 40 cycles of 95 °C for 10 s and 60 °C for 1 min,

followed by a melting curve analysis performed using the default program of the ABI StepOne

qPCR machine (Applied Biosystems). The real-time PCR amplification was run on the ABI step-one

qPCR instrument (Applied Biosystems) with software version 2.3. The relative gene expression

analysis was conducted by using the 2ΔCt method[65] with the *gyrA* gene as a housekeeping gene to

normalize mRNA levels between different samples. The target genes in this study were *dfnA*, *baeJ*

and *acnA*.

Secondary metabolite analysis

For detection of BSMs, *B. velezensis* and *Pseudomonas* sp. were cultured in EM and CAA as

described above. After an incubation period of 24 h for *B. velezensis*, if not differentially indicated,

and 48 h for *Pseudomonas* sp., supernatants of the bacteria were collected and analyzed by UPLC

MS and UPLC qTOF MS/MS. Metabolites were identified using Agilent 1290 Infinity II coupled with

DAD detector and Mass detector (Jet Stream ESI-Q-TOF 6530) in both negative and positive mode

with the parameter set up as follows: parameters: capillary voltage: 3.5 kV; nebulizer pressure: 35

psi; drying gas: 8 l/min; drying gas temperature: 300 °C; flow rate of sheath gas: 11 l/min; sheath

gas temperature: 350 °C; fragmentor voltage: 175 V; skimmer voltage: 65 V; octopole RF: 750 V.

Accurate mass spectra were recorded in the range of m/z = 40-250. An C18 Acquity UPLC BEH

column (2.1 × 50 mm × 1.7 µm; Waters, milford, MA, USA) was used at a flow rate of 0.3 ml/min

and a temperature of 40 °C. The injection volume was 20 µl and the diode array detector (DAD)

scanned a wavelength spectrum between 190 and 600 nm. A gradient of 0.1% formic acid water

(solvent A) and acetonitrile acidified with 0.1% formic acid (solvent B) was used as a mobile phase

with a constant flow rate at 0.45 ml/min starting at 10% B and raising to 100% B in 20 min. Solvent

B was kept at 100% for 2 min before going back to the initial ratio. Secondary metabolite

quantification was performed by using UPLC–MS with UPLC (Acquity H-class, Waters) coupled to

a single quadrupole mass spectrometer (SQD mass analyzer, Waters) using a C18 column

(Acquity UPLC BEH C18 2.1 mm × 50 mm, 1.7 μm). Elution was performed at 40 °C with a

constant flow rate of 0.6 ml/min using a gradient of Acetonitrile (solvent B) and water (solvent A)

both acidified with 0.1% formic acid as follows: 2 min at 15% B followed by a gradient from 15% to

95% during 5 min and maintained at 95% up to 9.5 min before going back to initial conditions at 10

min during 2 min before next injection. Compounds were detected in both electrospray positive and

negative ion mode by setting SQD parameters as follows: cone voltage: 60V; source temperature

130 °C; desolvation temperature 400 °C, and nitrogen flow: 1000 l/h with a mass range from m/z

300 to 2048. 3D chromatograms were generated using the open-source software MzMine 2[66].

Bioguided fractionation

Pseudomonas CFS were concentrated with a C18 cartridge 'Chromafix, small' (Macherey-Nagel,

Düren, Germany). The column was conditioned with 10 ml of MeOH followed by 10 ml of milliQ

water. Then, 20 ml of supernatant flowed through the column. The metabolites were eluted with 1

ml of a solution of increasing acetonitrile/water ratio from 5:95 to 100:0 (v/v). The triggering effect of

these fractions on *Bacillus* 2H-bae production was tested in 48 wells microplate containing 1 ml of

EM medium inoculated with *B. velezensis* GA1 (OD_{600nm} = 0.1) and 4% v/v of aforementioned

Pseudomonas fractions, growing for 24 h, with shaking at 300 rpm and 30 °C. Afterward, the

production of 2H-bae was quantified compared to controls and crude supernatant.

Purification of E-PCH and PVD

PVD and E-PCH were purified in two steps. Firstly, *Pseudomonas* CFS were concentrated with a

C18 cartridge (as indicated in section Bioguided fractionation) and eluted with 2 times 2 ml of a

solution of water and ACN (15 and 30% of ACN (v/v)). Secondly, the fractions were injected on

HPLC for purification performed on an Eclipse+ C18 column (L = 150 mm, D = 3.0 mm, Particles

diameter 5 µm) (Agilent, Waldbronn, Germany). The volume injected was 100 µl. The UV-Vis

absorbance was measured with a VWD Agilent technologies 1100 series (G1314A) detector

(Agilent, Waldbronn, Germany). The lamp used was a Deuterium lamp G1314 Var Wavelength Det.

(Agilent, Waldbronn, Germany). Two wavelengths were selected: 320 nm, used for the detection of

E-PCH, and 380 nm, used for the detection of PVD. The fractions containing the PVD and E-PCH

were collected directly at the detector output. Further, the purity of the samples was verified by two

detectors, a diode array detector (DAD) 190 to 601 nm (steps: 1 nm) and a Q-TOF (tandem mass

spectrometry, quadrupole and Time of flight detector combined) (Agilent, Waldbronn, Germany).

Electrospray ionization was performed in positive mode (ESI+) (Dual AJS ESI) (Vcap = 3500 V,

Nozzle Voltage = 1000 V), with a mass range from m/z 200 to 1500. Finally, the concentration of

PVD and E-PCH were estimated by utilization of Beer-Lambert law formula, $A = \mathcal{E}lc$ (A:

absorbance; $\mathcal{E}$: molar attenuation coefficient or absorptivity of the attenuating species; l: optical

path length and c: concentration of molecule). l value for E-PCH and PVD is 1 cm while $\mathcal{E}$ is 4000

$L.mol^{-1}.cm^{-1}$ or 16000 $L.mol^{-1}.cm^{-1}$, respectively[67]. The absorbance was measured with VWR, V-

1200 Spectrophotometer, at 320 nm (pH = 8) for E-PCH and 380nm (pH = 5) for PVD[67]. Further,

the absorbance value was used for calculating the final concentration. The fragmentation pattern of

Pseudomonas sp. CMR12a PVD was obtained by UPLC MS/MS analysis of m/z = 1288.5913 ion in

positive mode with fragmentation energy at 75 V and compared to the one described in *P.*

protegens Pf-5[40].

Confrontation, white line formation and motility test

For confrontation assays on agar plates*, Bacillus* and *Pseudomonas* strains were grown overnight

in EM and CAA liquid mediums, respectively. After bacterial washing in peptone water and

adjustment of OD_{600nm} to 0.1, 5 µl of bacterial suspension was spotted at 1 mm, 5 mm and 7.5 mm

distance onto an EM agar plate. For the white line formation experiments, *B. velezensis* line was

applied with a cotton stick and 5 µl of *Pseudomonas* sp. cell suspensions were spotted at a 5 mm

distance onto CAA agar plates. Plates were incubated at 30 °C and images taken after 24 h.

Photographs were captured using CoolPix camera (NiiKKOR 60x WIDE OPTICAL ZOOM EDVR

4.3-258 mm 1:33-6.5).

MALDI-FT-ICR MS imaging

Mass spectrometry images were obtained as recently described[68] using a FT-ICR mass

spectrometer (SolariX XR 9.4T, (Bruker Daltonics, Bremen, Germany)) mass calibrated from 200

m/z to 2,300 m/z to reach a mass accuracy of 0.5 ppm. Region of interest from agar microbial

colonies was directly collected from the Petri dish and transferred onto an ITO Glass slide (Bruker,

Bremen, Germany), previously covered with double-sided conductive carbon tape. The samples

were dried under vacuum and covered with an α-cyano-4-hydroxycinnamic acid (HCCA) matrix

solution at 5 mg/mL (70 : 30 acetonitrile : water v/v). In total, 60 layers of HCCA matrix were

sprayed using the SunCollect instrument (SunChrom, Friedrichsdorf, Germany). FlexImaging 5.0

(Bruker Daltonics, Bremen, Germany) software was used for MALDI-FT-ICR MS imaging

acquisition, with a pixel step size for the surface raster set to 100 µm.

***In planta* competition**

For *in planta* studies, tomato seeds (*Solanum lycopersicum* var. Moneymaker) were sterilized in

75% ethanol with shaking for 2 min. Subsequently, ethanol was removed and seeds were added to

the 50 ml sterilization solution (8.5 ml of 15% bleach, 0.01 g of Tween 80 and 41.5 ml of sterile

ultra-pure water) and shaken for 10 min. Seeds were thereafter washed five times with water to

eliminate stock solution residues. Further, seeds were placed on square Petri dishes (5

seeds/plate) containing Hoagland solid medium (14 g/l agar, 5 ml stock 1 (EDTA 5,20 mg/l;

$FeSO_4x7H_2O$ 3,90 mg/l; H_3BO_3 1,40 mg/l; $MgSO_4x7H_2O$ 513 mg/l; $MnCl_2x4H_2O$ 0,90 mg/l,

$ZnSO_4x7H_2O$ 0,10 mg/l; $CuSO_4x5H_2O$ 0,05 mg/l; 1 ml in 50 ml stock 1, $NaMo0_4x2H_2O$ 0,02 mg/l 1

ml in 50 ml stock 1), 5 ml stock 2 (KH_2PO_4 170 mg/l; 5 ml stock 3: $KN0_3$ 316 mg/l, $Ca(NO_3)_2$ $4H_2O$

825 mg/l), pH = 6,5) and placed in the dark for three days. Afterwards, 10 seeds were inoculated

with 2 µl of overnight culture (OD_{600} = 0.1) of the appropriate strains (control) or with a mix of

Bacillus and *Pseudomonas* cells (95:5 ration) (interaction) and grown at 22 °C under a 16/8 h

night/day cycle with constant light for three days. After the incubation period, to determine bacterial

colonization levels, bacteria from roots of six plants per condition were detached from roots by

vortexing for 1 min in peptone water solution supplemented with 0.1% of Tween. Serial dilutions

were prepared and 200 µl of each were plated onto LB medium using plating beads. After 24 h of

incubation at 30 °C for *Pseudomonas* and at 42 °C for *Bacillus*, colonies were counted.

Colonization results (six plants per strain) were log-transformed and statistically analyzed. Three

independent assays were performed with six plants each for *in planta* competition assays. To

measure bacterial BSMs production *in planta*, a rectangle part (1 x 2.5 cm) of medium close to the

tomato roots was sampled. BSMs were extracted for 15 min, with 1.5 ml of acetonitrile (85%). After

centrifugation for 5 min at 4000 rpm, the supernatant was recovered for UPLC-MS analysis as

previously described.

Statistical analysis

Statistical analyses were performed using GraphPad PRISM software with Student paired T-test or

Mann-Whitney test. For multiple comparisons, one-way ANOVA and Tukey tests were used in

RStudio 1.1.423 statistical software environment (R language version 4.03)[69].

 Figure legends

 Figure 1. Stimulation of BSMs production by *B. velezensis* GA1 and enhanced anti-bacterial

 activities in response to *Pseudomonas* sp. CMR12a secreted metabolites. a. UPLC-MS

 extracted ions chromatograms (EIC) illustrating the relative abundance of ions corresponding to

 non-ribosomal metabolites produced by *B. velezensis* GA1 in CFS-supplemented (2% v/v) EM

 medium (blue) compared to un-supplemented cultures used as control (red). Red-coloured parts in

 the representation of lipopeptides and macrolactin illustrate the variable structural traits explaining

 the occurrence of naturally co-produced variants (multiple peaks) **b**. Fold increase in GA1 BSM

 production upon addition of CMR12a CFS (2% v/v) compared to un-supplemented cultures (fold

 change = 1, red line). Data were calculated based on the relative quantification of the compounds

 by UPLC-MS (peak area) in both conditions. Mean values were calculated from data obtained in

 three cultures (repeats) from two independent experiments (n = 6). Statistical significance was

 calculated using Mann–Whitney test where ''****'' represents significant difference at *P*<0.0001. **c.**

 Enhanced Anti-*Xanthomonas campestris* (**I** and **II**) and anti-*Clavibacter michiganensis* (**III** and **IV**)

 activities of GA1 extracts (cell-free culture supernatant) after growth in CMR12a CFS-

 supplemented medium (GA1+CFS) compared to control (GA1). It was assessed both on plates by

 the increase in inhibition zone around paper disc soaked with 5 µl the GA1 extracts (**I** and **III**) and in

 liquid cultures of the pathogens by reduction of growth upon addition of 4% (v/v) of GA1 extracts (**II**

 and **IV**). Data are from one representative experiment and similar results were obtained in two

 independent replicates. **d.** Antibacterial activities of extracts from GA1 WT and mutants impaired in

 production of specific BSMs. Metabolites not produced by the different mutants are illustrated with

 red boxes in the table below. All values represent means with error bars indicating SD calculated

 on data from three cultures (repeats) in two independent experiments (n = 6). Letters a to d indicate

 statistically significant differences according to one-way analysis of variance (ANOVA) and Tukey's

 HSD test (Honestly significantly different, α = 0.05). **e.** Differential expression of the *acnA* gene

encoding the amylocylicin precursor, upon supplementation with CMR12a CFS compared to GA1

un-supplemented culture. Mean and SD values, n = 6, "**" indicates statistical significance

according to Mann–Whitney test, *P*<0.01.

Figure 2: E-PCH as main *Pseudomonas* trigger of anti-bacterial activity boosted in *B.*

***velezensis* GA1. a,** Effect of GA1 culture supplementation with CFS (2% v/v) from CMR12a WT

and various mutants on dihydrobacillaene (2H-bae) production. Metabolites specifically repressed

in the different CMR12a mutants are illustrated by red boxes. Fold changes were calculated based

on relative quantification of the compounds by UPLC-MS (peak area) in treated cultures compared

to un-supplemented controls (fold change = 1, red line). Data are means and SE calculated from

three replicate cultures in two (n = 6) or three (n = 9) independent experiments and different letters

indicate statistically significant differences (ANOVA and Tukey's test, α = 0.05). **b,** Differential

production of 2H-bae after addition of 0.35 µM pure PVD, 1.4 µM pure E-PCH, 4% v/v

Pseudomonas sp. CMR12a CFS (CFS CAA), CMR12a CFS from iron supplemented culture (CFS

CAA+Fe) and different concentration of the iron-chelating agent 2,2'-dipyridyl (DIP). Data are

expressed and were statistically treated as described in **a** with n = 6 in all treatments. **c,** Dose-

dependent effect of pure PVD and E-PCH on bacillibactin and 2H-bae production. GA1 cultures

were supplemented with the indicated concentrations of HPLC-purified CMR12a siderophores.

Experiments were replicated and data statistically processed as described in **b**. **d,** Impact of the

addition of pure PVD and E-PCH on the growth of GA1 WT and its Δ*dhbC* mutant repressed in

bacillibactin synthesis. *Pseudomonas* siderophores were added at a final concentration similar to

the one obtained by adding CMR12a CFS at 4% v/v. Means and SD are from three replicates. See

Supplementary Figure 8 for detailed data and statistical significance.

Figure 3: Distance- and surfactin-dependent enhanced motility of *B. velezensis* GA1 in interaction with *Pseudomonas* CMR12a. a, GA1 motility phenotype on EM gelified medium when cultured alone (left panel) or in confrontation with CMR12a at a short distance (1 cm) (right panel). **b,** Motility pattern of GA1 or his Δ*srfaA* surfactin deficient mutant in confrontation with CMR12a at a short distance. **c,** MALDI FT-ICR MSI (Mass spectrometry Imaging) heatmaps showing spatial localization and relative abundance of ions ([M+Na]$^+$) corresponding to the C_{14} surfactin homolog (most abundant) when *B. velezensis* GA1 is in confrontation with CMR12a at increasing distances (one biological replicate).

Figure 4: Surfactin attenuates sessilin-mediated toxicity via white-line formation. a, I. Polarized inhibition of GA1 micro-colony development upon co-cultivation at close contact with CMR12a colonies on EM plates. **II.** Inhibition of GA1 cell growth in EM liquid culture supplemented with 6% v/v of CFS prepared from CMR12a wild-type or mutants repressed in the synthesis of orfamides and phenazines (Δ*ofaBC-phz*), sessilins (Δ*sesA*), sessilins and orfamides (Δ*sesA-ofaBC*), sessilins and phenazines (Δ*sesA-phz*), or of all compounds (Δ*sesA-ofaBC-phz*). Data show mean and SD calculated from two independent experiments each with three culture replicates (n = 6) and different letters indicate statistically significant differences (ANOVA and Tukey's test, α = 0.05). **b,** Growth inhibition of GA1 WT and Δ*srfaA* mutant upon delayed supplementation (added 6 h after incubation start) with CFS from CMR12a WT alone or together with pure surfactin as chemical complementation) and with CFS from the sessilin mutant (Δ*sesA*). Un-supplemented cultures of GA1 were used as control**.** Experiments were replicated and data statistically processed as described in **a**. **c,** White line formation and/or *Bacillus* inhibition observed upon confrontation of GA1 WT or the surfactin mutant Δ*srfaA* with (**I**) CMR12a or its Δ*sesA* derivative, (**II**) *P. tolaasii* CH36 or its tolaasin defective mutant Δ*tolA* and (**III**) other *Pseudomonas* CLP producers WCU-84,

SS101, BW11M1, RW10S2. CLPs produced by the individual *Pseudomonas* strains are mentioned

in the chart below. **d,** 3D representation of UPLC-MS analysis of CLPs that are present in the

white-line zone between GA1 and CMR12a (**I**). It shows the specific accumulation of sessilin and

surfactin molecular ions (one biological replicate).

Figure 5: Competitive colonization assays support the roles of BSMs in *Bacillus-*

***Pseudomonas* interaction *in planta*.** **a,** GA1 and CMR12a cell populations as recovered from

roots at 3 days post-inoculation (dpi) of tomato plantlets when inoculated alone (GA1, CMR12a) or

co-inoculated (co-inoculation). Box plots were generated based on data from three independent

assays each involving at least 4 plants per treatment (n=16). The whiskers extend to the minimum

and maximum values, and the midline indicates the median. Statistical differences between the

treatments were calculated using Mann–Whitney test and ''****'' and ''***'' represent significant

differences at *P*<0.0001 and *P*<0.001, respectively. **b,** *In planta* (3 dpi on tomato roots) relative

expression of the *dfnA, baeJ* and *acnA* genes responsible for the synthesis of respectively

(oxy)difficidin, 2H-bae and amylocyclicin. Graphs show the mean and SD calculated from three

biological replicates (n = 3) each involving six plants. Fold change = 1 as red line corresponds to

gene expressions in GA1 inoculated alone on roots used as control conditions. Statistical

comparison between data in co-colonization setting and control conditions was performed based on

T-test (*, *P*<0.05; **, *P*<0.01; ***, *P*<0.001; ****, *P*<0.0001). **c.** UPLC-MS EIC illustrating relative *in*

planta production of sessilins and surfactins by monocultures of *B. velezensis* GA1 (GA1) and co-

cultures of wild-types (GA1+CMR12a) and *B. velezensis* GA1 and *Pseudomonas* sp. CMR12a

impaired in sessilins production (GA1+Δ*sesA*). **d,** Cell populations recovered at 3 dpi for GA1 WT

(GA1) or the surfactins impaired mutant (Δ*srfaA*) co-inoculated with CMR12a WT (CMR12a) or its

sessilins KO mutant (Δ*sesA*). See **a** for replicates and statistics (**, *P*<0.01).

References

1.   Nayfach, S. *et al.* A genomic catalog of Earth's microbiomes. *Nat. Biotechnol.* (2020)
doi:10.1038/s41587-020-0718-6.

2.   Fierer, N. Embracing the unknown: disentangling the complexities of the soil microbiome.
Nat. Rev. Microbiol. **15**, 579–590 (2017).

3.   Schmidt, R., Ulanova, D., Wick, L. Y., Bode, H. B. & Garbeva, P. Microbe-driven chemical
ecology: past, present and future. *ISME Journal.* **13,** 2656–2663 (2019).

4.   Tyc, O., Song, C., Dickschat, J. S., Vos, M. & Garbeva, P. The ecological role of volatile and
soluble secondary metabolites produced by soil bacteria. *Trends Microbiol.* **25**, 280–292
(2017).

5.   Bernal, P., Llamas, M. A. & Filloux, A. Type VI secretion systems in plant-associated
bacteria. *Environ. Microbiol.* **20**, 1–15 (2018).

6.   Abreu, N. A. & Taga, M. E. Decoding molecular interactions in microbial communitiesa.
FEMS Microbiol. Rev. **40**, 648–663 (2016).

7.   Andrić, S., Meyer, T. & Ongena, M. *Bacillus* responses to plant-sssociated fungal and
bacterial communities. *Front. Microbiol.* **11**, (2020).

8.   Mendes, R., Garbeva, P. & Raaijmakers, J. M. The rhizosphere microbiome: significance of
plant beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiol.*
Rev. **37**, 634–663 (2013).

9.   Müller, D. B., Vogel, C., Bai, Y. & Vorholt, J. A. The plant microbiota: systems-level insights
and perspectives. *Annu. Rev. Genet.* **50**, 211–234 (2016).

10. Penha, R. O., Vandenberghe, L. P. S., Faulds, C., Soccol, V. T. & Soccol, C. R. *Bacillus*
lipopeptides as powerful pest control agents for a more sustainable and healthy agriculture:

recent studies and innovations. *Planta* **251**, 1–15 (2020).

11.   Grubbs, K. J. *et al.* Large-Scale bioinformatics analysis of *Bacillus* genomes uncovers

677   conserved roles of natural products in bacterial physiology. *mSystems* **2**, 1–18 (2017).

678   12.   Harwood, C. R., Mouillon, J.-M. M., Pohl, S. & Arnau, J. Secondary metabolite production

679   and the safety of industrially important members of the *Bacillus subtilis* group. *FEMS*

680   *Microbiol. Rev.* **42**, 721–738 (2018).

681   13.   Ye, M. *et al.* Characteristics and application of a novel species of *Bacillus* : *Bacillus*

682   *velezensis*. *ACS Chem. Biol.* **13**, 500–505 (2018).

683   14.   Rabbee, M. *et al.* Bacillus velezensis: A valuable member of bioactive molecules within plant

684   microbiomes. *Molecules* **24**, 1–13 (2019).

685   15.   Pieterse, C. M. J. *et al.* Induced systemic resistance by beneficial microbes. *Annu. Rev.*

686   *Phytopathol.* **52**, 347–375 (2014).

687   16.   Köhl, J., Kolnaar, R. & Ravensberg, W. J. Mode of action of microbial biological control

688   agents against plant diseases: relevance beyond efficacy. *Front. Plant Sci.* **10**, 845 (2019).

689   17.   Traxler, M. F. & Kolter, R. Natural products in soil microbe interactions and evolution. *Nat.*

690   *Prod. Rep.* **32**, 956–970 (2015).

691   18.   Raaijmakers, J. M. & Mazzola, M. Diversity and natural functions of antibiotics produced by

692   beneficial and plant pathogenic bacteria. *Annu. Rev. Phytopathol.* **50**, 403–424 (2012).

693   19.   Li, Y. & Rebuffat, S. The manifold roles of microbial ribosomal peptide-based natural

694   products in physiology and ecology. *J. Biol. Chem.* **295**, 34–54 (2020).

695   20.   Raaijmakers, J. M., de Bruijn, I., Nybroe, O. & Ongena, M. Natural functions of lipopeptides

696   from *Bacillus* and *Pseudomonas*: More than surfactants and antibiotics. *FEMS Microbiol.*

697   *Rev.* **34**, 1037–1062 (2010).

698   21.   Blin, K. *et al.* antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline.

699   *Nucleic Acids Res.* **47**, W81–W87 (2019).

700 22. Nihorimbere, V. *et al.* Impact of rhizosphere factors on cyclic lipopeptide signature from the

701    plant beneficial strain *Bacillus amyloliquefaciens* S499. *FEMS Microbiol. Ecol.* **79**, 176–191

702    (2012).

703 23. D'aes, J. *et al.* To settle or to move? The interplay between two classes of cyclic lipopeptides

704    in the biocontrol strain *Pseudomonas* CMR12a. *Environ. Microbiol.* **16**, 2282–2300 (2014).

705 24. Hua, G. K. H. & Höfte, M. The involvement of phenazines and cyclic lipopeptide sessilin in

706    biocontrol of *Rhizoctonia* root rot on bean (*Phaseolus vulgaris*) by *Pseudomonas* sp.

707    CMR12a is influenced by substrate composition. *Plant Soil* **388**, 243–253 (2015).

708 25. Perneel, M. *et al.* Characterization of CMR5c and CMR12a, novel fluorescent *Pseudomonas*

709    strains from the cocoyam rhizosphere with biocontrol activity. *J. Appl. Microbiol.* **103**, 1007–

710    1020 (2007).

711 26. Olorunleke, F. E. *et al.* Coregulation of the cyclic lipopeptides orfamide and sessilin in the

712    biocontrol strain *Pseudomonas* sp. CMR12a. *Microbiologyopen* **6**, 1–12 (2017).

713 27. Olorunleke, F. E., Hua, G. K. H., Kieu, N. P., Ma, Z. & Höfte, M. Interplay between

714    orfamides, sessilins and phenazines in the control of *Rhizoctonia* diseases by *Pseudomonas*

715    sp. CMR12a. *Environ. Microbiol. Rep.* **7**, 774–781 (2015).

716 28. Geudens, N. & Martins, J. C. Cyclic lipodepsipeptides from *Pseudomonas* spp. - Biological

717    Swiss-Army knives. *Front. Microbiol.* **9**, 1–18 (2018).

718 29. Omoboye, O. O. *et al. Pseudomonas* cyclic lipopeptides suppress the rice blast fungus

719    *Magnaporthe oryzae* by induced resistance and direct antagonism. *Front. Plant Sci.* **10**, 901

720    (2019).

721 30. Mansfield, J. *et al.* Top 10 plant pathogenic bacteria in molecular plant pathology. *Mol. Plant*

722    *Pathol.* **13**, 614–629 (2012).

723 31. Molinatto, G. *et al.* Complete genome sequence of *Bacillus amyloliquefaciens* subsp.

724    *plantarum* S499, a rhizobacterium that triggers plant defences and inhibits fungal

phytopathogens. *J. Biotechnol.* **238**, 56–59 (2016).

32. Fan, B. *et al.* *Bacillus velezensis* FZB42 in 2018: The gram-positive model strain for plant growth promotion and biocontrol. *Front. Microbiol.* **9**, 3389 (2018).

33. Pandin, C. *et al.* Complete genome sequence of *Bacillus velezensis* QST713: A biocontrol agent that protects *Agaricus bisporus* crops against the green mould disease. *J. Biotechnol.* **278**, 10–19 (2018).

34. Scholz, R. *et al.* Amylocyclicin, a novel circular bacteriocin produced by *Bacillus amyloliquefaciens* FZB42. *J. Bacteriol.* **196**, 1842–1852 (2014).

35. Youard, Z. A., Mislin, G. L. A., Majcherczyk, P. A., Schalk, I. J. & Reimmann, C. *Pseudomonas fluorescens* CHA0 produces enantio-pyochelin, the optical antipode of the *Pseudomonas aeruginosa* siderophore pyochelin. *J. Biol. Chem.* **282**, 35546–35553 (2007).

36. Grandchamp, G. M. *et al.* Pirated siderophores promote sporulation in *Bacillus subtilis*. *Appl. Environ. Microbiol.* **83**, 1–17 (2017).

37. Miethke, M. *et al.* Ferri-bacillibactin uptake and hydrolysis in *Bacillus subtilis*. *Mol. Microbiol.* **61**, 1413–1427 (2006).

38. Adler, C. *et al.* Catecholate siderophores protect bacteria from pyochelin toxicity. *PLoS One* **7**, (2012).

39. Trottmann, F., Franke, J., Ishida, K., García-Altares, M. & Hertweck, C. A pair of bacterial siderophores releases and traps an intercellular signal molecule: An unusual case of natural nitrone bioconjugation. *Angew. Chemie* **131**, 206–210 (2019).

40. Hartney, S. L. *et al.* Ferric-pyoverdine recognition by Fpv outer membrane proteins of *Pseudomonas protegens* Pf-5. *J. Bacteriol.* **195**, 765–776 (2013).

41. Rokni-Zadeh, H. *et al.* Genetic and functional characterization of cyclic lipopeptide white-line-inducing principle (WLIP) production by rice rhizosphere isolate *Pseudomonas putida* RW10S2. *Appl. Environ. Microbiol.* **78**, 4826–4834 (2012).

42.   van Gestel, J., Vlamakis, H. & Kolter, R. From cell differentiation to cell collectives: *Bacillus subtilis* uses division of labor to migrate. *PLoS Biol.* **13**, 1–29 (2015).

43.   Munsch, P. & Alatossava, T. The white-line-in-agar test is not specific for the two cultivated mushroom associated pseudomonads, *Pseudomonas tolaasii* and *Pseudomonas 'reactans'*. *Microbiol. Res.* **157**, 7–11 (2002).

44.   Salari, F. *et al.* Draft Genome Sequence of *Pseudomonas aeruginosa* strain LMG 1272, an atypical white line reaction producer. *Microbiol. Resour. Announc.* **9**, (2020).

45.   Xu, Z. *et al.* Antibiotic bacillomycin D affects iron acquisition and biofilm formation in *Bacillus velezensis* through a Btr-mediated FeuABC-dependent pathway. *Cell Rep.* **29**, 1192-1202.e5 (2019).

46.   Powers, M. J., Sanabria-Valentín, E., Bowers, A. A. & Shank, E. A. Inhibition of cell differentiation in *Bacillus subtilis* by *Pseudomonas protegens*. *J. Bacteriol.* **197**, 2129–2138 (2015).

47.   Molina-Santiago, C. *et al.* The extracellular matrix protects *Bacillus subtilis* colonies from *Pseudomonas* invasion and modulates plant co-colonization. *Nat. Commun.* **10**, (2019).

48.   Liu, Y., Kyle, S. & Straight, P. D. Antibiotic stimulation of a *Bacillus subtilis* migratory response. *mSphere* **3**, 1–13 (2018).

49.   Rosenberg, G. *et al.* Not so simple, not so subtle: The interspecies competition between *Bacillus simplex* and *Bacillus subtilis* and its impact on the evolution of biofilms. *npj Biofilms Microbiomes* **2**, 15027 (2016).

50.   Straight, P. D., Willey, J. M. & Kolter, R. Interactions between *Streptomyces coelicolor* and *Bacillus subtilis*: Role of surfactants in raising aerial structures. *J. Bacteriol.* **188**, 4918–4925 (2006).

51.   Lee, N. *et al.* Iron competition triggers antibiotic biosynthesis in *Streptomyces coelicolor* during coculture with *Myxococcus xanthus*. *ISME J.* **14**, 1111–1124 (2020)

52. Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host
interactions. *Nat. Rev. Microbiol.* **18**, 152–163 (2020).

53. Niehus, R., Picot, A., Oliveira, N. M., Mitri, S. & Foster, K. R. The evolution of siderophore
production as a competitive trait. *Evolution (N. Y).* **71**, 1443–1455 (2017).

54. Hibbing, M. E., Fuqua, C., Parsek, M. R. & Peterson, S. B. Bacterial competition: Surviving
and thriving in the microbial jungle. *Nature Reviews Microbiology* **8** 15–25 (2010).

55. Gu, S. *et al.* Competition for iron drives phytopathogen control by natural rhizosphere
microbiomes. *Nat. Microbiol.* **5**, 1002–1010 (2020).

56. Youard, Z. A., Wenner, N. & Reimmann, C. Iron acquisition with the natural siderophore
enantiomers pyochelin and enantio-pyochelin in *Pseudomonas* species. *BioMetals* **24**, 513–
522 (2011).

57. Cornelis, P. Iron uptake and metabolism in pseudomonads. *Appl. Microbiol. Biotechnol.* **86**,
1637–1645 (2010).

58. Ronnebaum, T. A. & Lamb, A. L. Nonribosomal peptides for iron acquisition: pyochelin
biosynthesis as a case study. *Current Opinion in Structural Biology* **53**, 1–11 (2018).

59. Esmaeel, Q. *et al. Burkholderia* genome mining for nonribosomal peptide synthetases
reveals a great potential for novel siderophores and lipopeptides synthesis.
Microbiologyopen **5**, 512–526 (2016).

60. Seipke, R. F. *et al.* The plant pathogen *Streptomyces scabies* 87-22 has a functional
pyochelin biosynthetic pathway that is regulated by TetR- and AfsR-family proteins.
Microbiology **157**, 2681–2693 (2011).

61. Inahashi, Y. *et al.* Watasemycin biosynthesis in *Streptomyces venezuelae*: thiazoline C-
methylation by a type B radical-SAM methylase homologue. *Chem. Sci.* **8**, 2823–2831
(2017).

62. Jarmer, H., Berka, R., Knudsen, S. & Saxild, H. H. Transcriptome analysis documents

induced competence of *Bacillus subtilis* during nitrogen limiting conditions. *FEMS Microbiol.*

Lett. **206**, 197–200 (2002).

63.   Martínez-García, E. & de Lorenzo, V. Engineering multiple genomic deletions in Gram-

negative bacteria: analysis of the multi-resistant antibiotic profile of *Pseudomonas putida*

KT2440. *Environ. Microbiol.* **13**, 2702–2716 (2011).

64.   Vacheron, J. *et al.* T6SS contributes to gut microbiome invasion and killing of an herbivorous

pest insect by plant-beneficial *Pseudomonas protegens*. *ISME J.* **13**, 1318–1329 (2019).

65.   Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time

quantitative PCR and the 2-ΔΔCT method. *Methods* **25**, 402–408 (2001).

66.   Pluskal, T., Castillo, S., Villar-Briones, A. & Orešič, M. MZmine 2: Modular framework for

processing, visualizing, and analyzing mass spectrometry-based molecular profile data.

BMC Bioinformatics **11**, (2010).

67.   Hoegy, F., Mislin, G. L. A. & Schalk, I. J. Pyoverdine and pyochelin measurements. *Methods*

Mol. Biol. **1149**, 293–301 (2014).

68.   Kune, C. *et al.* Rapid visualization of chemically related compounds using kendrick mass

defect as a filter in mass spectrometry imaging. *Anal. Chem.* **91**, 13112–13118 (2019).

69.   R Core Team (2020). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.r-project.org/ (2020).

Acknowledgments

We gratefully acknowledge Sébastien Rigali, Alexandre Jousset and Loïc Ongena for critically

reading the manuscript. We thank C. Keel for the kind gift of strains and J. Vacheron for the very

helpful indications on *Pseudomonas* mutagenesis. This work was supported by the EU Interreg V

France-Wallonie-Vlaanderen portfolio SmartBiocontrol (Bioprotect and Bioscreen projects, avec le

soutien du Fonds européen de développement régional - Met steun van het Europees Fonds voor

Regionale Ontwikkeling), by the European Union Horizon 2020 research and innovation program

under grant agreement No. 731077 and by the EOS project ID 30650620 from the FWO/F.R.S.-

FNRS. The MALDI FT-ICR SolariX XR was funded by FEDER BIOMED HUB Technology Support

(number 2.2.1/996). AA is recipient of a F.R.I.A. fellowship (Formation à la Recherche dans

l'Industrie et l'Agriculture) and MO is senior research associate at the F.R.S.-F.N.R.S.

Author contributions

SA, TM, AR and AA performed most of the co-culture and *in planta* experiments. SA and TM

performed most of molecular biology experiments with help of GH and SS for mutant generation

and of SS for transcriptomics. TM and GH did genome mining. AR and AA were involved in all

aspects of metabolomics using UPLC-MS. Data analysis was done by SA, TM, AA and AR. AM, AA

and EDP performed the MALDI FT-ICR experiments and analyzed the data. MH and RDM provided

Pseudomonas strains/mutants and also supported the study by providing intellectual input. SA, TM

and MO mainly wrote the manuscript. All of the authors commented on the manuscript and

contributed to the final form. MO supervised the study.

Competing interests

The authors declare no competing interests.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5