Chelator sensing and lipopeptide interplay mediates molecular interspecies

interactions between soil bacilli and pseudomonads

Sofija Andric´[1*], Thibault Meyer[1,2*], Augustin Rigolet[1], Anthony Argüelles Arias[1], Sébastien Steels[1], Grégory

Hoff[1#], Monica Höfte[3], René De Mot[4], Andrea McCann[5], Edwin De Pauw[5] and Marc Ongena[1]

[1]Microbial Processes and Interactions Laboratory, Terra Teaching and Research Center, Gembloux Agro-Bio Tech,

University of Liège, Gembloux, Belgium

[2]UMR Ecologie Microbienne, F-69622, University of Lyon, Université Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup,

Villeurbanne, France

[3]Laboratory of Phytopathology, Department of Plants and Crops, Faculty of Bioscience engineering, Ghent University,

Gent, Belgium

[4] Centre of Microbial and Plant Genetics, Faculty of Bioscience Engineering, University of Leuven, Heverlee, Belgium

[5] Mass Spectrometry Laboratory, , MolSys Research Unit, Department of Chemistry, University of Liège, Belgium

[#]Current address: Ecology and Biodiversity, Department of Biology, Utrecht University, Padualaan 8, 3584 CH, Utrecht,

The Netherlands

* Equal contribution

Abstract

Some bacterial species are important members of the rhizosphere microbiome and confer protection to the

host plant against pathogens. However, our knowledge is still limited about the multitrophic interactions

determining the ecological fitness of these biocontrol bacteria in their highly competitive natural niche. In

this work, we have investigated the molecular mechanisms underlying interactions between *B. velezensis,*

considered as model plant-associated and beneficial species in the *Bacillus* genus, and *Pseudomonas* as a

rhizosphere-dwelling competitor. Our data show that *B. velezensis* boosts its arsenal of specialized

antibacterials upon the perception of the secondary siderophore pyochelin produced by phylogenetically

distinct pseudomonads and some other genera. We postulate that *B. velezensis* has developed some

chelator sensing systems to learn about the identity of its surrounding competitors. Illustrating the

multifaceted molecular response of *Bacillus*, surfactin is another crucial component of the secondary

metabolome mobilized in interbacteria competition. Its accumulation not only enhances motility capability

but, unexpectedly, this lipopeptide also acts as a chemical trap that reduce the toxicity of other lipopeptides

released by *Pseudomonas* challengers. This favors the persistence of *Bacillus* populations upon competitive

root colonization. Our work thus highlights new ecological roles for bacterial secondary metabolites acting

as key drivers of social interactions.

INTRODUCTION

Soil is one of the richest ecosystems in terms of microbial diversity and abundance[1]. However, the scarcity

of resources makes it one of the most privileged environment for competitive interspecies interactions [2,3].

A subset of the bulk soil microbes has evolved to dwell in the rhizosphere compartment surrounding roots

due to continued nutrient-enriched exudation from the plant. Compared to bulk soil, microbial warfare in

the rhizosphere is presumably even more intense as the habitat is spatially restricted and more densely

populated[3]. Besides rivalry for nutrients (exploitative), interference competition is considered as a key

factor driving microbial interactions and community assembly. It can involve signal interference or toxins

deployed by contact-dependent delivery systems[4,5] but is mainly mediated at distance through the emission

of various molecular weapons. The molecular basis of interference interactions and their phenotypic

outcomes between diverse soil bacterial species have been amply investigated in the last decade[6,7].

Like some genera of the Proteobacteria, Actinobacteria, and Bacteroidetes taxa, bacilli belonging to the *B.*

subtilis complex are ubiquitous members of the rhizosphere microbiome[2,8,9]. Among these species, *B.*

velezensis has emerged as a model for plant-associated bacilli and displays strong potential as biocontrol

agents reducing diseases caused by phytopathogens[10]. *B. velezensis* distinguish from other species of the *B.*

subtilis group regarding their richness in biosynthetic gene clusters (BGCs, representing up to 13% of the

whole genome) responsible for the synthesis of bioactive secondary metabolites (BSMs) [11,12]. This

chemically-diverse secondary metabolome includes volatiles, terpenes, non-ribosomal (NR) dipeptides,

cyclic lipopeptides (CLPs) and polyketides (PKs), but also ribosomally synthesized lantibiotics and larger

bacteriocins (RiPP) [13,14]. BSMs are involved in the biocontrol activity via direct inhibition of infectious

microbes and/or via stimulation of the plant immune system[15,16]. From an ecological viewpoint, BSMs also

contribute to competitiveness in the rhizosphere niche thanks to multiple and complementary functions as

drivers of developmental traits, as antimicrobials, or as signals initiating cross-talk with the host plant[17–19 20–

26].

Mostly guided by practical concerns for use as biocontrol agents, research on BSMs has mainly focused on

the characterization of their biological activities. However, the impact of environmental factors that may

modulate their expression under natural conditions has been much poorly investigated. It includes

interactions with other organisms sharing the niche. Some recent reports illustrate how soil bacilli may

adapt their behavior upon sensing bacterial competitors but almost exclusively focusing on developmental

traits (sporulation, biofilm formation, or motility)[6]. Unlike other genera such as *Streptomyces*, it remains

largely unknown to what extent bacilli in general and *B. velezensis* in particular, may modulate the

expression of their secondary metabolome upon interaction with other bacteria[2,8,9,27]. Through this work,

we wanted to investigate the molecular outcomes of interspecies interactions that *B. velezensis* may

engage. We selected *Pseudomonas* as challenger considering that species of this genus are also highly

competitive and commonly encountered in rhizosphere microbiomes[8]. We performed experiments under

nutritional conditions mimicking the oligotrophic rhizosphere environment and used contact-independent

settings for pairwise interaction which also obviously best reflect the real situation in the soil. Our data

revealed that the two bacteria initiate a multifaceted interactions mostly mediated by the non-ribosomally

synthesized components of their secondary metabolome. We pointed out unsuspected roles for some of

these BSMs in the interaction context. Beyond its role as a metal chelator, the *Pseudomonas* secondary

siderophore pyochelin acts as a signal triggering dual production of PKs and RiPP in *Bacillus* while specific

lipopeptides modulate the inhibitory interaction between the two species. This resulted in marked

phenotypic changes in *B. velezensis* such as higher antibacterial potential, enhanced motility and protective

effect via chemical trapping. We also illustrate the relevance of these outcomes in the context of

competitive root colonization.

RESULTS

***B. velezensis* modulates its secondary metabolome and boosts antibacterial activity upon sensing**
***Pseudomonas* metabolites**

In this study, we used *B. velezensis* strain GA1 as BSM-rich and genetically amenable isolate representative
of the species. Genome mining with AntiSMASH 5.0 [28] confirmed the presence of all gene clusters necessary
for the biosynthesis of diverse known BSMs typically formed by this bacterium (Supplementary Table 1).
Based on the exact mass and absence of the corresponding peaks in deletion mutants, most of the
predicted non-ribosomal (NR) secondary metabolites were identified in cell-free crude supernatants via
optimized UPLC-MS (Supplementary Fig. 1). It includes the whole set of cyclic lipopeptides (CLPs of the
surfactin, fengycin and iturin families) and polyketides (the PKs difficidin, macrolactin and bacillaene) with
their multiple co-produced structural variants, as well as the siderophore bacillibactin. We verified that all
these compounds are readily formed in the so-called exudate-mimicking (EM) medium reflecting the
specific content in major carbon sources typically released by roots of Solanaceae (such as tomato) plants[29].
In addition to these NR products, genes encoding RiPPs such as amylocyclicin and amylolysin are also
present in GA1 but these compounds could not be reliably detected in culture broths. The NR dipeptide
bacilysin was also predicted but not detected. We selected as the main interaction partner the plant-
associated *Pseudomonas* sp. strain CMR12a based on its biocontrol potential and its production of multiple
secondary metabolites [30–34]. Genome mining confirmed the potential of CMR12a to synthesize a range of
BSMs, including antimicrobial phenazines, the siderophores pyoverdin (structure confirmation in
Supplementary Fig. 2) and (enantio-)pyochelin as well as two structurally distinct CLPs, sessilin and orfamide
(Supplementary Table 1). In contrast to *Bacillus*, the capacity to co-produce two different CLPs is a quite rare
trait for non-phytopathogenic pseudomonads and represented an additional criterion for selecting strain
CMR12a for this study[30,34–36]. In the case of CMR12a, all these compounds were detected in EM culture
broth but most of them are more efficiently produced upon growth in Casamino Acids medium (CAA)
commonly used for *Pseudomonas* cultivation (Supplementary Fig. 1).

Our prime objective was to evaluate the intrinsic potential of *B. velezensis* to react to the perception of
Pseudomonas metabolites in an experimental setting avoiding interferences due to diffusion constraints in a
semi-solid matrix or due to the formation of impermeable biofilm structures. The first assays were
performed by growing GA1 in agitated liquid EM medium supplemented or not with (sterile) BSM-
containing spent medium of CAA-grown CMR12a (CFS, cell-free supernatant). At low dose (2% (v/v)), the

addition of this CFS extract led to a marked increase in the production of some GA1 NR metabolites.

Significantly higher amounts were measured for surfactin, bacillaene and its dehydrated variant, difficidin

and its oxidized form, and bacillibactin (Fig. 1a, b) but not for fengycin, iturin nor macrolactin (Fig 1a).

Such boost in BSM synthesis triggered by *Pseudomonas* CFS was associated with an increase in the

antibacterial activity of the corresponding extracts when tested for growth inhibition of *Xanthomonas*

campestris and *Clavibacter michiganensis* used respectively as representative of Gram-negative and Gram-

positive plant pathogenic bacteria of agronomical importance[37] (Fig. 1c). Since most of the BSMs are not

commercially available and our attempt to purify PKs failed due to chemical instability, we could not use

individual compounds for their specific involvement in bacterial inhibition. As an alternative, we generated

and tested a range of GA1 knockout mutants including the Δ*sfp* derivative specifically repressed in 4'-

phosphopantetheinyl transferase essential for the proper functioning of the PK and NRP biosynthesis

machinery. Full loss of anti-*Xanthomonas* activity in Δ*sfp* extracts indicated a key role for NR BSMs and ruled

out the possible involvement of other chemicals known for their antibacterial activity such as bacilysin or

RiPPs (Fig. 1d). Loss of function of mutants specifically repressed in the synthesis of individual compounds

pointed out the key role of (oxy)difficidin and to a lower extent of (dihydro)bacillaene in *Xanthomonas*

inhibition (Fig. 1d). These two polyketides are also responsible for GA1 inhibitory activity toward other

important bacterial phytopathogens such as *Pectobacterium carotovorum*, *Agrobacterium tumefaciens* and

Rhodococcus fasciens but are not involved in the inhibition of plant pathogenic *Pseudomonas* species for

which bacilysin may be the active metabolite (Supplementary Fig. 3). However, as illustrated below in the

next sections, *B. velezensis* does not display significant toxicity against CMR12a and other non-pathogenic

soil *Pseudomonas* isolates tested here. Stimulation of PKs synthesis upon sensing CMR12a is not specific to

GA1 and was also observed in other *B. velezensis* strains such as S499, FZB42 and QST713 with well-known

biocontrol potential[38-40] (Supplementary Fig. 4).

In contrast to *Xanthomonas*, the enhanced antibiotic activity against *Clavibacter* is not mediated by NR

products as shown by the fully conserved activity in the Δ*sfp* mutant (Fig. 1d). We, therefore, suspected

from genomic data that some RiPPs such as amylocyclicin could be involved in inhibition. This hypothesis

was supported by the 80% reduction in antibiotic potential observed for the Δ*acnA* mutant knocked out for

the corresponding biosynthesis gene (Fig. 1d). Besides, RT-qPCR data revealed a highly induced expression

of the *acnA* in GA1 cells upon supplementation with CMR12a CFS (Fig. 1e) but we were not able to provide

evidence for higher accumulation of the mature peptide in the medium. Enhanced expression of the *acnA*
gene in presence of *Pseudomonas* products was also observed for strain S499 (Supplementary Fig. 5).

Pyochelin acts as a signal sensed by *Bacillus* to stimulate polyketide production

We next wanted to identify the signaling molecules secreted by *Pseudomonas* that are sensed by *Bacillus*
cells and lead to improved BSMs production. For that purpose, we used dihydrobacillaene (2H-bae) as an
indicator of the *Bacillus* response because it represents the most consistent and highly boosted polyketide
species. We first compared the triggering potential of CFS obtained from knockout mutants of CMR12a
specifically lacking the different metabolites identified (Supplementary Fig. 1). Only extracts from mutants
impaired in the production of siderophores and more specifically pyochelin were significantly affected in
PKs-inducing potential (Fig. 2a). Possible involvement of this compound was supported by the drastic
reduction in the activity of CFS prepared from CMR12a culture in CAA medium supplemented with Fe^{3+}
where siderophore expression is repressed (Supplementary Fig. 6). We also performed bioactivity-guided
fractionation and data showed that only extracts containing pyoverdine and/or pyochelin displayed
consistent PKs-triggering activity (Supplementary Fig. 7). HPLC-purified compounds were also tested
independently at a concentration similar to the one used in CFS CAA extract and it revealed a much higher
PK-triggering activity for pyochelin compared to the main pyoverdine isoform (Fig. 2b). Dose-dependent
assays further indicated that supplementation with pyoverdine, as strong chelator[41], caused some iron
limitation in the medium which is sensed by GA1. It is reflected first by the marked increase in production of
the siderophore bacillibactin in GA1 wild-type (Fig. 2c) and by the reduced growth of a Δ*dhbC* mutant
repressed in bacillibactin synthesis upon pyoverdin addition (Fig. 2d, Supplementary Fig. 8 for statistics).
This last result indicates that pyoverdine in its ferric form cannot be taken up by GA1 despite the presence
of several transporters for exogenous siderophores in *B. velezensis* similar to those identified in *B.*
subtilis[42,43] based on genome comparison (Supplementary Table 2). We, therefore, assumed that iron stress
mediated by pyoverdine only induces a rather limited boost in PKs production. We validate that such
response is not due to iron starvation by supplementing GA1 culture with increasing doses of the 2,2'-
dipyridyl chemical chelator that cannot be taken up by Bacillus cells (Fig. 2b). By contrast, the addition of
pyochelin with a much lower affinity for iron does not activate bacillibactin synthesis (Fig. 2c) and does not
affect Δ*dhbC* growth at the concentrations used (Fig. 2d). We concluded that the activity of this compound
referred to as secondary siderophore is not related to iron-stress. If internalized, pyochelin can cause
oxidative stress and damage in other bacteria as reported for *E. coli* [44,45] . However, the absence of toxicity

toward GA1 indicates that pyochelin is not taken up by *Bacillus* cells and thus clearly acts as a signal

molecule perceived at the cell surface. PKS boost also occurred upon addition of CFS obtained from other

Pseudomonas isolates such as *P. protegens* Pf-5[46] and *P. aeruginosa* PAO1 [47] producing pyochelin-type

siderophores. However, PKS stimulation was also observed in response to *P. tolaasi* CH36[48] and

Pseudomonas sp*.* A214[49] that do not form pyochelin indicating that some other unidentified BSMs may act

as triggers in other strains (Supplementary Figure 9).

Enhanced motility as distance-dependent and surfactin-mediated response of *Bacillus*

Surfactin production is also stimulated by CMR12a CFS and by pure pyochelin (Fig. 1a and Supplementary

Fig. 10). Based on the mutant analysis, this multifunctional CLP does not contribute to the antibacterial

potential of *B. velezensis* (Fig 1d and Supplementary Fig. 3) but is notably involved in developmental

processes of multicellular communities such as biofilm formation and motility[50]. We, therefore, wanted to

test a possible impact of *Pseudomonas* on the motile phenotype of *B. velezensis* upon co-cultivation on

plates. We observed distance-dependent enhanced motility on gelified medium containing high agar

concentrations (1.5% m/v) which phenotypically resembles the sliding-type of motility illustrated by typical

"van Gogh bundles"[50] (Fig. 3a). This migration pattern is flagellum independent but depends on multiple

factors including the synthesis of surfactin which reduces friction at the cell-substrate interface[50]. We thus

suspected such enhanced motility to derive from an increased formation of the lipopeptide. This was

supported by the almost full loss of migration of the Δ*srfaA* mutant in these interaction conditions (Fig. 3b).

Moreover, spatial mapping via Maldi-FT-ICR MS imaging confirmed a higher accumulation of surfactin ions

in the interaction zone and around the *Bacillus* colony when it is growing at a short or intermediate distance

from the *Pseudomonas* challenger compared to the largest distance where the motile phenotype is much

less visible (Fig. 3c). These data indicate that *Bacillus* cells in the microcolony perceive some soluble signal

diffusing from the *Pseudomonas* colony to a limited distance.

Interplay between CLPs drives antagonistic interactions

Furthermore, we observed that, besides modulating secondary metabolite synthesis, confrontation with

Pseudomonas may also lead to some antagonistic outcomes. GA1 growth as planktonic cells is slightly

inhibited upon supplementation of the medium with 2% v/v CMR12a CFS but this inhibition is much marked

at higher doses (Supplementary Fig. 11). To identify the *Pseudomonas* compound retaining such antibiotic

activity, we tested the effect of CFS from various CMR12a mutants impaired in the synthesis of lipopeptides

and/or phenazines. Even if some contribution of other compounds cannot be ruled out, it revealed that the

CLP sessilin is mainly responsible for toxic activity toward GA1 grown in liquid cultures but also when the

two bacteria are grown at close proximity on gelified EM medium (Fig 4a). Nevertheless, we observed that

the sessilin-mediated inhibitory effect is markedly reduced by delaying CFS supplementation at 6h of culture

instead of adding at the beginning of incubation (Fig 4b). This suggested that some early secreted *Bacillus*

compounds may counteract the toxic effect of sessilin. We hypothesized that surfactin can play this role as

it is the first detectable BSM to accumulate in significant amounts in the medium at early growth. We tested

the surfactin-deficient mutant in the same conditions and observed that its growth is still strongly affected

indicating that no other GA1 compound may be involved in toxicity alleviation. Chemical complementation

with purified surfactin allowed to restore growth to a large extent, providing further evidence for a

protective role of the surfactin lipopeptide (Fig. 4b). Such sessilin-dependent inhibition also occurred when

bacteria were confronted on solid CAA medium favoring *Pseudomonas* BSM production (Fig. 4c-I). In these

conditions, the formation of a white precipitate in the interaction zone was observed with CMR12a wild

type but not when GA1 was confronted with the Δ*sesA* mutant (Fig. 4c). UPLC-MS analysis of ethanol

extracts from this white-line area confirmed the presence of sessilin ions but also revealed an accumulation

of surfactin from GA1 in the confrontation zone (Fig. 4d). The involvement of surfactin in precipitate

formation was confirmed by the absence of this white-line upon testing the Δ*srfaA* mutant of GA1 (Fig. 4c).

The loss of surfactin production and white-line formation was associated with a higher sensitivity of the

Bacillus colony to the toxin secreted by *Pseudomonas*. Altogether, these data indicate that surfactin acts as

a chemical trap and inactivates sessilin via co-aggregation into insoluble complexes.

A similar CLP-dependent antagonistic interaction and white-line formation were observed upon co-

cultivation of GA1 with *P. tolaasii* strain CH36 producing tolaasin (Fig. 4c), a CLP structurally very similar to

sessilin (only differing by a single amino acid residue, Supplementary Fig. 12). However, this chemical

aggregation is quite specific regarding the type of CLP involved since it was not visible upon the interaction

of GA1 with other *Pseudomonas* strains forming different CLP structural groups that are not toxic for

Bacillus (Fig. 4c, see Supplementary Fig. 12 and 13 for identification and structures). Sessilin/tolaasin-

dependent toxicity and white-line formation were also observed when other surfactin-producing *B.*

velezensis isolates are confronted with CMR12a and CH36 (Supplementary Figs. 14 and 15, respectively).

Although the chemical basis and the steochiometry of such molecular interaction remain to be determined,

it probably follows the same rules as observed for the association between sessilin/tolaasin and other

endogenous *Pseudomonas* CLPs such as WLIP or orfamide[35] or between CLPs and other unknown

metabolites[51,52].

BSM-mediated interactions drive competitive root colonization

Our *in vitro* data thus point out how *B. velezensis* may modulate its secondary metabolome when

confronted to *Pseudomonas*. To appreciate the relevance of our findings in a more realistic context, we next

evaluated whether such BSM interplay may also occurred upon root co-colonization of tomato plantlets and

its possible impact on *Bacillus* fitness. When inoculated independently, CMR12a colonized roots more

efficiently than GA1 within the first 5 days, most probably due to a higher intrinsic growth rate[53]. Upon co-

inoculation, CMR12a colonization rate was not affected but GA1 populations were reduced compared to

mono-inoculated plantlets (Fig. 5a). UPLC-MS analysis of methanolic extracts prepared from co-bacterized

roots (and surrounding medium) revealed substantial amounts of pyochelin (Supplementary Fig. 16)

indicating that the molecule is readily formed under these conditions and could therefore also act as a

signal *in planta.* Probably due to the low populations of GA1, we could not detect Bacillus PKs and RiPPs in

these extracts. However, a significantly enhanced expression of gene clusters responsible for the synthesis

of bacillaene, difficidin and amylocyclicin was observed in GA1 cells confronted to *Pseudomonas* compared

to single inoculation (Fig 5b). It indicated that the metabolite response observed in GA1 cell cultures in EM

medium may also occur upon competitive colonization where the bacteria feed exclusively on root

exudates.

Lipopeptides involved in the interference interaction described in the previous section are also readily

formed upon single and dual root colonization (Fig. 5d). We hypothesized that the inhibitory effect of

sessilin may impact the colonization potential of GA1 in presence of CMR12a which was confirmed by the

increase in GA1 populations co-inoculated with the Δ*sesA* mutant (Fig 5c). Moreover, colonization by the

Δ*srfaA* mutant is more impacted compared to WT when co-cultivated with CMR12a and a significant gain in

root establishment is recovered upon co-colonization with the Δ*sesA* mutant (Fig. 5d). The sessilin-surfactin

interplay thus also occurs *in planta.* Sessilin would confer a competitive advantage to CMR12a during

colonization by inhibiting GA1 development but efficient surfactin production on roots, however, may

provide some protection to *Bacillus* cells.

DISCUSSION

Our current knowledge on the molecular interaction between soil bacilli and pseudomonads is rather limited, except the fact that the Type VI secretion system and the antibiotic 2,4-diacetylphloroglucinol may impact key developmental traits in *B. subtilis* such as biofilm formation and/or sporulation rate[54,55]. Here we show that the model species *B. velezensis* can mobilize a substantial part of its secondary metabolome to face *Pseudomonas* competitors. Our data provide the first evidence for the enhanced synthesis of broad-spectrum PKs (bacillaene and difficidin-type acting independently or in synergy) together with stimulation of bacteriocin production in *Bacillus* upon perception of other bacteria both in contact-independent *in vitro* settings and upon competitive root colonization. This correlates with an enhanced antibacterial potential which is of interest for biocontrol but which can also be considered as a defensive strategy to persist in its natural competitive niche. *B. velezensis* calls its antibiotic arsenal upon sensing *Pseudomonas* but we also show that the bacterium also recruits its multifunctional surfactin CLP to improve multicellular mobility. It may represent an escape mechanism enabling *Bacillus* cells to relocate after detecting harmful challengers. Improved motility of *B. subtilis* has been already described upon sensing competitors such as *Streptomyces venezuelae*[6,56] but no relationship was established with enhanced production of BSMs potentially involved in the process. We also highlight a new role for surfactin acting as a chemical shield to counteract the toxicity of exogenous CLPs. Intraspecies CLP co-precipitation has been reported[30] but our results make sense of this phenomenon in the context of interference interaction between two different genera. In planta, this new function of surfactin contributes to *Bacillus* competitiveness for root invasion. This has to be added to other previously reported implications of surfactin in *B. subtilis* interspecies interactions such as interfering with the growth of closely related species in synergy with cannibalism toxins [57], inhibiting the development of *Streptomyces* aerial hyphae [58] or participating in the expansion and motility of the interacting species [59].

Furthermore, we exemplify that PKs stimulation in *B. velezensis* is mainly mediated by pyochelin (even if other secreted products can also play some role) produced by *Pseudomonas* as secondary siderophore. Our data show that *Bacillus* perceives pyochelin in a way independent of iron stress and piracy indicating that beyond its iron-scavenging function, this siderophore may thus also act as infochemical in interspecies cross-talk. In the pairwise system used here, pyochelin signaling superimposes the possible effect of iron limitation in the external medium which may also result in enhanced production of antibacterial metabolites by *Bacillus* as occasionally reported[60]. That said, due to the limitation in bioavailable iron, almost all known rhizobacterial species have adapted to produce their siderophores to compete for this

essential element. Siderophore production is thus probably the most widely conserved trait among soil
bacteria. It also means that upon recognition of exogenous siderophores, any isolate may somehow identify
surrounding competitors. However, some of these siderophores are structurally very variable and almost
strain-specific (such as pyoverdins from fluorescent pseudomonads) while some others are much more
widely distributed across species and even genera (enterobactin-like, citrate). In both cases, their
recognition would not provide proper information about the producer because too specific or too general
respectively. Interestingly, the synthesis of pyochelin and its structurally very close enantio form is
conserved in several but not all *Pseudomonas* sp. as well as in a limited number of species belonging to
other genera such as *Burkholderia* and *Streptomyces*. We,therefore, hypothesize that *Bacillus* may have
evolved some chelator-sensing systems targeting siderophores that are conserved enough to be detected
but restricted to specific microbial phylogenetic groups. With this mechanism, soil bacilli would rely on
siderophores as public goods to accurately identify competitors and respond in an appropriate way like
remodeling its BSM secretome. This novel concept of chelator sensing represents a new facet of the
siderophore-mediated social interactions. Whether it is used for other secondary siderophores than
pyochelin and if yes whether this adaptive trait can be generalized to other soil-dwelling species remains
to be determined.

In summary, we provide the first evidence that the model *B. velezensis* species can mobilize a substantial
part of his secondary metabolome upon a perception of *Pseudomonas* competitors. Beyond the notion of
specialized metabolites, we point out unsuspected additional functions for some of these small molecules in
the context of interactions between bacterial clades that are important members of the plant-associated
microbiome. This metabolite response is viewed as a way to mount a multi-faceted defensive strategy to
gain fitness and persistence in its natural competitive niche.

MATERIAL AND METHODS

Bacterial strains and growth conditions

Strains and plasmids used in this study are listed in Supplementary Table S1. *B. velezensis* strains were
grown on recomposed exudate (EM) solid medium [29] or with shaking (160 rpm), at 30°C in EM liquid
medium. Deletion mutants of *B. velezensis* were selected on chloramphenicol (5 μg/ml) and/or phleomycin

(4 µg/ml) on Lysogeny broth (LB) (10 g l^{-1} NaCl, 5 g l^{-1} yeast extract and 10 g l^{-1} tryptone). *Pseudomonas* sp.

strains were grown on King B (20 g l^{-1} of bacteriological peptone, 10 g l^{-1} of glycerol and 1.5 g l^{-1} of K_2HPO_4,

1.5 g l^{-1} of $MgSO_4.7H_2O$, pH=7) and casamino acid (CAA) solid and liquid medium (10 g l^{-1} Casamino acid, 0.3

g l^{-1} K_2HPO_4, 0.5 g l^{-1} $MgSO_4$ and pH=7) with shaking (120 rpm), at 30°C. The phytopathogenic bacterial

strains were grown on LB and EM solid and liquid media and with shaking (150 rpm), at 30°C.

Construction of deletion mutants of *B. velezensis* GA1

All deletion mutants were constructed by marker replacement. Briefly, 1 kb of the upstream region of the

targeted gene, an antibiotic marker (chloramphenicol or phleomycine cassette) and downstream region of

the targeted gene was PCR amplified with specific primers (listed in Supplementary Table S2). The three

DNA fragments were linked by overlap PCR to obtain a DNA fragment containing the antibiotic marker

flanked by the two homologous recombination regions. This latter fragment was introduced into *B.*

velezensis GA1 by natural competence induced by nitrogen limitation[61]. Homologous recombination event

was selected by chloramphenicol resistance (or phleomycin resistance for double mutants) on LB medium.

All gene deletions were confirmed by PCR analysis with the corresponding UpF and DwR specific primers

and by the loss of the corresponding BSMs production.

Transformation of the *B. velezensis* GA1 strain was performed following the protocol described by Jarmer et

al., [61] with certain modifications. One fresh GA1 colony was inoculated into LB liquid medium at 37°C (160

rpm) until reaching an OD_{600nm} of 1.0. Afterwards, cells were washed one time with peptone water and one

time with a modified Spizizen minimal salt medium called MMG liquid medium (19 g l^{-1} K_2HPO_4 anhydrous; 6

g l^{-1} KH_2PO_4; 1 g l^{-1} Na_3 citrate anhydrous; 0.2 g l^{-1} $MgSO_4$ $7H_2O$; 2 g l^{-1} Na_2SO_4; 50 µM $FeCl_3$ (sterilized by

filtration at 0.22 µm); 2µM $MnSO_4$; 8 g l^{-1} glucose; 2 g l^{-1} L-glutamic acid; pH 7.0), 1µg of DNA recombinant

fragment was added to the GA1 cells suspension adjusted to an OD_{600nm} of 0.01 into MMG liquid medium.

One day after incubation at 37°C with shaking at 165 rpm, bacteria were spread on LB plates supplemented

with the appropriated antibiotic to select positive colonies.

Construction of deletion mutants of *Pseudomonas* sp. CMR12a

Pyochelin and pyoverdine mutants of *Pseudomonas* sp. CMR12a were constructed using the I-SceI system

and the pEMG suicid vector [62,63]. Briefly, the upstream and downstream region flanking the *pchA*

(*C4K39_5481*) or the *pvdI* (*C4K39_6027*) genes were PCR amplified (with primers listed in the

Supplementary Table S2), linked via overlap PCR and inserted into the pEMG vector. The resulting plasmid

(listed in Supplementary Table S1) was integrated by conjugation into the *Pseudomonas* sp. CMR12a

chromosome via homologous recombination. Kanamycin (25µg/mL) resistant cells were selected on King B

agar plates and transformed by electroporation with the pSW-2 plasmid (harboring I-SceI system).

Gentamycin (20µg/ml) resistant colonies on agar plates were transferred to King B medium with and

without kanamycine to verify the loss of the antibiotic (kanamycin) resistance. *Pseudomonas* mutants were

identified by PCR with the corresponding UpF and DwR specific primers and via the loss of pyochelin or/and

pyoverdin production (see section Secondary metabolites analysis).

***Pseudomonas* sp. cell-free supernatant**

Pseudomonas sp. strains were grown overnight on LB solid medium, at 30°C. Cells were adjusted to OD_{600nm}

of 0.05 by resuspension with 100 ml of CAA and when appropriate supplemented with 20 µg/l of $FeCl_3.6H_2O$

(iron supplementation). Cultures were shaken at 120 rpm at 30°C for 48 h and then centrifuged at 5000 rpm

at room temperature (22°C) with a Universal 320 centrifuge (Hettich) for 20 min. The supernatant was

sterilized with a 0.22 µm pore size Sterile PES Syringe (ROCC S.A.) filters and stored at -20°C until needed.

Dual interaction and time course experiment

B. velezensis strains were grown overnight on LB solid medium, at 30°C. Cells were resuspended in 2 ml of

$EM_{1/2}$ liquid medium to a final OD_{600nm} of 0.1 in which 1, 2, 4, or 8% v/v (depending on the experiment) of

Pseudomonas CFS were added while the control remained without the addition of CFS. *B. velezensis* liquid

cultures were shaken in an incubator at 300 rpm at 30°C for 24 h. During the culture time course

experiment, at 8, 24, and 36 h, the OD_{600nm} was measured with VWR, V-1200 Spectrophotometer to

correlate it with *Pseudomonas* CFS effect on *Bacillus* growth and BSMs production. In addition, 2 ml of the

(co-)culture supernatants were sampled, centrifugated at 5000 rpm at room temperature (approx. 22°C) for

10 min to extract supernatants and collect the cells. Cell-free (co-)culture supernatants were obtained by

sterilization with a 0.22µm pore size Sterile CA Syringe filters and thus used for analytical analysis of

secondary metabolites (as mentioned above) and antibacterial assays. For some experiments using 2H-bae

as a marker, the CFS obtained from the double mutant sessilin and orfamide (Δ*sesA-ofaBC*) was used as a

control instead of CFS from CMR12a wild-type because it yielded a higher response and lower inhibition

interferences by CLPs. The remaining cells after supernatant collection were stored at -80°C to avoid RNA

degradation, until performing RT-qPCR analysis.

Antimicrobial activity assays

Antibacterial activity of the *B. velezensis* supernatant generated after dual interaction with *Pseudomonas*

CFS was tested against *X. campestris* pv. *campestris* and *C. michiganensis* subsp. *michiganensis.* The activity

of co-culture supernatants was quantified in microtiter plates (96-well) filled with 250 µl of LB liquid

medium, inoculated at OD_{600nm}= 0.1 with *X. campestris* pv. *campestris* and *C. michiganensis* subsp.

michiganensis and supplemented with 2% or 6% v/v of *B. velezensis* supernatants, respectively. The activity

of (co-)culture supernatants was estimated by measuring the pathogen OD_{600nm} every 30min during 24h

with a Spectramax® (Molecular Devices, Wokingham, UK), continuously shaken, at 30°C. For estimating the

activity of co-culture supernatants on solid medium, 5 µl supernatantwas applied to a sterile paper disk (5

mm diameter). After drying, diskswere placed on LBAsquare plates previously inoculated with a confluent

layer of *X. campestris* pv. *campestris* or *C. michiganensis* subsp. *michiganensis*. LB liquid medium was used

as negative control. Plates were incubated at 25°C for 48 h. Three repetitions were done and the inhibition

zones from the edge of the paper discs to the edge of the zone were measured.

RNA isolation and RT-qPCR.

RNA extraction and Dnase treatment were carried out using the NucleoSpin RNA Kit (Macherey Nagel,

Germany), following the Gram + manufacturer's protocol. RNA quality and quantity were performed with

Thermo scientific NanoDrop 2000 UV-vis Spectrophotometer. The gyrase gene was chosen as a

housekeeping gene to normalize mRNA levels between different samples. The target gene in this study was

acnA. Primer 3 program available online was used for primer design and primers were synthesized by

Eurogentec. The primer efficiency was evaluated and primer pairs showing an efficiency between 90 and

110% in qPCR analysis were selected. Reverse transcriptase and RT-qPCR reactions were conducted using

the Luna® Universal One-Step RT-qPCR Kit (New England Biolabs, Ipswich, MA, United States). The reaction

was performed with 50 ng of total RNA in a total volume of 20 µL: 10 µL of luna universal reaction mix, 0.8

µL of each primer (10 µM), 5 µL of cDNA (50ng), 1µL of RT Enzyme MIX, 2.4 µl of Nuclease-free water. The

thermal cycling program applied on the ABI StepOne was: 55°c for 10 min, 95C °C for 1 min, 40 cycles of 95

°C for 10 s and 60 °C for 1 min, followed by a melting curve analysis performed using the default program of

the ABI StepOne qPCR machine (Applied Biosystems). The real-time PCR amplification was run on the ABI

step-one qPCR instrument (Applied Biosystems) with software version 2.3. Analysis was conducted on gene

expression level and has been quantified using the $2^{-\Delta\Delta CT}$ method[64].

Secondary metabolite analysis

For detection of BSMs, *B. velezensis* and *Pseudomonas* sp. were cultured in EM and CAA as described above. After an incubation period of 24 h for *B. velezensis*, if not differentially indicated, and 48 h for *Pseudomonas* sp., supernatants of the bacteria were collected and analyzed by UPLC MS/MS and UPLC qTOF MS/MS. Analysis of *Bacillus* metabolites was performed by UPLC MS/MS using an Acquity UPLC® BEH C 18 column (L=50mm, D= 2.1mm, Particles diameter= 1.7µm) (Waters, milford, MA, USA). The volume injected was 10 µl. Mass spectrometer (Acquity UPLC ® Class H SQD (Water, milford, MA, USA)) was set in negative mode ESI- and positive more ES+ (cone voltage: 60V) with a mass range from m/z 300.00 to 2048.00. Analysis of *Pseudomonas* metabolites was performed by Q-TOF (tandem mass spectrometry, quadrupole and Time of flight detector combined) (Agilent, Waldbronn, Germany) and a diode array detector (DAD) 190 to 601 nm (steps: 1nm). The volume injected was 10µl. The mass analysis was performed in positive mode ES+ (Dual AJS ESI) (Vcap= 3500V, Nozzle Voltage= 1000V), with a mass range from m/z 180 to 1700.

Bioguided fractionation

Pseudomonas CFS were concentrated with a C18 cartridge 'Chromafix, small' (Macherey-Nagel, Düren, Germany). The column was conditioned with 10 ml of MeOH followed by 10ml of milliQ water. Then, 20 ml of supernatant flowed through the column. The metabolites were eluted with 1 ml of a solution of increasing acetonitrile/water ratio from 5:95 to 100:0 (v/v). The triggering effect of these fractions on *Bacillus* 2H-bae production was tested in 48 wells microplate containing 1 ml of EM medium inoculated with *B. velezensis* GA1 (OD_{600nm} = 0.1) and 4% v/v of aforementioned *Pseudomonas* fractions, growing for 24h, with shaking at 300 rpm and 30°C. Afterward, the production of 2H-bae was quantified compared to controls and crude supernatant.

Purification of pyochelin and pyoverdine

Pyoverdine and pyochelin were purified in two steps. Firstly, *Pseudomonas* CFS were concentrated with a C18 cartridge (as indicated in section Bioguided fractionation) and eluted with 2 times 2ml of a solution of water and ACN (15 and 30% of ACN (v/v)). Secondly, the fractions were injected on HPLC for purification performed on an Eclipse+ C18 column (L=150mm, D= 3.0mm, Particles diameter 5µm) (Agilent, Waldbronn, Germany). The volume injected was 100 µl. The UV-Vis absorbance was measured with a VWD Agilent technologies 1100 series (G1314A) detector (Agilent, Waldbronn, Germany). The lamp used was a

Deuterium lamp G1314 Var Wavelength Det. (Agilent, Waldbronn, Germany). Two wavelengths were

selected: 320 nm, used for the detection of pyochelin, and 380 nm, used for the detection of pyoverdine.

The fractions containing the pyoverdine and pyochelin were collected directly at the detector output.

Further, the purity of the samples was verified by two detectors, a diode array detector (DAD) 190 to 601

nm (steps: 1nm) and a Q-TOF (tandem mass spectrometry, quadrupole and Time of flight detector

combined) (Agilent, Waldbronn, Germany). Electrospray ionization was performed in positive mode (ESI+)

(Dual AJS ESI) (Vcap= 3500V, Nozzle Voltage= 1000V), with mass range from m/z 200 to 1500. Finally, the

concentration of pyoverdine and pyochelin were estimated by utilization of Beer Lambert law formula,

A=Ɛlc (A: absorbance; Ɛ: molar attenuation coefficient or absorptivity of the attenuating species; l: optical

path length and c: concentration of molecule). l value for pyochelin and pyoverdine is 1 cm while Ɛ is 4000

$L.mol^{-1}.cm^{-1}$ or 16000 $L.mol^{-1}.cm^{-1}$, respectively [65]. The absorbance was measured with VWR, V-1200

Spectrophotometer, at 320nm (pH=8) for pyochelin and 380nm (pH=5) for pyoverdine[65]. Further, the

absorbance value was used for calculating the final concentration. The fragmentation pattern of

Pseudomonas sp. CMR12a pyoverdine was obtained by UPLC MS/MS analysis of m/z = 1288.5913 ion in

positive mode with a fragmentation energy at 75V and compared to the one described in *P. protegens* Pf-5

[1].

Confrontation, white line formation and motility test

For confrontation assays on agar plates*, Bacillus* and *Pseudomonas.* strains were grown overnight in $EM_{1/2}$

and CAA liquid mediums, respectively. After bacterial washing in peptone water and adjustment of OD_{600nm}

to 0.1, 5 µl of bacterial suspension was spotted at 1 mm, 5 mm and 7.5 mm distance onto an $EM_{1/2}$ agar

plate. For the white line formation experiments, *B. velezensis* line was applied with a cotton stick and 5 µl of

Psuedomonas sp. cell suspensions were spotted at a 5 mm distance onto CAA agar plates. Plates were

incubated at 30 °C and images taken after 24 h. Photographs were captured using CoolPix camera (NiiKKOR

60x WIDE OPTICAL ZOOM EDVR 4.3-258 mm 1:33-6.5), photos were thus proceeded in Adobe Photoshop CC

2019 or by direct edition.

Maldi-FT-ICR MS imaging

Mass spectrometry images were obtained as recently described (Kune et al 2019) using a FT-ICR mass

spectrometer (SolariX XR 9.4T, (Bruker Daltonics, Bremen, Germany)) mass calibrated from 200 m/z to

2,300 m/z to reach a mass accuracy of 0.5 ppm. Region of interest from agar microbial colonies were

directly collected from the Petri dish and transferred onto an ITO Glass slide (Bruker, Bremen, Germany),

previously covered with double sided conductive carbon tape. The samples were dried under vacuum and

covered with an α-cyano-4-hydroxycinnamic acid (HCCA) matrix solution at 5mg/mL (70:30

acetonitrile:water v/v). In total, 60 layers of HCCA matrix were sprayed using the SunCollect instrument

(SunChrom, Friedrichsdorf, Germany). FlexImaging 5.0 (Bruker Daltonics, Bremen, Germany) software was

used for MALDI MS imaging acquisition, with a pixel step size for the surface raster set to 100 μm.

***In planta* competition**

For *in planta* studies, tomato seeds (*Solanum lycopersicum* var. Moneymaker) were sterilized in 75%

ethanol with shaking for 2 min. Subsequently, ethanol was removed and seeds were added to the 50 ml

sterilization solution (8.5 ml of 15% bleach, 0.01 g of Tween 80 and 41.5 ml of sterile ultra-pure water) and

shaken for 10 min. Seeds were thereafter washed five times with water to eliminate stock solution residues.

Further, seeds were placed on square Petri dishes (5 seeds/plate) containing Hoagland solid medium (14 g/l

agar, 5 ml stock 1 (EDTA 5,20 mg/l; $FeSO_4x7H_2O$ 3,90 mg/l; H_3BO_3 1,40 mg/l; $MgSO_4x7H_2O$ 513 mg/l;

$MnCl_2x4H_2O$ 0,90 mg/l, $ZnSO_4x7H_2O$ 0,10 mg/l; $CuSO_4x5H_2O$ 0,05 mg/l; 1 ml in 50 ml stock 1, $NaMoO_4x2H_2O$

0,02 mg/l 1 ml in 50 ml stock 1), 5 ml stock 2 (KH_2PO_4 170 mg/l; 5 ml stock 3 : KNO_3 316 mg/l, $Ca(NO_3)_2$ $4H_2O$

825 mg/l), pH=6,5) and placed in the dark for three days. Afterwards, 6 seeds were inoculated with 2 μl of

overnight culture (OD_{600}=0.1) of the appropriate strains (control) or with a mix of *Bacillus* and *Pseudomonas*

cells (95:5 ration) (interaction), and grown at 22°C under a 16h/8h night/day cycle with constant light for

three days. After the incubation period, to determine bacterial colonization levels, bacteria from roots of six

plants per condition were detached from roots by vortexing for 1 min in peptone water solution

supplemented with 0.1% of Tween. Serial dilutions were prepared and 200 μl of each were plated onto LBA

medium using plating beads. After 24 h of incubation at 30 °C for *Pseudomonas* and at 42°C for *Bacillus*,

colonies were counted. Colonization results (six plants *per* strain) were log-transformed and statistically

analyzed. Three independent assays were performed with six plants each for *in planta* competition assays.

To measure bacterial BSMs production *in planta*, a rectangle part (1 x 2.5 cm) of medium close to the

tomato roots was sampled. BSMs were extracted for 15 min, with an acetonitrile (85%)/agar solution (1:1).

After centrifugation for 5 min at 4000 rpm, the supernatant was recovered for the UPLC analysis (section

Secondary metabolites analysis).

Statistical analysis

Statistical analyses were performed using GraphPad PRISM version 6 software (GraphPad) by using a paired

T-test or Mann-Whitney test. For multiple comparisons, one-way ANOVA and Tukey test were used in

RStudio 1.1.423 statistical software environment (R Core Team, 2020). Data represent mean ± s.d. of three

independent experiments unless stated otherwise. P < 0.05 was considered statistically significant.

Figure 1. Enhanced production of BSMs and related anti-*bacterial* activities of *B. velezensis* GA1 as a

response to interaction with *Pseudomonas* sp. CMR12a secreted metabolites. a. LCMS extracted ions

chromatograms (EIC) illustrating relative abundance of ions corresponding to dihydrobacillaene (**I**),

bacillaene (**II**), bacillibactin (**III**), oxydifficidin (**IV**), difficidin (**V**), surfactins (**VI**), macrolactins (**VII**), iturins (**VIII**)

and fengycins (**IX**) produced by *B. velezensis* GA1, after 24h growth in CFS-supplemented (2 % v/v) EM

medium (blue) compared to un-supplemented control (red). **b**. Production of *B. velezensis* GA1 BSMs co-

cultivated with 2 % (v/v) of *Pseudomonas* sp. CMR12a CFS. Fold change equals to 1 is represented as a red

line and corresponds to the production of metabolites in *B. velezensis* GA1 unsupplemented. Statistical

significance of each condition (n=9 within three biological repetitions and three technical replicates) was

calculated as a comparison of BSMs production by *B. velezensis* GA1 in treated cultures with the production

in related monoculture, by using Mann–Whitney test where ''****'' represents significant difference

(P<0.0001) **c.** Enhanced Anti-*Xanthomonas campestris* (**I** and **II**) and anti-*Clavibacter michiganensis* (**III** and

IV) activities of *B. velezensis* GA1 after growth in CFS-supplemented (4 %) medium (GA1+CFS) compared to

control (GA1) assessed by inhibition zone (**I** and **III**) and liquid pathogens culture growth kinetic (**II** and **IV**) **d.**

Anti-*Xanthomonas campestris* (I) and Anti- *Clavibacter michiganensis* (II) activities of *B. velezensis* GA1 (GA1

wt) BSMs and its derived mutants impaired in production of different BSMs. Metabolites not produced by

the different mutants are illustrated with red boxes in the table. Error bars indicate standard error (n=6

within two biological repetitions and three technical replicates). Different letters indicate groups of

statistically different conditions (one-way ANOVA and Tukey test; P<0.05). **e.** Differential expression of

acnA, encoding the amylocylicin precursor, upon supplementation with CMR12a CFS compared to control

Error bars indicate standard error (n=6 within two biological repetitions and three technical replicates). The

statistical difference, in *acnA* gene expression, between the two conditions was calculated by using Mann–

Whitney test where ''**'' represents a statistically significant difference between the two conditions

(0.001>P<0.05).

Figure 2: Enantio-pyochelin as the main *Pseudomonas* sp. CMR12a triggers of the anti-*bacterial* activity of

***B. velezensis* GA1. a,** Effect of 2% v/v *Pseudomonas* sp. CMR12a mutants CFS on dihydrobacillaene (2H-bae)

production by *B. velezensis* GA1. Metabolites not produced by the different mutants are illustrated by red

boxes in the table below. Fold change equals to 1 is represented as a red line and corresponds to the

production of 2H-bae in *B. velezensis* GA1 unsupplemented culture. Error bars indicate standard error (n= 9

with three biological repetitions and three technical replicates). Different letters indicate groups of

statistically different conditions (one-way ANOVA and Tukey test; P<0.05). **b,** Differential production of 2H-

bae after addition of 0.35µM pure pyoverdine (PVD), 1.4µM pure enantio-pyochelin (PCH), 4% v/v

Pseudomonas sp. CMR12a cell-free supernatant (CFS CAA) , CMR12a cell-free supernatant from iron

supplemented culture (CFS CAA+Fe) and different concentration of iron-chelating agent 2,2'-dipyridyl (DIP).

Fold change equals to 1 is represented as a red line and corresponds to the production of 2H-bae in *B.*

velezensis GA1 unsupplemented culture. Error bars indicate standard error (n=6 within two biological

repetitions and three technical replicates). Different letters indicate groups of statistically different

conditions (one-way ANOVA and Tukey test; P-value < 0.05). **c,** Effect of pure PVD and PCH on bacillibactin

and 2H-bae production by *B. velezensis* GA1. *B. velezensis* GA1 culture was supplemented with different

concentration of pure PVD or pure PCH. Fold change equals to 1 is represented as a red line and

corresponds to the production of 2H-bae in *B. velezensis* GA1 unsupplemented culture. Error bars indicate

standard error (n=6 within two biological repetitions and three technical replicates). Different letters

indicate groups of statistically different conditions (one-way ANOVA and Tukey test; P<0.05). **d.** Growth

curve of GA1 WT and bacillibactin KO mutant (ΔdhbC) upon supplementation with pyoverdine (pvd) and

pyochelin (pch) at concentration corresponding to the ones in co-culture experiment (4% (v/v))

pseudomonas CFS . Error bars indicate standard error (n=6 with two biological repetitions and three

technical replicates).

Figure 3: Enhanced motility of *Bacillus* is driven by distance and surfactin production . a, *B. velezensis* GA1

and P. sp. CMR12a phenotypes when cultured alone (left panel) or in confrontation at short distance

(1cm)(right panel). **b,** Phenotype and motile of GA1 in confrontation with CMR12a at short distance (1cm)

compared to his surfactin defficient mutant (Δ*srfaA*) **c,** Maldi FT-ICR MSI (Mass spectrometry Imaging)

heatmaps showing spatial localisation and relative abundance of C14-Surfactin ([M+Na]$^+$) when *B.*

velezensis GA1 is in confrontation with CMR12a at different distances.

Figure 4: Overview of the unsuspected roles of bacterial CLPs in *Bacillus-Pseudomonas* interaction: **a,**

Illustration of *B. velezensis* GA1 growth inhibition cultured in gelosed media in confrontation (**I**) or in liquid

with 6%v/v (**II**) with *Pseudomonas* sp. CMR12a wild-type (CFS) or mutants devoid of production of

orfamides and phenazines (Δ*ofaBC-phz*), sessilins (Δ*sesA*), sessilins and orfamides (Δ*sesA-ofaBC*), sessilins

and phenazines (Δ*sesA-phz*), or sessilins, orfamides and phenazines (Δ*sesA-ofaBC-phz*). Error bars indicate

standard error (n=6 within two biological repetitions and three technical replicates). Different letters

indicate groups of statistically different conditions (one-way ANOVA and Tukey test; P<0.05). **b, Left** :

Optical density of B. velezensis GA1 WT (GA1) after 7h culture supplemented with CMR12a WT CFS

(CMR12a) or sessilin KO mutant (Δ*sesA*).**Right :** Optical density of B. velezensis GA1 surfactin KO mutant

(Δ*srfaA*) after 7h culture supplemented with CMR12a WT CFS (CMR12a) and pure surfactin (CMR12a +

surfactin). Growth inhibition of CMR12 sessilin mutant CFS (Δ*sesA*). Unsupplemented culture of both B.

velezensis strains are mentioned as control culture**. c-** White line formation obtained by confrontation with

surfactin and sessilin/tolaasin producing strains (*B. velezensis* GA1 confronted to *Pseudomonas* sp. CMR12a

(**I**) or *P.* tolaasii CH36 (**II**). CLPs impaired mutants (Δ*srfaA,* and Δ*sesA* or Δ*tolA*) do not show any white-line.

Other *Pseudomonas* CLPs producers do not cause white-line formation or impact the *B. velezensis* GA1

growth (WCU-84, SS101, BW11M1, RW10S2).. CLPs produced by the aforementioned *Pseudomonas* strains

are listed in the table below. **d.** 3D LC-MS chromatogram after solid-liquid extraction of white-line showing

the presence of sessilin and surfactin.

Figure 5: *Bacillus-Pseudomonas in planta* competition and CLPs role. **a.** *Bacillus* (GA1) or *Pseudomonas*

(CMR12a) population recovered at 3 days post inoculation (dpi) when inoculated alone (GA1, CMR12a) or

coinoculated (co-inoculation) . Error bars indicate standard error (n=16) of 3 independent experiments

including at least 4 plants per condition. Statistical difference between the conditions was calculated by

using Mann–Whitney test where ''****'' and ''***'' represent significant differences (P<0.0001 and

0.0001>P<0.001, respectively). **b.** Relative gene expression of *srfaA, dfnA, baeJ* and *acnA*a fter 3 days of co-

inoculation on tomato plant roots. Error bars indicate standard error (n=3, each dot represents the mean of

three technical repetitions). Fold change equals to 1 is represented as a red line and corresponds to the

gene expressions in *B. velezensis* GA1 monoculture. Statistical significance of each group was calculated as a

comparison of genes expressions by *B. velezensis* GA1 in treated cultures with the one in monocultures, by

using T-test where ''*'', ''***'' and ''****'' represent significant difference (0.01>P<0.05, 0.0001>P<0.001

and P<0.0001, respectively). **c.** Recoverd population at 3dpi of *B. velezensis* GA1 WT (GA1) or his surfactin

impaired mutant (Δ*srfA*) co-inoculatied with *P. sp.* CMR12a (CMR12a) or his sessilin KO mutant (Δ*sesA*) .

Error bars represent standard deviations (n=16 to 18) of 3 independent experiments including at least 4

plants per condition. Statistical difference between two conditions was calculated by using Mann–Whitney

test where ''**'' indicates a significant difference (0.001>P<0.05). **d.** LC-MS EIC illustrating relative *in planta*

production of sessilin and surfactin by monocultures of *B. velezensis* GA1 (GA1) and co-cultures of wild

types (GA1+CMR12a) and *B. velezensis* GA1 and *Pseudomonas* sp. CMR12a impaired in sessilin production

(GA1+Δsessilin).

Funding

This work was supported by the EU Interreg V France-Wallonie-Vlaanderen portfolio SmartBiocontrol

(Bioprotect and Bioscreen projects, avec le soutien du Fonds européen de développement régional - Met

steun van het Europees Fonds voor Regionale Ontwikkeling), by the European Union Horizon 2020 research

and innovation program under grant agreement No. 731077 and by the EOS project ID 30650620 from the

FWO/F.R.S.-FNRS. FB is recipient of a F.R.I.A. fellowship (Formation à la Recherche dans l'Industrie et

l'Agriculture) and MO is senior research associate at the F.R.S.-F.N.R.S. We are grateful to the KU Leuven

HPC infrastructure and the Flemish Supercomputer Center (VSC) for providing the computational resources

and services to perform the RNA-seq analysis. The MALDI FT-ICR SolariX XR were funded by FEDER BIOMED

HUB Technology Support (number 2.2.1/996).

Acknowledgments

We gratefully acknowledge Sébastien Rigali, Alexandre Jousset and Loïc Ongena for critically reading the

manuscript. We thank C. Keel for the kind gift of strains and J. Vacheron for the very helpful indications on

Pseudomonas mutagenesis.

REFERENCES

1.   Nayfach, S. *et al.* A genomic catalog of Earth's microbiomes. *Nat. Biotechnol.* (2020).
doi:10.1038/s41587-020-0718-6

2.   Fierer, N. Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat. Rev.*
Microbiol. **15**, 579–590 (2017).

3.   Schmidt, R., Ulanova, D., Wick, L. Y., Bode, H. B. & Garbeva, P. Microbe-driven chemical ecology:
past, present and future. *ISME Journal* **13**, 2656–2663 (2019).

4.   Tyc, O., Song, C., Dickschat, J. S., Vos, M. & Garbeva, P. The Ecological Role of Volatile and Soluble
Secondary Metabolites Produced by Soil Bacteria. *Trends Microbiol.* **25**, 280–292 (2017).

5.   Bernal, P., Llamas, M. A. & Filloux, A. Type VI secretion systems in plant-associated bacteria. *Environ.*
Microbiol. **20**, 1–15 (2018).

6.   Andrić, S., Meyer, T. & Ongena, M. Bacillus Responses to Plant-Associated Fungal and Bacterial
Communities. *Front. Microbiol.* **11**, (2020).

7.   Abreu, N. A. & Taga, M. E. Decoding molecular interactions in microbial communitiesa. *FEMS*
Microbiol. Rev. **40**, 648–663 (2016).

8.   Mendes, R., Garbeva, P. & Raaijmakers, J. M. The rhizosphere microbiome: significance of plant
beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiol. Rev.* **37**, 634–
663 (2013).

9.   Müller, D. B., Vogel, C., Bai, Y. & Vorholt, J. A. The Plant Microbiota: Systems-Level Insights and
Perspectives. *Annu. Rev. Genet.* **50**, 211–234 (2016).

10.   Penha, R. O., Vandenberghe, L. P. S., Faulds, C., Soccol, V. T. & Soccol, C. R. Bacillus lipopeptides as
powerful pest control agents for a more sustainable and healthy agriculture: recent studies and
innovations. *Planta* **251**, 1–15 (2020).

11.   Grubbs, K. J. *et al.* Large-Scale Bioinformatics Analysis of *Bacillus* Genomes Uncovers Conserved
Roles of Natural Products in Bacterial Physiology. *mSystems* **2**, 1–18 (2017).

12.   Harwood, C. R., Mouillon, J.-M. M., Pohl, S. & Arnau, J. Secondary metabolite production and the
safety of industrially important members of the Bacillus subtilis group. *FEMS Microbiol. Rev.* **42**, 721–
738 (2018).

13.   Ye, M. *et al.* Characteristics and Application of a Novel Species of *Bacillus* : *Bacillus velezensis*. *ACS*
Chem. Biol. **13**, 500–505 (2018).

14.   Rabbee, M. *et al. Bacillus velezensis*: A Valuable Member of Bioactive Molecules within Plant
Microbiomes. *Molecules* **24**, 1–13 (2019).

15.   Pieterse, C. M. J. *et al.* Induced Systemic Resistance by Beneficial Microbes. *Annu. Rev. Phytopathol.*

52, 347–375 (2014).

16.   Köhl, J., Kolnaar, R. & Ravensberg, W. J. Mode of Action of Microbial Biological Control Agents

Against Plant Diseases: Relevance Beyond Efficacy. *Front. Plant Sci.* **10**, 845 (2019).

17.   Traxler, M. F. & Kolter, R. Natural products in soil microbe interactions and evolution. *Nat. Prod. Rep.*

32, 956–970 (2015).

18.   Raaijmakers, J. M. & Mazzola, M. Diversity and Natural Functions of Antibiotics Produced by

Beneficial and Plant Pathogenic Bacteria. *Annu. Rev. Phytopathol.* **50**, 403–424 (2012).

19.   Li, Y. & Rebuffat, S. The manifold roles of microbial ribosomal peptide-based natural products in

physiology and ecology. *J. Biol. Chem.* **295**, 34–54 (2020).

20.   Arguelles Arias, A. *et al.* Characterization of Amylolysin, a Novel Lantibiotic from Bacillus

amyloliquefaciens GA1. *PLoS One* **8**, e83037 (2013).

21.   Scholz, R. *et al.* Amylocyclicin, a novel circular bacteriocin produced by Bacillus amyloliquefaciens

FZB42. *J. Bacteriol.* **196**, 1842–1852 (2014).

22.   Torres Manno, M. A. *et al.* GeM-Pro: a tool for genome functional mining and microbial profiling.

Appl. Microbiol. Biotechnol. **103**, 3123–3134 (2019).

23.   Dutta, S. *et al.* Structure of a modular polyketide synthase. *Nature* **510**, 512–517 (2014).

24.   Winn, M., Fyans, J. K., Zhuo, Y. & Micklefield, J. Recent advances in engineering nonribosomal

peptide assembly lines. *Nat. Prod. Rep.* **33**, 317–347 (2016).

25.   Bozhüyük, K. A., Micklefield, J. & Wilkinson, B. Engineering enzymatic assembly lines to produce new

antibiotics. *Curr. Opin. Microbiol.* **51**, 88–96 (2019).

26.   Zhao, X. & Kuipers, O. P. O. P. Identification and classification of known and putative antimicrobial

compounds produced by a wide variety of Bacillales species. *BMC Genomics* **17**, (2016).

27.   Raaijmakers, J. M., de Bruijn, I., Nybroe, O. & Ongena, M. Natural functions of lipopeptides from

Bacillus and Pseudomonas: More than surfactants and antibiotics. *FEMS Microbiol. Rev.* **34**, 1037–

1062 (2010).

28.   Blin, K. *et al.* antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic*

Acids Res. **47**, W81–W87 (2019).

29.   Nihorimbere, V. *et al.* Impact of rhizosphere factors on cyclic lipopeptide signature from the plant

beneficial strain Bacillus amyloliquefaciens S499. *FEMS Microbiol. Ecol.* **79**, 176–191 (2012).

30.   D'aes, J. *et al.* To settle or to move? The interplay between two classes of cyclic lipopeptides in the

673   biocontrol strain Pseudomonas CMR12a. *Environ. Microbiol.* **16**, 2282–2300 (2014).

674 31. Hua, G. K. H. & Höfte, M. The involvement of phenazines and cyclic lipopeptide sessilin in biocontrol

675   of Rhizoctonia root rot on bean (Phaseolus vulgaris) by Pseudomonas sp. CMR12a is influenced by

676   substrate composition. *Plant Soil* **388**, 243–253 (2015).

677 32. Perneel, M. *et al.* Characterization of CMR5c and CMR12a, novel fluorescent Pseudomonas strains

678   from the cocoyam rhizosphere with biocontrol activity. *J. Appl. Microbiol.* **103**, 1007–1020 (2007).

679 33. Olorunleke, F. E. *et al.* Coregulation of the cyclic lipopeptides orfamide and sessilin in the biocontrol

680   strain Pseudomonas sp. CMR12a. *Microbiologyopen* **6**, 1–12 (2017).

681 34. Olorunleke, F. E., Hua, G. K. H., Kieu, N. P., Ma, Z. & Höfte, M. Interplay between orfamides, sessilins

682   and phenazines in the control of Rhizoctonia diseases by Pseudomonas sp. CMR12a. *Environ.*

683   *Microbiol. Rep.* **7**, 774–781 (2015).

684 35. Geudens, N. & Martins, J. C. Cyclic lipodepsipeptides from Pseudomonas spp. - Biological Swiss-Army

685   knives. *Front. Microbiol.* **9**, 1–18 (2018).

686 36. Omoboye, O. O. *et al.* Pseudomonas cyclic lipopeptides suppress the rice blast fungus Magnaporthe

687   oryzae by induced resistance and direct antagonism. *Front. Plant Sci.* **10**, 901 (2019).

688 37. Mansfield, J. *et al.* Top 10 plant pathogenic bacteria in molecular plant pathology. *Mol. Plant Pathol.*

689   **13**, 614–629 (2012).

690 38. Molinatto, G. *et al.* Complete genome sequence of Bacillus amyloliquefaciens subsp. plantarum

691   S499, a rhizobacterium that triggers plant defences and inhibits fungal phytopathogens. *J.*

692   *Biotechnol.* **238**, 56–59 (2016).

693 39. Fan, B. *et al.* Bacillus velezensis FZB42 in 2018: The gram-positive model strain for plant growth

694   promotion and biocontrol. *Front. Microbiol.* **9**, 3389 (2018).

695 40. Pandin, C. *et al.* Complete genome sequence of Bacillus velezensis QST713: A biocontrol agent that

696   protects Agaricus bisporus crops against the green mould disease. *J. Biotechnol.* **278**, 10–19 (2018).

697 41. Youard, Z. A., Mislin, G. L. A., Majcherczyk, P. A., Schalk, I. J. & Reimmann, C. Pseudomonas

698   fluorescens CHA0 produces enantio-pyochelin, the optical antipode of the pseudomonas aeruginosa

699   siderophore pyochelin. *J. Biol. Chem.* **282**, 35546–35553 (2007).

700 42. Grandchamp, G. *et al.* Pirated siderophores promote sporulation in Bacillus subtilis . *Appl. Environ.*

701   *Microbiol.* **83**, 1–17 (2017).

702 43. Miethke, M. *et al.* Ferri-bacillibactin uptake and hydrolysis in Bacillus subtilis. *Mol. Microbiol.* **61**,

703   1413–1427 (2006).

704   44.   Adler, C. *et al.* Catecholate Siderophores Protect Bacteria from Pyochelin Toxicity. *PLoS One* **7**,
705   (2012).

706   45.   Trottmann, F., Franke, J., Ishida, K., García-Altares, M. & Hertweck, C. A Pair of Bacterial
707   Siderophores Releases and Traps an Intercellular Signal Molecule: An Unusual Case of Natural
708   Nitrone Bioconjugation. *Angew. Chemie* **131**, 206–210 (2019).

709   46.   Hartney, S. L. *et al.* Ferric-pyoverdine recognition by Fpv outer membrane proteins of pseudomonas
710   protegens pf-5. *J. Bacteriol.* **195**, 765–776 (2013).

711   47.   Youard, Z. A., Wenner, N. & Reimmann, C. Iron acquisition with the natural siderophore enantiomers
712   pyochelin and enantio-pyochelin in Pseudomonas species. *BioMetals* **24**, 513–522 (2011).

713   48.   Rokni-Zadeh, H. *et al.* Genetic and functional characterization of cyclic lipopeptide white-line-
714   inducing principle (WLIP) production by rice rhizosphere isolate pseudomonas putida RW10S2. *Appl.*
715   *Environ. Microbiol.* **78**, 4826–4834 (2012).

716   49.   Buyer, J. S., Wright, J. M. & Leong, J. Structure of Pseudobactin A214, a Siderophore from a Bean-
717   Deleterious Pseudomonas. *Biochemistry* **25**, 5492–5499 (1986).

718   50.   van Gestel, J., Vlamakis, H. & Kolter, R. From Cell Differentiation to Cell Collectives: Bacillus subtilis
719   Uses Division of Labor to Migrate. *PLoS Biol.* **13**, 1–29 (2015).

720   51.   Munsch, P. & Alatossava, T. The white-line-in-agar test is not specific for the two cultivated
721   mushroom associated pseudomonads, Pseudomonas tolaasii and Pseudomonas 'reactans'.
722   *Microbiol. Res.* **157**, 7–11 (2002).

723   52.   Salari, F. *et al.* Draft Genome Sequence of Pseudomonas aeruginosa Strain LMG 1272, an Atypical
724   White Line Reaction Producer. *Microbiol. Resour. Announc.* **9**, (2020).

725   53.   Xu, Z. *et al.* Antibiotic Bacillomycin D Affects Iron Acquisition and Biofilm Formation in Bacillus
726   velezensis through a Btr-Mediated FeuABC-Dependent Pathway. *Cell Rep.* **29**, 1192-1202.e5 (2019).

727   54.   Powers, M. J., Sanabria-Valentín, E., Bowers, A. A. & Shank, E. A. Inhibition of cell differentiation in
728   Bacillus subtilis by Pseudomonas protegens. *J. Bacteriol.* **197**, 2129–2138 (2015).

729   55.   Molina-Santiago, C. *et al.* The extracellular matrix protects Bacillus subtilis colonies from
730   Pseudomonas invasion and modulates plant co-colonization. *Nat. Commun.* **10**, (2019).

731   56.   Liu, Y., Kyle, S. & Straight, P. D. Antibiotic Stimulation of a Bacillus subtilis Migratory Response.
732   *mSphere* **3**, 1–13 (2018).

733   57.   Rosenberg, G. *et al.* Not so simple, not so subtle: The interspecies competition between Bacillus
734   simplex and Bacillus subtilis and its impact on the evolution of biofilms. *npj Biofilms Microbiomes* **2**,

15027 (2016).

58.   Straight, P. D., Willey, J. M. & Kolter, R. Interactions between Streptomyces coelicolor and Bacillus

subtilis: Role of surfactants in raising aerial structures. *J. Bacteriol.* **188**, 4918–4925 (2006).

59.   Molina-Santiago, C. *et al.* The extracellular matrix protects Bacillus subtilis colonies from

Pseudomonas invasion and modulates plant co-colonization. *Nat. Commun.* **10**, 1919 (2019).

60.   Lee, N. *et al.* Iron competition triggers antibiotic biosynthesis in Streptomyces coelicolor during

coculture with Myxococcus xanthus. *ISME J.* (2020). doi:10.1038/s41396-020-0594-6

61.   Jarmer, H., Berka, R., Knudsen, S. & Saxild, H. H. Transcriptome analysis documents induced

competence of *Bacillus subtilis* during nitrogen limiting conditions. *FEMS Microbiol. Lett.* **206**, 197–

200 (2002).

62.   Martínez-García, E. & de Lorenzo, V. Engineering multiple genomic deletions in Gram-negative

bacteria: analysis of the multi-resistant antibiotic profile of Pseudomonas putida KT2440. *Environ.*

Microbiol. **13**, 2702–2716 (2011).

63.   Vacheron, J. *et al.* T6SS contributes to gut microbiome invasion and killing of an herbivorous pest

insect by plant-beneficial Pseudomonas protegens. *ISME J.* **13**, 1318–1329 (2019).

64.   Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative

PCR and the 2-ΔΔCT method. *Methods* **25**, 402–408 (2001).

65.   Hoegy, F., Mislin, G. L. A. & Schalk, I. J. Pyoverdine and pyochelin measurements. *Methods Mol. Biol.*

1149, 293–301 (2014).

66.   R Core Team (2020). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. (2020). Available at: https://www.r-project.org/.

Figure 1

Figure 2

Figure 3

1 cm

0% ▬▬▬▬▬▬▬▬▬▬ 100%
m/z=1044.6

Figure 4

Figure 5