Scientific modelling can be accessible, interoperable and user friendly: An example for pasture and livestock modelling

Alba Marquez Torres[1], Stefano Balbi[1,2], Ferdinando Villa[1,2]

[1] Basque Centre for Climate Change, Bilbao, Biscay, Spain
[2] Ikerbasque Foundation for Science, Bilbao, Biscay, Spain

Corresponding Author:
Alba Marquez Torres[1]
Edificio Sede 1, 1º Planta 1 | Parque Cientifico UPV/EHU, 48940 Leioa, Biscay
Email address: alba.marquez@bc3research.org

Abstract

This article describes the adaptation of a non-spatial model of pastureland dynamics, including vegetation life cycle, livestock management and nitrogen cycle, for use in a spatially explicit and modular modelling platform (k.LAB) dedicated to make data and models more interoperable. The aim is to deliver an existing, locally successful monolithic model, into a more modular, transparent and accessible approach to potential end users, regional managers, farmers and other stakeholders. This allows better usability and adaptability of the model beyond its originally intended geographical scope (the Cantabrian Region in the North of Spain). The original model, named Puerto, is developed in the R language and includes 1,491 lines of code divided into 13 script files and linked to 19 input tables. The spatiotemporal rewrite is structured around a set of 10 namespaces called PaL (Pasture and Livestock), which includes 198 interoperable but independent models. The end user chooses the spatial and temporal context of the analysis through an intuitive web-based user interface called k.Explorer. Each model can be called individually or in conjunction with the others, by querying any PaL-related concepts in a search bar. A scientific workflow is built as a response, which is run to produce result datasets and a report with information on the data sources and modelling processes used, delivering results with full transparency. We argue that this work demonstrates key steps needed to create more Findable, Accessible, Interoperable and Reusable (FAIR) models. This is particularly essential in environments as complex as agricultural systems, where multidisciplinary knowledge needs to be integrated across diverse spatial and temporal scales in order to understand complex and changing problems.

Introduction

Extensive farming, when paired with the conservation of natural vegetation, has historically been
capable of sustaining food production in agricultural areas while maintaining ecosystems in good
condition (European Environment Agency, 2004; Hendrickson et al., 2008; Lemaire et al., 2014).
Since the 1950s, the increase of labour costs and beginning of widespread mechanisation and
fertilizer application in the developed countries (Hayami & Ruttan, 1971; Billen, Lassaletta &
Garnier, 2014) led to important changes such as the intensification of land use and the expansion
of farming scale. This paradigm shift benefited from the European Union's Common Agricultural
Policy (CAP) subsidies scheme. These policies simultaneously contributed to disincentivizing
low-input land uses, causing land abandonment and afforestation in extensive agricultural areas,
while also decreased agricultural commodity prices due to overproduction of intensive farming
(Strijker, 2005; Pe'er et al., 2014).

Today, multiple human activities, such as urban development and tourism, are adding further
pressures to ecosystems in addition to the increased productivity of intensive agriculture. These
activities are moving pasture from mountain areas (Daugstad, Mier & Peña-Chocarro, 2014) to
more accessible locations closer to urban centres (Fernández-Giménez & Fillat Estaque, 2012).
Agricultural intensification in concentrated areas is threatening ecological sustainability and the
provision of ecosystem services (van Zanten et al., 2014; Balbi et al., 2015). Such pressures are
leading to ecosystem degradation by reducing biodiversity and threatening species linked to low-
intensive agricultural production (Pimentel et al., 1992; Donald, Green & Heath, 2001; Henle et
al., 2008), and by depleting plant resources, increasing contamination by leachate and soil erosion
(Pimentel et al., 1995; Kumar et al., 2005; Tan, Lal & Wiebe, 2005; Rosegrant, Ringler & Zhu,
2009; Cordell, Drangert & White, 2009).

At the same time, farmland abandonment in rural areas can cause: (i) loss of woodland clearings,
(ii) increased fuel loads and fire hazards and (iii) negative impacts on biological diversity
(MacDonald et al., 2000; Casasús et al., 2007). The improvement of farmers' socio-economic
conditions, extensive farming evolution and the balance with the environment require more
efficient use of pastoral vegetation, including proper livestock management (grazing rotations by
species and across time and space (Gibon, 2005)) and the controlled use of fire to preserve
pasture availability (López, 2002; López-Sáez et al., 2016). The lack of quantitative tools for the
analysis of such processes has been a major limitation for smarter and more sustainable
management of mountain pastureland (Bernués et al., 2011).

Agricultural production systems have benefited from technological advances primarily
developed for other industries such as mechanization, synthetic fertilizers, genetic engineering
and automation. The information age brings new technology that can transform agriculture to
low-input, high-efficiency and sustainable systems (Zhang, Wang & Wang, 2002; Balbi et al.,
2015), such as cloud computing, remote sensing and artificial intelligence (Putfarken et al., 2008;
Janssen et al., 2017; Kamilaris, Kartakoullis & Prenafeta-Boldú, 2017). The agricultural industry
is now capable of gathering more comprehensive data on production variability across both
space and time (Angelov, Iglesias & Corrales, 2018). Data and models can play an important role
in sustainable agriculture, optimizing resources, providing key spatial-temporal information and

identifying the most appropriate and effective practices for better management (Vries, Teng &
Metselaar, 1993).

One main issues preventing the full use of these new technologies in agricultural modelling
arises from the multidimensional nature of needed data and models that are produced by different
scientific domains from climatology to ecology and social sciences (Farina, 2000). Although an
agricultural system can be designed for a specific purpose, such as crop production or animal
breeding, understanding it requires knowledge from diverse fields (e.g., agricultural production,
natural resources and human factors) (Argent, 2004; Jones et al., 2017). These components
cannot be studied in isolation (Hieronymi, 2013), since they interact with each other and with
their environment (Wallach et al., 2019).

The Puerto model (Busqué, 2014) was created in response to some of the above-mentioned
agricultural systems challenges. The Puerto model was developed at the Centre for Agricultural
Research and Training of Cantabria (CIFA) as part of its research on the structure, growth and
utilization of pastures in the Cantabrian rangeland (Marcos, Lodos & Rodríguez-Arango, 2003).
Puerto is an empirical dynamic model based on established biophysical relationships and
constants between vegetation's life cycle (including growth, senescence death and litterfall),
livestock grazing process (livestock ingestion, digestion, excretion and weight change) and the
nitrogen cycle (nitrogen uptake, soil cycling and leaching). It evaluates existing nitrogen and
grazing imbalances (under- or overgrazing) and their relationship with animal productivity.
Puerto's four main goals are to: (i) provide a tool to support pastoral management; (ii) quantify
and assess grazing system and nitrogen cycle imbalances; (iii) enable managers to develop
strategies to resolve imbalances; and (iv) visualize the effects of management actions through
scenarios. This model has proven to be a valuable tool for modelling pastureland in Cantabria
and was used in several regional projects (Busqué, Fernández & Fernández, 2006; Bedia,
Cabañas & Busqué, 2009) at different temporal and spatial scales. Although its reliability and
usefulness have been validated and improved over the years, this model is essentially
inaccessible to a non-initiated programming audience, and Cantabrian land managers must rely
on technical consultancies to use it. Further, Puerto has always been used in isolation, never
contributing to more comprehensive computational modelling workflows. We argue that these
limitations arise from three choices made in Puerto's modelling philosophy, which are typical to
modern environmental modelling:

1. the model's interface is not user friendly, it is coded in R and is only usable by advanced
R users, with each run requiring the modification of source files to point to input data;
2. it is monolithic and cumbersome (1,491 lines of code divided into 13 script files and
linked to 19 input tables), which makes understanding of its computational workflow
difficult;
3. it lacks transparency in the definition of multiple parameters, which lack semantics and
appear as acronyms defined as fixed values in the code.

These limitations are common practice in most current scientific modelling exercises, which are
not developed as Findable, Accessible, Interoperable and Reusable (FAIR) scientific artifacts
(Parker et al., 2002; Wilkinson et al., 2016).At the same time, the importance of accessibility,
interoperability and reusability of models and resources is increasingly recognized by modelling
communities. While novel approaches are available to facilitate that (van Ittersum et al., 2008;
Verburg, Eickhout & van Meijl, 2008; Ewert et al., 2009; Peckham, Hutton & Norris, 2013),
none has yet reached the necessary levels of practicality, generality and community acceptance
to make a dent into a still widespread model and data curation malpractice.

The aim of this article is to demonstrate the implementation of Puerto into a semantic-first
modelling approach, which aims to better achieve the FAIR criteria. This redesign makes the
models, from now on referred to as the Pasture and Livestock (PaL) namespace(s), and their
results more accessible to end users such as farmers and policy-makers. PaL is written in k.IM, a
semantic modelling language designed for the k.LAB modelling platform. K.LAB is powered by
artificial intelligence, and in particular by machine reasoning, for the interoperability of data and
models (Villa, 2007; Villa et al., 2017). PaL will be part of ARIES (ARtificial Intelligence for
Environment and Sustainability), the best known application of k.LAB (Villa et al., 2014;
Bagstad et al., 2014). ARIES is an open-source platform for interoperable models and data
backed by an international and multidisciplinary community, producing a web-based platform
linking, synthesizing and providing easy access to integrated knowledge to address a wide range
of sustainability problems. In this article, we describe the PaL implementation and its application
to a study area in Northern Spain. In the methods section, we describe the key requirements and
distinctions of the semantic modelling approach as applied to Puerto and PaL. Our results
compare the outputs of Puerto and PaL when applied to a region in eastern Cantabria and
illustrate key end-user features of the k.LAB modelling environment. Finally, our discussion and
conclusions describe implications of this approach for environmental modelling more generally
and agricultural modelling specifically.

Materials & Methods

Study area

The study area was selected to match the location where the original model has been most
frequently applied. The Pas, Miera, and Ason watersheds (43°20′36″N, 3°44′28″W) are adjacent
to the Cantabrian mountain range in the eastern Cantabria region, covering a terrestrial, riverine
and estuarine system of 173,700 ha (Fig. 1). This study area, with its river basins draining into
the Cantabrian Sea, has a temperate hyper-oceanic climate, defined mainly by mild temperatures
and high humidity due to regular precipitation and fog. Although the average annual temperature
is 14º C, snow is common in the mountains from late autumn to early spring.

152
Figure 1 Location of the case study, the Pas, Miera and Ason in watersheds in northern Spain.

This unique landscape is a product of the combined use of fire and livestock grazing for over 400
years (Montserrat & Fillat, 1990). As a consequence, almost 75% of the landscape consists of
managed grasslands and shrublands, relegating mature forests to headwater basins and marginal
lands with low agricultural value on steeper slopes. The pastoral lands are dominated by nine
pastureland types and multiple livestock types, including cattle and mares (Fig. 2).

Legend:
- Livestock
- *Helictotrichon cantabricum*
- *Senecio jacobea*
- *Ulex europeas*
- *Festuca rubra*
- *Carex sempervirens*
- *Calluna Vulgaris*
- *Ulex galli*
- Gramineae
- *Polypodiopsida*
- *Phragmites*

0 2 4 8 12 16 Kilometers

159
Figure 2 Distribution of livestock and pastureland types in the case study.

There are three climatic sub-regions influenced by the mountain ranges (including the "Picos de
Europa" mountain range) and the ocean. The coastal zone, which is under high human pressure,
has widespread grasslands and eucalyptus plantations (*Eucalyptus globulus*) on gentle slopes.
The central part is the most rugged, with elevation ranging between 100-1200 m a.s.l., dominated
by semi-extensive pastures grazed by livestock. Large areas are occupied by *Ulex europaeus*,
Erica tetralix, *Pteridium aquilinum* or *Carex asturica* and productive plantations of *Pinus*
radiata. The mountain ranges in the south, with steep slopes and a more complex management,
have a great diversity of plant communities used by livestock.

Model description

The Puerto Model

Puerto's main code can be divided into four components. The first component consists of
climate, topography and soil, which affect vegetation growth and livestock ingestion of forage.
The second part captures the entire life cycle of vegetation including growth, senescence and
litterfall. The third component focuses on the nitrogen cycle, including mineralization, plant
reabsorption and leaching. Finally, the last component describes grazing, ingestion, weight
variation, and excretion of manure and urine of livestock.

Puerto needs with a substantial set of input data and parameters to be initialized, derived from
literature or field measurements, related to vegetation, soil, climate, the nitrogen cycle and,
optionally, livestock management. In total, it requires 56 data inputs as tables and 27 constant
parameters. After initialization, it executes dynamic transitions over a modeller-defined temporal
horizon, by daily timestep for entire years, simulating the management of pastoral systems.
Outputs are produced as R tables associated with each management unit of pastureland, such
tables can be transformed into vector data by a technician proficient in Geographical Information
System (GIS) software so that the results can be displayed spatially.

The Pasture and Livestock namespaces

The Pasture and Livestock (PaL) namespaces provide an integrated modelling framework for the
Puerto model designed to better adhere to the FAIR Principles while making the model more
accessible for nontechnical users. They operate under the k.LAB open-source software platform
and k.IM semantic annotation and modelling language (Table 1) (Villa, 2007; Villa et al., 2017)
the only programming language using semantics as the primary organizational principle. The
k.LAB platform connects a network of data, models, and semantic resources distributed globally
on the semantic web. The code of PaL in k.IM language is available online at the Bitbucket
repository (https://bitbucket.org/integratedmodelling/im.ecology.grassland.livestock/).

Semantics are used to annotate all resources (i.e., data and model components) in PaL
namespaces, using a well-established and expert-vetted vocabulary (Arp, Smith & Spear, 2015).
The concepts used to build the model components and to represent data are not built specifically
for a model, but come from a shared, network-accessible worldview which provides uniform
definitions encompassing concepts and the relationships between them. The use of semantics to
describe data and models enables an artificial intelligent algorithm to build meaningful
connections between inputs and outputs by making inferences and ranking each model
component for the best fit to the concepts required as input. Any resource available in k.LAB can
be automatically and accurately interpreted by a receiving system (Martínez-López et al., 2019)
as a response to a query. Such a modelling approach is modular by design, parsimonious and
logically consistent, which makes the knowledge contained in the resources unambiguously and
more transparently sharable while making the model more accessible for non-technical users. By
providing a web-based query tool with intuitive spatial and temporal context selection
(k.Explorer), the scientific information in models and data can be displayed in understandable
and accessible fashion, without compromising on rigor and machine-readability of results.

Table 1 Comparison between R and k.IM language for the "Potential above ground biomass caused by growth"
model. Colour coding in the k.IM language denotes different types of semantic meaning: brown denotes processes,
green qualities, and blue attributes that can be combined to describe semantically meaningful scientific observables
(Villa et al., 2017).

Concept	Language	Code
Potential above ground biomass caused by growth	R	setkey(Fhijt,com2);setkey(pl1$B3,com) T1<-pl1$B3[Fhijt][,.(IDMancha,com=i.com,com2=com,t,diay,FT,FR,FH,FTRH,xi,ph,prPerc, crecpot=FTRH*xi*ph)]

	k.IM	**model** im:Potential ecology:AboveGroundBiomass **caused by** biology:Growth **in** g/m^2
		'AboveGroundBiomass caused by Potential Growth'
		    observing
		        im:Maximum ecology:Biomass **caused by** biology:Growth **in** g/m^2 **named** xf,
		        percentage of ecology:Vegetation biology:Growth **caused by** *ecology:VegetationLimitingFactor*
		          named ftrh,
		        occurrence of ecology.incubation:PhenologyActivity **named** ph
		    set to [xf*ftrh*ph];

PaL is structured into 10 k.IM code files (namespaces), which integrate multiple data and models
related to climatic growth limitations, vegetation's life cycle, livestock grazing and the nitrogen
cycle. PaL generates spatially explicit outputs at user-specified temporal and spatial scales. In
case the user does not want to change the output characteristics, a set of default output properties
is defined. These features are: spatial resolution of 50 meters, daily time step, time period
between 2018 and 2050. Each model finds its input data on the network, previously annotated
from international and recognized data providers from regional to global scale and from the
literature; the choice of data is done by the k.LAB AI based on fit to the context and the scale
chosen by the user. The user can also provide data to override any of the PaL components, be
them input datasets or computational logics for each of the concepts involved in the model.
Outputs include multiple open-source models, algorithms and spatial outputs of primary interest
to pastureland managers. For example, selected results include the amount of above and below
ground biomass of vegetation, concentration of nitrogen leaching or livestock weight gain. These
outputs can be used for quantitative analysis of pastureland sustainability (or assessment of
farmland requirements and tradeoffs). The set of namespaces in PaL consists of 10 thematic
namespaces that describe the interactions between vegetation, animals and their environment
(Fig. 3, Table 2).

Figure 3 Workflow of PaL namespaces related to climatic growth limitations, vegetation life's cycle, livestock
grazing and nitrogen cycle

Table 2 Description of PaL namespaces related to climatic growth limitations, vegetation life cycle, livestock
grazing and nitrogen cycle.

Namespace	Description
Moisture	All processes involving the limitation of vegetation growth due to soil moisture.
Radiation	All processes involving the limitation of vegetation growth due to solar radiation.
Temperature	All processes involving the limitation of vegetation growth due to atmospheric temperature.
Ingestion	All processes related to grazing and digestion.
Livestock mass	Set of models related to livestock weight change.
Excretion	The process of livestock solid and liquid manure.

Nitrogen	Nitrogen concentration and nitrogen proportions in the N-cycle (including leaching and nitrogen uptake)
Vegetation Growth	Calculation of potential and actual vegetation growth depending on limiting abiotic factors
Senescence	Senescence process and quantity of the remaining, living biomass
Litterfall	Process related to dead plant material (harvesting, litterfall and dead biomass)

Each namespace, in turn, is composed of several model components that each describe one
concept involved in the PaL logical structure, for a total of 198 models that are logically
consistent, self-contained and can run independently. The dependencies between models are
defined at the purely logical level as concepts, and are resolved at the moment of execution by
the k.LAB engine: if needed, the modeler can influence the choice using well-defined scoping
rules. When dependencies cannot be satisfied within the same namespace or project, or within
user-provided data and models, the k.LAB engine will look for ways to satisfy them by looking
up models from the network and ranking them for appropriateness to the context. The ability to
access the entire k.LAB semantic web enacts a fully distributed, interoperable chain of
computation that minimizes the effort involved in producing results without compromising on
quality, transparency or traceability.

In this particular implementation, all models are deterministic, using equations and look-up
tables derived from the literature and expert knowledge. For example, the simplest namespace,
the Radiation module (Table 2), is composed of the "Solar Radiation over Vegetation" model,
which includes three different component models. Each of these sub-models generates an output
and, at the same time, is interoperable with others to generate more complex models, such as the
"Solar Radiation limiting factor causing Vegetation Growth" model (Fig 4).

252
Figure 4 Workflows of "Solar Radiation over Vegetation" and "Solar Radiation limiting factor causing Vegetation
Growth" (diagram on the left). The figure shows the interaction between them and the models that act as inputs. A
total of six models are involved in this process.

In addition, each of these models interact with other namespaces. For example, Figure 5 shows
the model of "*Nitrogen in living aboveground biomass caused by cattle solid manure*" from the
'Excretion' namespace (Table 2), which is composed of three different models that are developed
within other namespaces. For example, "*Proportion of Living AboveGround Biomass in Cattle*
Digestion" is located within the "*Livestock mass*" namespace while '*Proportion of Nitrogen in*
Living AboveGround Biomass 'is in the "*Nitrogen*" namespace and '*Living AboveGround*
Biomass causing Cattle Ingestion' in "*Ingestion.*"

Figure 5 Workflow of "Nitrogen in living above ground biomass caused by cattle solid manure" model from
excretion namespace composed of three models related to namespaces of livestock mass, nitrogen and ingestion.

In this way, each namespace is composed of models that can run independently, unlike Puerto's
original monolithic structure. This semantic-driven interoperability allows each model to
interoperate with models from the same namespace or from different ones, according to the
projects available in the k.LAB resource network and ARIES project. For example, the nitrogen
leaching model can interoperate with a runoff model from an independently developed
hydrological modelling project, automatically connecting knowledge across these projects.
Consistency is maintained through the semantic infrastructure, generating an integrated response
to user queries and scenarios.

PaL namespaces use spatially explicit data (raster and vector) and look-up tables as input files.
Most of the data come from field-validated expert knowledge, including for instance the raster
dataset of main pastureland species. Open-source data from global to local scale with different
temporalities can complement the model when local parameters are missing, such as the raster
data describing soil texture. Based on the user-defined spatial and temporal context, k.LAB
changes the spatial resolution and harmonizes the spatial reference of input data on the fly. Each
input dataset can thus have different spatial and temporal resolution, which are automatically
mediated by the system based on a given user query.

To use the model in dynamic mode, PaL requires climate data for the entire model timeline. The
rest of the inputs are only needed at initialization, because PaL generates the transitions based on
the declared algorithms.

Results
The main result of PaL, the k.LAB-compatible recoded version of Puerto, is the ability to
calculate any of the 198 component models independently and quickly; making them reliably
available to stakeholders with minimal work (depending on the model, from seconds to 6

minutes at 50 meters' spatial resolution). The results generate parameters with self-explanatory
variable names, thanks to the k.IM semantic language (Table 1). Both the data sources and the
algorithms used as inputs for the results are automatically generated, and are publicly available
and downloadable, giving the users additional information to interpret and communicate model
results and maintain quality control (see "End-user features" below).

In the following sections, we describe the main outputs of each PaL namespace for the
Cantabrian Pas, Miera, and Ason watersheds, thus emphasizing the importance of taking a
systems approach in agricultural modelling. The main outputs are temporally explicit raster data
produced on demand for the context of analysis (including the selected spatial and temporal
scales). As the graphical outputs of the Puerto model are limited, predetermined and based on a
monolithic code structure, it is difficult to directly compare all the PaL model results with those
of the original Puerto model. However, we can validate some of the PaL results that directly
match the final Puerto outputs. For this, the Puerto results had to be postprocessed to give them
spatial dimension, the R-generated outputs are not spatially explicit.

The PaL models outputs have been run at the default spatial resolution of 50 meters using mean
climate values for May 2018. The entire list of the models is in the Supplementary Material 1
and the code in Bitbucket repository
(https://bitbucket.org/integratedmodelling/im.ecology.grassland.livestock/).

Model outputs

Factors limiting Vegetation Growth

The "Factors limiting Vegetation Growth" model is composed of three main models: Moisture,
Radiation, Temperature and Nitrogen (Fig. 6). These dynamic models quantify climatic and soil
conditions' control of potential vegetation growth. Vegetation growth follows an annual cycle
influenced by seasonal patterns and extreme weather events. These models thus depend on time
and can help to forecast changes in vegetation behaviour with climate change, as seasons shift
and extreme events become more frequent. Moreover, factors limiting vegetation growth are
affected by the spatial distribution of vegetation, which is influenced for example by the
presence of mountain ranges. These effects are complex: soil characteristics affect water content,
aspect affects shade patterns and the incidence of radiation, and elevation affects the
temperatures and precipitation levels to which plants are exposed.

Figure 6 shows the influence of each variable managed vegetation growth in May 2018. While soil moisture (Fig. 6B) and solar radiation incidence (Fig. 6D) positively affect vegetation growth (except in some shaded areas in the case of solar radiation incidence), temperature (Fig. 6A) has an increasing influence with elevation and nitrogen is the most uniformly limiting factor (Fig. 6C).

Puerto does not provide these results in spatial form. An expert in R can extract the R internal table (Table 3), which contains outputs of the limiting factor for climate. The table indicates:

- the observed plot ("IDMancha"),
- the code of the main ("com") and overstory ("com2") vegetation, in case there is one,
- the timeline, starting at the first of January of the year determined by the modeller,
- the mean parameter corresponding to Temperature ("FT"), Radiation ("FR"), Moisture ("FH") and Nitrogen ("FN") as vegetation limiting factor for each observed plot, vegetation type, time.

336 In this case, users can link the plot identification to a vector dataset to know where the plots are
337 located. We do not know the distribution of vegetation within each plot.

338 *Table 3 Sample Vegetation Limiting Factor output from R table of Puerto model.*

IDMancha	com	com2	t	FT	FR	FH	FN
442	9	9	1	0.701909	0.164	1	0.5
442	14	14	1	0.701909	0.164	1	0.65
442	28	28	1	0.701909	0.164	1	0.65
458	7	7	1	0.701909	0.116	1	0.65
458	13	13	1	0.701909	0.116	1	0.5

Vegetation

The entire vegetation life cycle - including growth, senescence, and litterfall - is composed of
three different namespaces which include 44 component models. Vegetation life cycle is affected
not only by climate, but also by livestock activity, nutrient uptake and human intervention, in
particular by harvesting or fertilization cycles. With PaL, we can estimate the evolution of the
parameters in each grid cell over time, depending on the type of vegetation. This group of
models can be run with or without human and animal influence.

Figure 4 A) Potential Growth causing AboveGround Biomass model in grams/day and B) Growth causing
AboveGround Biomass model in grams/day (Cantabrian case study, May 2018).

Figure 7A shows the potential vegetation growth under climatic factors (temperature, solar
radiation and soil moisture). The results of Figure 7B are the actual growth model, based on
potential growth but also taking into account nitrogen limitation and the influence of livestock on
the grazing areas. Two notable trends emerge – first, that maximum potential daily vegetation
growth is 5.29 grams per day, while actual growth is 1.58 grams per day. Second, the distribution

of vegetation growth is heterogeneous, decreasing in mountainous areas than flatter areas (Figure
7B).

Puerto vegetation growth outputs include tables in R or graphical bar and line graph outputs (Fig.
8). The information on monthly average vegetation growth (bars) and livestock ingestion (line)
are shown for a period of years determined by the modeller, in this case, 5 years. Results are
aspatial, as compared to the spatially explicit outputs for a flexible, user-defined time period in
PaL.

Figure 5 Puerto output (not spatially explicit). Mean monthly vegetation Growth and Ingestion model outputs of
Puerto, for nine pasture types (see Fig. 2).

Livestock

The namespaces related to intake (Ingestion namespace), excretion (Excretion namespace) and
variation of body mass (Livestock mass) of livestock include a total of 55 component models,
including both cattle and mares. Key outputs include sustainability of the exploitation of
pastures, biomass intake, the variation of livestock weight and the amount of excrement returned
to the environment. Based on modelled livestock mass variation for cattle (Fig. 9A) and mares
(Fig. 9B), cattle are more affected by altitude and vegetation availability than mares.

Figure 6 Livestock mass variation of (A) Cattle and (B) Mares in kg/day, May 2018.

Results depend not only on vegetation type and life cycle, but also on the estimated number of
animals on each hectare of land, competition between them, accessibility to the vegetation, and
topography, among other influences.

The Puerto version result for livestock (Figure 10) is the cumulative livestock mass variation per
hectare and year for both mares ("Equino") and cattle ("Vacuno"). Results are aggregated by
grazing unit but are not spatially distributed as in the PaL model.

Figure 7 Cumulative livestock mass variation per hectare and year in Puerto output (no spatially explicit).

Nitrogen Cycle

The nitrogen cycle namespace includes all the models related to nitrogen in its different states
and forms. The calculation of the nitrogen content in senesced leaves, mineral nitrogen present in
the soil, nitrogen in livestock excrement and that used for plants are some of the models called
on by this namespace. An interesting part of this namespace is the "Nitrogen leaching" model
(Fig. 11), which can interact with the models related to the water cycle within k.LAB for future
studies of water quality and pasture management. The output of Puerto is an internal R table as
Table 3.

Figure 8 Nitrogen Leaching output model in grams of nitrogen mass.

<u>End-user features</u>

Output maps

The first set of outputs provided to the end-user is a series of temporally explicit maps.
Temporally dynamic outputs can be viewed using the "play" button at the bottom of the menu on
the left side of Figure 12. A user can also view all the models computed as dependencies of the
requested model. All results (main model and dependent models) can be downloaded in geotiff
format or as an image. In addition, basic information is provided such as total grid size, cell size,
temporality, total observed model area, symbology and colour ramp style with labelling and a
histogram for each of the model's inputs and outputs (Fig. 13).
400

401
402   *Figure 9 Output of the model "Potential above ground biomass caused by growth" and its model dependencies.*

403
Figure 10 Information related to each model output or dependency.

Data flow

k.LAB creates an interactive data flow of the requested model that is built on the fly (Fig. 14).
Thus, all the models and dependencies are shown. By clicking on each block of the data flow,
more information is provided describing:
1) for resources (data sources), basic information about the data source. This is based on
metadata contributed by users who have previously contributed data resources to the
k.LAB network, including links back to the original data source; Fig. 15.A);
2) for tables, each table's composition (Fig. 15.B); and
3) for parameterised models, the expression or algorithm used (Fig. 15.C).

415
Figure 11 Workflow of "Above ground biomass caused by growth" model created on the fly as the model runs.

Figure 12 Additional workflow component views. A) Information related to resources (i.e., data),
B) information related to a look-up table, C) information related to a model algorithm.

Report

A printable report (Fig.14) is also created on the fly, collecting documentation from each model
being run and adapting it to the results being calculated. Basic documentation about each model
component is entered by each model's contributor in k.LAB, which is called when the model is
run and assembled into the report; the modellers' documentation uses a template language that
makes it possible to "react" to the results. This reporting facility complements the workflow
graph in making the system transparent and reliable. The report follows the standard structure of
a scientific article (introduction, methods, results, discussion, conclusion and references). It can
include tables, figures or other elements, depending on the model, and can be downloaded in .pdf
format.

Figure 13 Report describing the model, created on the fly as the model runs.

Discussion

A sustainable balance between agricultural production and healthy ecosystems in agricultural
landscapes has been challenging to achieve. The main difficulties can be linked to population
growth and people's increased demands for food, water and energy, the limited area of arable land
to expand food production and increasing pressure on natural resources from various human
activities (Zhang, Wang & Wang, 2002; Kitzes et al., 2008). These factors are further compounded
by land degradation and water contamination, climate change, sub-optimal agricultural and land-
use policies and market fluctuations (Kendall & Pimentel, 1994; Laurance, Sayer & Cassman,
2014). The PaL models developed in k.LAB can be used to improve the management of
agricultural systems by:

• integrating all the components of agricultural systems modelling in one platform,
• simulating the effects of alternative resource use strategies,
• improving the efficiency of low-input and intensive agricultural systems, and
• improving accessibility and transparency of simulation models to stakeholders.

The divergence in the time scales between farmer choices and environmental goals is a
substantial management challenge. While farmers often need or want to fulfil their financial and
land management objectives in the short term (i.e., months and seasons in this and the following
year), environmental goals may take much longer to be reached (potentially years to decades).
The temporal flexibility in modelling plays a key role to quantify short- and long-term processes
in both the agricultural system and the environment. As we show in this article, the k.LAB
approach ensures semantic consistency in temporal data, from historical observations to future
scenarios, to respond to these needs in different situations. Moreover, the PaL namespaces could
be expanded to simulate environmental disturbances, disease spread, climatic change and
simulated management plans to deal with such challenges, building on the existing PaL

namespaces and without having to change any of them and its models. For example, providing a
model for "change in X" is all it takes to make a previously static model of concept X dynamic,
as the k.LAB engine will automatically insert it in the workflow whenever the context is
computed over multiple timesteps.

Because environmental modelling, including pasture and livestock simulation, tends to be driven
by the need to address case-specific issues, data and model reuse recommendations are often
unclearly defined. Moreover, the collected data are often not made available to other researchers;
when they are placed in public repositories data are often findable and accessible but lag in their
interoperability and reusability (Borycz & Carroll, 2020). As a result, in the best case substantial
manual GIS processing is required before a user can work with previously generated data; in the
worst case data may be lost entirely after the modelling results are published. In this article, we
demonstrate a semantics-first approach to harmonize data and models of livestock and
pastureland, in order to make them interoperable (Villa et al., 2014). Thus, PaL's modular
approach allows models and data to be combined for specific purposes in one platform, making
the simulation process more efficient by representing diverse pieces of knowledge in the same
system, which is a common difficulty in agricultural modelling systems (Harrison et al., 2016).
This is a significant improvement in dealing with the complex interdependencies between
humans and nature in agricultural systems, where data come from different sources and
knowledge domains as in the case study presented.

The models, algorithms, data sources, and results described in this article are accessible to non-
technical users through a web browser application, k.Explorer – a substantial improvement from
the previous edition of the Puerto model, which was only available to technical modelers
proficient in the R programming language. As described in the "End-user features" section of the
Results, this makes scientific information more easily understandable and accessible, bringing
scientific research closer to society with greater transparency (Figs. 12-16). k.LAB is an open
and collaborative technology aiming to expand and improve the availability of interoperable data
and models across disciplines (Willcock et al., 2018; Martínez-López et al., 2019). This
technology can be used to substantially improve agricultural data and models' accessibility,
harmonize them in order to facilitate their wider reuse, improve their quality and consistency.
PaL namespaces are made available to both farmers and policy makers as an open, reusable and
efficient toolbox. Modellers can contribute new data and models and the knowledge to ensure
their appropriate reuse through a dedicated interface (Villa et al., 2014) - a collective effort to
provide stakeholders with the needed tools to face the new challenges in agriculture systems
(Matthews et al., 2007; Verburg, Eickhout & van Meijl, 2008).

The versatility and flexibility of this approach encourages model reusability, which is
particularly valuable to iteratively update assessments as newer or more reliable information
becomes available. Data inputs made available in the k.LAB system can affect PaL modelling
outputs and other ecosystem services models connected through semantics (Fig. 5). Both inputs

and outputs from the PaL namespaces can be automatically reused at different temporal and
spatial scales, ranging from local analysis to national scales.

While this article offers an integrated and semantic modelling adaptation of the original Puerto
model, we note three limitations and complexities for the benefit of future investigations. First,
input data needed to run PaL outside the Cantabrian case study region are available on the
k.LAB network but may not have the same quality or resolution due when relying on global data.
This could affect the reliability of PaL outputs when run outside the Cantabrian region. Hence,
we recommend further validation of model outputs in future applications. Second, the types of
modelled pastureland vegetation and livestock are currently limited to certain classes (Figure 2).
Third, some excessively complicated models (Sun et al., 2016) could be replaced by simpler
ones. This would require more accessible cloud-hosted data, but would simultaneously decrease
computational needs.

This article demonstrated how agricultural modelling can be made more transparent and
accessible. In particular, we showed how to run and produce results from the Pasture and
Livestock (PaL) namespaces in the k.LAB modelling platform, capitalizing on a semantics-first
approach (Villa et al., 2017). We applied this set of models to a case study in the Cantabrian
region of Spain, where complex interactions among vegetation, livestock, and nitrogen need to
be disentangled for improved agroecosystem management. Additional agricultural models can be
incorporated and connected with the currently available PaL namespaces in the future. Some of
these models may expand on other ecological aspects, such as pest, weed and disease spread or
carbon and phosphorus cycling, which are closely linked to nitrogen. Others might expand on the
microeconomics of farm operations, taking into account the cost-efficiency of management
activities given farmers' current economic status. Similarly, the existing namespaces can
incorporate new input data related to vegetation and livestock species. Moreover, further
research could analyse the interactions between PaL namespaces and other ecosystem service
models, to fully capture the complex implications of pasture management patterns (Bagstad et
al., 2014; Balbi et al., 2015; Martínez-López et al., 2019).

Conclusions

The evolution of agriculture and the challenges it faces, both in terms of productivity and
ecological impacts, require focused efforts to design more sustainable agricultural systems. The
case study in Cantabria addresses a set of environmental and agricultural management changes
over the past decades. The current pressure of tourism and the trend of farmland abandonment
are risking the balance between nature and society in these systems. One of the main challenges
of this study was to combine, using a unified yet highly flexible and accessible approach, the
biophysical, technical and management knowledge needed to analyse the current conditions and
explore future trends.

In this article, we break down the original monolithic Puerto model, developed for managing rangelands in the Cantabrian region of Spain, into ten Pasture and Livestock k.LAB namespaces, composed of 198 models. We applied these a fine temporal and spatial scale over the case study area, the Pas, Miera and Ason watersheds in Cantabria, responding to the needs for modelling their extensive agricultural systems. To do so, we first provided insights into current and past agricultural trends derived from literature and expert knowledge regarding to the Cantabrian agroecosystem situation. Next, we developed an open and semantic modelling application for pasture and livestock modelling in the k.LAB platform. This provides stakeholders with an accessible and user-friendly web-browser with that better bridges the gap between technical scientific modelling and land managers. Accessible and context-dependent models can provide solutions for different needs, such as those of i) policy-makers, who can better monitor landscape performance and health, ii) farmers, who can simulate alternative management strategies and potential risks to farming production and devise adaptation strategies, and iii) scientists, who can contribute to greater knowledge reuse and application to on-the-ground decision making.

This article elaborated the importance of overall modelling strategy and design for interoperability and reusability, showing how to improve the ease of use of scientific models and their application to decision making. Within a collaborative modelling system like k.LAB, all models are enhanced through wider community testing, reuse, and application to different contexts. Through wider reuse, models can become increasingly realistic, reliable and useful. This approach is applicable for a wide range of environmental modelling problems, though it is especially suitable for agricultural systems, where underlying data are gathered from different sources and domains, as it facilitates a transdisciplinary scientific approach to complex modelling and management problems.

Acknowledgements
This research is supported by the Basque Government through the BERC 2018-2021 program, by the Ikertzaile Doktoreentzako Hobekuntzarako doktoretza-ondoko Programa, by FPI grant from Spanish Ministry of Economy and Competitiveness MINECO through BC3 María de Maeztu excellence accreditation MDM-2017-0714 and by the EU Interreg Atlantic Area Programme 2014–2020 (EAPA_261/2016 ALICE).

The authors would like to thank Joan Busqué who created and shared the original Puerto model and the team lead by José Barquín at the Hydrological Institute of Cantabria (IHC). Special thanks to Simone Langhans and Ken Bagstad who suggested revisions to the article.

References
Angelov PP, Iglesias JA, Corrales JC (eds.). 2018. *Advances in Information and Communication Technologies for Adapting Agriculture to Climate Change: Proceedings of the International Conference of ICT for Adapting Agriculture to Climate Change (AACC'17),*

November 22-24, 2017, Popayán, Colombia. Springer International Publishing. DOI:

10.1007/978-3-319-70187-5.

Argent RM. 2004. An overview of model integration for environmental applications—

components, frameworks and semantics. *Environmental Modelling & Software* 19:219–

234. DOI: 10.1016/S1364-8152(03)00150-6.

Arp R, Smith B, Spear AD. 2015. *Building Ontologies with Basic Formal Ontology.* MIT Press.

Bagstad K, Villa F, Batker D, Harrison-Cox J, Voigt B, Johnson G. 2014. From theoretical to

actual ecosystem services: mapping beneficiaries and spatial flows in ecosystem service

assessments. *Ecology and Society* 19. DOI: 10.5751/ES-06523-190264.

Balbi S, Prado A del, Gallejones P, Geevan CP, Pardo G, Pérez-Miñana E, Manrique R,

Hernandez-Santiago C, Villa F. 2015. Modeling trade-offs among ecosystem services in

agricultural production systems. *Environmental Modelling & Software* 72:314–326. DOI:

10.1016/j.envsoft.2014.12.017.

Bedia J, Cabañas S, Busqué J. 2009. Productivity and plant diversity are related to a community

functional signature in mountain grasslands. In: 79–82.

Bernués A, Ruiz R, Olaizola A, Villalba D, Casasús I. 2011. Sustainability of pasture-based

livestock farming systems in the European Mediterranean context: Synergies and trade-

offs. *Livestock Science* 139:44–57. DOI: 10.1016/j.livsci.2011.03.018.

Billen G, Lassaletta L, Garnier J. 2014. A biogeochemical view of the global agro-food system:

Nitrogen flows associated with protein production, consumption and trade.

Borycz J, Carroll B. 2020. Implementing FAIR Data for People and Machines: Impacts and

Implications - Results of a Research Data Community Workshop. *Information Services &*

Use 40:71–85. DOI: 10.3233/ISU-200083.

Busqué J. 2014. De la investigación a la práctica: herramientas para gestionar la ganadería de

montaña y los pastos comunales de Cantabria dentro de la política agraria común.

Revista Pastos 6:6–42.

Busqué J, Fernández N, Fernández B. 2006. A decision support tool to design rangeland

sustainable grazing systems. *Sustainable grassland productivity: Proceedings of the*

21st General Meeting of the European Grassland Federation, Badajoz, Spain, 3-6 April,

2006:682–684.

Casasús I, Bernués A, Sanz A, Villalba D, Riedel JL, Revilla R. 2007. Vegetation dynamics in

Mediterranean forest pastures as affected by beef cattle grazing. *Agriculture,*

Ecosystems & Environment 121:365–370. DOI: 10.1016/j.agee.2006.11.012.

Cordell D, Drangert J-O, White S. 2009. The story of phosphorus: Global food security and food

for thought. *Global Environmental Change* 19:292–305. DOI:

10.1016/j.gloenvcha.2008.10.009.

Daugstad K, Mier MF, Peña-Chocarro L. 2014. Landscapes of transhumance in Norway and

Spain: Farmers' practices, perceptions, and value orientations. *Norsk Geografisk*

Tidsskrift - Norwegian Journal of Geography 68:248–258. DOI:

10.1080/00291951.2014.927395.

Donald PF, Green RE, Heath MF. 2001. Agricultural intensification and the collapse of Europe's

farmland bird populations. *Proceedings of the Royal Society B-Biological Sciences*

268:25–29. DOI: 10.1098/rspb.2000.1325.

European Environment Agency. 2004. *High nature value farmland: Characteristics, trends and*

policy challenges. Luxembourg: Office for Official Publications of the European

Communities.

Ewert F, van Ittersum MK, Bezlepkina I, Therond O, Andersen E, Belhouchette H, Bockstaller C,

Brouwer F, Heckelei T, Janssen S, Knapen R, Kuiper M, Louhichi K, Olsson JA, Turpin

N, Wery J, Wien JE, Wolf J. 2009. A methodology for enhanced flexibility of integrated

assessment in agriculture. *Environmental Science & Policy* 12:546–561. DOI:

10.1016/j.envsci.2009.02.005.

Farina A. 2000. The Cultural Landscape as a Model for the Integration of Ecology and

Economics. *BioScience* 50:313–320. DOI: 10.1641/0006-

3568(2000)050[0313:TCLAAM]2.3.CO;2.

Fernández-Giménez ME, Fillat Estaque F. 2012. Pyrenean Pastoralists' Ecological Knowledge:

Documentation and Application to Natural Resource Management and Adaptation.

Human Ecology 40:287–300. DOI: 10.1007/s10745-012-9463-x.

Gibon A. 2005. Managing grassland for production, the environment and the landscape.

Challenges at the farm and the landscape level. *Livestock Production Science* 96:11–31.

DOI: 10.1016/j.livprodsci.2005.05.009.

Harrison PA, Dunford RW, Holman IP, Rounsevell MDA. 2016. Climate change impact

modelling needs to include cross-sectoral interactions. *Nature Climate Change* 6:885–

890. DOI: 10.1038/nclimate3039.

Hayami Y, Ruttan VW. 1971. Agricuitural development: an international perspective. *Johns

Hopkins University Press*:367.

Hendrickson JR, Hanson JD, Tanaka DL, Sassenrath G. 2008. Principles of integrated

agricultural systems: Introduction to processes and definition. *Renewable Agriculture

and Food Systems* 23:265–271. DOI: 10.1017/S1742170507001718.

Henle K, Alard D, Clitherow J, Cobb P, Firbank L, Kull T, McCracken D, Moritz RFA, Niemelä J,

Rebane M, Wascher D, Watt A, Young J. 2008. Identifying and managing the conflicts

between agriculture and biodiversity conservation in Europe–A review. *Agriculture,

Ecosystems & Environment* 124:60–71. DOI: 10.1016/j.agee.2007.09.005.

Hieronymi A. 2013. Understanding Systems Science: A Visual and Integrative Approach.

Systems Research and Behavioral Science 30:580–595. DOI: 10.1002/sres.2215.

van Ittersum MK, Ewert F, Heckelei T, Wery J, Alkan Olsson J, Andersen E, Bezlepkina I,

Brouwer F, Donatelli M, Flichman G, Olsson L, Rizzoli AE, van der Wal T, Wien JE, Wolf

J. 2008. Integrated assessment of agricultural systems – A component-based framework

for the European Union (SEAMLESS). *Agricultural Systems* 96:150–165. DOI:

10.1016/j.agsy.2007.07.009.

Janssen SJC, Porter CH, Moore AD, Athanasiadis IN, Foster I, Jones JW, Antle JM. 2017.

Towards a new generation of agricultural system data, models and knowledge products:

Information and communication technology. *Agricultural Systems* 155:200–212. DOI:

10.1016/j.agsy.2016.09.017.

Jones JW, Antle JM, Basso B, Boote KJ, Conant RT, Foster I, Godfray HCJ, Herrero M, Howitt

RE, Janssen S, Keating BA, Munoz-Carpena R, Porter CH, Rosenzweig C, Wheeler TR.

2017. Brief history of agricultural systems modeling. *Agricultural Systems* 155:240–254.

DOI: 10.1016/j.agsy.2016.05.014.

Kamilaris A, Kartakoullis A, Prenafeta-Boldú FX. 2017. A review on the practice of big data

analysis in agriculture. *Computers and Electronics in Agriculture* 143:23–37. DOI:

10.1016/j.compag.2017.09.037.

Kendall HW, Pimentel D. 1994. Constraints on the Expansion of the Global Food Supply. *Ambio*

23:198–205.

Kitzes J, Wackernagel M, Loh J, Peller A, Goldfinger S, Cheng D, Tea K. 2008. Shrink and

share: humanity's present and future Ecological Footprint. *Philosophical Transactions of*

the Royal Society B: Biological Sciences 363:467–475. DOI: 10.1098/rstb.2007.2164.

Kumar K, Gupta SC, Chander Y, Singh AK. 2005. Antibiotic use in agriculture and its impact on

the terrestrial environment. In: Sparks DL ed. *Advances in Agronomy, Vol 87*. San

Diego: Elsevier Academic Press Inc, 1–54. DOI: 10.1016/S0065-2113(05)87001-4.

Laurance WF, Sayer J, Cassman KG. 2014. Agricultural expansion and its impacts on tropical

nature. *Trends in Ecology & Evolution* 29:107–116. DOI: 10.1016/j.tree.2013.12.001.

Lemaire G, Franzluebbers A, Carvalho PC de F, Dedieu B. 2014. Integrated crop–livestock

systems: Strategies to achieve synergy between agricultural production and

environmental quality. *Agriculture, Ecosystems & Environment* 190:4–8. DOI:

10.1016/j.agee.2013.08.009.

López MM. 2002. Landscapes of northern Spain and pastoral systems. In: Redecker B, Härdtle

W, Finck P, Riecken U, Schröder E eds. *Pasture Landscapes and Nature Conservation*.

Berlin, Heidelberg: Springer, 67–86. DOI: 10.1007/978-3-642-55953-2_5.

López-Sáez JA, Alba-Sánchez F, Robles-López S, Pérez-Díaz S, Abel-Schaad D, Sabariego-

Ruiz S, Glais A. 2016. Exploring seven hundred years of transhumance, climate

dynamic, fire and human activity through a historical mountain pass in central Spain.

Journal of Mountain Science 13:1139–1153. DOI: 10.1007/s11629-016-3885-7.

MacDonald D, Crabtree JR, Wiesinger G, Dax T, Stamou N, Fleury P, Gutierrez Lazpita J,

Gibon A. 2000. Agricultural abandonment in mountain areas of Europe: Environmental

consequences and policy response. *Journal of Environmental Management* 59:47–69.

DOI: 10.1006/jema.1999.0335.

Marcos JB, Lodos SM, Rodríguez-Arango BH. 2003. Estructura, crecimiento y aprovechamiento

de pastos de Puerto Cantábricos invadidos o no por Lecherina (Euphoria Polygalifolia).

Pastos: Revista de la Sociedad Española para el Estudio de los Pastos 33:283–303.

Martínez-López J, Bagstad KJ, Balbi S, Magrach A, Voigt B, Athanasiadis I, Pascual M,

Willcock S, Villa F. 2019. Towards globally customizable ecosystem service models.

Science of The Total Environment 650:2325–2336. DOI:

10.1016/j.scitotenv.2018.09.371.

Matthews RB, Gilbert NG, Roach A, Polhill JG, Gotts NM. 2007. Agent-based land-use models:

a review of applications. *Landscape Ecology* 22:1447–1459. DOI: 10.1007/s10980-007-

9135-1.

Montserrat P, Fillat F. 1990. The systems of grassland management in Spain. In: 37–70.

Parker P, Letcher R, Jakeman A, Beck MB, Harris G, Argent RM, Hare M, Pahl-Wostl C, Voinov

A, Janssen M, Sullivan P, Scoccimarro M, Friend A, Sonnenshein M, Barker D,

Matejicek L, Odulaja D, Deadman P, Lim K, Larocque G, Tarikhi P, Fletcher C, Put A,

Maxwell T, Charles A, Breeze H, Nakatani N, Mudgal S, Naito W, Osidele O, Eriksson I,

Kautsky U, Kautsky E, Naeslund B, Kumblad L, Park R, Maltagliati S, Girardin P, Rizzoli

A, Mauriello D, Hoch R, Pelletier D, Reilly J, Olafsdottir R, Bin S. 2002. Progress in

integrated assessment and modelling1A Summary of a workshop on Integrated

Assessment and Modelling, held at EcoSummit 2000: Integrating the Sciences, Halifax,

June 18–22, 2000. See Costanza and Jorgensen (2001) for a further report on

Ecosummit.1. *Environmental Modelling & Software* 17:209–217. DOI: 10.1016/S1364-

8152(01)00059-7.

Peckham SD, Hutton EWH, Norris B. 2013. A component-based approach to integrated

modeling in the geosciences: The design of CSDMS. *Computers & Geosciences* 53:3–

12. DOI: 10.1016/j.cageo.2012.04.002.

Pe'er G, Dicks LV, Visconti P, Arlettaz R, Báldi A, Benton TG, Collins S, Dieterich M, Gregory

RD, Hartig F, Henle K, Hobson PR, Kleijn D, Neumann RK, Robijns T, Schmidt J,

Shwartz A, Sutherland WJ, Turbé A, Wulf F, Scott AV. 2014. EU agricultural reform fails

on biodiversity. *Science* 344:1090–1092. DOI: 10.1126/science.1253425.

Pimentel D, Harvey C, Resosudarmo P, Sinclair K, Kurz D, Mcnair M, Crist S, Shpritz L, Fitton

L, Saffouri R, Blair R. 1995. Environmental and Economic Costs of Soil Erosion and

Conservation Benefits. *Science* 267:1117–1123. DOI: 10.1126/science.267.5201.1117.

Pimentel D, Stachow U, Takacs DA, Brubaker HW, Dumas AR, Meaney JJ, O'Neil JAS, Onsi

DE, Corzilius DB. 1992. Conserving Biological Diversity in Agricultural/Forestry Systems.

BioScience 42:354–362. DOI: 10.2307/1311782.

Putfarken D, Dengler J, Lehmann S, Härdtle W. 2008. Site use of grazing cattle and sheep in a

large-scale pasture landscape: A GPS/GIS assessment. *Applied Animal Behaviour*

Science 111:54–67. DOI: 10.1016/j.applanim.2007.05.012.

Rosegrant MW, Ringler C, Zhu T. 2009. Water for Agriculture: Maintaining Food Security under

Growing Scarcity. *Annual Review of Environment and Resources* 34:205–222. DOI:

10.1146/annurev.environ.030308.090351.

Strijker D. 2005. Marginal lands in Europe—causes of decline. *Basic and Applied Ecology* 6:99–

106. DOI: 10.1016/j.baae.2005.01.001.

Sun Z, Lorscheid I, Millington JD, Lauf S, Magliocca NR, Groeneveld J, Balbi S, Nolzen H,

Müller B, Schulze J, Buchmann CM. 2016. Simple or complicated agent-based models?

A complicated issue. *Environmental Modelling & Software* 86:56–67. DOI:

10.1016/j.envsoft.2016.09.006.

Tan ZX, Lal R, Wiebe KD. 2005. Global Soil Nutrient Depletion and Yield Reduction. *Journal of*

Sustainable Agriculture 26:123–146. DOI: 10.1300/J064v26n01_10.

Verburg PH, Eickhout B, van Meijl H. 2008. A multi-scale, multi-model approach for analyzing

the future dynamics of European land use. *The Annals of Regional Science* 42:57–77.

DOI: 10.1007/s00168-007-0136-4.

Villa F. 2007. A semantic framework and software design to enable the transparent integration,

reorganization and discovery of natural systems knowledge. *Journal of Intelligent*

Information Systems 29:79–96. DOI: 10.1007/s10844-006-0032-x.

Villa F, Bagstad KJ, Voigt B, Johnson GW, Portela R, Honzák M, Batker D. 2014. A

Methodology for Adaptable and Robust Ecosystem Services Assessment. *PLOS ONE*

9:e91001. DOI: 10.1371/journal.pone.0091001.

Villa F, Balbi S, Athanasiadis IN, Caracciolo C. 2017. Semantics for interoperability of

distributed data and models: Foundations for better-connected information.

F1000Research 6:686. DOI: 10.12688/f1000research.11638.1.

Vries FWTP de, Teng P, Metselaar K (eds.). 1993. *Systems approaches for agricultural*

development: Proceedings of the International Symposium on Systems Approaches for

Agricultural Development, 2–6 December 1991, Bangkok, Thailand. Springer

Netherlands. DOI: 10.1007/978-94-011-2840-7.

Wallach D, Makowski D, Jones J, Brun F. 2019. *Working with Dynamic Crop Models*. Elsevier.

DOI: 10.1016/C2016-0-01552-8.

Wilkinson MD, Dumontier M, Aalbersberg IjJ, Appleton G, Axton M, Baak A, Blomberg N, Boiten

J-W, Santos LB da S, Bourne PE, Bouwman J, Brookes AJ, Clark T, Crosas M, Dillo I,

Dumon O, Edmunds S, Evelo CT, Finkers R, Gonzalez-Beltran A, Gray AJG, Groth P,

Goble C, Grethe JS, Heringa J, Hoen PAC 't, Hooft R, Kuhn T, Kok R, Kok J, Lusher SJ,

Martone ME, Mons A, Packer AL, Persson B, Rocca-Serra P, Roos M, Schaik R van,

Sansone S-A, Schultes E, Sengstag T, Slater T, Strawn G, Swertz MA, Thompson M,

Lei J van der, Mulligen E van, Velterop J, Waagmeester A, Wittenburg P, Wolstencroft

K, Zhao J, Mons B. 2016. The FAIR Guiding Principles for scientific data management

and stewardship. *Scientific Data* 3:1–9. DOI: 10.1038/sdata.2016.18.

Willcock S, Martínez-López J, Hooftman DAP, Bagstad KJ, Balbi S, Marzo A, Prato C,

Sciandrello S, Signorello G, Voigt B, Villa F, Bullock JM, Athanasiadis IN. 2018. Machine

learning for ecosystem services. *Ecosystem Services* 33:165–174. DOI:

10.1016/j.ecoser.2018.04.004.

van Zanten BT, Verburg PH, Espinosa M, Gomez-y-Paloma S, Galimberti G, Kantelhardt J,

Kapfer M, Lefebvre M, Manrique R, Piorr A, Raggi M, Schaller L, Targetti S, Zasada I,

Viaggi D. 2014. European agricultural landscapes, common agricultural policy and

ecosystem services: a review. *Agronomy for Sustainable Development* 34:309–325.

DOI: 10.1007/s13593-013-0183-4.

Zhang N, Wang M, Wang N. 2002. Precision agriculture—a worldwide overview. *Computers

and Electronics in Agriculture* 36:113–132. DOI: 10.1016/S0168-1699(02)00096-0.

768