1 Increased biting rate and decreased *Wolbachia* density in irradiated *Aedes* mosquitoes
2

3 Riccardo Moretti[1]*[%], Elena Lampazzi[1%], Claudia Damiani[2,3], Giulia Fabbri[1,2], Giulia
4 Lombardi[1,2], Claudio Pioli[4], Angiola Desiderio[1], Aurelio Serrao[2,3], Maurizio Calvitti[1]
5
6
1. Biotechnology and Agroindustry Division, ENEA (Italian National Agency for New
Technologies, Energy and Sustainable Economic Development), Casaccia Research Center,
Rome, Italy
2. School of Biosciences and Medical Veterinary, University of Camerino, Camerino (MC), Italy
3. Biovecblok S.r.l, Camerino (MC), Italy
4. Laboratory of Biomedical Technologies, ENEA (Italian National Agency for New
Technologies, Energy and Sustainable Economic Development), Casaccia Research Center,
Rome, Italy

%These authors contributed equally to this work.

*Corresponding author: riccardo.moretti@enea.it

RM: riccardo.moretti@enea.it

EL: elena.lampazzi@enea.it

CD: claudia.damiani@unicam.it

GF: giulia03.fabbri@unicam.it

GL: giulia.lombardi.1993@gmail.com

CP: claudio.pioli@enea.it

AD: angiola.desiderio@enea.it

AS: aurobros@hotmail.it

MC: maurizio.calvitti@enea.it

28
29
30
31
32 **Key words:** *Aedes albopictus*, *Aedes aegypti*, X-ray irradiation, radiation biology, vectorial
33 capacity, blood feeding, host-seeking, biting rate, *Wolbachia*, qPCR, FISH analysis, *Wolbachia*
34 density, SIT
35

Abstract

Background: Releasing considerable numbers of radiation-sterilized males is a promising strategy to suppress mosquito vectors. However, releases may also include small percentages of biting females which translate to large numbers when releases are large.

Currently, the effects of irradiation on the host-seeking and host-biting behaviors have not been exhaustively investigated. Information is also lacking regarding the effects of sterilizing treatment on the endosymbiotic bacterium *Wolbachia*, which is known to affect the vector competence of infected mosquitos.

Methods: To ascertain the effects of irradiation on females, the pupae of two *Aedes albopictus* strains, differing in their natural or artificial *Wolbachia* infection type, and *Ae. aegypti*—which is not infected by *Wolbachia*—were treated with various doses of X-rays and monitored for key fitness parameters and biting behavior over a period of two weeks. The effect of radiation on *Wolbachia* was investigated by qPCR and FISH analysis.

Results: Partial *Ae. albopictus* female sterility was achieved at 28 Gy but the number of weekly bites more than doubled compared to that of the controls. Radiation doses of 35 and 45 Gy completely inhibited progeny production but did not significantly affect the survival or flight ability of *Ae. albopictus* females and caused a tripling of the number of bites per female per week (compared to untreated controls). These results were also confirmed in *Ae. aegypti* after treatment at 50 Gy. *Wolbachia* density decreased significantly in 45-Gray-irradiated females, with the greatest decreases in the early irradiation group (26 ± 2-hour-old pupae). *Wolbachia* density also decreased as adults aged. This trend was confirmed in ovaries but not in extra-ovarian tissues. FISH analysis showed a strongly reduced *Wolbachia*-specific fluorescence in the ovaries of 13 ± 1-day-old females.

Conclusions: These results suggest that, under SIT programs, the vector capacity of a target population could increase with the frequency of the irradiated females co-released with the sterile males due to an increased biting rate. In the context of a successful suppression, the related safety issues could be generally negligible, but they should be conservatively evaluated when large scale programs relying on imperfect sexing and high overflooding release ratios are run for long time in areas endemic for arboviral diseases. Also, the effects of irradiation on the vector competence deserve further investigation.

Background

Despite countless attempts, efforts to eliminate vector-borne diseases rarely produce lasting results. In recent years, some vectors have rapidly adapted to the dramatic environmental changes driven by global warming, urbanization, and deforestation, have increased in invasiveness, and have rapidly developed resistance to most pesticides [1–3].

Owing to their reproductive potential and ability to rapidly spread both by migration and passive transportation, mosquitoes are prominent vectors. Substantial investment and research are necessary for the development of prevention and control measures.

The negative effects of widespread insecticide use have caused many researchers to focus on developing innovative control strategies characterized by high specificity and eco-compatibility, and targeting reductions in reproductive potential or vector competence of mosquito populations. This often involves the release of modified conspecifics who mate with wild types and introduce factors that induce sterility, lethality, or virus resistance in the progeny. To establish such modifications, which may or may not be heritable, advanced biotechnological methods based on genetic modification or the exploitation of beneficial microorganisms are used [4]. Methods based on genetic modification can be targeted to specific genes using various biotechnological tools [5–7] but such approaches can be only employed in countries where the release of GMOs is legal. Sterile Insect Technique (SIT) involves the use of optimized doses of mutagenic radiation to sterilize laboratory-reared males (through non-specific genetic modifications), and releasing large numbers of sterilized males to decrease reproduction in wild populations [8, 9].

A similar result can be obtained via a natural phenomenon of egg inviability induced by the common endosymbiotic bacterium *Wolbachia* [10]. Insect males infected by certain strains of this bacterium are only reproductively compatible with females harboring the same *Wolbachia* strain, whereas the lack of the infection or infection with a non-compatible strain causes the production of inviable offspring. This natural post-mating reproductive barrier is known as Cytoplasmic Incompatibility (CI) and may contribute to the spread of infected individuals in the wild, as infected females can produce viable progeny with both infected and uninfected males (Unidirectional CI, Uni-CI) [11,12]. The possibility of transferring the bacterium horizontally between species allowed researchers to exploit *Wolbachia* to produce incompatible males that can be released in a target area to reduce the reproductive potential of wild populations (Incompatible Insect Technique, IIT) [13–18].

Large-scale implementation of SIT and IIT must be combined with an efficient method of

sexing since the presence of residual females inadvertently released with males can result in the

occurrence of temporary spikes in vector density [13]. On the long term, the suppression effect of

control programs is expected to counterbalance the contribution of irradiated females to the mean

biting rate in the target area. However, even small percentages of co-released females may translate

to large numbers of individuals when high overflooding release ratios (10:1 or even 50:1) are

applied for long time [19]. Furthermore, especially when *Wolbachia*-induced Uni-CI occurs,

released females may be invasive, possibly leading to undesired population replacement [12, 20].

The implementation of SIT using mosquito strains capable of producing incompatible males

has been suggested as a method to mitigate the unpredictable effects of female co-release. Mosquito

females are more sensitive to radiation than males and this approach could utilize CI to achieve full

male sterilization, with the radiation dose lowered to levels that exert no effect on male fitness but

which are sufficient to induce complete sterility in females escaping the sexing procedure [21].

However, although recent trials based on this approach have been successful in terms of local *Ae.*

albopictus suppression, the strategy has involved the higher radiation doses typical of SIT alone

applied on a large scale. This has led to considerably reduced male mating competitiveness without

avoiding the release of a few fertile females [19, 22, 23].

It should be noted that all studies aimed at implementing SIT against vector mosquitoes

have been mainly focused on radiation doses aimed at the sterilization of the males [9, 24].

However, the radiation doses needed to achieve female sterilization are known to damage the

ovarian tissues considerably [21] and previous studies have reported that oogenesis plays a role in

the regulation of the tendency to bite and blood-feed [25, 26], which are critical factors for

determining the vectorial capacity of mosquitoes [27]. Therefore, an appropriate study on the

effects of radiation on these physiological and behavioral traits is warranted.

A thorough investigation on the effects of irradiation on *Wolbachia* would also be necessary

in the case of SIT or SIT-IIT combined strategy. In fact, the bacterium *Wolbachia* has been shown

to be affected by radiation [28, 29]; thus, this aspect should also be considered before conducting

large-scale control programs which involve *Wolbachia* infected mosquitoes as the titer of this

endosymbiont may be related to the vector competence of the host [30] and the level of induced CI

[31].

Together with certain anopheline species, *Aedes* mosquitoes represent the main concern for

human health as serve as vectors for several arboviruses. The impressive spread of diseases

associated with these pathogens in recent decades highlights the inefficacy of the current control

methods [32–34]. *Ae. albopictus* and *Ae. aegypti* have been the target of several SIT, IIT or SIT/IIT

combined experimental trials in recent years [14, 16-19, 35]. The first species is naturally infected

with two *Wolbachia* strains that have been demonstrated to interfere with pathogen transmission

(compared to the conditions in uninfected individuals) [36] while the latter is not infected by

Wolbachia in nature.

Herein, we aimed to determine whether irradiation interfered with the host-biting and host-

seeking behaviors of *Ae. albopictus* and *Ae. aegypti*. Furthermore, the effects of radiation on the

titer of *Wolbachia* in *Ae. albopictus* were evaluated in whole mosquito bodies using qPCR

(quantitative polymerase chain reaction), after applying the treatment at different pupal ages and

analyzing the effects at two adult ages. Additionally, ovaries and extra-ovarian tissues were

similarly studied. Finally, FISH (fluorescence in situ hybridization) analysis was conducted on the

irradiated ovaries to visualize the effects of the treatment on the bacterial population and to acquire

further information for interpreting the results. Two *Ae. albopictus* lines have been used in the

experiments to determine whether different *Wolbachia* strains might affect the results. The resultant

data will be useful for enhancing SIT and SIT-IIT strategies in terms of safety and sustainability,

and to better evaluate their large-scale applicability.

Materials and methods

Ae. albopictus **and** *Ae. aegypti* **strains and rearing**

Two *Ae. albopictus* strains and one *Ae. aegypti* strain were used in the experiments. S_{ANG} *Ae.*

albopictus originated from wild-type individuals collected from Anguillara Sabazia (Rome) in 2006

and harbors *w*AlbA and *w*AlbB *Wolbachia*. AR*w*P *Ae. albopictus* was established in 2008 through

the transinfection of *Wolbachia*-cured S_{ANG} individuals with *w*Pip *Wolbachia* from *Culex pipiens*

molestus [37] and is characterized by a bidirectional incompatibility pattern with wild-type *Ae.*

albopictus [38]. Both lines were reared under laboratory conditions at ENEA-Casaccia Research-

Center (Rome) and were periodically outcrossed with wild-type individuals from the same area to

preserve genetic variability [39]. The *Ae. aegypti* line (New Orleans, LA 2011) was provided by the

University of Camerino (Camerino, MC, Italy) where it had been laboratory-reared since 2014 and

is not infected with *Wolbachia*.

Colonies were maintained by raising larvae to adulthood inside 1-L larval trays at a density

of 1 larva/mL, provided with liquid food according to the methods described in a previous study

[39]. Adult mosquitoes were maintained inside $40 \times 40 \times 40$ cm cages at T = 28 ± 1 C°, RH =

70%±10%, and L:D = 14:10 h, and were supplied with water and 10% sucrose.

Blood meals were provided via anesthetized mice in agreement with the Bioethics

Committee for Animal Experimentation in Biomedical Research and in accordance with procedures

approved by the ENEA Bioethical Committee according to the EU directive 2010/63/EU. The mice

belonged to a colony housed at CR ENEA Casaccia and maintained for experimentation based on

the authorization N. 80/2017-PR released (on February 2, 2017) by the Italian Ministry of Health.

Feeding female mosquitoes on the blood of human hosts (i.e., the Authors RM, EL, GL, and MC)

during the experiments was also approved by the ENEA Bioethical Committee.

Radiation methods

Cohorts of *Ae. albopictus* and *Ae. aegypti* females belonging to the strains described above were

irradiated with X-rays to enable comparison with sham-exposed individuals (control).

Ae. albopictus pupae were sexed mechanically using a specific sieving procedure described

previously [16].

X-ray irradiation was performed using the Gilardoni CHF 320G X-ray generator (Gilardoni

S.p.A.; Mandello del Lario, Lecco, Italy) operated at 250 kVp, and 15 mA, with filters of 2.0 mm of

Al and 0.5 mm of Cu, furnished by the Physical Technologies for Security and Health Division of

ENEA. Depending on the experiment and according to doses already tested for the radiation-based

sterilization of the two species [9, 19, 21], *Ae. albopictus* pupae were subjected to 28, 35, and 45 Gy

(dose rate: 0.868 ± 0.004 Gy/min, mean ± SD), while a single dose of 50 Gy was used to treat *Ae.*

aegypti, as this dosage is known to fully inhibit egg production in the species [9]. Time of sample

exposure was determined according to dose rate in order to obtain the pre-established doses. Doses

were confirmed by monitoring the exposure with a PTW 7862 large-size plane parallel transmission

chamber connected to a PTW IQ4 electrometer. Groups of 100 female pupae were transferred to a

Petri dish (d= 4 cm) at 36±4 h of age (unless specified differently) and then transported to the

irradiation facility. Immediately before commencement of irradiation, most of the residual water

was removed using a glass pipette; irradiated pupae were then transferred to a larger water container

to facilitate complete development and allow for adult emergence inside the experimental cages.

Sham-exposed pupae, i.e., pupae treated in a manner similar to the exposed pupae except for the X-

ray exposure, were used as controls.

Survival, fecundity, and fertility in irradiated *Ae. albopictus* females

S_{ANG} or ARwP *Ae. albopictus* females irradiated at 28, 35, and 45 Gy, and untreated counterparts were allowed to emerge inside $30 \times 30 \times 30$ cm plastic cages, which were checked for the presence of males that escaped the sexing procedure. Thirty virgin females and thirty untreated males—characterized by the same *Wolbachia* infection type—were placed in each cage.

These cages were used to monitor survival, fecundity, and fertility in the same treatments during the two weeks of observation, and for the biting rate studies described below. Five repetitions were conducted.

Mortality was recorded daily by removing dead individuals. During the two observation weeks, egg collection was initiated three days after the first blood meal and collection was stopped on the fifth day after the last meal to limit overlap with the second gonotrophic cycle. Paper-lined cups for egg collection were replaced every three days to avoid uncontrolled egg hatching. Eggs were maintained under wet conditions for three days, allowed to dry, and counted to determine the fecundity rate. Egg fertility was assessed by counting the hatched eggs after immersion in a nutrient broth [40].

Engorgement rate in irradiated *Ae. albopictus* and *Ae. aegypti* females under small enclosures

Starting from the fifth day after emergence and continuing for five days, irradiated S_{ANG} and ARwP females were offered a daily blood meal to monitor their feeding behavior. To allow for the study of a second gonotrophic cycle, this same procedure was also conducted from the 12th to the 16th day. The number of engorged females was tallied within 20 min of placement of the blood meal in the cage. Results obtained with the two populations of females treated at the three radiation doses were compared with those observed using control untreated females. Each of these eight treatments was repeated five times.

Although the present study was focused on *Ae. albopictus*, to generalize the experiment, *Ae. aegypti* females irradiated at 50 Gy were similarly tested (in triplicate) in comparison with untreated controls. The hatch rate of irradiated *Ae. aegypti* was also measured and compared to that of control females to identify a successful radiation treatment.

Host-seeking ability and biting rate of irradiated *Ae. albopictus* females in large enclosures

The host-seeking behavior of ARwP and S$_{ANG}$ females irradiated at 45 Gy was studied in large

enclosures and compared with that of untreated individuals based on methods described in a

previous study [41]. The trials were conducted outdoor in two large cellular polycarbonate

experimental units (LEU, $8.5 \times 5 \times 5$ m L : W : H) with two large lateral openings (L = 8 m; H = 1

m) protected by a mosquito net to promote ventilation and ensure natural climatic conditions. The

latter were constantly monitored by a CR-10 data logger (Campbell Scientific, Logan, UT, USA)

which registered a mean temperature of 34.0 ± 1.0 °C and RH averaging $50.0\% \pm 5.0\%$. LEUs

contained benches with wet soil and potted plants which provided refugees for the females and

higher local levels of humidity (averaging $61.0\% \pm 5.0\%$).

For each experiment, an experimenter (the host) as acted as a source of blood and a second

experimenter (the collector) released, recovered, and counted the *Ae. albopictus* females after they

had fed on the host. The host wore a long-sleeved shirt and short pants, exposing only the lower

legs to limit the area for the mosquitoes to land on, whereas the collector wore a white tracksuit and

white shoes. Both researchers wore mosquito net hats. Four groups, differing in their infection type

(S$_{ANG}$ or ARwP) and treatment (irradiated and non-irradiated individuals), were compared. Each

group consisted of 30 starved females aged 6 ± 1 days. Females were released at the side of the LEU

by removing the cover of the cage. The second experimenter then immediately approached the host

on the opposite side of the LEU to collect the females that had started feeding. The proportion of

blood-fed females was tallied along with the time taken by each individual to reach the host, at 30-

second intervals. Female mosquitoes that landed on the host, but did not bite were excluded from

the count. The experiments lasted for a period of 15 min, and for each group, six repetitions were

performed, alternating the experimental units and the host. At the end of each experiment, an

electric mosquito swatter and a powerful aspirator were used to eliminate any mosquitoes remaining

inside the LEU. The experiments were conducted in the late afternoon on sequential days during

June 2020. A schematic describing the experiment is provided in the supplemental materials

(Additional file 1: Figure S1).

In addition to the methods described above, 6 ± 1-day-old females from each treatment group

were engorged in the laboratory and used to study the effect of the radiation treatment on their

willingness to seek and bite a host 48 h after the first blood meal. After their release in the LEU, the

proportion of feeding females and time to reach the host were again noted.

Quantitative PCR analysis of *Wolbachia* titer in irradiated *Ae. albopictus* females

Considering the importance of *Wolbachia* in modulating the vector competence of infected

mosquitoes [42], a qPCR was performed on the strains of this bacterium present in S_{ANG} and AR*w*P

females after irradiation at 45 Gy (a dose known to induce full female sterilization in the species)

[9]. Results were compared with those obtained using untreated counterparts. In operational SIT

programs, mosquitoes are generally irradiated at the pupal stage (24-48 h); however, it is known

that pupal age is one of the critical factors that affects the biological response to the radiation dose

[43]. Therefore, the effects of the irradiation were investigated by analyzing DNA extracts of whole

bodies of 6±1-day-old females developed from pupae irradiated at three ages (26±2, 36±2 and 46±2

h).

The density of *Wolbachia* is known to vary with the age of *Ae. albopictus* females [44].

Therefore, to investigate the effects of radiation on the bacterium in the germ line and somatic

tissues, the titer of the bacterium was measured in ovaries and bodies lacking ovaries of 6±1- and

13±1-day-old females after they were treated with 45 Gy as pupae aged 36±4 h. Untreated females

were used as a control.

DNA was extracted from single individuals using the ZR Tissue & Insect DNA Kit

MicroPrep (Zymo Research, Irvine, CA, USA), according to manufacturer's instructions. When

necessary, females were chilled on ice and dissected in phosphate-buffered saline (PBS) to isolate

the ovaries from the other tissues. Here we excluded individuals with ovaries that were not intact.

Real-time PCR was performed using the Roche LightCycler 96 Instrument (Roche

Molecular Systems, Inc., Rotkreuz, Switzerland). Each reaction was performed in triplicate with a

final reaction volume of 20 μL (10 μL of the Luna Universal qPCR Master Mix (New England

BioLabs, Ipswich, MA, USA), 0.03 μL of 150 nM each primer, and 2 μL of purified DNA) using

the following amplification program: initial activation at 95 °C for 120 s, followed by 40 cycles at

95 °C for 15 s and 60 °C for 40 s. The presence of specific amplification products was verified

using dissociation curves [44].

Strain-specific primers were used to amplify the *wsp* loci [45], namely, the *w*AlbA-*wsp* and

*w*AlbB-*wsp* loci in the case of S_{ANG} females, and the *w*Pip-*wsp* loci in the case of AR*w*P specimens.

Wsp plasmid standards were used to generate a standard curve [44]. The *Ae. albopictus* actin gene

was used as a reference for whole body extracts and extracts obtained after removal of the ovaries

and was amplified with the primer pair actAlbqPCR [46]. Owing to the marked decrease in actin

gene copies observed preliminarily in irradiated ovaries (Additional file 2: Figure S2), the

normalization of qPCR data related to these organs was performed using total DNA (2 μL of

purified DNA per reaction, corresponding to 200-300 ng) as reference [47]. For this purpose,

NanoDrop 2000 spectrophotometer (Thermo Fisher, Waltham, MA, USA) was used.

FISH analysis of the ovaries in irradiated *Ae. albopictus* females

Based on the results of qPCR, fluorescent in situ-hybridization (FISH) analysis was conducted on

the ovaries of 13±1-day-old irradiated and untreated females according to the protocol described by

previous authors [48]. Two hundred nanogram of the *Wolbachia* specific 16S rRNA probe (W2: 5'-

CTTCTGTGAGTACCGTCATTATC-3') was added in the hybridization buffer [43]. Tissues were

placed on a slide containing a drop of the VECTASHIELD Antifade Mounting Medium with DAPI

(Vector Laboratories, Burlingame, CA, USA) and visualized using the Nikon Eclipse E800

confocal microscope and NIS-Elements 4.0 software (Nikon, Tokyo, Japan). An aposymbiotic

population of *Ae. albopictus* [37] was used as the negative control.

Data analysis

Results were expressed as mean ± SE, and the arcsine square root transformation was applied to

analyze proportional data. The Levene's test and the Shapiro-Wilk test were performed to assess

equality of variances and normality, respectively. Statistical analysis was performed using the

software PASW statistics (PASW Statistics for Windows, Version 18.0. SPSS Inc., Chicago, IL,

USA) with the level of significance set at $P < 0.05$.

The survival curves of the four different treatments for each *Wolbachia* infection type, were

compared using the Kaplan-Meier method and the log-rank (Mantel-Cox) test. The Kruskal-Wallis

H-test followed by the Conover-Iman test was used to compare fecundity and egg hatch data

between treatments within each infection type.

Repeated measures ANOVA (Analysis of Variance) was used to analyze the bite data

obtained between treatments over each week. Cages were considered the experimental units and

data have been expressed as the mean percentage of engorged females per day, adjusted for cage-

specific mortality. If Mauchly's test indicated a violation of the sphericity assumption, the degrees

of freedom were corrected by using Huynh-Feldt estimates. Multiple comparisons between

treatments were assessed using Tukey's HSD post hoc test. Additionally, one-way ANOVA was

performed to compare the mean number of bites per female per treatment during each week of

study.

Data regarding the host-seeking behavior under large enclosures were analyzed by assigning

a value to each mosquito based on the median time of each catching interval to compute the average

time to landing. The proportion of engorged mosquitoes was measured based on the ones retrieved

within the defined 15-minute interval. Two-way ANOVA was used to analyze the differences

between groups in terms of the proportions of females that had bitten the host and to compare

average landing times. The Shapiro-Wilk test was conducted to ascertain that the proportions and

average landing times were normally distributed.

One-way ANOVA was used to compare within each *Wolbachia* strain, the qPCR data

obtained from *Ae. albopictus* females treated at the three tested pupal ages or untreated. The effect

of female aging on the overall titer of *Wolbachia* was also analyzed by performing one-way

ANOVA within data of each strain of the bacterium. Data regarding ovaries and extra-ovarian

tissues of the treated or untreated counterparts were similarly analyzed at the two tested female

ages. In the case of rejection of the assumptions of equality of variance and/or normality, the

Kruskal-Wallis rank-sum test was performed.

Results

Female survival, fecundity, and fertility in irradiated *Ae. albopictus* females
345
Regardless of the infection type, irradiation slightly reduced the survival of female mosquitoes but

this difference was not significant between treatments during the two weeks of observation (Fig 1;

Log rank test for S_{ANG}: $\chi^2 = 2.43$; df = 3; P = 0.49; Log rank test for ARwP: $\chi^2 = 1.847$; df = 3; P =

0.61).

A 45-Gray dose completely inhibited egg laying in *Ae. albopictus*, whereas a few eggs were

oviposited by females irradiated at 35 Gy. Among the latter, fertile females were very rare (Table

1). A dose of 28 Gy induced considerable reduction in the number of oviposited eggs compared to

that in untreated controls and markedly affected egg fertility, but did not fully inhibit progeny

production (Table 1). These results were confirmed in both *Ae. albopictus* strains and both

gonotrophic cycles.

Table 1. Fecundity and fertility of ARwP and S_{ANG} *Ae. albopictus* females irradiated at 28, 35, and 45 Gy compared to those of untreated counterparts over two gonotrophic cycles. Fecundity and egg hatching data related to *Ae. aegypti* irradiated at 50 Gy are also reported.

Mosquito line	Blood meal	Treatment	Mean (±SE) eggs per female	Mean (±SE) egg hatch (%)
ARwP	1st	untreated	31.54 ± 0.77 a	85.87 ± 2.37 a
		28 Gy	5.67 ± 0.36 b	13.72 ± 1.77 b
		35 Gy	0.11 ± 0.04 c	0.00 ± 0.00 c
		45 Gy	0.00 ± 0.00 d	0.00 ± 0.00 c
	2nd	untreated	30.89 ± 0.79 a	85.61 ± 2.28 a
		28 Gy	5.04 ± 0.46 b	15.03 ± 1.91 b
		35 Gy	0.15 ± 0.04 c	0.00 ± 0.00 c
		45 Gy	0.00 ± 0.00 d	0.00 ± 0.00 c
S_{ANG}	1st	untreated	32.24 ± 0.82 a	87.44 ± 2.38 a
		28 Gy	5.45 ± 0.44	14.75 ± 2.28
		35 Gy	0.13 ± 0.04 c	0.00 ± 0.00c
		45 Gy	0.00 ± 0.00 d	0.00 ± 0.00 d
	2nd	untreated	31.44 ± 0.79 a	86.77 ± 2.40 a
		28 Gy	5.27 ± 0.38 b	15.25 ± 1.60 b
		35 Gy	0.13 ± 0.03 c	2.86 ± 2.86 c
		45 Gy	0.00 ± 0.00 d	0.00 ± 0.00 d
Ae. aegypti	1st	50 Gy	0.00 ± 0.00	0.00 ± 0.00
	2nd	50 Gy	0.00 ± 0.00	0.00 ± 0.00

ARwP = wPip *Wolbachia* infected *Ae. albopictus*;

S_{ANG} = wild-type *Ae. albopictus*;

Within each group and column, values followed by a different letter are significantly different (Kruskal-Wallis test followed by the Conover-Iman test (P < 0.05)).

Increased engorgement rate in irradiated *Ae. albopictus* and *Ae. aegypti* females

Repeated measures ANOVA revealed significant differences between treatments with respect to the weekly biting activity (Fig. 2A; S_{ANG}-1st wk: Huynh-Feldt correction, $F_{(10.19, 54.35)} = 2.99$, $P < 0.05$; S_{ANG}-2nd wk: Sphericity assumed, $F_{(12, 64)} = 4.52$, $P < 0.05$; ARwP-1st wk: Sphericity assumed, $F_{(12, 64)} = 5.90$, $P < 0.05$; ARwP-2st wk: Huynh-Feldt correction, $F_{(9.12, 48.64)} = 3.36$, $P < 0.05$). Regardless of the week and the infection type, untreated mosquitoes generally showed most biting activity during the first day of the blood meal and then they generally rested while their eggs underwent maturation. In contrast, irradiated females exhibited an anomalous blood feeding behavior and the percentage of the daily engorged females only decreased slightly over each week (Fig. 2A).

For the 35- and 45-Gray treatments, total bites per female approximately tripled weekly over

those recorded for untreated females (Fig. 2B). During the first week, the mean number of bites per

S_{ANG} female averaged 1.15 ± 0.07 (untreated), 2.14 ± 0.06 (28 Gy), 2.69 ± 0.07 (35 Gy), and $2.77 \pm$

0.07 (45 Gy). Tukey's test showed significant differences among untreated and treated females, and

among the females treated with 28 Gy and the other three groups ($F_{3,16} = 118.01$; $P < 0.05$). During

the second week, the above-mentioned values were 1.07 ± 0.05 (untreated), 1.89 ± 0.05 (28 Gy),

2.63 ± 0.03 (35 Gy), and 2.84 ± 0.06 (45 Gy) and all treatments differed significantly from one

another ($F_{3,16} = 282.23$; $P < 0.05$). With respect to the AR*w*P line, the mean number of bites per

female was 1.10 ± 0.03 (untreated), 2.15 ± 0.02 (28 Gy), 2.68 ± 0.03 (35 Gy), and 2.84 ± 0.02 (45

Gy) during the first week ($F_{3,16} = 371.96$; $P < 0.05$), with the Tukey's test indicating significant

differences between all treatments. Values of 1.05 ± 0.03 (untreated), 2.12 ± 0.02 (28 Gy), $2.70 \pm$

0.03 (35 Gy), and 2.76 ± 0.04 (45 Gy) were obtained during the second week ($F_{3,16} = 642.89$; $P <$

0.05), with significant differences observed between all treatments except those at 35 and 45 Gy.

Similarly, 50 Gy-irradiated *Ae. aegypti* females showed an enhanced propensity to bite

compared to the untreated mosquitoes during both weeks of observation (Fig. 2C; 1st wk:

Sphericity assumed, $F_{(4, 16)} = 8.38$, $P < 0.05$; 2nd wk: Sphericity assumed, $F_{(4, 16)} = 11.51$, $P < 0.05$).

Overall, the mean number of bites for the 50-Gray-treated females almost tripled that of untreated

mosquitoes during the first (2.81 ± 0.03 and 1.03 ± 0.02; $F_{1,4} = 3315.67$; $P < 0.05$) and second

weeks (2.86 ± 0.04 and 1.05 ± 0.03; $F_{1,4} = 1522.09$; $P < 0.05$) (Fig. 2D). The applied radiation dose

was found to induce complete sterility in *Ae. aegypti* females (Table 1).

Host-seeking behavior of irradiated *Ae. albopictus* females in large enclosures

The host-biting behavior exhibited by the irradiated females in large enclosures reinforced the

results obtained in small cages. The applied X-ray dose was not sufficient to reduce the ability of

Ae. albopictus to reach the host in the LEU (compared to untreated females; Fig. 3A, B). The

average times to landing were not significantly affected by the infection type (two-way ANOVA:

$F_{1,20} = 0.66$, $P = 0.43$), nor were they affected by the treatment (two-way ANOVA: $F_{1,20} = 0.93$, $P =$

0.35) in starved females. On average, landing times ($\pm$SE) were 346.31 ± 15.98 and 354.30 ± 10.74

in irradiated and untreated S_{ANG} females and 331.11 ± 12.74 and 348.29 ± 12.30 in irradiated and

control AR*w*P individuals, respectively.

In contrast, the biting behavior was affected by irradiation. Starved females did not reveal a

significant difference in the proportion of biting females between untreated and 45-Gray-treated

females (two-way ANOVA: $F_{1,20} = 0.59$, $P = 0.45$) or between infection types (two-way ANOVA:

$F_{1,20} = 0.80$, $P = 0.38$) (Fig. 3A). Instead, the proportion of fed females that repeated the blood meal

after 48 h was significantly higher in irradiated mosquitoes than that in untreated mosquitoes (two-

way ANOVA: $F_{1,20} = 276.53$, $P < 0.05$), while the infection type showed no significant effect (two-

way ANOVA: $F_{1,20} = 1.13$ $P = 0.30$) (Fig. 3B). Consequently, a comparison between average

landing times related to the engorged females was not performed due to the reduced number of

biting individuals among the untreated controls (0 in certain repetitions). To verify this result in

older females that might have experienced the occurrence of age-related damage due to irradiation,

the test was also repeated in triplicate with treated and untreated individuals aged 13±1 days and

similar results were obtained (Additional file 3: Figure S3).

Decreased titer of *Wolbachia* in irradiated *Ae. albopictus* females

The age at which *Ae. albopictus* pupae had been irradiated at 45 Gy was found to significantly affect the density of *Wolbachia* (*w*AlbA: $F_{(3, 36)}$=2.96, $P < 0.05$; *w*AlbB: $F_{(3, 36)}$=4.05, $P < 0.05$; *w*Pip: $F_{(3, 36)}$=6.76, $P < 0.05$) regardless of the *Wolbachia* infection type. An early irradiation (26 ± 2-hour-old pupae) resulted in significantly decreased the whole-body *Wolbachia* titer, by a third in *w*AlbA, and by half in *w*AlbB and *w*Pip strains (Fig. 4A) compared to the levels observed in the controls. The *Wolbachia* titer did not significantly differ in pairwise comparisons between treated and untreated individuals when irradiation was performed on older pupae (36 and 46 ± 2-hours old).

Aging of adult females was also found to affect the titer of all the tested *Wolbachia* strains when a 45-Gray treatment was performed (Fig. 4B). The difference between treatments was not significant when the analysis had been performed on 6 ± 1-day-old females (one-way ANOVA; *w*AlbA: $F_{(1, 18)} = 1.54$, $P = 0.23$; *w*AlbB: $F_{(1, 18)} = 0.82$, $P = 0.89$; *w*Pip: $F_{(1, 18)} = 3.28$, $P = 0.09$), but the differences increased significantly in females aged 13 ± 1 days (one-way ANOVA; *w*AlbA: $F_{(1, 18)} = 62.11$, $P < 0.05$; *w*AlbB: $F_{(1, 18)}$=38.93, $P < 0.05$; *w*Pip: $F_{(1, 18)}$=4.60, $P < 0.05$).

In agreement with the data obtained from the analysis of whole bodies, qPCR revealed a correlation between aging and a reduced titer of the bacteria in the ovaries of irradiated *Ae. albopictus* females compared to that in untreated *Ae. albopictus* females (Fig. 5A). Differences were significant in females at 13 ± 1 days after the irradiation (one-way ANOVA; *w*AlbA: $F_{(1, 18)} = 62.11$, $P < 0.05$; *w*AlbB: $F_{(1, 18)} = 5.91$, $P < 0.05$; *w*Pip: $F_{(1, 18)} = 4.60$, $P < 0.05$), whereas the density of *w*AlbA and *w*AlbB *Wolbachia* strains did not differ between treatments in 6 ± 1 day-old females (one-way ANOVA; *w*AlbA: $F_{(1, 18)} = 0.39$, $P = 0.54$; *w*AlbB: $F_{(1, 18)} = 0.50$, $P = 0.49$). A significant difference was found between the *w*Pip *Wolbachia* titer in treated and untreated 6 ± 1-day-old AR*w*P females ($F_{(1, 18)} = 5.25$, $P < 0.05$). The titer of *w*AlbA and *w*AlbB *Wolbachia* also increased with aging in the ovaries of control females and confirmed the results of previous studies (Fig. 5A) [40]. With respect to *w*Pip, the controls exhibited an age-dependent decrease in titer and irradiation further intensified this trend in the density of the bacteria.

When testing females without ovaries, the qPCR amplification results of *Wolbachia* (somatic fraction) did not reveal differences between treatments related to the titers of *w*AlbA, *w*AlbB, and *w*Pip *Wolbachia* when irradiated females had been tested at 6 ± 1 days (Fig. 5B) (Kruskal Wallis test; *w*AlbA: $\chi2 = 1.85$, $df = 1$, $P = 0.17$; *w*AlbB: $\chi2 = 0.32$, $df = 1$; $P = 0.57$; *w*Pip: $\chi2 = 1.37$, $df = 1$, $P = 0.24$). In contrast with the results obtained in the ovaries, the titer of *Wolbachia* in the extra-ovarian tissues exhibited an apparent age-related increase in irradiated females compared to the controls but

447 differences between treatments were not significant (Kruskal Wallis test; wAlbA: $\chi2 = 2.17$, df = 1, P =

448 0.14; wAlbB: $\chi2 = 2.52$, df = 1, P = 0.11; wPip: $\chi2 = 0.24$, df = 1, P =0.62).

FISH

The results of quantitative PCR were consistent with FISH images that revealed evident histological

damage due to radiation and reduction in the mean size of the organs (Additional file 4: Figure S4),

coupled with a markedly reduced *Wolbachia*-specific fluorescence intensity (Fig. 6; Additional files 5,

6, 7: Figures S5, S6, S7). Analysis of the *Wolbachia*-infected individuals revealed that fluorescent

signals differed markedly between treatments (Fig. 6). In most of the ovaries dissected from S_{ANG} and

ARwP females irradiated at 45 Gy, a weak fluorescent signal was observed (Fig. 6B, 6D; Additional

files 6, 7: Figures S6, S7) compared to that in untreated controls (Fig. 6A, 6C; Additional files 8, 9:

Figures S8, S9). In certain cases, the fluorescence was only localized in the ovarioles that possibly had

not been degenerated by the irradiation (Fig. 6D, red circle). In contrast, the organs maintained their

typical ovariole structure in untreated samples and the signal was abundant, specific, and confined

within each oocyte (Fig. 6; Additional files 8, 9: Figures S8, S9).

Discussion

Multiple blood-feeding is known to naturally occur in *Ae. albopictus* as not all of the ovarian follicles commence the gonotrophic cycle with a single blood meal [49]. Studies conducted on engorged *Ae. aegypti* females demonstrated that they were generally inhibited from host-seeking by the occurrence of distension-induced and oocyte-induced mechanisms of regulation [24, 50]; however, this inhibition does not apply to all individuals [49]. Results presented here demonstrate that the radiation doses generally used in the framework of SIT programs significantly attenuate this inhibition in both *Ae. albopictus* and *Ae. aegypti*. In fact, most females irradiated at 35 and 45 Gy fed several times a week, even if the highest number of bites was generally observed on the first day of the blood meal. Additionally, the dose of 28 Gy was found to be sufficient to induce a doubled biting rate (compared to the untreated controls). Tissue alterations in ovaries in response to irradiation might explain this phenomenon as the host-seeking behavior is interrupted through the release of hormones produced by the ovaries in the hemocoel during oogenesis [51, 52]. In virus-endemic areas, multiple blood feeding increases the likelihood of a female mosquito being infected by a suitable arbovirus and this possibility may significantly increase the chances of virus transmission to other hosts. Furthermore, the comparison of the survival curves of irradiated and untreated females revealed that the tested irradiation doses were not adequate to induce a marked decrease in life expectancy within the first 20 days in *Ae. albopictus*, which is an age that only a minority of adults are expected to reach in nature [53]. This suggests that irradiated *Ae. albopictus* females escaping the sexing procedure may have a stronger vectorial capacity than wild-type females.

The vectorial capacity of a species describes its potential to transmit a pathogen and is dependent on the ratio of mosquitoes to humans, the extrinsic incubation period of the parasite, the mosquitos' rate of biting humans, and the survival of mosquito females [54-56]. SIT and SIT/IIT programs have the specific goal of reducing the vectorial capacity (together with the mosquito nuisance) via the suppression of a target population. In this context, the co-release of relatively small percentages of irradiated females with enhanced biting activity could be viewed as a negligible and transient side effect considering that the benefits are supposed to outweigh the potential risks. However, large scale releases of sterile males do not always adapt to the ever-decreasing wild population [19, 35] and if releases rely on high overflooding release ratios and a not perfect sexing, the increasing frequency of co-released irradiated females might represent an issue to be carefully evaluated in areas endemic for arboviral diseases.

Host-seeking is another behavioral trait that determines the biting rate and that could be
affected by the radiation treatment. Previously, flight ability in radiation-sterilized males was not
reported to be affected when compared to that in untreated males [9]. Our experiments testing the
host-seeking ability in large enclosures confirm that flying ability is not affected by radiation doses
that are capable of fully suppressing fecundity in females. Indeed, starved females irradiated at 45
Gy did not exhibit a significantly different ability to reach and bite the host compared to the
controls, but when engorged, their behavior was anomalous and they continued to seek hosts, unlike
the untreated females, leading to a marked increase in the mean number of bites per individual.

Overall, dose response curves related to fecundity and fertility were in agreement with the
results of other studies [9] and verified the success of the radiation treatment. However, differently
from previous reports [21], a dose of 28 Gy was inadequate to induce full sterilization in eggs. This
result may indicate that this dose approaches the minimum threshold necessary to achieve full
sterilization of *Ae. albopictus* females and that slight modifications of the experimental set up are
sufficient to allow the viability of a small percentage of fertile eggs [19]. This issue also becomes
evident when moving from laboratory conditions to large-scale operational conditions, and
highlights the need for studies specific to the latter prior to conducting open field trials [22]. This is
particularly necessary in relation to releases involving mosquito strains with altered *Wolbachia*
infection types that have the ability to spread in the wild population through a Uni-CI pattern [15].
In these cases, a few partially fertile females could initiate a local population replacement with
unpredictable consequences [20]. In this context, for safety purposes, adding a perfect sexing
protocol to Uni-CI based IIT programs [17, 18, 35] could be preferable to combining IIT with SIT
[19, 21, 22]. In fact, in order to avoid the release of fertile and potentially invasive females in a
combined IIT/SIT approach, the required radiation doses may be so high as to seriously
compromise male mating competitiveness [19] and may induce an enhanced biting behavior in the
co-released females as evidenced in this study.

The evaluation of the vectorial capacity of a mosquito population also builds on the
measurement of the vector competence which is mainly determined by genetic factors [57]. Due to
the ability exhibited by *Wolbachia* of modulating these factors in infected mosquitoes, the decrease
in density of these bacteria caused by the radiation treatment may introduce further concerns
regarding the safe application of SIT in insect species infected by this bacterium when perfect
sexing methods are not employed.

Wolbachia-mediated reduction of vector competence in mosquitoes has been suggested to
depend on the bacterial titer and the specific strain of the bacterium [30, 58]. The density of
Wolbachia in naturally infected *Ae. albopictus* is an individual feature that may vary substantially

within a population [44] and that may be affected by the environmental conditions (temperature and food availability) under which the larvae develop [59-61]. Although distribution in somatic tissues has been observed in various cases, *Wolbachia* is mainly present within the germ line [62], and previous studies have already shown that the ovarian tissues of irradiated females are severely affected by the treatment [9, 21]. Bacteria are generally less sensitive to radiation than eukaryotes [63]; nevertheless, the density of *Wolbachia* is markedly affected by radiation doses employed for other insect species to achieve sterilization [28]. The results presented in this study highlighted an overall decrease in the density of *Wolbachia* in irradiated females, and this decrease was greatest when the pupae were irradiated at younger ages, and was positively correlated with female aging. This downward trend was common in all the studied *Wolbachia* strains (compared to the untreated controls).

Early irradiation is generally associated with a stronger efficiency of the sterilization treatment [43]; however, negative effects on fitness have been found to increase in younger treated pupae. For this reason, SIT programs are generally targeted at an intermediate pupal age to achieve full sterilization and to sufficiently preserve the fitness of adult males [9]. The damage experienced by the ovaries is consistent with complete sterilization that occurs when sufficiently high doses are employed. Our results highlighted that the destruction of these tissues—which are relatively sensitive to radiation due to intense mitotic activity [64]—seems to also affect the survival and/or the reproduction of the hosted *Wolbachia*. The endosymbiont may be involved—together with the host cells—in the mechanisms of apoptosis usually characterizing the oocytes subjected to irradiation [65]. However, further studies should be conducted to investigate this phenomenon and to determine why the reduction in the density of the bacterium becomes evident by qPCR only over a week after the treatment. This occurrence may be partly explained by a gradual clearance of the bacterial DNA corresponding to dead individuals or present in extracellular form [66] in the damaged ovarian tissues of irradiated females. Selection of an effective method to normalize the qPCR data allowed us to study an otherwise difficult-to-study phenomenon, as irradiation has previously been demonstrated to compromise the suitability of various common housekeeping genes when tissues are damaged [67]. A comparison of the actin gene copy number in the ovaries of irradiated and control individuals confirms this (Additional file2: Figure S2), and suggests the need for specific studies to identify the best candidate housekeeping genes when performing similar experiments. Investigating the levels of expression of specific *Wolbachia* genes, or host genes under *Wolbachia* control, may provide useful information to measure the activity of the bacteria, and as a consequence, to estimate their response to the irradiation treatment [68].

FISH supported the qPCR data related to ovaries, as demonstrated by the markedly reduced *Wolbachia*-specific fluorescence intensity associated with these tissues in irradiated females aged 13±1 days. This evidence reinforces the idea that the population of this endosymbiotic bacterium is severely affected by a 45-Gray radiation treatment, at least in this organ, and that qPCR is only partially capable of highlighting this phenomenon.

Overall, our results are insufficient to determine whether radiation treatment can be associated with a loss of specific *Wolbachia*-controlled biological activities such as pathogen interference. *Wolbachia* endosymbionts mainly exert their limiting action on arbovirus dissemination in the bodies of *Aedes* species by acting in the midgut [69], and similar to ovarian cells, midgut cells are characterized by intense mitotic activity and their structure and physiology have been previously shown to be affected by radiation [64, 70]. Therefore, *Wolbachia* and the related host-symbiont mechanisms of regulation may be affected by radiation in these tissues as well. However, this study does not provide evidence of such phenomena. Furthermore, as reported above, the detection of *Wolbachia* DNA does not necessarily indicate that the bacterium is alive or capable of fully exerting its effects on host physiology, because nucleic-acid-based analytical methods provide only limited information regarding the activities and physiological state of microorganisms in samples. These aspects can be detected retrospectively, but only after sufficient time has elapsed for the degradation and removal of DNA associated with inactivated cells [66, 71].

qPCR analysis specifically targeting the *Wolbachia* density in the gut of irradiated females may provide useful information to investigate bacterial population dynamics in these tissues. However, specific vector-competence studies will be necessary to ascertain whether a sterilizing radiation treatment leads to increased risk of virus acquisition or transmission per single bite. Coupled with the enhanced biting activity shown in the present study, an increased vector competence would boost further the vectorial capacity of female mosquitos. For such a study to be viable, oral infection trials should be conducted on mass-reared mosquito females after applying a radiation treatment at the doses necessary for large-scale operational programs, and all *Wolbachia*-infected vector species should be included [22].

In the case of programs based on the combination of SIT and IIT, investigation of the effects of irradiation on the induced level of CI in treated males should also be performed [31], because based on the results of the ovaries, a decrease in the titer of *Wolbachia* in the testes following the irradiation is a reasonably possible scenario.

Conclusions

The results presented in this work stress the need for more thorough scientific investigations on the

radiation biology of female mosquitoes. Such studies should be particularly opportune prior to

conducting area-wide SIT or SIT/IIT programs that rely on imperfect sexing in areas endemic for

arboviral diseases. Indeed, even a low percentage of contaminating mosquito females may result in

the release of thousands of boosted vectors under large scale settings. Even if this safety issue

should be negligible in the context of a successful population suppression program, conservatively,

the effect of increasing frequencies of irradiated females in the target area should be thoroughly

evaluated with the support of opportune models analyzing the epidemiological risks [55, 72, 73].

Measuring the vector competence of irradiated females that are infected with *Wolbachia*

should be also opportune because the irradiation-induced decrease in the density of this bacterium

may be consistent with effects on biological phenomena such as pathogen interference. Applying

advanced and more efficient systems of sex separation capable of preventing the escape of females

during release of sterile males [17, 18, 35, 74] would be sufficient to mitigate risks. Additionally,

release protocols could use constant monitoring of the wild-type population to limit the number of

mosquitoes released to a minimum threshold to guarantee efficacy [75]. Certainly, in the case of

release programs that involve *Wolbachia* infections with pathogen interference phenotypes, the

irradiation of pupae at young ages should be avoided because this treatment would maximize the

biting activity and the *Wolbachia* depletion in adult females.

The data presented here may furnish useful cues to enhance the safety level of SIT-based

control programs against *Aedes* mosquitoes and encourage a careful comparison between the

various genetic control methods in search of the most efficient, sustainable, and safe strategy for

mosquito vector control.

List of Abbreviations

SIT: Sterile Insect Technique

CI: Cytoplasmic Incompatibility

Uni-CI: Unidirectional Cytoplasmic Incompatibility

IIT: Incompatible Insect Technique

qPCR: quantitative polymerase chain reaction

FISH: Fluorescence in situ hybridization

AR*w*P: *Aedes albopictus* population obtained through the replacement of the native *w*AlbA-*w*AlbB

Wolbachia infection with *w*Pip *Wolbachia*

S$_{ANG}$: Wild-type *Ae. albopictus* from Anguillara Sabazia (Rome)

LEU: large cellular polycarbonate experimental units

ANOVA: Analysis of viariance

Declarations

Ethics approval and consent to participate

Blood meals were provided via anesthetized mice in agreement with the Bioethics Committee for

Animal Experimentation in Biomedical Research and in accordance with procedures approved by

the ENEA Bioethical Committee according to the EU directive 2010/63/EU. The mice belonged to

a colony housed at CR ENEA Casaccia and maintained for experimentation based on the

authorization N. 80/2017-PR released (on February 2, 2017) by the Italian Ministry of Health.

Feeding female mosquitoes on the blood of human hosts (i.e., the Authors RM, EL, GL, and MC)

during the experiments was also approved by the ENEA Bioethical Committee.

Consent for publication

Two of the Authors (EL and MC) are present in Figure S1 and consent the publication of the image.

Availability of data and materials

All relevant data are within the paper and its Supporting Information files. The datasets used and/or

analysed during the current study are available from the corresponding author on reasonable request

Competing interests

The Authors declare that they have no competing interests

Funding

The Authors received no specific funding for this work.

Authors' contributions

RM and MC conceptualized the research. The methodology was planned by RM, EL, MC, AD, CD,

and CP. RM, EL, GL, CD, CP, AD, and MC conducted the investigation. RM, EL, GF and MC

organized and managed the datasets. RM, GF and MC conducted the statistical analysis of the data.

MC and AS took care of funding acquisition. MC and CD supervised the experimentation. RM

wrote the original draft of the manuscript. RM, MC, CD, EL, AS, GF, and GL contributed to

writing, reviewing, and editing the manuscript.

Acknowledgments

We thank Marta Piscitelli (Division for Health Protection Technologies, ENEA-Casaccia Research

Center, Rome, Italy) as responsible of the facility for the housing and care of the mice used for

blood feeding. We also thank Alessia Cappelli (School of Biosciences and Medical Veterinary,

University of Camerino, MC, Italy) for her assistance with the FISH analysis and Alessia Fiore

(Biotechnology and Agroindustry Division, ENEA-Casaccia Research Center, Rome, Italy) for her

contribution to the normalization of the qPCR data through the measurement of the DNA content in

the extracts from the ovarian tissues.

Figure Captions

Figure 1. Kaplan-Meier survival curves comparing, respectively, irradiated and untreated S_{ANG} (A)

and ARwP (B) *Ae. albopictus* over two weeks since the emergence. Differences between treatments

were not statistically significant (Log rank test analysis: $P < 0.05$).

Figure 2. Altered biting behavior in irradiated *Ae. albopictus* and *Ae. aegypti*. The experiment was

carried out in laboratory cages by offering daily a blood meal over two sets of 5 subsequent days

interrupted by 2 days of rest and starting with 5 days old females. **A:** Daily mean percentage of

biting S_{ANG} and ARwP *Ae. albopictus*; **B:** Mean weekly number of bites per S_{ANG} and ARwP *Ae.

albopictus female over two weeks; **C:** Daily mean percentage of biting *Ae. aegypti*; **D:** Mean

weekly number of bites per *Ae. aegypti* female over two weeks; A and C: within each mosquito

population, different letters indicate statistically significant differences between treatments

(Repeated measures ANOVA followed by Tukey's HSD test: $P < 0.05$); B and D: within each

mosquito population, different letters indicate statistically significant differences between

treatments (One-Way ANOVA followed by Tukey's test: $P < 0.05$).

Figure 3. Altered host seeking and biting behavior in irradiated S_{ANG} and ARwP *Ae. albopictus*

under large enclosures setting. The experiment was carried out in Large Experimental Units

(8.5×5×5 m) under open field climatic conditions and involved females irradiated at 45-Gy

compared to untreated counterparts. Average landing time and biting proportions were compared

between treatments within a 15-minute interval. **A:** Comparison between irradiated and untreated

starved females (6±1 days old); **B:** Comparison between irradiated and untreated females 48 h after

the engorgement (8±1 days old). Two-way ANOVA demonstrated that the difference between

treatments was statistically significant in the case of the engorged females (P < 0.05).

Figure 4. Decreased *Wolbachia* titer in irradiated S_{ANG} and ARwP *Ae. albopictus.* **A:** A 45-Gy dose

was applied at three different pupal ages (26, 36, and 46±2 hours); quantitative PCR analysis was

then performed by using primers targeting the specific *Wolbachia* strains characterizing the two *Ae.*

albopictus populations and analyzing 6±1 days old females. **B**: A 45-Gy dose was applied to pupae

aged 36±4 hours; qPCR analysis was then carried out studying two female ages (6 and 13±1 days).

Data were normalized by using the actin gene as host reference. Within each infection type and

female age, different letters indicate statistically significant differences between treatments (One-

way ANOVA followed by Tukey's test: P<0.05).

Figure 5. Decreased *Wolbachia* titer in the ovaries of irradiated S_{ANG} and ARwP *Ae. albopictus.* A

45-Gy dose was applied to 36±4 hours old pupae. **A:** qPCR analysis was performed on the dissected

ovaries of 6 and 13±1 days old females by using primers targeting the specific *Wolbachia* strains

characterizing the two *Ae. albopictus* populations. Data were normalized by using DNA (2 ul of

purified DNA per reaction). **B**: qPCR analysis was performed on bodies lacking ovaries of 6 and

13±1 days old females by using primers targeting the specific *Wolbachia* strains characterizing the

two *Ae. albopictus* populations. Data were normalized by using the actin gene as host reference.

Within each infection type and female age, different letters indicate statistically significant

differences between treatments (One-way ANOVA followed by Tukey's test: P<0.05).

Figure 6. Fluorescence In Situ Hybridization of *Ae. albopictus* ovaries in irradiated females. A 45-

Gy dose was applied to 36±4 hours old pupae of S_{ANG} and ARwP *Ae. albopictus*. Ovaries from

sample individuals aged 13±1 days were then subjected to FISH analysis and compared with

untreated counterparts. The distribution of *Wolbachia* is evidenced in green while blue stain is

DAPI. **A**: Ovaries of untreated S_{ANG} females; **B**: Ovaries of irradiated S_{ANG} females **C**: Ovaries of

untreated ARwP females; **D**: Ovaries of irradiated ARwP females.

Supplementary Information

Additional file 1: Figure S1. Schematic of host seeking behavior trials conducted under large enclosures

Additional file 2: Figure S2. Actin gene copies in the ovaries of *Ae. albopictus* females irradiated at 45 Gy in comparison with untreated counterparts

Additional file 3: Figure S3. Host seeking and biting behavior of irradiated S_{ANG} and AR*w*P *Ae. albopictus* under large enclosures compared to untreated controls. Biting proportions and average times to landing were compared between treatments within a 15-minute interval. **A:** Comparison between untreated and irradiated starved females aged 13±1 days; **B:** Comparison between irradiated and untreated females engorged females 48 hours after the engorgement (i.e., 15±1 days old). Two-way ANOVA demonstrated that the difference between treatments was statistically significant in the case of the engorged females ($P < 0.05$).

Additional file 4: Figure S4. Structural damage induced by irradiation at 45 Gy in the ovaries of 13±1 days old *Ae. albopictus* females in bright field**. A**: Ovaries of untreated S_{ANG} females; **B**: Ovaries of irradiated S_{ANG} females **C**: Ovaries of untreated AR*w*P females; **D**: Ovaries of irradiated AR*w*P females.

Additional file 5: Figure S5. FISH analysis of the ovaries of 13±1 days old *Ae. albopictus* belonging to a line (AR) cured from *Wolbachia* infection. **A** and **D**: DAPI stained; **B** and **E**: FITC stained; **C** and **F**: bright field. No specific green-fluorescent signal was detected in the aposymbiotic line.

Additional file 6: Figure S6. Additional images related to the FISH analysis of the ovaries of 13±1 days old S_{ANG} *Ae. albopictus* irradiated at 45 Gy. The distribution of *Wolbachia* is evidenced in green while blue stain is DAPI. **A** and **D**: DAPI stained; **B** and **E**: FITC stained; **C** and **F**: bright field. The green-fluorescent signal related to *Wolbachia* is weak and not homogeneously distributed.

Additional file 7: Figure S7. Additional images related to the FISH analysis of the ovaries of 13±1

days old ARwP *Ae. albopictus* irradiated at 45 Gy. The distribution of *Wolbachia* is evidenced in

green while blue stain is DAPI. **A** and **D**: DAPI stained; **B** and **E**: FITC stained; **C** and **F**: bright

field. The green-fluorescent signal related to *Wolbachia* is weak and not homogeneously

distributed.

Additional file 8: Figure S8. Additional images related to the FISH analysis of the ovaries of 13±1

days old untreated ARwP *Ae. albopictus*. Blue stain is DAPI. **A** and **D**: DAPI stained; **B** and **E**:

FITC stained; **C** and **F**: bright field. The green-fluorescent signal related to *Wolbachia* is strong and

regularly distributed.

Additional file 9: Figure S9. Additional images related to the FISH analysis of the ovaries of 13±1

days old untreated S$_{ANG}$ *Ae. albopictus*. Blue stain is DAPI. **A** and **D**: DAPI stained; **B** and **E**: FITC

stained; **C** and **F**: bright field. The green-fluorescent signal related to *Wolbachia* is strong and

regularly distributed.

References

Article within a journal (no page numbers)

Rohrmann S, Overvad K, Bueno-de-Mesquita HB, Jakobsen MU, Egeberg R, Tjønneland A, et al. Meat consumption and
mortality - results from the European Prospective Investigation into Cancer and Nutrition. BMC Medicine. 2013;11:63.

Article within a journal by DOI

Slifka MK, Whitton JL. Clinical implications of dysregulated cytokine production. Dig J Mol Med. 2000;
doi:10.1007/s801090000086.

Article within a journal supplement

Frumin AM, Nussbaum J, Esposito M. Functional asplenia: demonstration of splenic activity by bone marrow scan. Blood
1979;59 Suppl 1:26-32.

Book chapter, or an article within a book

Wyllie AH, Kerr JFR, Currie AR. Cell death: the significance of apoptosis. In: Bourne GH, Danielli JF, Jeon KW, editors.
International review of cytology. London: Academic; 1980. p. 251-306.

OnlineFirst chapter in a series (without a volume designation but with a DOI)

Saito Y, Hyuga H. Rate equation approaches to amplification of enantiomeric excess and chiral symmetry breaking. Top
Curr Chem. 2007. doi:10.1007/128_2006_108.

Complete book, authored

Blenkinsopp A, Paxton P. Symptoms in the pharmacy: a guide to the management of common illness. 3rd ed. Oxford:
Blackwell Science; 1998.

1. Iwamura T, Guzman-Holst A, Murray KA. Accelerating invasion potential of disease
vector *Aedes aegypti* under climate change. Nat Commun. 2020;11(1):2130.

2. Moyes CL, Vontas J, Martins AJ, Ng LC, Koou SY, Dusfour I, et al. Contemporary status
of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans. PLoS
Negl Trop Dis. 2017;11(7):e0005625.

3. Garcia GA, Hoffmann AA, Maciel-de Freitas R, Villela DA. *Aedes aegypti* insecticide
resistance underlies the success (and failure) of *Wolbachia* population replacement. Sci Rep.
2020;10:63.

4. Alphey L, Benedict M, Bellini R, Clark GG, Dame DA, Service MW, et al. Sterile-insect
methods for control of mosquito-borne diseases: an analysis. Vector Borne Zoonotic Dis.
2010;10(3):295-311.

5. Hammond A, Galizi R, Kyrou K, Simoni A, Siniscalchi C, Katsanos D, et al. A CRISPR-Cas9
gene drive system targeting female reproduction in the malaria mosquito vector *Anopheles*
gambiae. Nat Biotechnol. 2016;34(1):78-83.

6. Thomas DD, Donnelly CA, Wood RJ, Alphey LS. Insect population control using a dominant,
repressible, lethal genetic system. Science. 2000;287(5462):2474-2476.

7. Massonnet-Bruneel B, Corre-Catelin N, Lacroix R, Lees RS, Hoang KP, Nimmo D, et al. Fitness
of transgenic mosquito *Aedes aegypti* males carrying a dominant lethal genetic system. PLoS One.
2013;8(5):e62711.

8. Lees RS, Gilles JR, Hendrichs J, Vreysen MJ, Bourtzis K. Back to the future: the sterile insect

technique against mosquito disease vectors. Curr Opin Insect Sci. 2015;10:156-162.

9. Bond JG, Osorio AR, Avila N, Gomez-Simuta Y, Marina CF, Fernandez-Salas I, et al.

Optimization of irradiation dose to *Aedes aegypti* and *Ae. albopictus* in a sterile insect technique

program. PloS One. 2019;14(2):e0212520.

10. Werren JH, Baldo L, Clark ME. *Wolbachia*: master manipulators of invertebrate biology. Nat

Rev Microbiol. 2008;6(10):741-751.

11. Sinkins S, O'Neill S. *Wolbachia* as a vehicle to modify insect populations. In: Handler AM

JA, editor. Insect Transgenesis. Boca Raton: CRC Press; 2000. p. 271-287.

12. Dobson SL, Fox CW, Jiggins FM. The effect of *Wolbachia*-induced cytoplasmic

incompatibility on host population size in natural and manipulated systems. Proc Royal Soc B.

2002;269(1490):437-445.

13. Bourtzis K, Dobson SL, Xi Z, Rasgon JL, Calvitti M, Moreira LA, et al. Harnessing mosquito-

Wolbachia symbiosis for vector and disease control. Acta Trop. 2014;132:S150-S163.

14. Mains J, Brelsfoard C, Rose R, Dobson S. Female adult *Aedes albopictus* suppression by

Wolbachia-infected male mosquitoes. Sci. Rep. UK. 2016;6:33846.

15. Moretti R, Marzo GA, Lampazzi E, Calvitti M. Cytoplasmic incompatibility management to

support Incompatible Insect Technique against *Aedes albopictus*. Parasites Vectors. 2018;11(2):649.

16. Caputo B, Moretti R, Manica M, Serini P, Lampazzi E, Bonanni M, et al. A bacterium against

the tiger: preliminary evidence of fertility reduction after release of *Aedes albopictus* males with

manipulated *Wolbachia* infection in an Italian urban area. Pest Manag. 2020;76(4):1324-1332.

17. Mains JW, Kelly PH, Dobson KL, Petrie WD, Dobson SL. Localized control of *Aedes aegypti*

(Diptera: Culicidae) in Miami, FL, via inundative releases of *Wolbachia*-infected male mosquitoes.

J Med Entomol. 2019;56(5):1296-1303.

18. Beebe NW, Pagendam D, Trewin, BJ, Boomer A, Bradford M, et al. Releasing incompatible

males drives strong suppression across populations of wild and *Wolbachia*-carrying *Aedes*

aegypti in Australia. PNAS. 2021;118(41):e2106828118.

19. Zheng X, Zhang D, Li Y, Yang C, Wu Y, Liang X, et al. Incompatible and sterile insect

techniques combined eliminate mosquitoes. Nature. 2019;572(7767):56-61.

20. Yen PS, Failloux AB. A review: *Wolbachia*-based population replacement for mosquito

control shares common points with genetically modified control approaches. Pathogens.

2020;9(5):404.

21. Zhang D, Lees RS, Xi Z, Gilles JR, Bourtzis K. Combining the sterile insect technique with

Wolbachia-based approaches: II-a safer approach to *Aedes albopictus* population suppression

programmes, designed to minimize the consequences of inadvertent female release. PloS One.

2015;10(8):e0135194.

22. Moretti R, Calvitti M. Issues with combining incompatible and sterile insect techniques. Nature.

2021;590:E1–E2.

23. Li Y, Baton LA, Zhang D, Bouyer J Parker AG, Hoffmann AA, et al. Reply to: Issues with

combining incompatible and sterile insect techniques. Nature. 2021;590;E3–E5.

24. Helinski ME, Parker AG, Knols BG. Radiation biology of mosquitoes. Malar J.

2009;8(S2):S6.

25. Klowden MJ, Lea AO. Humoral inhibition of host-seeking in *Aedes aegypti* during oocyte

maturation. J Insect Physiol. 1979;25(3):231–235.

26. Beach R. Mosquitoes: biting behavior is inhibited by ecdysone. Science. 1979;205:829-831.

27. Kramer LD, Ciota AT. Dissecting vectorial capacity for mosquito-borne viruses. Curr Opin

Virol. 2015;15:112-118.

28. Demirbas-Uzel G, De Vooght L, Parker AG, Vreysen MJ, Mach RL, Van Den Abbeele J, et

al. Combining paratransgenesis with SIT: impact of ionizing radiation on the DNA copy number

of *Sodalis glossinidius* in tsetse flies. BMC Microbiol. 2018;18(1):160.

29. Balestrino F, Puggioli A, Bellini R, Petric D, Gilles J. Mass production cage for *Aedes*

albopictus (Diptera: Culicidae). J Med Entomol. 2014;51(1):155-163.

30. Lu P, Bian G, Pan X, Xi Z. *Wolbachia* induces density-dependent inhibition to dengue virus in

mosquito cells. PLoS Negl Trop Dis. 2012;6(7):e1754.

31. Ross PA, Ritchie SA, Axford JK, Hoffmann AA. Loss of cytoplasmic incompatibility in

Wolbachia-infected *Aedes aegypti* under field conditions. PLoS Negl Trop Dis.

2019;13(4):e0007357

32. Mayer SV, Tesh RB, Vasilakis N. The emergence of arthropod-borne viral diseases: A

global prospective on dengue, chikungunya and Zika fevers. Acta Trop. 2017;166:155-163.

33. Wilder-Smith A, Gubler DJ, Weaver SC, Monath TP, Heymann DL, Scott TW. Epidemic

arboviral diseases: priorities for research and public health. Lancet Infect Dis. 2017;17(3):101-106.

34. Baldacchino F, Caputo B, Chandre F, Drago A, della Torre A, Montarsi F, et al. Control

methods against invasive *Aedes* mosquitoes in Europe: a review. Pest Manag Sci.

2015;71(11):1471-1485.

35. Crawford JE, Clarke DW, Criswell V, Desnoyer M, Cornel D, Deegan B, et al. Efficient

production of male *Wolbachia*-infected *Aedes aegypti* mosquitoes enables large-scale suppression

of wild populations. Nat Biotechnol. 2020;38(4):482-492.

36. Mousson L, Zouache K, Arias-Goeta C, Raquin V, Mavingui P, Failloux A-B. The native

Wolbachia symbionts limit transmission of dengue virus in *Aedes albopictus*. PLoS Negl Trop

Dis. 2012;6(12):e1989.

37. Calvitti M, Moretti R, Lampazzi E, Bellini R, Dobson SL. Characterization of a new *Aedes*

albopictus (Diptera: Culicidae)-*Wolbachia pipientis* (Rickettsiales: Rickettsiaceae) symbiotic

association generated by artificial transfer of the *w*Pip strain from *Culex pipiens* (Diptera:

Culicidae). J Med Entomol. 2014;47(2):179-187.

38. Calvitti M, Moretti R, Skidmore AR, Dobson SL. *Wolbachia* strain *w*Pip yields a pattern of

cytoplasmic incompatibility enhancing a *Wolbachia*-based suppression strategy against the

disease vector *Aedes albopictus*. Parasites Vectors. 2012;5:254.

39. Moretti R, Calvitti M. Male mating performance and cytoplasmic incompatibility in a wPip

Wolbachia trans-infected line of *Aedes albopictus* (*Stegomyia albopicta*). Med Vet Entomol.

2013;27(4):377-386.

40. Bellini R, Calvitti M, Medici A, Carrieri M, Celli G, Maini S. Use of the sterile insect

technique against *Aedes albopictus* in Italy: first results of a pilot trial. In: Vreysen MJB,

Hendrichs J, Robinson AS (eds.) Area-wide control of insect pests. Dordrecht, the Netherlands:

Springer; 2007. pp. 505-515.

41. Lau M-J, Endersby-Harshman NM, Axford JK, Ritchie SA, Hoffmann AA, Ross PA.

Measuring the host-seeking ability of A*edes aegypti* destined for field release. Am J Trop Med

and Hyg. 2020;102(1):223–231.

42. Rainey SM, Shah P, Kohl A, Dietrich I. Understanding the *Wolbachia*-mediated inhibition of

arboviruses in mosquitoes: progress and challenges. J Gen Virol. 2014;95(3):517-530.

43. Yamada H, Maiga H, Juarez J, Carvalho DDO, Mamai W, Ali A, et al. Identification of

critical factors that significantly affect the dose-response in mosquitoes irradiated as pupae.

Parasites Vectors. 2019;12(1):435.

44. Calvitti M, Marini F, Desiderio A, Puggioli A, Moretti R. *Wolbachia* density and cytoplasmic
incompatibility in *Aedes albopictus:* concerns with using artificial *Wolbachia* infection as a vector
suppression tool. PloS One. 2015;10(3):e0121813.

45. Zhou W, Rousset F, O'Neill S. Phylogeny and PCR-based classification of *Wolbachia* strains
using wsp gene sequences. Proc Royal Soc B. 1998;265(1395):509-515.

46. Moretti R, Yen P-S, Houé V, Lampazzi E, Desiderio A. Failloux A-B. Calvitti M. Combining
Wolbachia-induced sterility and virus protection to fight *Aedes albopictus*-borne viruses. PLoS
Negl Trop Dis. 2018;12:e0006626.

47. Braquart-Varnier C, Altinli M, Pigeault R, Chevalier FD, Greve P, Bouchon D, et al. The
mutualistic side of *Wolbachia*-isopod interactions: *Wolbachia* mediated protection against
pathogenic intracellular bacteria. Front Microbiol. 2015;6:1388.

48. Mancini MV, Herd CS, Ant TH, Murdochy SM, Sinkins SP. *Wolbachia* strain *w*Au efficiently
blocks arbovirus transmission in *Aedes albopictus*. PLoS Negl Trop Dis. 2020;14(3):e0007926.

49. Farjana T, Tuno N. Multiple blood feeding and host-seeking behavior in *Aedes aegypti* and
Aedes albopictus (Diptera: Culicidae). J Med Entomol. 2013;50(4):838-846.

50. Klowden MJ, Lea AO. Abdominal distention terminates subsequent host-seeking behaviour of
Aedes aegypti following a blood meal. J Insect Physiol. 1979;25(7):583-585.

51. Klowden MJ. Endocrine aspects of mosquito reproduction. Arch Insect Biochem Phys:
Published in Collaboration with the Entomological Society of America. 1997;35(4):491-512.

52. Klowden MJ. Initiation and termination of host-seeking inhibition in *Aedes aegypti* during
oocyte maturation. J Insect Physiol. 1981;27(11):799-803.

53. Brady OJ, Johansson MA, Guerra CA, Bhatt S, Golding N, Pigott DM, et al. Modelling adult
Aedes aegypti and *Aedes albopictus* survival at different temperatures in laboratory and field
settings. Parasites Vectors. 2013;6:351.

54. Brady OJ, Godfray HC, Tatem AJ, Gething PW, Cohen JM, McKenzie FE, et al. Vectorial capacity and vector control: reconsidering sensitivity to parameters for malaria elimination. Trans R Soc Trop Med Hyg. 2016;110(2):107-17.

55. Catano-Lopez A, Rojas-Diaz D, Laniado H, Arboleda-Sánchez S, Puerta-Yepes MA, Lizarralde-Bejarano DP. An alternative model to explain the vectorial capacity using as example *Aedes aegypti* case in dengue transmission. Heliyon. 2019;5(10):e02577.

56. Wu SL, Sánchez CHM, Henry JM, Citron DT, Zhang Q, Compton K, et al. Vector bionomics and vectorial capacity as emergent properties of mosquito behaviors and ecology. PLoS Comput Biol. 2020;16(4):e1007446.

57. Beerntsen BT, James AA, Christensen BM. Genetics of mosquito vector competence. Microbiol Mol Biol Rev. 2000;64(1):115-37.

58. Johnson KN. The impact of *Wolbachia* on virus infection in mosquitoes. Viruses. 2015;7(11):5705-5717.

59. Ross PA, Wiwatanaratanabutr I, Axford JK, White VL, Endersby-Harshman NM, Hoffmann AA. *Wolbachia* infections in *Aedes aegypti* differ markedly in their response to cyclical heat stress. PLoS Pathog. 2017;13(1):e1006006.

60. Wiwatanaratanabutr I, Kittayapong P. Effects of crowding and temperature on *Wolbachia* infection density among life cycle stages of *Aedes albopictus*. J Invertebr Pathol. 2009;102(3):220-224.

61. Foo IJH, Hoffmann AA, Ross PA. Cross-generational effects of heat stress on fitness and *Wolbachia* density in *Aedes aegypti* mosquitoes. Trop Med Infect Dis. 2019;4(1):13.

62. Dobson SL, Bourtzis K, Braig HR, Jones BF, Zhou W, Rousset F, et al. *Wolbachia* infections are distributed throughout insect somatic and germ line tissues. Insect Biochem Mol Biol. 1999;29(2):153-160.

1019

1020 63. Confalonieri F, Sommer S. Bacterial and archaeal resistance to ionizing radiation. J Phys,

1021 Conf. Ser. 2011;261:012005.

1022

1023 64. Bakri A, Mehta K, Lance DR. Sterilizing insects with ionizing radiation, In: V.A. Dyck, J.

1024 Hendrichs and A.S. Robinson (eds.), Sterile insect technique. Principles and practice in area-wide

1025 integrated pest management. Dordrecht, the Netherlands: Springer; 2005. pp. 233-268.

1026

1027 65. Shim HJ, Lee EM, Nguyen LD, Shim J, Song YH. High-dose irradiation induces cell cycle

1028 arrest, apoptosis, and developmental defects during *Drosophila* oogenesis. PLoS One.

1029 2014;9(2):e89009.

1030

1031 66. Rogers G, Marsh P, Stressmann A, Allen C, Daniels T, Carroll M, et al. The exclusion of dead

1032 bacterial cells is essential for accurate molecular analysis of clinical samples. Clin Microbiol

1033 Infect. 2010;16(11):1656-1658.

1034

1035 67. Iyer G, Wang AR, Brennan SR, Bourgeois S, Armstrong E, Shah P, et al. Identification of

1036 stable housekeeping genes in response to ionizing radiation in cancer research. Sci Rep.

1037 2017;7(1):43763.

1038

1039 68. Lindsey ARI, Bhattacharya T, Hardy RW, Newton ILG. *Wolbachia* and virus alter the host

1040 transcriptome at the interface of nucleotide metabolism pathways. mBio. 2021;12(1):e03472-20.

1041

1042 69. Chouin-Carneiro T, Ant TH, Herd C, Louis F, Failloux AB, Sinkins SP. *Wolbachia* strain

1043 *w*AlbA blocks Zika virus transmission in *Aedes aegypti*. Med Vet Entomol. 2020;34(1):116-119.

1044

1045 70. Kheirallah DA, El-Samad LM. Midgut cells alteration in gamma-irradiated beetles (*Blaps*

1046 *polycresta*, Coleoptera: Tenebrionidae). Braz J Biol. 2020;80(2):465-473.

1047

1048 71. Cangelosi GA, Meschke JS. Dead or alive: molecular assessment of microbial viability. Appl

1049 Environ Microbiol. 2014;80(19):5884-5891.

1050

1051 72. Mayton EH, Tramonte AR, Wearing HJ, Christofferson RC. Age-structured vectorial capacity

1052 reveals timing, not magnitude of within-mosquito dynamics is critical for arbovirus fitness

1053 assessment. Parasites Vectors. 2020;13:310.

1054

1055 73. Velázquez-Castro J, Anzo-Hernández A, Bonilla-Capilla B, Soto-Bajo M, Fraguela-Collar A.

1056 Vector-borne disease risk indexes in spatially structured populations. PLoS Negl Trop Dis.

1057 2018;12(2):e0006234.

1058

1059 74. Koskinioti P, Augustinos AA, Carvalho DO, Misbah-Ul-Haq M, Pillwax G, de la Fuente LD,

1060 Salvador-Herranz G, Herrero RA, Bourtzis K. Genetic sexing strains for the population

1061 suppression of the mosquito vector *Aedes aegypti*. Philos Trans R Soc, B, Biol Sci.

1062 2021;376(1818):20190808.

1063

1064 75. Pagendam DE, Trewin BJ, Snoad N, Ritchie SA, Hoffmann AA, Staunton KM, Paton C,

1065 Beebe N. Modelling the *Wolbachia* incompatible insect technique: strategies for effective

1066 mosquito population elimination. BMC Biol. 2020;18:161.