macroH2A2 shapes chromatin accessibility at enhancer elements in glioblastoma to modulate a targetable self-renewal epigenetic network

Ana Nikolic[1,2,3], Anna Bobyn[1,2,4], Francesca Maule[1,2], Katrina Ellestad[1,2], Xueqing Lun[1,2], Michael Johnston[1,2], Christopher J Gafuik[1,2], Franz J Zemp[1,2], Seungil Paik[1,2], Nicoletta Ninkovic[1,2], Sajid A Marhon[5], Parinaz Mehdipour[5], Yaoqing Shen[6,7], N. Daniel Berger[1,2,3], Duncan K Brownsey[1,2,8], Peter B Dirks[9,10], Darren J Derksen[1,2,8], Steven JM Jones[6,7], Daniel de Carvalho[5,11], Donna L Senger[1,2,12], Jennifer A Chan[1,2,12], Douglas J Mahoney[1,2,13], Marco Gallo[1,2,3]*

[1]Arnie Charbonneau Cancer Institute, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada
[2]Alberta Children's Hospital Research institute, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada
[3]Department of Biochemistry and Molecular Biology, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada
[4]Department of Biological Sciences, Faculty of Science, University of Calgary, Calgary, AB, Canada
[5]Princess Margaret Cancer Centre, Toronto, ON, Canada
[6]Canada's Michael Smith Genome Sciences Centre, BC Cancer, Vancouver, BC, Canada
[7]Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada
[8]Department of Chemistry, Faculty of Science, University of Calgary, Calgary, AB, Canada
[9]Program in Developmental and Stem Cell Biology, Hospital for Sick Children, Toronto, ON, Canada
[10]Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada
[11]Department of Medical Biophysics, Faculty of Science, University of Toronto, Toronto, ON, Canada
[12]Department of Oncology, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada
[13]Department of Microbiology, Immunology and Infectious Diseases, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada

*Correspondence to: Marco Gallo (marco.gallo@ucalgary.ca)

SUMMARY

Self-renewal is a crucial property of glioblastoma cells and is enabled by the choreographed function of chromatin regulators and transcription factors. Identifying targetable epigenetic mechanisms of self-renewal could represent an important step toward developing new and effective treatments for this universally lethal cancer. Here we uncover a targetable epigenetic axis of self-renewal mediated by the histone variant macroH2A2. Using patient-derived *in vitro* and *in vivo* models, we show that macroH2A2 has a direct role in shaping chromatin accessibility at enhancer elements to antagonize transcriptional programs of self-renewal. Pharmaceutical inhibition of the chromatin remodeler Menin increased macroH2A2 levels and repressed self-renewal. Our results reveal a targetable epigenetic mechanism of self-renewal controlled by macroH2A2 and suggest new treatment approaches for glioblastoma patients.

KEYWORDS

Glioblastoma, cancer, brain tumor, epigenomics, chromatin, histone variants, macroH2A2, self-renewal, patient-derived models.

SIGNIFICANCE

Glioblastoma is an incurable brain cancer. Malignant self-renewing cells have been shown to drive tumor growth, to be refractory to current treatment approaches and to seed relapses, which ultimately prove lethal. Identifying new and targetable mechanisms associated with self-renewal could be a fundamental first step in designing effective therapies that slow or prevent glioblastoma relapses. Using patient-derived models of glioblastoma, we deployed epigenomic approaches and functional assays to define the role of the histone variant macroH2A2 in repressing self-renewal. We identified compounds that increase macroH2A2 levels and repress self-renewal, including a Menin inhibitor. As Menin inhibitors are being tested in clinical trials, these compounds could be used in new therapeutic paradigms to target self-renewing cell populations in glioblastoma.

INTRODUCTION

Glioblastoma (GBM) is the most common malignant primary brain tumor in adults and carries abysmal prognoses, even in cases treated with gross total resection and adjuvant chemoradiotherapy (Stupp et al., 2005). Several factors are thought to contribute to the aggressiveness of GBM, especially at recurrence: The activation of drug resistance mechanisms (Chen et al., 2012), hypermutation (Cahill et al., 2007; Hunter et al., 2006; Yip et al., 2009), a highly immunosuppressive tumor microenvironment (Alvarado et al., 2017; Antonios et al., 2017), and underlying disease characteristics, most notably a high degree of intratumoral heterogeneity (Neftel et al., 2019; Suvà et al., 2014). Recent studies have shown that the cellular heterogeneity observed in GBM occurs both at the genetic and epigenetic level (Gallo et al., 2015; Lan et al., 2017; Neftel et al., 2019; Suvà et al., 2014). It is thought that epigenetic heterogeneity in GBM reflects the co-existence of functionally non-equivalent populations of cells, similarly to what has been observed in normal developmental paradigms. This functional interpretation of epigenetic heterogeneity in GBM well aligns with the experimentally-validated concept that GBM is populated by cell populations with different functional properties, including slow-cycling self-renewing cells, proliferative progenitor-like cells with limited self-renewal and differentiated progeny that are destined to die (Cusulin et al., 2015a; Lan et al., 2017; Lathia et al., 2010; Miller et al., 2017; Singh et al., 2004; Suvà et al., 2014) .

GBM self-renewing cells (GSCs) are expandable in culture (Pollard et al., 2009), can differentiate into non-self-renewing cell types *in vitro* (Galli et al., 2004; Singh et al., 2003) and are capable of tumor initiation and serial propagation of the tumor in immunocompromised mice *in vivo* (Singh et al., 2004). Studies have shown GSCs are more resistant to the standard of care treatment for GBM than non-self-renewing cells, and they represent a larger fraction of tumor cells at relapse than at diagnosis (Bao et al., 2006; Liu et al., 2006). GSCs therefore play crucial roles in tumor growth, therapy resistance and in seeding relapses. However, therapeutic targeting of self-renewing cells is currently an unmet clinical need.

Pro-self-renewal epigenetic programs are maintained by the choreographed function of several chromatin remodelers. These include the histone methyltransferase DOT1L and arginine demethylase JMJD6 (MacLeod et al., 2019), members of the mixed-lineage leukemia (MLL) family (Gallo et al., 2013; Heddleston et al., 2012) and of the polycomb-repressive complexes 1 and 2 (PRC1/2) (Abdouh et al., 2009; Suvà et al., 2009), among others. These chromatin remodelers promote self-renewal by activating transcriptional networks associated with stemness and by repressing differentiation. This can be achieved through the activation of master transcription factors (Rheinbay et al., 2013; Suvà et al., 2014) or by modulating the expression of histone variants, which ultimately shape chromatin organization and downstream transcriptional programs .

Histone variants differ from core histones at key amino acids that determine their patterns of incorporation into chromatin and their effects on transcription of downstream genes (Maze et al., 2014). For instance, *H2AFY* and *H2AFY2* encode two closely related variants of core histone H2A, namely macroH2A1 and macroH2A2, respectively. They share only about 60% sequence identity in their histone domain with core histone H2A (Buschbeck and Hake, 2017), and contain two additional domains: A basic linker region with putative DNA binding function, and a "macro"

domain with little sequence conservation between the two paralogs (Costanzi and Pehrson, 2001).
Because of the large size of the macro domain, substitution of H2A with macroH2A1 or
macroH2A2 is predicted to have large effects on chromatin organization (Cantariño et al., 2013).
Loss or downregulation of macroH2A1 and macroH2A2 has been described in a number of
malignancies, including bladder cancer, melanoma, lung cancer, and gastric cancer (Kapoor et al.,
2010; Novikov et al., 2011; Sporn et al., 2009).
145
MacroH2A1 was first studied for its role in X chromosome inactivation in female mammalian cells
(Costanzi and Pehrson, 1998). It localizes to large chromatin domains throughout the genome that
encompass developmentally regulated genes and regions of imprinting, and it accumulates at
senescence-associated heterochromatic foci (Chen et al., 2015; Choo et al., 2006; Gamble et al.,
2010; Zhang et al., 2005, 2007). Both macroH2A1 and macroH2A2 are involved in the repair of
double-strand breaks, and knockdown of macroH2A1 impairs DNA double strand repair (Kim et
al., 2017, 2018; Kozlowski et al., 2018; Timinszky et al., 2009). MacroH2A2 loss has been
described in invasive melanoma (Kapoor et al., 2010). Knockdown or knockout of macroH2A
paralogs increase reprogramming efficiency of somatic cells into induced pluripotent stem cells
(Barrero et al., 2013), suggesting that these histone variants might be involved in repression of
stemness properties, including self-renewal.
157
The molecular mechanisms exploited by macroH2A paralogs in modulating stemness and self-
renewal are not currently understood, especially in the context of cancer. Given the current
inability to target self-renewing cells in most cancer types, dissection of these mechanisms could
be important to identify new treatment options for GBM and other malignancies characterized by
intratumoral functional heterogeneity. Here we deploy epigenomic and functional assays to
unravel the mechanisms utilized by macroH2A2 to control self-renewal programs in GBM, and
pursue potential translational avenues with our preclinical models.
165
166
167
168 **RESULTS**
169
Low expression of *H2AFY2* is a negative prognostic factor for GBM patients
We embarked on a systematic analysis to identify genes encoding histone variants that may be
involved in GBM biology. Our approach consisted in looking for histone variant genes whose
expression could stratify GBM patients based on overall survival. One of our best candidates was
H2AFY2, the gene encoding the histone H2A variant macroH2A2. Median expression levels were
used to classify patients as high- or low-expressors. We found that low levels of *H2AFY2*
transcription were associated with shorter overall survival in IDH-wildtype GBM in the
Gravendeel cohort (Gravendeel et al., 2009) (log-rank p = 0.0069), as well as in the GBM cohort
collected by The Cancer Genome Atlas (TCGA, 2015) (**Figure 1A, 1B**). Age is a potent
confounder for overall survival, with younger glioma patients usually performing better than older
patients. A multivariate Cox regression model also showed that increased *H2AFY2* expression had
prognostic significance in both cohorts (hazard ratio: 0.54 [0.38 – 0.78]; **Figure 1C; Figure S2A-
S2C**), even when adjusted for known prognostic factors such as IDH mutation status, age, and
treatment status **(Figure S2C)**. In the TCGA cohort, the survival benefit associated with high
H2AFY2 transcription was also observed in patients with recurrent disease, and these patients were

noted to have significantly higher levels of *H2AFY2* expression on average (**Figure S2D; Figure**
S2E). Transcription levels of *H2AFY* (which encodes macroH2A1) had no prognostic significance
in either patient cohort **(Figure S1B, S1C).** Moreover, the effects of *H2AFY2* appeared to be
specific to IDH-wildtype GBM, as no effect was seen in IDH-mutant tumours in the Gravendeel
dataset **(Figure S1D).** When RNA-seq data from the TCGA are separated by transcriptional
subtype, the survival benefit of *H2AFY2* expression is significant specifically in proneural tumours
(p = 0.0024; log-rank test) and not in classical or mesenchymal tumours (**Figure 1D**).
When expression of *H2AFY2* is compared to expression of *H2AFY* in the TCGA glioblastoma
cohort, the former has a clear bimodal distribution, a pattern not seen for *H2AFY* (macroH2A1)
(Figure S1E). These data point to a specific role for macroH2A2 in GBM chromatin biology.
We used transcriptome datasets from the Allen Brain Atlas BrainSpan database (Miller et al.,
2014) to extract information on *H2AFY2* expression during human brain development in samples
collected between 8 weeks post-conception and 40 years of age. We found that *H2AFY2* expression
peaks at 8 weeks post-conception, steadily declines during fetal development and stabilizes at 2-3
years of age into adulthood (**Figure 1E**). Our analyses of data from BrainSeq2 also show that
expression of *H2AFY2* is particularly elevated in fetal astrocytes compared to other brain cell types
(**Figure 1F**) (Zhang et al., 2016). Single-cell expression data from mouse striatum highlights a
fetal expression pattern for *H2afy2*, with preferential expression largely in neuronal precursor cells
and oligodendrocyte precursor cells **(Figure S1C)** (Saunders et al., 2018). Transcription of
H2AFY2 in the brain therefore appears to be temporally-regulated, with the highest expression
observed during early fetal development in both humans and mice.
Because *H2AFY2* levels have prognostic significance and are dynamically regulated during
brain development, we decided to investigate whether this gene plays a direct role in the biology
of GBM. We used our patient-derived GSC cultures to generate stable inducible *H2AFY2*
knockdown models of GBM. We transplanted GSCs carrying control scrambled shRNA constructs
(shScr) or either of two shRNAs targeting *H2AFY2* (shH2AFY2a/b) into the forebrains of NSG
mice. We observed reproducible negative effects of *H2AFY2* knockdown on mouse survival (log-
rank p < 0.001 for shH2AFY2a and p = 0.0027 for shH2AFY2b) (**Figure 1G**). Therefore our
patient-derived knockdown models recapitulate the association between low *H2AFY2* levels and
poor prognosis we observed in glioma patient cohorts (**Figures 1A,1B**). Importantly, our results
indicate that this gene plays a direct functional role in GBM biology and deserves further
investigation.
217

218
219
Figure 1. Low transcription levels of *H2AFY2*/macroH2A2 contribute to GBM aggressiveness.
(A, B) Kaplan-Meier survival analysis of adult *IDH*-wildtype glioblastoma patients (GSE16011; Gravendeel et al
2009; TCGA glioblastoma cohort) based on *H2AFY2* transcription levels. *H2AFY2*-low and -high groups were
determined by median gene expression. Shaded region represents 95% confidence interval. P value was obtained by
log-rank test. **(C)** Hazard ratios for macroH2A2 expression in a multivariate Cox regression model adjusting for
other factors relevant for glioblastoma (age, IDH mutation status). Error bars represent 95% confidence intervals.
(D) Kaplan-Meier survival analysis of adult glioblastoma patients separated by transcriptional subtype (TCGA).

Shaded region represents 95% confidence interval. P values obtained by log-rank test.**E)** Expression of *H2AFY2* at
different timepoints of brain development. Datasets were accessed through the BrainSpan database.
(F) Transcriptional levels of *H2AFY2* were assessed in different human brain cell types. Datasets were accessed
through the BrainSeq2 database.
(G) Orthotopic xenograft experiments to assess the effects of *H2AFY2* knockdown on survival of transplanted mice.
Patient-derived GSCs carrying either scrambled control shRNA constructs (shScr; n = 9) or independent shRNAs
targeting *H2AFY2* (shH2AFY2a/b; n = 10 mice per group) were transplanted orthotopically in immunocompromised
mice. P values were calculated with the log-rank test.
235
236

macroH2A2 antagonizes self-renewal in GBM
We previously reported that the histone H3 variant H3.3 has a strong role in repressing self-renewal
in GBM (Gallo et al., 2015). Self-renewal, or self-replication, is a key property of a population of
cells that propagate tumor growth (Magee et al., 2012). It was shown that the fraction of GBM
cells that can self-renew, as well as transcriptional signatures associated with self-renewal, are
negative prognostic factors for this brain cancer (Murat et al., 2008). Given the significance of
self-renewal in GBM, we investigated whether macroH2A2 plays a role in modulating this crucial
functional property. The frequency of self-renewing cells in a population can be estimated with *in*
vitro limiting dilution assays (LDAs), which use sphere-forming frequency as a measure of self-
renewal. We used our doxycycline-inducible *H2AFY2* knockdown models (**Figure 2A,B; Figure**
S2A) for *in vitro* LDAs in three different primary patient-derived cultures (G523, GSC3,GSC5),
and found that knocking down *H2AFY2* resulted in increased self-renewal (**Figure 2C-2E**). These
H2AFY2 knockdown constructs were specific to this H2A variant and showed no effect on protein
levels of the related variant macroH2A1 **(Figure S2B)**. In order to cross-check these results with
an independent experimental system, we decided to overexpress *H2AFY2* using a fusion protein
composed of catalytically dead Cas9 (dCas9) and the transcriptional activator VPR (dCas9-VPR
system (Chavez et al., 2015); **Figure 2F**). We transduced GSCs with a lentivirus carrying dCas9-
VPR to generate a stable inducible overexpression model. We then transfected this GSC line with
a pool of five single guide RNAs (sgRNAs) targeting *H2AFY2*. RT-qPCR showed that this
experimental system resulted in a 10-fold overexpression of *H2AFY2* over control cells transfected
with scrambled sgRNAs (sgScr), with no effects on transcription levels of the close paralog *H2AFY*
encoding the H2A variant macroH2A1 (**Figure 2G; Figure S2C, S2D**). We therefore employed
this *H2AFY2*-specific overexpression models in *in vitro* LDAs, which showed that *H2AFY2*
overexpression causes a ~50% reduction in sphere-forming frequency (**Figure 2H**). Therefore, our
knockdown and overexpression systems concordantly show an antagonistic effect of macroH2A2
on self-renewal in patient-derived GSC cultures.
To further validate these findings, we performed *in vivo* LDAs. We transplanted GBM cells
carrying our doxycycline-inducible stable *H2AFY2* system or shScr controls into the forebrains of
NSG mice (**Figure 2I**). We transplanted 10 mice at each cell dose (100,000, 10,000 and 1,000
cells) for knockdown and control cells. At the lowest dose, 10 out of 10 mice transplanted with
H2AFY2-knockdown cells developed tumors, whereas only 6 out of 10 control mice did (**Figure**
2J). Overall, we observed a significant difference in engraftment potential between GBM cells
with *H2AFY2* knockdown and control cells (χ^2 p = 0.0105). Our *in vivo* LDAs therefore confirm
our *in vitro* experiments and support an antagonistic role of macroH2A2 on self-renewal.
271
272
273
274

275
276
Figure 2. macroH2A2 antagonizes self-renewal in GBM.
(A) qRT-PCR to assess the transcriptional levels of *H2AFY2* in cells carrying control (shScr) or knockdown
(shH2AFY2a/b) constructs 48 hours after doxycycline induction. Expression normalized to actin and GAPDH. P
value were determined with a two-tailed t-test. The experiment was repeated 4 times.
(B) Western blot to compare macroH2A2 protein levels in control and knockdown cells after 14 days of shRNA
induction. The experiment was repeated 3 times.
(C-E) Limiting dilution assay results after 14 days of doxycycline induction in G523 glioma cells **(C)**, GSC3 **(D)**,
and GSC5 **(E)**. P value was determined by Chi-square test with the tool ELDA (see Methods). Error bars: 95%
confidence interval. Statistics from 6 technical replicates; the experiment was repeated 3 times.
(F) Schematic of the dCas9-VPR overexpression system.

(G) Validation of overexpression system by qRT-PCR after 48 hours of induction by sgRNA. Transcription was
normalized to actin and GAPDH. P value was determined with a two-tailed t-test. The experiment was repeated 2
times.
(H) Limiting dilution assay of G523 cells with dCas9-VPR and guide RNA targeting macroH2A2 or a non-targeting
guide control at 7 days. P value was determined by Chi-square test with the tool ELDA (see Methods). Error bars:
95% confidence interval. Statistics from 6 technical replicates; the experiment was repeated 3 times.
(I) Schematic of *in vivo* limiting dilution assay.
(J) Overview of *in vivo* limiting dilution assay results. Mice were transplanted orthotopically with either shScr or
shH2AFY2a-transduced GSCs. P value and chi square value obtained by Chi-square test.
296
297
298
299
macroH2A2 represses a transcriptional signature of self-renewal in GBM
Given that our *in vitro* and *in vivo* functional assays showcased a role for macroH2A2 in repressing
self-renewal properties of GBM cells, we hypothesized that this histone variant might regulate
expression of stemness genes. To test this hypothesis, we performed RNA-seq on GBM cells with
H2AFY2 knockdown (two independent hairpins) or shScr constructs (two biological replicates per
condition). We then looked at the effects of *H2AFY2* knockdown on a recently published GBM
stemness signature composed of 17 genes (Suvà et al., 2014). We found that overall, *H2AFY2*
knockdown resulted in increased transcription of these stemness genes, with strong and consistent
effects especially on *VAX2*, *SALL2*, *SOX8*, *SOX21*, *POU3F*3, *PDGFRA*, *ASCL1* and *OLIG2*
(**Figure 3A**). These data support a role for macroH2A2 in repressing self-renewal by antagonizing
a stemness transcriptional network in GBM.
Gene set enrichment analysis (GSEA) (Subramanian et al., 2005) of differentially expressed
genes upon *H2AFY2* knockdown revealed positive enrichment for signatures associated with the
proneural molecular subtype of GBM (Verhaak et al., 2010) (**Figure 3B**) and with
oligodendrocytic gene signatures (**Figure 3C**). Volcano plots further illustrated that some of the
most upregulated genes upon *H2AFY2* knockdown are oligodendrocyte markers, including
COL20A1 and *PDGFRA* (**Figure 3D**) (Rivers et al., 2008). When gene expression is compared to
the single-cell transcriptional subtypes identified by Neftel et al, we see an increase in the OPC
and NPC1 lineages by ssGSEA, and a reduction in mesenchymal (MES1) and astrocytic (AC)
lineages (**Figure 3E**). Interestingly, with shH2AFY2a, which has a stronger knockdown, we see
more of the MES and NPC2 phenotypes, which are not seen with the shH2AFY2b knockdown.
This suggests that macroH2A2 may act as a rheostat regulating the transitions between distinct
cellular fates. We also found that *CSPG4/NG2*, a widely-used marker of the oligodendrocyte
lineage, is transcribed at higher levels following *H2AFY2* knockdown (**Figure 3F**). Overall, our
data point to a role for macroH2A2 in repressing gene expression signatures associated with self-
renewal and the oligodendrocytic lineage. This is a particularly intriguing finding because
oligodendrocyte progenitor cells have been proposed as the cell of origin of GBM based on
experimental work with murine (Alcantara Llaguno et al., 2015; Liu et al., 2011) and patient-
derived (Yuan et al., 2018) models.

334
335   **Figure 3. MacroH2A2 knockdown leads to widespread gene expression changes in stemness and**
336   **oligodendroglial genes.**

(A) RNA-seq was used to determine transcriptional levels of an established stemness gene signature at 7 days following *H2AFY2*/macroH2A2 knockdown.Two biological replicates were used per condition.
(B,C) Two signatures enriched by GSEA in knockdown versus control cells.
(D) Volcano plot highlighting differentially expressed genes after 7 days of *H2AFY2*/macroH2A2 knockdown in G523 cells.
(E) Comparative ssGSEA analysis of the Neftel scRNA-seq GBM transcriptional subtypes (Neftel et al., 2019) in control versus *H2AFY2* knockdown samples.
(F) Heatmap showing differential expression of oligodendrocyte precursor cell-associated genes in knockdown cells versus control. Two biological replicates per condition.

MacroH2A2 maintains chromatin organization at developmentally regulated genes
MacroH2A variants have been mostly studied in the context of their roles in chromatin compaction, including X inactivation. However, there is emerging evidence that macroH2A variants can also be found at sites of open chromatin (Sun et al., 2018; Yildirim et al., 2014). To disambiguate the function of macroH2A2 in epigenetic programs in GBM, we performed the sequencing-based assay for transposase-accessible chromatin (ATAC-seq) in our patient-derived knockdown models. ATAC-seq was performed on two biological replicates for both shScr control cells and *H2A2FY2* knockdown cells. We found that *H2AFY2* knockdown caused both gains and losses of chromatin accessibility (**Figure 4A**), although losses were more frequent and globally resulted in fewer base pairs of DNA in accessible chromatin in knockdown cells (**Figure 4B**). The data support the notion that macroH2A2 contributes to the organization of both accessible and inaccessible chromatin regions, although its contributions to accessible regions are more prominent in our GBM models.

Gene ontology analysis of genes located in differential chromatin accessibility regions upon *H2AFY2* knockdown identified significant enrichment for genes involved in neurodevelopmental pathways, including the terms "nervous system development" and "axonal guidance" (**Figure 4C**). HOMER motif analysis (Heinz et al., 2010) revealed that chromatin accessibility changes occur at sites with significant enrichment for binding sites of transcription factors associated with neurodevelopment. For instance, sites that lost chromatin accessibility had an enrichment of c-JUN motifs (identified at 32.89% of targets compared to 2.04% of background genomic sites, p = 1e-43, hypergeometric test; **Figure 4D**), whereas sites that gained accessibility upon macroH2A2 knockdown had enrichment for SOX10 motifs (identified at 31.71% of targets compared to 13.93% of background genome-wide sites, p = 1e-4; hypergeometric test; **Figure 4E**), a transcription factor associated with neural crest development and oligodendrocyte differentiation (Claus Stolt et al., 2002; Finzsch et al., 2008; Stolt et al., 2004). These findings, together with the developmental regulation of *H2AFY2* we showed above (**Figures 1D,E**), implicate macroH2A2 in fine tuning brain-specific epigenetic and transcriptional programs of self-renewal by modulating chromatin organization.

Next, we investigated whether macroH2A2 levels had a greater impact on *global* chromatin architecture, as we have reported for the histone variant H3.3, compared to its effects on the *local* chromatin environment. We have previously shown that ATAC changepoint analysis provides a measure of global changes in chromatin architecture. We performed changepoint analysis using our ATAC-seq datasets generated with shScr and shH2AFY2 GBM cells and we did not observe large structural differences with the exception of chromosomes 3, 5 and 10 (**Figure 4F**). We conclude that macroH2A2 has dual roles in maintaining compacted and accessible regions without

384 causing large-scale chromatin reorganization. We therefore decided to investigate potential roles
385 of macroH2A2 in shaping the local chromatin environment.

396
397
Figure 4. MacroH2A2 contributes to both compacted and accessible chromatin at neurodevelopmental genes
in GBM.
(A) Heatmap of differentially accessible regions (n = 270) in two biological replicates of *H2AFY2*/macroH2A2
knockdown versus control cells.
(B) Total number of base pairs in peaks of accessible chromatin, as determined by ATAC-seq, in control versus
knockdown cells. P value was determined by one-way ANOVA.
(C) Top process terms resulting from Gene Ontology term analysis of differentially accessible regions.
(D) Example of a transcription factor motif overrepresented in regions of reduced accessibility in knockdown cells.
P value calculated by hypergeometric test.

macroH2A2 represses enhancer function in GSCs

macroH2A2 represses enhancer function in GSCs
There is very little known on the effects of macroH2A2 on chromatin organization and
transcriptional control, and this is particularly true in GBM. We initially hypothesized that
macroH2A2 might exert its effects on transcription by modulating chromatin accessibility at gene
bodies. Surprisingly, permutation analyses revealed a clear depletion of both gained and lost
ATAC peaks in gene bodies and their promoters upon *H2AFY2* knockdown (p = 0.002, **Figure
S3**). On the other hand, ATAC peaks that were gained upon knockdown showed only a slight but
significant depletion (p = 0.0039) at enhancer elements **(Figure 5A)**, while those lost upon
H2AFY2 knockdown were significantly over-represented at enhancer elements (p = 0.002; **Figure
5B**). Specific peaks gained upon knockdown were identified at putative enhancer regions
associated with developmental genes, including *FOXP1* and the oligodendrocyte lineage gene
COL20A1 **(Figure 5C,D)**.
Active enhancers are transcribed by RNA polymerase II, resulting in the production of enhancer
RNA (eRNA) (Li et al., 2016). eRNA levels often closely track the transcription levels of their
associated genes (Azofeifa et al., 2018; Hah et al., 2013). We reasoned that if macroH2A2 affects
the function of enhancer elements, changes in enhancer activity associated with depletion of
macroH2A2 should result in alterations in eRNA transcription levels. Analysis of RNA-seq data
from control and *H2AFY2*/macroH2A2 knockdown GBM cells showed significant differential
transcription of 33 distinct eRNA transcripts **(Figure S4A-S4E)**. Interestingly, we observed a
significant transcriptional increase of an eRNA at the *COL20A1* enhancer locus in our knockdown
cells **(Figure 5E)**, suggesting that macroH2A2 may play a role in inhibiting this enhancer site.
These data are consistent with the effects of *H2AFY2* knockdown on *COL20A1* transcription, and
suggest a mechanism by which macroH2A2 represses this gene associated with the
oligodendrocytic lineage by compacting the chromatin at its cognate enhancer element.
To further validate the differentially accessible regions we identified in our knockdown cells,
we analysed single-cell ATAC-seq (scATAC-seq) we had previously generated on 4 adult GBM
primary patient samples (Guilhamon et al., 2021). Adult GBM resections showed accessibility in
the majority of differentially accessible peaks gained and lost upon *H2AFY2* knockdown,
confirming the relevance of the results generated with our patient-derived models. Gained and lost
peaks were accessible in two distinct clusters of cells, suggesting that they define populations
characterized by different chromatin states **(Figure 5F)**. The *COL20A1* enhancer was also
identified in the primary tumor specimens, and its accessibility was restricted to a single cluster of
tumour cells in 2 out of 4 primary samples profiled by scATAC-seq **(Figure 5G; Figure S5A-C)**.
All together, our data show that *H2AFY2*/macroH2A2 has an important role in mediating enhancer
accessibility in GBM cells, and appears to modulate a regulatory network driving GSC cell
identity.

Figure 5. macroH2A2 represses enhancer elements linked to neurodevelopmental genes.
(A-B) Permutation analysis of accessible regions gained **(A)** and lost **(B)** upon *H2AFY2*/macroH2A2 knockdown at enhancer elements genome-wide.
(C, D) Examples of ATAC-seq enhancer peaks gained upon knockdown of *H2AFY2*/macroH2A2.

(E) Expression of eRNA at the *COL20A1* locus shown in **(D)** in control versus knockdown cells. Two biological
replicates were used to generate RNA-seq libraries. P value was obtained by two-tailed t-test..
(F) Differentially accessible chromatin regions identified upon *H2AFY2*/macroH2A2 knockdown in a scATAC-seq
sample from four primary glioblastoma resections. The horizontal axis represents individual cells in the specimens,
with differentially accessible regions listed along the Y axis.
(**G**) Accessibility at the *COL20A1* enhancer in a scATAC-seq dataset generated from four primary GBM surgical
specimens. Samples are separated by arcs.

High-content chemical screen identified compounds that increase macroH2A2 levels
We performed further analyses to explore associations between *H2AFY2* levels and clinical
parameters in previously published patient cohorts. Interestingly, we observed that *H2AFY2* levels
stratify survival only in patients that received radiation with or without chemotherapy (**Figures
6A**). In all cases, we observed that higher transcription levels of *H2AFY2* were associated with
better response to therapy, possibly because of its repressing effects on self-renewal programs,
which have been linked to radiation resistance. Because of the positive association between
macroH2A2 levels and response to radiation, we reasoned that increasing macroH2A2 levels could
lead to "differentiation" of GBM cells and could sensitize tumor cells to standard of care therapy.
We therefore set out to perform a chemical screen with the goal of identifying compounds that
could increase levels of macroH2A2 (**Figure 6B; Figure S6A**). We screened 182 compounds from
the Selleckchem Epigenetic Drug Library using the GE InCell automated confocal microscopy
system, coupled with immunofluorescence to assess macroH2A2 levels (**Figure 6C, Figure 6D**).
We were able to identify 35 compounds that led to a greater than two-fold increase in the
percentage of macroH2A2$^+$ GBM cells, including the MLL-menin inhibitor MI-3 and the HDAC
inhibitor RGFP966 (**Figure 6E**). We have previously reported that MLL-menin inhibition is a
potent antagonist of self-renewal and other stemness-associated properties (Gallo et al., 2015; Lan
et al., 2017), and we confirmed here that it has a potent negative effect on cell viability (**Figures
6F**). We validated that MI-3 increases macroH2A2 levels by western blot (**Figure 6G**). Similarly,
we found that RGFP966 could reduce cell viability with IC$_{50}$ < 1 μM (**Figure S6B**), thus
supporting the robustness of our screen. We decided to focus our follow up studies on MI-3
because of (1) its potent role at reducing self-renewal in GBM and (2) the translational potential
associated with Menin inhibition, as evidenced by the recently-reported development of clinical-
grade analogs (Krivtsov et al., 2019) and the active use of the Menin inhibitor KO-539 in a phase
1 clinical trial for patients with refractory or relapsed acute myelogenous leukemia
(ClinicalTrials.gov identifier: NCT04067336).

Figure 6. Drug screen to identify epigenetic compounds that elevate macroH2A2 levels in GBM cells.
(A) Survival analysis of patients in the TCGA GBM cohort segregated by median *H2AFY2* expression and treatment received (none, chemotherapy, radiotherapy, or both). Shaded region represents 95% confidence interval. P value was calculated by log-rank test.

(B) Diagram summarizing our screening strategy to identify compounds that increase macroH2A2 levels.
(C) Normalized density of the log fold change of macroH2A2 positive cells for all compounds in the screen. The
green shaded region represents compounds with greater than 2-fold change of macroH2A2 positive cells.
(D) Effects of MI-3 and vehicle control (DMSO) on macroH2A2 protein levels were assessed by
immunofluorescence. Scale bars: 50 μm.
(E) List of all compounds with greater than two fold increase of macroH2A2-positive viable cells upon treatment.
(F) Dose-response curve for MI-3 using GSC cultures, as measured by the Alamar Blue assay. Experiment
performed with six technical replicates per concentration, normalized to DMSO control. Error bars represent
standard deviation. The experiment was repeated 2 times.
(G) Western blot of macroH2A2 levels after 7 days of treatment with 200 nM of MI-3. Three replicates per
condition.
518
519
520

Menin inhibition activates viral mimicry in GBM cells
We then investigated potential mechanisms of action of this epigenetic compound. We treated
GBM cells with sub-lethal concentrations (200 nM) of MI-3 or DMSO vehicle control and
performed transcriptional studies by RNA-seq (3 biological replicates per condition). Our
transcriptomic studies showed that MI-3 treatment results in repression of markers of the
oligodendrocytic lineage, including *PDFGRA* and *COL20A1* (**Figure 7A**).
Repression of *PDGFRA* through MI-3 treatment was confirmed by RT-qPCR (**Figure 7B**) and
at the protein level by western blot (**Figure 7C**). Both *PDGFRA* and *COL20A1* were negatively
regulated by macroH2A2 (**Figure 3E**), and our data are therefore consistent with MI-3 acting
through macroH2A2 to execute chromatin and transcriptional programs, thus validating the
premises of our chemical screen.
We performed GSEA and found that differentially-regulated genes following MI-3 treatment
were positively associated with signatures of interferon gamma, alpha and beta signaling (**Figures
7D; Figure S7A**) and negatively associated with signatures linked to DNA methylation (**Figure
7E**) and PRC2-mediated methylation of histones (**Figure S7B**). Modulation of interferon
signalling related to altered DNA methylation has been reported with epigenetic treatments that
elicit viral mimicry in cancer cells (Chiappinelli et al., 2015; Roulois et al., 2015). Viral mimicry
is dependent on the transcription of repetitive elements across the genome, particularly ERVL
elements (Krug et al., 2019). We therefore re-analyzed our RNA-seq datasets to specifically look
at the effects of MI-3 treatment on the transcription of repetitive elements. We found that most
differentially transcribed repetitive elements were upregulated by MI-3 treatment, including ERVL
family members (**Figure 7F; Figure S7C, S7D**). Our data therefore reveal a role for Menin
inhibition in repressing self-renewal programs by upregulating macroH2A2 and potentially
activating viral mimicry pathways in GBM **(Figure 7G)**.
545
546
547
548

Figure 7. Menin inhibition induces differentiation of GSCs and activates viral mimicry pathways in GBM.
(**A**) Heatmap displaying the top 50 differentially expressed genes between MI-3 and DMSO treated cells based on
RNA-seq data (3 biological replicates per treatment).
(**B**) RT-qPCR results showing expression levels of *PDGFRA* in GBM cells after 7 days of MI-3 treatment.
Transcription was normalized to actin and GAPDH. P value was calculated by unpaired two-tailed T test. Error bars
represent standard deviation. The experiment was repeated two times.
557

(C) Western blot showing levels of PDGFRA and TNFR in MI-3 and DMSO treated GBM cells after 7 days of treatment in vitro. Three biological replicates per condition. The experiment was repeated two times.
(D,E) GSEA analyses of differentially-expressed genes in MI-3-treated versus vehicle-treated cells.
(F) Transcriptional levels of repeat elements upon MI-3 treatment were determined by RNA-seq.
(G) Proposed model for the mechanisms of action of macroH2A2 and Menin inhibition in GSCs.

DISCUSSION

Self-renewal is a fundamental property of cancer cells that are therapy-resistant and responsible for long-term tumor propagation. The identification of molecular mechanisms that are responsible for the attainment and maintenance of self-renewal could be key to significantly improve patient outcomes for difficult to treat cancers like GBM (Clarke, 2019). Self-renewal in GBM cells is dependent on the achievement of specific chromatin states and transcriptional profiles through the regulated function of chromatin remodelers, transcription factors and other epigenetic regulators (see for instance Bender et al., 2013; Gallo et al., 2013, 2015; Heddleston et al., 2012; Jin et al., 2017; Miller et al., 2017; Suvà et al., 2009). Interfering with chromatin and epigenetic factors that have key roles in the self-renewal program can destabilize these chromatin states and result in partial or complete loss of self-renewal.

In the present work, we identify macroH2A2 as a key antagonist of self-renewal properties in GBM. This conclusion is supported by multiple orthogonal lines of evidence. First, high *H2AFY2* transcript levels are associated with better response to radiotherapy and longer overall survival in GBM patient cohorts. These effects of *H2AFY2* were recapitulated in our *in vivo* orthotopic patient-derived models, showing a direct effect of this gene in regulating tumor aggressiveness. Second, our *in vitro* and *in vivo* functional assays demonstrated that macroH2A2 has a direct functional role in curbing self-renewal. Our genomic approaches showcased a new function for macroH2A2 in shaping chromatin organization at enhancer elements and regulating the expression of a self-renewal gene network. Overall these results indicate that macroH2A2 has an important effect on the regulation of chromatin and transcriptional dynamics that inhibit self-renewal in GBM.

The development of compounds that target chromatin and epigenetic factors – known as "epidrugs" – has opened new opportunities to pharmaceutically inhibit self-renewal networks in cancer cells. Some of these compounds have been shown to be effective and are currently in use in a subset of patients with certain types of hematological malignancies (Estey, 2013; Fenaux et al., 2009; San-Miguel et al., 2014) We reasoned that if high macroH2A2 levels are associated with better response to radiotherapy, treatment approaches that elevate macroH2A2 levels prior to radiation therapy could improve patient outcomes. We therefore performed a screen to identify compounds that could increase macroH2A2 levels. The Menin inhibitor MI-3 was one of our best hits.

We previously reported that Menin inhibition was a potent repressor of self-renewal in GBM (Lan et al., 2017). Other groups reported effectiveness of Menin inhibition in pediatric glioma models (Funato et al., 2014). However, the epigenetic mechanisms underlying the

effectiveness of Menin inhibition in glioma models have not yet been established. Here we show
that Menin inhibition acts through at least two complementary but converging mechanisms. First,
Menin inhibition results in increased macroH2A2 levels, which in turn repress self-renewal
transcriptional networks. Second, Menin inhibition causes activation of transcription of repetitive
elements that have been linked to the onset of viral mimicry pathways and lead to the demise of
cancer cells (Chiappinelli et al., 2015; Krug et al., 2019; Roulois et al., 2015). Importantly, the
Menin inhibitor KO-539 is being tested in a phase 1 clinical trial for patients with refractory or
relapsed acute myelogenous leukemia (ClinicalTrials.gov identifier: NCT04067336). A second
highly bioavailable Menin inhibitor was recently described and was shown to be effective against
patient-derived preclinical models of leukemia (Krivtsov et al., 2019). Targeting Menin is
therefore becoming an achievable reality in the clinic, and our findings suggest that this treatment
approach should further be tested in preclinical models of GBM.
617
Because repression of macroH2A2 contributes to aggressive phenotypes in several solid
malignancies, it is possible low levels of this histone variant could be used as a biomarker to
identify patients that could benefit from more aggressive treatment approaches. These patients
could receive clinical-grade Menin inhibitors to increase the levels of macroH2A2, which in turn
would repress self-renewal states and curb the aggressiveness of the disease. More studies will be
required to explore the translational opportunities presented by Menin inhibition in the context of
solid malignancies.
625
626 In conclusion, our work characterizes a previously unknown and targetable epigenetic
627 mechanism regulated by the histone variant macroH2A2. Our data underscore the close connection
628 between chromatin and functional states in cancer cells.
629
630
631
632 **ACKNOWLEDGMENTS**
633

Funding for this work was provided by a Canadian Institutes of Health Research (CIHR) grant
(PJT-156278) to MG; a Natural Sciences and Engineering Research Council (NSERC) Discovery
grant to MG; a Clinician Investigator Program fellowship to AN; an NSERC studentship to AB.
Our research was supported by SU2C Canada Cancer Stem Cell Dream Team Research Funding
(SU2C-AACR-DT-19-15) to PBD provided by the Government of Canada through Genome
Canada and CIHR, with supplemental support from the Ontario Institute for Cancer Research,
through funding provided by the Government of Ontario. SU2C Canada is a Canadian Registered
Charity (Reg.# 80550 6730 RR0001). Research funding is administered by the AACR
International - Canada, the Scientific Partner of SU2C Canada.

643
644
645
646 **MATERIALS AND METHODS**
647
Experimental Model and Subject Details
All specimens and primary cultures generated and used in this study were approved by the Health
Research Ethics Board of Alberta and the research ethics board of the Hospital for Sick Children.

GSC primary cultures G523, GSC3 and GSC5 were generated using previously described
methods; in brief, primary tumor sample was minced in Accutase (StemCell Technologies), and
dissociated with glass beads on a nutator for 30 minutes, followed by centrifugation and
resuspension in NS media (Pollard et al. 2009). Cultures were STR genotyped and confirmed to
match the patient tissue.
656
Cell Culture
Primary glioblastoma cultures were grown in adherent culture on Corning Primaria dishes coated
with poly-L-ornithine (Sigma-Aldrich, P4957) and laminin (Sigma-Aldrich, L2020), under
standard temperature, oxygen and humidity conditions. Cells were kept in NeuroCult NS-A Basal
Medium and Proliferation Supplement (StemCell Technologies, #05751), supplemented with 20
µg/mL rhEGF (Peprotech, AF-100-15), 10 µg/mL bFGF (StemCell Technologies, #78003), and 2
µg/mL heparin (StemCell Technologies, #07980). All cultures were used within the first 20
passages of generation. Adherent cells were disassociated with Accutase (Stemcell Technologies,
#07920) and plated onto fresh, coated plates when confluence reached 80/90%. Cell numbers and
viability were determined using Countess II (Thermo Fisher Scientific, AMQAX1000).
667
668 **Generation of macroH2A2 knockdown cultures**
669 Commercial inducible shRNA constructs (3 for *H2AFY2*, and non-targeting control 1) were
670 obtained from Dharmacon (Horizon Biosciences) and packaged into lentiviral particles. Primary
671 cells were infected with lentiviral particles in the presence of polybrene followed by selection
672 using 1.5 µg/mL puromycin for 72 hours.
673
MacroH2A2 overexpression with CRISPRa
Generation of an inducible overexpression line. The PB-TRE-dCas9-VPR construct (Addgene
#63800) (Chavez et al., 2015) was used to create a stable line using Piggybac transposase (System
Biosciences). A total of 500,000 cells were transfected using the using the mouse neural stem cell
nucleofection kit (Lonza, VPG-1004) with 0.66 µg of the construct and 0.33 µg of transposase
using the Lonza Amaxa Nucleofector I with protocol A-33, followed by selection with hygromycin
B at 50 ug/mL for 7 days.
681
682 *Generation of sgRNA plasmids.* Guide RNAs were designed using the crispr.mit.edu guide design
683 tool, using as templates the DNAse accessible region directly upstream of the TSS of
684 *H2AFY2*(Kent et al., 2002). The plasmid backbone pLKO-sgRNA-GFP (Addgene #57822) was
685 used and sgRNA plasmids were constructed using BsmBI digestion followed by ligation, as
686 previously described (Heckl et al 2014).
687
688 *Transfection for overexpression experiments.* Two million cells containing the dCas9-VPR
689 construct were transfected with a pool of multiple guides targeting *H2AFY2* or a non-targeting
690 control using the Amaxa Nucleofector. Doxycycline (2 µg/mL) was added to media 24 hours after
691 transfection.
692
***In vitro* limiting dilution analysis**
Cells were plated on uncoated low-adhesion 96-well plates in a two-fold dilution series spanning
from 2000 down to 4 cells per well in NeuroCult NS-A media (StemCell Technologies, #05751)
containing doxycycline at 2 µg/mL, with 6 replicates per concentration. Sphere formation

frequency was estimated using ELDA (Hu and Smyth 2009). Sphere formation was scored on day
7 and day 14.
699
700 **RT-qPCR**
701 RNA samples were used to generate cDNA using the SuperScript II kit (Invitrogen) and poly-A
702 primers. PCR was performed using the SSOFast EvaGreen Supermix (BioRad # 1725201) on the
703 BioRad CFX with all samples in triplicate. Results were analysed using the delta Ct method.
704
Western blot
Protein concentration of samples was determined using the DC (detergent compatible) protein
assay (Bio-Rad, #5000112). Samples were prepared in a total volume of 20 μL at 15 μg/μL in
Laemmli loading buffer. Samples were run on 7.5% Mini-PROTEAN gels for cytoplasmic
proteins and 12.5% Mini-PROTEAN gels for histones (Bio-Rad, #4568025). Primary antibodies
used: Rabbit anti-mH2A2 (Invitrogen; PA5-57437), mouse monoclonal anti-β-ACTIN (Sigma-
Aldrich, A5441, Lot# 127M4866V, Clone AC-15) at 1:1000), rabbit anti-PDGFRα (CST #3164),
rabbit monoclonal TNF-R1 (CST #3736). Secondary antibodies used: Goat anti-rabbit IgG H&L
(HRP) (Abcam, #6721, Lot# GR3192725-6) 1:20,000, Goat anti-mouse IgG H&L (HRP) (Abcam,
#6789).
715
Mouse intracranial orthotopic xenografts
Mouse survival. For each mouse, 100,000 tumor cells (control or knockdown) in PBS were
stereotactically injected into the forebrain (Location: 2.0-3.0 mm to the right of bregma, 1.0 mm
anterior to coronal suture) of 3 month old female NSG mice (Jackson Stock no 005557), using a
30 gauge needle. Mice were fed 2 mg/mL doxycycline in 2% sucrose water solution. Endpoint was
reached once mice showed signs of disease, including ataxia, hunching, domed heads, kyphosis,
paresis and lethargy.
723
724 *Orthotopic limiting dilution assay.* Stereotactic injections of 100,000, 10,000 or 1,000 cells were
725 performed into the right forebrain of NSG mice as described above. Mice were sacrificed at
726 endpoint as in the previous experiment.
727
ATAC-seq
Experimental method and sequencing. ATAC seq was performed using the Omni-ATAC protocol
(Corces et al., 2017). In brief, 50,000 cells were harvested fresh from culture from two biological
replicates, lysed on ice, spun at 4 C, and treated with Tn5 transposase (Illumina) for 30 minutes at
37 C. Libraries were amplified using the NebNext HiFi polymerase mastermix (New England
Biolabs) and standard Illumina Nextera primers, and DNA was purified using the MinElute kit
(Qiagen). Sequencing was performed on a NextSeq 500 using 150 cycles of paired end 75 bp
sequencing at the Centre for Health Genomics and Informatics (University of Calgary)..
736
ATAC-seq data analysis. ATAC-seq data was aligned to hg38 using bwa mem (Li and Durbin,
2009). Following this, samtools was used to remove the mitochondrial and Y chromosome reads,
and reads mapping to genome blacklists (Li et al., 2009). Duplicates were removed using picard.
Peak assignments were generated using macs2 (Zhang et al., 2008). Differential peak calls were
performed by generating a pileup of counts of all consensus peaks from all samples, transforming
the counts to counts per million using edgeR, and running these normalized counts through

DESeq2 (Love et al., 2014). Peaks with a fold change of > 1.5 were considered significantly
altered. Changepoint analysis was run on a pileup of ATAC signal reads binned into 1 mb bins
using bedtools, using the R package *changepoint*, as previously described (Gallo et al., 2015).
Permutation analysis was performed using the R package regioneR (Gel et al., 2016). Gene
ontology analysis was performed using DAVID 6.8 (Huang et al., 2009b, 2009a).
748
RNA extraction and RNA sequencing
Sample preparation and sequencing. For macroH2A2 knockdown RNA-seq, stably transduced
cells were induced with 2 µg/mL doxycycline and grown for 7 days in culture and processed in
biological duplicates. For MI-3 vs DMSO treated cells, G523 cells were treated with either 200
nM MI-3 or DMSO for 7 days and processed in biological triplicate. Cells were harvested using
Accutase, and RNA was extracted using the RNEasy Mini kit (Qiagen) as per manufacturer
instructions. Libraries were constructed at the Center for Health Genomics and Informatics using
the NEBNext Ultra II Directional RNA Library prep kit (New England Biolabs) with ribosomal
RNA depletion. Samples were sequenced on a NextSeq 500 for 150 cycles in single-end mode for
mH2A2 knockdown, and paired-end mode for MI-3.
759
RNA-seq analysis. Samples were pseudoaligned to the human transcriptome (GRCh38.rel79) using
kallisto, and differential analysis was performed using sleuth (Bray et al., 2016; Pimentel et al.,
2017). GSEA was performed using a ranked list of all genes generated using sleuth. For analysis
of eRNAs, the Fantom5 CAGE (Andersson et al., 2014) consensus list of enhancers, as well as a
custom list of predicted enhancers based on G523 ATAC-seq and H3K27ac consensus peaks, was
used to construct a custom pseudotranscriptome, which was analysed using kallisto and sleuth in
a similar fashion. Analysis of repeat expression was performed using REdiscoverTE (Kong et al.,
2019). Heatmaps were constructed with R using the heatmap.2 function from the package gplots.
768
Dose-response curves
Cells were plated at 4000 cells/well in NS media into laminin-poly-L-ornithine coated 96-well
plates (Corning). The compounds MI-3, RGFP-966 and AZD6102 (Selleckchem) were tested at
different concentrations (range: 2 µM to 8 nM in serial dilutions) with six technical replicates per
dose, and a DMSO control. Cell viability was assessed on day 7. Alamar blue (Thermo Fisher
Scientific, Cat# DAL1025) was added and cells were incubated at 37°C in the dark for 4 hours.
Fluorescence was measured on the Spectramax spectrophotometer and was normalized to the
DMSO control.
777
Immunofluorescence High-Content Drug Screening
Screen procedure. G523 cells were plated at a density of 5,000 cells per well in a 96-well optical
plate coated with PLO-laminin as per protocol. An epigenetic drug library (Z195677-L1900;
Selleckchem) was added at 1 µM to plates in triplicate, with DMSO used as a control. Cells were
incubated for 10 days and fixed in 4% PFA for 10 minutes followed by storage at 4°C until
imaging.
784
Immunocytochemistry and imaging. Plates were blocked in 5% BSA in PBS with 0.1% Triton-
X100 for 1 hour at room temperature. Staining was performed overnight at 4°C in the same media
with the macroH2A2 antibody (Invitrogen; PA5-57437) at a dilution of 1:250, and a fluorescently
conjugated secondary antibody (Invitrogen Goat anti-rabbit IgG A568), and a DAPI counterstain

(1:1000; Thermo Fisher #62248). Washes were performed using PBS with 0.1% Triton-X100.
Samples were left in the plates in PBS and imaged using the GE InCell 6000 with a 60x objective
in 3 different planes of section.
792
Image analysis. Images for each well were stitched together using Fiji and the grid/collection
stitching plugin (Preibisch et al., 2009). The DAPI channel was used to generate a list of nuclei
and Fiji was used to measure intensity for both DAPI and the GFP channel using an automated
custom script. Data was then further analysed using R, where data for all compounds on one plate
was pooled together, and k-means clustering using brightness and shape parameters was used to
stratify live/dead cells and mH2A2 positive or negative cells. These categories were then used to
separate cells in each well as alive and mH2A2-negative, alive and mH2A2-positive, and dead for
subsequent analysis. Percentages of mH2A2-positive cells were then compared to DMSO control
to calculate a fold change.
802
803 **Data availability**
804 All RNA-seq and ATAC-seq data will be deposited to the Gene Expression Omnibus (GEO) upon
805 publication of the manuscript. The scATAC-seq data of adult GBM has been published and is
806 available in GEO (accession: GSE139136) (Guilhamon et al., 2021). Analysis code is available
807 upon request.
808
809
810

REFERENCES
Abdouh, M., Facchino, S., Chatoo, W., Balasingam, V., Ferreira, J., and Bernier, G. (2009). BMI1
Sustains Human Glioblastoma Multiforme Stem Cell Renewal. J. Neurosci. *29*, 8884–8896.
Alcantara Llaguno, S.R., Wang, Z., Sun, D., Chen, J., Xu, J., Kim, E., Hatanpaa, K.J., Raisanen,
J.M., Burns, D.K., Johnson, J.E., et al. (2015). Adult Lineage-Restricted CNS Progenitors
Specify Distinct Glioblastoma Subtypes. Cancer Cell *28*, 429–440.
Alvarado, A.G., Thiagarajan, P.S., Mulkearns-Hubert, E.E., Silver, D.J., Hale, J.S., Alban, T.J.,
Turaga, S.M., Jarrar, A., Reizes, O., Longworth, M.S., et al. (2017). Glioblastoma Cancer Stem
Cells Evade Innate Immune Suppression of Self-Renewal through Reduced TLR4 Expression.
Cell Stem Cell.
Andersson, R., Gebhard, C., Miguel-Escalada, I., Hoof, I., Bornholdt, J., Boyd, M., Chen, Y.,
Zhao, X., Schmidl, C., Suzuki, T., et al. (2014). An atlas of active enhancers across human cell
types and tissues. Nature.
Antonios, J.P., Soto, H., Everson, R.G., Moughon, D., Orpilla, J.R., Shin, N.P., Sedighim, S.,
Treger, J., Odesa, S., Tucker, A., et al. (2017). Immunosuppressive tumor-infltrating myeloid
cells mediate adaptive immune resistance via a PD-1/PD-L1 mechanism in glioblastoma. Neuro.
Oncol.
Azofeifa, J.G., Allen, M.A., Hendrix, J.R., Read, T., Rubin, J.D., and Dowell, R.D. (2018).
Enhancer RNA profiling predicts transcription factor activity. Genome Res. *28*, 334–344.
Bao, S., Wu, Q., McLendon, R.E., Hao, Y., Shi, Q., Hjelmeland, A.B., Dewhirst, M.W., Bigner,
D.D., and Rich, J.N. (2006). Glioma stem cells promote radioresistance by preferential activation
of the DNA damage response. Nature *444*, 756–760.
Barrero, M.J., Sese, B., Martí, M., and Belmonte, J.C.I. (2013). Macro histone variants are critical
for the differentiation of human pluripotent cells. J. Biol. Chem. *288*, 16110–16116.

Bender, S., Tang, Y., Lindroth, A.M., Hovestadt, V., Jones, D.T.W., Kool, M., Zapatka, M.,
Northcott, P.A., Sturm, D., Wang, W., et al. (2013). Reduced H3K27me3 and DNA
Hypomethylation Are Major Drivers of Gene Expression in K27M Mutant Pediatric High-Grade
Gliomas. Cancer Cell *24*, 660–672.
Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
quantification. Nat. Biotechnol.
Buschbeck, M., and Hake, S.B. (2017). Variants of core histones and their roles in cell fate
decisions, development and cancer. Nat. Rev. Mol. Cell Biol. *18*, 299–314.
Cahill, D.P., Levine, K.K., Betensky, R.A., Codd, P.J., Romany, C.A., Reavie, L.B., Batchelor,
T.T., Futreal, P.A., Stratton, M.R., Curry, W.T., et al. (2007). Loss of the mismatch repair protein
MSH6 in human glioblastomas is associated with tumor progression during temozolomide
treatment. Clin. Cancer Res. *13*, 2038–2045.
Cantariño, N., Douet, J., and Buschbeck, M. (2013). MacroH2A - An epigenetic regulator of
cancer. Cancer Lett. *336*, 247–252.
Chavez, A., Scheiman, J., Vora, S., Pruitt, B.W., Tuttle, M., P R Iyer, E., Lin, S., Kiani, S.,
Guzman, C.D., Wiegand, D.J., et al. (2015). Highly efficient Cas9-mediated transcriptional
programming. Nat. Methods *12*, 326–328.
Chen, H., Ruiz, P.D., McKimpson, W.M., Novikov, L., Kitsis, R.N., and Gamble, M.J. (2015).
MacroH2A1 and ATM Play Opposing Roles in Paracrine Senescence and the Senescence-
Associated Secretory Phenotype. Mol. Cell *59*, 719–731.
Chen, J., Li, Y., Yu, T.S., McKay, R.M., Burns, D.K., Kernie, S.G., and Parada, L.F. (2012). A
restricted cell population propagates glioblastoma growth after chemotherapy. Nature *488*, 522–
526.
Chiappinelli, K.B., Strissel, P.L., Desrichard, A., Li, H., Henke, C., Akman, B., Hein, A., Rote,
N.S., Cope, L.M., Snyder, A., et al. (2015). Inhibiting DNA Methylation Causes an Interferon
Response in Cancer via dsRNA Including Endogenous Retroviruses. Cell *162*, 974–986.
Choo, J.H., Kim, J. Do, Chung, J.H., Stubbs, L., and Kim, J. (2006). Allele-specific deposition of
macroH2A1 in imprinting control regions. Hum. Mol. Genet. *15*, 717–724.
Clarke, M.F. (2019). Clinical and therapeutic implications of cancer stem cells. N. Engl. J. Med.
380, 2237–2245.
Claus Stolt, C., Rehberg, S., Ader, M., Lommes, P., Riethmacher, D., Schachner, M., Bartsch, U.,
and Wegner, M. (2002). Terminal differentiation of myelin-forming oligodendrocytes depends
on the transcription factor Sox10. Genes Dev. *16*, 165–170.
Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong, N.A., Vesuna,
S., Satpathy, A.T., Rubin, A.J., Montine, K.S., Wu, B., et al. (2017). An improved ATAC-seq
protocol reduces background and enables interrogation of frozen tissues. Nat. Methods.
Costanzi, C., and Pehrson, J.R. (1998). Histone macroH2A1 is concentrated in the inactive X
chromosome of Female Mammals. Nature *628*, 1997–1999.
Costanzi, C., and Pehrson, J.R. (2001). MACROH2A2, a New Member of the MACROH2A Core
Histone Family. J. Biol. Chem. *276*, 21776–21784.
Cusulin, C., Chesnelong, C., Bose, P., Bilenky, M., Kopciuk, K., Chan, J.A., Cairncross, J.G.,
Jones, S.J., Marra, M.A., Luchman, H.A., et al. (2015a). Precursor States of Brain Tumor
Initiating Cell Lines Are Predictive of Survival in Xenografts and Associated with Glioblastoma
Subtypes. Stem Cell Reports *5*, 1–9.
Cusulin, C., Chesnelong, C., Bose, P., Bilenky, M., Kopciuk, K., Chan, J.A., Cairncross, J.G.,
Jones, S.J., Marra, M.A., Luchman, H.A., et al. (2015b). Precursor States of Brain Tumor

Initiating Cell Lines Are Predictive of Survival in Xenografts and Associated with Glioblastoma
Subtypes. Stem Cell Reports.

Estey, E.H. (2013). Epigenetics in clinical practice: The examples of azacitidine and decitabine in
myelodysplasia and acute myeloid leukemia. Leukemia.

Fenaux, P., Mufti, G.J., Hellstrom-Lindberg, E., Santini, V., Finelli, C., Giagounidis, A., Schoch,
R., Gattermann, N., Sanz, G., List, A., et al. (2009). Efficacy of azacitidine compared with that
of conventional care regimens in the treatment of higher-risk myelodysplastic syndromes: a
randomised, open-label, phase III study. Lancet Oncol.

Finzsch, M., Stolt, C.C., Lommes, P., and Wegner, M. (2008). Sox9 and Sox10 influence survival
and migration of oligodendrocyte precursors in the spinal cord by regulating PDGF receptor α
expression. Development *135*, 637–646.

Funato, K., Major, T., Lewis, P.W., Allis, C.D., and Tabar, V. (2014). Use of human embryonic
stem cells to model pediatric gliomas with H3.3K27M histone mutation. Science *346*, 1529–
1533.

Galli, R., Binda, E., Orfanelli, U., Cipelletti, B., Gritti, A., De Vitis, S., Fiocco, R., Foroni, C.,
Dimeco, F., and Vescovi, A. (2004). Isolation and characterization of tumorigenic, stem-like
neural precursors from human glioblastoma. Cancer Res. *64*, 7011–7021.

Gallo, M., Ho, J., Coutinho, F., Vanner, R., Lee, L., Head, R., Ling, E., Clarke, I., and Dirks, P.
(2013). A Tumorigenic MLL-Homeobox Network in Human Glioblastoma Stem Cells. Cancer
Res. *73*, 417–427.

Gallo, M., Coutinho, F.J., Vanner, R.J., Gayden, T., Mack, S.C., Murison, A., Remke, M., Li, R.,
Takayama, N., Desai, K., et al. (2015). MLL5 Orchestrates a Cancer Self-Renewal State by
Repressing the Histone Variant H3.3 and Globally Reorganizing Chromatin. Cancer Cell *28*,
715–729.

Gamble, M.J., Frizzell, K.M., Yang, C., Krishnakumar, R., and Kraus, W.L. (2010). The histone
variant macroH2A1 marks repressed autosomal chromatin, but protects a subset of its target
genes from silencing. Genes Dev. *24*, 21–32.

Gel, B., Díez-Villanueva, A., Serra, E., Buschbeck, M., Peinado, M.A., and Malinverni, R. (2016).
RegioneR: An R/Bioconductor package for the association analysis of genomic regions based on
permutation tests. Bioinformatics.

Gravendeel, L.A.M., Kouwenhoven, M.C.M., Gevaert, O., De Rooi, J.J., Stubbs, A.P., Duijm, J.E.,
Daemen, A., Bleeker, F.E., Bralten, L.B.C., Kloosterhof, N.K., et al. (2009). Intrinsic gene
expression profiles of gliomas are a better predictor of survival than histology. Cancer Res. *69*,
9065–9072.

Guilhamon, P., Chesnelong, C., Kushida, M.M., Nikolic, A., Singhal, D., Macleod, G.,
Tonekaboni, S.A.M., Cavalli, F.M.G., Arlidge, C., Rajakulendran, N., et al. (2021). Single-cell
chromatin accessibility profiling of glioblastoma identifies an invasive cancer stem cell
population associated with lower survival. Elife *10*, e64090.

Hah, N., Murakami, S., Nagari, A., Danko, C.G., and Lee Kraus, W. (2013). Enhancer transcripts
mark active estrogen receptor binding sites. Genome Res. *23*, 1210–1223.

Heddleston, J.M., Wu, Q., Rivera, M., Minhas, S., Lathia, J.D., Sloan, A.E., Iliopoulos, O.,
Hjelmeland, A.B., and Rich, J.N. (2012). Hypoxia-induced mixed-lineage leukemia 1 regulates
glioma stem cell tumorigenic potential. Cell Death Differ. *19*, 428–439.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh,
H., and Glass, C.K. (2010). Simple Combinations of Lineage-Determining Transcription Factors
Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. Mol. Cell *38*,

576–589.

Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009a). Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. Nat. Protoc. *4*, 44–57.

Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009b). Bioinformatics enrichment tools: Paths
toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res. *37*, 1–13.

Hunter, C., Smith, R., Cahill, D.P., Stephens, P., Stevens, C., Teague, J., Greenman, C., Edkins,
S., Bignell, G., Davies, H., et al. (2006). A hypermutation phenotype and somatic MSH6
mutations in recurrent human malignant gliomas after alkylator chemotherapy. Cancer Res. *66*,
3987–3991.

Jin, X., Kim, L.J.Y., Wu, Q., Wallace, L.C., Prager, B.C., Sanvoranart, T., Gimple, R.C., Wang,
X., Mack, S.C., Miller, T.E., et al. (2017). Targeting glioma stem cells through combined BMI1
and EZH2 inhibition. Nat. Med. *23*, 1352–1361.

Kapoor, A., Goldberg, M.S., Cumberland, L.K., Ratnakumar, K., Segura, M.F., Emanuel, P.O.,
Menendez, S., Vardabasso, C., LeRoy, G., Vidal, C.I., et al. (2010). The histone variant
macroH2A suppresses melanoma progression through regulation of CDK8. Nature *468*, 1105–
1111.

Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., and Haussler,
a. D. (2002). The Human Genome Browser at UCSC. Genome Res.

Kim, B.J., Chan, D.W., Jung, S.Y., Chen, Y., Qin, J., and Wang, Y. (2017). The Histone Variant
MacroH2A1 Is a BRCA1 Ubiquitin Ligase Substrate. Cell Rep. *19*, 1758–1766.

Kim, J., Sturgill, D., Sebastian, R., Khurana, S., Tran, A.D., Edwards, G.B., Kruswick, A., Burkett,
S., Hosogane, E.K., Hannon, W.W., et al. (2018). Replication Stress Shapes a Protective
Chromatin Environment across Fragile Genomic Regions. Mol. Cell *69*, 36-47.e7.

Kong, Y., Rose, C.M., Cass, A.A., Williams, A.G., Darwish, M., Lianoglou, S., Haverty, P.M.,
Tong, A.J., Blanchette, C., Albert, M.L., et al. (2019). Transposable element expression in tumors
is associated with immune infiltration and increased antigenicity. Nat. Commun.

Kozlowski, M., Corujo, D., Hothorn, M., Guberovic, I., Mandemaker, I.K., Blessing, C., Sporn,
J., Gutierrez-Triana, A., Smith, R., Portmann, T., et al. (2018). MacroH2A histone variants limit
chromatin plasticity through two distinct mechanisms. EMBO Rep. *19*, e44445.

Krivtsov, A. V., Evans, K., Gadrey, J.Y., Eschle, B.K., Hatton, C., Uckelmann, H.J., Ross, K.N.,
Perner, F., Olsen, S.N., Pritchard, T., et al. (2019). A Menin-MLL Inhibitor Induces Specific
Chromatin Changes and Eradicates Disease in Models of MLL-Rearranged Leukemia. Cancer
Cell *36*, 660–673.

Krug, B., De Jay, N., Harutyunyan, A.S., Deshmukh, S., Marchione, D.M., Guilhamon, P.,
Bertrand, K.C., Mikael, L.G., McConechy, M.K., Chen, C.C.L., et al. (2019). Pervasive H3K27
Acetylation Leads to ERV Expression and a Therapeutic Vulnerability in H3K27M Gliomas.
Cancer Cell *35*, 782–797.

Lan, X., Jörg, D.J., Cavalli, F.M.G., Richards, L.M., Nguyen, L. V., Vanner, R.J., Guilhamon, P.,
Lee, L., Kushida, M.M., Pellacani, D., et al. (2017). Fate mapping of human glioblastoma reveals
an invariant stem cell hierarchy. Nature *549*, 227–232.

Lathia, J.D., Gallagher, J., Heddleston, J.M., Wang, J., Eyler, C.E., MacSwords, J., Wu, Q.,
Vasanji, A., McLendon, R.E., Hjelmeland, A.B., et al. (2010). Integrin Alpha 6 regulates
glioblastoma stem cells. Cell Stem Cell *6*, 421–432.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and

Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*,
2078–2079.

Li, W., Notani, D., and Rosenfeld, M.G. (2016). Enhancers as non-coding RNA transcription units:
Recent insights and future perspectives. Nat. Rev. Genet. *17*, 207–223.

Liu, C., Sage, J.C., Miller, M.R., Verhaak, R.G.W., Hippenmeyer, S., Vogel, H., Foreman, O.,
Bronson, R.T., Nishiyama, A., Luo, L., et al. (2011). Mosaic analysis with double markers reveals
tumor cell of origin in glioma. Cell *146*, 209–221.

Liu, G., Yuan, X., Zeng, Z., Tunici, P., Ng, H., Abdulkadir, I.R., Lu, L., Irvin, D., Black, K.L., and
Yu, J.S. (2006). Analysis of gene expression and chemoresistance of CD133+ cancer stem cells
in glioblastoma. Mol. Cancer *5*, 1–12.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

MacLeod, G., Bozek, D.A., Rajakulendran, N., Monteiro, V., Ahmadi, M., Steinhart, Z., Kushida,
M.M., Yu, H., Coutinho, F.J., Cavalli, F.M.G., et al. (2019). Genome-Wide CRISPR-Cas9
Screens Expose Genetic Vulnerabilities and Mechanisms of Temozolomide Sensitivity in
Glioblastoma Stem Cells. Cell Rep. *27*, 971-986.e9.

Magee, J.A., Piskounova, E., and Morrison, S.J. (2012). Cancer Stem Cells: Impact,
Heterogeneity, and Uncertainty. Cancer Cell *21*, 283–296.

Maze, I., Noh, K.M., Soshnev, A.A., and Allis, C.D. (2014). Every amino acid matters: Essential
contributions of histone variants to mammalian development and disease. Nat. Rev. Genet. *15*,
259–271.

Miller, J.A., Ding, S.L., Sunkin, S.M., Smith, K.A., Ng, L., Szafer, A., Ebbert, A., Riley, Z.L.,
Royall, J.J., Aiona, K., et al. (2014). Transcriptional landscape of the prenatal human brain.
Nature.

Miller, T.E., Liau, B.B., Wallace, L.C., Morton, A.R., Xie, Q., Dixit, D., Factor, D.C., Kim, L.J.Y.,
Morrow, J.J., Wu, Q., et al. (2017). Transcription elongation factors represent in vivo cancer
dependencies in glioblastoma. Nature *547*, 355–359.

Murat, A., Migliavacca, E., Gorlia, T., Lambiv, W.L., Shay, T., Hamou, M.F., De Tribolet, N.,
Regli, L., Wick, W., Kouwenhoven, M.C.M., et al. (2008). Stem cell-related "self-renewal"
signature and high epidermal growth factor receptor expression associated with resistance to
concomitant chemoradiotherapy in glioblastoma. J. Clin. Oncol. *26*, 3015–3024.

Neftel, C., Laffy, J., Filbin, M.G., Hara, T., Shore, M.E., Rahme, G.J., Richman, A.R., Silverbush,
D., Shaw, M.L., Hebert, C.M., et al. (2019). An Integrative Model of Cellular States, Plasticity,
and Genetics for Glioblastoma. Cell *178*, 835-849.e21.

Novikov, L., Park, J.W., Chen, H., Klerman, H., Jalloh, A.S., and Gamble, M.J. (2011). QKI-
Mediated Alternative Splicing of the Histone Variant MacroH2A1 Regulates Cancer Cell
Proliferation. Mol. Cell. Biol. *31*, 4244–4255.

Pimentel, H., Bray, N.L., Puente, S., Melsted, P., and Pachter, L. (2017). Differential analysis of
RNA-seq incorporating quantification uncertainty. Nat. Methods.

Pollard, S.M., Yoshikawa, K., Clarke, I.D., Danovi, D., Stricker, S., Russell, R., Bayani, J., Head,
R., Lee, M., Bernstein, M., et al. (2009). Glioma Stem Cell Lines Expanded in Adherent Culture
Have Tumor-Specific Phenotypes and Are Suitable for Chemical and Genetic Screens. Cell Stem
Cell *4*, 568–580.

Preibisch, S., Saalfeld, S., and Tomancak, P. (2009). Globally optimal stitching of tiled 3D
microscopic image acquisitions. Bioinformatics.

Rheinbay, E., Suvà, M.L., Gillespie, S.M., Wakimoto, H., Patel, A.P., Shahid, M., Oksuz, O.,

Rabkin, S.D., Martuza, R.L., Rivera, M.N., et al. (2013). An Aberrant Transcription Factor
Network Essential for Wnt Signaling and Stem Cell Maintenance in Glioblastoma. Cell Rep. *3*,
1567–1579.

Rivers, L.E., Young, K.M., Rizzi, M., Jamen, F., Psachoulia, K., Wade, A., Kessaris, N., and
Richardson, W.D. (2008). PDGFRA/NG2 glia generate myelinating oligodendrocytes and
piriform projection neurons in adult mice. Nat. Neurosci. *11*, 1392–1401.

Roulois, D., Loo Yau, H., Singhania, R., Wang, Y., Danesh, A., Shen, S.Y., Han, H., Liang, G.,
Jones, P.A., Pugh, T.J., et al. (2015). DNA-Demethylating Agents Target Colorectal Cancer Cells
by Inducing Viral Mimicry by Endogenous Transcripts. Cell *162*, 961–973.

San-Miguel, J.F., Hungria, V.T.M., Yoon, S.S., Beksac, M., Dimopoulos, M.A., Elghandour, A.,
Jedrzejczak, W.W., Günther, A., Nakorn, T.N., Siritanaratkul, N., et al. (2014). Panobinostat plus
bortezomib and dexamethasone versus placebo plus bortezomib and dexamethasone in patients
with relapsed or relapsed and refractory multiple myeloma: A multicentre, randomised, double-
blind phase 3 trial. Lancet Oncol.

Saunders, A., Macosko, E.Z., Wysoker, A., Goldman, M., Krienen, F.M., de Rivera, H., Bien, E.,
Baum, M., Bortolin, L., Wang, S., et al. (2018). Molecular Diversity and Specializations among
the Cells of the Adult Mouse Brain. Cell.

Singh, S.K., Hawkins, C., Clarke, I.D., Squire, J.A., Bayani, J., Hide, T., Henkelman, R.M.,
Cusimano, M.D., Dirks, P.B., Terasaki, M., et al. (2003). Identification of a cancer stem cell in
human brain tumors. Cancer Res. *63*, 5821–5828.

Singh, S.K., Hawkins, C., Clarke, I.D., Squire, J.A., Bayani, J., Hide, T., Henkelman, R.M.,
Cusimano, M.D., and Dirks, P.B. (2004). Identification of human brain tumour initiating cells.
Nature *432*, 396–401.

Sporn, J.C., Kustatscher, G., Hothorn, T., Collado, M., Serrano, M., Muley, T., Schnabel, P., and
Ladurner, A.G. (2009). Histone macroH2A isoforms predict the risk of lung cancer recurrence.
Oncogene *28*, 3423–3428.

Stolt, C.C., Lommes, P., Friedrich, R.P., and Wegner, M. (2004). Transcription factors Sox8 and
Sox10 perform non-equivalent roles during oligodendrocyte development despite functional
redundancy. Development *131*, 2349–2358.

Stupp, R., Mason, W.P., Van Den Bent, M.J., Weller, M., Fisher, B., Taphoorn, M.J.B., Belanger,
K., Brandes, A.A., Marosi, C., Bogdahn, U., et al. (2005). Radiotherapy plus concomitant and
adjuvant temozolomide for glioblastoma. N. Engl. J. Med. *352*, 987–996.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment
analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proc.
Natl. Acad. Sci. U. S. A.

Sun, Z., Filipescu, D., Andrade, J., Gaspar-Maia, A., Ueberheide, B., and Bernstein, E. (2018).
Transcription-associated histone pruning demarcates macroH2A chromatin domains. Nat. Struct.
Mol. Biol. *25*, 958–970.

Suvà, M.L., Riggi, N., Janiszewska, M., Radovanovic, I., Provero, P., Stehle, J.C., Baumer, K., Le
Bitoux, M.A., Marino, D., Cironi, L., et al. (2009). EZH2 is essential for glioblastoma cancer
stem cell maintenance. Cancer Res. *69*, 9211–9218.

Suvà, M.L., Rheinbay, E., Gillespie, S.M., Patel, A.P., Wakimoto, H., Rabkin, S.D., Riggi, N.,
Chi, A.S., Cahill, D.P., Nahed, B. V., et al. (2014). Reconstructing and reprogramming the tumor-
propagating potential of glioblastoma stem-like cells. Cell *157*, 580–594.

TCGA (2015). Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas.

N. Engl. J. Med. *372*, 2481–2498.

Timinszky, G., Till, S., Hassa, P.O., Hothorn, M., Kustatscher, G., Nijmeijer, B., Colombelli, J.,
Altmeyer, M., Stelzer, E.H.K., Scheffzek, K., et al. (2009). A macrodomain-containing histone
rearranges chromatin upon sensing PARP1 activation. Nat. Struct. Mol. Biol. *16*, 923–929.

Verhaak, R.G.W., Hoadley, K.A., Purdom, E., Wang, V., Qi, Y., Wilkerson, M.D., Miller, C.R.,
Ding, L., Golub, T., Mesirov, J.P., et al. (2010). Integrated Genomic Analysis Identifies
Clinically Relevant Subtypes of Glioblastoma Characterized by Abnormalities in PDGFRA,
IDH1, EGFR, and NF1. Cancer Cell *17*, 98–110.

Yildirim, O., Hung, J.H., Cedeno, R.J., Weng, Z., Lengner, C.J., and Rando, O.J. (2014). A System
for Genome-Wide Histone Variant Dynamics In ES Cells Reveals Dynamic MacroH2A2
Replacement at Promoters. PLoS Genet. *10*, e1004515.

Yip, S., Miao, J., Cahill, D.P., Iafrate, A.J., Aldape, K., Nutt, C.L., and Louis, D.N. (2009). MSH6
mutations arise in glioblastomas during temozolomide therapy and mediate temozolomide
resistance. Clin. Cancer Res. *15*, 4622–4629.

Yuan, J., Levitin, H.M., Frattini, V., Bush, E.C., Boyett, D.M., Samanamud, J., Ceccarelli, M.,
Dovas, A., Zanazzi, G., Canoll, P., et al. (2018). Single-cell transcriptome analysis of lineage
diversity in high-grade glioma. Genome Med. *10*, 57.

Zhang, R., Poustovoitov, M. V., Ye, X., Santos, H.A., Chen, W., Daganzo, S.M., Erzberger, J.P.,
Serebriiskii, I.G., Canutescu, A.A., Dunbrack, R.L., et al. (2005). Formation of macroH2A-
containing senescence-associated heterochromatin foci and senescence driven by ASF1a and
HIRA. Dev. Cell *8*, 19–30.

Zhang, R., Chen, W., and Adams, P.D. (2007). Molecular Dissection of Formation of Senescence-
Associated Heterochromatin Foci. Mol. Cell. Biol. *27*, 2343–2358.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nussbaum, C.,
Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS).
Genome Biol. *9*, R137.

Zhang, Y., Sloan, S.A., Clarke, L.E., Caneda, C., Plaza, C.A., Blumenthal, P.D., Vogel, H.,
Steinberg, G.K., Edwards, M.S.B., Li, G., et al. (2016). Purification and Characterization of
Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences
with Mouse. Neuron.

1111 **Supplementary Figures**

1112
Supplementary Figure S1, related to Figure 1.
(A) Overall survival of the TCGA GBM cohort stratified by median *H2AFY2* level. P value
calculated by log-rank test. Shaded region represents 95% confidence interval.

(B) Overall survival for *IDH* mutant gliomas in GSE 16011 stratified by median *H2AFY2*. P
value calculated by log-rank test. Shaded region represents 95% confidence interval.
(C) and **(D)** Overall survival of *IDH* wildtype GBM (GSE 16011) and TCGA GBM stratified by
median *H2AFY* level. P value calculated by log-rank test. Shaded region represents 95%
confidence interval.
(E) Expression data for *H2AFY* and *H2AFY2* in the TCGA GBM cohort.
(F) Single cell expression data for *H2afy2* in mouse striatum (DropViz mouse datasets).
1123

1124
Supplementary Figure 2, related to Figure 1. Multivariate cox regression of *H2AFY2*
expression in GSE 16011 **(A)** and TCGA GBM **(B)**. **(C)** Cox multivariate regression and hazard
ratio for *H2AFY2* expression status with age and chemotherapy and radiotherapy status in TCGA
GBM. **(D)** Kaplan-Meier survival status for patients with recurrent glioblastoma in TCGA cohort
(shaded region represents 95% confidence interval; p value calculated by log-rank test). **(E)**
Comparison of expression levels for *H2AFY2* in TCGA data for untreated versus treated primary
glioblastoma (p value: two-tailed T test with Welch's correction; whiskers represent 95%
confidence interval).
1133
1134

1135
1136

Supplementary Figure S2, related to Figure 2.
(A) RT-qPCR of macroH2A2 expression after 48 hours of doxycycline induction for two distinct hairpins. Expression normalized to actin and GAPDH. P value calculated by unpaired two-tailed T test. Error bars represent standard deviation. Experiment repeated three times.
(B) Western blot of macroH2A2 and macroH2A1 after 7 or 14 days of activation with doxycycline. Experiment repeated two times.
(C) RT-qPCR of macroH2A1 and macroH2A2 CRISPR activation in a second primary GSC culture with Expression normalized to actin and GAPDH. P value calculated by unpaired two-tailed T test. Error bars represent standard deviation. Experiment repeated two times.
(D) Western blot of macroH2A2 levels after 7 days of CRISPR activation in a macroH2A2-low GSC culture. Experiment repeated two times.

1148
1149

1150
1151 **Supplementary Figure S3, related to Figure 5.**
1152 Permutation analysis of regions with at least 1.5 fold $\log_2$ change in accessibility upon *H2AFY2*
1153 knockdown with **(A-B)** DNA loop boundaries (Johnston et al 2019); **(C-D)** Gene bodies; **(E-F)**
1154 repeat regions from the RepeatMasker database; **(G-H)** Introns. Regions of accessibility gain and
1155 loss were analysed separately. P value from 500 permutations.
1156

Supplementary Figure S4, related to Figure 5.
(A) Differential expression of top 200 differentially expressed GBM-specific putative eRNAs
between *H2AFY2* knockdown and control cells.
(B) Top 2000 differentially transcribed GBM-specific eRNAs between *H2AFY2*/macroH2A2
knockdown and control cells, ranked by fold change.
(C) Close up of significantly differentially transcribed eRNA, ranked by p value.
(D) Example of a representative ATAC-seq enhancer peak lost upon *H2AFY2*/macroH2A2
knockdown.
(E) Example of eRNA at an enhancer locus with increased accessibility. P value calculated by
unpaired T test. Error bars represent standard deviation.

1169
1170
1171 **Supplementary Figure S5, related to Figure 5.**
1172 **(A-C)** tSNE plots of scATAC-seq data from three different primary GBM
1173 resectionsAccessibility at the *COL20A1* enhancer locus is highlighted in orange.
1174

1175

A

| Screen GSCs (epigenetic drug library; 1 mM for 10 days; 3 replicates) | → | Stain for: macroH2A2 DAPI and image on InCell | → | Image analysis (ImageJ) Object detection and quantification | → | Data integration (R) |

| Cluster individual cells DAPI macroH2A2 morphology | → | Categorize clusters: dead alive macroH2A2+ | → | Summarize each well | → | Plot and save data |

B

RGFP-966

1176
1177
1178 **Supplementary Figure S6, related to Figure 6.**
1179 **(A)** Overview of high-content screening strategy.
1180 **(B)** Alamar blue *in vitro* dose-response curve for RGFP-966 in G523 cells. Six technical
1181 replicates per concentration. Error bars represent standard deviation. Experiment repeated two
1182 times.
1183
1184

1185
Supplementary Figure S7, related to Figure 7.
(A, B) GSEA results showing altered interferon signalling and methylation signatures upon MI-3
treatment compared to DMSO control.
(C) Volcano plot showing expression of repeat elements in MI-3 treated cells versus control; points
in red represent statistically significant changes (q value < 0.01). Three biological replicates per
condition.
(D) Heatmap showing differential expression of repeat families between vehicle and MI-3 treated
cells. Three biological replicates per condition.
1194
1195