SARS-CoV-2 causes brain inflammation and induces Lewy body formation in macaques

Ingrid H.C.H.M. Philippens[1,*†], Kinga P. Böszörményi[1†], Jacqueline A. Wubben[1], Zahra C.

Fagrouch[1], Nikki van Driel[1], Amber Q. Mayenburg[1], Diana Lozovagia[1], Eva Roos[2], Bernadette

Schurink[2], Marianna Bugiani[2], Ronald E. Bontrop[1,3], Jinte Middeldorp[1], Willy M. Bogers[1], Lioe-Fee

de Geus-Oei[5,6], Jan A.M. Langermans[1,4], Marieke A. Stammes[1‡], Babs E. Verstrepen[1‡], Ernst J.

Verschoor[1‡]

[1] Biomedical Primate Research Centre (BPRC), Lange Kleiweg 161, 2288GJ Rijswijk, The

Netherlands

[2] Department of Pathology, Amsterdam UMC, De Boelelaan 1117, 1081HV Amsterdam, The

Netherlands

[3] Department of Biology, Theoretical Biology and Bioinformatics, Utrecht University, Padualaan 8,

Utrecht, The Netherlands

[4] Department Population Health Sciences, Division of Animals in Science and Society, Faculty of

Veterinary Medicine, Utrecht University, Yalelaan 1, 3584CL Utrecht, The Netherlands

[5] Department of Radiology, Leiden University Medical Center, Albinusdreef 2, 2333ZA Leiden, The

Netherlands

[6] Biomedical Photonic Imaging Group, University of Twente, Drienerlolaan 5, 7522ND Enschede,

The Netherlands

† These authors contributed equally

‡ These authors jointly supervised this work

* Corresponding author. E-mail: philippens@bprc.nl

Keywords: SARS-CoV-2; macaques; brain inflammation; Lewy body formation; neuropathology;

PET-CT

Abstract:

SARS-CoV-2 may cause acute respiratory disease, but the infection can also initiate neurological symptoms. Here we show that SARS-CoV-2 infection causes brain inflammation in the macaque model. An increased metabolic activity in the pituitary gland of two macaques was observed by longitudinal positron emission tomography-computed tomography (PET-CT). Post-mortem analysis demonstrated infiltration of T-cells and activated microglia in the brain, and viral RNA was detected in brain tissues from one animal. We observed Lewy bodies in brains of all rhesus macaques. These data emphasize the virus' capability to induce neuropathology in this nonhuman primate model for SARS-CoV-2 infection. As in humans, Lewy body formation is an indication for the development of Parkinson's disease, this data represents a warning for potential long-term neurological effects after SARS-CoV-2 infection.

Teaser:

SARS-CoV-2 causes brain inflammation and Lewy bodies, a hallmark for Parkinson, after an asymptomatic infection in macaques.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) causes a multi-system inflammatory disease syndrome, COVID-19 (*1*). Although SARS-CoV-2 predominantly affects the respiratory organs, over 30% of the hospitalized COVID-19 patients also suffer from neurological manifestations, including loss of smell or taste, delirium, diminished consciousness, epilepsy, and psychosis (*2-5*). Besides these general neurological symptoms, some patients additionally endure Parkinsonism (*6-8*). The mechanisms behind this process are poorly understood. Neurological symptoms may be triggered by infection of the brain tissue, or indirectly, via virus-induced immune cell activation (*9*). In humans, a direct link between brain inflammation and the presence of SARS-CoV-2 RNA has not been established yet (*10*), and thus, formal proof that central nervous system (CNS)-related symptoms of COVID-19 are directly caused by the infection, or indirectly due to overactivation of the immune system, is lacking. Additionally, the long-term effects on the CNS after a mild to moderate SARS-CoV-2 infection, likely the vast majority of human cases, are unknown, and post-mortem brain samples from these individuals are not expected to become available for research in the near future. Controlled infection studies in a standardized experimental setting are crucial to investigate SARS-CoV-2-induced brain pathology (*11*). To address this issue, a study was performed in two macaques species, rhesus macaques (*Macaca mulatta*) and cynomolgus macaques (*Macaca fascicularis)* (Table 1), both well-accepted animal models for COVID-19 (*12*). Four male rhesus and four male cynomolgus macaques were inoculated with 10^5 TCID$_{50}$ of SARS-CoV-2 strain BetaCoV/BavPat1/2020 via a combined intratracheal and intranasal route (*13, 14*). Following infection, SARS-CoV-2 genomic material was

detected in tracheal and nasal swabs up to ten days, and based on clinical signs and thorax CTs,

all animals showed mild to moderate disease symptoms (*13, 15*).

Weekly [18]F-FDG PET-CTs of the brains of all macaques were initiated at the time point that the

virus became undetectable in nasal and tracheal swabs. The uptake of tracer renders a marker

for metabolic activity. Two of four cynomolgus macaques (C1 and C2) displayed an increased

uptake of [18]F-FDG in the pituitary gland at multiple time points. In animal C1, an increased uptake

of [18]F-FDG in the pituitary gland was seen at days 30 and 36 post-infection (Fig. 1). In animal C2,

increased metabolic activity was already visible on day 8 and continued through day 35 when the

final scan was obtained (Table S1).

In humans and macaques, the volume of the pituitary gland is small, and under physiological

conditions, its metabolic activity is comparable to the background level of the entire brain (*16,*

17). The [18]F-FDG uptake of the pituitary gland may even be underestimated due to the partial-

volume-effects that affect the emission signal recovery (*16*). Because pituitary gland tissue

expresses angiotensin-converting enzyme 2 (ACE2) (*18*), the increased [18]F-FDG uptake may be a

direct effect of the infection or an indirect effect due to either a (reversible) hypophysitis, or

transient hypothalamic-pituitary dysfunction (*19*). Hypocortisolism has been reported as a

delayed complication of SARS and has also been described in a SARS-CoV-2 patient (*20*). For

animal C1, which showed increased [18]F-FDG uptake 30 days after infection, it is likely that the

hypothalamic-pituitary axis was activated, leading to hypocortisolism similar to what has been

found in patients with both a SARS and SARS-CoV-2 infection.

To further investigate the consequences of SARS-CoV-2-infection on macaque brain tissue, the

animals were euthanized 5-6 weeks after experimental infection. Sections of the whole brain

were systematically collected for further examination. As several regions of the brain express the

SARS-CoV-2 receptor ACE2, and inflammation was found in the human brain (*10, 21, 22*), we used

various immunological markers for innate and adaptive immune activation to investigate for signs

of immune activation, and also explored the localization of virus particles (Fig. 2 and Table 2).

Viral RNA was detected by real-time quantitative polymerase chain reaction (RT-qPCR) in multiple

brain areas of the right hemisphere of cynomolgus macaque C3 (Fig. 2). More precisely,

cerebellum (1.48×10^5 RNA genome equivalents (GE)/gram), medial motor cortex (2.09×10^5

GE/gram), sensory cortex (2.07×10^5 GE/gram) and frontal basal cortex (8.29×10^4 GE/gram), as well

as hippocampus (1.24×10^5 GE/gram), hypothalamus (1.05×10^6 GE/gram), and globus pallidus

(5.45×10^4 GE/gram) all tested positive in the RT-qPCR. No viral RNA was detected in samples

collected from the pituitary gland or olfactory bulb, substantia nigra, medulla oblongata, pons,

nucleus caudatus, and putamen. Of interest, other tissues collected from macaque C3, including

tracheobronchial lymph nodes, heart, liver, spleen, and kidney, also tested positive in the RT-

qPCR, with comparable (lymph nodes, heart), or lower (kidney, liver, spleen) viral RNA loads (*13*).

Subgenomic messenger RNA analysis did not show evidence for active virus replication in the

brains at the time point of euthanasia. Additionally, SARS-CoV-2 antigen was not detectable by

immunohistochemistry the brains of all macaques.

The brains of all SARS-CoV-2-infected macaques showed evidence of inflammation. Presence of

T-cells was visualized by CD3 staining in intraparenchymal brain tissue, suggesting the infiltration

of T-cells that passed the blood-brain barrier after SARS-CoV-2 infection (Fig. 3A, top panel).

Additionally, activation of microglia cells in different areas of the brain, including the olfactory

bulb and pituitary gland, was confirmed by Mamu-DR staining (upregulation of MHC class II

expression) (Fig. 3A, middle panel). However, nodule formation, which is a measure for severity

of activation, was rarely present (Table 2). No B-lymphocytic infiltration was found as evidenced

by lack of CD20 staining (not shown). Hematoxylin and eosin (HE) staining did not show any

abnormalities in the brain tissue of the virus-exposed macaques, including the absence of

ischemic/necrotic lesions. For comparison, post-mortem brain tissues from two healthy, age-

matched macaques of each species were used as controls (Table 1), none of the four control

animals displayed obvious signs of immune activation (T-cells and microglia).

Brain tissues were screened for α-synuclein deposits, known as Lewy bodies, by

immunohistochemistry. In humans, Lewy body formation is linked to the development of

Parkinson's disease or Lewy body dementia (*23, 24*). It has been hypothesized that certain

neurotropic viruses, including MERS and SARS coronaviruses, can trigger formation of Lewy

bodies and cause Parkinsonism (*25-30*).

The formation of intracellular Lewy bodies was clearly shown in the ventral midbrain region next

to the caudate nucleus of all infected rhesus macaques (Table 2, Fig. 3B), and in one aged

cynomolgus macaque (C4), while Lewy bodies were absent in the brains of all four control

animals. For cynomolgus macaque C4, an age-dependent factor related to Parkinsonism cannot

be excluded as this animal was older (16 years) than the other panel members (5-7 years), but

the data from the rhesus macaques provide clear evidence for SARS-CoV-2-driven inflammation

in the brain of macaques. In humans, neuropathology has been described in moribund COVID-19

patients, but we report of SARS-CoV-2-related brain involvement in macaques without displaying

overt clinical signs. In general, macaque models for SARS-CoV-2 infection typically represent mild

to moderate COVID-19 symptoms on the CT scan compared to humans (*12, 14, 31*). Detection of

viral RNA in the brain of an animal demonstrates the virus' neuroinvasive capability. This matches

a recent study describing neuroinvasion in mouse brains and in human brain organoids (*32*).

How exactly SARS-CoV-2 caused widespread brain inflammation and induced Lewy body

formation remains unknown. Viruses can enter the brain via different pathways. In this study,

infiltration of T-cells was found perivascular and in the brain parenchyma, which indicates that

the blood-brain barrier integrity may have been disturbed, offering the virus the opportunity to

enter the brain. Alternatively, we hypothesize that SARS-CoV-2 gained access to the brain via

neuronal pathways, such as the retrograde and anterograde neuronal transport through infected

motor or sensory neurons (*33*), and entered the pituitary gland via binding to the ACE2 receptor

protein expressed on its cell surfaces (*18, 34*). Such neural connection also exists between the

olfactory bulb and the nasal mucosa (*35*), and the loss of taste and smell, a characteristic of

COVID-19, can thus be explained by nasal infection and subsequent inflammation in the olfactory

bulb. Such a scenario is in line with the finding that in all the SARS-CoV-2-exposed macaques

immune system activation in the olfactory bulb was evidenced by the presence of T-cells and/or

activated microglia.

Neuronal transport can also explain why some COVID-19 patients develop Parkinson's disease-

like symptoms. Viruses can also, via retrograde transport in parasympathic motor neurons of the

nervus vagus to the medulla, pons, and midbrain, reach the substantia nigra in the midbrain (*27,*

36). Notably, the α-synuclein inclusions were found in the ventral midbrain region of the animals.

In humans, these inclusions of accumulated misfolded proteins are associated with Parkinson's

disease or Lewy body dementia (*37*).

There is a growing concern that symptomatic COVID-19 patients may suffer from long-term

consequences (*9, 38*). In this light the finding of Lewy bodies in brains of infected macaques

without overt clinical signs is intriguing. <mark>Together with signs of inflammation and immune

activation in the brains of the macaques this finding may point to a not yet earlier described SARS-

CoV-2-induced neurodegenerative process that can explain the neurological symptoms that

COVID-19 survivors experience (*39*).</mark>

Lewy bodies are considered a hallmark for the development of Parkinson's disease, or Lewy body

dementia. More confirmation is required, but the observations in the translational macaque

models for COVID-19 (*12-14, 40, 41*) can be regarded as a serious warning as they may be

predictive for COVID-19-related dementia cases in humans in the future, even after an

asymptomatic infection or mild disease process.

Materials and Methods

Animals and SARS-CoV-2 exposure

Four cynomolgus macaques (*Macaca fascicularis*) and four Indian-origin rhesus macaques

(*Macaca mulatta*) (Table 1) were selected for this study. All macaques were mature, outbred

animals, purpose-bred and socially housed at the BPRC. The animals were in good physical health

with normal baseline biochemical and hematological values. All were pair-housed with a socially

compatible cage-mate. The animals were offered a daily diet consisting of monkey food pellets

(Ssniff, Soest, Germany) supplemented with vegetables and fruit. Enrichment was provided daily

in the form of pieces of wood, mirrors, food puzzles, and a variety of other homemade or

commercially available enrichment products. Drinking water was available *ad libitum* via an

automatic system. Animal Care staff provided daily visual health checks before infection, and

twice-daily after infection. The animals were monitored for appetite, general behavior, and stool

consistency. All possible precautions were taken to ensure the welfare and to avoid any

discomfort to the animals. All experimental interventions (intratracheal and intranasal infection,

swabs, blood samplings, and PET-CTs) were performed under anesthesia. The research protocol

was approved by national authorities (CCD, Central Committee for Animal Experiments; license

number AVD5020020209404). Approval to start was obtained after further assessment of the

detailed study protocol by the institutional animal welfare body (AWB) (in Dutch: Instantie voor

Dierenwelzijn, IvD). The BPRC is accredited by the American Association for Accreditation of

Laboratory Animal Care (AAALAC) International and is compliant with European directive

2010/63/EU as well as the "Standard for Humane Care and Use of Laboratory Animals by Foreign

Institutions" provided by the Department of Health and Human Services of the US National

Institutes of Health (NIH, identification number A5539-01).

On day 0, all animals were exposed to a dose of 10^5 TCID$_{50}$ of SARS-CoV-2 (strain

BetaCOV/BavPat1/2020), diluted in 5 ml phosphate-buffered saline (PBS). The virus was

inoculated via a combination of the intratracheal route (4.5 ml) and intranasal route (0.25 ml per

nostril). For the histological examination brains from naive control macaques from the same age

were obtained from the BPRC biobank, two cynomolgus, and two rhesus macaques.

Positron Emission Tomography – Computed Tomography

Positron Emission Tomography (PET)-computed tomography (CT) data were acquired on multiple

time points post-infection using a MultiScan Large Field of View Extreme Resolution Research

Imager (LFER) 150 PET-CT (Mediso Medical Imaging Systems Ltd., Budapest, Hungary) as

described before (*42*). Animals were fasted overnight (glucose level > 8.5 mmol/l). The animals

were sedated with ketamine (10 mg/kg ketamine hydrochloride (Alfasan Nederland BV, Woerden,

The Netherlands)) combined with medetomidine hydrochloride (0.05 mg/kg (Sedastart; AST

Farma B.V., Oudewater, The Netherlands)) to induce sedation and muscle relaxation, both applied

intramuscularly (IM). The animals were positioned head first supine (HFS) with the arms up. The

scans were acquired under mechanical ventilation in combination with a forced breathing

pattern. For anesthetic maintenance, a minimum alveolar concentration of isoflurane (iso-MAC)

of around 0,80%-1.00% was used. The body temperature of the animal was maintained by using

the Bair Hugger (3M™, St Paul, MN, USA) supplied with 43˚C airflow. Typically, around 100 MBq

of ^{18}F-FDG was applied intravenously (GE Healthcare, Leiderdorp, NL). Thirty minutes after

injection the plateau in tracer activity uptake is reached, subsequently a PET of 15 minutes was

acquired. After the scan, upon return to their home cage, atipamezole hydrochloride (Sedastop,

ASTFarma B.V., Oudewater, NL, 5 mg/ml, 0.25 mg/kg) was administrated IM to antagonize

medetomidine. Afterward the emission data was iteratively reconstructed (OSEM3D, 8 iterations,

and 9 subsets) into a single frame PET image normalized and corrected for attenuation, scatter,

and random coincidences using a reference CT and corrected for radioactive decay. The analysis

was performed in VivoQuant 4.5 (Invicro, Boston, USA). Based on repeatability parameters for

correct interpretation of the results, a standardized uptake value (SUV) ratio was used for

robustness (*17, 42*). An increased uptake, and pituitary gland hypermetabolism is defined as a

SUV$_{mean}$ ratio above 1.5 for the pituitary gland over the surrounding brain in combination with a

SUV$_{peak}$ ratio above background levels (>1.0). A group of non-infected control rhesus macaques

(n=6) were used to calculate average background uptake of ^{18}F-FDG.

Brain tissue collection

Five weeks after virus exposure the macaques were euthanized and the brains were collected for

further examination. The right hemisphere was used for RT-qPCR analysis and the left hemisphere

was fixed in formalin for histology. Fifteen different regions were collected from the right

hemisphere for RT-qPCR analysis: 1) part of the pituitary gland, 2) the olfactory bulb, 3) substantia

nigra, 4) medulla oblongata, 5) pons, 6) anterior part of the cerebellum, 7) motor cortex medial,

8) sensory cortex, 9) frontal basal cortex, 10) hippocampus, 11) caudate nucleus, 12)

hypothalamus, 13) globus pallidus, 14) putamen, and 15) thalamus. For the preparation of

paraffin-embedded sections of the formalin-fixed left hemisphere, the cerebrum and cerebellum

were dissected in 3-4 mm parts on the anterior-posterior axis. Pituitary gland and olfactory bulb

were also embedded. From each part, sections (4 μm) were prepared for different staining

methods. Immunohistochemistry stains were used for T-cells (CD3), B-cells (CD20), activated

microglia (Mamu-DR), Lewy bodies (α-synuclein ab), and for SARS-CoV-2. Hematoxyline-eosine

(HE) staining was used for general morphology.

Viral RNA detection in brain tissue

Brain tissue samples were weighed and placed in gentleMACS M tubes (30 mg in 100 μl PBS) and

treated using a gentleMACS Tissue Dissociator (protein01 program)(Miltenyi Biotec B.V., Leiden,

The Netherlands). Next, the homogenized tissue was centrifuged, and 100 µl supernatant was

used for RNA isolation. Viral RNA was isolated from using a QIAamp Viral RNA Mini kit (Qiagen

Benelux BV, Venlo, The Netherlands) following the manufacturer's instructions. Viral RNA was

reverse-transcribed to cDNA using a Transcriptor First Strand cDNA Synthesis kit (Roche

Diagnostics BV, Almere, The Netherlands). Viral RNA was quantified by RT-qPCR specific for RdRp

gene of SARS-CoV-2, as described by Corman *et al.* (*43*). The lower detection limit of the RT-qPCR

was 3.6 viral RNA copies per reaction. Viral sub-genomic RNA was detected essentially as

described by Wölfel *et al.* (*44*). For both assays, RNA standard curves were generated by *in vitro*

transcription of the target regions.

Tissue preparation for histology

The left hemisphere of the brains, part of the pituitary gland, and one olfactory bulb were fixed

in formalin for 24 hours and thereafter stored in buffered PBS. Preserved brains were

cryoprotected in 30% w/v sucrose in PBS. The cerebrum was dissected in 12 different parts cut

anterior-posterior axis at +10, +8, +5, +1, -3, -6, -8, -11, -14, -18, -22 from Bregma (*45*), the

cerebellum and pons were cut in 4 parts. These part were embedded in paraffin. From the eight

brain parts in which viral RNA was detected by RT-qPCR, strips of brain sections were sliced into

12-series of 4 µm sections for different stains. These parts included the frontal cortex, midbrain

parts, cerebellum, pituitary gland, and olfactory bulb. Sections were stained with virus antibody

staining for virus detection and immunohistochemistry for immune reaction such as T-cell

staining (CD3), B-cell staining (CD20), MHC-II cell staining (HLA-DR). Mirror sections were analyzed

with a HE staining for brain morphology.

Immunohistochemistry

The optimal concentration was determined for each antibody: CD3 (polyclonal rabbit – anti-

human CD3 IgG, cat. no. A045201-2, Agilent Technologies), 1:60; CD20 (monoclonal mouse – anti-

human CD20 IgG2a, clone L26, cat. no. M075501-2, Agilent Technologies), 1:800; Mamu-DR

(monoclonal mouse – anti-human HLA-DR/DQ- IgG1, clone CR3/43, cat. no. M077501-2, Agilent

Technologies), 1:150. For antigen retrieval, a steamer was used. Antigen Retrieval solution: IHC-

TEK epitope retrieval solution, ready to use (catno IW-1100, IHC world). All incubation steps were

at room temperature unless mentioned otherwise. Additionally, hematoxylin was used as a

counterstaining in all protocols. After a dehydration sequence, the slides were mounted in

Malinol. The counting of cells was performed in a blind matter.

CD3 and CD20 staining

The slides were deparaffinized by putting the slides sequentially in xylene, 100% ethanol, 96%

ethanol, 70% ethanol, and PBS. Subsequently, an epitope antigen retrieval was executed in a

steamer for 1h. After cooling down, the slides were placed in cuvettes (Sequenza cover plate

system productnr 36107 Ted Pella inc.). Endogenous peroxidase (PO) activity was blocked by the

PO blocking solution from DAKO (S2023) for 15 minutes. After a washing step (PBS with 0.05%

Tween) Avidin was added from the DAKO kit (X0590) for 10 minutes. Thereafter, another washing

step and biotin was added from the same DAKO kit (10 min) for blocking endogenous biotine.

After washing a blocking step was executed for 20 minutes (PBS with 0.1% BSA and 1% normal

human serum, NHS). The primary antibody was added (diluted in 0.1% BSA in PBS) and the slices

were left overnight at 4°C. After washing a secondary antibody (Rabbit-anti-mouse IgG

biotinylated (E0354), Agilent Technologies; 1:200 diluted in PBS + 1% BSA + 1% NHS) was added

and, after washing, the slides were incubated with Vectastain ABC-peroxidase (ABC-PO, from

Vector Laboratories; PK-4000; diluted 1:100 in PBS) for 30 minutes. After washing, 3,3'-

diaminobenzidine (DAB) with 0.02% H_2O_2 was added to visualize the antigen-antibody binding (20

min).

Mamu-DR staining

The EnVision™ staining kit (G|2 Double-stain System, Rabbit/Mouse, DAB+/Permanent RED code

K5361; Agilent technologies, Dako DK) was used for the immunohistochemical stain of Mamu-DR.

The slides were deparaffinized by putting them sequentially in xylene, 100% ethanol, 96%

ethanol, 70% ethanol, and PBS. Subsequently, an epitope antigen retrieval was executed and the

slides were put in a steamer for 1 hour. The cooled down slides were placed in cuvettes and the

endogenous peroxidase activity was blocked by the envision kit. The slides were washed and

thereafter a blocking step was used consisting of 1% NHS + 1% BSA + 0.2% triton x100 in PBS for

10 minutes. Subsequently, the primary antibody was added (diluted in 0.1% BSA/PBS) for 30

minutes. Thereafter a washing step was implemented and the EnVision™ polymer/HRP

(secondary antibody) was added for 10 minutes. Polymer HRP was added for 10 minutes followed

by a washing step. Thereafter DAB+ was added for 15 minutes to visualize the antigen-antibody

binding.

α-Synuclein staining

The slides were deparaffinized by immersing them sequentially in xylene, 100% ethanol, 96%

ethanol, 70% ethanol, and PBS. Subsequently, an epitope antigen retrieval was executed by

putting the slides for 15 minutes in Formic acid (100%) diluted 1:10 in demineralized water. After

2 washing steps in PBS with 0,05% Tween, the slides were placed in cuvettes (Sequenza cover

plate system product no. 36107 Ted Pella inc.). Endogenous PO activity was blocked by the PO

blocking solution from DAKO (S2023) for 20 minutes. After washing (PBS with 0.05% Tween),

avidin was added from the DAKO kit (X0590) for 10 minutes. Then, after another washing step,

biotin was added from the same DAKO kit (10 min) for blocking of endogenous biotine. After

washing, a blocking step was executed for 30 minutes (PBS with 0.1% BSA and 1% NHS and 0.02%

Triton-X100). The primary antibody, α-synuclein clone 4D6 (Biolegend SIG-39720), was added

(diluted in 0.1% BSA in PBS) and the slides were left overnight at 4°C. After washing, a secondary

antibody (rabbit anti-mouse IgG Biotinylated (E0354), Agilent Technologies; 1:200 diluted in PBS

+ 1% BSA) was added for 30 minutes and the slides were incubated with Vectastain ABC-PO kit

from Vector Laboratories (PK-4000; diluted 1:100 in PBS) for 30 minutes. After a final washing step,

DAB with 0.02% H_2O_2 was added to visualize the Antigen-antibody binding (20 min).

SARS-CoV-2 staining

The Roche Optiview DAB IHC kit was used in a Ventana Benchmark Ultra (Roche, Basel

Switzerland) immunostainer to immunohistochemically stain SARS-CoV-2. Two monoclonal

antibodies of ThermoFisher raised to SARS-CoV-2 Nucleocapsid (clone B46F, catno MA1-7404,

and E16C, catno. MA1-7403) were validated on formaldehyde-fixed and paraffin-embedded

SARS-CoV-2 and mock-infected Vero E6 cells (*10*), as well as lung tissue sections of human SARS-

CoV-2 patients. The clone E16C was superior to B46F and was further used in this study. Antigen

retrieval took place with cell conditioning 1 (CC1 , Ventana Medical Systems) (pH 8,5) for 24

minutes at 100°C, 1/5.000 diluted. Thereafter, incubation took place with the primary antibody

for 48 minutes at 36°C followed by standard Optiview detection/visualization with DAB and

Copper. After immunohistochemical staining, the sections were dehydrated with grades of

ethanol and cleared with xylene. All sections were mounted with TissueTek® coverslipping film

(Sakura Finetek Europe B.V., Alphen aan den Rijn, The Netherlands).

References

1.　V. Coronaviridae Study Group of the International Committee on Taxonomy of, The
species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV
and naming it SARS-CoV-2. *Nat Microbiol* **5**, 536-544 (2020).
2.　L. Mao, H. Jin, M. Wang, Y. Hu, S. Chen, Q. He, J. Chang, C. Hong, Y. Zhou, D. Wang, X.
Miao, Y. Li, B. Hu, Neurologic Manifestations of Hospitalized Patients With Coronavirus
Disease 2019 in Wuhan, China. *JAMA Neurol* **77**, 683-690 (2020).
3.　R. W. Paterson, R. L. Brown, L. Benjamin, R. Nortley, S. Wiethoff, T. Bharucha, D. L.
Jayaseelan, G. Kumar, R. E. Raftopoulos, L. Zambreanu, V. Vivekanandam, A. Khoo, R.
Geraldes, K. Chinthapalli, E. Boyd, H. Tuzlali, G. Price, G. Christofi, J. Morrow, P.
McNamara, B. McLoughlin, S. T. Lim, P. R. Mehta, V. Levee, S. Keddie, W. Yong, S. A. Trip,
A. J. M. Foulkes, G. Hotton, T. D. Miller, A. D. Everitt, C. Carswell, N. W. S. Davies, M.
Yoong, D. Attwell, J. Sreedharan, E. Silber, J. M. Schott, A. Chandratheva, R. J. Perry, R.
Simister, A. Checkley, N. Longley, S. F. Farmer, F. Carletti, C. Houlihan, M. Thom, M. P.
Lunn, J. Spillane, R. Howard, A. Vincent, D. J. Werring, C. Hoskote, H. R. Jager, H. Manji,
M. S. Zandi, The emerging spectrum of COVID-19 neurology: clinical, radiological and
laboratory findings. *Brain* **143**, 3104-3120 (2020).
4.　M. Fotuhi, A. Mian, S. Meysami, C. A. Raji, Neurobiology of COVID-19. *J Alzheimers Dis*
76, 3-19 (2020).
5.　A. Giacomelli, L. Pezzati, F. Conti, D. Bernacchia, M. Siano, L. Oreni, S. Rusconi, C.
Gervasoni, A. L. Ridolfo, G. Rizzardini, S. Antinori, M. Galli, Self-reported Olfactory and
Taste Disorders in Patients With Severe Acute Respiratory Coronavirus 2 Infection: A
Cross-sectional Study. *Clin Infect Dis* **71**, 889-890 (2020).
6.　M. E. Cohen, R. Eichel, B. Steiner-Birmanns, A. Janah, M. Ioshpa, R. Bar-Shalom, J. J. Paul,
H. Gaber, V. Skrahina, N. M. Bornstein, G. Yahalom, A case of probable Parkinson's
disease after SARS-CoV-2 infection. *Lancet Neurol* **19**, 804-805 (2020).

7.   I. Faber, P. R. P. Brandao, F. Menegatti, D. D. de Carvalho Bispo, F. B. Maluf, F. Cardoso,
Coronavirus Disease 2019 and Parkinsonism: A Non-post-encephalitic Case. *Mov Disord*
35, 1721-1722 (2020).
8.   A. Mendez-Guerrero, M. I. Laespada-Garcia, A. Gomez-Grande, M. Ruiz-Ortiz, V. A.
Blanco-Palmero, F. J. Azcarate-Diaz, P. Rabano-Suarez, E. Alvarez-Torres, C. P. de
Fuenmayor-Fernandez de la Hoz, D. Vega Perez, R. Rodriguez-Montalban, A. Perez-
Rivilla, J. Sayas Catalan, A. Ramos-Gonzalez, J. Gonzalez de la Aleja, Acute hypokinetic-
rigid syndrome following SARS-CoV-2 infection. *Neurology* **95**, e2109-e2118 (2020).
9.   M. T. Heneka, D. Golenbock, E. Latz, D. Morgan, R. Brown, Immediate and long-term
consequences of COVID-19 infections for the development of neurological disease.
Alzheimers Res Ther **12**, 69 (2020).
10. B. Schurink, E. Roos, T. Radonic, E. Barbe, C. S. C. Bouman, H. H. de Boer, G. J. de Bree, E.
B. Bulle, E. M. Aronica, S. Florquin, J. Fronczek, L. M. A. Heunks, M. D. de Jong, L. Guo, R.
du Long, R. Lutter, P. C. G. Molenaar, E. A. Neefjes-Borst, H. W. M. Niessen, C. J. M. van
Noesel, J. Roelofs, E. J. Snijder, E. C. Soer, J. Verheij, A. P. J. Vlaar, W. Vos, N. N. van der
Wel, A. C. van der Wal, P. van der Valk, M. Bugiani, Viral presence and immunopathology
in patients with lethal COVID-19: a prospective autopsy cohort study. *Lancet Microbe* **1**,
e290-e299 (2020).
11. S. Natoli, V. Oliveira, P. Calabresi, L. F. Maia, A. Pisani, Does SARS-Cov-2 invade the
brain? Translational lessons from animal models. *Eur J Neurol* **27**, 1764-1773 (2020).
12. C. Munoz-Fontela, W. E. Dowling, S. G. P. Funnell, P. S. Gsell, A. X. Riveros-Balta, R. A.
Albrecht, H. Andersen, R. S. Baric, M. W. Carroll, M. Cavaleri, C. Qin, I. Crozier, K.
Dallmeier, L. de Waal, E. de Wit, L. Delang, E. Dohm, W. P. Duprex, D. Falzarano, C. L.
Finch, M. B. Frieman, B. S. Graham, L. E. Gralinski, K. Guilfoyle, B. L. Haagmans, G. A.
Hamilton, A. L. Hartman, S. Herfst, S. J. F. Kaptein, W. B. Klimstra, I. Knezevic, P. R.
Krause, J. H. Kuhn, R. Le Grand, M. G. Lewis, W. C. Liu, P. Maisonnasse, A. K. McElroy, V.
Munster, N. Oreshkova, A. L. Rasmussen, J. Rocha-Pereira, B. Rockx, E. Rodriguez, T. F.
Rogers, F. J. Salguero, M. Schotsaert, K. J. Stittelaar, H. J. Thibaut, C. T. Tseng, J. Vergara-
Alert, M. Beer, T. Brasel, J. F. W. Chan, A. Garcia-Sastre, J. Neyts, S. Perlman, D. S. Reed,
J. A. Richt, C. J. Roy, J. Segales, S. S. Vasan, A. M. Henao-Restrepo, D. H. Barouch, Animal
models for COVID-19. *Nature* **586**, 509-515 (2020).
13. K. P. Böszörményi, M. A. Stammes, Z. C. Fagrouch, G. Kiemenyi-Kayere, H. Niphuis, D.
Mortier, N. van Driel, I. Nieuwenhuis, E. Zuiderwijk-Sick, L. Meijer, P. Mooij, E. J.
Remarque, G. Koopman, A. C. R. Hoste, P. Sastre, B. L. Haagmans, R. E. Bontrop, J. A. M.
Langermans, W. M. Bogers, E. J. Verschoor, B. E. Verstrepen, Comparison of SARS-CoV-2
infection in two non-human primate species: rhesus and cynomolgus macaques. *bioRxiv*,
11.05.369413 (2020).
14. B. Rockx, T. Kuiken, S. Herfst, T. Bestebroer, M. M. Lamers, B. B. Oude Munnink, D. de
Meulder, G. van Amerongen, J. van den Brand, N. M. A. Okba, D. Schipper, P. van Run, L.
Leijten, R. Sikkema, E. Verschoor, B. Verstrepen, W. Bogers, J. Langermans, C. Drosten,
M. Fentener van Vlissingen, R. Fouchier, R. de Swart, M. Koopmans, B. L. Haagmans,
Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate model.
Science **368**, 1012-1015 (2020).

15.   F. Pan, T. Ye, P. Sun, S. Gui, B. Liang, L. Li, D. Zheng, J. Wang, R. L. Hesketh, L. Yang, C.
412   Zheng, Time Course of Lung Changes at Chest CT during Recovery from Coronavirus
413   Disease 2019 (COVID-19). *Radiology* **295**, 715-721 (2020).

414   16.   M. Meyer, G. Allenbach, M. Nicod Lalonde, N. Schaefer, J. O. Prior, S. Gnesin, Increased
415   (18)F-FDG signal recovery from small physiological structures in digital PET/CT and
416   application to the pituitary gland. *Sci Rep* **10**, 368 (2020).

417   17.   S. Goutal, N. Tournier, M. Guillermier, N. Van Camp, O. Barret, M. Gaudin, M.
418   Bottlaender, P. Hantraye, S. Lavisse, Comparative test-retest variability of outcome
419   parameters derived from brain [18F]FDG PET studies in non-human primates. *PLoS One*
420   **15**, e0240228 (2020).

421   18.   R. Pal, M. Banerjee, COVID-19 and the endocrine system: exploring the unexplored. *J*
422   *Endocrinol Invest* **43**, 1027-1031 (2020).

423   19.   R. Pal, COVID-19, hypothalamo-pituitary-adrenal axis and clinical implications. *Endocrine*
424   **68**, 251-252 (2020).

425   20.   M. W. J. Chua, M. P. W. Chua, Delayed Onset of Central Hypocortisolism in a Patient
426   Recovering From COVID-19. *AACE Clin Case Rep* **7**, 2-5 (2021).

427   21.   I. Hamming, W. Timens, M. L. Bulthuis, A. T. Lely, G. Navis, H. van Goor, Tissue
428   distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in
429   understanding SARS pathogenesis. *J Pathol* **203**, 631-637 (2004).

430   22.   M. Hoffmann, H. Kleine-Weber, S. Schroeder, N. Kruger, T. Herrler, S. Erichsen, T. S.
431   Schiergens, G. Herrler, N. H. Wu, A. Nitsche, M. A. Muller, C. Drosten, S. Pohlmann, SARS-
432   CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven
433   Protease Inhibitor. *Cell* **181**, 271-280 e278 (2020).

434   23.   Z. Walker, K. L. Possin, B. F. Boeve, D. Aarsland, Lewy body dementias. *Lancet* **386**, 1683-
435   1697 (2015).

436   24.   M. Sezgin, B. Bilgic, S. Tinaz, M. Emre, Parkinson's Disease Dementia and Lewy Body
437   Disease. *Semin Neurol* **39**, 274-282 (2019).

438   25.   T. Yamada, Viral etiology of Parkinson's disease: Focus on influenza A virus. *Parkinsonism*
439   *Relat Disord* **2**, 113-121 (1996).

440   26.   R. L. Doty, The olfactory vector hypothesis of neurodegenerative disease: is it viable?
441   *Ann Neurol* **63**, 7-15 (2008).

442   27.   I. Mori, Viremic attack explains the dual-hit theory of Parkinson's disease. *Med*
443   *Hypotheses* **101**, 33-36 (2017).

444   28.   E. M. Gatto, J. Fernandez Boccazzi, COVID-19 and neurodegeneration: what can we learn
445   from the past? *Eur J Neurol* **27**, e45 (2020).

446   29.   A. Y. Vittor, M. Long, P. Chakrabarty, L. Aycock, V. Kollu, S. T. DeKosky, West Nile Virus-
447   Induced Neurologic Sequelae-Relationship to Neurodegenerative Cascades and
448   Dementias. *Curr Trop Med Rep* **7**, 25-36 (2020).

449   30.   S. Ait Wahmane, A. Achbani, Z. Ouhaz, M. Elatiqi, A. Belmouden, M. Nejmeddine, The
450   Possible Protective Role of alpha-Synuclein Against Severe Acute Respiratory Syndrome
451   Coronavirus 2 Infections in Patients With Parkinson's Disease. *Mov Disord* **35**, 1293-1294
452   (2020).

453   31.   D. K. Singh, B. Singh, S. R. Ganatra, M. Gazi, J. Cole, R. Thippeshappa, K. J. Alfson, E.
454   Clemmons, O. Gonzalez, R. Escobedo, T. H. Lee, A. Chatterjee, Y. Goez-Gazi, R. Sharan,

M. Gough, C. Alvarez, A. Blakley, J. Ferdin, C. Bartley, H. Staples, L. Parodi, J. Callery, A. Mannino, B. Klaffke, P. Escareno, R. N. Platt, 2nd, V. Hodara, J. Scordo, S. Gautam, A. G. Vilanova, A. Olmo-Fontanez, A. Schami, A. Oyejide, D. K. Ajithdoss, R. Copin, A. Baum, C. Kyratsous, X. Alvarez, M. Ahmed, B. Rosa, A. Goodroe, J. Dutton, S. Hall-Ursone, P. A. Frost, A. K. Voges, C. N. Ross, K. Sayers, C. Chen, C. Hallam, S. A. Khader, M. Mitreva, T. J. C. Anderson, L. Martinez-Sobrido, J. L. Patterson, J. Turner, J. B. Torrelles, E. J. Dick, Jr., K. Brasky, L. S. Schlesinger, L. D. Giavedoni, R. Carrion, Jr., D. Kaushal, Responses to acute infection with SARS-CoV-2 in the lungs of rhesus macaques, baboons and marmosets. *Nat Microbiol* **6**, 73-86 (2021).

32. E. Song, C. Zhang, B. Israelow, A. Lu-Culligan, A. V. Prado, S. Skriabine, P. Lu, O. E. Weizman, F. Liu, Y. Dai, K. Szigeti-Buck, Y. Yasumoto, G. Wang, C. Castaldi, J. Heltke, E. Ng, J. Wheeler, M. M. Alfajaro, E. Levavasseur, B. Fontes, N. G. Ravindra, D. Van Dijk, S. Mane, M. Gunel, A. Ring, S. A. J. Kazmi, K. Zhang, C. B. Wilen, T. L. Horvath, I. Plu, S. Haik, J. L. Thomas, A. Louvi, S. F. Farhadian, A. Huttner, D. Seilhean, N. Renier, K. Bilguvar, A. Iwasaki, Neuroinvasion of SARS-CoV-2 in human and mouse brain. *J Exp Med* **218**, (2021).

33. P. A. Swanson, 2nd, D. B. McGavern, Viral diseases of the central nervous system. *Curr Opin Virol* **11**, 44-54 (2015).

34. E. Lazartigues, M. M. F. Qadir, F. Mauvais-Jarvis, Endocrine Significance of SARS-CoV-2's Reliance on ACE2. *Endocrinology* **161**, bqaa108 (2020).

35. J. Meinhardt, J. Radke, C. Dittmayer, J. Franz, C. Thomas, R. Mothes, M. Laue, J. Schneider, S. Brünink, S. Greuel, M. Lehmann, O. Hassan, T. Aschman, E. Schumann, R. L. Chua, C. Conrad, R. Eils, W. Stenzel, M. Windgassen, L. Rößler, H.-H. Goebel, H. R. Gelderblom, H. Martin, A. Nitsche, W. J. Schulz-Schaeffer, S. Hakroush, M. S. Winkler, B. Tampe, F. Scheibe, P. Körtvélyessy, D. Reinhold, B. Siegmund, A. A. Kühl, S. Elezkurtaj, D. Horst, L. Oesterhelweg, M. Tsokos, B. Ingold-Heppner, C. Stadelmann, C. Drosten, V. M. Corman, H. Radbruch, F. L. Heppner, Olfactory transmucosal SARS-CoV-2 invasion as a port of central nervous system entry in individuals with COVID-19. *Nature Neuroscience* **24**, 168-175 (2021).

36. C. H. Hawkes, K. Del Tredici, H. Braak, Parkinson's disease: a dual-hit hypothesis. *Neuropathol Appl Neurobiol* **33**, 599-614 (2007).

37. J. P. Taylor, J. Hardy, K. H. Fischbeck, Toxic proteins in neurodegenerative disease. *Science* **296**, 1991-1995 (2002).

38. M. Marshall, The lasting misery of coronavirus long-haulers. *Nature* **585**, 339-341 (2020).

39. J. P. Rogers, A. S. David, A longer look at COVID-19 and neuropsychiatric outcomes. *Lancet Psychiatry* **8**, 351-352 (2021).

40. H. Zheng, H. Li, L. Guo, Y. Liang, J. Li, X. Wang, Y. Hu, L. Wang, Y. Liao, F. Yang, Y. Li, S. Fan, D. Li, P. Cui, Q. Wang, H. Shi, Y. Chen, Z. Yang, J. Yang, D. Shen, W. Cun, X. Zhou, X. Dong, Y. Wang, Y. Chen, Q. Dai, W. Jin, Z. He, Q. Li, L. Liu, Virulence and pathogenesis of SARS-CoV-2 infection in rhesus macaques: A nonhuman primate model of COVID-19 progression. *PLoS Pathog* **16**, e1008949 (2020).

41. V. J. Munster, F. Feldmann, B. N. Williamson, N. van Doremalen, L. Perez-Perez, J. Schulz, K. Meade-White, A. Okumura, J. Callison, B. Brumbaugh, V. A. Avanzato, R. Rosenke, P.

W. Hanley, G. Saturday, D. Scott, E. R. Fischer, E. de Wit, Respiratory disease in rhesus
macaques inoculated with SARS-CoV-2. *Nature* **585**, 268-272 (2020).

42. M. A. Stammes, J. Bakker, R. A. W. Vervenne, D. G. M. Zijlmans, L. van Geest, M. P. M.
Vierboom, J. A. M. Langermans, F. A. W. Verreck, Recommendations for Standardizing
Thorax PET–CT in Non-Human Primates by Recent Experience from Macaque Studies.
Animals **11**, 204 (2021).

43. V. M. Corman, O. Landt, M. Kaiser, R. Molenkamp, A. Meijer, D. K. Chu, T. Bleicker, S.
Brunink, J. Schneider, M. L. Schmidt, D. G. Mulders, B. L. Haagmans, B. van der Veer, S.
van den Brink, L. Wijsman, G. Goderski, J. L. Romette, J. Ellis, M. Zambon, M. Peiris, H.
Goossens, C. Reusken, M. P. Koopmans, C. Drosten, Detection of 2019 novel coronavirus
(2019-nCoV) by real-time RT-PCR. *Euro Surveill* **25**, (2020).

44. R. Wolfel, V. M. Corman, W. Guggemos, M. Seilmaier, S. Zange, M. A. Muller, D.
Niemeyer, T. C. Jones, P. Vollmar, C. Rothe, M. Hoelscher, T. Bleicker, S. Brunink, J.
Schneider, R. Ehmann, K. Zwirglmaier, C. Drosten, C. Wendtner, Virological assessment
of hospitalized patients with COVID-2019. *Nature* **581**, 465-469 (2020).

45. G. Paxinos, X. F. Huang, M. Petrides, A. Toga, *The Rhesus Monkey Brain: in Stereotaxic*
Coordinates. (Elsevier Science Publishing Co Inc, San Diego, Ca., ed. 2nd, 2008), pp. 416.
515

Acknowledgements

We want to thank Francisca van Hassel for assistance with editing of figures, Yolanda Kap

for reviewing the manuscript, Wim Vos for the excellent histology, and the Animal Science

Department of the BPRC, the veterinarians and animal caretakers for all the experimental

support.

Funding

This study was supported by funding from the Biomedical Primate Research Centre. KPB

was supported by the European Union's Marie Skłodowska-Curie Innovative Training

Network HONOURs; grant agreement no. 721367.

Author contributions

Conceptualization: IHCHMP, EJV, MAS, BEV

Data curation: MAS, BEV, IHCHMP

Formal Analysis: MAS, IHCHMP, BEV

Funding acquisition: EJV

Investigation: KPB, JAW, ZCF, NvD, AQM, DL

533 Methodology: MAS, MB, BS, IHCHMP

534 Project administration: EJV, BEV

535 Resources: REB

536 Supervision: IHCHMP, MAS, BEV, EJV

537 Validation: MB, ER, L-FG-O,

538 Writing – original draft: IHCHMP, KPB

539 Writing – review & editing: MAS, EJV, BEV, JAML, REB, JM, WMB, L-FG-O

541 **Competing interests:**

542 The authors declare that they have no competing interests.

544 **Data and materials availability:**

545 All data needed to evaluate the conclusions in the paper are present in the paper and/or

546 the Supplementary Materials. Correspondence should be addressed to IHCHMP

547 (philippens@bprc.nl).

A

B

Fig. 1. ¹⁸F-FDG PET-CT image of a SARS-CoV-2 infected cynomolgus macaque (C1).

Representative images of cynomolgus macaque C1 **(A)** on day 29, and a healthy control animal

(B) are shown. The pituitary gland is indicated by the cross-hairs in all three directions and is

boxed (right pictures). Similar window-level settings are applied for all sections. The pituitary

gland-brain ratio of animal C1 was calculated with both the mean and peak standard uptake value

(SUV); for the SUV$_{mean}$ this ratio was 1.9, and for the SUV$_{peak}$ 1.3. The average values calculated

from a group of non-infected control animals (n=6) were 1.1 (std 0.3) for the SUV$_{mean}$, and 0.6 (std

0.1) for the SUV$_{peak}$ ratio. The uptake values of the brain, defined as background, were similar

with SUV$_{mean}$ values of 2.2 and 1.9 for the SARS-CoV-2-infected animals and the non-infected

controls, respectively. After the relevant corrections for attenuation, scatter and decay, our

results are indeed indicative for pituitary hypermetabolism after SARS-CoV-2 infection in animals

C1 and C2.

1. pituitary gland
2. olfactory bulb
3. substantia nigra
4. medulla oblongata
5. pons
6. cerebellum
7. motor cortex
8. sensory cortex
9. frontal basal cortex
10. hippocampus
11. caudate nucleus
12. hypothalamus
13. globus pallidus
14. putamen
15. thalamus

Fig. 2. Overview of CNS effects by SARS-CoV-2 exposure in a macaque brain. Presence of viral

RNA was investigated in multiple regions of the brain as indicated by the numbers. Viral RNA-

positive regions in cynomolgus macaque C3 are indicated by a yellow background. The analysed

brain regions are indicated with a number. Brain areas with T-cells (CD3+) and activated microglia

(Mamu-DR+) are shown in light blue (mild expression) and dark blue (moderate expression),

respectively. Brain areas with Lewy bodies (α-synuclein+) are shown in orange.

Fig. 3. SARS-CoV-2 causes brain inflammation and Lewy body formationin brains of macaques.

Immunohistochemical stainings of macaque brain tissues. **(A)** top panel: T-cells (CD3+) in the

pituitary gland (left), perivascular (middle) and in the brain parenchyma (right) from a rhesus

macaque (20x). **(A)** middle panel: activated microglia (Mamu-DR+) in the pituitary gland (left),

olfactory bulb (middle) and in the brain parenchyma (right) from a cynomolgus macaque (20x).

(B) bottom panel: Lewy bodies (α-synuclein +) were found in the ventral midbrain region next to

the caudate nucleus indicated with a circle (left) in the coronal section of one hemisphere

(anterior-posterior 0), presence of Lewy bodies in a SARS-CoV-2-infected rhesus macaque (20x)

(middle), a magnified image (40x) of the square in the left image (right).

581 **Table 1: Animals featuring in this study**

Macaque species	Monkey ID	Infected Y/N	Age (Year)	Body weight[#] (kg)	Sex M/F
Rhesus	R1	Y	6	8.2	M
Rhesus	R2	Y	5	7.9	M
Rhesus	R3	Y	5	7.8	M
Rhesus	R4	Y	5	8.7	M
Rhesus	control R5	N	5	8.5	M
Rhesus	control R6	N	6	5.1	F
Cynomolgus	C1	Y	4	5.7	M
Cynomolgus	C2	Y	4	3.3	M
Cynomolgus	C3	Y	4	4.9	M
Cynomolgus	C4	Y	16	9.7	M
Cynomolgus	control C5	N	5	5.3	M
Cynomolgus	control C6	N	7	5.1	M

582 [#] at start of study

583 **Table 2**: **Histological findings.**

Marker		Brain area	Rhesus				Cynomolgus				Controls			
			R1	R2	R3	R4	C1	C2	C3	C4	R5	R6	C5	C6
T-cells (CD3+)	intra-parenchymal	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
	perivascular	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
	nodules	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
	meninges	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
Activated microglia (Mamu-DR+)	presence	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
	morphology: ramified/ amoeboid	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
	nodules	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
	meninges	pituitary gland												
		olfactory bulb												
		front brain												
		dorsal												
		ventral												
		cerebellum												
Lewy bodies (α-synuclein+)	inclusions	ventral midbrain												

Table 2 outlines the presence of 1) T-cells (CD3+ cells) in the brain tissue (intraparenchymal),

around blood vessels (perivascular), in group formation (nodules), or in the meninges, 2) activated

microglia (Mamu-DR+ cells) in different parts of the brain, the morphology as a measure for the

severity of activation (ramified or amoeboid), in group formation (nodules) or in the meninges, 3)

α-synuclein/Lewy bodies (α-synuclein + cells with inclusions) in the ventral midbrain region next

to the caudate nucleus. The last column shows the absence of most of these markers in the

control animals. Light grey: mild observation; dark grey: moderate observation (including

amoeboid microglia cells); black: moderate to severe observation of these markers.

Supplementary material:

C1	SUVmean					SUVpeak				
Tissue/day	*D8/10*	*D16/17*	*D22/23*	*D28/29*	*D35/36*	*D8/10*	*D16/17*	*D22/23*	*D28/29*	*D35/36*
Pituitary gland	3.1	2.5	2.5	3.5	3.9	5.3	3.9	4.1	6.1	6.4
Brain	2.2	1.9	2.4	1.9	2.6	5.0	4.6	4.7	4.6	4.6
Ratio	1.4	1.3	1.0	1.9	1.5	1.1	0.8	0.9	1.3	1.4
C2	SUVmean					SUVpeak				
Tissue/day	*D8/10*	*D16/17*	*D22/23*	*D28/29*	*D35/36*	*D8/10*	*D16/17*	*D22/23*	*D28/29*	*D35/36*
Pituitary gland	3.1	2.6	2.7	3.8	3.2	7.1	6.3	4.5	6.5	6.1
Brain	2.0	1.8	2.1	2.2	2.1	5.1	3.6	3.8	5.0	3.8
Ratio	1.5	1.5	1.3	1.7	1.5	1.4	1.8	1.2	1.3	1.6

Controls	SUVmean						SUVpeak					
Tissue/ animal	*NHP1*	*NHP2*	*NHP3*	*NHP4*	*NHP5*	*NHP6*	*NHP1*	*NHP2*	*NHP3*	*NHP4*	*NHP5*	*NHP6*
Pituitary gland	2.3	2.6	2.1	2.7	1.3	1.3	3.3	3.7	2.9	3.9	1.9	1.6
Brain	2.3	2.0	1.9	1.9	1.5	1.8	5.6	6.1	4.9	3.9	3.7	3.5
Ratio	1.0	1.3	1.1	1.4	0.9	0.7	0.6	0.6	0.6	1.0	0.5	0.5

Supplementary Table S1: Quantitative PET-CT analysis of the macaque brains

The standardized uptake values (SUVs) of animals C1 and C2 are represented together with the SUVs of six non-infected control rhesus macaques. Of these animals the average and peak uptake are determined for the pituitary gland, and the brain minus the pituitary gland. By dividing these the pituitary gland/brain ratio is calculated. For the SUVmean values above 1.0 are defined as slightly increased (light grey) and increased (dark grey) when equal or above 1.5. For the SUVpeak values above 1.0 are demarcated as slightly increased and above 1.2 as increased.