GPI-anchored FGF directs cytoneme-mediated bidirectional signaling to self-regulate tissue-specific dispersion

Lijuan Du [1], Alex Sohr [1,2], Yujia Li [1], Sougata Roy [1]*

[1] Department of Cell Biology and Molecular Genetics; University of Maryland, College Park, MD 20742, USA

[2] Present Address: Division of Cell and Gene Therapy, Center for Biologics Evaluation and Research, Food and Drug Administration, Silver Spring, MD 20993, USA.

*Corresponding author: sougata@umd.edu

Keywords: FGF, bidirectional signaling, GPI anchor, cytonemes, synapse, self-organization, Bnl, cell-adhesion-molecule (CAM), *Drosophila*

ABSTRACT

How signaling proteins generate a multitude of information to organize tissue patterns is critical to understanding morphogenesis. In *Drosophila*, FGF produced in wing-disc cells regulates the development of the disc-associated air-sac-primordium/ASP. Here, we show that FGF is Glycosylphosphatidylinositol-anchored to the producing cell surface and that this modification both inhibits free FGF secretion and activates target-specific bidirectional FGF-FGFR signaling through cytonemes. FGF-source and recipient ASP cells extend cytonemes that present FGF and FGFR on their surfaces and reciprocally recognize each other over distance by contacting through CAM-like FGF-FGFR binding. Contact-mediated FGF-FGFR binding induces bidirectional signaling, which, in turn, promotes ASP and source cells to polarize cytonemes toward each other and reinforce signaling contacts. Subsequent un-anchoring of FGFR-bound-FGF from the source cell membrane dissociates cytoneme contacts and delivers FGF target-specifically to ASP cytonemes for paracrine functions. Thus, GPI-anchored FGF organizes both source and recipient cells and self-regulates its cytoneme-mediated tissue-specific dispersion and signaling.

INTRODUCTION

During development, intercellular communication of morphogens is critical for embryonic cells to determine their positional identity, directionality, and interactions in an organized pattern to sculpt tissue. These conserved families of secreted morphogens/signals, such as Fibroblast Growth Factor (FGF), Hedgehog (Hh), Wingless (Wg)/Wnt, Epidermal Growth Factor (EGF), and Decapentaplegic (Dpp - a BMP homolog), act away from their sources and, upon binding to receptors, activate gene regulatory pathways to induce functions in recipient cells [1,2]. Strikingly, each signal and signaling pathway can generate a wide range of cell types and organizations in diverse contexts [3]. Understanding how signals might inform cells of their positional identity, directionality, and interactions and organize these functions in diverse tissue-specific patterns is critical to understanding morphogenesis.

The discrete tissue-specific organization of morphogen signaling is known to be dependent on the ability of signal-receiving cells to selectively sense and respond to a specific signal [3]. In contrast, traditional models predict that the signal presentation from the source via free secretion and extracellular diffusion is a non-selective process. However, recent advances in microscopy revealed that both signal-producing and receiving cells could extend signaling filopodia named cytonemes and selectively deliver or receive signals through cytoneme-cell contact sites [4–9]. Essential roles of cytonemes or cytoneme-like filopodia have been discovered in many vertebrate and invertebrate systems and are implicated in most signaling pathways, including Hh, Dpp, FGF, EGF, Ephrin, and Wnt under various contexts [4–18]. The prevalence and similarities of these signaling filopodia suggest that the polarized target-specific morphogen exchange through filopodial contacts is an evolutionarily conserved signaling mechanism.

These findings bring along a paradox - not only do signals instruct cells and organize discrete cellular patterns, but cells also control the spatiotemporal patterns of signal presentation and reception by organizing the distribution of cytonemes and cytoneme contacts [6,9]. The functioning of this interdependent relationship of signals and signaling cells through cytonemes, however, would require precise spatiotemporal coordination

between cytoneme contact formation and signal release. We started the current

investigation with the premise that a better understanding of the processes that produce

cytoneme contacts and control contact-driven signal release is essential to

understanding morphogenesis. We asked: (1) How do cytonemes recognize a specific

target cell and form signaling contacts? (2) How are secreted signals controlled for

polarized target-specific release, exclusively at the cytoneme contact sites? (3) Do

cytoneme contact formation and signal release spatiotemporally coordinate with each

other? If so, how?

To address these questions, we focused on the inter-organ dispersion of a *Drosophila*

FGF, Branchless (Bnl), during the development of the wing imaginal disc-associated air-

sac primordium (ASP) [19,20]. Bnl is expressed in a discrete group of wing disc cells, and it

induces morphogenesis of the tubular ASP epithelium that expresses the Bnl receptor,

Breathless (FGFR/Btl) [9,19,21]. Epithelial cells at the ASP tip extend polarized Btl-

containing cytonemes to contact Bnl-producing wing disc cells and directly take up Bnl

in a contact- and receptor-dependent manner [5,9]. The formation of Bnl-specific polarity

and contacts of ASP cytonemes are self-sustained by Bnl-signaling feedbacks [9].

Consequently, Bnl reception and signaling via cytonemes can precisely adapt and

dynamically coordinate with ASP growth. With increasing distance from the Bnl-source,

ASP cells extend gradually fewer polarized Bnl-receiving cytonemes, leading to the

emergence of asymmetric Bnl dispersion and signaling patterns within the ASP [9].

However, how ASP cytonemes might recognize the *bnl*-source for signaling contacts,

and, on the other hand, how Bnl producing cells might both inhibit free Bnl secretion and

facilitate Bnl release selectively at the cytoneme contact sites are unknown.

Here, we report that Bnl is post-translationally lipid-modified by the addition of a

glycosylphosphatidylinositol (GPI) moiety, which anchors Bnl to the outer leaflet of its

source cell membrane, and selectively presents Bnl to only Btl-expressing cells via

cytonemes. We provide evidence that GPI-anchored Bnl can act as a cell adhesion

molecule (CAM) [22–28] and produce target-specific cytoneme contacts and contact-

dependent CAM-like bidirectional Btl-Bnl signaling between source and recipient cells.

We show that this bidirectional signaling enables the source and recipient cells to

coordinate over distance and reciprocally promote each other to polarize cytonemes to

establish contacts. Our results further provide evidence that although the GPI anchor

inhibits free Bnl secretion, it facilitates the release of Bnl exclusively at the cytoneme

contact sites, ensuring its tissue-specific morphogen-like dispersion and signaling

patterns. These findings suggest that while cytonemes are critical for organizing tissue-

specific Bnl signaling, the GPI-anchored Bnl programs the spatiotemporal distribution of

cytoneme contacts to self-regulate its dispersion.

RESULTS

Bnl presentation and reception are reciprocally polarized toward each other.

Bnl is produced in the wing disc and transported target-specifically to the overlaying

ASP via Btl-containing ASP cytonemes across a layer of interspersed myoblasts (Fig.

1a,b) [9]. The *bnl*-specific polarity of ASP cytonemes might be determined by the extrinsic

patterns of Bnl presentation from the source. Previously, non-permeabilized anti-Bnl

immunostaining (αBnl^{ex}) designed to detect secreted externalized Bnl (Bnl^{ex}) [9,29]

showed that the Bnl^{ex} was not randomly dispersed in the source-surrounding

extracellular space (Fig.1a). Instead, Bnl^{ex} was restricted exclusively to the basal

surface of Bnl-producing cells and on the ASP cytonemes. Importantly, even within the

bnl-expressing disc area, Bnl^{ex} puncta were asymmetrically congregated near the

contact sites of Btl-containing ASP cytonemes that received Bnl^{ex} (Fig.1a,c). These

results indicated that the Bnl presentation is likely to be spatially polarized.

To examine if Bnl distribution in source cells is spatially biased toward the ASP, we co-

expressed Bnl:GFP with mCherryCAAX (prenylated mCherry for membrane marking)

under *bnl-Gal4*. Strikingly, although *bnl-Gal4*-driven mCherryCAAX equally labeled all

source cells, Bnl:GFP was asymmetrically enriched at the ASP-proximal source area

(Figs.1d-d"; Supplementary Figure 1a,a'). Bnl:GFP puncta were also displayed on short

polarized cytonemes emanating from the ASP-proximal disc cells (Fig.1d;

Supplementary Figure 1a). To further verify if the Bnl presentation is polarized via

cytonemes, we imaged the distribution of endogenous Bnl:GFP^{endo}, expressed from a

bnl:gfp^{endo} knock-in allele [9], in the mCherryCAAX-marked *bnl*-source. Bnl:GFP^{endo}

puncta represented all Bnl isoforms. Indeed, Bnl:GFP^{endo} puncta were selectively

enriched in source cell cytonemes that oriented toward the ASP (Fig.1e,e';

Supplementary Figure 1b).

To examine the organization of Bnl-presenting source cytonemes, we observed live

wing discs that expressed a fluorescent membrane marker (e.g., CD8:GFP or

CherryCAAX) either in all of the *bnl*-expressing cells (Fig.1f,g) or in small clones of cells

within the Bnl-expressing area (see Methods; Fig.1h-h"). Three-dimensional image

projections of live discs revealed that each of the Bnl-expressing columnar cells

proximal to the ASP extended ~2-4 short (<15 μm) cytonemes perpendicularly from

their basal surface (Fig.1g-h"; Supplementary Figure 1d, Movie 1). The organization of

source cells, therefore, can be described as polarized for Bnl presentation with basal

cytonemes extending toward the ASP or ASP cytonemes. This organization is mirrored

in the ASP, which is known to exhibit polarized Btl synthesis/localization, Bnl reception,

and cytoneme orientation toward source cells [9]. Thus, the cellular components

responsible for Bnl presentation in the disc source and for its reception in the ASP are

likely to be reciprocally polarized toward each other.

Reciprocal guidance of Bnl-presenting and -receiving cytonemes.

To examine if Bnl-presenting and -receiving cytonemes could reciprocally guide each

other's polarity, we examined live wing discs harboring the CD8:GFP-marked ASP and

mCherryCAAX-marked source. Time-lapse imaging of ex vivo cultured discs revealed

that ASP and source cytonemes orient toward each other and transiently contact each

other's tips, bases, or shafts as they dynamically extend and retract (Fig. 1i-k'';

Supplementary Movie 2). Both cytoneme types had short lifetimes and repeated cycles

of contact association-dissociation (Fig. 1l-n; Supplementary Figure 1e-h, Movie 2,

Table 1). We also examined the inter-cytoneme interactions during the development of

the ASP from the early-to-late L3 larval stages. Despite dynamic morphological changes

in the growing ASP and disc, the relative positions of the ASP, *bnl*-source, and the site

of inter-cytoneme interactions were maintained throughout the development

(Supplementary Figure 1i-l"). Thus, interacting cells in the ASP and *bnl*-source polarize

to face each other and apparently maintain a tissue-level niche at the ASP:source

interface to promote cytoneme-mediated interactions.

Based on our previous observations [5,9], Bnl is exchanged at the cytoneme contact sites.

However, it was technically challenging to visualize Bnl exchange during dynamic inter-

cytoneme interactions. Therefore, we sought to genetically ablate source cytonemes in

bnl:gfp[endo] larvae and analyze if the levels of Bnl:GFP[endo] uptake in the ASP are

reduced. An actin modulator formin, Diaphanous (Dia), could influence source

cytonemes. Overexpression of Dia:GFP or a constitutively active Dia:GFP[ca] induced

cytonemes (Fig.2a-e). Asymmetric enrichment of Dia:GFP[ca] puncta in source cytoneme

tips suggested localized Dia activity (Fig.2c). In contrast, *dia* knockdown (*dia-i*) in the

mCherryCAAX-marked source (*bnl-Gal4 x UAS-dia-i,UAS-mCherryCAAX*) suppressed

cytoneme formation without any visible effects in *bnl* expression (Fig.2a-e;

Supplementary Figure 2a). Importantly, the *dia-i* mediated ablation of source cytonemes

in *bnl:gfp*[endo] larvae significantly reduced Bnl:GFP[endo] uptake in the ASP. These ASPs

were abnormally stunted, suggesting a reduction in Bnl signaling (Fig.2f-h). Thus,

source cytonemes are required to deliver Bnl to the ASP.

Inter-cytoneme Bnl exchange is consistent with reports that Hh and Wg are both sent

and received by cytonemes[10,30,31]. However, how do source and ASP cytonemes find

and adhere to each other? Dynamic interactions of Bnl-exchanging cytonemes that are

convergently polarized toward each other suggested a possibility of contact-dependent

reciprocal guidance of source and recipient cytonemes. To test this possibility, we first

ablated source cytonemes by *dia-i* expression and analyzed the non-autonomous

effects on CD2:GFP-marked ASP cytonemes. The ablation of source cytonemes

significantly reduced the long, polarized ASP tip cytonemes (Fig.2i-k). In contrast, short,

randomly oriented ASP cytonemes were unaffected. Thus, Bnl-presenting cytonemes

are required for the formation of the polarized Bnl-receiving ASP cytonemes.

We next removed ASP cytonemes by expressing *dia-i* under *btl-Gal4* and recorded non-
autonomous effects on mCherryCAAX-marked source cytonemes. The *dia-i* expression
had to be controlled with Gal80[ts] to avoid lethality (see Methods). Tracheal *dia-i*
expression not only reduced ASP cytonemes but also non-autonomously reduced
source cytonemes (Fig.2l-n'). Similarly, tracheal expression of a dominant-negative form
of Btl (Btl-DN) was known to suppress ASP growth and cytoneme formation without
affecting the wing disc development [19]. When both source and Btl-DN-expressing
tracheal cells were marked, the complete loss of ASP and ASP cytonemes was found to
produce a corresponding loss of *bnl*-source cytonemes (Fig.2o,p). Thus, Btl-presenting
ASP cytonemes are required to produce source cytonemes that polarize toward the
ASP. Collectively, these results suggested that the source and recipient cytonemes
reciprocally guide each other to form signaling contacts.
220
CAM-like Btl-Bnl binding drives bidirectional contact matchmaking of cytonemes.
The above results also suggested that the inter-cytoneme interactions might recruit and
activate a bidirectional signaling mechanism, responses of which could induce ASP
cells to extend Btl-containing cytonemes toward the source and activate source cells to
extend Bnl-containing cytonemes toward the ASP. We hypothesized that such selective
matchmaking between source and ASP cytonemes could be mediated by the binding of
surface-displayed Btl and Bnl. In this model, Btl and Bnl are analogous to cell-
recognition or cell-adhesion molecules (CAMs), physical interaction of which can
produce selective cell-cell adhesion and contact-mediated bidirectional signaling [23,32,33].
The initiation of CAM-like interactions might not require Btl to activate canonical
transcriptional outputs [34]. An alternative possibility is that the Bnl-Btl binding activates
MAPK signaling and transcription of target genes in the ASP, and these gene products,
in turn, non-autonomously act on the wing disc *bnl*-source to induce a response.
234
Notably, Btl-DN can bind to Bnl via its extracellular ligand-binding domain
(Supplementary Figure 2b) and can heterodimerize with *WT* Btl, but cannot activate
nuclear MAPK signaling due to the lack of its intracellular kinase domain [19,21]. As
observed before[13,19], while most wing discs with Btl-DN-expressing trachea (TC)

completely suppressed ASP/ASP cytonemes, occasionally a few produced partially

suppressed nascent ASP with variable numbers of ASP cytonemes orienting toward the

source. Strikingly, in each of these discs, the appearance of polarized ASP cytonemes

correlated with the concomitant appearance of similar numbers of polarized source

cytonemes, forming direct cytoneme:cytoneme contacts (Fig.3a-c). These results were

consistent with the idea that source and ASP cells reciprocally promote each other's

cytoneme-forming polarity by contacting each other via cytonemes and that the Btl-

DN::Bnl binding is likely to induce the reciprocal cytoneme-mediated matchmaking.

Indeed, non-permeabilized αBnl[ex] staining showed that Bnl[ex] was selectively enriched at

these inter-cytoneme contact sites (Fig.3d,d', Supplementary Figure 2c-c'''). Similarly,

when we expressed Btl-DN:Cherry in the trachea, ASP cytonemes were enriched with

Btl-DN:Cherry puncta that colocalized with Bnl[ex] (Fig.3e,e'; Supplementary Figure 2d,d').

Thus, Btl-DN localized on the ASP cytonemes could bind to Bnl[ex] presented on the

source membrane. We predicted that this Btl-DN::Bnl interaction was capable of

inducing the bidirectional cytoneme::cytoneme matchmaking for contacts. This result

provided a clue that surface-localized Btl and Bnl might act like heterophilic cell

adhesion/recognition molecules (CAM). CAM-CAM interactions are known to induce

contact-dependent bidirectional signaling that controls cell shape, polarity, and dynamic

interactions by modulation the local organization of subcellular regulators such as the

actomyosin complex [22–28].

To verify if Btl and Bnl can act as CAMs, we performed an in vitro cell culture-based

assay using *Drosophila* S2 cells (embryonic hemocyte lineage). S2 cells lack

endogenous Btl and Bnl expression (modENCODE), but when S2 cells ectopically

expressed Bnl, αBnl[ex] immunostaining detected enrichment of Bnl[ex] on the expressing

cell surface similar to what was found on the wing disc Bnl source (Supplementary

Figure 2e,e'). The lack of polarity and cell junctions in S2 cells was another suitable

feature for ectopic induction of these properties. As illustrated in Figure 3f, we mixed

and co-incubated Bnl:GFP-expressing cells (S2-Bnl:GFP) with cells that expressed

either Btl:Cherry (S2-Btl:Cherry), Btl-DN:Cherry (S2-Btl-DN:Cherry), or a secreted

Btl:Cherry (sBtl:Cherry) that lacked transmembrane and intracellular domains of Btl (S2-

sBtl:Cherry, see Methods).

S2-Bnl:GFP and S2-Btl:Cherry cells alone did not show homophilic cell-cell adhesion,

but, when mixed and co-cultured, S2-Bnl:GFP cells selectively trans-paired with S2-

Btl:Cherry by forming polarized trans-synaptic receptor-ligand clusters (Fig.3g-h;

Supplementary Figure 2h; Supplementary Movie 3). Moreover, the binding of Btl:Cherry

and Bnl:GFP induced reciprocally polarized congregation of the receptors and ligands at

the contact interface of the trans-paired cells, thereby stabilizing cell contacts. We also

observed localized enrichment of cortical f-actin (phalloidin-stained) at the synaptic

interface (Supplementary Movies 3-5), similar to what is observed during the formation

of receptor-ligand-dependent immunological synapse [35]. These results are consistent

with the CAM-like Btl-Bnl interactions inducing bidirectional recognition of cells for

contacts.

Secondly, almost all (averaging 98%) of S2-Btl:Cherry cells that were trans-paired to

S2-Bnl:GFP cells had nuclear-localized dpERK. In the same co-culture experiment,

unpaired (non-adhering) S2-Btl:Cherry lacked nuclear dpERK, indicating inactive

FGF/MAPK signaling in these cells (Fig. 3i-k'; Supplementary Figure 2i,i'). These

unpaired S2-Btl:Cherry cells were similar to control S2-Btl:Cherry cells cultured alone or

to non-transfected S2 cells that rarely had nuclear dpERK (average ~2-5% of cells),

(Fig.3j,j'). In contrast, when S2-sBtl:Cherry cells were cocultured with S2-Bnl:GFP, they

did not trans-pair with S2-Bnl:GFP and lacked nuclear dpERK (Fig.3h,i,n;

Supplementary Figure 2g,j,j'). Therefore, we considered that the Btl-Bnl-mediated trans-

pairing of S2 cells is a successful in vitro recapitulation of contact-dependent Btl-Bnl

signaling between the ASP and Bnl source via cytoneme::cytoneme contacts [9].

Strikingly, when co-cultured, S2-Btl-DN:Cherry cells showed strong selective trans-

pairing with S2-Bnl:GFP similar to the S2-Btl:Cherry control. However, the trans-paired

S2-Btl-DN:Cherry did not activate nuclear localization of dpERK due to the lack of its

intracellular kinase domain (Fig.3h,i,l-m'; Supplementary Figure 2f,h). Therefore,

surface-localization and physical interactions of Btl/Btl-DN and Bnl were sufficient to

induce the bidirectional contact matchmaking of the Bnl exchanging cells. This is

consistent with the idea that Btl:DN (or Btl) and Bnl can act as heterophilic CAMs.

These results suggested a possibility that the contact-dependent Bnl-Btl binding might

produce two distinct signaling responses. First, a non-canonical CAM-like bidirectional

signaling response that mediates contact-dependent reciprocal matchmaking of Bnl

source and recipient cells/cytonemes. Second, the release and exchange of Btl-bound

Bnl molecules at the inter-cellular/cytoneme contact sites that promote the canonical

nuclear MAPK signaling in recipient cells.

Bnl is tethered to the source cell surface by a GPI anchor.

However, to drive heterophilic CAM-like bidirectional recognition for synapse, Bnl needs

to be tightly associated to the source cell membrane. How might a secreted protein be

associated exclusively on the source cell surface to act as a CAM without being

randomly dispersed in the extracellular space? A probable mechanism emerged while

exploring post-translational Bnl modifications during its intracellular trafficking [36]. We

knew that a small N-terminal portion (residue 1-164) upstream of the central 'FGF

domain' of Bnl is cleaved off in the source cell Golgi by Furin1 to facilitate polarized

trafficking of the remaining C-terminal signaling portion of Bnl to the basal side of the

source cell (Fig.4a; [36]). When cells expressed a Furin-sensor $HA_1Bnl:GFP_3$ construct

with HA (site 1) and GFP (site 3) flanking the Furin cleavage site, the cleaved HA-

tagged portion was retained in the Golgi and the truncated $Bnl:GFP_3$ fragment was

externalized for dispersal [36]. Therefore, we hypothesized that cells expressing a triple-

tagged $HA_1Bnl:GFP_3Cherry_c$ construct with a C-terminal mCherry fusion (Fig.4a) would

externalize a truncated $Bnl:GFP_3Cherry_c$ portion marked with both GFP and mCherry.

However, when we expressed $HA_1Bnl:GFP_3Cherry_c$ (hereafter called $Bnl:GFP_3Cherry_c$)

in S2 cells, GFP and mCherry tags were separated, and, importantly, while the

$Bnl:GFP_3$ portion was localized on the cell surface (detected with αGFP^{ex}

immunostaining), the C-terminal mCherry remained intracellular (Fig.4b-b'''). The C-

terminal mCherry tag did not alter the predicted topology and physico-chemical

properties of Bnl (see Supplementary information). In fact, when Bnl:GFP$_3$Cherry$_c$ was

expressed in the wing disc source under *bnl-Gal4*, the mCherry-tag was retained in

source cells and the Bnl:GFP$_3$ portion was efficiently delivered to the ASP (Fig.4c).

These results indicated an intracellular Bnl:GFP$_3$Cherry$_c$ cleavage, which separated the

C-terminal mCherry prior to the secretion of the truncated Bnl:GFP$_3$ portion. Cleavages

at multiple locations in the Bnl backbone was consistent with the detection of multiple

Bnl:GFP bands in Western blots of expressing cell lysates [36].

Bioinformatic analyses revealed that the Bnl C-terminus has a short 20 amino acid

hydrophobic tail preceded by a hydrophilic spacer (Fig.4a,a'). A 15-20 residue long

hydrophobic C-terminal tail together with an immediately upstream hydrophilic spacer

commonly constitutes the signal sequence (SS) of a pro-GPI-anchored protein (pro-

GPI-APs) [25,37–39]. The C-terminal hydrophobic portion of the SS is cleaved off and

replaced with a GPI moiety via a transamidation reaction in the endoplasmic reticulum

(ER). GPI-APs are then trafficked to the cell surface and anchored to the outer leaflet of

the plasma membrane by the phosphatidylinositol (PI) portion of the GPI moiety [25,37,38]

(Fig.4d). Because the presence of C-terminal tags does not prevent glypiation of pro-

GPI-APs [39], we surmised that GPI-anchoring of Bnl might explain the intracellular

cleavage of mCherry from Bnl:GFP$_3$Cherry$_c$ prior to the surface display of its truncated

Bnl:GFP$_3$ portion.

To detect Bnl glypiation, we followed the phosphoinositide phospholipase C (PI-PLC)-

dependent shedding assay. Since PI-PLC specifically cleaves the GPI moiety, PI-PLC-

dependent shedding of a cell surface protein confirms its GPI anchoring [40]. Using the

Gal4/UAS-based expression in S2 cells, we ectopically expressed GFP-GPI [41] (positive

control), untagged Bnl (co-transfected with CD8:GFP for detecting transfection and

expression), Bnl:GFP$_3$, HA$_1$Bnl:GFP$_3$ (henceforth referred as Bnl:GFP), and a

palmitoylated cell-surface protein, the constitutively active *Drosophila* EGF, cSpitz:GFP

[42] (negative control). The levels of cell surface proteins with and without PI-PLC

treatment were probed with non-permeabilized αGFP^{ex} or αBnl^{ex} (for untagged Bnl)

immunostaining, and the ratio of the surface proteins to the total protein per cell was

compared between cells/conditions (see Methods). These analyses showed that the PI-

PLC treatment specifically removed source-surface Bnl^{ex} and $Bnl:GFP^{ex}$, like control

GPI-GFP, but PIPLC did not remove $cSpitz:GFP^{ex}$ (Fig.4d,d',f, Supplementary Figure

3a-e). Thus, Bnl is a GPI-AP.

An in-silico analysis predicted $Bnl-S^{741}$ as a probable glypiation site (ω-site). Therefore,

to verify if Bnl's C-terminal region acts as a SS, we generated - (i) Bnl:GFPΔC, lacking

the C-terminal 40 amino acid residues including the putative ω site; (ii) Bnl:GFPΔC-TM,

where the transmembrane domain from the mammalian CD8a was added to the C-

terminus of Bnl:GFPΔC; (iii) Bnl:GFP-ω^{m}, Bnl:GFP with mutated ω, $\omega+1$, and $\omega+2$ sites;

and (iv) bGFP-GPI, a secreted super-folder GFP (secGFP [9]) fused to Bnl's C-terminal

53 amino acids region (see Methods) (Fig.4e). Bnl:GFPΔC and Bnl:GFP-ω^{m} were not

localized on the producing cell surface, even without the PI-PLC treatment (Fig.4d',g;

Supplementary Figure 3f,h,i). However, when a TM domain was added to Bnl:GFPΔC

(i.e., Bnl:GFPΔC-TM), the protein was surface localized in a PI-PLC-resistant manner

(Fig.4d'-f; Supplementary Figure 3g,h).

Usually, readily secreted secGFP was not detectable on the expressing S2 cell surface

(Supplementary Figure 3j,j'), but when the same secGFP was added with Bnl's SS to

make bGFP-GPI, it showed PI-PLC-sensitive surface-localization (Fig.4d'-f). Thus, Bnl's

SS is required for glypiation. Secreted Bnl can bind to GPI-anchored glypicans, and the

binding site occurs at its conserved FGF domain (Fig.4a) [43]. Therefore, we examined if

PIPLC-dependent removal of surface Bnl could be due to the removal of Bnl-bound cell

surface glypicans. We generated a Bnl:GFP$^{\Delta FGF}$ construct, where the core 'FGF domain'

of Bnl was replaced with the sfGFP sequence. This chimeric protein showed clear

PIPLC-sensitive surface localization (Fig.4e; Supplementary Figure 3k), indicating that

PIPLC directly removes GPI-anchored Bnl.

The GPI-AP signal sequences (including ω-sites) are known to have little sequence

conservation, and its extreme C-terminal positioning is not an absolute requirement [39,44].

The Bnl constructs described here, were derived from the well-characterized *bnl-PA*

isoform [20] that has been used in all previous reports of ectopic Bnl expression. Bnl also

has a shorter splice variant (PC) (FlyBase) with altered C-terminal hydrophobicity

(Supplementary Figure 4a-a'''). Therefore, we generated a Bnl:GFP-PC construct

(Methods) and expressed it in S2 cells. PI-PLC treatment of S2 cells expressing

Bnl:GFP-PC removed the surface-localized Bnl:GFP-PC, indicating its GPI-anchored

display (Supplementary Figure 4b-g). Thus, both Bnl-PC and Bnl-PA isoforms are

glypiated, but strikingly, with two distinct signal sequences.

Next, to detect Bnl's GPI-anchoring in vivo, we developed a PI-PLC assay on live ex

vivo cultured wing discs (see Methods). First, we detected native extracellular Bnl by

non-permeabilized immunostaining of ex vivo cultured w^- wing discs with a Bnl antibody

that detects all Bnl isoforms (Supplementary Figure 4a,a'). Bnl^{ex} that normally was

asymmetrically enriched on the disc source was significantly reduced with PI-PLC

treatment (Fig.5a-c). When Bnl, Bnl:GFP, Bnl:GFPΔC-TM, Bnl:GFPΔC, and Bnl:GFP-ω^m

constructs were expressed under bnl-$Gal4$, PI-PLC treatment significantly reduced Bnl^{ex}

and Bnl:GFP^{ex} on the source surface, but not Bnl:GFPΔC-TM^{ex} (Fig.5d-m). As observed

in S2 cells, Bnl:GFPΔC^{ex} and Bnl:GFP^{ex}-ω^m puncta were not detected on source cells

irrespective of the PI-PLC treatment (Fig. 5i,j,m; Supplementary Figure 5a-b).

Although Bnl:GFPΔC^{ex} was absent from the source membrane, it was broadly spread

through the extracellular disc areas surrounding the source and was also received by

the ASP, suggesting that the protein was readily secreted and randomly dispersed from

the source (Supplementary Figure 5a,a'). Externalized Bnl:GFPΔC^{ex} contains the

conserved glypican binding FGF domain, yet it was absent on the source surface,

indicating that the secreted Bnl:GFPΔC^{ex} was not restricted on the source surface by

glypican binding. In contrast to Bnl:GFPΔC^{ex}, Bnl:GFP-ω^m showed severely reduced

externalization (Supplementary Figure 5b). This was consistent with previous reports of

ER retention of the uncleaved pro-GPI-APs, in contrast to the normal trafficking of the

same protein with deleted SS [45,46]. These results indicated that Bnl is cleaved at its C

terminus and added with a GPI moiety, which both facilitated Bnl externalization and

inhibited its free secretion.

GPI-anchored Bnl acts as a CAM to produce target-specific cytoneme contacts.
To test if GPI anchoring is required for Bnl's CAM-like activity, we employed the cell
culture-based assay to compare the trans-pairing efficiency of S2-Btl:Cherry with either
S2-Bnl:GFP∆C, S2-Bnl:GFP∆C-TM, or S2-Bnl:GFP (control). Notably, when Btl:Cherry
was co-transfected with either Bnl:GFP, Bnl:GFP∆C, or Bnl:GFP∆C-TM in same cells,
almost all (at least 90%) of the cells expressing both ligands and receptors had nuclear
dpERK (Supplementary Figure 5c, Table 3a). Thus all Bnl variants could efficiently
activate Btl:Cherry when co-expressed in the same cell. When co-cultured, S2-Bnl:GFP
and S2-Btl:Cherry were trans-paired with each other and the trans-paired S2-Btl:Cherry
activated nuclear dpERK localization as shown before (Fig.3). In contrast, co-cultured
S2-Bnl:GFP∆C and S2-Btl:Cherry cells were rarely trans-paired (only ~1% frequency of
juxtaposition) (Fig.6a,c; Supplementary Figure 5d, Table 3b). Even when S2-Btl:Cherry
were juxtaposed to S2-Bnl:GFP∆C they lacked polarized Btl-Bnl co-clustering at the
contact interface. Moreover, almost 85% of S2-Btl:Cherry cells that were nearby to the
S2-Bnl:GFP∆C source lacked dpERK (Fig.6a; Supplementary Figure 5d, Table 3b). A
few dpERK-positive S2-Btl:Cherry that were found, had unpredictable random locations
relative to the S2-Bnl:GFP∆C.

In contrast to S2-Bnl:GFP∆C, S2-Bnl:GFP∆C-TM cells selectively trans-adhered to S2-
Btl:Cherry as efficiently as S2-Bnl:GFP by forming polarized trans-synaptic receptor-
ligand co-clusters (Fig.6b,c; Supplementary Table 3b). Polarized trans-pairing of
Bnl:GFP∆C-TM and Btl:Cherry also induced MAPK signaling in the Btl:Cherry-
expressing cells, but at a lower frequency than the control S2-Bnl:GFP::S2-Btl:Cherry
interactions (Fig.6b; Supplementary Table 3b). Notably, MAPK signaling was activated
only in those trans-paired S2-Btl:Cherry cells that had high numbers of internalized
receptor-bound Bnl:GFP∆C-TM puncta (Fig.6b). It is possible that although being TM-
tethered, Btl:Cherry-bound Bnl:GFP∆C-TM could somehow be internalized into the
recipient cells through the contact sites to activate nuclear MAPK signaling. However,
irrespective of the activation of MAPK signaling, the trans-synaptic binding of

Bnl:GFPΔC-TM with Btl:Cherry were sufficient to induce reciprocal polarity of signal

delivery and reception between the interacting cells. This is consistent with the CAM-like

activity of membrane-tethered Bnl.

To test if CAM-like bidirectional Btl::Bnl interactions occur through cytonemes in vivo,

we compared how GPI-modified (Bnl:GFP) and non-GPI-modified Bnl:GFP variants

affect source and ASP cytonemes. Despite the Bnl:GFP overexpression, both ASP and

source cytonemes retained their reciprocal polarity toward each other (Figs.6d-f;

Supplementary Figure 6g). An increase in extension-retraction rates of ASP cytonemes

in this condition suggested an increase in signaling activity in the ASP (Supplementary

Movie 6, Table 1). In contrast, overexpressed Bnl:GFPΔC significantly suppressed the

formation of polarized cytonemes from both source and ASP cells (Fig.6g-i;

Supplementary Figure 6a,b,g). Short cytonemes, when detectable, lacked any

directional bias and Bnl:GFPΔC localization.

Importantly, unlike Bnl:GFPΔC, Bnl:GFPΔC-TM induced both ASP and source cells to

extend large numbers of long polarized cytonemes that were adhered to each other

(Fig.6j-l; Supplementary Figure 6c-h, Movies 7-9). Bnl:GFPΔC-TM puncta populated at

multiple inter-cytoneme contact interfaces (Figs.6j, 7a-a''). To visualize the CAM-like

Bnl-Btl binding at the inter-cytoneme contacts, we expressed Bnl:GFPΔC-TM from the

CD4:IFP$_2$-marked wing disc source in *btl:cherry*[endo] larvae. These larvae expressed

endogenous Btl:Cherry[endo] in the ASP. Btl:Cherry[endo] puncta on the ASP cytonemes

were co-clustered with Bnl:GFPΔC-TM puncta at multiple contact sites along the length

of the source and recipient cytonemes (Fig.7b-c). Bnl:GFPΔC-TM-exchanging

cytonemes showed higher stability and longer contact lifetime than WT or Bnl:GFP-

exchanging cytonemes (Fig.7d,e; Supplementary Figure 6h, Movies 10,11, Table 1).

The increased stability of inter-cytoneme adhesion might account for the higher intensity

of bidirectional responses with Bnl:GFPΔC-TM than with Bnl:GFP. These results

supported CAM-like bidirectional Btl:Bnl interactions at cytoneme contacts.

To verify if bidirectional Btl:Bnl interactions can induce reciprocal guidance of source

and recipient cytonemes, we produced randomly-localized mCherryCAAX-marked wing

disc clones that expressed Bnl:GFPΔC-TM (Fig.7f-j). Clones in the wing disc pouch that

occurred far away from the ASP were unable to establish contact with the ASP. These

clones had only short, randomly oriented signal-containing cytonemes (Fig.7f,g,h). In

contrast, ASP-proximal clones extended long polarized cytonemes and establish

contacts with the ASP (Fig.7f,i,j). These results were consistent with the contact-

dependent activation of a retrograde signaling response in the Bnl-source that

reinforced the ASP-specific source cytoneme polarity. ASP cells also can extend

cytonemes toward ectopic Bnl-expressing clones [19] and the source-specific polarity can

be reinforced by Bnl signaling feedbacks [9]. Moreover, when we produced randomly-

localized Btl:GFP-expressing wing disc clones, these clones and the mCherryCAAX-

marked Bnl-source were found to extend polarized cytonemes toward each other

(Fig.7k-m'). These results showed that the cytoneme-mediated Bnl:Btl signaling is

bidirectional and that this bidirectional signaling is required for the reciprocal guidance

of source and recipient cytonemes.

Bnl's GPI anchor ensures contact-dependent tissue-specific Bnl release.

Although Bnl:GFPΔC-TM induced strong bidirectional responses that were manifested

in cytoneme polarity and inter-cytoneme contacts, Bnl:GFPΔC-TM-exchanging

cytonemes had a significantly longer lifetime than WT or Bnl:GFP-exchanging

cytonemes (Fig.7d,e; Supplementary Figure 6h, Supplementary Table 1). Moreover,

unlike Bnl:GFP, Bnl:GFPΔC-TM puncta were often abnormally internalized into the ASP

with colocalized source cell membrane, indicating a defect in release of the TM-tethered

signal from the source cell membrane (Fig.8a,d,e; Supplementary Figure 7a-c';

Supplementary Movie 14). When both source and ASP cells were simultaneously

marked and imaged in time-lapse, Bnl:GFPΔC-TM-exchanging cytonemes appeared to

resist contact dissociation, leading to cytoneme breakage and absorption of the source

membrane in the ASP (Supplementary Movie 11). We predicted that GPI anchoring is

critical for efficient tissue-specific Bnl release and contact disassembly.

To investigate if GPI-anchoring of Bnl facilitates its target-specific release, we compared

the spatial distribution of GPI-modified (Bnl:GFP) and non-GPI-modified Bnl:GFP

constructs expressed from the mCherryCAAX-marked wing disc Bnl source. As

observed before [9,36], despite overexpression in the disc *bnl*-source, Bnl:GFP puncta

were exclusively transferred from the disc source to the ASP (Fig.8a; Supplementary

Movie 12). In contrast, two ΔC variants (Bnl:GFPΔC and Bnl:GFPΔC_{168}) showed

dispersion in the non-specific disc areas surrounding the source (Fig.8b,c;

Supplementary Movies 13, 14). Importantly, the corresponding Bnl:GFPΔC-TM and

Bnl:GFPΔC_{168}-TM variants regained the exclusive ASP-specific distribution, but their

range of distribution was restricted only to the ASP tip, indicating reduced levels of

signal exchange (Fig.8d,e; Supplementary Figure 7a-c'; Supplementary Movies 15, 16).

These results suggested that GPI anchoring might be required for both inhibiting free

Bnl secretion/dispersion and facilitating target-specific contact-dependent Bnl release.

To better understand the dual roles of GPI anchoring, we compared the ASP-specific

and non-specific spreading of Bnl:GFP variants over time in live ex vivo cultured discs

(see Methods). To accurately estimate the levels of signal uptake in the ASP, we took

advantage of the dual-tagged Furin-sensors - HA_1Bnl:GFP$_3$, HA_1Bnl:GFP$_3\Delta$C-TM, and

HA_1Bnl:GFP$_3\Delta$C (Fig.8f-s). As expected, in ex-vivo cultured larval discs, the N-terminal

HA-tagged portion (αHA-probed) of all three constructs were cleaved in source cells

and only their truncated Bnl:GFP$_3$ portions were transferred to the ASP (Fig.8f,g,j,m).

However, when Furin inhibitors were added to the culture media, uncleaved signals

(αHA-probed) were received by the ASP[36]. Therefore, the fraction of the HA-probed

uncleaved signal relative to the total levels of the GFP-probed signal (pre-existing

Bnl:GFP$_3$+ HA_1Bnl:GFP$_3$) accumulated in the ASP during a Furin-inhibited period

provided a semi-quantitative estimate of the rate of signal uptake in the ASP (Fig.8f').

As observed before [36], the levels of HA_1Bnl:GFP$_3$ (control) uptake in the ASP gradually

increased with increasing duration of the culture. In comparison, the levels of

HA_1Bnl:GFP$_3\Delta$C-TM and HA_1Bnl:GFP$_3\Delta$C in the ASP did not change dramatically,

indicating a slow rate of ASP-specific transfer of these variants (Fig.8g-p). Notably, even

after a 5 h of culture, $HA_1Bnl:GFP_3$ dispersed exclusively to the ASP (Fig.8q). In

contrast, within a 5 h of incubation period, $HA_1Bnl:GFP_3\Delta C$ was randomly localized in

the source-surrounding disc areas, but was barely received by the ASP from the same

disc (Fig.8j-l,q-s). These results suggested that GPI anchoring is required to both inhibit

free Bnl secretion/dispersion and activate its directed contact-dependent release.

Bnl's GPI-anchor is required for organized signaling patterns.

To investigate if GPI anchoring and contact-dependent release are important for Bnl

signaling, we generated small (2-3 cell) gain-of-function (GOF) clones expressing GPI-

modified and non-GPI-modified Bnl:GFP variants directly within the ASP epithelium, as

described earlier [9]. To distinguish the ectopically induced signaling patterns from the

endogenous signaling patterns, we analyzed clones within the ASP stalk and transverse

connective (TC), which lack Bnl uptake from the original disc source and MAPK

signaling [9] (Supplementary Figure 7d). In consistence with earlier reports [9], all cells

within 3 cell diameter area surrounding a Bnl:GFP GOF clone received Bnl:GFP, and all

Bnl:GFP-receiving cells also induced dpERK (Fig.9a,a',d; Supplementary Figure 7g). In

comparison, Bnl:GFPΔC was received by many cells surrounding its clonal source, but

only a few randomly located Bnl:GFPΔC-receiving cells induced dpERK (Fig.9b,b',d;

Supplementary Figure 7e,g).

Apparently, the normal spatial correlation between signal uptake and signaling was lost

with Bnl:GFPΔC. The correlation of signal uptake and signaling was regained with

Bnl:GFPΔC-TM. However, Bnl:GFPΔC-TM activity was restricted to only a few source-

juxtaposed ASP cells (Fig.9c,d; Supplementary Figure 7f,g). Similarly, when either

Bnl:GFP, Bnl:GFPΔC, or Bnl:GFPΔC-TM was overexpressed from the disc *bnl*-source,

unlike Bnl:GFP or Bnl:GFPΔC-TM, a significant number of Bnl:GFPΔC-receiving ASP

cells lacked nuclear MAPK signaling (Fig.9d; Supplementary Figure 7h-l). These results

suggested that GPI anchoring and contact-dependent Bnl release are required for the

normal coordination between signal dispersion and interpretation.

Bnl was also known to chemoattract tracheal migration toward its source [47,48]. To further

assess the morphogenetic potency of Bnl variants, we examined their ability to

chemoattract tracheal branches to an ectopic expressing source, such as larval salivary

gland, a non-essential, trachea-free organ, which normally does not express *bnl*[36,48].

bnl-Gal4 was reported to be non-specifically expressed in the salivary glands, and *bnl-*

Gal4-driven Bnl expression in the salivary glands induced tracheal invasion into this

trachea-free organ [36]. Therefore, we expressed comparable levels of Bnl:GFP,

Bnl:GFPΔC, and Bnl:GFPΔC-TM in salivary glands under the *bnl-Gal4* control (see

Methods and Supplementary Information). Strikingly, both Bnl:GFP and Bnl:GFPΔC-TM

induced extensive tracheal invasion and branching into the expressing salivary gland,

but Bnl:GFPΔC did not (Fig.9e-h).

Source surface distribution of Bnl was known to attract tracheal branching [36,49].

Therefore, we predicted that the extent of tracheal invasion/branching might be

correlated with the levels of Bnl variants available on the source surface. Although all

three constructs were expressed at an equivalent level under *bnl-Gal4*, the level of

extra-cellular Bnl:GFPΔC on the salivary gland surface was significantly less than that of

Bnl:GFP or Bnl:GFPΔC-TM (Fig.9i-l). Thus, although Bnl:GFPΔC can activate receptors,

its unavailability on the source cell surface caused the loss of its ability to guide tracheal

branching into its source. In contrast, the same signal, when was tethered to the source

membrane by a TM-domain (i.e., Bnl:GFPΔC-TM), its chemotactic function was

regained. These results suggested that Bnl retention on the source surface is critical for

its morphogenetic potency.

DISCUSSION

This study uncovered an elegant program of reciprocal inter-organ communication that

is encoded by the lipid-modification of FGF/Bnl and orchestrated by cytoneme-mediated

contact-dependent signaling. It provided mechanistic insights into how cytonemes find a

specific target to establish signaling contacts, exchange signals at their contact sites,

and inform cells where they are, what they should do, and when. It also discovered the

first lipid-modified FGF and characterized how lipidation enables the signal to program

cytoneme-mediated signaling to self-regulate its diverse tissue-specific dispersion and
interpretation.
610
We discovered that Bnl is GPI anchored to the source cell surface, and this modification
facilitates the signal to simultaneously act as a local CAM and a long-range paracrine
morphogen/growth factor. Based on our results, mechanistically, GPI-anchored Bnl
directs three cellular events, as summarized in Figure 10a. These include - cytoneme-
mediated target selection, target-specific signal release, and feedback regulations of
these processes. GPI-anchored Bnl on the source cell surface and Btl on the receiving
cell surface act as heterophilic CAMs. Bnl source and recipient cells extend cytonemes
to present Bnl and Btl on their surfaces, and when Btl-Bnl binding occurs, it leads to
selective cell-cell cytoneme contact formation over distance. The CAM-like Btl-Bnl
binding induces forward signaling in the ASP and retrograde signaling in the source,
responses of which promote these cells to reinforce the polarity of Bnl-receiving and
Bnl-sending cytonemes toward each other. This explains how the CAM-like bidirectional
Btl-Bnl signaling might drive reciprocal guidance of source and ASP cytonemes and
ensure target-specific cytoneme contact formation. However, the purpose of this CAM-
like function of Bnl is to promote its subsequent release through cytoneme contacts in a
target-specific manner. Bnl release promotes receptor-mediated endocytosis and
activation of MAPK signaling asymmetrically only in the source-adhering recipient cells.
628
Traditionally, secreted signals are presumed to signal unidirectionally, only in recipient
cells, by activating transcription of cascades of target genes that, in turn, control various
functions. In this model, signaling ligands themselves do not physically shape
cells/tissues, but control gene expression required for morphogenesis. In contrast to this
general idea, our results provided evidence that lipidated Bnl acts as a CAM and can
directly control cell/cytoneme polarity by inducing contact-dependent bidirectional
signaling to instruct both source and recipient cells. Thus, GPI-anchoring can enable
Bnl to simultaneously act as both a ligand and a receptor for Btl and transmit
information inside-out and outside-in across the source cell membrane by binding to Btl.
Although our results provided evidence for the contact-mediated response of a reverse

Btl-Bnl signaling in the source, the components of the intracellular pathway that induces

source cells to extend polarized cytonemes toward the ASP are unknown. Based on our

results, we predict that the CAM-like Btl-Bnl signaling might control local cytoskeletal re-

organization required to produce polarized cytonemes. The crosslinking of GPI-APs on

the outer cell surface are known to induce polarized congregation of GPI-APs in

membrane microdomains and this dynamic activity on the outer cell surface is known to

generate a mechanical cue across cell membrane that can transiently reorganize

actomyosin structures in the inner membrane leaflet [24,27]. However, the possibility of

such mechanochemical signaling feedback [50] at the cytoneme-contact sites needs to be

investigated in the future.

The contact-dependent bidirectional signaling via cytonemes is reminiscent of synaptic

communication in neurons. Filopodia-mediated bidirectional matchmaking for synapse

has been reported in *Drosophila* neuromuscular junctions [32]. Bidirectional transmission

of information is the basis for neuronal synaptic assembly, plasticity, and functions. An

important implication of the bidirectional Btl-Bnl signaling is that the cause and effect of

the signaling process become interdependent. For instance, the same cytoneme

contacts that the Btl-Bnl binding helps to form also bring Btl and Bnl molecules together

to interact. Consequently, not only is the signal exchange cytoneme/contact-dependent,

but the cytoneme contacts are also formed signal- or tissue-specifically.

Free secretion and dispersion of paracrine signals are presumed to be required for long-

range morphogen-like signaling. In this general paradigm, source surface retention is

inhibitory to long-range dispersion and activity. In contrast, we found that surface

localized display of Bnl is critical for its morphogenetic potency. This is consistent with

our previous finding showing that the Bnl availability on the source surface controls the

levels and range of Bnl signaling in the ASP [9,36]. How signal retention might facilitate

recipient-specific long-range patterning can be explained by our previous findings of

self-generation of cytoneme-mediated signaling by Bnl-signaling feedbacks [9,36].

Although Bnl release is critical for its long-range morphogen like functions, we do not

know how GPI-anchored Bnl is released from the source membrane and how the

release mechanism is specifically activated at the cytoneme contact sites. We speculate
that an enzymatic shedding [51] of Bnl might be activated at the cytoneme contact sites.
672
Thus, GPI-anchored Bnl apparently provides a balance between two functions -
free/random secretion and CAM-like membrane-anchored display (i.e., inhibition of
secretion). This dual strategy of inhibition and activation of signal release can induce
diverse context-specific morphogenic outcomes via contact-dependent signaling. For
instance, we showed that Bnl's CAM-like cell surface display (e.g., GPI-anchored Bnl or
TM-tethered BnlΔC) is critical for inducing tracheal chemotaxis toward the Bnl source as
well as the reciprocal organization of source and recipient cells via cytoneme-cytoneme
contacts. Whereas the Bnl release through cytoneme contacts can produce diverse
recipient-specific dispersion and signaling patterns [9,36] . Bnl release also can dissolve
the inter-cytoneme adhesive contacts. Dissociation of cytoneme:cytoneme contacts
might be required for dynamic context-specific adaptation of growing tracheal branches.
Repeated contact association and dissociation might ensure the reciprocal guidance of
the Bnl-expressing embryonic cells in synchrony and close association with the growing
tips of each tracheal branch [52].
687
Bnl's CAM-like local and morphogen-like global organizing functions are likely to be
interdependent. For instance, previously, we showed that the Bnl's morphogen-like
transcriptional responses in the ASP maintain steady-state gradients of the Btl
expression and the numbers and polarity of Bnl-receiving ASP cytonemes [9,36]. On the
other hand, here we found that the CAM activity of Bnl is a prerequisite for its
subsequent target-specific release and morphogen-like roles. Consequently, a readily
secreted/dispersed non-GPI Bnl, although can activate receptors, fails to induce both
CAM-like local and morphogen-like large-scale organization (Fig.10b). Whereas, TM-
tethered Bnl efficiently acts as a CAM, but due to its poor release from the source,
causes only a scaled-down ASP patterning within a narrow range (Fig.10c).
698
The membrane association and the dual strategy of selective inhibition and activation of
signal release might also be present in other signals. For instance, a *Drosophila* FGF,

Pyramus (Pyr) is a transmembrane protein, and the TM-tethering is predicted to be

required for its spatiotemporal functions [53]. Similar to Bnl, Ephrins are GPI-/TM-tethered

signals, and their membrane tethering causes contact-dependent bidirectional

signaling[22]. Lipid modifications are critical for the activity of Hh, Wnt, and EGF/Spi

[22,42,54,55]. Analogous to Bnl, engineered TM-tethered forms of Hh, Spi, and Wnt can

induce morphogenesis within a narrow range, and removal of lipid-modification and

unrestricted spreading of non-lipidated Hh, Spi, and Wnt reduce their tissue organizing

potency [42,56–60]. Moreover, all signals, including those that are not known to be lipidated

(e.g., BMPs and many FGFs), can interact with membrane-anchored proteoglycans,

which can restrict free signal dispersion and induce biphasic signaling activation and

inhibition [61–63]. Glypicans also can control cytoneme stability [64–66]. Therefore, our

findings showing how GPI anchoring enables Bnl to direct source and recipient cells to

reciprocally coordinate with each other by cytonemes provide important new insights

into how other signal retention strategies might control signaling and morphogenesis.

716
METHODS

Fly genetics

All fly lines and their sources are described in Supplementary Table 4. Flies were raised

at 25 °C with a 12 h/12 h light/dark cycle, except tracheal *dia-RNAi* expression. All the

experiments were performed under non-crowded situations. The sequence-verified DNA

constructs were used to generate transgenic flies by P-element mediated germline

transformation as described in Du et al [9]. Transgenic injections were performed by

Rainbow Transgenic Flies, Inc.

Mosaic analyses

(i) To generate ectopic clones in the ASP, *hsFlp; btlenh>y+>Gal4,btlenh-mRFPmoe*

females were crossed to males carrying *UAS-Bnl:GFP, UAS-Bnl:GFP∆C, UAS-*

Bnl:GFP∆C-TM, or *UAS-Btl-DN.* Flip-out clones were generated by heat shocking early

3rd instar larvae at 37 C foreither 5 or 10 min. arvae, then , were incubated at 25 C until

they reached the mid-late 3rd instar stages and dissected for further analysis.

(ii) To generate CD8:GFP-expressing clones in the *bnl* source, *hs-mFlp;bnlGal4*

females were crossed to FlyBow *FB2.0* flies (see Supplementary Table 4) and clones

were induced in the progenies by heat-shock. Only CD8:GFP-marked cells were

visualized in live tissues.

(iii) Ectopic Bnl:GFP-TM-expressing clones in the wing disc were induced in progenies

of *hs-Flp;UAS-bnl:GFP-TM* (females) x *mCherryCAAX;act>CD2>Gal4* (males) cross.

iv) Ectopic Btl:GFP-expressing clones in the wing disc were induced in progenies of *hs-*

Flp;;UAS-Btl:GFP (females) x *act>CD2>Gal4;;bnlLexA,LexO-mCherryCAAX* (males)

cross.

Tissue-specific transgene expression

For the transgene expression in the trachea, *btl-Gal4/UAS* or *btl-LexA/LexO* systems

were used. To express transgenes in the wing disc *bnl*-source, *bnl-Gal4/UAS* or *bnl-*

LexA/LexO systems were used. Comparable levels of *bnl-Gal4*-driven expression of

GPI-modified Bnl:GFP and non-GPI modified Bnl:GFPΔC and Bnl:GFPΔC-TM were

determined as described in the Supplementary Information (Supplementary Notes,

section C). Although *bnl* is not expressed in the salivary gland, *bnl-Gal4* is non-

specifically expressed in the larval salivary gland [37]. Therefore, *bnl-Gal4* was used to

ectopically express Bnl:GFP variants in the larval salivary glands. Thus, phenotypic

consequences of *bnl-Gal4*-driven expression of Bnl:GFP variants were recorded in two

distinct tissue contexts of the same larva: wing disc (for native Bnl source and ASP

interactions) and salivary glands (for ectopic source and tracheal invasion into the

ectopic source).

Cytoneme removal from the ASP and *bnl*-source

To remove source cytonemes, *UAS-dia-RNAi* was expressed under *bnlGal4* and larvae

were reared at 25°C. In trachea, a high-level *dia-RNAi* expression (at 25°C) caused

larval lethality. Therefore, *tub-Gal80^{ts}; UAS-diaRNAi* males were crossed to

btlGal4,UAS-CD8:GFP; bnlLexA,LexO-mCherryCAAX/TM6 females; The *btl-Gal4-*

driven expression of *dia-RNAi* was suppressed by *Gal80^{ts}* at 18°C until L3 stage and

activated by shifting the temperature to 29°C (that inactivated *Gal80^ts^*), 24 hr prior to

harvesting the L3 larvae for imaging.

Cell lines and cell culture

Schneider's 2 (S2) cells (S2-DGRC) were cultured and transfected following standard

protocols [36]. Cells were transfected either with Lipofectamine 3000 or Mirus TransIT-

Insect Transfection Reagent for CAM assays following manufacturer's protocol.

Transient ectopic expression of various constructs in S2 cells was achieved by co-

transfecting *act-Gal4* and *UAS-x* constructs (x = various cDNA or cDNA fusions) and

analyzed after 48 hrs of incubation at 25°C.

Immunohistochemistry

773 The standard immunostaining and the extracellular immunostaining under live-cell non-

774 permeabilized condition (αGFP^{ex} for GFP or αBnl^{ex} for Bnl) was carried out following

775 standard protocols [9,36]. Supplementary Table 4 lists all antibodies and dilutions used.

DNA constructs

778 All constructs generated and used here are described in Supplementary Table 4.

Bioinformatic analysis

781 DNA sequences were analyzed with SnapGene, Protein sequences were analyzed with

782 MacVector, ProtScale (ExPASy), EMBOSS Pepinfo (www.ebi.ac.uk), PredGPI

783 (http://gpcr.biocomp.unibo.it/predgpi).

Flow cytometric analyses

S2 cells expressing various constructs were immunostained and scanned using a BD

CantoII (BD Biosciences) flow cytometer and the data were analyzed using FACSDiva

(BD Biosciences). For quantitative assays as shown in Supplementary Figure 3b-h,

Supplementary Figure 4e-g, the number of cells detected in Q2 (GFP+ cells with

αGFP^{ex}+) was divided by the number of cells in either Q2 or Q4 (total GFP+ cells) to

obtain the Y-axis value. These values were obtained from three independent

experimental repeats. An example of the gating strategy for FACS analyses is shown in

the Supplementary Information (Supplementary Note, section D).

Ex vivo organ culture and Furin inhibition

Ex vivo wing disc culture in WM1 media, pharmacological inhibition of Furin in cultured

discs, and analyses of ASP-specific uptake of Bnl were carried out following standard

protocols described in Sohr et al. [36,67]. In brief, late third instar larval tissues were *ex*

vivo cultured in 2 ml of WM1 medium in the presence or absence of a cocktail of Furin

inhibitor I and II (50 μM final concentration each; Calbiochem; 344930 and 344931).

Cultured discs were removed from a single pool of culture-media after 0, 1, 2.5, and 5h

of incubation at 25°C, followed by fixation and αHA immunostaining of the tissues. The

temporal increase in the levels of GFP-tagged Bnl in the ASP over time was difficult to

assess due to the pre-existing Bnl:GFP$_3$ in the L3 ASP used for culturing. Therefore,

Furin-sensors (HA$_1$Bnl:GFP$_3$, HA$_1$Bnl:GFP$_3\Delta$C-TM, and HA$_1$Bnl:GFP$_3\Delta$C) that were

detectable by both αHA immunostaining and GFP were used. The time when tissues

were transferred to the Furin-inhibited media, was considered as t=0 for the appearance

of intact Furin sensors (HA$_1$Bnl:GFP$_3$). For a comparative analyses among samples, a

semi-quantitative estimate was obtained by measuring the ratio of the uncleaved sensor

(αHA immunofluorescence intensity) to the total GFP signal (pre-existing Bnl:GFP$_3$ +

post-inhibition HA$_1$Bnl:GFP$_3$) per ASP for t=1h or 2.5h or 5 h.

CAM assay using S2 cells

S2 cells ectopically expressing either Btl variants (*UAS-Btl:Cherry, -BtlDN:Cherry*, or -

sBtl:Cherry) or ligand variants (*UAS-Bnl:GFP* or *-Bnl:GFPΔC* or *-Bnl:GFPΔC-TM*) (48 h

after transfection) were resuspended in 1 ml of fresh M3 media. 200 μl of the receptor-

expressing cells was gently mixed with 200 μl of the ligand-expressing cells for 10 min

in a sterile tube. The well-mixed cell suspension was plated to the center of a sterile

cover slip within a 6-well plate and incubated at 25°C for 16 hrs before fixing them with

4% PFA following standard protocols. Coverslips were carefully mounted with cells

facing down to 10 μl of the VECTASHIELD on microscopic slides. For comparative

analyses, co-culture assays were performed in identical conditions. Cells were analyzed

from more than three transfection repeats, with at least 30 random frames/experiment

under 20X and 40X objectives. Regions with comparable cell density were analyzed.

Adjacent cells with the ring-like heterophilic receptor-ligand co-clusters were considered

as trans-paired cells and those without the receptor-ligand co-clusters were considered

as juxtaposed. Homophilic Btl-Btl or Bnl-Bnl clusters between adjacent cells were rarely

observed as indicated in Supplementary Figure 2h. Cells were imaged in both 20X and

40X to thoroughly verify Btl-Bnl trans-pairing in the mixed cell population.

Autocrine and paracrine Bnl-Btl signaling in S2 cells

For autonomous MAPK signaling, S2 cells were co-transfected with *act-Gal4*, *UAS-*

Btl:Cherry, and *UAS-X* (*X* = various Bnl:GFP variants) and prepared on cover-slips as

described before. Cover slips with cells were processed with standard fixation and anti-

dpERK immunostaining. The percentage of Btl:Cherry expressing cells with nuclear

dpERK signals was scored with confocal microscope (20X/40X). For non-autonomous

dpERK signaling, cells were prepared following the CAM assay, followed by PFA

fixation and anti-dpERK staining. Both trans-paired and unpaired Btl:Cherry variants

were recorded to estimate the contact-dependent non-autonomous dpERK signaling.

PIPLC treatment of transfected S2 cells and wing imaginal discs

Transfected S2 cells (1 mL) were harvested (700 g, 5 min) in a 1.5 mL Eppendorf tube.

Cells were washed twice in 1XPBS (500 μL each) and incubated either in 500 μL

1 PBS (control) or in PIP C containing 1 PBS solution (1 U m PIP C) at 2 -25 C for

30 min with gentle rotation. Cells were harvested and prepared for the standard non-

permeabilized extracellular staining before imaging or FACS. To reliably compare the

levels of surface-localized proteins with and without PIPLC treatment, the ratio of the

surface:total Bnl levels per cell was measured using Fiji (at least 3 independent repeats).

Note that the surface levels of GFP-tagged proteins per cell was measured with αGFP^{ex}

immuno-fluorescence and the total GFP fluorescence of the same protein measured the

total expression in the same cell. For untagged Bnl, CD8:GFP was co-transfected and

the Bnl^{ex} level was normalized with CD8:GFP in the same cell.

For PIPLC assay in wing discs, third instar larvae were prepared following ex-vivo organ

culture method [36] and transferred to 1.5 ml Eppendorf tubes containing 1 ml of either

WM1 media (control) or WM1 media with PIPLC (1U/mL). Tissues were incubated for

30 min at 20-25°C with gentle rotation. Then the PIPLC reaction was stopped by

removing the solution and washing the tissues 3 times with WM1 media. Tissues were

then prepared for extracellular staining as described before.

Live imaging of cytonemes

Wing imaginal discs were prepared and imaged in WM1 medium as described in Du et

al. [9]. Time-lapse imaging of cytonemes was carried out in ex vivo cultured wing discs in

Grace's insect culture medium as described in Barbosa and Kornberg [68]. A spinning

disc confocal microscope under 40X/60X magnifications was used to capture ~30-50

μm Z-sections with 0.2 μm step size of wing discs. For Figure 1e,e', images were

captured using the Zeiss LSM900 confocal with an Airyscan-2 detector in 60X

magnifications. The images were processed and analyzed with Fiji. For 3D-rendering,

Andor iQ3 and Imaris software were used.

Quantitative analyses of cytoneme number, orientation, and dynamics

Cyonemes were manually counted and plotted by methods described in Du et al [9]. For

ASP cytonemes, cytonemes were recorded across a 100 μm arc centered at the tip

(Figs.2i-k,l,m,m',n; 6d-l; Supplementary Figure 6g). Wing disc *bnl* source cytonemes

were recorded from the 3D projections across a 100 μm perimeter surface centering at

the ASP tip contact as a reference (Figs. 2a-e,l',m'',m''',n'; 6e-l; Supplementary Figure

6g). For Figure 2l',n', cytonemes were not grouped based on the length as all source

cytonemes were less than 15 μm. For Figures 1l-n,7d-e, different parameters of

cytoneme dynamics were measured following previous reports (see Supplementary

Table 1) [65,69].

Quantitative analyses of fluorescence intensities in tissues

For intracellular and extracellular surface Bnl levels, all fluorescent intensity

measurements were background corrected. The density of fluorescence intensity (e.g.,

spatial range and density of signals) was measured from maximum-intensity projections

encompassing the wing disc, ASP, or salivary gland sections from a selected region of

interest (ROI) using Fiji. For each genotype, at least 3 samples were used to obtain the

average plot profile. Quantitative estimates of levels of Bnl:GFP variants and signaling

outcomes are normalized with internal controls to avoid variations among samples. For

example, to compare between Bnl variants, we compared the ratio of surface levels of

each protein (red, anti-GFP non-permeabilized immunofluorescence) to total expression

(total GFP fluorescence) in the same ROI of wing disc source (Fig.5) and the salivary

glands (Fig. 9i-l). Similarly, to assess MAPK signaling patterns of different Bnl variants

(Fig. 9a-d), we measured the percentage of signal recipient cells (cells with Bnl:GFP

variant puncta) that induced MAPK. The correlated patterns between signal reception

and signaling per cell/tissue were then compared between conditions.

Sholl analysis for tracheation in salivary gland

The extent and frequency of tracheal branching on the larval salivary glands expressing

equivalent levels of Bnl:GFP, Bnl:GFPΔC-TM, or Bnl:GFPΔGPI was quantitated using

Sholl analysis in Fiji as described in [36]. The analysis created 20 concentric circles in

increments of 5-μm radius from the point of origin up to 100 μm and counted the

number of times any tracheal branch crossed these circles. These values were

averaged across multiple samples and compared between the different Bnl variants

expressed in the salivary gland. For Fig. 9h total number of branch interconnection

within 100 μm radius, $p < 0.01$: TM (n = 5) vs either Bnl:GFP (n = 6) or ΔC (n = 4), $p <$

0.05: Bnl:GFP vs ΔC; p values were calculated using one way-ANOVA followed by

Tukey's honestly significant different test.

Statistics and Reproducibility

Statistical analyses were performed using VassarStat and GraphPad Prism 8, MS

Excel. P values were determined using the unpaired two-tailed t-test for pair-wise

comparisons, or the one-way ANOVA followed by Tukey's honestly significant different

(HSD) test for comparison of multiple groups. $p < 0.05$ is considered significant. All

experimental results were analyzed from at least three independent experiments. The

sample size (n) for each data analysis is indicated in the figures/figure

legends/methods/source data. Graphs in Figure 4f and g show intensity analyses from

randomly selected cells from a large pool of cells from three experimental repeats. All

cells showed consistent patterns. The same results were confirmed using FACS

analyses of the same cell populations (Supplementary Figure 3b-h). For Figure 4f,

number of randomly selected cell images analyzed for intensity: Bnl + CD8:GFP- control

(n=14) & PIPLC (n=14); GFP-GPI - control (n=9) & PIPLC (n=8); cSpi:GFP - control

(n=14) & PIPLC (n=14); Bnl:GFP - control (n=14) & PIPLC (n=12); bGFP-GPI - control

(n=13) & PIPLC (n=11); Bnl:GFPΔC-TM - control (n=9) & PIPLC (n=10). For Figure 4g,

number of randomly selected cell images analyzed for intensity - Bnl:GFP (n=14),

Bnl:GFP-ω^m (n=16) and Bnl:GFPΔC (n=9). Rose plots were generated by R software as

described in [9].

RNA isolation and RT-PCR

929 Total RNA was extracted from 20 wing discs of the w^{1118} L3 larvae using TRI reagent

930 (Sigma-Aldrich) followed by Direct-zol RNA purification kits (Zymo Research).

931 Expression analyses of *bnl PA* and *PC* isoforms are described in Supplementary

932 Information.

DATA AVAILABILITY

935 All data generated and analyzed are included in the manuscript and supporting files.

936 Source data are provided with this paper.

CODE AVAILABILITY

939 The code for R plots is provided in the Supplementary Information.

940
941

REFERENCES

1. Shilo, B.-Z. New Twists in Drosophila Cell Signaling*. *J Biol Chem* 291, 7805–7808 (2016).

2. Perrimon, N., Pitsouli, C. & Shilo, B.-Z. Signaling mechanisms controlling cell fate and embryonic patterning. *Cold Spring Harbor Perspectives in Biology* 4, a005975–a005975 (2012).

3. Housden, B. E. & Perrimon, N. Spatial and temporal organization of signaling pathways. *Trends Biochem Sci* 39, 457–464 (2014).

4. Zhang, C. & Scholpp, S. Cytonemes in development. *Curr Opin Genet Dev* 57, 25–30 (2019).

5. Roy, S., Huang, H., Liu, S. & Kornberg, T. B. Cytoneme-mediated contact-dependent transport of the Drosophila decapentaplegic signaling protein. *Science (New York, N.Y.)* 343, 1244624 (2014).

6. Bischoff, M. *et al.* Cytonemes are required for the establishment of a normal Hedgehog morphogen gradient in Drosophila epithelia. *Nat Cell Biol* 15, 10.1038/ncb2856 (2013).

7. González-Méndez, L., Gradilla, A.-C. & Guerrero, I. The cytoneme connection: direct long-distance signal transfer during development. *Development* 146, dev174607 (2019).

8. Kornberg, T. B. & Roy, S. Communicating by touch--neurons are not alone. *Trends in cell biology* 24, 370–376 (2014).

9. Du, L., Sohr, A., Yan, G. & Roy, S. Feedback regulation of cytoneme-mediated transport shapes a tissue-specific FGF morphogen gradient. (2018).

10. Chen, W., Huang, H., Hatori, R. & Kornberg, T. B. Essential basal cytonemes take up Hedgehog in the Drosophila wing imaginal disc. *Development (Cambridge, England)* 144, 3134–3144 (2017).

11. Hall, E. T. *et al.* Cytoneme delivery of Sonic Hedgehog from ligand-producing cells requires Myosin 10 and a Dispatched-BOC/CDON co-receptor complex. *eLife* 10, (2021).

12. Huang, H. & Kornberg, T. B. Myoblast cytonemes mediate Wg signaling from the wing imaginal disc and Delta-Notch signaling to the air sac primordium. *Elife* 4, e06114 (2015).

13. Roy, S., Hsiung, F. & Kornberg, T. B. Specificity of Drosophila Cytonemes for
Distinct Signaling Pathways. *Science (New York, N.Y.)* 332, 354–358 (2011).

14. Stanganello, E. *et al.* Filopodia-based Wnt transport during vertebrate tissue
patterning. *Nat Commun* 6, 5846 (2015).

15. Sanders, T. A., Llagostera, E. & Barna, M. Specialized filopodia direct long-range
transport of SHH during vertebrate tissue patterning. *Nature* 497, 628–632 (2013).

16. Inaba, M., Buszczak, M. & Yamashita, Y. M. Nanotubes mediate niche-stem cell
signaling in the Drosophila testis. *Nature* 523, 329–332 (2015).

17. Fereres, S., Hatori, R., Hatori, M. & Kornberg, T. B. Cytoneme-mediated signaling
essential for tumorigenesis. *Plos Genet* 15, e1008415 (2019).

18. Krupke, O. A., Zysk, I., Mellott, D. O. & Burke, R. D. Eph and Ephrin function in
dispersal and epithelial insertion of pigmented immunocytes in sea urchin embryos.
Elife 5, e16000 (2016).

19. Sato, M. & Kornberg, T. B. FGF Is an Essential Mitogen and Chemoattractant for
the Air Sacs of the Drosophila Tracheal System. *Dev Cell* 3, 195–207 (2002).

20. Sutherland, D., Samakovlis, C. & Krasnow, M. A. branchless encodes a Drosophila
FGF homolog that controls tracheal cell migration and the pattern of branching. *Cell* 87,
1091–1101 (1996).

21. Klämbt, C., Glazer, L. & Shilo, B. Z. breathless, a Drosophila FGF receptor
homolog, is essential for migration of tracheal and specific midline glial cells. *Gene Dev*
6, 1668–1678 (1992).

22. Egea, J. & Klein, R. Bidirectional Eph–ephrin signaling during axon guidance.
Trends in cell biology 17, 230–238 (2007).

23. Hynes, R. O. Integrins: bidirectional, allosteric signaling machines. *Cell* 110, 673–
687 (2002).

24. Kalappurakkal, J. M. *et al.* Integrin Mechano-chemical Signaling Generates Plasma
Membrane Nanodomains that Promote Cell Spreading. *Cell* 177, 1738-1756.e23
(2019).

25. Saha, S., Anilkumar, A. A. & Mayor, S. GPI-anchored protein organization and
dynamics at the cell surface. *J Lipid Res* 57, 159–175 (2016).

26. Gowrishankar, K. *et al.* Active Remodeling of Cortical Actin Regulates
Spatiotemporal Organization of Cell Surface Molecules. *Cell* 149, 1353–1367 (2012).

27. Raghupathy, R. *et al.* Transbilayer Lipid Interactions Mediate Nanoclustering of
Lipid-Anchored Proteins. *Cell* 161, 581–594 (2015).

28. Um, J. W. & Ko, J. Neural Glycosylphosphatidylinositol-Anchored Proteins in
Synaptic Specification. *Trends in cell biology* 27, 931–945 (2017).

29. Schwank, G. *et al.* Formation of the long range Dpp morphogen gradient. *PLoS*
Biology 9, e1001111 (2011).

30. González-Méndez, L., Seijo-Barandiarán, I. & Guerrero, I. Cytoneme-mediated cell-
cell contacts for Hedgehog reception. *Elife* 6, e24045 (2017).

31. Mattes, B. & Scholpp, S. Emerging role of contact-mediated cell communication in
tissue development and diseases. *Histochemistry and cell biology* 150, 431–442 (2018).

32. Kohsaka, H. & Nose, A. Target recognition at the tips of postsynaptic filopodia:
accumulation and function of Capricious. *Development (Cambridge, England)* 136,
1127–1135 (2009).

33. Sanes, J. R. & Zipursky, S. L. Synaptic Specificity, Recognition Molecules, and
Assembly of Neural Circuits. *Cell* 181, 1434–1435 (2020).

34. Gilmour, D., Rembold, M. & Leptin, M. From morphogen to morphogenesis and
back. *Nature* 541, 311–320 (2017).

35. Dustin, M. L. The Immunological Synapse. *Cancer Immunol Res* 2, 1023–1033
(2014).

36. Sohr, A., Du, L., Wang, R., Lin, L. & Roy, S. Drosophila FGF cleavage is required
for efficient intracellular sorting and intercellular dispersal. *J Cell Biol* 218, 1653–1669
(2019).

37. Kinoshita, T. & Fujita, M. Biosynthesis of GPI-anchored proteins: special emphasis
on GPI lipid remodeling. *Journal of Lipid Research* 57, 6–24 (2016).

38. Zurzolo, C. & Simons, K. Glycosylphosphatidylinositol-anchored proteins:
Membrane organization and transport. *Biochimica Et Biophysica Acta Bba - Biomembr*
1858, 632–639 (2016).

39. Caras, I. W. An internally positioned signal can direct attachment of a
glycophospholipid membrane anchor. *J Cell Biology* 113, 77–85 (1991).

40. Ferguson, M. A., Kinoshita, T. & Hart, G. W. *Glycosylphosphatidylinositol Anchors*.
(Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press, 2015). doi:doi:
10.1101/glycobiology.3e.012.

41. Greco, V., Hannus, M. & Eaton, S. Argosomes: a potential vehicle for the spread of
morphogens through epithelia. *Cell* 106, 633–645 (2001).

42. Miura, G. I. *et al.* Palmitoylation of the EGFR ligand Spitz by Rasp increases Spitz
activity by restricting its diffusion. *Developmental Cell* 10, 167–176 (2006).

43. Yan, D. & Lin, X. Drosophila glypican Dally-like acts in FGF-receiving cells to
modulate FGF signaling during tracheal morphogenesis. *Dev Biol* 312, 203–216 (2007).

44. Böhme, U. & Cross, G. A. M. Mutational analysis of the variant surface glycoprotein
GPI-anchor signal sequence in Trypanosoma brucei. *J Cell Sci* 115, 805–16 (2002).

45. Delahunty, M. D., Stafford, F. J., Yuan, L. C., Shaz, D. & Bonifacino, J. S.
Uncleaved signals for glycosylphosphatidylinositol anchoring cause retention of
precursor proteins in the endoplasmic reticulum. *J Biol Chem* 268, 12017–12027
(1993).

46. Incardona, J. P. & Rosenberry, T. L. Replacement of the glycoinositol phospholipid
anchor of Drosophila acetylcholinesterase with a transmembrane domain does not alter
sorting in neurons and epithelia but results in behavioral defects. *Mol Biol Cell* 7, 613–
630 (1996).

47. Ochoa-Espinosa, A. & Affolter, M. Branching Morphogenesis: From Cells to Organs
and Back. *Csh Perspect Biol* 4, a008243 (2012).

48. Jarecki, J., Johnson, E. & Krasnow, M. A. Oxygen Regulation of Airway Branching
in Drosophila Is Mediated by Branchless FGF. *Cell* 99, 211–220 (1999).

49. Peterson, S. J. & Krasnow, M. A. Subcellular Trafficking of FGF Controls Tracheal
Invasion of Drosophila Flight Muscle. *Cell* 160, 313–323 (2015).

50. Gilmour, D., Rembold, M. & Leptin, M. From morphogen to morphogenesis and
back. *Nature* 541, 311–320 (2017).

51. Lichtenthaler, S. F. Sheddase Gets Guidance. *Science (New York, N.Y.)* 335, 179–
180 (2012).

52. Du, L. *et al.* Unique patterns of organization and migration of FGF-expressing cells
during Drosophila morphogenesis. *Dev Biol* 427, 35–48 (2017).

53. Stepanik, V., Sun, J. & Stathopoulos, A. FGF Pyramus Has a Transmembrane
Domain and Cell-Autonomous Function in Polarity. *Current biology : CB* (2020).

54. Lee, J. J. *et al.* Autoproteolysis in hedgehog Protein Biogenesis. *Science* 266,
1528–1537 (1994).

55. Willert, K. *et al.* Wnt proteins are lipid-modified and can act as stem cell growth
factors. *Nature* 423, 448–452 (2003).

56. Taylor, F. R. *et al.* Enhanced potency of human Sonic hedgehog by hydrophobic
modification. *Biochemistry* 40, 4359–4371 (2001).

57. Gallet, A., Ruel, L., Staccini- ave nant, . Th rond, P. P. Cholesterol modification
is necessary for controlled planar long-range activity of Hedgehog in Drosophila
epithelia. *Development* 133, 407–418 (2006).

58. Callejo, A., Torroja, C., Quijada, L. & Guerrero, I. Hedgehog lipid modifications are
required for Hedgehog stabilization in the extracellular matrix. *Development* 133, 471–
483 (2006).

59. Speer, K. F. *et al.* Non-acylated Wnts Can Promote Signaling. *CellReports* 26, 875-
883.e5 (2019).

60. Alexandre, C., Baena-Lopez, A. & Vincent, J.-P. Patterning and growth control by
membrane-tethered Wingless. *Nature* 505, 180–185 (2014).

61. Yan, D. & Lin, X. Shaping morphogen gradients by proteoglycans. *Cold Spring
Harbor Perspectives in Biology* 1, a002493–a002493 (2009).

62. Dejima, K., Kanai, M. I., Akiyama, T., Levings, D. C. & Nakato, H. Novel contact-
dependent bone morphogenetic protein (BMP) signaling mediated by heparan sulfate
proteoglycans. *The Journal of biological chemistry* 286, 17103–17111 (2011).

63. Balasubramanian, R. & Zhang, X. Mechanisms of FGF gradient formation during
embryogenesis. *Semin Cell Dev Biol* 53, 94–100 (2016).

64. Huang, H. & Kornberg, T. B. Cells must express components of the planar cell
polarity system and extracellular matrix to support cytonemes. *Elife* 5, e18979 (2016).

65. Gonz lez- endez, ., Seijo -Barandia an, I. Guer rero, I. Cytoneme-mediated cell-
cell contacts for Hedgehog reception. *eLife* 6, 605 (2017).

66. Simon, E. *et al.* Glypicans define unique roles for the Hedgehog co-receptors boi
and ihog in cytoneme-mediated gradient formation. *Elife* 10, e64581 (2021).

67. Sohr, A., Du, L. & Roy, S. Ex vivo Drosophila Wing Imaginal Disc Culture and Furin
Inhibitor Assay. *Bio-protocol* 9, e3336 (2019).

68. Barbosa, G. O. & Kornberg, T. B. Protocol for ex vivo time lapse imaging of
Drosophila melanogaster cytonemes. *Star Protoc* 3, 101138 (2022).

69. Chen, W., Huang, H., Hatori, R. & Kornberg, T. B. Essential basal cytonemes take
up Hedgehog in the Drosophila wing imaginal disc. *Development* 144, 3134–3144
(2017).

1105
1106 **ACKNOWLEDGMENTS**
1107

1108 We thank Drs. T.B. Kornberg and G.O. Barbosa for sharing the design of the culture

1109 chamber for live imaging; Ge Yan, for R plot analyses; Drs. N. Andrews, T.B. Kornberg,

1110 W. Snell, and S. Ogden lab for comments on the manuscript; the Bloomington Stock

1111 Center for *Drosophila* lines; the DSHB for antibodies; A.E. Beaven for UMD imaging

1112 core facility. Funding: NIH grant R35GM124878 to S.R.

1113

1114 **AUTHOR CONTRIBUTIONS**

1115 A.S. discovered GPI-anchoring of Bnl and L.D. discovered bidirectional signaling and

1116 roles of GPI-anchored Bnl; S.R. supervised the work and designed the project; L.D.,

1117 A.S, Y.L. conducted experiments: S.R, L.D, and A.S. wrote the paper.

1118

COMPETING INTERESTS

None declared.

FIGURE LEGENDS

Fig. 1. The reciprocal polarity of Bnl presentation and reception via cytonemes.

a Drawing depicting the organization of the ASP, wing disc, myoblasts, and Btl-

containing ASP cytonemes receiving Bnl from the disc *bnl* source. **b** Spatial

organization of the wing disc *bnl*-source (*bnl-LexA,LexO-mCherryCAAX*) and ASP

(*btlGal4,UAS-CD8:GFP*) cytonemes (arrow). **c-e'** Polarized Bnl presentation from the

source orienting toward the ASP; c, polarized clusters of externalized Bnl^{ex} (red; αBnl^{ex})

at the contact sites of the unmarked source (dashed lined area) and Btl:GFP-containing

ASP cytonemes (Btl:GFP - *btl:GFP* fTRG, see Supplementary Table 4); d,d',e,e',

polarized presentation and cytoneme-mediated delivery of overexpressed Bnl:GFP (d-

d"; *UAS*-mCherryCAAX/+*; bnl-Gal4/ UAS*-Bnl:GFP) and endogenous Bnl:GFP^{endo} (e,e';

UAS-mCherryCAAX/+*; bnl-Gal4/bnl:gfp^{endo}*) from the mCherryCAAX-marked *bnl*-source,

orienting toward the overlaying ASP (dashed line); dashed arrow, Bnl:GFP puncta in

internalized vesicles [9] within the ASP (dashed line); arrow, Bnl:GFP puncta on source

cytonemes; e,e', airyscan image; d", Bnl:GFP intensity plot within the boxed source area

in d' along the proximal (p)-to-distal (d) direction (arrows) relative to the ASP, showing

selective enrichment of the overexpressed signal toward the ASP. **f-h"** 3D-rendered

images showing the ASP-specific polarity of source cytonemes (arrows); f,g,

mCherryCAAX-marked source and nlsGFP-marked ASP (*btl-Gal4,UAS*-nlsGFP/+; *bnl-*

LexA,LexO-mCherryCAAX/+); h,h', CD8:GFP-expressing mosaic clones within the *bnl*

source area (see Methods); h", violin plot displaying the source cytoneme length

distribution (see Supplementary Figure 1d). **i-n** Contact-dependent reciprocal guidance

of source (red) and ASP (green) cytonemes (*btl-Gal4,UAS*-CD8:GFP/+; *bnl-LexA,LexO*-

mCherryCAAX/+); arrowhead, contact site; j, illustration of i-k''; l-n, violin plots displaying

ASP (green) and source (red) cytoneme dynamics as indicated (also see

Supplementary Figure 1e-h and Supplementary Table 1 for statistics). All except c-e',

live imaging. Violin plots: black dotted lines - median and 25^{th} and 75^{th} percentiles.

Source data are provided as a Source Data file. Scale bars, 20 μm; 5 μm (e,e',h,h').

1149

Fig. 2. Bidirectional matchmaking of Bnl presenting and receiving cytonemes for

contacts.

a-e 3D-rendered views of mCherryCAAX-marked *bnl*-source, showing autonomous

effects of Dia:GFP, Dia:GFP^{CA}, and *diaRNAi* expression on source cytoneme numbers

(*UAS-mCherryCAAX;bnl-Gal4* X w$^-$ for control, or *UAS-"X"*); a-d, arrow, source

cytonemes, arrowhead, missing source cytonemes; e, violin plots showing numerical

values, as indicated; p values - one way-ANOVA followed by Tukey's honestly

significant different test. **f-h** Levels of Bnl:GFP^{endo} uptake (dashed arrow) in the ASP

(dashed line) from wild type source cells (f; control: *bnl:gfp^{endo}* X *bnl-Gal4*) and from *dia-*

i-expressing source cells (g; cytoneme-depleted condition: *UAS-dia-i,bnl:gfp^{endo}* X *bnl-*

Gal4); h, violin plots showing numerical values, as indicated; p value – unpaired two-

tailed t test. **i-k** Comparison of numbers of CD2:GFP-marked ASP cytonemes (dashed

arrow) of various length produced under the control condition and under *dia-i*-induced

source cytoneme depleted conditions as indicated; k, violin plots showing numerical

values as indicated; p value – unpaired two-tailed t test. e,h,k, violin plots: black dotted

lines show the median as well as 25[th] and 75[th] percentiles. **l-n'** Non-autonomous effects

of *dia-i*-induced depletion of ASP cytonemes (arrows; m,m') on source cytonemes

(dashed arrows; m",m'''); l,l',n,n', R-plots, showing the correlation of ASP and source

cytoneme number, length and orientation in control (l,l'; *w-) and btl>dia-i* condition (n,n');

see Supplementary Table 2 for statistical values. **o,p** Non-autonomous effects of *Btl-*

DN-induced depletion of ASP cytonemes (arrow) on source cytonemes (dashed arrow):

genotypes - *btl-Gal4,UAS*-CD8GFP/+; *bnl-LexA,LexO*-mCherryCAAX/+ (o); *btl-*

Gal4,UAS-CD8GFP/+; *bnl-LexA,LexO*-mCherryCAAX/*UAS-Btl-DN* (p). All panels

(except f,g), live imaging. Graphs (e,h,k,l,l',n,n'): n represents numbers of biologically

independent samples; Source data are provided as a Source Data file. Scale bars, 20

μm.

1176

Fig. 3. CAM-like Btl-Bnl signaling drives bidirectional contact matchmaking of

source and recipient cells.

a-c Correlation of numbers of polarized Btl-DN-expressing ASP cytonemes (arrow) with

the polarized *bnl*-source cytoneme (dashed arrow) (*btl-Gal4,UAS*-CD8GFP/+; *bnl-*

LexA,LexO-mCherryCAAX/*UAS-Btl-DN*; see Methods); arrowheads, contact sites

between ASP and source cytonemes; c, graph showing the correlation between source

and ASP cytoneme numbers in each disc (n=19). **d,d'** Bnl^{ex} (blue, αBnl^{ex}) is

asymmetrically enriched at the contact sites (arrowheads) between the source and Btl-

DN-expressing ASP cytonemes; d', only αBnl^{ex} channel from d. **e,e'** Btl-DN:Cherry

colocalized with Bnl^{ex} (green; αBnl^{ex}) on tracheal cytonemes; e', only αBnl^{ex} channel

from e. **f** Drawing illustrating the CAM assay using S2 cells and the binding of cell

surface Btl:Cherry and Bnl:GFP that could induce reciprocally polarized synaptic co-

clustering of receptors and ligands and trans-pairing of the interacting cells. **g-n** Trans-

pairing of S2-Bnl:GFP with either S2-Btl:Cherry (g-g",k,k') or S2-Btl-DN:Cherry (l-m') but

not with S2-sBtl:Cherry control (n); dashed lines, cell outline; arrow, trans-synaptic

receptor-ligand co-clusters; dashed arrow, nucleus; dpERK (blue, αdpERK). **h** Bar

graphs comparing the mean frequency (± SD) of receptor-ligand trans-pairing from

three independent experiments (see Methods); n, total GFP-positive + mCherry-positive

cells analyzed: 1916 (Btl:Cherry + Bnl:GFP), 1868 (BtlDN:Cherry + Bnl:GFP), 1532

(sBtl:Cherry + Bnl:GFP); p values: one way-ANOVA followed by Tukey's honestly

significant different test. **i** Graphs comparing % of dpERK-positive Btl-expressing cells

that are either trans-paired (for Btl:Cherry and Btl-DN:Cherry) or nearby (for sBtl:Cherry

& Unpaired Btl:Cherry) to S2-Bnl:GFP cells; All panels except a,b: fixed cells/tissues.

Source data are provided as a Source Data file. Scale bars, 20 μm; 10 μm (g-n).

1201

Fig. 4. Post-translational Bnl cleavage and lipidation.

a,a' Schematic map of the Bnl protein (a) and hydrophobicity plot (a'); SP, signal

peptide; conserved FGF-domain, HSPG+FGFR binding sites; arrow, furin cleavage site;

sites for HA- (site #1), GFP- (site # 3), and mCherry- tags and putative signal sequence

(SS; hydrophilic spacer - blue; hydrophobic tail - yellow) shown. **b-b'''** Representative

optical sections of S2 cells expressing Bnl:GFP$_3$Cherry$_c$. arrowheads, uncleaved

intracellular protein, arrow, cleaved externalized Bnl:GFP$_3$^{ex} portion probed with αGFP^{ex}

(blue); dashed line, cell outline; split channels - as indicated. **c** Wing disc expressing

Bnl:GFP$_3$Cherry$_c$ under *bnl-Gal4*; arrow, intact protein (GFP+mCherry); open arrow,

cleaved Bnl:GFP3; long and short dashed line, ASP and *bnl*-source, respectively; blue,

αDlg. **d** Illustration of a GPI-AP and the PIPLC cleavage site. **d'-g** PIPLC-based cell-

surface shedding of various constructs expressed from S2 cells; d', surface-localized

fractions of signals relative to the total signals expressed in cells with and without PIPLC

treatments; total GFP-tagged protein - GFP intensity; surface protein levels (red) -

αGFP^{ex}-probing of GFP-tagged signals (αBnl^{ex} used for native Bnl); e, schematic maps

of Bnl:GFP constructs used; f,g, box plots comparing the ratio of cell surface (red,

αGFP^{ex} or αBnl^{ex} immuno-fluorescence) to total proteins (GFP fluorescence) from cell

images as shown in d'; For untagged Bnl, surface Bnl level was normalized with co-

expressed CD8:GFP levels/cell. For number of cells used for analyses, see Methods;

Box plots: box shows the median and 1st and 3rd quartile, and whiskers are minimum

and maximum. ***, p<0.001 (unpaired two-tailed t-test). S2 cells: co-transfected with

actin-Gal4 and *UAS-"X"*. Source data are provided as a Source Data file. Scale bars: 10

μm.

1225

Fig. 5. Bnl is GPI-anchored to the wing disc source cell surface.

a-c Wing discs (*w⁻*) showing surface-localized native Bnl^{ex} (red, αBnl^{ex}) levels on the

bnl-source area (arrows) before (a) and after (b) PIPLC treatment; c, box plots

comparing numerical values as indicated. **d-f** Wing discs overexpressing Bnl under *bnl-*

Gal4 showing surface Bnl^{ex} (red, αBnl^{ex}) levels on the *bnl*-source (arrows) before (d)

and after (e) PIPLC treatment; f, box plots comparing numerical values as indicated. c,f,

n = biologically independent samples; p = 0.0013 for c and p <0.0001 for f (unpaired

two-tailed t-test). **g-m** Source surface levels (red, αGFP^{ex}) of Bnl:GFP, Bnl:GFPΔC and

Bnl:GFPΔC-TM on wing discs when expressed under *bnl-Gal4* before and after PIPLC;

asterisks, non-expressing source-surrounding disc area; dashed line, ASP or *bnl-*

source; m, box plots comparing the fraction of surface localized (red, αGFP^{ex}) to total

protein (probed by GFP) of Bnl:GFP variants on the source before and after PIPLC

treatment; n = 5 biologically independent samples for each; p values were obtained by

one way-ANOVA followed by Tukey's honestly significant different test. All box plots:

box shows the median, 1st quartile, 3rd quartile, and whiskers show minimum and

maximum. Source data are provided as a Source Data file. Scale bars, 30 μm.

1242

Fig. 6. GPI-anchored Bnl acts as a CAM for bidirectional matchmaking of

cytonemes.

a-c CAM-like polarized trans-pairing of S2-Btl:Cherry with either S2-Bnl:GFPΔC-TM

(b,c) or S2-Bnl:GFP (c; see Figure 3g,k) but not with S2-Bnl:GFPΔC (a,c); arrow,

polarized receptor-ligand co-clusters at the synaptic site; open arrow, nucleus in source

proximal S2-Btl:Cherry (a) and trans-paired S2-Btl:Cherry (b); arrowhead, Bnl:GFP

signal uptake into the juxtaposed or trans-paired S2-Btl:Cherry cell; blue, nuclear

dpERK (αdpERK); c, bar graphs comparing the mean ($\pm$ SD) frequency of receptor-

ligand trans-pairing for GPI-modified and non-GPI modified Bnl:GFP variants from three

independent experiments (see Methods); n, total GFP-positive + Cherry-positive cells

analyzed: 1916 (Bnl:GFP + Btl:Cherry), 2664 (Bnl:GFPΔC + Btl:Cherry), 2192

(Bnl:GFPΔC-TM + Btl:Cherry); p values were obtained by one way-ANOVA followed by

Tukey's honestly significant different test. **d-l** Comparison of Bnl:GFP (control),

Bnl:GFPΔC, or Bnl:GFPΔC-TM signals for induction of reciprocal polarity of ASP and

source cytonemes (arrows), when expressed from the disc source; genotypes, as
indicated; d, inset, ROI (dashed box) in green and blue channels; d,g,j, extended Z
projection; e,e',h,k, 3D-rendered views; dashed lines, ASP; g,h, dashed arrows,
randomly oriented short cytonemes; f,i,l, R-plots comparing numbers, length, and
directionality of ASP and source cytonemes as indicated; n = number of discs analyzed
(also see Supplementary Figure 6a-g). Source data are provided as a Source Data file.
All panels except a,b, live imaging. Scale bars, 10μm (a,b), 20μm (d-k).
1264

Fig. 7. CAM-like Btl-Bnl-interactions organize cytonemes and cytoneme contacts.
a-a'' 3D projection (a,a'') and orthogonal view (a') of wing discs showing cytoneme-
mediated adhesion between the ASP (*btl>CherryCAAX*) and the Bnl:GFPΔC-TM-
expressing disc *bnl*-source (blue, *bnl>CD4:IFP2,*Bnl:GFPΔC-TM); arrows, Bnl:GFPΔC-
TM-localized contact sites. **b-c** 3D-projected images of Bnl:GFPΔC-TM-expressing wing
disc *bnl*-source (blue, *bnl>CD4:IFP2*) in *btl:cherry*[endo] knock-in background (ASP
expressed endogenous Btl:Cherry [9]), showing trans-synaptic co-clustering of
Btl:Cherry[endo] and Bnl:GFPΔC-TM puncta (arrows) at the inter-cytoneme contact sites.
d,e Violin plots showing dynamics of Bnl:GFPΔC-TM-exchanging recipient and source
cytonemes (compared to control in Figure 1l-n; also see Supplementary Table 1 for
statistics); in violin plots, black dotted lines show the median and 25[th] and 75[th]
percentiles. **f-j** Randomly-localized wing disc clones (mCherryCAAX-marked)
expressing Bnl:GFPΔC-TM, and their cytoneme-dependent interactions (arrow) with the
ASP (dashed line); f, an approximate map of clones in g-j; open arrows, randomly
oriented cytonemes. **k-m'** Randomly-localized wing disc clones expressing Btl:GFP (*),
and their cytoneme-dependent polarized interactions (arrow) with mCherryCAAX-
marked *bnl*-source (only the basal-most section of disc columnar cells shown);
genotypes: see Method; m', zoomed in ROI of m. All panels, live imaging. Source data
are provided as a Source Data file. Scale bars, 20 μm.
1284

Fig. 8. GPI anchor ensures Bnl release at the cytoneme contact sites.
a-e Distribution patterns of Bnl:GFP variants expressed from the mCherry-marked *bnl*
source (*UAS-mCherryCAAX;bnl-Gal4* x *UAS-"X"*); extended Z-projections from the

basal disc area and disc ASP interface shown (for 3D projections, see Supplementary

Movies 12-16); b, inset, extended z-stack includes ASP and disc basal area; d, Inset,

split-colors of ROI (box); arrowhead, source cell membrane containing Bnl:GFPΔC-TM

puncta in the ASP. **f** Schematic map of a Furin-sensor HA$_1$Bnl:GFP$_3$; red arrow, Furin

cleavage site; black arrow, ΔC or TM modification sites; double-sided arrows, uncleaved

HA$_1$Bnl:GFP$_3$ and cleaved Bnl:GFP$_3$ that were transferred to the ASP in the presence

and absence of Fur inhibition, respectively. **f'** Illustration depicting experimental strategy

to detect ASP-specific dispersal rate of uncleaved signals in Furin-inhibited media. **g-p**

Comparison of the ASP-specific uptake of HA$_1$Bnl:GFP$_3$, HA$_1$Bnl:GFP$_3$$\Delta$C-TM, and

HA$_1$Bnl:GFP$_3$$\Delta$C (yellow puncta: red-$\alpha$HA + green-GFP) from the disc source; p, Graphs

comparing the levels of uptake over time (see Methods); for each time point, values

represent the mean $\pm$ SD from multiple biologically independent samples; HA$_1$Bnl:GFP$_3$:

n = 12 (1 h), 11 (2.5 h), 10 (5 h); HA$_1$Bnl:GFP$_3$$\Delta$C-TM: n = 16 (1 h), 18 (2.5 h), 15 (5 h);

HA$_1$Bnl:GFP$_3$$\Delta$C: n = 8 (1 h), 14 (2.5 h), 11 (5 h). $p < 0.01$ for HA$_1$Bnl:GFP$_3$ vs

HA$_1$Bnl:GFP$_3$$\Delta$C-TM or HA$_1$Bnl:GFP$_3$$\Delta$C at 2.5 h and 5 h; p values were calculated

using one way-ANOVA followed by Tukey's honestly significant different test. **q-s**

Comparison of non-specific dispersion between HA$_1$Bnl:GFP$_3$ (q) and HA$_1$Bnl:GFP$_3$$\Delta$C

(r,s) in ex vivo wing disc; All panels: dashed outline, ASP; arrow, cleaved Bnl:GFP$_3$;

arrowhead, uncleaved signal; dashed arrow, source cells; αDlg, cell outlines; asterisk,

non-specific disc areas; g-s, only merged and corresponding red channels were shown.

Source data are provided as a Source Data file. Scale bars, 20 μm.

1309

Fig. 9. Contact-dependent release controls context-specific Bnl signaling

patterns.

a-c Comparison of the dispersion and signaling (dpERK, red; arrows) patterns of

Bnl:GFP (control), Bnl:GFPΔC, and Bnl:GFPΔC-TM expressed from ectopic GOF

clones within the ASP stalk and TC; approximate clone locations and dpERK patterns

indicated in inset; genotype: *hsFlp*; *btlenh>y+>Gal4,btlenh-mRFPmoe* x *UAS-"X"*; arrow

and arrowhead (b,b'), signal-recipient cells with and without nuclear dpERK,

respectively; **d** Violin plots comparing the percentage of signal-receiving ASP cells with

nuclear dpERK from clonal analyses in a-c as well as overexpression of signal under

bnl-Gal4 (OE; *bnlGa4 x UAS-X*; see Supplementary Figure 7h-k); for OE: p < 0.01 for

ΔC (n = 16) vs either Bnl:GFP (n = 17) or TM (n = 13) (one way-ANOVA followed by

Tukey's HSD test); for clonal analyses: see Supplementary Figure 7g for statistics. **e-h**

Levels of tracheal branch invasion (arrows) into larval salivary glands ectopically

expressing either Bnl:GFP (control), TM, or ΔC under *bnl-Gal4* (*bnl-Gal4* X *UAS-X*); e-g,

brightfield images, 10X magnification; h, graphs comparing frequency of terminal

branching using Sholl analysis (see Methods); values represent mean ± SD. **i-l** Levels

of ectopically expressed Bnl:GFP, TM, and ΔC displayed on the basal surface of the

expressing salivary glands; l, violin plots comparing the fraction of surface-displayed

signals (red/arrowhead, αGFP^{ex}) relative to the total protein (GFP level); arrow, cell

junctions; n, number of salivary glands from different larvae; p < 0.05: Bnl:GFP vs. TM,

p < 0.01: ΔC vs. either Bnl:GFP or TM; p values - one way-ANOVA followed by Tukey's

honestly significant different test. Violin plots, black dotted lines show the median as

well as 25th and 75th percentiles. Source data are provided as a Source Data file. Scale

bars, 30 μm; 100 μm (e-g); 20 μm (i-k').

1334

Fig.10. Bnl's GPI anchoring and cytoneme-mediated bidirectional signaling.

a Model illustrating mechanisms of GPI anchoring and tissue-specific cytoneme contact

assembly by CAM-like bidirectional Btl-Bnl signaling, followed by the contact-dependent

signal release and MAPK signaling feedbacks in the recipient ASP cell; dashed arrows,

putative CAM-like signaling feedbacks at the cell membrane cortex. **b-c** Schematic

models, illustrating the loss of CAM-like bidirectional signaling with BnlΔC (b) and the

rescue of its contact-dependent bidirectional signaling by the addition of a TM domain

(BnlΔC-TM) (c).

1343

a Cytonemes | Btl | Bnl | ASP | Myoblasts | Bnl source | Wing disc

b *btl*>CD8:GFP   *bnl*>Cherry

c Btl:GFP   αBnl^ex

d *bnl*>Bnl:GFP

d' | d | p

d'' source area
p → d
Levels (arb. unit) 140
Distance (μm) 0 — 20

e Bnl:GFP^endo

e'

f *btl*>nlsGFP
bnl>CherryCAAX
wing disc   Z ↑ X ←

g

i *btl*>CD8:GFP

j

k | **k'** | **k''** Z ↑ X →

h *bnl-Gal4 UAS>FRT>CD8:GFP*

h'

Z ↑ X →   5 μm

h'' Length (μm) 30 — 0

l Av. velocity of cytonemes (nm/s) 60
ASP   Source
Extend   Retract   Extend   Retract

m Max. cytoneme extension (μm) 40

n Cytoneme Lifetime (min) 30

a Control **b** Dia:GFP **c** Dia:GFP^CA **d** *dia-i*

*bnl>*CherryCAAX

e
source cytonemes
p<0.01
p<0.05
w⁻ (n=6) Dia:GFP (n=6) Dia:GFP^CA (n=8) *bnl>dia-i* (n=6)

f *(x) w⁻* **g** *(x) dia-i*
bnlGal4/ Bnl:GFP^endo Phalloidin
ASP
Control source
ASP
dia-i source

h
ASP Bnl:GFP^endo levels (arb. unit)
p<0.0001
w⁻ (n=7) *bnl>dia-i* (n=17)

Source>*dia-i*

i *bnlGal4* *btl-LexA>*CD2:GFP
Control

j *bnl>dia-i*
dia-i source

k
#ASP cytonemes
w⁻ (n=6) *bnl>dia-i* (n=14)
p<0.0001
<15 >15
Cytoneme length (μm)

ASP>*dia-i*

ASP cytonemes Source cytonemes

l *(x) w⁻* **l'**
< 15 μm > 15 μm
n=11

*btl-Gal4>*CD8:GFP, Gal80^ts *bnl-LexA>*mCherryCAAX

m *(x) btl>dia-i* **m''**
m' **m'''**

n **n'**
n=8

o Control
TC
Control ASP

ASP>*btl-DN*

p *btl>*BtlDN, CD8:GFP
TC
*bnl>*mCherryCAAX
TC
Btl:DN

btl>BtlDN,CD8:GFP   **bnl>mCherryCAAX**

c   R² = 0.7324

Source cytoneme

ASP cytonemes

d   αBnl^ex

d'   αBnl^ex

e   BtlDN:Cherry   αBnl^ex

e'

f   S2-Btl:Cherry   Mix & co-incubation

S2-Bnl:GFP

Trans-paired cells

Btl:Cherry   Bnl:GFP   Btl   Bnl

CAM-like Btl-Bnl binding and reciprocally polarized co-clustering at the contact site

S2-Btl:Cherry + S2-Bnl:GFP

g   **g'**   **g''**

h   p<0.01

% of Btl-Bnl-mediated trans-coupled cells

Btl:Cherry + Bnl:GFP   BtlDN:Cherry + Bnl:GFP   sBtl:Cherry + Bnl:GFP

j   dpERK   **j'**   **k**   **k'**

i   % of Btl-expressing cells with nuclear dpERK

n=83   n=25   n=68   n=28

Trans-paired   Unpaired Btl:Cherry   BtlDN:Cherry + Bnl:GFP   sBtl:Cherry + Bnl:GFP

Btl:Cherry + Bnl:GFP

l   S2-Btl-DN:Cherry + S2-Bnl:GFP   **l'**   **l''**

m   S2-Btl-DN:Cherry + S2-Bnl:GFP   **m'**   dpERK   **n**   S2-sBtl:Cherry + S2-Bnl:GFP   Phalloidin

a S2-Bnl:GFPΔC

dpERK

b S2-Btl:Cherry

S2-Bnl:GFPΔC-TM

c
% of Btl-Bnl-mediated trans-coupled cells

p<0.01   p<0.01

Bnl:GFP +Btl:Cherry | Bnl:GFPΔC +Btl:Cherry | Bnl:GFPΔC-TM +Btl:Cherry

ASP cytonemes
btl>mCherryCAAX   *bnl*>CD4:IFP2

Source cytonemes
bnl>mCherryCAAX

d *bnl*>Bnl:GFP

e   **e'**

g *bnl*>Bnl:GFPΔC

h

j *bnl*>Bnl:GFPΔC-TM

k

ASP cytonemes # (av)   Source cytonemes # (av)

■ < 15 μm   ■ <10 μm ■ 10-20 μm
■ > 15 μm   ■ >20 μm

f   n=12   n=13

i   n=7   n=11

l   n=11   n=11

Bnl:GFP **Bnl:GFPΔC** **Bnl:GFPΔC-TM**

a dpERK

a' btl>FRT>Gal4

btl-RFPmoe

b

b'

c

btl>FRT>Gal4

d

Clone OE Clone OE Clone OE

% signal receiving ASP cells inducing dpERK

Bnl:GFP TM ΔC

e bnl-Gal4 (salivary gland)

100 µm

f

100 µm

g

100 µm

h

branch interconnection

Bnl:GFPΔC-TM
Bnl:GFP
Bnl:GFPΔC

0 Distance from the central branch point (µm) 100

i bnl-Gal4 (salivary gland)

i' Bnl:GFPˣ

j

j'

k

k'

l

Ratio of surface Bnl:GFP variants to total levels

n=17 n=26 n=29

Bnl:GFP TM ΔC

a GPI-modified Bnl

Signaling feedback   **Btl-expressing ASP cell**

PntP1 ← MAPK

Btl
Cytoskeleton

CAM-like bi-directional signaling

Basal

Signaling feedback

Polarized Bnl trafficking

Golgi   Fur1

ER

Transamidation

Bnl-expressing wing disc cell

Apical

b BnlΔC

MAPK

Lack of bidirectional signaling responses

c BnlΔC-TM

MAPK Signaling

Bi-directional signaling

Golgi   Fur1

ER

⇢ CAM-like signaling feedack for cytoneme polarity

━ Polymerized f-actin   Pre-processed pro-Bnl

- - - Cortical actin   GPI-anchored pro-Bnl

Endosome   GPI-anchored mature Bnl

Y Btl   Btl-Bnl   BnlΔC   BnlΔC-TM