1   ***VAL* genes regulate vegetative phase change via miR156-dependent and**

2   **independent mechanisms**

Jim P. Fouracre[1], Jia He[1,3], Victoria J. Chen[1], Simone Sidoli[2] and R. Scott Poethig[1]*

[1] Biology Department, University of Pennsylvania, Philadelphia, Pennsylvania, 19104,

USA

[2] Department of Biochemistry, Albert Einstein College of Medicine, Bronx, New York

10461, USA

[3] Current address: Cold Spring Harbor Laboratory, Cold Spring Harbor, New York

11724, USA

* Corresponding author: Poethig, R. S. (spoethig@sas.upenn.edu)

Summary

How organisms control when to transition between different stages of development is a key question in biology. In plants, epigenetic silencing by Polycomb repressive complex 1 (PRC1) and PRC2 plays a crucial role in promoting developmental transitions, including from juvenile-to-adult phases of vegetative growth. It is well established that PRC1/2 repress the master regulator of vegetative phase change, miR156, leading to the transition to adult growth, but how this process in temporally regulated is unknown. Here we investigate whether transcription factors in the *VIVIPAROUS/ABI3-LIKE* (*VAL*) gene family provide the temporal signal for the epigenetic repression of miR156. Exploiting a novel *val1* allele, we found that *VAL1* and *VAL2* redundantly regulate vegetative phase change by controlling the overall level, rather than temporal dynamics, of miR156 expression. Furthermore, we discovered that *VAL1* and *VAL2* also act independently of miR156 to control this important developmental transition.

Keywords

Developmental transitions; vegetative phase change; miR156; *SPL*; *VAL1*; epigenetic silencing

Introduction

Flowering plant development is underpinned by transitions between stereotypical stages of growth: embryogenesis, seed maturation, juvenile and adult phases of vegetative development and flowering (Huijser and Schmid, 2011). The correct timing of these transitions is critical to plant survival and, ultimately, reproductive success. Vegetative phase change describes the transition from juvenile-to-adult vegetative growth and is associated with changes to multiple traits, including leaf morphology, light-use efficiency, herbivore resistance and shoot physiology (Gou et al., 2011; Lawrence et al., 2020; Leichty and Poethig, 2019; Mao et al., 2017). In *Arabidopsis thaliana*, the juvenile phase is characterized by small round leaves that lack both trichomes on the abaxial surface and serrations. Adult leaves, on the other hand, are larger, more elongated, serrated and produce abaxial trichomes (Telfer et al., 1997).

Vegetative phase change is triggered by activity of members of the *SQUAMOSA PROMOTER BINDING PROTEIN-LIKE* (*SPL*) family of transcription factors, which are post-transcriptionally repressed during juvenile development by the microRNAs miR156/miR157 (Fouracre and Poethig, 2019; Wu and Poethig, 2006; Wu et al., 2009; Xu et al., 2016a). miR156/miR157 are encoded by multiple genes of which *MIR156A* and *MIR156C* are the most functionally significant (He et al., 2018). The expression of *MIR156A* and *MIR156C* declines during juvenile growth (Yang et al., 2013b; Yu et al., 2013), leading to the de-repression of their *SPL* targets and the transition to adult growth. Elucidating what controls the decline in *MIR156A/C* expression is therefore critical to understanding how the juvenile-to-adult transition is regulated in plants.

The molecular mechanisms that lead to the temporal repression of *MIR156A/C*

are only beginning to be understood. The activity of Polycomb Group (PcG)

transcriptional repressors appears critical. There are two functional complexes of PcG

proteins in plants, both of which repress gene expression through covalent histone

modifications. PcG repressive complex 1 (PRC1) consists of a H2A E3 ubiquitin ligase

module containing one AtBMI1 protein (AtBMI1A/B/C) and one AtRING1 protein

(RING1A/B). PRC1 represses gene expression through ubiquitination of H2A

(H2AK119ub) (Bratzel et al., 2010; Calonje, 2014; Yang et al., 2013a). The PRC2

complex includes histone methyltransferases such as *CURLY LEAF* (*CLF*) and

SWINGER (*SWN*) and promotes H3 trimethylation (H3K27me3) (Chanvivattana et al.,

2004; Goodrich et al., 1997).

We have previously found that H3K27me3 increases at *MIR156A/C* in a PRC2-

dependent manner during juvenile development, and that vegetative phase change is

delayed in *swn* mutants (Xu et al., 2016b). The temporal deposition of H3K27me3 is

accompanied by depletion of the antagonistic H3K27ac mark that is associated with

active transcription. miR156 accumulation is also repressed by PRC1, as *atbmi1a/b*

mutants exhibit delayed vegetative phase change (Pico et al., 2015). In addition, we

have found that accumulation of the active histone mark H3K4me3 decreases at

MIR156A/C during vegetative development (Xu et al., 2018).

The findings that H3K27me3 replaces H3K27ac and H3K4me3 at *MIR156A/C*

over time, and that PRC1/PRC2-activity promotes vegetative phase change, led us to

propose that the temporal dynamics of miR156 accumulation are coordinated by

antagonistic patterns of active (H3K27ac, H3K4me3) and repressive (H3K27me3)

histone modifications (Xu et al., 2018, 2016b). In this model the stochastic removal of

H3K27ac/H3K4me3 facilitates the deposition of H3K27me3 and the gradual epigenetic

silencing of miR156. Similar mechanisms have been reported to function at other

developmental transitions (Yang et al., 2014). For example, during flowering, H3K27

deacetylation is a pre-requisite for PRC2-mediated H3K27me3 deposition at

FLOWERING LOCUS C (*FLC*) (Zeng et al., 2020), and during seed maturation, PRC1

promotes the exchange of H3K4me3 for H3K27me3 at *DELAY OF GERMINATION1*

(*DOG1*) and *ABSCISIC ACID INSENSITIVE3* (*ABI3*) (Molitor et al., 2014).

Although there is good evidence that *MIR156A/C* are epigenetically silenced

during vegetative development*,* how this mechanism is regulated temporally remains

unknown. *VIVIPAROUS/ABI3-LIKE* (*VAL*) genes are excellent candidates for temporal

effectors in this model. *VAL* genes encode B3 domain transcription factors that are

closely related to the *ABI3/FUSCA3* (*FUS3*)/*LEAFY COYTLEDON2* (*LEC2*) clade of

embryogenesis regulators. There are three *VAL* genes in *Arabidopsis*, of which *VAL1*

and *VAL2* (also known as *HSI2* and *HSL2* respectively) are the most functionally

important (Suzuki et al., 2007). VAL proteins repress their targets by binding to 6 base

pair RY-sequence motifs (CATGCA) via their B3 domain (Chen et al., 2018, 2020; Guo

et al., 2013; Jing et al., 2019; Qüesta et al., 2016; Suzuki et al., 1997; Yuan et al.,

2016).

A number of observations suggest that *VAL* genes might provide the temporal

information that coordinates vegetative phase change: 1) *VAL* genes regulate other

developmental transitions, i.e. seed maturation (Suzuki et al., 2007; Yang et al., 2013a)

and flowering (Qüesta et al., 2016; Yuan et al., 2016); 2) *MIR156A/C* expression is

elevated in *val1/2* mutants (Pico et al., 2015); 3) VAL1/2 physically interact with several

histone deacetylases (*HDA6/9/19*) (Chhun et al., 2016; Qüesta et al., 2016; Zeng et al.,

2020; Zhou et al., 2013); and 4) *VAL* genes promote PRC1 and PRC2-binding (Chen et

al., 2018; Pico et al., 2015; Qüesta et al., 2016; Yuan et al., 2020, 2016)

In this study we investigated whether *VAL* genes function as temporal regulators

of vegetative phase change. We report that reduced *VAL* activity significantly delays the

timing of vegetative phase change through both miR156-dependent and independent

mechanisms. We find that the temporal decline in miR156 expression is remarkably

robust and is insensitive to loss of *VAL* function, inhibition of VAL1-binding and the

combined loss of *VAL1* and PRC2 components. Finally, we show that the effects of

VAL1 on the timing of vegetative phase cannot be explained by temporal changes in its

interactions with other proteins.

Results

***VAL* genes promote vegetative phase change**

To investigate the role of *VAL* genes in vegetative phase change, we exploited a novel

mutant we identified in an EMS screen for plants exhibiting prolonged juvenile

development. Mapping-by-sequencing revealed a substitution at the *VAL1* locus,

resulting in the conversion of a highly conserved arginine residue to a cysteine in the N-

arm of the VAL1 B3 DNA-binding domain (Figure S1A). This arginine residue is critical

for VAL1 binding to target RY-motifs (Sasnauskas et al., 2018). The mutation in *VAL1*

was confirmed to be the cause of the late juvenile phenotype by its failure to

complement the null *val1-2* T-DNA insertion allele, and by the ability of the *VAL1*

genomic sequence to rescue this phenotype (Figure S1B,C). Unlike *val1-2*, the novel

val1 allele is semi-dominant, and delays vegetative phase change when heterozygous

(Figure S1D). We therefore named this new allele *val1-5(sd)*, consistent with the

nomenclature of existing *val1* alleles (Veerappan et al., 2012).

Both *val1-5(sd)* and *val1-2* exhibit delayed vegetative phase change, with *val1-*

5(sd) having a stronger effect on the timing of abaxial trichome production than *val1-2*

(Figure S1B). As *VAL1* functions redundantly with *VAL2* to regulate other

developmental transitions (Qüesta et al., 2016; Suzuki et al., 2007; Tsukagoshi et al.,

2007; Yuan et al., 2016), we tested the effects of *val1; val2* double mutants on

vegetative phase change. Previous analyses of *VAL* gene function have utilized *val1-2;*

val2-1 and *val1-2; val2-3* double mutants. However, seedling development is so strongly

perturbed in *val1-2; val2-1* and *val1-2; val2-3* plants (Suzuki et al., 2007; Tsukagoshi et

al., 2007; Yang et al., 2013a; Yuan et al., 2020) that analyses of vegetative growth is

problematic in these backgrounds. Therefore, we generated new *val1; val2*

combinations using *val2-3* and a previously uncharacterized T-DNA insertion allele we

named *val2-4* (Figure 1A). Consistent with previous studies (Suzuki et al., 2007;

Tsukagoshi et al., 2007; Yuan et al., 2016), *val2* single mutants had no discernible

effect on vegetative phase change (Figure 1B, S1E). However, loss of *VAL2* activity

enhanced the phenotypes of *val1-2* and *val1-5(sd)*. *val1-2; val2-4* and *val1-5(sd); val2-3*

both exhibited delayed abaxial trichome production relative to *val1-2* and *val1-5(sd)*,

respectively (Figure 1C). *val1-5(sd); val2-3* flowered significantly later than *val1-5(sd)*

(Figure 1D) and *val1-2; val2-4* produced leaves that were significantly more juvenile in

shape (i.e. rounder) than *val1-2* (Figure 1E). Neither double mutant combination was as

Figure 1. *VAL* genes redundantly regulate vegetative phase change

(A)   Schematic of *val1* and *val2* alleles used in this study – grey boxes represent

          UTRs, black boxes represent exons, red triangles represent T-DNA insertions,

          red line represents EMS-induced base substitution.

(B)   Phenotypes at 21 DAG in LD conditions, scale bar = 5mm.

(C-E) Quantitative analysis of vegetative development. Statistically distinct genotypes

          were identified by one-way ANOVA with *post hoc* Tukey multiple comparison test

          (letters indicate statistically distinct groups $P < 0.05$; for (E) comparisons were

          made at leaf 5), all plants grown in LD. (C,D) Boxes display the interquartile

          range (IQR) (boxes), median (lines) and values beyond 1.5*IQR (whiskers);

          mean values are marked by ◆. (E) Colored lines represent the mean and black

          bars the SEM. Sample sizes (C, D) 21-46, (E) 13-46.

phenotypically severe as *val1-2; val2-1* (Figure 1C-E). The weaker phenotype of *val1-*

5(sd); val2-3 than *val1-2; val2-3* (Yang et al., 2013a; Yuan et al., 2020) suggests that

the semi-dominant phenotype of *val1-5(sd)* is mediated by interaction with *VAL2*.

Importantly, the rate of germination was higher in *val1-2; val2-4* and *val1-5(sd); val2-3*

relative to existing *val1; val2* double mutants. *val1-2; val2-4* and *val1-5(sd); val2-3* thus

provide a balance between phenotypic strength and experimental viability and are

useful tools for investigating the role of *VAL* genes in developmental timing.

VAL genes function predominantly as quantitative – rather than temporal –

regulators of miR156 expression

Vegetative phase change results from a temporal decline in miR156 expression (Wu

and Poethig, 2006). A previous analysis of *val1-2; val2-1* revealed elevated expression

of *MIR156A/C* at a single time point (Pico et al., 2015). To determine whether the delay

in vegetative phase change we observed in *val* mutants is associated with a general

increase in the level of miR156, or with a delay in the decline in this miRNA, we

quantified miR156 expression in the shoot apex and in isolated leaf primordia at

different times in shoot development. The primary transcripts of *MIR156A* and *MIR156C*

were expressed at similar levels, and exhibited a similar temporal expression pattern, in

wild type, *val1-5(sd)* (Figure 2A,B) and *val1-2* shoot apices and leaf primordia (Figure

S2). However, the abundance of the mature miR156 miRNA transcript was significantly

higher in *val1-5(sd)* leaf primordia than in wild type (Figure 2B), and it was also

marginally higher in *val1-5(sd)* shoot apices than in wild type (Figure 2A). The *val1-*

5(sd); val2-3 (Figure 2A,B) and *val1-2; val2-1* (Figure S2) double mutants had stronger

Figure 2. *VAL* genes function predominantly as quantitative regulators of miR156 expression

(A, B) qRT-PCR analyses of gene expression. (A) Shoot apices with leaf primordia (LP) ≥ 1mm removed at 1, 2 and 3 weeks. (B) Isolated LP 0.5-1mm in size. All plants were grown in SD conditions. Each data point represents a biological replicate and is the average of three technical replicates. Coloured lines represent the mean and black bars mean±s.e.m. Asterisks represent significant differences between WT and *val* mutants at the same time point, calculated by an unpaired two-tailed *t*-test with a Bonferroni correction for multiple comparisons (* $P <$ 0.025; ** $P <$ 0.005; *** $P <$ 0.0005).

effects on *MIR156A* and *MIR156C* expression than the respective *val1* single mutants,

suggesting that *VAL1* and *VAL2* function redundantly to repress *MIR156A* and

MIR156C transcription. Mature miR156 was elevated throughout development in both

the shoot apices (Figure 2A) and the leaf primordia (Figure 2B) of *val1-5(sd); val2-3*

double mutants. Loss of *VAL* activity also produced a slight increase in miR157 levels

(Figure 2A,B, S2). Although *val1-5(sd); val2-3* increased the abundance of

miR156/miR157, it had only a minor effect on the temporal expression patterns of these

miRNAs. For example, miR156 expression decreased 2.36-fold between 1W and 2W

and 2.31-fold between 2W and 3W in wild type plants, but decreased 1.77 and 1.82-fold

between the same time points in *val1-5(sd); val2-3* plants (Figure 2A). Taken together,

these data suggest that *VAL* genes function primarily as general, rather than temporal,

regulators of miR156 expression.

VAL genes coordinate PRC1 and PRC2 recruitment at *MIR156* loci

We have previously demonstrated that the temporal decline in *MIR156A* and *MIR156C*

expression is associated with PRC2-dependent and progressive deposition of

H3K27me3 at these loci (Xu et al., 2016b). To determine if *VAL* genes contribute to this

process, we examined PRC2 activity in the *val1-5(sd); val2-3* double mutant. As

previously reported (Xu et al., 2016b), H3K27me3 levels increased at the *MIR156A* and

MIR156C loci during vegetative development in wild type plants (Figure 3A,B). Although

there was no difference in the temporal pattern of H3K27me3 deposition at *MIR156A* in

val1-5(sd); val2-3, the rate of H3K27me3 deposition at *MIR156C* was significantly

slower in this double mutant (Figure 3B). These results are consistent with a recent

Figure 3. *VAL* genes regulate miR156 activity via-PRC2 and PRC1

(A)   Schematics of the primer locations used for ChIP-qPCR. Blue and green bars represent sequences encoding the mature miRNA and miRNA hairpin respectively.

(B)   Temporal analysis of H3K27me3 by ChIP-qPCR. Each data point represents a biological replicate and is the average of three technical replicates. Lines represent the mean, bars represent the mean±s.e.m., asterisks represent significant differences between WT and *val1-5(sd); val2-3* at the same time point, calculated by an unpaired two-tailed *t*-test (* *P* < 0.05; ** *P* < 0.01; *** *P* < 0.001). H3K2me3 values are relative to H3 and normalised to *STM* as an internal control. Plants were grown in SD conditions.

(C, D) Phenotypes in LD. (D) Photographs taken at 21DAG, scale bar = 5mm. (E)
Asterisks represent significant differences to either WT or *val1-2*, calculated by
unpaired two-tailed *t*-test with Bonferroni correction for multiple comparisons (***
$P < 0.00033$). Sample size 22-46.
(E)   qRT-PCR analyses of gene expression in shoot apices with LP $\geq$ 1mm removed
at 1, 2 and 3 weeks. Each data point represents a biological replicated and is the
average of three technical replicates. Coloured lines represent the mean and
black bars represent the mean±s.e.m. Asterisks represent significant differences
between *val1-2* and *val1-2; clf-29/val1-2; swn-3/val1-2; hda19-3* at the same time
point, calculated by unpaired two-tailed *t*-test with Bonferroni correction for
multiple comparisons (** $P < 0.0033$). Plants were grown in SD conditions.
(F)   Temporal analysis of H2AK119ub by ChIP-qPCR, values are relative to input and
normalised to *ACT7* as an internal control. See (B) for details.
257

genome-wide study that revealed a decrease in H3K27me3 levels at *MIR156C*, but not

MIR156A, in *val1-2; val2-3* plants (Yuan et al., 2020). As a control, we measured

H3K27me3 deposition at the floral regulator *FLC.* In the absence of vernalization, we

observed no change in the level of H3K27me3 during vegetative development in wild

type plants (Figure 3B). However, consistent with previous reports (Qüesta et al., 2016;

Yuan et al., 2016), there was a significant decrease in H3K27me3 at *FLC* in *val1-5(sd);*

val2-3.

VAL1 is thought to act by recruiting PRC1 which, in turn, promotes the activity of

PRC2 (Baile et al., 2020; Calonje, 2014; Zhou et al., 2017). As a genetic test of this

hypothesis, we examined the interaction between *val1-2* and *clf-29* and *swn-3,* loss-of-

function mutations in the functionally redundant genes that encode the histone

methyltransferase activity of PRC2. As we have shown previously (Xu et al., 2016b),

swn-3 had a larger effect on the timing of vegetative phase change than *clf-29* (Figure

3C,D). Consistent with the hypothesis that *VAL1* regulates vegetative phase via its

effect on PRC2 activity, *clf-29* and *swn-3* interacted synergistically with *val1-2,* in that

the double mutants had a much more severe vegetative phase change phenotype than

the single mutants (Figure 3C,D). Notably, *val1-2* suppressed the curling leaf phenotype

of *clf-29* (Figure 3C), presumably because it enhances *FLC* expression (Lopez-Vernaza

et al., 2012). We also examined the interaction between *val1-2* and *hda19-3*, a putative

null allele of the histone deacetylase HDA19. *hda19-3* is predicted to delay vegetative

phase change because *HDA19* is required for PRC2-mediated repression (Zeng et al.,

2020), HDA19 physically interacts with VAL1 (Qüesta et al., 2016) and loss-of *HDA19*

enhances H3K4me3 levels (Jang et al., 2011). As predicted, *hda19-3* produced abaxial

trichomes significantly later than wild type (Fig. 3D), and *val1-2; hda19-3* double

mutants produced abaxial trichomes significantly later than either single mutant.

Together, these results suggest that *val* mutations delay vegetative phase change by

interfering with the activity of PRC2.

To determine whether the synergistic interaction between *clf-29, swn-3, hda19-3*

and *val1-2* is due to enhanced miR156/miR157 expression, we quantified expression of

the mature miR156 and miR157 miRNAs, and the primary *MIR156A* and *MIR156C*

transcripts, in these mutant backgrounds. The overall level and expression pattern of

the mature miR156/miR157 transcripts were not affected by *val1-2*, or by *val1-2; clf-29*,

val1-2; swn-3, and *va1-2; hda19-3* double mutants (Figure 3E). However, *pri-MIR156A*

and *pri-MIR156C* transcripts were significantly elevated in *val1-2; hda19-3,* although the

rate of decline in the expression of these transcripts was normal. The lack of correlation

between the expression of miR156 and the level of *pri-MIR156A* and *pri-MIR156C* is

surprising given that these loci are the major sources of miR156. One possibility is that

the miRNA processing pathway is compensating for elevated levels of the primary

transcripts. In any case, these results suggest that *VAL* genes temporally regulate the

deposition of H3K27me3 at specific *MIR156* loci, but are not necessary for the temporal

decline in miR156 expression.

VAL genes also repress gene expression by promoting H2AK119ub deposition

via recruitment of PRC1 (Yang et al., 2013a). Unlike H3K27me3 (Figure 3B) (Xu et al.,

2016b), we found no evidence that H2AK119ub increases over time at *MIR156A* and

MIR156C (Figure 3F). However, we did find that *val1-5(sd); val2-3* had significantly

lower levels of H2AK119ub than wild type plants. This effect was observed throughout

development at *MIR156C*, but seemed to be limited to later in development at

MIR156A. Importantly, *val1-5(sd); val2-3* had significantly reduced levels of H2AK119ub

at *MIR156C* one week after planting, which is before a difference in H3K27me3 is

detectable at this locus (Figure 3B). *val1-5(sd); val2-3* also had reduced levels of

H2AK119ub at *FLC* (Figure 3F).

VAL1 regulates vegetative phase change via miR156-dependent and miR156-

independent mechanisms

To investigate whether the effects of *val1-5(sd); val2-3* on the chromatin state of

MIR156A/C are due to a direct regulatory interaction, we carried out chromatin-

immunoprecipitation qPCR using an HA-tagged version of VAL1 (Qüesta et al., 2016).

Confirming the results of a recent ChIP-seq study (Yuan et al., 2020), we found VAL1-

binding at specific locations within both the *MIR156A* and *MIR156C* loci (Figure 4A,B,

S3). The affinity of VAL1 for *MIR156A/C* appeared consistent throughout vegetative

development. VAL1/2 bind to RY-sequence motifs, of which there are multiple copies in

both *MIR156A* and *MIR156C* (Figure 4A). To determine if these RY-sites are required

for the regulation of vegetative phase change, we mutated 5 RY-sites in *MIR156C*

individually and in combination. We selected *MIR156C* because it is more sensitive to

VAL activity than *MIR156A* (Figure 2,3, S2). A C>T substitution that eliminates VAL1

binding (Sasnauskas et al., 2018) was introduced in one or all of these sites in a

genomic construct of *MIR156C*. Wild type and mutant constructs were then transformed

into a *mir156a; mir156c; mir157a; mir157c quadruple mutant* (*qm*) background. We

chose this background because endogenous miR156/miR157 activity has been largely

327
Figure 4. Loss of RY VAL-binding motifs at the *MIR156C* locus delays vegetative phase change

(A)   Schematic depicting the location of primers used for ChIP-qPCR, the sequences encoding the miR156 hairpin and mature miRNA are coloured green and blue respectively.

(B)   Anti-HA ChIP-qPCR of WT Col control plants at 2W and *VAL1::VAL1-HA; val1-2; FRI-Sf2* plants at 1, 2 and 3W of growth. The data is presented as percentage input normalized to *ACT7* and is displayed relative to WT. Each data point represents a biological replicate and is the average of three technical replicates, bars represent the mean and error bars the mean±s.e.m.

(C, D) Phenotypes of T1 plants transformed with *MIR156C* RY variants. Statistically
distinct genotypes were identified by one-way ANOVA with *post hoc* Tukey
multiple comparison test (letters indicate statistically distinct groups $P < 0.05$;
comparison in (C) made at leaf 5). *qm = mir156a mir156c mir157a mir157c*
quadruple mutant, tm = mir156a mir157a mir157c triple mutant, EV = empty
vector. (C) Colored lines represent the mean and black bars the mean±s.e.m.
Sample size (B) 26-52, (C) 37-51.
(E)   qRT-PCR analyses of gene expression in shoot apices with LP ≥ 1mm removed
at 1, 2 and 3 weeks. Each data point represents an independent homozygous T3
line and is the average of three technical replicates. Colored lines represent the
mean and black bars the mean±s.e.m. Asterisks represent significant differences
between *qm; MIR156C* and *qm; MIR156C -RY* lines at the same time point,
calculated by an unpaired two-tailed *t*-test with a Bonferroni correction for
multiple comparisons (* $P < 0.025$, ** $P < 0.005$).
(F)   Genetic interaction between *val1-2* and *MIR156C* RY deletions. Asterisks
represent significant differences between plants with wild type or null *VAL1*
alleles calculated by an unpaired two-tailed *t*-test (*** $P < 0.001$). Sample size
24-36. Phenotyping analyses were carried out in LD conditions, gene expression
and ChIP analyses were carried out in SD conditions.
357

eliminated and it is therefore sensitive to small changes in the level of miR156 (He et

al., 2018). Plants transformed with a wild type *MIR156C* construct (*+RY*) produced

leaves with abaxial trichomes at the same node as the *mir156a; mir157a; mir157c triple*

mutant tm (Figure 4C). *tm* has an endogenous copy of *MIR156C*, confirming that the

transgenic *MIR156C* sequence is fully functional. Deletion of individual RY-sites

produced a significant delay in the timing of abaxial trichome production relative to

MIR156C +RY, and deletion of all 5 RY-sites produced a more significant delay than

deletion of any single site (Figure 4C). A similar result was obtained in the case of the

angle of the leaf base (Figure 4D). These results demonstrate that all five RY-sites are

important for the expression of *MIR156C*, and that they function additively. Individual

RY-sites have also been shown to interact additively to repress the VAL1-PRC2 targets

FLC and *DOG1* (Chen et al., 2020; Yuan et al., 2016).

To determine if these phenotypic effects are due to altered *MIR156C* expression,

we quantified miR156 levels in *MIR156C +RY, MIR156C -RY2* and *MIR156C -RY12345*

plants. *MIR156C -RY2* was selected because it has a marginally stronger effect than

other individual -RY deletions (Figure 4C,D). Although there was considerable variation

in miR156 levels between independent transgenic lines, *MIR156C -RY2* and *MIR156C -*

RY12345 plants had significantly more miR156 than plants transformed with *MIR156C*

+RY (Figure 4E, S4). However, the temporal expression pattern of *MIR156C* was

identical in *-RY* and *+RY* plants.

To establish whether the effects of RY-deletion are *VAL1*-dependent, we crossed

val1-2 into the *qm; MIR156C +/-RY* lines. If the delay in vegetative phase change in

MIR156C -RY lines is a consequence of reduced VAL1 binding, *val1-2* should have less

effect in *MIR156C -RY* lines than in *MIR156C +RY* lines or the *tm*, in which RY sites are

intact. Surprisingly, we found that loss-of *VAL1* significantly delayed abaxial trichome

production in *MIR156C -RY* as well as *MIR156C +RY* and *tm* plants (Figure 4F). It is

possible that *MIR156C* RY-sites are bound by other B3 domain transcription factors.

However, RY-binding is restricted to the *ABI3/FUS3/LEC2* and *VAL* clade of B3 domain

genes (Kagaya et al., 1999; Sasnauskas et al., 2018; Yuan et al., 2016). The

expression of *ABI3/FUS3/LEC2* is largely restricted to seed development (Carbonero et

al., 2017) and therefore these genes are unlikely to regulate vegetative shoot identity.

Moreover, FUS3 and ABI3 directly promote the expression of *MIR156C* (Tian et al.,

2020; Wang and Perry, 2013). A role for these genes in the regulation of *MIR156C* post-

germination is thus inconsistent with the juvenilized phenotype and elevated miR156

expression we found in *MIR156C -RY* plants. (Figure 4C-E). With regard to the potential

effects of other *VAL* genes, we observed no vegetative phase change phenotype in *val2*

single mutants (Figure 1B, S1E) and *VAL3* has limited expression and functionality

relative to *VAL1* and *VAL2* (Suzuki et al., 2007). Alternatively, this result suggests that

VAL1 may regulate vegetative phase change through both miR156-dependent and

miR156-independent mechanisms. This interpretation is supported by the observation

that loss-of *VAL1* and PRC2-components strongly delayed vegetative phase change but

had only minor effects on miR156 expression (Figure 3C-E).

To test this hypothesis, we introgressed *val1-2* into the *qm* genetic background,

which has very low levels of miR156/miR157 (He et al., 2018). Although the gross

morphology (Figure 5A) of *val1-2*; *qm* seedlings was indistinguishable from that of *qm*

plants, *val1-2* partially suppressed the effect of the *qm* genotype on the morphology of

404
Figure 5. *VAL1* regulates vegetative phase change by miR156-dependent and independent mechanisms

(A-C)   Phenotypes of *val1-2* and *mir156a mir156c mir157a mir157c quadruple mutant* lines. (A) Photographs taken at 17 DAG. Scale bar = 1mm. (B, C) Statistically distinct genotypes were identified by one-way ANOVA with *post hoc* Tukey multiple comparison test (letters indicate statistically distinct groups $P < 0.05$; (B) comparisons made at leaf 5). Bars represent the mean±s.e.m. Sample size (B) 17-60, (C) 18-36. Silhouettes in B show representative leaf 5 shapes.

(D)   Expression of a miR156-resistant (*rSPL9*) reporter construct in WT, *val1-2* and *val1-5(sd)* backgrounds. Scale bars = 1mm.

(E)   qRT-PCR analysis of gene expression in shoot apices with LP ≥ 1mm removed at 1, 2 and 3 weeks. Each data point represents a biological replicate and is the average of three technical replicates. Coloured lines represent the mean and gray bars represent the mean±s.e.m. Asterisk represents significant difference between WT and *val* mutant lines at the same time point, calculated by an unpaired two-tailed *t*-test with a Bonferroni correction (* $P < 0.025$; ** $P < 0.005$). All phenotypic analyses were carried out in LD conditions, the gene expression analysis was carried out in SD conditions.

leaves 3 and 5 (Figure 5B). This confirms that *VAL1* functions through a

miR156/miR157-independent mechanism to regulate vegetative phase change.

Surprisingly, a survey of abaxial trichome production revealed that *val1-2* enhanced this

aspect of the precocious *qm* phenotype (Figure 5C)*.* This result can be explained given

that reduced histone deacetylation in a *toe* loss-of-function background accelerates

abaxial trichome formation (Wang et al., 2019), that *SPL* genes repress *TOE* activity

(Wu et al., 2009), and that *VAL* genes promote histone deacetylation (Zeng et al., 2020;

Zhou et al., 2013).

To determine if *VAL1* regulates *SPL* gene expression independently of miR156,

we crossed a miR156-resistant (*rSPL9*) *SPL9::rSPL9-GUS* reporter construct into *val1-*

5(sd). The expression of this reporter was visibly and strongly suppressed by *val1-2* and

val1-5(sd) (Figure 5D), implying that miR156 is not required for the regulation of *SPL9*

by *VAL1*. The transcript levels of *SPL9*, and its target miR172, were also decreased in

val mutant plants (Figure 5E). However, it is difficult to know if this decrease is

dependent or independent of miR156 because miR156 induces cleavage of the *SPL9*

transcript (German et al., 2008; He et al., 2018; Ronemus et al., 2006).

VAL1/2 act as transcriptional repressors. As *SPL9* transcription decreases in *val*

loss-of function mutants it is therefore unlikely that VAL1/2 regulate *SPL9* directly. To

test this prediction, we quantified H2AK119ub and H3K27me3 at *SPL9.* Consistent with

previous studies, we found high levels of H2AK119ub but negligible H3K27me3 at *SPL9*

(Figure S5) (Li et al., 2017; Zhou et al., 2017)*.* We also found no difference in the

abundance of these modifications in wild type and *val1-5(sd); val2-3* plants. Taken

together these results suggest that in addition to repressing *SPL9* via their effect on

miR156 levels, VAL1/2 repress *SPL9* indirectly through one or more miR156-

independent mechanisms. The observation that *val1-5(sd); val2-3* has no effect on

H2AK119ub at *SPL9* also suggests that *VAL1/2* are not universally required for PRC1-

activity.

The effects of *VAL1* on developmental timing may be partly explained by its

expression pattern

Our results show that *VAL* genes control the timing of vegetative phase change (Figure

1C,E), and have subtle effects on the expression pattern of miR156 during vegetative

development (Figure 2, S2). To determine if these effects are attributable to changes in

the expression level of *VAL1*, we measured the abundance of *VAL1* transcript levels

during vegetative growth. We observed a small but significant increase in *VAL1*

transcripts in shoot apices between 2 and 3 weeks of growth (Figure 6A) but there was

no significant change in *VAL1* levels in leaf primordia (Figure 6B). To further investigate

VAL1 expression over time, we generated a *VAL1* transcriptional reporter by fusing a

2.3kb sequence containing the *VAL1* promoter and 5' UTR, and a 2kb sequence

containing the *VAL1* 3'UTR and terminator, to the GUS coding sequence (*VAL1::GUS-*

VAL1 3' UTR). We generated a *VAL1* translational fusion by inserting a 3.8kb *VAL1*

genomic sequence upstream of *GUS* in this construct (*VAL1::VAL1-GUS-VAL1 3'UTR*).

Because the expression of the transcriptional fusion was consistently more diffuse,

variable, and weaker than the expression of the translational fusion (Figure S6A,B), we

used plants containing the translational fusion for subsequent studies.

Figure 6. *VAL1* is expressed throughout vegetative development

(A, B) qRT-PCR analyses of gene expression. (A) Shoot apices with leaf primordia (LP) ≥ 1mm removed at 1, 2 and 3 weeks. (B) Isolated LP 0.5-1mm in size.

(C-F) Analyses of a *VAL1-GUS* translational fusion. (C) Expression in whole plants, scale bars = 1mm. (D) Expression in shoot apices following wax sectioning, scale bars = 0.1mm. (E, F) VAL1 levels quantified by MUG assay in two independent homozygous T3 lines. (A, B, E, F) Each data point represents a biological replicate and is the average of three technical replicates. Lines represent the mean and grey bars mean±s.e.m. Asterisks represent significant differences between two continuous time points, calculated by an unpaired two-tailed *t*-test (* $P < 0.05$; ** $P < 0.01$). All plants except the 3 DAG sample (C – long days) were grown in SD conditions.

During embryogenesis, the translational fusion was expressed in the root and shoot

apical meristems and provasculature (Figure 6C, S6A). Following germination,

expression became restricted to the shoot and root apices and initiating lateral root

primordia (Figure 6C, S6B). Throughout the rest of shoot development, the translational

fusion was expressed in the shoot apex and during the early stages of leaf development

(Figure 6C). Histological inspection indicated that *VAL1* expression increases in the

shoot apex during vegetative development. This was validated by a quantitative

analysis of GUS expression, which demonstrated that VAL1 accumulates more strongly

in the shoot apex than leaf primordia. Further, that VAL1 levels increase over time in the

shoot apex but not in older leaf primordia (Figure 6E,F). Taken together, these data

indicate that *VAL1* expression is restricted to apical meristems and the early stages of

root and leaf development. Our data also indicate that *VAL1* expression increases

during shoot development in very young leaf primordia, quickly declines to a uniform

level as the leaf develops, and ceases before the leaf is fully expanded. However, the

functional significance of increased VAL1 accumulation during vegetative development

is unclear, as we did not detect a concomitant increase in VAL1-binding to *MIR156A/C*

(Figure 4B, S3). Finally, the difference in the staining patterns and stability of our

transcriptional and translational reporters suggest that cis-regulatory elements within the

VAL1 coding sequence regulate the level and site of its expression.

VAL1 has previously been found to physically interact with VAL2, multiple PRC1

and PRC2 components, and the transcriptional repressor SAP18 (Chen et al., 2018,

2020; Chhun et al., 2016; Jing et al., 2019; Qüesta et al., 2016; Xie et al., 2018; Yang et

al., 2013a; Yuan et al., 2016). However, these studies do not provide information about

VAL1's *in planta* protein interactions over time. To determine whether VAL1 interacts

with different proteins at different stages of vegetative development, we carried out a

mass spectrometry analysis of proteins bound to HA-tagged VAL1 at 1, 2 and 3 weeks

of growth. Immunoprecipitations were carried out on a *VAL1-HA; val1-2; FRI-Sf2* line,

using a *val1-2; fri* line as a control. The difference in the *FRI* genotype of these lines is a

consequence of the genotypes available at the time the experiments were performed

and may have had an effect on our results.

ATBMI1A was significantly enriched in the combined experimental samples

relative to the control samples (Figure 7A), which is consistent with a previous mass

spectrometry analysis of proteins bound to VAL1-HA (Qüesta et al., 2016). We did not

detect other proteins that have been identified by mass spectrometry in

immunoprecipitation experiments with VAL1-HA. We also observed a highly significant

enrichment of the chloroplast binding protein CRB. However, this is explained by the

enrichment of CRB in the total proteome of the experimental versus control samples

(Figure 7B). Comparisons of the proteins present in samples harvested from 1, 2 and 3

weeks old plants revealed that ATBMI1A was consistently among the most abundant

proteins immunoprecipitated with VAL1-HA (Figure 7C-E). The abundance of ATBMI1A

in the immunoprecipitated sample increased significantly from 1 week to 2 weeks, as

indicated by both the increase in the fold change between experimental and control

samples, and the statistical significance of the enrichment. This is probably a result of

the increase in the abundance of VAL1 between 1 and 2 weeks (Figure 6D,E), a result

which was confirmed by the increasing abundance of VAL1 in the immunoprecipitated

samples from different time points (Figure 7C-E). The parallel changes in the

531
532
Figure 7. VAL1-protein interactions are consistent during vegetative development
(A-E) Protein enrichment calculated via mass spectrometry. Fold change represents
the ratio of proteins purified from experimental (*VAL1-HA; val1-2; FRI-Sf2*) to
control (*val1-2; fri*) samples, a t-test *P*-value is represented on the y-axis. Red
dotted line indicates a *P*-value < 0.05 (resulting from the -log2 transformation of
the actual p-value; proteins above the line have a significant enrichment). Each
bubble represents an individual protein, the size or the bubble represents the
protein abundance averaged across the experimental and control samples. (A,
C-E) Proteins immunoprecipitated using an anti-HA antibody, (B) total proteome
samples.
543

abundance of VAL1-HA and ATBMI1A, and the absence of any major change in the

proteins associated with VAL1-HA in different samples, suggest that the binding

partners of VAL1 do not change significantly during shoot development.

To investigate overall trends in protein accumulation during vegetative

development we conducted a supervised clustering analysis of the *val1-2; VAL1-HA;*

FRI-Sf2 total proteome sample. We designed two clusters in which proteins either

increased or decreased from 1 to 2 to 3 weeks of development (see Methods for

details). The 50 proteins with the strongest and most consistent decreasing

developmental trend were significantly enriched for Gene Ontology terms related to

photosynthesis and carbon fixation (Figure S7 A,B). In contrast, the 50 proteins with the

highest increasing trend score were enriched for GO terms relating to water stress and

translation. The finding that younger plants invest more resources in photosynthesis is

consistent with a transcriptomic analysis of vegetative development in maize (Beydler et

al., 2016), and the enhanced photosynthetic capacity of juvenile plants at low light levels

(Lawrence et al., 2020).

Discussion

Plant life cycles are characterized by transitions between distinct developmental

phases. *VAL* genes have previously been shown to promote the switch from

embryogenesis to seed maturation (Suzuki et al., 2007; Yang et al., 2013a), and from

vegetative to reproductive growth (Qüesta et al., 2016; Yuan et al., 2016). The results

presented here demonstrate that *VAL1* and *VAL2* also regulate the intervening

transition from juvenile to adult stages of vegetative growth. *VAL* genes thus function as

a regulatory hub that coordinates developmental transitions throughout plant life cycles.

Regulation of miR156 expression by *VAL1/2*

Vegetative phase change is promoted by a temporal decline in miR156/miR157

expression (Wu and Poethig, 2006). When the level of miR156/miR157 falls below a

specific threshold, the de-repression of *SPL* genes initiates a switch to adult identity.

Previous work has shown that – with the exception of *SPL3* – the increase in *SPL*

transcript levels during development is entirely attributable to post-transcriptional

regulation by miR156/miR157 (He et al., 2018). Factors that control the timing of

vegetative phase change can therefore act in three ways: 1) by modifying the rate of

decline in miR156; 2) by constitutively increasing or decreasing the level of miR156; and

3) by constitutively increasing or decreasing the rate of transcription of *SPL* genes. Our

results suggest that *VAL1* and *VAL2* regulate vegetative phase change both by

constitutively decreasing the level of miR156 and by repressing *SPL* gene expression

independently of miR156.

Evidence that *VAL1/VAL2* constitutively regulate the level of miR156 was

provided by the phenotype of plants deficient for *VAL1* and *VAL2*, VAL1-DNA binding

patterns and from the phenotype of plants expressing a *MIR156C* transgene lacking

VAL-binding sites. We found that although *VAL1* expression increases in the shoot apex

as plants develop, *val1; val2* double mutants displayed only a slight decrease in the rate

at which miR156 declines. Instead, *val1; val2* double mutants exhibited a significant

increase in the level of miR156 at every stage of vegetative development we examined.

Consistent with this result, *mir156a; mir157a; mir157c* triple mutants transformed with a

MIR156C transgene lacking VAL-binding sites had elevated levels of miR156 relative to

the wild type *MIR156C* control, but displayed the same temporal decrease in miR156 as

control plants.

We have proposed that the decrease in *MIR156A/C* expression during shoot

development may be attributable to the stochastic replacement of H3K27ac and

H3K4me3 by H3K27me3 (Xu et al., 2018, 2016b). The observation that the increase in

miR156 expression in *val1; val2* is associated with a decrease in the level of H3K27me3

at *MIR156C* supports this hypothesis, in that it shows that H3K27me3 is associated with

low levels of *MIR156C* expression. The difference in the phenotype of plants doubly

mutant for *val1-2* and mutations the PRC2 histone methyltransferases, *SWN* and *CLF*,

and between *val1-2* and a mutation in the histone deacetylase *HDA19*, supports the

hypothesis that the expression of *MIR156C* is regulated by a combination of histone

marks. *val1-2; swn-3* and *val1-2; clf-29* double mutants have approximately the same

amount of miR156 as *val1-2* single mutants, which is expected given that these genes

are involved in the same process, namely the deposition of H3K27me3. In contrast,

hda19-3 significantly enhanced the level of miR156 compared to *val1-2*. This type of

genetic interaction is predicted if miR156 expression is dependent on histone

modifications aside from H3K27me3.

The evidence that *VAL1/VAL2* regulate the level, but not the temporal expression

pattern, of miR156 leaves open the question of how its temporal pattern arises. Several

other chromatin regulators, such as the CHD3 nucleosome remodeler PICKLE (Xu et

al., 2016b), the PRC-accessory protein LHP1 (Cui et al., 2020), the SWI/SNF2

chromatin remodeler BRAHMA (Xu et al., 2016c), and the histone 2 regulators ARP6

and HTA9/11 (Choi et al., 2016; Xu et al., 2018) have been found to play a role in the

expression of miR156. Furthermore, transcription factors including AGL15/18

(Serivichyaswat et al., 2015), MYB33 (Guo et al., 2017), and members of the *NF-Y*

family (Wei et al., 2017; Zhao et al., 2020), also regulate miR156 expression. It is

possible that the temporal expression pattern of miR156 is a consequence of complex

interactions between these diverse factors, rather than being dependent on a single

class of regulator, such as *VAL1/2*.

<u>*VAL* genes and PRC1 activity</u>

VAL1/2 are thought to be necessary for the recruitment of PRC1 to target loci,

where it represses gene expression via PRC2-dependent and independent mechanisms

(Baile et al., 2020; Yuan et al., 2020; Zhou et al., 2017). We found that *VAL1/2*

accelerate vegetative phase change by repressing the expression of *MIR156A/C*, and

by indirectly promoting the expression of *SPL9*, a target of miR156. These results are

consistent with a previous study (Pico et al., 2015), which showed that *VAL1/2* and the

PRC1 components, AtBMI1A/B, repress *MIR156A/C* expression. However, the PRC1

components EMBRYONIC FLOWER1 (Pico et al., 2015) and RING1A/B (Li et al., 2017)

have also been reported to repress the expression of *SPL9* independently of miR156.

These latter effects delay vegetative phase change, which is the exact opposite of the

effect produced by PRC1-mediated repression of *MIR156A/C*. Together, these results

indicate that PRC1 can operate at different points within a regulatory pathway, or in

interacting regulatory pathways, to modulate the output of the pathway or pathways. If

the genes repressed by PRC1 have different functions—as in the case of miR156 and

its *SPL* targets—then the functional significance of a particular level of PRC1 activity at

a particular locus can be difficult to predict. These results also support the hypothesis

that there may be different forms of PRC1, which target different genes. Moreover, our

finding that H2AK119ub deposition at *SPL9* was unaffected in *val1-5(sd); val2-3*

suggests that VAL1/2 may not be universally required for PRC1-recruitment.

In this regard, it is interesting that although *MIR156A* and *MIR156C* are close

paralogs and have similar expression patterns, previous studies (Xu et al., 2018,

2016b), and the results presented here, indicate that these genes are differentially

sensitive to mutations that affect the activity of PRC2 and PRC1. For example, we found

that *val1; val2* mutants display a greater reduction in H2AK119ub and H3K27me3 at

MIR156C than at *MIR156A*. Our findings align with the results of ChIP-seq studies of

H2AK119ub and H3K27me3 in *atbmi1a/b/c* (Zhou et al., 2017) and *val1-2; val2-3* (Yuan

et al., 2020) mutants. The implication of these observations is that *MIR156C* expression

is more dependent on PRC1 and PRC2 activity than *MIR156A*. Defining the molecular

basis for this difference could provide important insights into factors that influence

epigenetic regulation in plants.

Unlike in animals, where PRC1 binding requires the prior deposition of

H3K27me3, in plants PRC1 is thought to promote PRC2 activity at the majority of its

targets (Calonje, 2014; Zhou et al., 2017). This model is supported by our results. In the

case of *MIR156C*, we found that in 1 week old plants loss of *VAL1/2* significantly

reduced H2AK119ub, but not H3K27me3. However, by 3 weeks both H2AK119ub and

H3K27me3 were reduced in *val1/2* mutants. These results suggest that VAL1/2 and, by

inference, PRC1 are present at *MIR156C* prior to PRC2, and promote the activity of

PRC2.

In addition to their roles in vegetative phase change, *VAL* genes regulate multiple

nodes of the flowering time (Jing et al., 2019; Qüesta et al., 2016; Yuan et al., 2016)

and seed development (Chen et al., 2018, 2020; Suzuki et al., 2007; Yang et al., 2013a)

networks. The co-option of *VAL* activity throughout genetic networks thus appears

critical to coordinating plant developmental transitions. Despite the centrality of *VAL*

function to the control of developmental timing, the persistent and robust pattern of

MIR156A/C expression in *val* mutant plants emphasizes the complexity of temporal

regulation in plants.

Acknowledgements

We thank members of the R.S.P laboratory and Doris Wagner (University of

Pennsylvania) for helpful discussions, Daniel Martinez and Ryan McCarthy (University

of Pennsylvania) for assistance with histology and chromatin sonication, the Arabidopsis

Biological Resource Center for T-DNA insertion lines and Julia Qüesta and Caroline

Dean (John Innes Center) for the kind donation of *VAL1-HA; val1-2; FRI-Sf2* seed. This

project was funded by NIH Grant GM051893 (to R.S.P). S.S. gratefully acknowledges

Merck/MSD for the Umberto Mortari Award (2019), the Japan Agency for Medical

Research and Development (AMED) and the New York Academy of Sciences (NYAS)

for supporting his lab in aging research.

Author Contributions

Conceptualization: J.P.F., R.S.P.; Methodology: J.P.F., R.S.P.; Investigation: J.P.F.,

J.H., V.J.C., S.S.; Formal analysis: J.P.F., S.S.; Resources: J.P.F., J.H.; Visualization:

J.P.F., S.S.; Writing - original draft: J.P.F.; Writing - review & editing: J.P.F., J.H., S.S.,

R.S.P.; Supervision: R.S.P.; Project administration: R.S.P.; Funding acquisition: R.S.P.

Declaration of Interests

The authors declare no competing interests.

Figure titles and legends

Figure 1. *VAL* genes redundantly regulate vegetative phase change

(C)　　Schematic of *val1* and *val2* alleles used in this study – grey boxes represent

UTRs, black boxes represent exons, red triangles represent T-DNA insertions,

red line represents EMS-induced base substitution.

(D)　　Phenotypes at 21 DAG in LD conditions, scale bar = 5mm.

(C-E)　Quantitative analysis of vegetative development. Statistically distinct genotypes

were identified by one-way ANOVA with *post hoc* Tukey multiple comparison test

(letters indicate statistically distinct groups $P < 0.05$; for (E) comparisons were

made at leaf 5), all plants grown in LD. (C,D) Boxes display the interquartile

range (IQR) (boxes), median (lines) and values beyond 1.5*IQR (whiskers);

mean values are marked by ◆. (E) Colored lines represent the mean and black

bars the SEM. Sample sizes (C, D) 21-46, (E) 13-46.

Figure 2. *VAL* genes function predominantly as quantitative regulators of miR156

expression

(A, B) qRT-PCR analyses of gene expression. (A) Shoot apices with leaf primordia (LP)

≥ 1mm removed at 1, 2 and 3 weeks. (B) Isolated LP 0.5-1mm in size. All plants

were grown in SD conditions. Each data point represents a biological replicate

and is the average of three technical replicates. Coloured lines represent the

mean and black bars mean±s.e.m. Asterisks represent significant differences

between WT and *val* mutants at the same time point, calculated by an unpaired

two-tailed *t*-test with a Bonferroni correction for multiple comparisons (* $P <$

0.025; ** $P < 0.005$; *** $P < 0.0005$).

Figure 3. *VAL* genes regulate miR156 activity via-PRC2 and PRC1

(B)   Schematics of the primer locations used for ChIP-qPCR. Blue and green bars

represent sequences encoding the mature miRNA and miRNA hairpin

respectively.

(B)   Temporal analysis of H3K27me3 by ChIP-qPCR. Each data point represents a

biological replicate and is the average of three technical replicates. Lines

represent the mean, bars represent the mean±s.e.m., asterisks represent

significant differences between WT and *val1-5(sd); val2-3* at the same time point,

calculated by an unpaired two-tailed *t*-test (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).

H3K2me3 values are relative to H3 and normalised to *STM* as an internal control.

Plants were grown in SD conditions.

(C, D) Phenotypes in LD. (D) Photographs taken at 21DAG, scale bar = 5mm. (E)

Asterisks represent significant differences to either WT or *val1-2*, calculated by

unpaired two-tailed *t*-test with Bonferroni correction for multiple comparisons (***

$P < 0.00033$). Sample size 22-46.

(G) qRT-PCR analyses of gene expression in shoot apices with LP $\geq$ 1mm removed

at 1, 2 and 3 weeks. Each data point represents a biological replicated and is the

average of three technical replicates. Coloured lines represent the mean and

black bars represent the mean±s.e.m. Asterisks represent significant differences

between *val1-2* and *val1-2; clf-29/val1-2; swn-3/val1-2; hda19-3* at the same time

point, calculated by unpaired two-tailed *t*-test with Bonferroni correction for

multiple comparisons (** $P < 0.0033$). Plants were grown in SD conditions.

(H) Temporal analysis of H2AK119ub by ChIP-qPCR, values are relative to input and

normalised to *ACT7* as an internal control. See (B) for details.

Figure 4. Loss of RY VAL-binding motifs at the *MIR156C* locus delays vegetative

phase change

(C) Schematic depicting the location of primers used for ChIP-qPCR, the sequences

encoding the miR156 hairpin and mature miRNA are coloured green and blue

respectively.

(D) Anti-HA ChIP-qPCR of WT Col control plants at 2W and *VAL1::VAL1-HA; val1-2;*

FRI-Sf2 plants at 1, 2 and 3W of growth. The data is presented as percentage

input normalized to *ACT7* and is displayed relative to WT. Each data point

represents a biological replicate and is the average of three technical replicates,

bars represent the mean and error bars the mean±s.e.m.

(C, D) Phenotypes of T1 plants transformed with *MIR156C* RY variants. Statistically

distinct genotypes were identified by one-way ANOVA with *post hoc* Tukey

multiple comparison test (letters indicate statistically distinct groups $P < 0.05$;

comparison in (C) made at leaf 5). *qm = mir156a mir156c mir157a mir157c*

quadruple mutant, tm = mir156a mir157a mir157c triple mutant, EV = empty

vector. (C) Colored lines represent the mean and black bars the mean±s.e.m.

Sample size (B) 26-52, (C) 37-51.

(E)   qRT-PCR analyses of gene expression in shoot apices with LP ≥ 1mm removed

at 1, 2 and 3 weeks. Each data point represents an independent homozygous T3

line and is the average of three technical replicates. Colored lines represent the

mean and black bars the mean±s.e.m. Asterisks represent significant differences

between *qm; MIR156C* and *qm; MIR156C -RY* lines at the same time point,

calculated by an unpaired two-tailed *t*-test with a Bonferroni correction for

multiple comparisons (* $P < 0.025$, ** $P < 0.005$).

(G)   Genetic interaction between *val1-2* and *MIR156C* RY deletions. Asterisks

represent significant differences between plants with wild type or null *VAL1*

alleles calculated by an unpaired two-tailed *t*-test (*** $P < 0.001$). Sample size

24-36. Phenotyping analyses were carried out in LD conditions, gene expression

and ChIP analyses were carried out in SD conditions.

Figure 5. *VAL1* regulates vegetative phase change by miR156-dependent and

independent mechanisms

(A-C)   Phenotypes of *val1-2* and *mir156a mir156c mir157a mir157c quadruple mutant*

lines. (A) Photographs taken at 17 DAG. Scale bar = 1mm. (B, C) Statistically

distinct genotypes were identified by one-way ANOVA with *post hoc* Tukey

multiple comparison test (letters indicate statistically distinct groups *P* < 0.05; (B)

comparisons made at leaf 5). Bars represent the mean±s.e.m. Sample size (B)

17-60, (C) 18-36. Silhouettes in B show representative leaf 5 shapes.

(F)   Expression of a miR156-resistant (*rSPL9*) reporter construct in WT, *val1-2* and

val1-5(sd) backgrounds. Scale bars = 1mm.

(G)   qRT-PCR analysis of gene expression in shoot apices with LP $\geq$ 1mm removed

at 1, 2 and 3 weeks. Each data point represents a biological replicate and is the

average of three technical replicates. Coloured lines represent the mean and

gray bars represent the mean±s.e.m. Asterisk represents significant difference

between WT and *val* mutant lines at the same time point, calculated by an

unpaired two-tailed *t*-test with a Bonferroni correction (* $P < 0.025$; ** $P < 0.005$).

All phenotypic analyses were carried out in LD conditions, the gene expression

analysis was carried out in SD conditions.

Figure 6. *VAL1* is expressed throughout vegetative development

(A, B) qRT-PCR analyses of gene expression. (A) Shoot apices with leaf primordia (LP)

$\geq$ 1mm removed at 1, 2 and 3 weeks. (B) Isolated LP 0.5-1mm in size.

(C-F) Analyses of a *VAL1-GUS* translational fusion. (C) Expression in whole plants,

scale bars = 1mm. (D) Expression in shoot apices following wax sectioning, scale

bars = 0.1mm. (E, F) VAL1 levels quantified by MUG assay in two independent

homozygous T3 lines. (A, B, E, F) Each data point represents a biological

replicate and is the average of three technical replicates. Lines represent the

mean and grey bars mean±s.e.m. Asterisks represent significant differences

between two continuous time points, calculated by an unpaired two-tailed *t*-test (*

$P < 0.05$; ** $P < 0.01$). All plants except the 3 DAG sample (C – long days) were

grown in SD conditions.

Figure 7. VAL1-protein interactions are consistent during vegetative development

(A-E) Protein enrichment calculated via mass spectrometry. Fold change represents

the ratio of proteins purified from experimental (*VAL1-HA; val1-2; FRI-Sf2*) to

control (*val1-2; fri*) samples, a t-test *P*-value is represented on the y-axis. Red

dotted line indicates a *P*-value < 0.05 (resulting from the -log2 transformation of

the actual p-value; proteins above the line have a significant enrichment). Each

bubble represents an individual protein, the size or the bubble represents the

protein abundance averaged across the experimental and control samples. (A,

C-E) Proteins immunoprecipitated using an anti-HA antibody, (B) total proteome

samples.

STAR methods

KEY RESOURCES TABLE

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to, and

will be fulfilled by, the Lead Contact Scott Poethig (spoethig@sas.upenn.edu).

Materials Availability

The *val1-5(sd), val1-5(sd); val2-3, val1-2; val2-4, val1-2; mir156ac mir157ac* and

VAL1::VAL1-GUS-VAL1 3'UTR lines described herein have been donated to the ABRC

(stock numbers CS72451-CS72455 respectively). All other plant lines and plasmids

generated during this project are available on request to Scott Poethig

(spoethig@sas.upenn.edu).

Data and Code Availability

The raw files from the experiments performed with mass spectrometry are deposited on

the free repository Chorus (https://chorusproject.org) at Project No. 1706.

EXPERIMENTAL MODELS AND SUBJECT DETAILS

Plant material and growth conditions

All stocks were grown in the Col-0 background. The following genetic lines have been

described previously: *val1-2* (SALK_088606)*, val2-1* (CS906036) (Suzuki et al., 2007)

(*val2-1* was backcrossed to Col-0 6 times from the original Wassilewskija parent); *val2-3*

(SALK_059568C) (Yang et al., 2013a); *clf-29* (SALK_021003) (Bouveret et al., 2006);

swn-3 (SALK_050195) (Chanvivattana et al., 2004); *hda19-3* (SALK_139445) (Kim et

al., 2008); *mir156a-2 mir156c-1 mir157a-1 mir157c-1* (He et al., 2018); *SPL9::rSPL9-*

GUS (Xu et al., 2016a); *val1-2 VAL1::VAL1-3xHA FRI-Sf2* (Qüesta et al., 2016). *val2-4*

(SALK_127961) was obtained from the Arabidopsis Biological Resources Center (Ohio

State University, OH, USA). Seeds were sown on fertilized Farfard #2 soil (Farfard) and

kept at 4°C for 3 days prior to transfer to a growth chamber, with the transfer day

counted as day 0 for plant age (0 DAG – days after germination). Plant were grown at

22°C under a mix of both white (USHIO F32T8/741) and red-enriched (Interlectric

F32/T8/WS Gro-Lite) fluorescent bulbs in either long day (16 hrs light/8 hrs. dark; 40

µmol m^{-2} s^{-1}) or short day (10 hrs light/14 hrs dark; 100 µmol m^{-2} s^{-1}) conditions.

Identification of the *val1-5(sd)* mutant

The *val1-5(sd)* allele was generated by exposing *mir157a-1; mir157c-1* seed to ethyl

methanesulfonate. An M2 mutant plant exhibiting delayed vegetative phase change was

backcrossed to the parental line and allowed to self. Tissue was pooled from 30 plants

exhibiting severely delayed vegetative phase change in the BC1F2 generation. DNA

was extracted via-SDS lysis and phenol-chloroform extraction and further purified using

Clean and Concentrator columns (Zymo Research). DNA concentration was determined

using a Qubit 2.0 Fluorometer (Invitrogen) and 1µg of DNA sheared using a Covaris S2

sonicator (Covaris) to produce 350bp inserts. Sequencing libraries were made following

the TruSeq DNA PCR-free LT Sample Prep Kit (Illumina) manufacturer's instructions.

Library quality and quantity was validated by Bioanalyser (Agilent) and KAPA analysis

(Kapa Biosystems). 100bp paired end reads were generated using a HiSeq 2500

(Illumina) and aligned to the TAIR10 reference genome following the default SHORE

pipeline (Ossowski et al., 2008). The SHOREmap backcross pipeline (Schneeberger et

al., 2009) using default options was employed to identify polymorphisms. Manual

inspection of allele frequencies in the mutant revealed a peak centered on the *VAL1*

locus. The causative mutation was confirmed by Sanger sequencing and

complementation assays. The mutant was backcrossed to a 'Traffic Line' (Wu et al.,

2015) with seed-fluorescent markers inserted adjacent to the *VAL1* locus to eliminate

additional closely linked polymorphisms. Consequently, the resultant *val1-5(sd)* plants

used in this study contain a linked *pNAP::RFP* insertion at 13,622,737bp on

Chromosome 2 (Crick strand).

Generation of transgenic plants

For RY-mutation lines the RY-site TGCATG was replaced by TGTATG. A 5kb *MIR156C*

genomic sequence including 2kb upstream of the transcriptional start site and 665bp

downstream of the end of the last exon was cloned into the binary vector pAGM4723

from the Golden Gate MoClo toolbox supplied by Addgene (www.addgene.org) (Weber

et al., 2011; Werner et al., 2012). For -RY3, -RY4 and -RY5 Q5 Site Directed

Mutagenesis Kit (New England Biolabs) was used to induce a substitution directly into

the expression vector. For -RY1, -RY2 and -RY12345 Gibson Assembly cloning (New

England Biolabs) was used to assemble individual fragments into the same backbone.

Golden Gate cloning was also used to generate *VAL1::VAL1-VAL1 3'UTR*,

VAL1::VAL1-GUS-VAL1 3'UTR, *VAL1::GUS-VAL1 3'UTR* lines. Promoter/5'UTR (*VAL1*

- 2.3kb), functional (*VAL1* – 3.8kb, *MIR156A* (Fouracre and Poethig, 2019)) and

3'UTR/terminator (*VAL1* – 2kb) sequences were cloned separately from *Arabidopsis*

gDNA, with Type II restriction sites removed where necessary. *GUS* and *AtuOCS*

sequences were obtained from the MoClo Plant Parts toolkit supplied by Addgene

(www.addgene.org) (Engler et al., 2014). Component parts were assembled using

Golden Gate cloning into the pAGM4723 binary vector, including green or red seed

fluorescent expression cassettes as selectable markers. Constructs were transformed

into *Arabidopsis* using the floral dip method. All primers used for cloning are included in

Table S1.

METHOD DETAILS

Quantification of gene expression

Tissue (either shoot apices with leaf primordia ≤1mm attached or isolated leaf primordia

0.5-1mm in size – as specified in the text) were ground in liquid nitrogen and total RNA

extracted using Trizol (Invitrogen) as per the manufacturer's instructions. RNA was

treated with RNAse-free DNAse (Qiagen) and 250ng-1µg of RNA was used for reverse

transcription using Superscript III (Invitrogen). Gene specific RT primers were used to

amplify miR156, miR157, miR172 and SnoR101 and a polyT primer for mRNA

amplification. Three-step qPCR of cDNA was carried out using SYBR-Green Master Mix

(Bimake). qPCR reactions were run in triplicate and an average was calculated.

Relative transcript levels were normalized to snoR101 (for miRNAs) and *ACT2* (for

mRNAs) and expressed as a ratio of expression to a specified control sample. The

qPCR primers used in this study are listed in Table S1.

Chromatin immunoprecipitation

Expanded leaves and roots were removed during tissue harvesting to produce samples

enriched for shoot apices and young leaves. For histone ChIP ~0.5g of fresh tissue per

antibody and for anti-HA ChIP ~5g of fresh tissue were harvested. Samples were fixed

in 1% formaldehyde under vacuum for 15 minutes. Cross-linked samples were ground

in liquid nitrogen and suspended in Honda buffer (0.44M sucrose, 1.25% ficoll, 2.5%

dextran 40, 20mM hepes pH 7.4, 10mM $MgCl_2$, 0.5% Triton, 5mM DTT, 1mM PMSF,

1% protease inhibitors), filtered through two layers of Miracloth (EMD Millipore), and

pelleted and washed thrice in Honda buffer. For histone ChIP, pellets were

resuspended in nuclei lysis buffer (50mM Tris-HCl pH 8, 10mM EDTA, 1% SDS, 1%

protease inhibitors), for anti-HA ChIP, pellets were resuspended in RIPA buffer (1X

PBS, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 1% protease inhibitors).

Samples were sonicated using a Fisherbrand Sonic Dismembrator (Fisher Scientific) 6x

10s at setting 3.2. ChIP samples were pre-cleared using Dynabeads Protein A

(Invitrogen). 2% was removed as input and samples were incubated overnight with 1%

antibody (for histone ChIP: anti-H3 (abcam ab1791), anti-H3K27me3 (EMD Millipore 07-

449), anti-H2AK119ub (Cell Signaling Technology 8240); for VAL1-HA ChIP: anti-HA

(Roche 11583816001)). Chromatin-antibody conjugates were purified with Dynabeads

Protein A and washed in low/high salt, lithium and TE buffers. Following reverse-

crosslinking DNA was isolated using a QIAquick PCR Purification Kit (Qiagen).

For ChIP-qPCR assays, three-step qPCR was carried out using SYBR-Green

Master Mix (Bimake). qPCR reactions were run in triplicate and an average was

calculated. Data were normalized and presented as follows: 1) For H3K27me3 – *STM*

was used as a control locus, data is presented as a ratio of (H3K27me3 gene of

interest/H3 gene of interest) to (H3K27me3 *STM*/H3 *STM*); 2) For H2AK119ub – *ACT7*

was used as a control locus, data is presented as a ratio of (H2AK119ub gene of

interest/input gene of interest) to (H2AK119ub *ACT7*/input *ACT7*); 3) For VAL1-HA –

UBQ10, ACT7 and *TA3* were used as control loci, data is presented as a ratio of

((VAL1-HA ChIP gene of interest/input gene of interest)/(VAL1-HA ChIP control/input

control)) relative to ((WT ChIP gene of interest/input gene of interest)/(WT ChIP

control/input control)). The qPCR primers used in this study are listed in Table S1.

Mass spectrometry

VAL1::VAL1-3xHA; val1-2; FRI-Sf2 and *val1-2* genotypes were used as experimental

and control samples respectively. Expanded leaves and roots were removed during

tissue harvesting to produce samples enriched for shoot apices and young leaves. 2-3g

of fresh tissue was harvested for immunoprecipitation, 0.2-0.6g was harvested for total

protein extraction. For immunoprecipitation, tissue was ground in liquid nitrogen,

suspended in IP buffer (20mM Tris-HCl pH 8, 150mM NaCl, 2.5mM EDTA, 0.5% Triton,

1% protease inhibitors, 1mM PMSF), rotated for 2 hours at 4°C and filtered through 2

layers of Miracloth. Anti-HA (Roche) conjugated Dynabeads Protein A (Invitrogen) were

added and samples were rotated overnight at 4°C. Beads were washed thrice with IP

buffer and proteins were purified for mass spectrometry using an S-Trap: Rapid

Universal MS Sample Prep (Protifi) following the manufacturer's instructions. For total

proteomes, ground tissue was suspended in 8M urea and rotated at room temperature

951 for 45 minutes. The samples were centrifuged thrice and the supernatant reduced with

952 DTT (final concentration 5mM) and alkylated with iodoacetamide 40 (final concentration

953 40mM) before overnight digest with trypsin. Samples were resuspended in 10 µl of

954 water + 0.1% TFA and loaded onto a Dionex RSLC Ultimate 300 (Thermo Scientific,

955 San Jose, CA, USA), coupled online with an Orbitrap Fusion Lumos (Thermo Scientific).

956 Chromatographic separation was performed with a two-column system, consisting of a

957 C_{18} trap cartridge (300 µm ID, 5 mm length) and a picofrit analytical column (75 µm ID,

958 25 cm length) packed in-house with reversed-phase Repro-Sil Pur C_{18}-AQ 3 µm resin.

959 Peptides were separated using a 90 min gradient (for the IP experiments) and 180 min

960 (for the full proteome experiment) from 2-28% buffer-B (buffer-A: 0.1% formic acid,

961 buffer-B: 80% acetonitrile + 0.1% formic acid) at a flow rate of 300 nl/min. The mass

962 spectrometer was set to acquire spectra in a data-dependent acquisition (DDA) mode.

963 Briefly, the full MS scan was set to 300-1200 m/z in the orbitrap with a resolution of

964 120,000 (at 200 m/z) and an AGC target of 5x10e5. MS/MS was performed in the ion

965 trap using the top speed mode (2 secs), an AGC target of 10e4 and an HCD collision

966 energy of 30. Raw files were searched using Proteome Discoverer software (v2.4,

967 Thermo Scientific) using SEQUEST as search engine using the SwissProt *Arabidopsis*

968 *thaliana* database. The search for total proteome included variable modifications of

969 methionine oxidation and N-terminal acetylation, and fixed modification of

970 carbamidomethyl cysteine. Trypsin was specified as the digestive enzyme. Mass

971 tolerance was set to 10 pm for precursor ions and 0.2 Da for product ions. Peptide and

972 protein false discovery rate was set to 1%. Data transformation, normalization and

973   statistical analysis using heteroscedastic t-test was performed as previously described

974   (Aguilan et al., 2020).

975   Proteins were sorted according to their descending or ascending linearity across

the three weeks. To do so, we used a custom score taking into account monotonic

trend, reproducibility across replicates and magnitude of change across weeks. The 50

proteins with the highest descending and ascending trend scores were used to identify

enriched GO terms for biological processes using the Fisher's Exact PANTHER

Overrepresentation Test (released 2020-07-28) and GO Ontology database DOI:

10.5281/zenodo.4081749 (released 2020-10-09). The *Arabidopsis thaliana* genome

was used as a reference list. Protein interaction maps for the same sets of 50 proteins

were made using the STRING app from Cytoscape (Shannon et al., 2003).

GUS staining and histology

Shoot apices and whole plants were fixed in 90% acetone on ice for 10 minutes,

washed with GUS staining buffer (5mM potassium ferricyanide and 5mM ferrocyanide in

0.1M PO_4 buffer) and incubated at 37°C overnight in GUS staining buffer with 2mM X-

Gluc. Embryos were placed directly in X-Gluc (GoldBio) GUS staining buffer and

incubated for 1hr. To quantify GUS activity, a 4-methylumbelliferyl b-D-glucuronide

(MUG) (Sigma-Aldrich) assay was carried out as previously described (He et al., 2018).

For histological observations individuals were fixed in FAA (3.7% formaldehyde),

dehydrated in an ethanol series and cleared using Histo-Clear (National Diagnostics).

Following embedding in Paraplast Plus (Sigma-Aldrich) 8µM sections were produced

using an HM 355 microtome (Microm) and visualized using an Olympus BX51

microscope with a DP71 camera attachment (Olympus).

QUANTIFICATION AND STATISTICAL ANALYSES

Details of all statistical analyses, including the type of statistical test, sample size,

replicate number and significance threshold, are included in the relevant figure legend.

For figures featuring boxplots, boxes display the IQR (boxes), median (lines), and

values beyond 1.5* IQR (whiskers); mean values are marked by a solid diamond (◆).

Statistical analyses were carried out using RStudio (RStudio Team, 2020) and Microsoft

Excel. Measurements of leaf shape were made using ImageJ (Schindelin et al., 2012).

1006

1007   **Supplemental Information titles and legends**

1008   **Figure S1. *val1-5(sd)* is an antimorphic allele**

1009   (A)   Sequence alignment of the B3 DNA-binding domain N-arm of *Arabidopsis* LAV

1010   family members and the maize ABI3 ortholog VP1. Numbers correspond to

1011   amino acid sequence numbers, colors correspond to the ClustalX amino acid

1012   color scheme. In the *val1-5(sd)* mutant a C>T base substitution converts an

1013   arginine to a cysteine.

1014   (B)   *val1-5(sd)* complementation test with the null *val1-2* allele

1015   (C)   Rescue of the *val1-5(sd)* abaxial trichome phenotype with a *VAL1* genomic

1016   sequence. Independent T1 lines are shown.

1017   (D)   Allele heterozygosity testing. (B-D) Boxes display the interquartile range (IQR)

1018   (boxes), median (lines) and values beyond 1.5*IQR (whiskers); mean values are

1019   marked by ◆. Samples sizes are displayed on the graph. Statistically distinct

1020   genotypes were identified by one-way ANOVA with *post hoc* Tukey multiple

1021   comparison test (letters indicate statistically distinct groups; $P < 0.05$), all plants

1022   grown in LD.

1023 (E)   Heteroblastic series of lines shown in Figure 1.

1024

Figure S2. *VAL* genes redundantly regulate miR156 expression

qRT-PCR analyses of gene expression in shoot apices with leaf primordia (LP) $\geq$

1mm removed at 1, 2 and 3 weeks. All plants were grown in SD conditions. Each

data point represents a biological replicate and is the average of three technical

replicates. Coloured lines represent the mean and black lines mean±s.e.m.

Asterisks represent significant differences between WT and *val* mutants at the

same time point, calculated by an unpaired two-tailed *t*-test with a Bonferroni

correction for multiple comparisons (* $P < 0.025$; ** $P < 0.005$; *** $P < 0.0005$).

1033

Figure S3. VAL1 binds consistently to *MIR156A* and *MIR156C* during vegetative

development

(A)   Schematic depicting the location of primers used for ChIP-qPCR, the sequences

encoding the miR156 hairpin and mature miRNA are coloured green and blue

respectively.

(B)   Anti-HA ChIP-qPCR of WT Col control plants at 2W and *VAL1::VAL1-HA; val1-2;*

FRI-Sf2 plants at 1, 2 and 3W of growth. The data is presented as percentage

input normalized to a control locus (*UBQ10* or *TA3*) and is displayed relative to

WT. Each data point represents a biological replicate and is the average of three

technical replicates, bars represent the mean and error bars the mean±s.e.m.

Plants were grown in SD conditions.

1045

Figure S4. RY sites are not required for the temporal decline of *MIR156C*

(A)   qRT-PCR analyses of gene expression in shoot apices with LP $\geq$ 1mm removed

at 1 and 2 weeks. Bars represent the average of three technical replicates for a

single biological replicate of pooled T1 plants, at least 15 independent T1 plants

were pooled for each sample. *qm = mir156a mir156c mir157a mir157c quadruple*

mutant, tm = mir156a mir157a mir157c triple mutant. Plants were grown in SD

conditions.

1053

Figure S5. *VAL* genes do not regulate *SPL9* chromatin state

(A, B) Temporal analysis of histone modification scalculated by ChIP-qPCR. Each data

point represents a biological replicate and is the average of three technical

replicates. Lines represent the mean and bars represent the mean±s.e.m., (A)

H3K2me3 values are relative to H3 and normalised to *STM* as an internal control.

(B) H2AK119ub values are relative to input and normalised to *ACT7* as an

internal control. Plants were grown in SD conditions.

1061

1062   **Figure S6. *VAL1* expression is dependent on genetic elements in the coding**

1063   **sequence of the gene**

1064   (A)   Torpedo-stage embryos of two-independent homozygous transgenic lines each

1065   expressing a transcriptional (top panel) or translational (bottom panel) *VAL1-*

1066   *GUS* reporter construct.

1067   (B)   Seedlings at 3 DAG. Each number designates an independent homozygous

1068   transgenic line. Arrow heads point to initiating lateral root primordia. All plants

1069   were grown in LD.

1070

1071   **Figure S7. Proteomic changes during vegetative development**

1072   (A,B) The 50 proteins with the strongest increasing or decreasing trend score during

1073   vegetative development in the experimental sample only (see Methods for

1074   details). (A) GO terms enriched within the 50 proteins that have the strongest

1075   increasing or decreasing trend score. (G) Interaction networks for each set of 50

1076   proteins, the darker the color the stronger the increasing (red) or decreasing

1077   (blue) trend score.

1078

1079   **Table S1. Primer sequences**

1080   This table includes all the primer sequences used in this study.

1081

1082   **Table S2. Proteins detected by mass spectrometry following anti-HA**

1083   **immunoprecipitation**

1084   Related to Figure S6.

1085   This table includes the raw mass spectrometry results and data processing for IP-

1086   samples.

1087

1088   **Table S3. Total proteome changes during vegetative development**

1089   Related to Figure S6.

1090   This table includes the raw mass spectrometry results and data processing for total

1091   proteome samples.

1092

REFERENCES

Aguilan, J.T., Kulej, K., and Sidoli, S. (2020). Guide for protein fold change and p-value
calculation for non-experts in proteomics. Mol. Omics *16*, 573–582.

Baile, F., Merini, W., Hidalgo, I., and Calonje, M. (2020). Dissection of PRC1 and PRC2
recruitment in Arabidopsis connects EAR repressome to PRC2 anchoring. BioRxiv
2020.08.28.271999.

Beydler, B., Osadchuk, K., Cheng, C.-L., Manak, J.R., and Irish, E.E. (2016). The Juvenile Phase of
Maize Sees Upregulation of Stress-Response Genes and Is Extended by Exogenous Jasmonic
Acid. Plant Physiol. *171*, 2648–2658.

Bouveret, R., Schönrock, N., Gruissem, W., and Hennig, L. (2006). Regulation of flowering time
by *Arabidopsis MSI1*. Development *133*, 1693–1702.

Bratzel, F., López-Torrejón, G., Koch, M., Del Pozo, J.C., and Calonje, M. (2010). Keeping Cell
Identity in Arabidopsis Requires PRC1 RING-Finger Homologs that Catalyze H2A
Monoubiquitination. Current Biology *20*, 1853–1859.

Calonje, M. (2014). PRC1 Marks the Difference in Plant PcG Repression. Molecular Plant *7*, 459–
471.

Carbonero, P., Iglesias-Fernández, R., and Vicente-Carbajosa, J. (2017). The AFL subfamily of B3
transcription factors: evolution and function in angiosperm seeds. Journal of Experimental
Botany *68*, 871–880.

Chanvivattana, Y., Bishopp, A., Schubert, D., Stock, C., Moon, Y.-H., Sung, Z.R., and Goodrich, J.
(2004). Interaction of Polycomb-group proteins controlling flowering in *Arabidopsis*.
Development *131*, 5263–5276.

Chen, N., Veerappan, V., Abdelmageed, H., Kang, M., and Allen, R.D. (2018). HSI2/VAL1 Silences
AGL15 to Regulate the Developmental Transition from Seed Maturation to Vegetative Growth
in Arabidopsis. Plant Cell *30*, 600–619.

Chen, N., Wang, H., Abdelmageed, H., Veerappan, V., Tadege, M., and Allen, R.D. (2020).
HSI2/VAL1 and HSL1/VAL2 function redundantly to repress DOG1 expression in Arabidopsis
seeds and seedlings. New Phytologist *227*, 840–856.

Chhun, T., Chong, S.Y., Park, B.S., Wong, E.C.C., Yin, J.-L., Kim, M., and Chua, N.-H. (2016). HSI2
Repressor Recruits MED13 and HDA6 to Down-Regulate Seed Maturation Gene Expression
Directly During Arabidopsis Early Seedling Growth. Plant and Cell Physiology *57*, 1689–1706.

Choi, K., Kim, J., Muller, S.Y., Oh, M., Underwood, C., Henderson, I., and Lee, I. (2016).
Regulation of microRNA-mediated developmental changes by the SWR1 chromatin remodeling
complex. Plant Physiol *171*, 1128–1143.

Cui, Y., Cheng, J., Ruan, S., Qi, P., Liu, W., Bian, H., Ye, L., Zhang, Y., Hu, J., Dong, G., et al. (2020).
The heterochronic gene Oryza sativa LIKE HETEROCHROMATIN PROTEIN 1 modulates
miR156b/c/i/e levels. Journal of Integrative Plant Biology *62*, 1839–1852.

Engler, C., Youles, M., Gruetzner, R., Ehnert, T.M., Werner, S., Jones, J.D., Patron, N.J., and
Marillonnet, S. (2014). A golden gate modular cloning toolbox for plants. ACS Synth. Biol. *3*,
839–843.

Fouracre, J.P., and Poethig, R.S. (2019). Role for the shoot apical meristem in the specification
of juvenile leaf identity in *Arabidopsis*. Proc Natl Acad Sci USA *116*, 10168–10177.

German, M.A., Pillay, M., Jeong, D.-H., Hetawal, A., Luo, S., Janardhanan, P., Kannan, V.,
Rymarquis, L.A., Nobuta, K., German, R., et al. (2008). Global identification of microRNA–target
RNA pairs by parallel analysis of RNA ends. Nature Biotechnology *26*, 941–946.

Goodrich, J., Puangsomlee, P., Martin, M., Long, D., Meyerowitz, E.M., and Coupland, G. (1997).
A Polycomb-group gene regulates homeotic gene expression in Arabidopsis. Nature *386*, 44–51.

Gou, J.Y., Felippes, F.F., Liu, C.J., Weigel, D., and Wang, J.W. (2011). Negative regulation of
anthocyanin biosynthesis in Arabidopsis by a miR156-targeted SPL transcription factor. Plant
Cell *23*, 1512–1522.

Guo, C., Xu, Y., Shi, M., Lai, Y., Wu, X., Wang, H., Zhu, Z., Poethig, R.S., and Wu, G. (2017).
Repression of miR156 by miR159 regulates the timing of the juvenile-to-adult transition in
Arabidopsis. Plant Cell *29*, 1293–1304.

Guo, X., Hou, X., Fang, J., Wei, P., Xu, B., Chen, M., Feng, Y., and Chu, C. (2013). The rice
GERMINATION DEFECTIVE 1, encoding a B3 domain transcriptional repressor, regulates seed
germination and seedling development by integrating GA and carbohydrate metabolism. The
Plant Journal *75*, 403–416.

He, J., Xu, M., Willmann, M.R., McCormick, K., Hu, T., Yang, L., Starker, C.G., Voytas, D.F.,
Meyers, B.C., and Poethig, R.S. (2018). Threshold-dependent repression of *SPL* gene expression

by miR156/miR157 controls vegetative phase change in *Arabidopsis thaliana*. PLoS Genet. *14*,
e1007337.

Huijser, P., and Schmid, M. (2011). The control of developmental phase transitions in plants.
Development *138*, 4117–4129.

Jang, I.-C., Chung, P.J., Hemmes, H., Jung, C., and Chua, N.-H. (2011). Rapid and Reversible Light-
Mediated Chromatin Modifications of Arabidopsis *Phytochrome A* Locus. Plant Cell *23*, 459–
470.

Jing, Y., Guo, Q., and Lin, R. (2019). The B3-Domain Transcription Factor VAL1 Regulates the
Floral Transition by Repressing *FLOWERING LOCUS T*. Plant Physiol. *181*, 236–248.

Kagaya, Y., Ohmiya, K., and Hattori, T. (1999). RAV1, a novel DNA-binding protein, binds to
bipartite recognition sequence through two distinct DNA-binding domains uniquely found in
higher plants. Nucleic Acids Research *27*, 470–478.

Kim, K.-C., Lai, Z., Fan, B., and Chen, Z. (2008). *Arabidopsis* WRKY38 and WRKY62 Transcription
Factors Interact with Histone Deacetylase 19 in Basal Defense. Plant Cell *20*, 2357–2371.

Lawrence, E.H., Springer, C.J., Helliker, B.R., and Poethig, R.S. (2020). MicroRNA156-mediated
changes in leaf composition lead to altered photosynthetic traits during vegetative phase
change. New Phytologist.

Leichty, A.R., and Poethig, R.S. (2019). Development and evolution of age-dependent defenses
in ant-acacias. Proc Natl Acad Sci USA *116*, 15596–15601.

Li, J., Wang, Z., Hu, Y., Cao, Y., and Ma, L. (2017). Polycomb Group Proteins RING1A and RING1B
Regulate the Vegetative Phase Transition in Arabidopsis. Frontiers in Plant Science *8*, 867.

Lopez-Vernaza, M., Yang, S., Müller, R., Thorpe, F., de Leau, E., and Goodrich, J. (2012).
Antagonistic Roles of SEPALLATA3, FT and FLC Genes as Targets of the Polycomb Group Gene
CURLY LEAF. PLOS ONE *7*, e30715.

Mao, Y.-B., Liu, Y.-Q., Chen, D.-Y., Chen, F.-Y., Fang, X., Hong, G.-J., Wang, L.-J., Wang, J.-W., and
Chen, X.-Y. (2017). Jasmonate response decay and defense metabolite accumulation
contributes to age-regulated dynamics of plant insect resistance. Nature Communications *8*,
13925.

Molitor, A.M., Bu, Z., Yu, Y., and Shen, W.-H. (2014). Arabidopsis AL PHD-PRC1 Complexes
Promote Seed Germination through H3K4me3-to-H3K27me3 Chromatin State Switch in
Repression of Seed Developmental Genes. PLOS Genetics *10*, e1004091.

Ossowski, S., Schneeberger, K., Clark, R.M., Lanz, C., Warthmann, N., and Weigel, D. (2008).
Sequencing of natural strains of Arabidopsis thaliana with short reads. Genome Res *18*, 2024–
2033.

Pico, S., Ortiz-Marchena, M.I., Merini, W., and Calonje, M. (2015). Deciphering the role of
Polycomb Repressive Complex 1 (PRC1) variants in regulating the acquisition of flowering
competence in Arabidopsis. Plant Physiology *168*, 1286–1297.

Qüesta, J.I., Song, J., Geraldo, N., An, H., and Dean, C. (2016). *Arabidopsis* transcriptional
repressor VAL1 triggers Polycomb silencing at *FLC* during vernalization. Science *353*, 485–488.

Ronemus, M., Vaughn, M.W., and Martienssen, R.A. (2006). MicroRNA-Targeted and Small
Interfering RNA–Mediated mRNA Degradation Is Regulated by Argonaute, Dicer, and RNA-
Dependent RNA Polymerase in *Arabidopsis*. Plant Cell *18*, 1559–1574.

RStudio Team (2020). RStudio: Integrated Development Environment for R (Boston, MA:
RStudio, PBC).

Sasnauskas, G., Kauneckaitė, K., and Siksnys, V. (2018). Structural basis of DNA target
recognition by the B3 domain of Arabidopsis epigenome reader VAL1. Nucleic Acids Research
46, 4316–4324.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
image analysis. Nature Methods *9*, 676–682.

Schneeberger, K., Ossowski, S., Lanz, C., Juul, T., Petersen, A.H., Nielsen, K.L., Jorgensen, J.E.,
Weigel, D., and Andersen, S.U. (2009). SHOREmap: simultaneous mapping and mutation
identification by deep sequencing. Nat Methods *6*, 550–551.

Serivichyaswat, P., Ryu, H.S., Kim, W., Kim, S., Chung, K.S., Kim, J.J., and Ahn, J.H. (2015).
Expression of the floral repressor miRNA156 is positively regulated by the AGAMOUS-like
proteins AGL15 and AGL18. Mol Cells *38*, 259–266.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski,
B., and Ideker, T. (2003). Cytoscape: A Software Environment for Integrated Models of
Biomolecular Interaction Networks. Genome Research *13*, 2498–2504.

Suzuki, M., Kao, C.Y., and McCarty, D.R. (1997). The conserved B3 domain of VIVIPAROUS1 has a
cooperative DNA binding activity. Plant Cell *9*, 799–807.

Suzuki, M., Wang, H.H.-Y., and McCarty, D.R. (2007). Repression of the *LEAFY COTYLEDON 1/B3*
Regulatory Network in Plant Embryo Development by *VP1/ABSCISIC ACID INSENSITIVE 3-LIKE* B3
Genes. Plant Physiol. *143*, 902–911.

Telfer, A., Bollman, K.M., and Poethig, R.S. (1997). Phase change and the regulation of trichome
distribution in *Arabidopsis thaliana*. Development *124*, 645–654.

Tian, R., Wang, F., Zheng, Q., Niza, V.M.A.G.E., Downie, A.B., and Perry, S.E. (2020). Direct and
indirect targets of the arabidopsis seed transcription factor ABSCISIC ACID INSENSITIVE3. The
Plant Journal *103*, 1679–1694.

Tsukagoshi, H., Morikami, A., and Nakamura, K. (2007). Two B3 domain transcriptional
repressors prevent sugar-inducible expression of seed maturation genes in *Arabidopsis*
seedlings. Proc Natl Acad Sci USA *104*, 2543–2547.

Veerappan, V., Wang, J., Kang, M., Lee, J., Tang, Y., Jha, A.K., Shi, H., Palanivelu, R., and Allen,
R.D. (2012). A novel HSI2 mutation in Arabidopsis affects the PHD-like domain and leads to
derepression of seed-specific gene expression. Planta *236*, 1–17.

Wang, F., and Perry, S.E. (2013). Identification of direct targets of FUSCA3, a key regulator of
Arabidopsis seed development. Plant Physiology *161*, 1251–1264.

Wang, L., Zhou, C.-M., Mai, Y.-X., Li, L.-Z., Gao, J., Shang, G.-D., Lian, H., Han, L., Zhang, T.-Q.,
Tang, H.-B., et al. (2019). A spatiotemporally regulated transcriptional complex underlies
heteroblastic development of leaf hairs in Arabidopsis thaliana. The EMBO Journal *38*, e100063.

Weber, E., Engler, C., Gruetzner, R., Werner, S., and Marillonnet, S. (2011). A Modular Cloning
System for Standardized Assembly of Multigene Constructs. PLOS ONE *6*, e16765.

Wei, Q., Ma, C., Xu, Y., Wang, T., Chen, Y., Lü, J., Zhang, L., Jiang, C.-Z., Hong, B., and Gao, J.
(2017). Control of chrysanthemum flowering through integration with an aging pathway.
Nature Communications *8*, 829.

Werner, S., Engler, C., Weber, E., Gruetzner, R., and Marillonnet, S. (2012). Fast track assembly
of multigene constructs using Golden Gate cloning and the MoClo system. Bioeng Bugs *3*, 38–
43.

Wu, G., and Poethig, R.S. (2006). Temporal regulation of shoot development in *Arabidopsis*
thaliana by miR156 and its target *SPL3*. Development *133*, 3539–3547.

Wu, G., Park, M.Y., Conway, S.R., Wang, J.W., Weigel, D., and Poethig, R.S. (2009). The
sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. Cell
138, 750–759.

Wu, G., Rossidivito, G., Hu, T., Berlyand, Y., and Poethig, R.S. (2015). Traffic lines: new tools for
genetic analysis in *Arabidopsis thaliana*. Genetics *200*, 35–45.

Xie, Y., Zhang, Y., Han, J., Luo, J., Li, G., Huang, J., Wu, H., Tian, Q., Zhu, Q., Chen, Y., et al.
(2018). The Intronic *cis* Element *SE1* Recruits *trans*-Acting Repressor Complexes to Repress the
Expression of *ELONGATED UPPERMOST INTERNODE1* in Rice. Molecular Plant *11*, 720–735.

Xu, M., Hu, T., Zhao, J., Park, M.Y., Earley, K.W., Wu, G., Yang, L., and Poethig, R.S. (2016a).
Developmental functions of miR156-regulated *SQUAMOSA PROMOTER BINDING PROTEIN-LIKE*
(*SPL*) genes in *Arabidopsis thaliana*. PLoS Genet. *12*, e1006263.

Xu, M., Leichty, A.R., Hu, T., and Poethig, R.S. (2018). H2A.Z promotes the transcription of
MIR156A and *MIR156C* in *Arabidopsis* by facilitating the deposition of H3K4me3. Development
145, dev152868.

Xu, M.L., Hu, T.Q., Smith, M.R., and Poethig, R.S. (2016b). Epigenetic regulation of vegetative
phase change in *Arabidopsis*. Plant Cell *28*, 28–41.

Xu, Y.M., Guo, C.K., Zhou, B.Y., Li, C.L., Wang, H.S., Zheng, B., Ding, H., Zhu, Z.J., Peragine, A.,
Cui, Y.H., et al. (2016c). Regulation of vegetative phase change by SWI2/SNF2 chromatin
remodeling ATPase BRAHMA. Plant Physiol *172*, 2416–2428.

Yang, C., Bratzel, F., Hohmann, N., Koch, M., Turck, F., and Calonje, M. (2013a). VAL- and
AtBMI1-Mediated H2Aub Initiate the Switch from Embryonic to Postgerminative Growth in
Arabidopsis. Current Biology *23*, 1324–1329.

Yang, H., Howard, M., and Dean, C. (2014). Antagonistic Roles for H3K36me3 and H3K27me3 in
the Cold-Induced Epigenetic Switch at Arabidopsis FLC. Current Biology *24*, 1793–1797.

Yang, L., Xu, M., Koo, Y., He, J., and Poethig, R.S. (2013b). Sugar promotes vegetative phase
change in *Arabidopsis thaliana* by repressing the expression of *MIR156A* and *MIR156C*. Elife *2*,
e00260.

Yu, S., Cao, L., Zhou, C.M., Zhang, T.Q., Lian, H., Sun, Y., Wu, J.Q., Huang, J.R., Wang, G.D., and
Wang, J.W. (2013). Sugar is an endogenous cue for juvenile-to-adult phase transition in plants.
Elife *2*, e00269.

Yuan, L., Song, X., Zhang, L., Yu, Y., Liang, Z., Lei, Y., Ruan, J., Tan, B., Liu, J., and Li, C. (2020). The
transcriptional repressors VAL1 and VAL2 recruit PRC2 for genome-wide Polycomb silencing in
Arabidopsis. Nucleic Acids Research.

Yuan, W., Luo, X., Li, Z., Yang, W., Wang, Y., Liu, R., Du, J., and He, Y. (2016). A cis cold memory
element and a trans epigenome reader mediate Polycomb silencing of FLC by vernalization in
Arabidopsis. Nature Genetics *48*, 1527–1534.

Zeng, X., Gao, Z., Jiang, C., Yang, Y., Liu, R., and He, Y. (2020). HISTONE DEACETYLASE 9
Functions with Polycomb Silencing to Repress *FLOWERING LOCUS C* Expression. Plant Physiol.
182, 555–565.

Zhao, H., Lin, K., Ma, L., Chen, Q., Gan, S., and Li, G. (2020). Arabidopsis NUCLEAR FACTOR Y A8
inhibits the juvenile-to-adult transition by activating transcription of MIR156s. Journal of
Experimental Botany *71*, 4890–4902.

Zhou, Y., Tan, B., Luo, M., Li, Y., Liu, C., Chen, C., Yu, C.-W., Yang, S., Dong, S., Ruan, J., et al.
(2013). HISTONE DEACETYLASE19 interacts with HSL1 and participates in the repression of seed
maturation genes in Arabidopsis seedlings. Plant Cell *25*, 134–148.

Zhou, Y., Romero-Campero, F.J., Gómez-Zambrano, Á., Turck, F., and Calonje, M. (2017). H2A
monoubiquitination in Arabidopsis thaliana is generally independent of LHP1 and PRC2 activity.
Genome Biology *18*, 69.

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299