Acute and chronic effects of exams week on cortisol production in undergraduate students

Running head: Cortisol in undergraduates

Borghi, Filipy[1]; da Silva, Priscila Cristina[1]; Canova, Fernando[1]; Souza, Aglecio Luiz[2]; Arouca,

Aline Barbedo[1]; Grassi-Kassisse, Dora Maria[1]*

[1]LABEEST - Laboratory of Stress Study, Department of Structural and Functional Biology.

Institute of Biology, University of Campinas – UNICAMP, Campinas, São Paulo, Brazil

[2]Unidade Metabólica, Faculdade de Ciências Médicas, Universidade Estadual de Campinas

(UNICAMP), São Paulo, Brazil

*Corresponding author

Laboratory of Stress Study, Department of Structural and Functional Biology, Biology Institute -

University of Campinas (UNICAMP), Cidade Universitária Zeferino Vaz, Monteiro Lobato

Street, 255, ZIP CODE: 13083-862, Campinas, SP, Brazil, PHONE: +55-19-3521-6186, FAX:

+55-19-3521-6185.

e-mail: doramgk@unicamp.br

ABSTRACT

Undergraduate students experience many levels of stress throughout their university trajectory.

The pressure to achieve good results highlight the exams week in the Universities as a stimulus

that triggers psychosocial stress and increased release of cortisol. Considering the increasingly

experience an undue amount of stress, we aimed in this study to investigate the short- and long-

term effects of exams week on HPA axis in undergraduates by the cortisol production. Twenty-

eight undergraduates aged 18-24 years from biological sciences during the final exams' week at

the end of the school year collected hair and saliva for cortisol measures. Hair cortisol was

significantly higher in exams month. They exhibited cortisol rhythmicity and preserved CAR

(cortisol awakening response) even under psychosocial stress. The exams week is a trigger for

psychosocial stress, however, did not generate significant short-term changes in cortisol

rhythmicity. Thus, it is evident that young people adapt to the stressful stimulus, avoiding a

possible trigger for mental illness.

Keywords: Cortisol; hair; rhythmicity, saliva; psychosocial stress; undergraduate students.

INTRODUCTION

Stress can be defined as a mismatch between demands and the perceived ability to cope with the stressor (Gallagher et al., 2014; Moore, Truscott, St Clair, & Klingborg, 2007). It is founded on an individual's perceptions and cognitive appraisal of his or her situation and cannot be predicted objectively (Moore et al., 2007). In general, the stress response is adaptive and has a crucial role in the adaptation to external demands (García-León, Pérez-Mármol, Gonzalez-Pérez, García-Ríos, & Peralta-Ramírez, 2019). The key mediator of this response is considered to be the neuroendocrine hypothalamus-pituitary-adrenal (HPA) axis and its main downstream effector, the glucocorticoid cortisol (Stalder et al., 2017).

Systemic cortisol levels are highly variable, which is due to the circadian rhythm, acute stress and pulsatile secretion (Wester & van Rossum, 2015). Measurements of cortisol in saliva, blood or urine only assess acute short sampling periods, usually 12 or 24 h, limiting the observation of long-term chronic effects, in addition to depending on the patient's correct adherence to the collection protocols (Stalder et al., 2017; Wester & van Rossum, 2015). Hair analysis has been used for decades in forensic science to measure exposure to environmental toxins and drugs and nowadays has gained attention as a viable biomarker for chronic stress and long-term activity of the HPA axis (D'Anna-Hernandez, Ross, Natvig, & Laudenslager, 2011; Sauvé, Koren, Walsh, Tokmakejian, & Van Uum, 2007; Wester & van Rossum, 2015). In addition, the hair cortisol evaluated hair segments grew in the past, making this technique particularly interesting to assess endocrine activity of past events (Stalder et al., 2017).

Undergraduate students experience many levels of stress throughout their university trajectory. There is empirical evidence that the stressful experience at the university was linked to reduced academic performance, low grade averages, and low rates of graduation and higher dropout, in addition to increased psychological distress (Dwyer & Cummings, 2001; Elias, Ping, & Abdullah, 2011; Portoghese et al., 2019). The pressure to achieve good results highlight the exams week in the Universities as a stimulus that triggers psychosocial stress and increased release of cortisol (Karatsoreos & McEwen, 2014; Liston, McEwen, & Casey, 2009). Considering that undergraduate students increasingly experience an undue amount of stress, we aimed in this

study to investigate the short- and long-term effects of exams week on HPA axis in undergraduates by the cortisol production.

METHODS

Subjects

Volunteers aged between 18 and 24 years were invited during class and poster for general inclusion criteria. They signed the free and informed consent, declaring understanding on the procedures that would be performed during the protocol. The study was conducted according to the guidelines laid down in the Declaration of Helsinki and ethical approval was granted from Research Ethics Committee of the School of Medical Sciences/University of Campinas (CAAE: 41225614.8.0000.5404). This study was performed with collaboration of biological sciences undergraduate students from University of Campinas (Unicamp). This undergraduate course was chosen because the full-time and evening courses have the same curriculum and practically all college subjects are taught in the same institute, avoiding the stress of displacement within the university. All samples were collected during the final exam/test week at the end of the school year.

Hair cortisol samples

The monthly production of cortisol was evaluated from each of the volunteers in the last 2 months. It was assumed that the average hair growth rate is 1 cm per month according to Dolnick *et al.* (Dolnick, Lindahl, Terrill, & Reynolds, 1969). Twenty-seven undergraduate students from biological sciences course in Unicamp provided hair samples, which were approximately 0.5 cm in diameter with a length between 3 and 5 cm near the scalp. The strands of hair were collected from the posterior vertex of the scalp and were cut off as close to the scalp as possible following Manenschijn *et al.* (Manenschijn, Koper, Lamberts, & van Rossum, 2011). The hair cortisol extraction was performed following Meyer et al. (Meyer, Novak, Hamel, & Rosenberg, 2014). The results are presented in nmol/L and are derived from 40 mg of hair samples sprayed and resuspended in 0.2 ml of assay buffer, and then the conversion factor in pg/mg of hair.

 Salivary cortisol samples

Participants were instructed to collect by themselves saliva samples at home, whose

Sallivetes® were to be stored under refrigeration. Twenty-eight undergraduate students collect

saliva to evaluate the cortisol rhythmicity. The cortisol rhythmicity was evaluated by saliva

collections at five times of the day during three consecutive days (Tuesday to Thursday), upon

awakening (6 a.m.), 30 minutes after awakening (6:30 a.m.), before lunch (12 a.m.), before dinner

(6 p.m.) and before bedtime (11 p.m.).

Cortisol quantification

The samples were centrifuged at 2,800 rpm, 20 min, 4 °C and analyzed using the

commercial ELISA kit (Diagnostics Biochem Canada Inc. - Ref CAN-C-290) at the Laboratory

of Stress Studies (LABEEST) - UNICAMP, as conducted by Madeira et al. (Madeira et al., 2019).

The results were expressed in nmol/L for each sample.

Statistical analysis

Data are presented as means ± SEM. The normality was confirmed by D'Agostino &

Pearson and then we performed paired or unpaired Student's t-test for parametric and Wilcoxon

or Mann-Whitney for nonparametric data. All statistical analysis was done with Graph Pad Prism

version 8.00 for Windows (Graph Pad Software, San Diego, California, USA). The acceptance

level of significance was set at p<0.05.

RESULTS

Hair cortisol and psychosocial stress

To verify the chronic effect of the test/exams week in cortisol release, we evaluated the

hair segments in months with two different expectations: one month without tests/exams

(October) and one month of tests/exams with possible emergence of psychosocial stress

(November). The results showed that the hair cortisol in November was significantly higher than

that observed in October (Figure 1).

Salivary cortisol

These volunteers also presented rhythmicity in cortisol release in the consecutive days, with higher values in the morning and a significantly decrease throughout the day. Even presenting adequate rhythmicity after the test marathon, we can observe an increase in the before bedtime sample at 2nd and 3th day when compared to the before dinner sample, since only 1st day exhibited significant difference from the same point. There was no significant change in the amount and rhythmicity of cortisol produced on different days (Figure 2). The cortisol awakening response (CAR) in these volunteers was preserved, with significantly higher values 30 min upon awakening in all days. CAR values increased throughout the days, but were not significant (Figure 3).

DISCUSSION

The hair cortisol is relevant to provide a retrospective reflection, for periods of several months, of cortisol release. In the last years, a number of evidences have been accumulated on several key features of hair cortisol measurements, including their validity as an indicator of long-term systemic cortisol concentrations, their reliability through repeated assessments, and their robustness over a series of influences. Based on these findings, researchers started to use this measurement to answer more specific questions about the role of stress in long-term cortisol production and its effect on different health-related conditions. The possibility to measure the long-term production of other steroid hormones (i.e. androgens or estrogens) in hair analysis provides a more intriguing prospect for future research. Taking in accounts its unique characteristics, the hair analysis is a great promise to significantly enhance current understanding about the role of steroid hormones (Braig et al., 2016; Serwinski, Salavecz, Kirschbaum, & Steptoe, 2016; Stalder et al., 2012; Stalder et al., 2017). The results obtained in this study are higher than those described in the literature for a healthy population, according to Parrot and Downey *et al.*. A healthy population presents an average of 13.8 pg/mg and our volunteers presents values of 58.32 and 109.11 for the months of October and November, respectively (Ouellet-Morin et al., 2016; Parrott & Downey, 2017; Short et al., 2016). However, hair cortisol

is known to be influenced by a number of factor that contribute to the variability of the hair-

cortisol results, as hair bleaching and dye treatment, hair texture and composition, exposure to

sun and hair washing (Mezentsef et al., 2017; Russell, Koren, Rieder, & Van Uum, 2012).

The daily variation of stressors, such as differences in workload levels between week days

and weekend days, can mask long-term changes in cortisol rhythms (Adam, Hawkley, Kudielka,

& Cacioppo, 2006; Karlamangla, Friedman, Seeman, Stawksi, & Almeida, 2013). The literature

already reported associations between cortisol parameters and psychosocial outcomes in studies

with 3-day sampling protocols (Hulett, Fessele, Clayton, & Eaton, 2019). Increased evening

cortisol over time may be linked to fatigue, as suggested by other studies (Schmidt et al., 2016;

Tell, Mathews, & Janusek, 2014; Weinrib et al., 2010).

The HPA axis can be assessed by quantifying salivary cortisol 30 min after awakening

(Fries, Dettenborn, & Kirschbaum, 2009; Pruessner et al., 1997; Wang et al., 2014). CAR

mechanism is independent of cortisol secretion throughout the day, being influenced by factors

not yet well clarified. However, that abnormalities in the rhythm of cortisol secretion, either in

CAR or in the rhythmicity, are associated with numerous diseases, as well as psychosocial

variables and chronic stress (Clow, Thorn, Evans, & Hucklebridge, 2004). In this way, cortisol

values 30 min after awakening significantly higher than those upon awakening indicate that the

HPA axis is intact. The cortisol produced in CAR also agrees with that described in the literature

for a healthy population and reflects a 38% increase in cortisol values upon awakening, being

considered healthy up to 100% of this baseline (Dedovic & Ngiam, 2015; Vreeburg et al., 2010).

Therefore, we demonstrated that the studied population, even living in a moment of psychosocial

stress, exhibited the rhythmicity of daily cortisol production as the CAR preserved and intact.

Regarding the evaluation of salivary cortisol, we can affirm that the studied population is healthy,

since it presented rhythmicity maintained during the week. However, the amounts of cortisol

produced in this undergraduate population are similar to the Canadian elderly population and

lower than the healthy third-year students at a US medical school during successive clinical

rotations (Souza-Talarico, Plusquellec, Lupien, Fiocco, & Suchecki, 2014; Tseng, Iosif, &

Seritan, 2011).

Biological instruments and techniques for the evaluation of chronic stress and the relation

with deleterious effects in the organism are fundamental. The possibility of biological

quantification of the effects of psychosocial stress in undergraduate students is necessary since

the incidence of mental illness in undergraduate students is high (Campos, Oliveira, Mello, &

Dantas, 2017). This pilot study presents limitations related to the reduced number of volunteers

who participated in the study because of adherence throughout protocol and the data were

obtained from a convenience sample offering reduced generalizability of our results.

The exams week is a reality in universities and can become a deleterious factor to the

undergraduate health since they maintain high cortisol levels throughout this period. The period

is a trigger for psychosocial stress, however, did not generate significant short-term changes in

cortisol production rhythmicity. Thus, it is evident that young people adapt to the stressful

stimulus, avoiding a possible trigger for mental illness. Beyond that, investigations with this same

population and the association or not with the previous experience of allostatic overload,

evaluated by cortisol in the hair, highlights information for possible preventive actions by mental

health strategy in the University.

ACKNOWLEDGEMENT

The authors wish to thank all the volunteers. The study was funded by Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES - Finance Code 001), Serviço de

Apoio ao Estudante da Unicamp (SAE-Unicamp), Fundo de Apoio ao Ensino, Pesquisa e

Extensão (Faepex-PRP), Conselho Nacional de Desenvolvimento Científico e Tecnológico

(CNPq) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). Part of these

results were presented in the XXIV Congresso de Iniciação Científica da Unicamp (PIBIC) –

2016 and in the 17ª Congresso de Stress do ISMA-BR – 2017, as a research reported poster.

AUTHORS' CONTRIBUTIONS

All authors had full access to all the data in the study and take responsibility for the

integrity of the data and the accuracy of the data analysis. Conceptualization, D.M.G.K. and

A.B.A.; Methodology, F.B., F.C. and D.M.G.K.; Investigation, F.B., P.C.S., F.C. and A.L.S.;

Formal Analysis, F.B. and D.M.G.K.; Resources: D.M.G.K; Writing – Original Draft, F.B., F.C.

P.C.S. and D.M.G.K.; Writing – Review & Editing, F.B. and D.M.G.K.; Visualization, F.B. and

D.M.G.K.; Supervision, F.B and D.M.G.K.; Funding Acquisition, D.M.G.K.

REFERENCES

Adam, E. K., Hawkley, L. C., Kudielka, B. M., & Cacioppo, J. T. (2006). Day-to-day Dynamics

of Experience--Cortisol Associations in a Population-Based Sample of Older Adults.

Proceedings of the National Academy of Sciences of the United States of America,

103(45). doi:10.1073/pnas.0605053103

Braig, S., Grabher, F., Ntomchukwu, C., Reister, F., Stalder, T., Kirschbaum, C., . . . Genuneit, J.

(2016). The Association of Hair Cortisol with Self-Reported Chronic Psychosocial Stress

and Symptoms of Anxiety and Depression in Women Shortly after Delivery. *Paediatr*

Perinat Epidemiol, 30(2), 97-104. doi:10.1111/ppe.12255

Campos, C. R., Oliveira, M. L., Mello, T. M., & Dantas, C. R. (2017). Academic performance of

students who underwent psychiatric treatment at the students' mental health service of a

Brazilian university. *Sao Paulo Med J, 135*(1), 23-28. doi:10.1590/1516-

3180.2016.017210092016

Clow, A., Thorn, L., Evans, P., & Hucklebridge, F. (2004). The awakening cortisol response:

methodological issues and significance. *Stress, 7*(1), 29-37.

doi:10.1080/10253890410001667205

D'Anna-Hernandez, K. L., Ross, R. G., Natvig, C. L., & Laudenslager, M. L. (2011). Hair cortisol

levels as a retrospective marker of hypothalamic-pituitary axis activity throughout

pregnancy: comparison to salivary cortisol. *Physiol Behav, 104*(2), 348-353.

doi:10.1016/j.physbeh.2011.02.041

Dedovic, K., & Ngiam, J. (2015). The cortisol awakening response and major depression:

examining the evidence. In *Neuropsychiatr Dis Treat* (Vol. 11, pp. 1181-1189).

Dolnick, E. H., Lindahl, I. L., Terrill, C. E., & Reynolds, P. J. (1969). Cyclophosphamide as a

227   chemical "defleecing" agent for sheep. *Nature, 221*(5179), 467-468.

228   Dwyer, A. L., & Cummings, A. L. (2001). Stress, Self-Efficacy, Social Support, and Coping

229   Strategies in University Students. *Canadian Journal of Counselling, 35*(3), 208-220.

230   Elias, H., Ping, W. S., & Abdullah, M. C. (2011). Stress and Academic Achievement among

231   Undergraduate Students in Universiti Putra Malaysia. *Procedia - Social and Behavioral*

232   *Sciences, 29*, 646-655. doi:10.1016/j.sbspro.2011.11.288

233   Fries, E., Dettenborn, L., & Kirschbaum, C. (2009). The cortisol awakening response (CAR):

234   facts and future directions. *Int J Psychophysiol, 72*(1), 67-73.

235   doi:10.1016/j.ijpsycho.2008.03.014

236   Gallagher, C. T., Mehta, A. N. V., Selvan, R., Mirza, I. B., Radia, P., Bharadia, N. S., & Hitch,

237   G. (2014). Perceived stress levels among undergraduate pharmacy students in the UK.

238   *Currents in Pharmacy Teaching and Learning, 6*(3), 437-441.

239   doi:10.1016/j.cptl.2014.02.004

240   García-León, M. Á., Pérez-Mármol, J. M., Gonzalez-Pérez, R., García-Ríos, M. D. C., & Peralta-

241   Ramírez, M. I. (2019). Relationship Between Resilience and Stress: Perceived Stress,

242   Stressful Life Events, HPA Axis Response During a Stressful Task and Hair Cortisol.

243   *Physiology & behavior, 202*. doi:10.1016/j.physbeh.2019.02.001

244   Hulett, J. M., Fessele, K. L., Clayton, M. F., & Eaton, L. H. (2019). Rigor and Reproducibility:

245   A Systematic Review of Salivary Cortisol Sampling and Reporting Parameters Used in

246   Cancer Survivorship Research. *Biological research for nursing, 21*(3).

247   doi:10.1177/1099800419835321

248   Karatsoreos, I. N., & McEwen, B. S. (2014). Timing is everything: a collection on how clocks

249   affect resilience in biological systems. *F1000Res, 3*, 273.

250   doi:10.12688/f1000research.5756.1

251   Karlamangla, A. S., Friedman, E. M., Seeman, T. E., Stawksi, R. S., & Almeida, D. M. (2013).

252   Daytime Trajectories of Cortisol: Demographic and Socioeconomic Differences.

Findings from The National Study of Daily Experiences. *Psychoneuroendocrinology,*

38(11). doi:10.1016/j.psyneuen.2013.06.010

Liston, C., McEwen, B. S., & Casey, B. J. (2009). Psychosocial stress reversibly disrupts

prefrontal processing and attentional control. *Proc Natl Acad Sci U S A, 106*(3), 912-917.

doi:10.1073/pnas.0807041106

Madeira, L. P., Borghi, F., Wilson, V. D., Souza, A. L., Pires-da-Rocha, M. C., Ferreira, H. A., .

. . Schwartz, G. M. (2019). Perceived stress and salivary cortisol on rock climbing.

Translational Sports Medicine, 2, 370-375. doi:10.1002/tsm2.104

Manenschijn, L., Koper, J. W., Lamberts, S. W., & van Rossum, E. F. (2011). Evaluation of a

method to measure long term cortisol levels. *Steroids, 76*(10-11), 1032-1036.

doi:10.1016/j.steroids.2011.04.005

Meyer, J., Novak, M., Hamel, A., & Rosenberg, K. (2014). Extraction and analysis of cortisol

from human and monkey hair. *J Vis Exp*(83), e50882. doi:10.3791/50882

Mezentsef, A., Jackson, P., Baig, U., Fairclough, J. L., Brook, C., & Mitroka, J. (2017). Hair

Cortisol and Perceived Stress in Health - studies Students during Summer Break and Fall

Term. *J Steroids Horm Sci, 8*(1). doi:10.4172/2157-7536.1000183

Moore, D. A., Truscott, M. L., St Clair, L., & Klingborg, D. J. (2007). Effects of a Veterinary

Student Leadership Program on Measures of Stress and Academic Performance. *Journal*

of veterinary medical education, 34(2). doi:10.3138/jvme.34.2.112

Ouellet-Morin, I., Laurin, M., Robitaille, M. P., Brendgen, M., Lupien, S. J., Boivin, M., & Vitaro,

F. (2016). Validation of an adapted procedure to collect hair for cortisol determination in

adolescents. *Psychoneuroendocrinology, 70*, 58-62. doi:10.1016/j.psyneuen.2016.05.002

Parrott, A. C., & Downey, L. A. (2017). Psychoactive drug influences on hair cortisol.

Psychoneuroendocrinology. doi:10.1016/j.psyneuen.2017.02.034

Portoghese, I., Galletta, M., Porru, F., Burdorf, A., Sardo, S., D'Aloja, E., . . . Campagna, M.

(2019). Stress Among University Students: Factorial Structure and Measurement

Invariance of the Italian Version of the Effort-Reward Imbalance Student Questionnaire.

BMC psychology, 7(1). doi:10.1186/s40359-019-0343-7

Pruessner, J. C., Wolf, O. T., Hellhammer, D. H., Buske-Kirschbaum, A., von Auer, K., Jobst, S.,

. . . Kirschbaum, C. (1997). Free cortisol levels after awakening: a reliable biological

marker for the assessment of adrenocortical activity. *Life Sci, 61*(26), 2539-2549.

Russell, E., Koren, G., Rieder, M., & Van Uum, S. (2012). Hair cortisol as a biological marker of

chronic stress: current status, future directions and unanswered questions.

Psychoneuroendocrinology, 37(5), 589-601. doi:10.1016/j.psyneuen.2011.09.009

Sauvé, B., Koren, G., Walsh, G., Tokmakejian, S., & Van Uum, S. H. (2007). Measurement of

Cortisol in Human Hair as a Biomarker of Systemic Exposure. *Clinical and investigative*

medicine. Medecine clinique et experimentale, 30(5). doi:10.25011/cim.v30i5.2894

Schmidt, M. E., Semik, J., Habermann, N., Wiskemann, J., Ulrich, C. M., & Steindorf, K. (2016).

Cancer-related Fatigue Shows a Stable Association With Diurnal Cortisol Dysregulation

in Breast Cancer Patients. *Brain, behavior, and immunity, 52.*

doi:10.1016/j.bbi.2015.10.005

Serwinski, B., Salavecz, G., Kirschbaum, C., & Steptoe, A. (2016). Associations between hair

cortisol concentration, income, income dynamics and status incongruity in healthy

middle-aged women. *Psychoneuroendocrinology, 67*, 182-188.

doi:10.1016/j.psyneuen.2016.02.008

Short, S. J., Stalder, T., Marceau, K., Entringer, S., Moog, N. K., Shirtcliff, E. A., . . . Buss, C.

(2016). Correspondence between hair cortisol concentrations and 30-day integrated daily

salivary and weekly urinary cortisol measures. *Psychoneuroendocrinology, 71*, 12-18.

doi:10.1016/j.psyneuen.2016.05.007

Souza-Talarico, J. N., Plusquellec, P., Lupien, S. J., Fiocco, A., & Suchecki, D. (2014). Cross-

country differences in basal and stress-induced cortisol secretion in older adults. *PLoS*

One, 9(8), e105968. doi:10.1371/journal.pone.0105968

Stalder, T., Steudte, S., Miller, R., Skoluda, N., Dettenborn, L., & Kirschbaum, C. (2012).

Intraindividual stability of hair cortisol concentrations. *Psychoneuroendocrinology,*

37(5), 602-610. doi:10.1016/j.psyneuen.2011.08.007

Stalder, T., Steudte-Schmiedgen, S., Alexander, N., Klucken, T., Vater, A., Wichmann, S., . . .
309   Miller, R. (2017). Stress-related and basic determinants of hair cortisol in humans: A
310   meta-analysis. *Psychoneuroendocrinology, 77,* 261-274.
311   doi:10.1016/j.psyneuen.2016.12.017

312 Tell, D., Mathews, H. L., & Janusek, L. W. (2014). Day-to-day Dynamics of Associations
313   Between Sleep, Napping, Fatigue, and the Cortisol Diurnal Rhythm in Women Diagnosed
314   as Having Breast Cancer. *Psychosomatic medicine, 76*(7).
315   doi:10.1097/PSY.0000000000000097

316 Tseng, T., Iosif, A.-M., & Seritan, A. L. (2011). Stress Effects: A Study of Salivary Cortisol
317   Levels in Third-year Medical Students. *Stress and Health, 27*(5), 436-440.
318   doi:10.1002/smi.1377

319 Vreeburg, S. A., Hartman, C. A., Hoogendijk, W. J., van Dyck, R., Zitman, F. G., Ormel, J., &
320   Penninx, B. W. (2010). Parental history of depression or anxiety and the cortisol
321   awakening response. *Br J Psychiatry, 197*(3), 180-185. doi:10.1192/bjp.bp.109.076869

322 Wang, X., Sánchez, B. N., Golden, S. H., Shrager, S., Kirschbaum, C., Karlamangla, A. S., . . .
323   Roux, A. V. (2014). Stability and predictors of change in salivary cortisol measures over
324   six years: MESA. *Psychoneuroendocrinology, 49,* 310-320.
325   doi:10.1016/j.psyneuen.2014.07.024

326 Weinrib, A. Z., Sephton, S. E., Degeest, K., Penedo, F., Bender, D., Zimmerman, B., . . .
327   Lutgendorf, S. K. (2010). Diurnal Cortisol Dysregulation, Functional Disability, and
328   Depression in Women With Ovarian Cancer. *Cancer, 116*(18). doi:10.1002/cncr.25299

329 Wester, V. L., & van Rossum, E. F. (2015). Clinical Applications of Cortisol Measurements in
330   Hair. *European journal of endocrinology, 173*(4). doi:10.1530/EJE-15-0313

    FIGURES

Figure 1 – Hair cortisol from undergraduate students in October (month without exams) and November (month of exams). Data are presented as mean ± SEM. *p<0.05 in the same group.

Figure 2 – Cortisol rhythmicity by saliva samples at five different times of the day during three consecutive days: upon awakening (6 a.m.), 30 minutes after awakening (6:30 a.m.), before lunch (12 a.m.), before dinner (6 p.m.) and before bedtime (11 p.m.). Data are presented as mean ± SEM. *p<0.05 vs. 6 a.m. in the same group; #p<0.05 vs. 11 p.m. in the same group; @p<0.05 1st day 6 p.m. vs. 1st day 11 p.m.

346 **Figure 3 -** CAR from undergraduate students in different consecutive days. Data are presented as

347 mean ± SEM. *p<0.05 in the same group.