Comparative genomics analysis of *Chryseobacterium* sp. KMC2 reveals metabolic

pathways involved in keratinous utilization and natural product biosynthesis

Dingrong Kang[abcd*], Saeed Shoaie[c], Samuel Jacquiod[e], Søren J. Sørensen[a], Rodrigo Ledesma-

Amaro[b,*]

[a]Section of Microbiology, Department of Biology, University of Copenhagen, Copenhagen,

Denmark

[b]Department of Bioengineering and Imperial College Centre for Synthetic Biology, Imperial

College London, London, SW7 2AZ, UK

[c]Centre for Host-Microbiome Interactions, Faculty of Dentistry, Oral & Craniofacial Sciences,

King's College London, SE1 9RT, UK

[d]TERRA Research and Teaching Centre, Microbial Processes and Interactions (MiPI),

Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium

[e]Agroécologie, AgroSup Dijon, INRAE, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, F-

21000 Dijon, France

[*]Corresponding authors: Rodrigo Ledesma-Amaro (r.ledesma-amaro@imperial.ac.uk) and

Dingrong Kang (dingrong.kang@hotmail.com)

Abstract

Several efforts have been made to valorize keratinous materials, an abundant and renewable resource. Despite these attempts to valorize products generated from keratin hydrolysate, either *via* chemical or microbial conversion, they generally remain with an overall low value. In this study, a promising keratinolytic strain from the genus *Chryseobacterium* (*Chryseobacterium* sp. KMC2) was investigated using comparative genomic tools against publicly available reference genomes to reveal the metabolic potential for biosynthesis of valuable secondary metabolites. Genome and metabolic features of four species were compared, shows different gene numbers but similar functional categories. We successfully mined eleven different secondary metabolite gene clusters of interest from the four genomes, including five common ones shared across all genomes. Among the common metabolites, we identified gene clusters involved in biosynthesis of flexirubin-type pigment, microviridin, and siderophore, all showing remarkable conservation across the four genomes. Unique secondary metabolite gene clusters were also discovered, for example, ladderane from *Chryseobacterium* sp. KMC2. Additionally, this study provides a more comprehensive understanding of the potential metabolic pathways of keratin utilization in *Chryseobacterium* sp. KMC2, with the involvement of amino acid metabolism, TCA cycle, glycolysis/gluconeogenesis, propanoate metabolism, and sulfate reduction. This work uncovers the biosynthesis of secondary metabolite gene clusters from four keratinolytic *Chryseobacterium* spp. and shades lights on the keratinolytic potential of *Chryseobacterium* sp. KMC2 from a genome-mining perspective, providing alternatives to valorize keratinous materials into high-value natural products.

Importance

Keratin is an abundant and renewable resource from slaughterhouses or the poultry industry.

Low-value products such as animal feed and fertilizer were generated from these feedstocks

based on conventional processing like chemical conversion. In fact, microorganisms possess the

potential to synthesize valuable natural products. In this work, we explored the metabolic

potential of *Chryseobacterium* sp. KMC2, which was isolated with efficient keratinolytic

capacity from a previous study. Comparative genomics analysis displayed similar functional

categories against three publicly available reference genomes of keratin-

degrading *Chryseobacterium* spp.. Eleven different secondary metabolite gene clusters of interest

were mined among four genomes, including five common and unique ones. Furthermore, we

provide a more comprehensive understanding of metabolic pathways on keratin utilization

in *Chryseobacterium* sp. KMC2, with the involvement of amino acid assimilation and sulfate

reduction. These findings contribute to expanding the application of *Chryseobacterium* sp.

KMC2 on the valorization of keratinous materials.

Keywords: keratinous materials, metabolic potential, genome mining, gene clusters,

degradation pathways

Introduction

Keratin is the most abundant proteins in epithelial cells, constituting the bulk of epidermal appendages such as hair and feather (1, 2). Keratinous materials represent an abundant protein source, particularly originating from the commercial slaughterhouses or poultry farms (3). They contain peptides and amino acids, which are renewable natural resources with great potential in sustainable development (4). However, keratin is an insoluble protein with highly cross-linked disulfide bonds giving it a tough and recalcitrant structure (5). Many attempts have been made to hydrolysis keratinous materials in terms of physicochemical treatment, enzymatic hydrolysis, and microbial conversion (6, 7). The hydrolysis products of keratinous materials have been used for animal feed (8) and fertilizer (6, 9) based on conventional processing.

Microorganisms represent one of the most important natural sources, which have the potential to generate bioactive compounds such as antibiotics, biofuels, and natural pigments derived from cellular metabolites (10, 11). For example, *Yarrowia lipolytica* has been used to convert different renewable feedstocks to high-value metabolites (12). Similarly, *Escherichia coli* has become one of the best cell reactors to produce alcohols, organic acids, biodiesel, even hydrogen by utilizing renewable resources (13). Other bacteria such as *Bacillus subtilis* (14), *Caldicellulosiruptor bescii* (15), *Corynebacterium glutamicum* (16), and *Ruminococcaceae* bacterium (17) were identified and evaluated with the capacity to generate different products by converting renewable carbon sources. Notably, some microorganisms were reported to degrade keratinous waste effectively (18). Exploring keratinolytic potential of these microbes to generate high value-added products is an important step to recycle and valorize keratinous materials.

Molecular mechanisms of microbial keratin degradation are still not fully understood, while

genome sequencing offers possibilities to reveal the metabolic potential behind efficient

microbial degradation (19). Novel keratinolytic enzymes were identified from the genome of

Bacillus pumilus 8A6, an efficient keratin degrader (20). Furthermore, going beyond the

degradation reaction itself, genomes can also be minded for valuable accessory functions of

interest, adding more values to the microbial conversion processes. For instance, gene clusters

and biosynthesis pathways of secondary metabolites could be disclosed from genomes *via*

adequate analysis tools (21, 22). A total of 104 putative biosynthetic gene clusters (BGCs) for

secondary metabolites were predicted from nine *Ktedonobacteria* genomes (23). Secondary

metabolites were identified and linked to gene clusters based on the comparison and mining of

six genomes belonging to diverse *Aspergillus* species, successfully fueling industrial

biotechnology initiatives and medical research (24). Therefore, using the genomes of

keratinolytic microbial species in a similar way would represent a promising approach to

discover biosynthetic gene clusters of secondary metabolites of interest, excavating the full

application potential of these microbes.

Recently, several studies based on different environments have revealed the remarkable potential

of representative taxa from the *Chryseobacterium* genus for keratin degradation using isolation,

activity tests and genome sequencing (25, 26). In this study, a novel strain *Chryseobacterium* sp.

KMC2 was previously obtained from an enrichment procedure, displaying a potent capacity of

keratin degradation (27). The genome of *Chryseobacterium* sp. KMC2 was analyzed and

compared with publicly available genomes of other keratinolytic *Chryseobacterium* spp. to

clarify the genomic basis of keratin degradation, and to unravel hidden biosynthetic gene clusters

of interest. Subsequently, the metabolic pathways associated with keratin degradation were

constructed, providing deeper insight into the yet obscure keratinolytic processes. This work

reveals the keratinolytic potential of *Chryseobacterium* spp. and mined potential accessory gene

clusters of secondary metabolites, which could i) contribute to optimizing the processes of

keratin degradation and ii) broaden the perspective to generate added-value products from

keratin hydrolysate.

Results and discussion

Genome feature comparison of four keratinolytic *Chryseobacterium* strains

Chryseobacterium sp. KMC2 originated from a river-bank soil sample, and displayed a potent

degradation ability toward milled pig bristle and hooves (27, 28). The genome of

Chryseobacterium sp. KMC2 was sequenced and compared to three reference genomes of

Chryseobacterium spp. Including: i) *Chryseobacterium camelliae* Dolsongi-HT, isolated from

green tea leaves (29); ii) *Chryseobacterium gallinarum* strain DSM 27622, isolated from chicken

(30); and iii) *Chryseobacterium* sp. strain P1-3 isolated from poultry waste (31), which all

display keratinolytic capacity (Table 1). *Chryseobacterium* sp. KMC2 showed distinct genome

feature from the other known keratinolytic strains. The genome size of *Chryseobacterium* sp.

KMC2 is 5.28 Mbp, larger than the other three genomes, which ranged from 4.38 Mbp to 4.63

Mbp. A total of 4,773 genes were predicted from *Chryseobacterium* sp. KMC2 genome, and

about 4,000 genes were annotated from the other three genomes. Besides, the GC content ranges

from 36.33% to 41.80% in *Chryseobacterium* spp. genomes. Furthermore, the whole-genome

phylogenetic tree was constructed with other eight publically available *Chryseobacterium* spp.

genomes (Fig. 1a), showing the highly similarity among *Chryseobacterium gallinarum* strain

DSM 27622 and *Chryseobacterium* sp. strain P1-3. Notably, *Chryseobacterium* sp. KMC2 and

Chryseobacterium camelliae Dolsongi-HT have closer evolutionary relationship with other

Chryseobacterium species. Although the keratinolytic capacity within these eight strains is still

not clear, more keratinolytic species are expected to be found in this genus according to the

phylogenic tree analysis. Four strains among this genus have documented with actual keratin

degradation capability (29-31), while *Chryseobacterium* sp. KMC2 may possess different

keratinolytic potential encoded in its larger genome compared to three other *Chryseobacterium*

strains.

Metabolic potential comparison of four keratinolytic *Chryseobacterium* genomes

About 40% of the genes from the four genomes were classified into various functional categories

based on the KEGG database. The vast majority of annotated genes belonged to metabolism,

genetic information processing, environmental information processing, and cellular processes

(Fig. 1b). The functional categories of the genomes were overall highly similar, with ~85% of

annotated genes assigned to "metabolism" (category A) which included ~1.000 genes into the

sub-category "global and overview maps". Additionally, about 8% and 4% annotated genes from

each genome were assigned to "genetic information processing" (category B) and

"environmental information processing" (category C), respectively. The remaining annotated

genes belonged to "cellular processes" (category D), which occupied 3% of the annotated

genomes approximately.

Remarkably, each genome had more than 200 genes assigned into the "amino acid metabolism"

sub-category. Keratin is mainly composed of amino acids (3), which is ultimately the operational

carbon nutrient source exploited for microbial growth. Numerous amino acid metabolism-related

enzymes were annotated, revealing the genetic potential of these *Chryseobacterium* strains for

using keratin materials as energy sources. Of particular interest, several biosynthesis genes of

secondary metabolites were detected from the genomes, of which more than 20 genes were

annotated as "metabolism of terpenoids and polyketides" and around 40 genes were annotated as

"biosynthesis of other secondary metabolites" sub-category (Fig. 1b). Terpenoids are a group of

natural products with diverse commercial applications, which have been produced from

microbial cell factories (32). Many polyketides are considered as significant natural products

with broad applications in the agriculture and pharmaceutical industry (33). The metabolic

pathways related to polyketides biosynthesis production are well understood in some

microorganisms like *Streptomyces* which play a crucial role in industrial bioproduction (34).

This result indicates that these *Chryseobacterium* strains could have the potential to synthesize

high-value secondary metabolites such as terpenoids and polyketides from keratinous materials.

Mining and comparing secondary metabolite gene clusters

Genome mining is an effective approach to discover new natural products from microorganisms

based on "signature genes" detection or searching for specific patterns in gene sequences (35).

To explore the potential of producing high value chemicals from these four *Chryseobacterium*,

secondary metabolite gene clusters were predicted by using antiSMASH 5.0 mining pipeline (Fig.

2). In total, eleven different secondary metabolite gene clusters were identified.

Chryseobacterium sp. KMC2 possesses the largest number (15), while *Chryseobacterium*

camelliae Dolsongi-HT1 has the fewest (8). Ten gene clusters were predicted from the other two

strains. Five clusters are present in the four genomes, which are flexirubin-type pigment

(resorcinol and arylpolyene), microviridin, lanthipeptide, NRPS-like, and siderophore.

Remarkably, the flexirubin-type pigment is a typical metabolite produced from *Flavobacterium*

(36). Several species from *Chryseobacterium* were previously designated and known

as *Flavobacterium* owes to similar characteristics including the presence of yellow pigments (37).

Flexirubin-type pigment was isolated and characterized from *Chryseobacterium* sp. UTM-3T

(38). In addition, *Chryseobacterium* sp. KMC2 owns a unique gene cluster to produce ladderane.

Another unique natural product is beta-lactone from *Chryseobacterium camelliae* Dolsongi-HT1.

Ladderanes are hydrocarbon chains which were regarded as membrane lipid components

produced by anammox (anaerobic ammonia-oxidizing) bacteria uniquely, but the production is

not affordable due to their extremely low growth (39, 40). These results demonstrate that various

secondary metabolite gene clusters including both expected and unusual candidates were

discovered from *Chryseobacterium* genomes, which could turn into novel natural product

sources.

Synteny analysis and features of secondary metabolite gene clusters

Comparative genomics can reveal unique cluster and distribution patterns of secondary

metabolites in species (41). Five secondary metabolite gene clusters, predicted to be present in

the four genomes, were selected to explore the evolutionary relationship among four

Chryseobacterium strains. Three of them including flexirubin-type pigment, microviridin, and

siderophore display a conserved gene cluster structure from synteny analysis (Fig. 3, 4, and 5),

while the other two showed no evident synteny relation (Fig. S1 and S2).

Flexirubin-type pigment. Natural pigments have increasing applications in food,

pharmaceutical, and textile industries, owing to their advantages such as non-toxic,

biodegradable, and low allergenic potential compared to synthetic pigments (42). In particular,

flexirubin-type pigment has a potential antimicrobial and anti-tumoral activities (43).

Biosynthesis gene clusters of flexirubin-type pigment are conserved across the four tested

genomes, especially within *Chryseobacterium gallinarum* strain DSM 27622 and

Chryseobacterium sp. strain P1-3 (Fig. 3a). A total of 61 biosynthesis-related genes of

flexirubin-type pigment were predicted from *Chryseobacterium* sp. KMC2, including four core

biosynthesis genes. One of the core biosynthesis genes was annotated as 3-oxoacyl-(acyl carrier

protein) synthase III (*Flex11*), and the other three were annotated as Beta-ketoacyl synthases

(*Flex21*, *Flex24*, and *Flex40*). Besides, transport-related genes and regulatory genes were

predicted from the gene cluster. A previous study identified the molecular structure of flexirubin-

type pigment isolated from a *Chryseobacterium* sp. UTM-3T (38). According to the products

from core biosynthesis genes and their molecular structures, a proposition of biosynthesis

pathway was established (Fig. 3b), where flexirubin-type pigment is generated from resorcinol

and arylpolyene. Further transcriptomics and metabolomics analysis would be required to

confirm the validity of this potential pathway discovery.

Microviridin. Microviridins represent a group of ribosomally synthesized peptides under post-

translational modifications, which have been mainly isolated from cyanobacteria and present

potent serine-type protease inhibitory activities (44, 45). These properties could make

microviridin serve as the natural antimicrobial agents for developing potential drugs.

Biosynthesis gene clusters of microviridin from four *Chryseobacterium* genomes show a highly

conserved structure with a similarity greater than 71% from most gene synteny analysis (Fig. 4a).

There are 23 biosynthesis genes of microviridin that were predicted from *Chryseobacterium* sp.

KMC2. Two core biosynthetic genes (A and B) were identified from genomes and transport-

related genes were also been discovered. Besides, amino acid sequences of mvdA and mvdB

were aligned, showing that multiple motifs from mvdA and mvdB are conserved (Fig. 4bc).

Interestingly, many keratinases were reported to be classified as serine proteases, acting on the

molecular structure of keratin (46). This suggests that microviridins may regulate keratinolytic

activity, further characterizing and manipulating the microviridin synthetic pathway could

contribute to improving the keratin degradation efficiency.

Siderophore. Siderophores are ferric ion-specific chelators to scavenge iron from the

extracellular environment, which play important roles in virulence and oxidative stress tolerance

in microorganisms (47). It has been designed as a Trojan horse antibiotic to enter and kill

pathogenic bacteria (48), and also has shown the potential to decrease the growth of cancerous

cells (49). Biosynthesis gene cluster of siderophore shows a high synteny conservation among

Chryseobacterium sp. KMC2 and *Chryseobacterium camelliae* Dolsongi-HT1,

Chryseobacterium gallinarum strain DSM 27622 and *Chryseobacterium* sp. Strain P1-3,

respectively (Fig. 5). A total of ten genes were predicted from siderophore biosynthesis cluster of

Chryseobacterium sp. KMC2, and eight genes from the other three *Chryseobacterium* strains

separately. Functional description of each gene related to siderophore biosynthesis in

Chryseobacterium sp. KMC2 shows two core biosynthesis genes, and includes one regulatory

gene and one transport-related gene. This further suggests that those siderophore are potentially

fully functional molecular features that can be regulated on-demand and exported outside the cell

when needed.

Metabolic pathways of keratin utilization in *Chryseobacterium* sp. KMC2 genome

The main metabolic pathways related to keratin utilization in *Chryseobacterium* sp. KMC2

genome were investigated. These pathways included amino acid metabolism, TCA cycle,

glycolysis/gluconeogenesis, propanoate metabolism, and sulfate reduction (Fig. 6). A previous

study suggested that abundant amino acids are released during microbial degradation and used as

nutrient sources, such as leucine and aspartate (28). The metabolic pathways of amino acid

utilization were mapped from the genome of *Chryseobacterium* sp. KMC2. Most of the amino

acids are converted into intermediates of the TCA cycle. For instance, arginine can be converted

to succinate, then enter to TCA cycle after a multiple-steps enzyme reaction. Aspartate, tyrosine,

phenylalanine, and glutamate could serve as the substrates to generate fumarate, thus being part

of the TCA cycle. Besides, isoleucine turns into the substrates of 2-methyl-acetoacetyl-CoA after

several enzymatic steps, which is then converted into acetyl-CoA and propanoyl-CoA *via* acetyl-

CoA C-acyltransferase. Acetyl-CoA is an important intermediate, which can entry to the TCA

cycle *via* citrate synthase (50). It is also the precursors of fatty acid and polyketides biosynthesis

(51). Propanoyl-CoA serves as the critical substrate within propanoate metabolism and can also

be used to make lipids (52, 53). On the other hand, methionine can be converted to 2-

oxobutanoate, which is also an intermediate of propanoate metabolism. Subsequently, the

methylmalonyl-CoA generated in propanoate metabolism enters into the TCA cycle *via* succinyl-

CoA. Besides, the key enzymes of glycolysis/gluconeogenesis were found, indicating the

potential to produce essential biomass components based on oxaloacetate from TCA cycle.

Evidence indicates that a source of redox is needed for complete keratin degradation with

keratinases (54, 55). Several metabolites such as sulfite were revealed to be associated with

efficient keratin degradation (56). Therefore, here the complete metabolic pathway of sulfate

reduction was mapped in the genome, which shows that the potential to create a redox

environment needed for keratinases is indeed present. Following the development of sequencing

technologies, increasing genomes of keratinolytic species have been unveiled, which provide a

genomic perspective to reveal the molecular keratinolytic mechanisms. For instance, metabolic

pathways related to keratin degradation such as enzymolysis and reduction of disulfide bonds

were clarified through uncovering the genetic basis of microbial genomes (57). The complex

keratinolytic processes of *Streptomyces* sp. included protease secretion, iron uptake, spore

formation, and resuscitation were recently revealed from a genome view (58). Our results are in

line with the notion that a redox environment is indeed required for efficient keratinolytic

activity to occur. It is expected that the integrated metabolic pathways associated with

keratinolytic processes will be deciphered along with more genomes sequencing and biochemical

studies of relevant metabolic pathways.

Conclusion

In this work, the genomes from four *Chryseobacterium* spp. with keratinolytic activity were

analyzed. Common and unique secondary metabolite gene clusters were mined from

Chryseobacterium spp. genomes, suggesting the potential to generate high value metabolites

using keratin-rich wastes as the nutrient sources. Therefore, the use of these microorganisms

could be an alternative way to valorize keratinous materials through microbial conversion.

Furthermore, the metabolic pathways of keratin degradation from *Chryseobacterium* sp. KMC2

was studied from a genomic viewpoint. Nevertheless, there are still unknowns to link both

metabolic pathways of keratinous utilization and the natural products biosynthesis.

Understanding these connected pathways and their regulation will contribute to developing

synthetic biology approaches to boost high value-added products from microbial keratin

degradation.

Materials and Methods

DNA preparation

Chryseobacterium sp. KMC2 was isolated and identified from a keratinolytic microbial

consortium enriched from a soil sample (19, 27, 28). The keratinolytic capacity of

Chryseobacterium sp. KMC2 was evaluated as the keystone strain among the microbial

consortium, which was able to degrade keratin materials efficiently (27). *Chryseobacterium* sp.

KMC2 was inoculated to LB medium, and cultured overnight (200 rpm, 30 °C). Two milliliters

of the suspension were centrifuged and collected to prepare the DNA extraction, performed by

using by FAST Soil DNA Kit (MP Biomedicals, United States) according to the manufacturer's

instructions.

Genome sequencing, assembling, and functional annotation

The genome sequencing was performed by an Illumina Miseq instrument at the University of

Copenhagen by using DNA Library Preparation Kits v2 (2×250 bp), according to the

manufacturer's instructions. Raw reads were treated and assembled to contigs on CLC Genomic

Workbench 8.5.1. The obtained contigs were validated using QUAST 4.5 (59). Genes were

predicted from the contigs and further annotated with Prokka v1.14.5 (60). Predicted genes from

contigs were submitted to eggNOG 5.0 database to obtain an integrated functional annotation

and classification (61).

Whole-genome phylogenetic analysis

To determine the phylogenetic origin of *Chryseobacterium* sp. KMC2 in the *Chryseobacterium*

genus, the whole-genome sequences of 11 publicly available *Chryseobacterium* spp. were

downloaded from NCBI database to construct a phylogenetic tree. The whole-genome sequence-

based phylogenetic tree was inferred by using an online pipeline: The Reference sequence

Alignment based Phylogeny builder (REALPHY 1.12) (62), based on the merge reference

alignments. Then the visualization of the phylogenetic tree was generated by iTOL v5 (63).

Secondary metabolite gene cluster detection and annotation

Assembled contigs of four *Chryseobacterium* spp. were uploaded to antiSMASH 5.0 secondary

metabolite genome mining web platform (21). Predicted secondary metabolites gene clusters

from *Chryseobacterium* sp. KMC2 were compared with other keratinolytic *Chryseobacterium*

strains. Gene annotation of each cluster from *Chryseobacterium* sp. KMC2 was performed by

Prokka v1.14.5 (60) and BLASTP with the NCBI database. The best match sequencing ID was

recorded for the annotated genes. Synteny and features of conservative secondary metabolite

gene clusters were analyzed by using Easyfig 2.2.2, showing the similarity of gene sequences

(64). Feature comparison of amino acid sequences and motifs from core synthetic genes were

analyzed by using Clustal Omega (65) to get the multiple sequence alignment and using

Seq2logo to generate sequence logo (66).

Metabolic networks construction

The genomes of *Chryseobacterium* sp. KMC2 and other three *Chryseobacterium* spp. were

submitted to GhostKOALA (67) to obtain the KO number for each gene, then genes were

assigned to different metabolic pathways and functional categories. Following the metabolic

networks construction of *Chryseobacterium* sp. KMC2 was achieved through mapping the

annotated enzyme genes to KEGG (68) reference pathway and Biocyc database (69) manually.

Data availability

Reference genomes were downloaded from the NCBI database: *Chryseobacterium camelliae*

strain Dolsongi-HT1 (GenBank: GCA_002770595.1), *Chryseobacterium gallinarum* strain DSM

27622 (GenBank: GCA_001021975.1), *Chryseobacterium* sp. P1-3 (GenBank:

GCA_000738495.1), *Chryseobacterium gleum* NCTC11432 (GenBank: GCA_900636535.1),

Chryseobacterium bernardetii H4638 (GenBank: GCA_003815955.1), *Chryseobacterium*

arthrosphaerae FDAAGOS 519 (GenBank: GCA_003812705.1), *Chryseobacterium indologenes*

FDAARGOS 337 (GenBank: GCA_002208925.2), *Chryseobacterium joostei* DSM 16927

(GenBank: GCA_003815775.1), *Chryseobacterium glaciei* IHBB 10212 (GenBank:

GCA_001648155.1), *Chryseobacterium carnipullorum* F9942 (GenBank: GCA_003815855.1),

and *Chryseobacterium* sp. SNU WT5 (GenBank: GCA_007362475.1). Raw sequencing data

were deposited in the Sequence Read Archive (SRA) database under the BioProject number

PRJNA686768 with an accession number SRR13278108. The assembled genome sequence

of *Chryseobacterium* sp. KMC2 has been deposited at DDBJ/ENA/GenBank under the accession

JAESIT000000000.

Acknowledgements

This research was funded by the Danish Innovation Fund (Grant Number 1308-00015B,

Keratin2Protein) and also under the support of the Chinese Scholarship Council (CSC)

Scholarship Program.

References

1.   Wang B, Yang W, McKittrick J, Meyers MA. 2016. Keratin: Structure, mechanical

properties, occurrence in biological organisms, and efforts at bioinspiration. Progress in

Materials Science 76:229-318.

2.   Coulombe PA, Omary MB. 2002. 'Hard'and 'soft'principles defining the structure,

function and regulation of keratin intermediate filaments. Current opinion in cell biology 14:110-

122.

3.   Korniłłowicz-Kowalska T, Bohacz J. 2011. Biodegradation of keratin waste: theory and

practical aspects. Waste management 31:1689-1701.

4.   Khosa M, Ullah A. 2013. A sustainable role of keratin biopolymer in green chemistry: a

review. J Food Processing & Beverages 1:8.

5.   Rajabi M, Ali A, McConnell M, Cabral J. 2020. Keratinous materials: Structures and

functions in biomedical applications. Materials Science and Engineering: C 110:110612.

6.   Holkar CR, Jain SS, Jadhav AJ, Pinjari DV. 2018. Valorization of keratin based waste.

Process Safety and Environmental Protection 115:85-98.

7.   Zhao W, Yang R, Zhang Y, Wu L. 2012. Sustainable and practical utilization of feather

keratin by an innovative physicochemical pretreatment: high density steam flash-explosion.

Green chemistry 14:3352-3360.

8.   Dios D. 2001. Fishmeal replacement with feather-enzymatic hydrolyzates co-extruded

with soya-bean meal in practical diets for the Pacific white shrimp (*Litopenaeus vannamei*).

Aquaculture Nutrition 7:143-151.

9.   Gaidau C, Epure D-G, Enascuta CE, Carsote C, Sendrea C, Proietti N, Chen W, Gu H.

2019. Wool keratin total solubilisation for recovery and reintegration-An ecological approach.

Journal of Cleaner Production 236:117586.

10.   Donadio S, Monciardini P, Alduina R, Mazza P, Chiocchini C, Cavaletti L, Sosio M,

Puglia AM. 2002. Microbial technologies for the discovery of novel bioactive metabolites.

Journal of Biotechnology 99:187-198.

11.   Smitha M, Singh S, Singh R. 2017. Microbial biotransformation: a process for chemical

alterations. J Bacteriol Mycol Open Access 4:85.

12.   Ledesma-Amaro R, Nicaud J-M. 2016. Metabolic engineering for expanding the

substrate range of *Yarrowia lipolytica*. Trends in Biotechnology 34:798-809.

13.   Zhao C, Zhang Y, Li Y. 2019. Production of fuels and chemicals from renewable

resources using engineered *Escherichia coli*. Biotechnology advances 37:107402.

14.   Chen C, Lin J, Wang W, Huang H, Li S. 2019. Cost-effective production of surfactin

from xylose-rich corncob hydrolysate using *Bacillus subtilis* BS-37. Waste and Biomass

Valorization 10:341-347.

15.   Straub CT, Bing RG, Wang JP, Chiang VL, Adams MW, Kelly RM. 2020. Use of the

lignocellulose-degrading bacterium *Caldicellulosiruptor bescii* to assess recalcitrance and

conversion of wild-type and transgenic poplar. Biotechnology for biofuels 13:1-10.

16.   Sgobba E, Blöbaum L, Wendisch VF. 2018. Production of food and feed additives from

non-food-competing feedstocks: Valorizing N-acetylmuramic acid for amino acid and carotenoid

fermentation with *Corynebacterium glutamicum*. Frontiers in Microbiology 9:2046.

17.   Zhu X, Zhou Y, Wang Y, Wu T, Li X, Li D, Tao Y. 2017. Production of high-

concentration n-caproic acid from lactate through fermentation using a newly isolated

Ruminococcaceae bacterium CPB6. Biotechnology for biofuels 10:102.

18.   Li Q. 2019. Progress in microbial degradation of feather waste. Frontiers in Microbiology

10:2717.

19.   Kang D, Huang Y, Nesme J, Herschend J, Jacquiod S, Kot W, Hansen LH, Lange L,

Sørensen SJ. 2020. Metagenomic analysis of a keratin-degrading bacterial consortium provides

insight into the keratinolytic mechanisms. Science of The Total Environment:143281.

20.   Huang Y, Łężyk M, Herbst F-A, Busk PK, Lange L. 2020. Novel keratinolytic enzymes,

discovered from a talented and efficient bacterial keratin degrader. Scientific reports 10:1-11.

21.   Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, Medema MH, Weber T. 2019.

antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. Nucleic acids

research 47:W81-W87.

22.   Weber T, Kim HU. 2016. The secondary metabolite bioinformatics portal: Computational

tools to facilitate synthetic biology of secondary metabolite production. Synthetic and Systems

Biotechnology 1:69-79.

23.   Zheng Y, Saitou A, Wang C-M, Toyoda A, Minakuchi Y, Sekiguchi Y, Ueda K, Takano

H, Sakai Y, Abe K. 2019. Genome features and secondary metabolites biosynthetic potential of

the class *Ktedonobacteria*. Frontiers in microbiology 10:893.

24.   Kjærbølling I, Vesth TC, Frisvad JC, Nybo JL, Theobald S, Kuo A, Bowyer P, Matsuda

Y, Mondo S, Lyhne EK. 2018. Linking secondary metabolites to gene clusters through genome

sequencing of six diverse *Aspergillus* species. Proceedings of the National Academy of Sciences

115:E753-E761.

25.   Fontoura R, Daroit DJ, Corrêa APF, Moresco KS, Santi L, Beys-da-Silva WO, Yates III

JR, Moreira JCF, Brandelli A. 2019. Characterization of a novel antioxidant peptide from feather

keratin hydrolysates. New biotechnology 49:71-76.

26.   Kshetri P, Roy SS, Sharma SK, Singh TS, Ansari MA, Prakash N, Ngachan S. 2019.

Transforming chicken feather waste into feather protein hydrolysate using a newly isolated

multifaceted keratinolytic bacterium *Chryseobacterium sediminis* RCM-SSR-7. Waste and

Biomass Valorization 10:1-11.

27.   Kang D, Jacquiod S, Herschend J, Wei S, Nesme J, Sørensen SJ. 2020. Construction of

simplified microbial consortia to degrade recalcitrant materials based on enrichment and

dilution-to-extinction cultures. Frontiers in microbiology 10:3010.

28.   Kang D, Herschend J, Al-Soud WA, Mortensen MS, Gonzalo M, Jacquiod S, Sørensen

SJ. 2018. Enrichment and characterization of an environmental microbial consortium displaying

efficient keratinolytic activity. Bioresource technology 270:303-310.

29.   Kim E-M, Hwang KH, Park J-S. 2018. Complete Genome Sequence of

Chryseobacterium camelliae Dolsongi-HT1, a Green Tea Isolate with Keratinolytic Activity.

Genome Announcements 6.

30. Park G-S, Hong S-J, Jung BK, Khan AR, Park Y-J, Park CE, Lee A, Kwak Y, Lee Y-J,

Lee D-W. 2015. Complete genome sequence of a keratin-degrading bacterium *Chryseobacterium*

gallinarum strain DSM 27622T isolated from chicken. Journal of biotechnology 211:66-67.

31. Park G-S, Hong S-J, Lee C-H, Khan AR, Ullah I, Jung BK, Choi J, Kwak Y, Back C-G,

Jung H-Y. 2014. Draft genome sequence of *Chryseobacterium* sp. strain P1-3, a keratinolytic

bacterium isolated from poultry waste. Genome announcements 2.

32. Lin P-C, Pakrasi HB. 2019. Engineering cyanobacteria for production of terpenoids.

Planta 249:145-154.

33. Barajas JF, Blake-Hedges JM, Bailey CB, Curran S, Keasling JD. 2017. Engineered

polyketides: synergy between protein and host level engineering. Synthetic and systems

biotechnology 2:147-166.

34. Wang W, Li S, Li Z, Zhang J, Fan K, Tan G, Ai G, Lam SM, Shui G, Yang Z. 2020.

Harnessing the intracellular triacylglycerols for titer improvement of polyketides in *Streptomyces*.

Nature Biotechnology 38:76-83.

35. Adamek M, Spohn M, Stegmann E, Ziemert N. 2017. Mining bacterial genomes for

secondary metabolite gene clusters, p 23-47, Antibiotics. Springer.

36. Reichenbach H, Kohl W, Böttger-Vetter A, Achenbach H. 1980. Flexirubin-type

pigments in *Flavobacterium*. Archives of Microbiology 126:291-293.

37. Vandamme P, Bernardet J-F, Segers P, Kersters K, Holmes B. 1994. New Perspectives in

the Classification of the Flavobacteria: Description of *Chryseobacterium* gen. nov., *Bergeyella*

gen. nov., and *Empedobacter* nom. rev. International Journal of Systematic and Evolutionary

Microbiology 44:827-831.

38. Venil CK, Zakaria ZA, Usha R, Ahmad WA. 2014. Isolation and characterization of

flexirubin type pigment from *Chryseobacterium* sp. UTM-3T. Biocatalysis and Agricultural

Biotechnology 3:103-107.

39. Javidpour P, Deutsch S, Mutalik VK, Hillson NJ, Petzold CJ, Keasling JD, Beller HR.

2016. Investigation of proposed ladderane biosynthetic genes from anammox bacteria by

heterologous expression in *E. coli*. PloS one 11:e0151087.

40. Mercer JA, Cohen CM, Shuken SR, Wagner AM, Smith MW, Moss III FR, Smith MD,

Vahala R, Gonzalez-Martinez A, Boxer SG. 2016. Chemical synthesis and self-assembly of a

ladderane phospholipid. Journal of the American Chemical Society 138:15845-15848.

41. Adamek M, Alanjary M, Sales-Ortells H, Goodfellow M, Bull AT, Winkler A, Wibberg

D, Kalinowski J, Ziemert N. 2018. Comparative genomics reveals phylogenetic distribution

patterns of secondary metabolites in *Amycolatopsis* species. BMC genomics 19:1-15.

42. Venil CK, Zakaria ZA, Ahmad WA. 2013. Bacterial pigments and their applications.

Process Biochemistry 48:1065-1079.

43. Aruldass CA, Dufossé L, Ahmad WA. 2018. Current perspective of yellowish-orange

pigments from microorganisms-a review. Journal of Cleaner Production 180:168-182.

44. Philmus B, Christiansen G, Yoshida WY, Hemscheidt TK. 2008. Post-translational

modification in microviridin biosynthesis. ChemBioChem 9:3066-3073.

45. Gatte-Picchi D, Weiz A, Ishida K, Hertweck C, Dittmann E. 2014. Functional analysis of

environmental DNA-derived microviridins provides new insights into the diversity of the

tricyclic peptide family. Applied and environmental microbiology 80:1380-1387.

46.   Hassan MA, Abol-Fotouh D, Omer AM, Tamer TM, Abbas E. 2020. Comprehensive

insights into microbial keratinases and their implication in various biotechnological and

industrial sectors: A review. International Journal of Biological Macromolecules.

47.   Khan A, Singh P, Srivastava A. 2018. Synthesis, nature and utility of universal iron

chelator–Siderophore: A review. Microbiological research 212:103-111.

48.   Boda SK, Pandit S, Garai A, Pal D, Basu B. 2016. Bacterial siderophore mimicking iron

complexes as DNA targeting antimicrobials. RSC advances 6:39245-39260.

49.   Nakouti I, Sihanonth P, Palaga T, Hobbs G. 2013. Effect of a siderophore producer on

animal cell apoptosis: a possible role as anti-cancer agent. International Journal of Pharma

Medicine and Biological Sciences 2:1-5.

50.   REMINGTON SJ. 1992. Structure and mechanism of citrate synthase, p 209-229,

Current topics in cellular regulation, vol 33. Elsevier.

51.   Cronan JE, Thomas J. 2009. Bacterial fatty acid synthesis and its relationships with

polyketide synthetic pathways. Methods in enzymology 459:395-433.

52.   Park Y-K, Dulermo T, Ledesma-Amaro R, Nicaud J-M. 2018. Optimization of odd chain

fatty acid production by *Yarrowia lipolytica*. Biotechnology for biofuels 11:1-12.

53.   Park Y-k, Ledesma-Amaro R, Nicaud J-M. 2020. De novo biosynthesis of odd-chain

fatty acids in *Yarrowia lipolytica* enabled by modular pathway engineering. Frontiers in

Bioengineering and Biotechnology 7:484.

54.   Nasipuri P, Herschend J, Brejnrod AD, Madsen JS, Espersen R, Svensson B, Burmølle M,

Jacquiod S, Sørensen SJ. 2020. Community-intrinsic properties enhance keratin degradation

from bacterial consortia. PloS one 15:e0228108.

55. Ramnani P, Gupta R. 2007. Keratinases vis-à-vis conventional proteases and feather

degradation. World Journal of Microbiology and Biotechnology 23:1537-1540.

56. Navone L, Speight R. 2018. Understanding the dynamics of keratin weakening and

hydrolysis by proteases. PLoS One 13:e0202608.

57. Huang Y, Busk PK, Herbst F-A, Lange L. 2015. Genome and secretome analyses provide

insights into keratin decomposition by novel proteases from the non-pathogenic fungus *Onygena*

corvina. Applied microbiology and biotechnology 99:9635-9649.

58. Li Z-W, Liang S, Ke Y, Deng J-J, Zhang M-S, Lu D-L, Li J-Z, Luo X-C. 2020. The

feather degradation mechanisms of a new *Streptomyces* sp. isolate SCUT-3. Communications

Biology 3:1-13.

59. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for

genome assemblies. Bioinformatics 29:1072-1075.

60. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics

30:2068-2069.

61. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H,

Mende DR, Letunic I, Rattei T, Jensen LJ. 2019. eggNOG 5.0: a hierarchical, functionally and

phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses.

Nucleic acids research 47:D309-D314.

62. Bertels F, Silander OK, Pachkov M, Rainey PB, van Nimwegen E. 2014. Automated

reconstruction of whole-genome phylogenies from short-sequence reads. Molecular biology and

evolution 31:1077-1088.

63. Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic acids research 47:W256-W259.

64.   Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison visualizer.

Bioinformatics 27:1009-1010.

65.   Sievers F, Higgins DG. 2018. Clustal Omega for making accurate alignments of many

protein sequences. Protein Science 27:135-145.

66.   Thomsen MCF, Nielsen M. 2012. Seq2Logo: a method for construction and visualization

of amino acid binding motifs and sequence profiles including sequence weighting, pseudo counts

and two-sided representation of amino acid enrichment and depletion. Nucleic acids research

40:W281-W287.

67.   Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG tools

for functional characterization of genome and metagenome sequences. Journal of molecular

biology 428:726-731.

68.   Kanehisa M, Araki M, Goto S, Hattori M, Hirakawa M, Itoh M, Katayama T, Kawashima

S, Okuda S, Tokimatsu T. 2007. KEGG for linking genomes to life and the environment. Nucleic

acids research 36:D480-D484.

69.   Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA, Holland TA,

Keseler IM, Kothari A, Kubo A. 2014. The MetaCyc database of metabolic pathways and

enzymes and the BioCyc collection of Pathway/Genome Databases. Nucleic acids research

42:D459-D471.

Fig. 1. Analysis of the *Chryseobacterium* spp. genomes. (a). The whole-genome sequence-based phylogenetic tree of *Chryseobacterium* spp., constructed by REALPHY 1.12 (62), based on the merge reference alignments of all genomes. Branch length represents divergence. (b). Comparison of KEGG function classification amongst four *Chryseobacterium* spp. genomes. Functional categories: Metabolism (A), Genetic information processing (B), Environmental

575   information processing (C), and Cellular processes (D). The stars show the sub-categories:

576   Amino acid metabolism, metabolism of terpenoids and polyketides, and biosynthesis of other

577   secondary metabolites.

Fig. 2. Composition of secondary metabolite gene clusters from four *Chryseobacterium* spp.

genomes.

Fig. 3. Flexirubin-type pigment gene cluster from four *Chryseobacterium* spp. genomes. (a).

Synteny analysis and features of flexirubin-type pigment gene cluster in *Chryseobacterium* spp.

genomes. (b). The proposed biosynthetic reaction of flexirubin-type pigment. The detailed

description of each gene can be found in Table S1.

Fig. 4. Microviridin gene cluster from four *Chryseobacterium* spp. genomes. (a). Synteny

analysis and features of microviridin gene cluster in *Chryseobacterium* spp. genomes. (b).

Amino acid sequence comparison of mdnA from four *Chryseobacterium* spp. genomes. (c).

Amino acid sequence comparison of mdnB from four *Chryseobacterium* spp. genomes. The

detailed description of each gene can be found in Table S2.

Fig. 5. Siderophore gene cluster from four *Chryseobacterium* spp. genomes. The detailed description of each gene can be found in Table S3.

Fig. 6. Metabolic pathways reconstruction of keratin utilization in *Chryseobacterium* sp.

KMC2 genome. It includes amino acid metabolism, TCA cycle, glycolysis/gluconeogenesis

propanoate metabolism, and sulfur metabolism. The number in the box represents the gene

related to the metabolic pathway. The detailed description of each gene can be found in Table S7.

Table 1. Features comparison of four *Chryseobacterium* spp. genomes.

Parameters	*Chryseobacterium* sp. KMC2	*Chryseobacterium camelliae* Dolsongi-HT1	*Chryseobacterium gallinarum* strain DSM 27622	*Chryseobacterium* sp. P1-3
Total length (bp)	5.276.159	4.376.354	4.633.632	4.628.764
Contigs	63	1	1	45
N50 (bp)	231.784	4.376.354	4.633.632	342.512
GC content (%)	36.33	41.80	37.30	37.02
Gene	4.773	3.999	4.182	4.087
CDS	4.692	3.881	4.033	3.119